

FIG. 1. Comparison of the amino acid sequence of GES-3 with those of the GES-1, GES-2, and IBC-1 β -lactamases. Only the substituted amino acid residues are indicated by the single-letter amino acid code. Dashes represent positions where no amino acid substitution was observed among the four enzymes. The amino acid residues conserved among class A β -lactamases are indicated with asterisks.

Genetic environment of *bla*_{GES-3}. Two distinct BamHI fragments carrying the *bla*_{GES-3} gene were cloned; their sizes were approximately 6.7 and 11.6 kb, respectively, and pKGB525 had the 6.7-kb fragment. Sequencing analysis of the entire insert on pKGB525 revealed that the *bla*_{GES-3} gene was located as a gene cassette within a class 1 integron structure, as was observed in the other GES-type β -lactamase genes, *bla*_{GES-1}, *bla*_{GES-2}, and *bla*_{IBC-1} (Fig. 2). The 59-base element downstream of the *bla*_{GES-3} gene was made up of 110 bp and was different from that of the truncated 59-base element of *bla*_{GES-1} on pTK1 (GenBank accession number AF156486), but shared it 99% nucleotide identity with those of *bla*_{GES-2} on pLAP-1 (GenBank accession number AF326355) and *bla*_{IBC-1} on pHT9-2 (GenBank accession number AF208529).

The second gene cassette adjacent to the *bla*_{GES-3} gene was a fused *aacA1-orfG* gene cassette. The results of the disk diffusion test indicated that the presence of an *aacA1* component, which encodes aminoglycoside-6'-N-acetyltransferase, conferred kanamycin resistance to the transformant *E. coli* XL1-Blue(pKGB525) (data not shown). The nucleotide sequence of this fused gene cassette shared 100% identity with that in a class 1 integron on plasmid pCMXR1 (GenBank accession number AB061794). The function of the product encoded by *orfG* has not been characterized in detail. The third gene cassette is 327 bp and was named *orfA*. The *orfA* gene was

suggested to be a gene cassette by recognition of the features typical of these elements: (i) the presence at the cassette boundaries of 7-bp core site sequences that completely fit the consensus sequence and (ii) the presence of a 59-base element of 78 bp downstream of the *orfA* gene. However, no remarkable similarity between the hypothetical protein encoded by *orfA* and any other known protein sequences was detected in a search performed with the BLAST program. The 3' conserved segment of this integron showed a characteristic organization. The *qacEΔ1* gene was truncated at nucleotide position 114 by the insertion sequence IS26. In the region downstream of IS26, an IS6100 element and two ORFs of unknown function, i.e., *orf5* and *orf6*, were found. The nucleotide sequence of the 1.4-kb region containing IS6100, *orf5*, and *orf6* was identical to that seen downstream of the *sulI* gene of the class 1 integron in the chromosomal multidrug resistance locus of *Salmonella enterica* subsp. *enterica* serovar Typhimurium (GenBank accession number AF261825). Sequencing analysis of pKGM525 carrying the second *bla*_{GES-3} gene on an 11.6-kb fragment was also done, and the *bla*_{GES-3} gene was also found in a class 1 integron structure with the same gene cassette configuration as in pKGB525. The nucleotide sequence of the region from *intI1* to IS26 was the same as that found in pKGB525 carrying a 6.7-kb BamHI fragment.

Construction of IBC-1 by site-directed mutagenesis. Only one amino acid substitution, M62T, was found between the sequences of GES-3 and IBC-1 (Fig. 1). Therefore, to examine whether this amino acid substitution affects the MICs of β -lactams for the *E. coli* clones producing each enzyme, we constructed plasmid pIBC1, which encodes the IBC-1 enzyme under the same promoter as that for GES-3, by site-directed mutagenesis of the *bla*_{GES-3} gene within the parental plasmid, pGES3. However, this single substitution did not markedly influence the MICs for the *E. coli* clones (data not shown).

PCR detection and genotypic comparison. The remaining five nonrepetitive ceftazidime-resistant *K. pneumoniae* strains, strains KG914, KG1116, KG545, KG502, and KG827, were all found to be *bla*_{GES} positive by PCR. The results of PFGE analysis of all six isolates are shown in Fig. 3. Their fingerprinting patterns were very similar but in some cases were distinct. We examined the fingerprinting patterns from 48.5 to 194 kb in detail under other conditions (data not shown). Overall, there were from three to seven band differences among the six strains examined. The ERIC-PCR patterns amplified with the ERIC-2 primer were indistinguishable from one another (data not shown). Taken together with the fact that these isolates were collected over a 1-year period, we speculate that they were genetically related and had probably spread via nosocomial transmission of an endemic clone.

Plasmid profiles and Southern hybridization. The plasmid profiles of the six *bla*_{GES}-positive strains showed the presence of a large plasmid of approximately similar size in five of the six strains (Fig. 4A), while some of them possessed additional plasmids which were smaller and more diverse in size. Hybridization analyses with the probe specific for the GES-type β -lactamase genes, including *bla*_{IBC-1}, revealed that the location of this gene varied among the strains tested. Hybridization signals for large plasmids were detected for strains KG914, KG1116, and KG502 (Fig. 4B). Hybridization signals for both plasmids and chromosomal positions were observed for

TABLE 3. MICs of antimicrobial agents for the parental strain, transconjugant, and transformant

β-Lactam ^a	MIC (μg/ml)				
	<i>K. pneumoniae</i> KG525	<i>E. coli</i> CSH- 2(pKGC525) ^b	<i>E. coli</i> CSH-2	<i>E. coli</i> XL1- Blue(pKGB525) ^c	<i>E. coli</i> XL1- Blue(pBCSK+)
Ampicillin	>128	>128	4	>128	4
Ampicillin + sulbactam	>128	2	2	2	2
Amoxicillin	>128	>128	8	>128	4
Amoxicillin + clavulanate	>128	32	4	32	4
Piperacillin	128	16	1	16	1
Piperacillin + tazobactam	128	0.5	1	0.5	0.5
Cefotaxime	64	2	0.13	2	0.13
Cefotaxime + clavulanate	8	0.06	0.06	0.06	0.06
Cefotaxime + sulbactam	32	0.06	0.06	0.06	0.06
Cefotaxime + tazobactam	64	0.06	0.06	0.06	0.06
Ceftazidime	>1,024	128	0.13	128	0.13
Ceftazidime + clavulanate	256	4	0.13	4	0.06
Ceftazidime + sulbactam	>128	0.25	0.13	0.5	0.13
Ceftazidime + tazobactam	>128	0.5	0.13	0.5	0.13
Cephaloridine	>128	16	2	16	2
Cefminox	8	0.5	0.5	1	0.5
Moxalactam	4	0.25	0.13	0.5	0.13
Cefpirome	>128	1	0.06	2	0.06
Cefepime	16	0.13	0.06	0.25	0.06
Aztreonam	64	4	0.06	4	0.06
Imipenem	0.25	0.5	0.25	0.13	0.13
Gentamicin	2	0.13	0.13	<0.06	<0.06
Amikacin	64	2	0.25	4	0.25
Levofloxacin	<0.06	0.13	0.13	<0.06	<0.06
Ciprofloxacin	<0.06	<0.06	<0.06	<0.06	<0.06

^a Clavulanate, tazobactam, and sulbactam were used at a fixed concentration of 4 μg/ml each.

^b pKGC525 is a resident plasmid found in *K. pneumoniae* strain KG525, and it carries the *bla*_{GES-3} gene.

^c pKGB525 is a recombinant plasmid that carries a 6.7-kb BamHI insert that mediates the *bla*_{GES-3} gene.

KG525. One of the hybridized plasmids from each of KG914, KG1116, and KG525 were similar in size. A single hybridization signal corresponding to the chromosomal position was detected for each of the strains KG545 and KG827 (Fig. 4C).

pIs of β-lactamases. The pI value of the purified GES-3 enzyme was determined to be 7.0 (Fig. 5A). IEF of crude extracts from six GES-type β-lactamase-producing clinical strains revealed two bands with pIs of 7.6 and 7.0 (Fig. 5B). The band with pI 7.0 was also detected in a GES-3-producing

E. coli transformant which harbored the *bla*_{GES-3} gene of *K. pneumoniae* strain KG525. The band with a pI of 7.6 corresponds to the chromosomally encoded LEN-1 (1) or SHV-type penicillinase of *K. pneumoniae*.

DISCUSSION

Considerable differences in the levels of resistance to various cephalosporins were observed among the *E. coli* clones pro-




FIG. 2. Schematic comparison of the class 1 integron on pKGB525 with those on pTK1 (GenBank accession number AF156486), pLAP-1 (GenBank accession number AF326355), and pAK33 (34). Filled circles indicate the positions of GTTRRRY (core site) or the 59-base elements around the gene cassettes. pKGM525, which carries the 11.6-kb BamHI insert, was also sequenced; and the nucleotide sequence from *intI1* to *IS26* was the same as that found in pKGB525.




FIG. 3. PFGE analysis of *K. pneumoniae* isolates. Lanes: 1 and 8, PFGE marker; 2, *K. pneumoniae* KG914; 3, *K. pneumoniae* KG1116; 4, *K. pneumoniae* KG525; 5, *K. pneumoniae* KG545; 6, *K. pneumoniae* KG502; 7, *K. pneumoniae* KG827.

ducing GES-1, GES-2, and IBC-1, although the level of production of each enzyme may differ in individual clones. For instance, the MIC of ceftazidime for an *E. coli* clone producing GES-1 was 128 $\mu\text{g/ml}$ (24), while that for an *E. coli* clone producing GES-2 was 8 $\mu\text{g/ml}$ (28). The single amino acid substitution in the Ω loop observed between GES-1 and GES-2, G170N, may well contribute to the difference in the




FIG. 4. Plasmid profiles and Southern hybridization analysis. (A) Plasmid profiles of each strain prepared by the method of Kado and Liu (16); (B) hybridization to large plasmids harbored by each strain; (C) hybridization to the chromosomal position of each strain. The photograph of the results of gel electrophoresis of chromosomal DNAs prepared by the method of Stauffer et al. (35) was omitted. The large plasmids and chromosomal DNA were separately extracted by using freshly prepared reagents to avoid cross contamination of nicked or physicochemically amputated DNA fragments. For strains KG545 and KG827, the bla_{GES} gene was suggested to be encoded by the chromosome. In strain KG525, the bla_{GES} gene was suggested to be encoded by both the plasmid and the chromosome. Lanes: M, HindIII-digested DNA marker; 1, *K. pneumoniae* KG914; 2, *K. pneumoniae* KG1116; 3, *K. pneumoniae* KG525; 4, *K. pneumoniae* KG545; 5, *K. pneumoniae* KG502; 6, *K. pneumoniae* KG827.

substrate specificities of these enzymes. On the other hand, the MIC of ceftazidime for an *E. coli* clone producing IBC-1 was $>256 \mu\text{g/ml}$ (11). The only amino acid substitution observed between GES-1 and IBC-1, E104K, might well also be attributed to a higher level of resistance of IBC-1 than that of GES-1 to ceftazidime, cefotaxime, and aztreonam (11, 19, 24). The MIC of ceftazidime for an *E. coli* clone producing GES-3 was 128 $\mu\text{g/ml}$, and GES-3 has a single M62T substitution compared with the sequence of IBC-1, which also confers high-level resistance to ceftazidime in an *E. coli* clone (MIC, $>256 \mu\text{g/ml}$). We investigated whether this one amino acid substitution observed between GES-3 and IBC-1 affects the MICs for *E. coli* clones expressing each enzyme. However, this substitution did not result in significant changes in the MICs for the *E. coli* clones. This finding suggests that the amino acid substitution at position 62 may not play a crucial role in the extended substrate specificity of GES-3 against ceftazidime and that those at positions 104 and 170 would be crucial for extended-spectrum enzyme activity.

In the present study, we also isolated a novel GES-type class A enzyme, GES-3, from *K. pneumoniae* strains which caused neonatal nosocomial infections in 2002 in Japan. Sequence analysis of the genetic environments of the $bla_{\text{GES-3}}$ genes on pKGB525 carrying a 6.7-kb insert and pKGM525 carrying a 11.6-kb insert revealed that the $bla_{\text{GES-3}}$ genes were located as gene cassettes in class 1 integrons, as observed in other GES-type β -lactamase genes, including $bla_{\text{IBC-1}}$ (Fig. 2). Integrons are very sophisticated site-specific recombination systems that capture various gene cassettes, including antibiotic resistance genes, between their 5' and 3' conserved segments (14, 15, 22). The gene cassettes for the GES-type enzymes with a very close phylogenetic relationship might have originated as a single clone and then disseminated worldwide with the help of class 1 integrons possessing very similar genetic organizations. These integrons are mediated by self-transmissible plasmids with a wide host range. Since very similar GES enzymes have so far been found in French Guiana, Greece, South Africa, Portugal, and Japan, these GES-type β -lactamase-producing strains might have been scattered globally by the recent extensive international travel or dissemination of humans, foods, and animals.

In the present study, we analyzed genetic relatedness using PFGE and ERIC-PCR of all six GES-type β -lactamase-producing *K. pneumoniae* strains isolated in a NICU over a 1-year period. Since the fingerprinting patterns obtained by PFGE and ERIC-PCR were very similar, these isolates were suggested to belong to the same genetic lineage that caused the nosocomial spread. The minor differences in the fingerprinting patterns obtained by PFGE might be due to the occurrence of genetic rearrangements over the course of the nosocomial spread. Interestingly, the results of Southern hybridization suggested that the locations of the bla_{GES} genes were multiple and diverse among the six strains studied. By consideration of the results of PFGE, ERIC-PCR, and Southern hybridization, it can be speculated that an endemic strain containing the bla_{GES} genes might have spread within the NICU setting over the 1-year period and might have undergone genetic rearrangements, including translocation and multiplication of the bla_{GES} gene.

The presence of multiple $bla_{\text{GES-3}}$ genes in strain KG525 is




FIG. 5. IEF. (A) IEF and staining with Coomassie blue. Lanes: M, pI marker; GES-3, purified GES-3 enzyme. (B) IEF and staining with nitrocefin. Lanes: 1, GES-3-producing *E. coli* transformant that harbors pKGB525 carrying *bla*_{GES-3}; 2, *K. pneumoniae* KG914; 3, *K. pneumoniae* KG1116; 4, *K. pneumoniae* KG525; 5, *K. pneumoniae* KG545; 6, *K. pneumoniae* KG502; 7, *K. pneumoniae* KG827. The bands of pI 7.63 (pI 7.6 in the text) are the chromosomally encoded LEN-1 or SHV-1 β -lactamase of *K. pneumoniae*, and the bands at pI 6.98 (pI 7.0 in the text) are GES-3. Several β -lactamases with activities at pIs lower than 7.0 were speculated to be partially unfolded GES-3 β -lactamase, because these bands were also found in the IEF gels of an *E. coli* clone harboring only the *bla*_{GES-3} gene (data not shown).

probably the result of these genetic rearrangements. Translocation and multiplication of the *bla*_{GES} gene might be facilitated by mediation of a site-specific recombination system of an integron or a transposon. A similar example of multiple locations of the same antibiotic resistance gene has been reported by Yagi et al. (38). In their study, a single clinical *E. coli* isolate was found to carry three distinguishable Toho-1-like β -lactamase genes, which were later identified as *bla*_{CTX-M-2} by their restriction digestion patterns on the chromosome. These multiple locations of the same β -lactamase gene would be beneficial to bacteria, since they increase the chance of amino acid substitutions necessary for extension of the substrate profiles of β -lactamases as well as the multicopy effect of gene expression. Bradford et al. (3) reported that point mutations leading to ESBLs (ESBLs TEM-1 to TEM-10 and TEM-12) occurred on the plasmids of a single *K. pneumoniae* clinical isolate. A notable finding presented in that report was the distinct hydrolyzing activity between TEM-10 and TEM-12. TEM-10 had hydrolyzing activity against ceftazidime, while TEM-12 also hydrolyzed cefotaxime and aztreonam, in addition to ceftazidime. A variety of susceptibility profiles for cephamycins were also observed among the six *K. pneumoniae* strains in the present study. For instance, strains KG914, KG1116, and KG525 were susceptible to cefmetazole (MICs, 16 μ g/ml), whereas strains KG545, KG502, and KG827 were resistant to this agent (MICs, \geq 128 μ g/ml). In particular, strain KG502 showed high-level resistance to other cephamycins, such as ceftoxitin (MIC, >128 μ g/ml), cefminox (MIC, >128 μ g/ml), and moxalactam (MIC, 128 μ g/ml). An evolutionary event similar to that observed in the TEM enzymes (3) might have occurred in these *K. pneumoniae* strains to give them further resistance to a broad range of antibiotics. The MIC of imipenem for strain KG502 was 8 μ g/ml, and this might be due to the hyperproduction of some β -lactamase with an extended

substrate specificity as well as the loss of some outer membrane protein, as reported by Bradford et al. (4). Further molecular characterization of the cephamycin resistance observed in strain KG502 will be undertaken in the next study.

ACKNOWLEDGMENTS

We are grateful to Leonidas S. Tzouveleakis, Laboratory of Bacteriology, Hellenic Pasteur Institute, Athens, Greece, for kindly providing *E. cloacae* HT9-producing IBC-1 and *E. coli* DH5 α containing pHIT9-2 and to Kumiko Kai for technical help.

This work was supported by grants H12-Shinko-19 and H12-Shinko-20 from the Ministry of Health, Labor and Welfare of Japan.

REFERENCES

1. Arakawa, Y., M. Ohta, N. Kido, Y. Fujii, T. Komatsu, and N. Kato. 1986. Close evolutionary relationship between the chromosomally encoded β -lactamase gene of *Klebsiella pneumoniae* and the TEM β -lactamase gene mediated by R-plasmids. *FEBS Lett.* 207:69-74.
2. Arakawa, Y., N. Shibata, K. Shibayama, H. Kurokawa, T. Yagi, H. Fujiwara, and M. Goto. 2000. Convenient test for screening metallo- β -lactamase-producing gram-negative bacteria by using thiol compounds. *J. Clin. Microbiol.* 38:40-43.
3. Bradford, P. A., C. E. Cherubin, V. Idemiyor, B. A. Rasmussen, and K. Bush. 1994. Multiply resistant *Klebsiella pneumoniae* strains from two Chicago hospitals: identification of the extended-spectrum TEM-12 and TEM-10 ceftazidime-hydrolyzing β -lactamases in a single isolate. *Antimicrob. Agents Chemother.* 38:761-766.
4. Bradford, P. A., C. Urban, N. Mariano, S. J. Projan, J. J. Rahal, and K. Bush. 1997. Imipenem resistance in *Klebsiella pneumoniae* is associated with the combination of ACT-1, a plasmid-mediated AmpC β -lactamase, and the loss of an outer membrane protein. *Antimicrob. Agents Chemother.* 41:563-569.
5. Burwen, D. R., S. N. Banerjee, R. P. Gaynes, et al. 1994. Ceftazidime resistance among selected nosocomial gram-negative bacilli in the United States. *J. Infect. Dis.* 170:1622-1625.
6. Bush, K., G. A. Jacoby, and A. A. Medeiros. 1995. A functional classification scheme for β -lactamases and its correlation with molecular structure. *Antimicrob. Agents Chemother.* 39:1211-1233.
7. Chaves, J., M. G. Ladona, C. Segura, A. Coira, R. Reig, and C. Ampurdanes. 2001. SHV-1 β -lactamase is mainly a chromosomally encoded species-specific enzyme in *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 45:2856-2861.

8. Doi, Y., N. Shibata, K. Shibayama, K. Kamachi, H. Kurokawa, K. Yokoyama, T. Yagi, and Y. Arakawa. 2002. Characterization of a novel plasmid-mediated cephalosporinase (CMY-9) and its genetic environment in an *Escherichia coli* clinical isolate. *Antimicrob. Agents Chemother.* 46:2427-2434.
9. Duarte, A., F. Boavida, F. Grosso, M. Correia, L. M. Lito, J. M. Cristino, and M. J. Saigado. 2003. Outbreak of GES-1 β -lactamase-producing multidrug-resistant *Klebsiella pneumoniae* in a university hospital in Lisbon, Portugal. *Antimicrob. Agents Chemother.* 47:1481-1482.
10. Dubois, V., L. Poirel, C. Marie, C. Arpin, P. Nordmann, and C. Quentin. 2002. Molecular characterization of a novel class 1 integron containing *bla*_{GES-1} and a fused product of *aac3-Ib/aac6'-Ib'* gene cassettes in *Pseudomonas aeruginosa*. *Antimicrob. Agents Chemother.* 46:638-645.
11. Giakkoupi, P., L. S. Tzouveleki, A. Tsakris, V. Loukova, D. Sofianou, and E. Tzelepi. 2000. IBC-1, a novel integron-associated class A β -lactamase with extended-spectrum properties produced by an *Enterobacter cloacae* clinical strain. *Antimicrob. Agents Chemother.* 44:2247-2253.
12. Girlich, D., L. Poirel, A. Leclaporn, A. Karim, C. Tribuddharat, M. Fennewald, and P. Nordmann. 2001. Molecular epidemiology of the integron-located VEB-1 extended-spectrum β -lactamase in nosocomial enterobacterial isolates in Bangkok, Thailand. *J. Clin. Microbiol.* 39:175-182.
13. Gniadkowski, M., I. Schneider, A. Palucha, R. Jungwirth, B. Mikiewicz, and A. Bauernfeind. 1998. Cefotaxime-resistant *Enterobacteriaceae* isolates from a hospital in Warsaw, Poland: identification of a new CTX-M-3 cefotaxime-hydrolyzing β -lactamase that is closely related to the CTX-M-1/MEN-1 enzyme. *Antimicrob. Agents Chemother.* 42:827-832.
14. Hall, R. M., D. E. Brookes, and H. W. Stokes. 1991. Site-specific insertion of genes into integrons: role of the 59-base element and determination of the recombination cross-over point. *Mol. Microbiol.* 5:1941-1959.
15. Hall, R. M., and C. M. Collis. 1995. Site-specific insertion of genes into integrons: role of 59-base element and determination of the recombination cross-over point. *Mol. Microbiol.* 15:593-600.
16. Kado, C. I., and S. T. Liu. 1981. Rapid procedure for detection and isolation of large and small plasmids. *J. Bacteriol.* 145:1365-1373.
17. Kartali, G., E. Tzelepi, S. Pournaras, C. Kontopoulou, F. Kontos, D. Sofianou, A. N. Muniatis, and A. Tsakris. 2002. Outbreak of infections caused by *Enterobacter cloacae* producing the integron-associated β -lactamase IBC-1 in a neonatal intensive care unit of a Greek hospital. *Antimicrob. Agents Chemother.* 46:1577-1580.
18. Matsumoto, Y., and M. Inoue. 1999. Characterization of SFO-1, a plasmid-mediated inducible class A β -lactamase from *Enterobacter cloacae*. *Antimicrob. Agents Chemother.* 43:307-313.
19. Mavroidi, A., E. Tzelepi, A. Tsakris, V. Miriagou, D. Sofianou, and L. S. Tzouveleki. 2001. An integron-associated β -lactamase (IBC-2) from *Pseudomonas aeruginosa* is a variant of the extended-spectrum β -lactamase IBC-1. *J. Antimicrob. Chemother.* 48:627-630.
20. Naas, T., F. Benabdou, S. Massuard, and P. Nordmann. 2000. Integron-located VEB-1 extended-spectrum β -lactamase gene in a *Proteus mirabilis* clinical isolate from Vietnam. *J. Antimicrob. Chemother.* 46:703-711.
21. National Committee for Clinical Laboratory Standards. 2003. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 5th ed. Document M7-A5. National Committee for Clinical Laboratory Standards, Wayne, Pa.
22. Paulsen, I. T., T. G. Littlejohn, P. Radstrom, L. Sundstrom, O. Skold, G. Swedberg, and R. A. Skurray. 1993. The 3' conserved segment of integrons contains a gene associated with multidrug resistance to antiseptics and disinfectants. *Antimicrob. Agents Chemother.* 37:761-768.
23. Petit, A., D. L. Strot, C. M. Chanal, J. L. Sirot, R. Labia, G. Gerband, and R. A. Cluzel. 1988. Novel plasmid-mediated β -lactamase in clinical isolates of *Klebsiella pneumoniae* more resistant to ceftazidime than to other broad-spectrum cephalosporins. *Antimicrob. Agents Chemother.* 32:626-630.
24. Poirel, L., I. Le Thomas, T. Naas, A. Karim, and P. Nordmann. 2000. Biochemical sequence analyses of GES-1, a novel class A extended-spectrum β -lactamase, and the class 1 integron In52 from *Klebsiella pneumoniae*. *Antimicrob. Agents Chemother.* 44:622-632.
25. Poirel, L., T. Naas, M. Guilbert, E. B. Chaibi, R. Labia, and P. Nordmann. 1999. Molecular and biochemical characterization of VEB-1, a novel class A extended-spectrum β -lactamase encoded by an *Escherichia coli* integron gene. *Antimicrob. Agents Chemother.* 43:573-581.
26. Poirel, L., T. Naas, D. Nicolas, L. Collet, S. Bellais, J. D. Cavallo, and P. Nordmann. 2000. Characterization of VIM-2, a carbapenem-hydrolyzing metallo- β -lactamase and its plasmid- and integron-borne gene from a *Pseudomonas aeruginosa* clinical isolate in France. *Antimicrob. Agents Chemother.* 44:891-897.
27. Poirel, L., G. F. Weldhagen, C. De Champs, and P. Nordmann. 2002. A nosocomial outbreak of *Pseudomonas aeruginosa* isolates expressing the extended-spectrum β -lactamase GES-2 in South Africa. *J. Antimicrob. Chemother.* 49:561-565.
28. Poirel, L., G. F. Weldhagen, T. Naas, C. De Champs, M. G. Dove, and P. Nordmann. 2001. GES-2, a class A β -lactamase from *Pseudomonas aeruginosa* with increased hydrolysis of imipenem. *Antimicrob. Agents Chemother.* 45:2598-2603.
29. Rasheed, J. K., C. Jay, B. Metchock, F. Berkowitz, L. Weigel, J. Crellin, C. Steward, B. Hill, A. A. Medeiros, and F. C. Tenover. 1997. Evolution of the extended-spectrum β -lactam resistance (SHV-8) in a strain of *Escherichia coli* during multiple episodes of bacteremia. *Antimicrob. Agents Chemother.* 41:647-653.
30. Riccio, M. L., N. Franceschini, L. Boschi, B. Caravelli, G. Cornaglia, R. Fontana, G. Amicosante, and G. M. Rossolini. 2000. Characterization of the metallo- β -lactamase determinant of *Acinetobacter baumannii* AC-54/97 reveals the existence of *bla*_{IMP} allelic variants carried by gene cassettes of different phylogeny. *Antimicrob. Agents Chemother.* 44:1229-1235.
31. Sabate, M., R. Tarrago, F. Navarro, E. Miro, C. Verges, J. Barbe, and G. Prats. 2000. Cloning and sequence of the gene encoding a novel cefotaxime-hydrolyzing β -lactamase (CTX-M-9) from *Escherichia coli* in Spain. *Antimicrob. Agents Chemother.* 44:1970-1973.
32. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. Molecular cloning: a laboratory manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
33. Senda, K., Y. Arakawa, S. Ichiyama, K. Nakashima, H. Ito, S. Ohsuka, K. Shimokata, N. Kato, and M. Ohta. 1996. PCR detection of metallo- β -lactamase gene (*bla*_{IMP}) in gram-negative rods resistant to broad-spectrum β -lactams. *J. Clin. Microbiol.* 34:2909-2913.
34. Shi, Z. Y., P. Y. Liu, Y. J. Lau, Y. H. Lin, and B. S. Hu. 1996. Epidemiological typing of isolates from an outbreak of infection with multidrug-resistant *Enterobacter cloacae* by repetitive extragenic palindromic unit b1-primed PCR and pulsed-field gel electrophoresis. *J. Clin. Microbiol.* 34:2784-2790.
35. Stauffer, G. V., M. D. Plamann, and L. T. Stauffer. 1981. Construction and expression of hybrid plasmids containing the *Escherichia coli* *glyA* genes. *Gene* 14:63-72.
36. van der Zee, A., N. Steer, E. Thijssen, J. Nelson, A. van't Veen, and A. Buiting. 2003. Use of multienzyme multiplex PCR amplified fragment length polymorphism typing in analysis of outbreaks of multiresistant *Klebsiella pneumoniae* in an intensive care unit. *J. Clin. Microbiol.* 42:798-802.
37. Vourli, S., L. S. Tzouveleki, E. Tzelepi, E. Lebesi, N. J. Legakis, and V. Miriagou. 2003. Characterization of In111, a class 1 integron that carries the extended-spectrum β -lactamase gene *bla*_{IBC-1}. *FEMS Microbiol. Lett.* 225: 149-153.
38. Yagi, T., H. Kurokawa, K. Senda, S. Ichiyama, H. Ito, S. Ohsuka, K. Shibayama, K. Shimokata, N. Kato, M. Ohta, and Y. Arakawa. 1997. Nosocomial spread of cephem-resistant *Escherichia coli* strains carrying multiple Toho-I-like β -lactamase genes. *Antimicrob. Agents Chemother.* 41:2606-2611.
39. Yagi, T., H. Kurokawa, N. Shibata, K. Shibayama, and Y. Arakawa. 2000. A preliminary survey of extended-spectrum β -lactamases (ESBLs) in clinical isolates of *Klebsiella pneumoniae* and *Escherichia coli* in Japan. *FEMS. Microbiol. Lett.* 184:53-56.
40. Yan, J. J., W. C. Ko, Y. C. Jung, C. L. Chuang, and J. J. Wu. 2002. Emergence of *Klebsiella pneumoniae* isolates producing inducible DHA-1 β -lactamase in a university hospital in Taiwan. *J. Clin. Microbiol.* 40:3121-3126.
41. Yan, J. J., W. C. Ko, S. H. Tsai, H. M. Wu, Y. T. Jin, and J. J. Wu. 2000. Dissemination of CTX-M-3 and CMY-2 β -lactamases among clinical isolates of *Escherichia coli* in southern Taiwan. *J. Clin. Microbiol.* 38:4320-4325.

Genetic Environments of the *rmtA* Gene in *Pseudomonas aeruginosa* Clinical Isolates

Kunikazu Yamane, Yohei Doi, Keiko Yokoyama, Tetsuya Yagi, Hiroshi Kurokawa, Naohiro Shibata, Keigo Shibayama, Haru Kato, and Yoshichika Arakawa*

Department of Bacterial Pathogenesis and Infection Control, National Institute of Infectious Diseases, Tokyo, Japan

Received 24 August 2003/Returned for modification 24 December 2003/Accepted 22 February 2004

Nine *Pseudomonas aeruginosa* strains showing very high levels of resistance to various aminoglycosides have been isolated from clinical specimens in seven separate Japanese hospitals in five prefectures since 1997. These strains harbor the newly identified 16S rRNA methylase gene (*rmtA*). When an *rmtA* gene probe was hybridized with genomic DNAs of the nine strains digested with EcoRI, two distinct patterns were observed. The 11.1- and 15.8-kb regions containing the *rmtA* genes of strains AR-2 and AR-11, respectively, were sequenced and compared. In strain AR-2, a transposase gene-like sequence (sequence 1) and a probable tRNA ribosyltransferase gene (*orfA*) were located upstream of *rmtA*, and a Na⁺/H⁺ antiporter gene-like sequence (sequence 2) was identified downstream of *rmtA*. This 6.2-kbp insert (the *rmtA* locus) was flanked by 262-bp *icy* elements. Part of the *orfQ* gene adjacent to an inverted repeat was found outside of the *rmtA* locus. In strain AR-11, the *rmtA* gene and sequence 2 were found, but the 5' end of the *orfA* gene was truncated and replaced with IS6100. An *orfQ*-*orfI* region was present on each side of the *rmtA* gene in strain AR-11. The G+C content of the *rmtA* gene was about 55%, and since the newly identified *rmtA* gene may well be mediated by some mobile genetic elements such as Tn5041, further dissemination of the *rmtA* gene could become an actual clinical problem in the near future.

Pseudomonas aeruginosa is an important opportunistic pathogen that is capable of causing chronic and severe invasive diseases in critically ill and immunocompromised patients. Aminoglycosides are clinically effective agents for treating infections caused by *P. aeruginosa* as well as other gram-negative bacilli. However, multidrug resistance is rapidly emerging in *P. aeruginosa*, whose spectrum of resistance often includes aminoglycosides as well as broad-spectrum β -lactams and fluoroquinolones (15). The most frequently encountered molecular mechanism for aminoglycoside resistance in *P. aeruginosa* is the production of aminoglycoside-modifying enzymes such as plasmid-dependent acetyltransferase (AAC), adenylyltransferase (AAD), and phosphotransferase (APH) (6, 17, 23). Among these, production of AAC(6')-II and AAD(2'')-I is the most common mechanism for resistance to aminoglycosides in *P. aeruginosa* (1), although ribosomal mutations also play some part in aminoglycoside resistance (20). Arbekacin, one of the semisynthetic aminoglycosides belonging to the kanamycin group, is very efficacious for treatment of infections caused by both gram-positive and gram-negative bacteria, and since 1990 it has been approved, for chemotherapy of methicillin-resistant *Staphylococcus aureus* (MRSA) infections only, by the Japanese health insurance system. Unlike the other aminoglycosides, arbekacin is not inactivated by most of the modifying enzymes listed above. Only the bifunctional modifying enzyme composed of aminoglycoside-6'-N-acetyltransferase and 2''-O-phosphotransferase activity [AAC(6')/APH(2'')] is able to inactivate arbekacin. How-

ever, such enzymes have not been found in gram-negative bacilli to date.

We recently reported a *P. aeruginosa* strain that was highly resistant to most aminoglycosides, including arbekacin. This strain harbors a novel aminoglycoside resistance gene named *rmtA*, which encodes a new 16S rRNA methylase (29). Production of 16S rRNA methylase had been reported among aminoglycoside-producing actinomycetes, including *Micromonospora* spp. and *Streptomyces* spp., but this novel aminoglycoside resistance mechanism had not been identified in clinical pathogens before, although a similar putative 16S rRNA methylase, ArMA, was found quite recently in *Klebsiella pneumoniae* in Europe (11). In the present study, we investigated the genetic environments of the *rmtA* genes harbored by two different *P. aeruginosa* strains isolated in separate Japanese hospitals.

(Some of the findings presented in this manuscript have been reported at the 102nd General Meeting of the American Society for Microbiology [abstr. A-28, 2002] by Y. Doi and at its 103rd General Meeting [abstr. A-105, 2003] by K. Yamane.)

MATERIALS AND METHODS

Screening of 16S rRNA methylase producers. In October 2001, a total of 903 nonrepetitive clinical strains of *P. aeruginosa* were collected from 278 medical institutions located in 22 prefectures across Japan. Potential producers of *rmtA* were first screened for a lack of susceptibility to gentamicin, amikacin, and arbekacin (MICs, ≥ 32 μ g/ml). Our bacterial stock of 210 *P. aeruginosa* strains isolated clinically since 1997 was also subjected to a screening test for the *rmtA* gene. Strains that formed colonies on aminoglycoside-containing Mueller-Hinton agar plates were subjected to PCR analyses to check whether or not they harbored the *rmtA* gene. Primers used for amplification of the *rmtA* gene were RMTA-F (5'-CTA GCG TCC ATC CTT TCC TC-3') and RMTA-R (5'-TTT GCT TCC ATG CCC TTG CC-3'), which amplify a 635-bp DNA fragment within the *rmtA* gene. Template DNAs used were prepared by boiling the bacterial suspension at 100°C for 10 min. Cycling parameters consisted of an initial cycle at 94°C for 5 min; 30 cycles of 94°C for 30 s, annealing at 60°C for 30 s, and extension at 74°C for 2 min; and a final 5-min incubation at 74°C. Detection of

* Corresponding author. Mailing address: Department of Bacterial Pathogenesis and Infection Control, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-Murayama, Tokyo 208-0011, Japan. Phone: 81-42-561-0771, ext. 500. Fax: 81-42-561-7173. E-mail: yarakawa@nih.go.jp.

TABLE 1. Bacterial strains and plasmids used in this study

Strain(s) or plasmid	Characteristics	Source or reference
Strains		
<i>P. aeruginosa</i> AR-2, AR-3, AR-11, AR-15, AR-26, AR-101, AR-105, AR-112, and AR-118	Clinical isolates carrying the <i>rmtA</i> gene	This study
<i>E. coli</i> XL1-Blue	<i>supE44 recA1 endA1 gyrA96 thi hsdR17(r_K⁻ m_K⁺) relA1 lac [F⁻ proAB⁺ lacI^q ZΔM15::Tn10(Tet^r)]</i>	Stratagene
Plasmids		
pBCSK+	Cloning vector; chloramphenicol resistant	Stratagene
pBCRMTH2	Recombinant plasmid carrying a 6.8-kb HindIII fragment containing the <i>rmtA</i> gene of <i>P. aeruginosa</i> strain AR-2	This study
pBCRMTE2	Recombinant plasmid carrying a 10.3-kb EcoRI fragment containing the <i>rmtA</i> gene of <i>P. aeruginosa</i> strain AR-2	This study
pBCRMTE11	Recombinant plasmid carrying a 15.8-kb EcoRI fragment containing the <i>rmtA</i> gene of <i>P. aeruginosa</i> strain AR-11	This study

AAC(6')/APH(2'') was carried out as described by Ida et al. (13). Clinical isolates and plasmids used in this study are listed in Table 1.

Antibiotics and susceptibility testing. Antibiotics were obtained from the following sources: amikacin, Bristol Pharmaceuticals K. K., Tokyo, Japan; arbekacin, kanamycin, and streptomycin, Meiji Seika Kaisha Ltd., Tokyo, Japan; chloramphenicol, Sankyo Co., Ltd., Tokyo, Japan; gentamicin and sisomicin, Schering-Plough K. K., Osaka, Japan; hygromycin B, Sigma-Aldrich Japan K. K., Tokyo, Japan; isepamicin, Asahi Kasei Corporation, Tokyo, Japan; neomycin, Nippon Kayaku Co., Ltd., Tokyo, Japan; rifampin, Daiichi Pharmaceutical Co., Ltd., Tokyo, Japan; tobramycin, Shionogi Pharmaceutical Co., Ltd., Osaka, Japan. MICs were determined by the agar dilution method according to the protocol recommended by the National Committee for Clinical Laboratory Standards in document M7-A5 (19).

PFGE analysis. SpeI (New England Biolabs, Beverly, Mass.)-digested genomic DNAs of *P. aeruginosa* isolates were subjected to pulsed-field gel electrophoresis (PFGE) analysis by using a CHEF-DR11 system (Bio-Rad Laboratories, Hercules, Calif.) under conditions described elsewhere (5). The pulses were increased linearly from 4 to 8 s for 10 h, after which the phase was 8 to 50 s for 12 h in this study. Banding patterns of the strains were compared visually; distinct patterns were defined by more than three fragment differences, in accordance with the criteria proposed by Tenover et al. (27).

Southern hybridization analysis of the *rmtA* gene. Total DNAs of all strains were digested with EcoRI (New England Biolabs), electrophoresed through 1.0% agarose gels, transferred to nylon membranes (Bio-Rad Laboratories) by the method of Southern (25), and then hybridized with digoxigenin-labeled *rmtA* gene fragments by use of the PCR DIG detection system (Roche Diagnostics, Tokyo, Japan).

Cloning of the *rmtA* gene. Basic recombinant-DNA techniques were carried out as described by Sambrook et al. (21). EcoRI and HindIII (New England Biolabs) were used for digestion of genomic DNA. The resultant fragments were ligated into the plasmid vector pBCSK+ (Stratagene, La Jolla, Calif.), and electrocompetent *Escherichia coli* XL1-Blue (Stratagene) was transformed with these recombinant plasmids. Transformants were selected on Luria-Bertani agar plates supplemented with 4 μg of arbekacin/ml and 30 μg of chloramphenicol/ml.

DNA sequencing. DNA sequences were determined as described by Sanger et al. (22) with BigDye Terminator Cycle Sequencing Ready Reaction kits and a model 3100 DNA sequence analyzer (Applied Biosystems, Foster City, Calif.). The sequences of the cloned fragments were determined with custom sequencing primers. Nucleotide sequence alignment was performed with GENETYX-MAC (version 10.1.1; Software Development Co., Ltd., Tokyo, Japan). The nucleotide sequence was analyzed by the FASTA service of the DNA Data Bank of Japan (DDBJ) homology search system.

Nucleotide sequence accession numbers. The nucleotide sequence data determined in this study will appear in the DDBJ database under nucleotide accession numbers AB083212 and AB120321.

RESULTS

Bacterial strains. Among 903 strains collected in October 2001, the MICs of arbekacin, gentamicin, and amikacin for 23 strains (2.5%) were greater than 32 μg/ml. Of these, four strains (AR-101, AR-105, AR-112, and AR-118), accounting for 0.4% of all isolates, were found to be positive for *rmtA* by PCR analysis. From our bacterial collection of 210 *P. aeruginosa* strains, 5 strains (AR-2, AR-3, AR-11, AR-15, and AR-26) were PCR positive for *rmtA*. AAC(6')/APH(2'') was not detected in any of these nine strains by PCR analysis. Strains AR-2 and AR-3 were isolated from a hospital, as were strains AR-101 and AR-105. These nine *rmtA*-positive strains have been isolated from seven separate medical institutions in five prefectures in Eastern and Central Japan since 1997.

Susceptibility to antimicrobial agents. MICs of representative aminoglycosides for these nine strains carrying the *rmtA* gene are shown in Table 2. All the strains were highly resistant

TABLE 2. Results of antibiotic susceptibility testing

Aminoglycoside	MIC (μg/ml) for the following <i>P. aeruginosa</i> strain:								
	AR-2	AR-3	AR-11	AR-15	AR-26	AR-101	AR-105	AR-112	AR-118
Kanamycin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Amikacin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Tobramycin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Arbekacin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Gentamicin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Sisomicin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Isepamicin	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024	>1,024
Neomycin	>1,024	>1,024	>1,024	128	>1,024	1,024	512	>1,024	1,024
Hygromycin B	>1,024	1,024	256	128	512	128	128	256	512
Streptomycin	128	128	128	>1,024	512	64	128	128	32




FIG. 1. PFGE fingerprinting of total DNAs from *P. aeruginosa* isolates digested with SpeI. M, PFGE molecular weight marker. The number above each lane indicates the AR strain number shown in Table 1.

to 4,6-disubstituted deoxystreptamines such as kanamycin, amikacin, tobramycin, and arbekacin, which belong to the kanamycin group, as well as to gentamicin, isepamicin, and sisomicin, belonging to the gentamicin group. In contrast, levels of resistance to neomycin, streptomycin, and hygromycin B varied. Strain AR-11 showed a multidrug-resistant profile to ceftazidime, imipenem, and ciprofloxacin as well as to most aminoglycosides.

PFGE profiles. The results of the PFGE analysis are shown in Fig. 1. The SpeI-digested patterns of the total DNAs of nine strains harboring the *rmtA* gene were apparently different from each other. This finding suggests not a clonal expansion of an *rmtA*-carrying strain but plasmid-mediated transmission of the *rmtA* gene among clinical strains with different genetic backgrounds by the help of some movable genetic elements such as a transposon and transferable plasmids.

Southern hybridization. DNA fragments digested with EcoRI showed two hybridization patterns. The *rmtA* probe hybridized with a 10.3-kbp EcoRI fragment for strains AR-2, AR-3, and AR-118 and with a 15.8-kbp fragment for strains AR-11, AR-15, AR-26, AR-101, AR-105, and AR-112 (Fig. 2).

Genetic environments harboring *rmtA* genes. A 6.8-kbp HindIII fragment and a 10.3-kbp EcoRI fragment containing the *rmtA* gene of AR-2 were cloned into the plasmid vector pBCSK+. The 6.8- and 10.3-kbp fragments were inserted into pBCRMTH2 and pBCRMTE2, respectively. The schematic structure of the 11.1-kbp sequenced region cloned from strain AR-2 is shown in Fig. 3. The *rmtA* gene was located within a 6.2-kbp genetic locus (the *rmtA* locus) flanked by a 262-bp sequence named the $\kappa\gamma$ element that was previously found in Tn5041 and predicted to be a relic of mobile genetic elements (Fig. 3). The elements of the 6.2-kbp *rmtA* locus, comprising *rmtA*, *orfA*, and two additional specific sequences, were located in the following order: transposase gene-like sequence (se-

quence 1), probable tRNA ribosyltransferase gene (*orfA*), *rmtA*, and Na^+/H^+ antiporter gene-like sequence (sequence 2) (Fig. 3). The 5' end of the HindIII fragment flanked *merR* of the *mer* operon found in Tn5041. However, the 3' end of the EcoRI fragment was located within a 17-bp sequence which was completely identical to a part of the terminal inverted repeat of Tn1721. This 17-bp sequence was within *orfQ*, located upstream of *orfI* in Tn5041. The G+C content of the 6.2-kbp *rmtA* locus was about 55%. The 15.8-kbp EcoRI fragment of AR-11 containing the *rmtA* gene was also cloned into the plasmid vector pBCSK+, and the resultant recombinant plasmid was designated pBCRMTE11. In the 15.8-kbp EcoRI fragment, a 5'-truncated *orfA* (*orfA'*), *rmtA*, and sequence 2 were found between IS6100 and a $\kappa\gamma$ element, and the sequence was completely identical to that of the corresponding region of the 6.2-kbp *rmtA* locus cloned from strain AR-2. The *orfQ* and *orfI* sequences of Tn5041 were present both upstream of IS6100 and downstream of a $\kappa\gamma$ element in the 15.8-kbp EcoRI fragment cloned from strain AR-11. In the sequenced areas, the fragments harboring the *rmtA* gene appeared to be inserted between the $\kappa\gamma$ sequences found in Tn5041 (Fig. 3).

DISCUSSION

Aminoglycoside-producing actinomycetes such as *Micromonospora* spp. and *Streptomyces* spp. protect their 30S ribosome through methylation of its 16S rRNA at the aminoglycoside-binding A site (10, 30). For example, Kgm, which was isolated from *Micromonospora purpurea* (28), methylates G1405, and Kam, which was isolated from *Streptomyces tenjimariensis* (24), methylates A1408 (2). The 16S rRNA methylases had been thought to exist among aminoglycoside-producing environmental actinomycetes such as *Micromonospora* spp. or *Streptomyces* spp (7). However, we recently reported a novel 16S rRNA methylase, RmtA, that was identified in a *P. aeruginosa* clinical strain, AR-2 (29). This strain demonstrated an extraor-




FIG. 2. Southern hybridization patterns of EcoRI-digested genomic DNAs. The number above each lane represents the AR strain number shown in Table 1. The nine strains tested appeared to be divided into two groups by the sizes of EcoRI-digested fragments (10.3 and 15.8 kbp, respectively).




FIG. 3. Comparison of the genetic organizations of AR-2 and AR-11. Double-headed striped arrows indicate the position of the *rmtA* locus and that of the region common to both sequenced areas. Inserts of pBCRMTH2, pBCRMTE2, and pBCRMTE11 are indicated by horizontal lines. Rectangles filled with wavy lines, sequences similar to part of Tn5041. Solid arrowheads in the 15.8-kbp EcoRI fragment, terminal inverted repeats. *mer*, the mercury resistance operon, includes *merR*. Sequence 1, transposase gene-like sequence; sequence 2, Na⁺/H⁺ antiporter-like sequence; *orfA*, probable tRNA ribosyltransferase gene; *orfQ'*, part of *orfQ*; *orfA'*, part of *orfA*; IR, probable inverted repeat. Restriction sites: H, HindIII; E, EcoRI. Sequences 1 and 2 encode no complete proteins due to several frameshifts and deletions.

dinarily high level of aminoglycoside resistance to various 4,6-disubstituted deoxystreptamines, including semisynthetic arbekacin, as well as to gentamicin and kanamycin. In the present study, we investigated the genetic environments mediating the *rmtA* genes found in two different strains of *P. aeruginosa*. The G+C content of the *rmtA* gene was 55%, and those of 16S rRNA methylase genes found in aminoglycoside-producing actinomycetes were 64 to 72%. These observations suggested that the *rmtA* gene might have been acquired by *P. aeruginosa* from some environmental bacteria such as aminoglycoside-producing actinomycetes, although the *armA* gene, with a 30% G+C content, was speculated to have originated from unknown bacteria other than actinomycetes. At any rate, lateral gene transfer across bacterial genera would become much more important for acquisition of new antibiotic resistance profiles hereafter.

Although the PFGE patterns of the nine RmtA-producing strains in this study were highly divergent, Southern hybridization showed only two hybridization patterns when genomic DNAs were digested with EcoRI. This finding indicated that the *rmtA* gene might be mediated by some mobile genetic elements sharing similar genetic environments and spreading among genetically unrelated strains in geographically separate hospitals. This speculation would be supported by the finding that even strains AR-2 and AR-3, isolated at the same hospital, showed different PFGE patterns. Strains AR-101 and AR-105 also demonstrated quite different PFGE profiles despite being isolated at the same hospital. Furthermore, the arbakacin resistance profile of AR-2 was transferable to another *P. aeruginosa* strain by conjugation (29). This suggested that *rmtA* was mediated by some transferable plasmids in strain AR-2, but we failed to visualize the plasmid either by the method of Kado and Liu (14) or by cesium chloride-ethidium bromide density

gradient ultracentrifugation (21). This is possibly due to the instability or the very low copy number of the plasmid which mediates the *rmtA* gene.

Tn5041 was previously identified in a strain of a *Pseudomonas* species as a mercury resistance transposon (3, 16). Tn5041 carries a 4-kbp insert of unknown origin between *orfQ* and the *mer* operon, and several nonfunctional pseudogenes and possible mobile elements such as the *κγ* element locate in this region. The 262-bp *κγ* element, containing 38 bp of imperfect inverted repeats starting with the sequence GGGG and terminating internally with the sequence TAAG, falls into the inverted repeats of Tn3 family (4). Transposons belonging to the Tn3 family usually contain transposase and resolvase genes and some additional genes encoding resistance to antimicrobial agents or heavy metals such as mercury between the terminal inverted repeats. The 6.2-kbp *rmtA* locus found in this study was flanked by an insertion element-like *κγ* element. Moreover, the *rmtA* locus had a transposase gene-like sequence (sequence 1) whose 5' part showed 80.2% identity with part of the transposase gene derived from *Pseudomonas putida* (accession number AF109307); the 3' part of sequence 1 had 67.2% identity with part of the transposase gene derived from *Pseudomonas pseudoalcaligenes* (accession number AF028594), but this sequence had no apparent initiation and stop codons. Thus, the 6.2-kb *rmtA* locus itself is unlikely to be an active transposon, although the nucleotide sequences outside of the two *κγ* elements were completely identical to the corresponding regions of Tn5041. The Na⁺/H⁺ antiporter gene-like sequences (sequence 2) found in strains AR-11 and AR-2 were completely identical, although they seemed nonfunctional. Multicopy expression of the intact transposase-like gene and the Na⁺/H⁺ antiporter-like gene might disturb systematic bacterial cell growth, so these genes might have been

inactivated during replication and translocation of the *rmtA* locus.

To examine whether strains other than AR-2 and AR-11 also carry part of the sequence found in Tn5041, Southern hybridization analysis was performed using a Tn5041-specific DNA probe containing a sequence between the right-hand *ky* element and the *orfQ* gene, which is conserved in both strains AR-2 and AR-11. The DNA probes and the *rmtA* gene probe hybridized to the same fragments in all nine strains (data not shown). This finding strongly suggests the probable implication of some mobile genetic elements such as Tn5041 in the dissemination of the *rmtA* gene among strains of *P. aeruginosa*.

The 5' end of the *rmtA* locus was replaced by IS6100 in strain AR-11. IS6100 was originally discovered in *Mycobacterium fortuitum* (accession number X53635) (18) and was subsequently found in several gram-negative and -positive bacteria (9, 26). It has been reported that transposition of IS6100 stimulates genetic rearrangement (12). Thus, it may be possible to speculate that the region containing *orfQ* and *orfI* found upstream of IS6100 might be duplicated during IS6100-mediated recombination in strain AR-11. The outside sequences of both inverted repeats had no DNA homology to the genomic DNA of *P. aeruginosa* PAO-1. This finding suggests that the 15.8-kb EcoRI fragment of strain AR-11 might be carried by a much longer mobile genetic element, since the arbekacin-resistant profile of AR-11 was not transferred to another *P. aeruginosa* strain by conjugation, and no apparent plasmid was detected in this strain by the method of Kado and Liu (14). Additionally, *rmtA* gene probes hybridized to the position of chromosomal DNA (data not shown). These findings strongly suggested that the *rmtA* gene and its adjacent regions might be integrated into the chromosomal DNA in strain AR-11.

P. aeruginosa strains harboring the *rmtA* gene have already been found in several separate clinical settings in Japan, and a gene encoding the same kind of 16S rRNA methylase, called *armA*, has also been identified in members of the family *Enterobacteriaceae*, such as *Citrobacter freundii* (accession number NC004464) and *K. pneumoniae* (11) (accession number AY220558), in Europe. ArmA shares 29% identity with RmtA at the amino acid sequence level. Moreover, a new plasmid-mediated 16S rRNA methylase, RmtB, that shares 82% identity with RmtA at the amino acid sequence level, has also been identified in *Serratia marcescens* in Japan (8) (accession number AB103506). From our preliminary study on a bacterial stock, the presence of these genes was also suggested in several strains of *K. pneumoniae*, *E. coli*, and *Acinetobacter* species isolated in Japan. Thus, further dissemination of these genetic determinants to various pathogenic gram-negative bacilli could become a serious concern in the near future.

In Japanese clinical settings, various aminoglycosides have been used in the treatment of bacterial infections, since these agents still have very high efficacies against both gram-positive and gram-negative bacteria. Arbekacin is a semisynthetic aminoglycoside belonging to the kanamycin-group. It has been approved, for MRSA infection only, since 1990, and it is still very efficacious for MRSA infection. Under such clinical circumstances, arbekacin has been preferentially used in many clinical settings, although arbekacin-resistant strains which produce the bifunctional enzyme AAC(6')/APH(2'') have emerged in MRSA. No such bifunctional enzymes, however, have been

found in gram-negative bacilli to date. Thus, acquisition of 16S rRNA methylase would give gram-negative bacteria a great advantage in coping with clinical environments where huge amounts of semisynthetic aminoglycosides, including arbekacin, are consumed. Hence, one should recall again that bacteria can survive and proliferate in clinical environments, given their natural hereditary capacity to overcome the hazards of any environment.

ACKNOWLEDGMENTS

This work was supported by grants H12-Shinkou-19, H12-Shinkou-20, H15-Shinkou-9, and H15-Shinkou-10 from the Ministry of Health, Labor, and Welfare of Japan. The travel costs for presenting part of the data reported in this paper at the 103rd General Meeting of the American Society for Microbiology was supported by a grant from The Japan Antibiotics Research Association, sponsored by the Pfizer Infectious Diseases Research Fund 2003.

REFERENCES

1. Aminoglycoside Resistance Study Groups. 1994. Resistance to aminoglycosides in *Pseudomonas*. Trends Microbiol. 2:347-353.
2. Beancleer, A. A. D., and E. Cundliffe. 1987. Sites of action of two ribosomal RNA methylases responsible for resistance to aminoglycosides. J. Mol. Biol. 193:661-671.
3. Bogdanova, E. S., S. Z. Mindlin, E. S. Kalyaeva, and V. G. Nikiforov. 1988. The diversity of mercury reductases among mercury-resistant bacteria. FEBS Lett. 234:280-282.
4. Craig, N. L., R. Craigie, M. Gellert, and A. M. Lambowitz. 2002. Mobile DNA II. ASM Press, Washington, D.C.
5. D'Agata, E., L. Venkataraman, P. DeGirolami, and M. Samore. 1997. Molecular epidemiology of acquisition of ceftazidime-resistant gram-negative bacilli in a nonoutbreak setting. J. Clin. Microbiol. 35:2602-2605.
6. Davies, J., and G. D. Wright. 1997. Bacterial resistance to aminoglycoside antibiotics. Trends Microbiol. 5:234-240.
7. Davies, J. D. 1983. Resistance to aminoglycosides: mechanisms and frequency. Rev. Infect. Dis. 5:S261-S267.
8. Doi, Y., K. Yokoyama, K. Yamane, J. Wachino, N. Shibata, T. Yagi, K. Shibayama, H. Kato, and Y. Arakawa. 2004. Plasmid-mediated 16S rRNA methylase in *Serratia marcescens* conferring high-level resistance to aminoglycosides. Antimicrob. Agents Chemother. 48:491-496.
9. Dubois, Y., C. Arpin, P. Nonry, and C. Quentin. 2002. Clinical strain of *Pseudomonas aeruginosa* carrying a *bla*_{TEM-21} gene located on a chromosomal interrupted Tn4 type transposon. Antimicrob. Agents Chemother. 46:3624-3626.
10. Fouemy, D., M. I. Recht, S. C. Blanchard, and J. D. Puglisi. 1996. Structure of the A site of *Escherichia coli* 16S ribosomal RNA complexed with an aminoglycoside antibiotic. Science 274:1347-1371.
11. Galimand, M., P. Courvalin, and T. Lambert. 2003. Plasmid-mediated high-level resistance to aminoglycosides in *Enterobacteriaceae* due to 16S rRNA methylation. Antimicrob. Agents Chemother. 47:2565-2571.
12. Gunos, G., B. Smith, and P. Dyson. 1999. Genetic instability associated with insertion of IS6100 into one end of the *Streptomyces lividans* chromosome. Microbiology 145:2203-2208.
13. Ida, T., R. Okamoto, C. Shimanchi, T. Okubo, A. Kuga, and M. Inoue. 2001. Identification of aminoglycoside-modifying enzymes by susceptibility testing: epidemiology of methicillin-resistant *Staphylococcus aureus* in Japan. J. Clin. Microbiol. 39:3115-3121.
14. Kado, C. I., and S. T. Liu. 1981. Rapid procedure for detection and isolation of large and small plasmids. J. Bacteriol. 145:1365-1373.
15. Karlowitsky, J. A., D. C. Draghi, M. E. Jones, C. Thornsberry, I. R. Friedland, and D. F. Sahn. 2003. Surveillance for antimicrobial susceptibility among clinical isolates of *Pseudomonas aeruginosa* and *Acinetobacter baumannii* from hospitalized patients in the United States, 1998 to 2001. Antimicrob. Agents Chemother. 47:1681-1688.
16. Kholdii, G. Y., O. V. Yurieva, Z. M. Gorlenko, S. Z. Mindlin, I. A. Bass, O. L. Lomovskaya, A. V. Kopteva, and V. G. Nikiforov. 1997. Tn5041: a chimeric mercury resistance transposon closely related to the toluene degradative transposon Tn4651. Microbiology 143:2549-2556.
17. Kotra, L. P., J. Haddad, and S. Mobashery. 2000. Aminoglycosides: perspectives on mechanisms of action and resistance and strategies to counter resistance. Antimicrob. Agents Chemother. 44:3249-3256.
18. Martin, C., J. Timm, J. Raugier, R. Gomez-Lus, J. Davies, and B. Gicquel. 1990. Transposition of an antibiotic resistance element in mycobacteria. Nature 345:739-743.
19. National Committee for Clinical Laboratory Standards. 1988. Methods for dilution antimicrobial susceptibility tests for bacteria that grow aerobically, 4th ed. Document M7-A5. National Committee for Clinical Laboratory Standards, Wayne, Pa.

20. Recht, M. I., and J. D. Puglisi. 2001. Aminoglycoside resistance with homogeneous and heterogeneous populations of antibiotic-resistant ribosomes. *Antimicrob. Agents Chemother.* 45:2414–2419.
21. Sambrook, J., E. F. Fritsch, and T. Maniatis. 1989. *Molecular cloning: a laboratory manual*, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
22. Sanger, F., S. Nicklen, and A. R. Coulson. 1977. DNA sequencing with chain-terminating inhibitors. *Proc. Natl. Acad. Sci. USA* 74:5463–5467.
23. Shaw, K. J., P. N. Rather, R. S. Hare, and G. H. Miller. 1993. Molecular genetics of aminoglycoside resistance genes and familial relationships of the aminoglycoside-modifying enzymes. *Microbiol. Rev.* 57:138–163.
24. Skeggs, P. A., J. Thompson, and E. Cundliffe. 1985. Methylation of 16S ribosomal RNA and resistance to the aminoglycoside antibiotics in clones of *Streptomyces lividans* carrying DNA from *Streptomyces tenjimariensis*. *Mol. Gen. Genet.* 200:415–421.
25. Southern, E. M. 1975. Detection of specific sequences among DNA fragments separated by gel electrophoresis. *J. Mol. Biol.* 98:503–517.
26. Tauch, A., S. Gotker, A. Puhler, J. Kalinowski, and G. Thierbach. 2002. The 27.8-kb R-plasmid pTET3 from *Corynebacterium glutamicum* encodes the aminoglycoside adenylyltransferase gene cassette *aadA9* and the regulated tetracycline efflux system Tet 33 flanked by active copies of the widespread insertion sequence IS6100. *Plasmid* 48:117–129.
27. Tenover, F. C., R. D. Arbeit, R. V. Goering, P. A. Mickelsen, B. E. Murray, D. H. Persing, and B. Swaminathan. 1995. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J. Clin. Microbiol.* 33:2233–2239.
28. Thompson, J., P. A. Skeggs, and E. Cundliffe. 1985. Methylation of 16S ribosomal RNA and resistance to the aminoglycoside antibiotics gentamicin and kanamycin determined by DNA from the gentamicin-producer, *Micromonospora purpurea*. *Mol. Gen. Genet.* 201:168–173.
29. Yokoyama, K., Y. Doi, K. Yamane, H. Kurokawa, N. Shibata, K. Shibayama, T. Yagi, H. Kato, and Y. Arakawa. 2003. Acquisition of 16S rRNA methylase gene in *Pseudomonas aeruginosa*. *Lancet* 362:1888–1893.
30. Yoshizawa, S., D. Fourmy, and J. D. Puglisi. 1998. Structural origins of gentamicin antibiotic action. *EMBO J.* 17:6437–6448.

Spread of Novel Aminoglycoside Resistance Gene *aac(6′)-Iad* among *Acinetobacter* Clinical Isolates in Japan

Yohei Doi, Jun-ichi Wachino, Kunikazu Yamane, Naohiro Shibata, Tetsuya Yagi,
Keigo Shibayama, Haru Kato, and Yoshichika Arakawa*

Department of Bacterial Pathogenesis and Infection Control, National Institute
of Infectious Diseases, Tokyo, Japan

Received 16 November 2003/Returned for modification 2 February 2004/Accepted 26 February 2004

A novel aminoglycoside resistance gene, *aac(6′)-Iad*, encoding aminoglycoside 6′-N-acetyltransferase, was identified in *Acinetobacter* genospecies 3 strain A-51. The gene encoded a 144-amino-acid protein, which shared modest identity (up to 36.7%) with some of the aminoglycoside 6′-N-acetyltransferases. The results of high-pressure liquid chromatography assays confirmed that the protein is a functional aminoglycoside 6′-N-acetyltransferase. The enzyme conferred resistance to amikacin, tobramycin, sisomicin, and isepamicin but not to gentamicin. The prevalence of this gene among *Acinetobacter* clinical isolates in Japan was then investigated. Of 264 *Acinetobacter* sp. strains isolated from geographically diverse areas in Japan in 2002, 16 were not susceptible to amikacin, and *aac(6′)-Iad* was detected in 7. Five of the producers of aminoglycoside 6′-N-acetyltransferase type Iad were identified as *Acinetobacter baumannii*, and two were identified as *Acinetobacter* genospecies 3. These results suggest that *aac(6′)-Iad* plays a substantial role in amikacin resistance among *Acinetobacter* spp. in Japan.

Acinetobacter spp., especially *Acinetobacter baumannii*, are emerging pathogens responsible for causing a variety of nosocomial infections, including pneumonia, urinary tract infections, and septicemia (1). Outbreaks have been increasingly reported in the past 2 decades, particularly from intensive care units, where patients undergo invasive procedures and receive broad-spectrum antimicrobial agents, resulting in higher mortality rates (5, 27). Furthermore, because *Acinetobacter* spp. have an ability to readily accept foreign DNA, including genetic determinants for antimicrobial resistance, so as to adapt to and survive in environments that are hazardous to bacterial growth (6, 17), they have a propensity for developing resistance to multiple classes of useful antimicrobial agents, including broad-spectrum cephalosporins, fluoroquinolones, and aminoglycosides (1).

Aminoglycosides are widely used to treat infections caused by gram-negative bacilli, including *Acinetobacter* spp. (1). However, resistance rates to classic aminoglycosides such as gentamicin and kanamycin are now high among *Acinetobacter* spp. in many geographic regions (15). The mechanisms of *Acinetobacter* sp. resistance to newer semisynthetic aminoglycosides such as amikacin, tobramycin, sisomicin, and isepamicin are diverse and commonly involve production of aminoglycoside-modifying enzymes such as aminoglycoside acetyltransferases (AAC), aminoglycoside nucleotidyltransferases (ANT, or AAD), and/or aminoglycoside phosphotransferases (APH). Production of AAC(3)-I, APH(3′)-VI, and ANT(3′′)-I was reported to be predominant by worldwide surveys on *Acinetobacter* spp., but there were considerable regional differences in their genotypes (14, 15, 21). In Japan, although the prevalence of ami-

kacin resistance was estimated to be high, especially among non-carbapenem-susceptible *Acinetobacter* strains (25), the overall prevalence of aminoglycoside resistance and the mechanisms of resistance among *Acinetobacter* spp. have not been elucidated to date.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. In March 2002, 264 nonrepetitive strains identified as belonging to *Acinetobacter* spp. were collected from 88 hospitals located in geographically diverse areas in Japan. Among these, 16 strains (6.1%) that were not susceptible to amikacin (MICs, >16 µg/ml) by preliminary susceptibility testing were selected for further study. Species identification was carried out with API 20NE (bioMérieux Japan, Ltd., Tokyo, Japan) complemented by a carbon source utilization test and growth at 41 and 44°C (2). *Escherichia coli* XLI-Blue was used as the host for cloning experiments with vector pBCSK+ (Stratagene, La Jolla, Calif.). *E. coli* BL21(DE3)pLysS was used with vector pET29a(+) (Novagen, Madison, Wis.) for expression of *aac(6′)-Iad*. The strains were grown in Luria-Bertani (LB) broth or medium (Becton Dickinson Diagnostic Systems, Sparks, Md.) supplemented with appropriate antimicrobial agents, unless described otherwise.

Antimicrobial agents and susceptibility testing. Antimicrobial agents were obtained from the following sources: amikacin, Bristol Pharmaceuticals K. K., Tokyo, Japan; arbekacin, kanamycin, ribostamycin, and streptomycin, Meiji Seika Kaisha Ltd., Tokyo, Japan; chloramphenicol, Sankyo Co., Ltd., Tokyo, Japan; gentamicin and sisomicin, Schering-Plough K. K., Osaka, Japan; isepamicin, Asahi Kasei Corporation, Tokyo, Japan; neomycin, Nippon Kayaku Co., Ltd., Tokyo, Japan; rifampin, Daiichi Pharmaceutical Co., Ltd., Tokyo, Japan; tobramycin, Shionogi Pharmaceutical Co., Osaka, Japan.

MICs were determined by the agar dilution method with Mueller-Hinton agar (Becton Dickinson Diagnostic Systems) according to the protocol recommended by the National Committee for Clinical Laboratory Standards (16).

Transfer of aminoglycoside resistance genes. Conjugation experiments were conducted by using rifampin-resistant *E. coli* CSH2 and *Acinetobacter calcoaceticus* DU1, a rifampin-resistant derivative of *A. calcoaceticus* ATCC 33305, as the recipients by the broth mating method (7). Transconjugants were selected on LB agar supplemented with rifampin (50 µg/ml) and kanamycin (10 µg/ml).

Cloning and sequencing of the aminoglycoside resistance gene. The genomic DNA of *Acinetobacter* genospecies 3 strain A-51 was partially digested with *Sau*3AI, and the resultant fragments were ligated to the *Bam*HI-cleaved cloning site of plasmid vector pBCSK+ (Stratagene). Electrocompetent *E. coli* XLI-Blue was transformed with these recombinant plasmids carrying total-DNA restriction fragments of various sizes prepared from the aminoglycoside-resistant

* Corresponding author. Mailing address: Department of Bacterial Pathogenesis and Infection Control, National Institute of Infectious Diseases, 4-7-1, Gakuen, Musashi-Murayama, Tokyo 208-0011, Japan. Phone: 81-42-561-0771, ext. 500. Fax: 81-42-561-7173. E-mail: yarakawa@nih.go.jp.

TABLE 1. Susceptibilities of *Acinetobacter* spp. and *E. coli* strains with *aac(6')*-*Iad* to various aminoglycosides

Strain	Hospital	Specimen	MIC ($\mu\text{g/ml}$) of the following aminoglycoside ^a :								
			KAN	TOB	AMK	ABK	GEN	SISO	ISP	NEO	STR
<i>Acinetobacter</i> genomic species 3, strain A-51	A	Sputum	>1,024	>1,024	1,024	1,024	>1,024	>1,024	>1,024	64	>1,024
<i>A. baumannii</i> A-67	B	Urine	>1,024	64	128	32	8	1,024	256	8	256
<i>A. baumannii</i> A-74	B	Pus	>1,024	512	128	32	8	512	256	8	256
<i>A. baumannii</i> A-87	C	Sputum	512	128	32	16	4	256	256	4	256
<i>A. baumannii</i> A-88	C	Sputum	256	64	128	32	4	128	128	8	256
<i>Acinetobacter</i> genomic species 3, strain A-178	D	Sputum	128	16	32	8	1	64	64	1	64
<i>A. baumannii</i> A-260	E	Sputum	512	256	128	16	4	256	128	8	128
<i>E. coli</i> XL1-Blue(pA51S3)			256	64	128	16	1	64	64	4	4
<i>E. coli</i> XL1-Blue(pA51SG5)			512	32	1	0.13	32	32	0.13	0.25	2
<i>E. coli</i> XL1-Blue(pBCSK+)			0.5	0.25	0.5	0.13	0.13	0.13	0.25	0.25	1

^a KAN, kanamycin; TOB, tobramycin; AMK, amikacin; ABK, arbekacin; GEN, gentamicin; SISO, sisomicin; ISP, isepamicin; NEO, neomycin; STR, streptomycin.

strain. Transformants were selected by their resistance to chloramphenicol (30 $\mu\text{g/ml}$) and kanamycin (25 $\mu\text{g/ml}$). The enzymes used for gene manipulation were purchased from New England Biolabs, Inc. (Beverly, Mass.), or TAKARA Bio, Inc. (Ohtsu, Japan). The DNA sequences were determined on both strands by using BigDye Terminator Cycle Sequencing Ready Reaction kits and an ABI 3100 DNA sequence analyzer (Applied Biosystems, Foster City, Calif.). Alignments of nucleotide and amino acid sequences were performed with the GENE-TYX-MAC computer program (version 10.1.1; Software Development Co., Ltd., Tokyo, Japan).

Purification of the acetyltransferase. For use in N-terminal sequencing and high-pressure liquid chromatography (HPLC) assays, AAC(6')-Iad was purified by using a histidine tag purification system. The entire coding region of *aac(6')*-*Iad* and its upstream sequence were amplified by PCR with primers AAC-F (5'-GCT CTA GAA GAC TGA CTT CGC ATT G-3') and AAC-R (5'-CCC AAG CTT GAG CTG CTT TGT AAA AC-3'). The product was double digested with XbaI and HindIII and then ligated with pET29a(+) (Novagen) digested with the same enzymes. Electrocompetent *E. coli* XL1-Blue was transformed with the recombinant plasmids, and transformants were selected on LB agar containing kanamycin (25 $\mu\text{g/ml}$). Several of the colonies obtained were found to harbor plasmids with inserts encoding AAC(6')-Iad tagged with six histidine residues at the C-terminal end. *E. coli* BL21(DE3)pLysS (Novagen) was transformed with one such plasmid, pA51H7. The transformants were cultured in 1 liter of LB broth supplemented with kanamycin (25 $\mu\text{g/ml}$) to an A_{620} of approximately 0.7. The pellet was washed once with 50 mM phosphate buffer (pH 7.0) and suspended in 20 mM phosphate buffer (pH 7.4) containing 10 mM of imidazole. The suspension was passed twice through a French pressure cell (Ohtake Works Co., Ltd., Tokyo, Japan) at 120 MPa and then centrifuged at 30,000 \times g for 30 min. Histidine-tagged AAC(6')-Iad contained in the supernatant was purified by using HiTrap Chelating HP, included in the HiTrap kit (Amersham Biosciences, K. K., Tokyo, Japan), according to the manufacturer's instructions. It was eluted at an imidazole concentration of 300 mM and was estimated to be more than 95% pure by sodium dodecyl sulfate-polyacrylamide gel electrophoresis. Finally, the enzyme was dialyzed twice at 4°C against 500 volumes of 50 mM phosphate buffer (pH 7.4) and was stored in aliquots at -80°C until use. N-terminal sequencing of the purified enzyme was performed by Shimadzu Corporation (Kyoto, Japan).

Preparation of crude extracts. As positive controls for acetylation reactions and HPLC assays, the following strains were used: AAC(2')-producing *Streptomyces lividans* TK21/pANT12-1', AAC(3)-producing *S. lividans* TK21/pANT3-1, and AAC(6')-producing *S. lividans* TK21/pANTS-2 (8). They were cultured in 100 ml of TS medium containing 10 μg of ribostamycin/ml and 10 μg of thio-strepton/ml (Sigma-Aldrich Japan K. K., Tokyo, Japan) for 48 h. The cells were then harvested, washed once with 50 mM phosphate buffer (pH 7.0), and suspended in the same buffer. The suspension was passed twice through a French pressure cell (Ohtake Works) at 120 MPa and then centrifuged at 30,000 \times g for 30 min. The supernatant was used as the crude enzyme.

Acetylation. Reaction mixtures for acetylation contained 25 μmol of Tris-hydrochloride buffer (pH 7.6), 7.5 μmol of MgCl_2 , 200 nmol of acetyl coenzyme A (acetyl-CoA), and 50 μmol of either tobramycin or neomycin in a final volume of 500 μl . Acetylation was initiated by adding 50 μl of the enzyme and was carried out at 37°C for 30 min. *ortho*-Phthalaldehyde derivatization was then performed by adding equal volumes of 2-propanol and the derivatization reagent to the reaction mixture and heating at 60°C for 10 min. The derivatization

reagent consisted of 80 mM *o*-phthalaldehyde, 1 M boric acid, and 250 mM thioglycolic acid with the pH adjusted to 10.4 with 40% potassium hydroxide.

HPLC assay. HPLC was performed to identify the site of acetylation of substrate aminoglycosides according to the methods described by Lovering et al. (12). The system consisted of a Separations module 2690 (Waters Corporation, Milford, Mass.), a Dual λ absorbance detector set at 330 nm (Waters), and a Chemobond 5-ODS-H column (4.6 by 100 mm; Chemco Scientific Co., Ltd., Osaka, Japan). The mobile phase consisted of methanol-water-acetic acid (61.25:33.75:5) plus 5 g of 1-heptanesulfonic acid sodium salt per liter at a flow rate of 2 ml/min.

PCR amplification. PCR analysis was performed for the 16 non-amikacin-susceptible *Acinetobacter* strains with primers ABA-F (5'-TTT GGC TAT GAT CCT ATG-3') and ABA-R (5'-CAT GTC GAA CAA GTA CGC-3') to amplify an internal fragment of the *aac(6')*-*Iad* gene. The conditions used have been described previously (7). When amplicons were obtained, they were directly sequenced with the same primers.

Nucleotide sequence accession number. The nucleotide sequence of *aac(6')*-*Iad* will appear in GenBank under accession no. AB119105.

RESULTS

Prevalence and resistance profile of *Acinetobacter* strains with *aac(6')*-*Iad*. Of the 16 non-amikacin-susceptible *Acinetobacter* strains included in this study, 7 were PCR positive for *aac(6')*-*Iad*. Five were phenotypically identified as *A. baumannii*, whereas the remaining two were identified as *Acinetobacter* genospecies 3. When the amplicons were sequenced, all were identical to *aac(6')*-*Iad*. The MICs of aminoglycosides for *Acinetobacter* strains possessing *aac(6')*-*Iad* are shown in Table 1. All the strains studied were resistant to kanamycin, amikacin, tobramycin, sisomicin, isepamicin, and streptomycin. In addition, strain A-51 was resistant to all of the aminoglycosides tested, including arbekacin, gentamicin, and neomycin.

Molecular characterization of aminoglycoside resistance genes. Several transformants were obtained by selection with kanamycin and chloramphenicol. When these colonies were inoculated onto plates containing either amikacin (5 $\mu\text{g/ml}$) or gentamicin (5 $\mu\text{g/ml}$), they grew only on one or the other plate. The colonies on the plates containing amikacin or gentamicin were found to harbor recombinant plasmids of various sizes with inserts originating from the genomic DNA of strain A-51. Among these, the smallest plasmids (pA51S3 from an amikacin-resistant colony and pA51SG5 from a gentamicin-resistant colony) were selected out for further study. The MICs of aminoglycosides for *E. coli* XL1-Blue(pA51S3) and XL1-Blue(pA51SG5) are listed in Table 1. pA51S3 conferred resistance to kanamycin, amikacin, tobramycin, sisomicin, and isepami-

AAC(6')-Iad	MIRKATVQDPPLLARLAMNVWKESLKLVAEFEQMTKNSD---AVAFILFIED	51
AAC(6')-Ic	MIVICDHNDLDAWLALRTALWPSGSPEDHRAEMREILASPH---HTAFMARGLD	51
AAC(6')-Id	MIEACHSVECPGWLQLRFLWPQDSADHLEMAIFVAEPNR--FAQFIAYDEA	52
AAC(6')-If	MDEASLSMWWGLRSQLWPDHSDHILDSQHILSCDPK--YVSFLAINNQ	48
AAC(6')-Ig	MNIKPASEASLKDWLELRNKLWS-DSEASHLQEMHQLLAEKY---ALQLLAYS-	50
AAC(6')-Ih	MNIMPISSEQLSDWLALRCLLWP-DHEDVHLQEMRQLITQAH---RLQLLAYTDT	51
AAC(6')-Ij	MNIMPVSESLMADWLGRLKLLWP-DHDEAHLQEMRQLLQQTQ---SLQLLAYS-	51
AAC(6')-Ik	MNIKPASEASLKDWLKLRIKLWN-DLEESHQEMHQLLAEKH---ALQLLVYSD-	50
AAC(6')-Il	MDSSPLVRPVETDSASWLSMRCELWPDGTCQEHQSEIAEFLSGKVARPAVLIAPD	59
AAC(6')-Ir	MKIMPVSEFLADWLQLRILLWP-DHEDAHLLEMRLLEQPH---TLQLLSYNDQ	51
AAC(6')-Is	MNIMPISSEQLSDWLALRCLLWP-DHEDAHLLEMRLVLRKQTD---TLQLLVYSET	51
AAC(6')-It	MNIMPISSEQLSDWLALRCLLWP-DHEDAHLLEMRLVLRKQTD---TLQLLVYSET	51
AAC(6')-Iu	MNIMPISSEQLSDWLALRCLLWP-DHEEAHLQEMRQLLQKTD---TLQLLAYSET	51
AAC(6')-Iv	MKIMPISSEQLSDWLALRCLLWP-DHEEAHLQEMRQLLQKTD---TLQLLAYSET	51
AAC(6')-Iw	MKIMPISSEQLSDWLALRCLLWP-DHEEAHLQEMRQLLQKTD---TLQLLAYSET	51
AAC(6')-Ix	MNIMPISSEQLSDWLALRCLLWP-DHDDAHLLEMHQLLQKTD---TLQLLAYTDS	51
AAC(6')-Iy	MDIRQMNKTHLEHWRGLRKLQWPGHPDDAHLADGEEILQA-DH--LAFIAMADG	52
AAC(6')-Iz	MIASAPTIRQATPADAAAWAQLRLGLWP--DADDPLEELTQSLADAE---GAVFLACAD	55
	: * :	:
AAC(6')-Iad	-QAVGFAQCQLRHDYVEGNTNSPVVGLYLEGIFVEKEFRHRGYASELCLKCEDVWTKGCLQ	110
AAC(6')-Ic	GAFVAFAEVALRYDYVNGCESSPVAFLEGIYTABRARRQGWAAARLIAQVQEWAKQGCSE	111
AAC(6')-Id	NKPLGFVEAALRSYVNGTNSPVAFLEGIYVLEARRRRIAHAALVGVAVIWARNRACTE	111
AAC(6')-If	SQAIAFADAARHDYVNGCESSPVVYLEGIFVPEQRGHGVAKLLVAAVQDVGWVAKGCTE	108
AAC(6')-Ig	HQAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	110
AAC(6')-Ih	QQAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Ij	QQAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Ik	DHAVGMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	110
AAC(6')-Il	GEALGFAEALSIR-PYAECEYSGNVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	118
AAC(6')-Ir	QQAIVAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Is	QLAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-It	QKAIGMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Iu	QHAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Iv	QQAIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Iw	QQPIAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Ix	QQAVAMLEASIRFEYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Iy	-VAIGFADASIRHDYVNGCDSSPVVLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	111
AAC(6')-Iz	GETVGFAEVRLRHDYVNGTETSPVAFLEGIYVLPVLRPAHRRSGVATMLIRQAEVWAKQFSCTE	115
	::: : * * : : : * * * : : * * * : : * * * :	
AAC(6')-Iad	FASDCELDNIDSLAFHLKVGFTANRMICFTKQL	144
AAC(6')-Ic	LASDTDIANLDSQRLHAALGFATERVVYFRKTLG	146
AAC(6')-Id	FASDASTDNPESHRFHQSLGFKETERVVYFRKMLAPE	149
AAC(6')-If	MASDAALDNHISYQMHQALGFEBETERVVYFRKRIAG	144
AAC(6')-Ig	FASDAALDNVISHAMHRLGFGQTEKVVYFSSKID	145
AAC(6')-Ih	FASDAALDNQISHAMHQLGPHETERVVYFKNIG	146
AAC(6')-Ij	FASDAALDNRI SHAMHQLGPHETERVVYFKNIG	146
AAC(6')-Ik	FASDAALDNVISHAMHRLGFGQTEKVVYFSSKID	145
AAC(6')-Il	FASDTQLTNSASTSAHLAAGFTEVAQVRCFRKPL	152
AAC(6')-Ir	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Is	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-It	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Iu	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Iv	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Iw	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Ix	FASDAALDNVISHAMHRLGFGQTEKVVYFKNIS	146
AAC(6')-Iy	MASDTSPENTISQKVHQLGFEETERVYFRKRC	145
AAC(6')-Iz	LASDSRVEDVQAAHAHRACGFEETERVVYFRMPLEPSA	153
	: * * * : : * * * : : :	

FIG. 1. Alignment of the deduced amino acid sequences of AAC(6')-Iad and other aminoglycoside acetyltransferases, including AAC(6')-Ic (GenBank accession no. M94066), AAC(6')-Id (X12618), AAC(6')-If (X55353), AAC(6')-Ig (L09246), AAC(6')-Ih (L29044), AAC(6')-Ij (L29045), AAC(6')-Ik (L29510), AAC(6')-Il (Z54241, U13880), AAC(6')-Ir (AF031326), AAC(6')-Is (AF031327), AAC(6')-It (AF031328), AAC(6')-Iu (AF031329), AAC(6')-Iv (AF031330), AAC(6')-Iw (AF031331), AAC(6')-Ix (AF031332), AAC(6')-Iy (AF144880), and AAC(6')-Iz (AF140221). Asterisks indicate identical amino acids. Conservative amino acid substitutions are indicated by dots.




FIG. 2. Dendrogram for aminoglycoside 6'-N-acetyltransferases belonging to the subfamily represented by AAC(6')-Ic. The dendrogram was calculated by the ClustalW computer program, available on the National Institute of Genetics website (<http://www.ddbj.nig.ac.jp/E-mail/clustalw-e.html>), and illustrated with the TreeViewPPC computer program (version 1.6.5 for Macintosh). Branch lengths correspond to the numbers of amino acid exchanges.

cin, while pA51SG5 conferred resistance to kanamycin, gentamicin, tobramycin, and sisomicin. Neither plasmid conferred resistance to streptomycin or neomycin.

pA51S3 contained a 1.0-kb insert with one open reading frame, though several possible start codons were recognized. Therefore, N-terminal sequencing of the purified protein was carried out. Consequently, it was confirmed that the open reading frame encodes 144 amino acids and has a G+C content of 36.1%. The deduced amino acid sequence displayed the highest identity with that of AAC(6')-Iy (36.7%) (13). It also showed moderate identities with the sequences of other aminoglycoside acetyltransferases [35.2% with AAC(6')-If, 34.6% with AAC(6')-Ic, 33.3% with AAC(6')-Iz, and 29.7% with AAC(6')-II] (3, 11, 23, 26). The motifs that are conserved among the aminoglycoside 6'-N-acetyltransferases (24) were also found in the newly identified enzyme. This novel aminoglycoside acetyltransferase gene was thus designated *aac(6')-Iad*. The deduced amino acid sequence of AAC(6')-Iad is shown in Fig. 1, along with those of known aminoglycoside acetyltransferases. The dendrogram of phylogenetic relationships among aminoglycoside acetyltransferases is shown in Fig. 2. The 1.1-kb insert of pA51SG5 contained an aminoglycoside (2') adenylyltransferase gene, *ant(2'')-Ia* (4).

Identification of site of modification. The results of HPLC assays are shown in Table 2. The retention times of *o*-phthalaldehyde derivatives of tobramycin and neomycin after the acetylation reaction with AAC(6')-Iad coincided only with those of positive controls for AAC(6'), confirming that AAC(6')-Iad is a functional acetyltransferase and modifies position 6' of aminoglycosides.

Transfer of aminoglycoside resistance. The amikacin resistance determinant of *A. baumannii* A-67 and A-74 could be transferred to the recipient *A. calcoaceticus* DU1 by conjugation at a frequency of approximately 5×10^{-4} to 1×10^{-3} and was confirmed by PCR to be *aac(6')-Iad*. It was not transferred to *E. coli* CSH2. For the rest of the strains, amikacin resistance was not transferable to *A. calcoaceticus* DU1 or *E. coli* CSH2. The DNA probes for detection of *aac(6')-Iad* hybridized with the large plasmids (>50 kb) harbored by all seven strains (data not shown).

DISCUSSION

A variety of aminoglycoside 6'-N-acetyltransferase genes from *Acinetobacter* species have been described to date (Fig. 2). *aac(6')-Ib* and *aac(6')-Ih* have been identified previously as

TABLE 2. Retention times of aminoglycoside modification products after acetylation reactions

Aminoglycoside acetyltransferase	Retention time (min) of aminoglycoside modification product			
	Tobramycin		Neomycin	
	With acetyl-CoA	Without acetyl-CoA	With acetyl-CoA	Without acetyl-CoA
AAC(6')-Iad	3.3	17.0	4.9	11.8
Positive controls				
AAC(6')	3.3	16.9	4.9	11.8
AAC(2')	11.3	16.9	10.5	11.9
AAC(3')	4.4	16.9	6.7	11.9

the most prevalent plasmid-mediated *aac(6')-I* genes among *A. baumannii* strains (18), while other genes have been associated with specific species. *aac(6')-I_g* is specific to *Acinetobacter haemolyticus* (10), whereas *aac(6')-I_j* and *aac(6')-I_k* are specific to *Acinetobacter* genospecies 13 and 6, respectively (9, 19). *aac(6')-I_r*, *aac(6')-I_s*, *aac(6')-I_t*, *aac(6')-I_u*, *aac(6')-I_v*, *aac(6')-I_w*, and *aac(6')-I_x* have also been described for various *Acinetobacter* species (20). However, *aac(6')-Iad* demonstrated considerable phylogenetic distance from these aminoglycoside-modifying enzymes (as shown in Fig. 2), suggesting the emergence of a novel subgroup of aminoglycoside 6'-*N*-acetyltransferases.

In the present study, we report identification of a novel aminoglycoside 6'-*N*-acetyltransferase gene, *aac(6')-Iad*, in seven clinical isolates belonging to *A. baumannii* and *Acinetobacter* genospecies 3. The spectrum of resistance conferred by the gene product included kanamycin, tobramycin, amikacin, isepamicin, and sisomicin, a pattern typical of AAC(6')-I (22). Preliminary sequencing results suggest that *aac(6')-Iad* is located on a transposon (data not shown); in view of this possibility, along with the fact that the gene is transferable by conjugation in some of the producers of the enzyme, it is likely that *aac(6')-Iad* is carried by a plasmid.

Three subgroups have been identified among aminoglycoside 6'-*N*-acetyltransferases (22). AAC(6')-Iad is closest to the largest subfamily, which contains the proteins mentioned above as identified in *Acinetobacter* species, but the amino acid sequence identity between AAC(6')-Iad and these proteins is limited ($\leq 36.7\%$) (Fig. 1). Considering the low G+C content (36.1%) of *aac(6')-Iad* for *Acinetobacter* species, we may speculate that the gene was acquired from some environmental species with an intrinsically low G+C content.

PFGE of the seven strains that produce AAC(6')-Iad showed five distinct digestion patterns, except for those isolated from the same hospital (data not shown). Taken together, it is likely that *aac(6')-Iad* was disseminated among *Acinetobacter* spp. via plasmid- and transposon-mediated lateral transfer, which is now responsible for reduced susceptibility to amikacin among *Acinetobacter* spp. in nearly half of the cases (7 out of 16 non-amikacin-susceptible strains) in Japan.

When the susceptibilities of the AAC(6')-Iad producers to other classes of antimicrobial agents were tested, we found that none were susceptible to ceftazidime, moxalactam, or aztreonam, and two were resistant to ciprofloxacin as well. Only imipenem and meropenem were uniformly effective in vitro among the agents tested. The emergence and spread of plasmid-mediated *aac(6')-Iad* genes could contribute to further acquisition of a multidrug-resistant phenotype among *Acinetobacter* spp. in Japan, thus limiting the treatment options in clinical settings in the near future.

ACKNOWLEDGMENTS

We thank all the participating medical institutions for providing bacterial strains. We also thank Kunimoto Hotta for the gift of AAC-producing strains and Kumiko Kai for technical assistance.

This work was supported by grants H12-Shinkou-19, H12-Shinkou-20, H15-Shinkou-9, and H15-Shinkou-10 from the Ministry of Health, Labor, and Welfare of Japan.

REFERENCES

- Bergogne-Berezin, E., and K. J. Towner. 1996. *Acinetobacter* spp. as nosocomial pathogens: microbiological, clinical, and epidemiological features. *Clin. Microbiol. Rev.* 9:148-165.
- Bouvet, P. J. M., and P. A. D. Grimont. 1986. Taxonomy of the genus *Acinetobacter* with the recognition of *Acinetobacter baumannii* sp. nov., *Acinetobacter haemolyticus* sp. nov., *Acinetobacter johnsonii* sp. nov., and *Acinetobacter junii* sp. nov., and emended descriptions of *Acinetobacter calcoaceticus* and *Acinetobacter lwoffii*. *Int. J. Syst. Bacteriol.* 36:228-240.
- Bunny, K. L., R. M. Hall, and H. W. Stokes. 1995. New mobile gene cassettes containing an aminoglycoside resistance gene, *aacA7*, and a chloramphenicol resistance gene, *catB3*, in an integron in pBWH301. *Antimicrob. Agents Chemother.* 39:686-693.
- Cameron, F. H., D. J. Groot Obbink, V. P. Ackerman, and R. M. Hall. 1986. Nucleotide sequence of the AAD(2⁺) aminoglycoside adenyltransferase determinant *aadB*. Evolutionary relationship of this region with those surrounding *aadA* in R538-1 and *dhfrII* in R388. *Nucleic Acids Res.* 14:8625-8635.
- Cisneros, J. M., and J. Rodríguez-Bano. 2002. Nosocomial bacteremia due to *Acinetobacter baumannii*: epidemiology, clinical features and treatment. *Clin. Microbiol. Infect.* 8:687-693.
- de Vries, J., and W. Wackernagel. 2002. Integration of foreign DNA during natural transformation of *Acinetobacter* sp. by homology-facilitated illegitimate recombination. *Proc. Natl. Acad. Sci. USA* 99:2094-2099.
- Doi, Y., K. Yokoyama, K. Yamane, J. Wachino, N. Shibata, T. Yagi, K. Shibayama, H. Kato, and Y. Arakawa. 2004. Identification of a plasmid-mediated 16S rRNA methylase in *Serratia marcescens* conferring high-level resistance to various aminoglycosides. *Antimicrob. Agents Chemother.* 48:491-496.
- Hotta, K., A. Sunada, Y. Ikeda, and S. Kondo. 2000. Double stage activity in aminoglycoside antibiotics. *J. Antibiot. (Tokyo)* 53:1168-1174.
- Lambert, T., G. Gerbaud, and P. Courvalin. 1994. Characterization of the chromosomal *aac(6')-I_j* gene of *Acinetobacter* sp. 13 and the *aac(6')-I_h* plasmid gene of *Acinetobacter baumannii*. *Antimicrob. Agents Chemother.* 38:1883-1889.
- Lambert, T., G. Gerbaud, M. Galimand, and P. Courvalin. 1993. Characterization of *Acinetobacter haemolyticus aac(6')-I_g* gene encoding an aminoglycoside 6'-*N*-acetyltransferase which modifies amikacin. *Antimicrob. Agents Chemother.* 37:2093-2100.
- Lambert, T., M. C. Ploy, F. Denis, and P. Courvalin. 1999. Characterization of the chromosomal *aac(6')-I_z* gene of *Stenotrophomonas maltophilia*. *Antimicrob. Agents Chemother.* 43:2366-2371.
- Lovering, A. M., L. O. White, and D. S. Reeves. 1984. Identification of aminoglycoside-acetylating enzymes by high-pressure liquid chromatographic determination of their reaction products. *Antimicrob. Agents Chemother.* 26:10-12.
- Magnet, S., P. Courvalin, and T. Lambert. 1999. Activation of the cryptic *aac(6')-I_y* aminoglycoside resistance gene of *Salmonella* by a chromosomal deletion generating a transcriptional fusion. *J. Bacteriol.* 181:6650-6655.
- Miller, G. H., F. J. Sabatelli, R. S. Hare, Y. Glupczyński, P. Mackey, D. Shlaes, K. Shimizu, K. J. Shaw, et al. 1997. The most frequent aminoglycoside resistance mechanisms—changes with time and geographic area: a reflection of aminoglycoside usage patterns? *Clin. Infect. Dis.* 24(Suppl. 1):S46-S62.
- Miller, G. H., F. J. Sabatelli, L. Naples, R. S. Hare, K. J. Shaw, et al. 1995. The most frequently occurring aminoglycoside resistance mechanisms—combined results of surveys in eight regions of the world. *J. Chemother.* 7(Suppl. 2):17-30.
- National Committee for Clinical Laboratory Standards. 2003. Performance standards for antimicrobial susceptibility testing; 13th informational supplement. M100-S13. National Committee for Clinical Laboratory Standards, Wayne, Pa.
- Nielsen, K. M., M. D. van Weerelt, T. N. Berg, A. M. Bones, A. N. Hagler, and J. D. van Elsas. 1997. Natural transformation and availability of transforming DNA to *Acinetobacter calcoaceticus* in soil microcosms. *Appl. Environ. Microbiol.* 63:1945-1952.
- Ploy, M. C., H. Giamarellou, P. Bourlioux, P. Courvalin, and T. Lambert. 1994. Detection of *aac(6')-I* genes in amikacin-resistant *Acinetobacter* spp. by PCR. *Antimicrob. Agents Chemother.* 38:2925-2928.
- Rudant, E., P. Bourlioux, P. Courvalin, and T. Lambert. 1994. Characterization of the *aac(6')-I_k* gene of *Acinetobacter* sp. 6. *FEMS Microbiol. Lett.* 124:49-54.
- Rudant, E., P. Bouvet, P. Courvalin, and T. Lambert. 1999. Phylogenetic analysis of proteolytic *Acinetobacter* strains based on the sequence of genes encoding aminoglycoside 6'-*N*-acetyltransferases. *Syst. Appl. Microbiol.* 22:59-67.
- Seward, R. J., T. Lambert, and K. J. Towner. 1998. Molecular epidemiology of aminoglycoside resistance in *Acinetobacter* spp. *J. Med. Microbiol.* 47:455-462.
- Shaw, K. J., P. N. Rather, R. S. Hare, and G. H. Miller. 1993. Molecular

- genetics of aminoglycoside resistance genes and familial relationships of the aminoglycoside-modifying enzymes. *Microbiol. Rev.* 57:138-163.
23. Shaw, K. J., P. N. Rather, F. J. Sabatelli, P. Mann, H. Mumayyer, R. Mierzwa, G. L. Petrikos, R. S. Hare, G. H. Miller, P. M. Bennett, and P. Downey. 1992. Characterization of the chromosomal *aac(6')*-Ic gene from *Serratia marcescens*. *Antimicrob. Agents Chemother.* 36:1447-1455.
 24. Shmara, A., N. Weisnetel, K. J. Dery, R. Chavideh, and M. E. Tolmusk. 2001. Systematic analysis of a conserved region of the aminoglycoside 6'-N-acetyltransferase type Ib. *Antimicrob. Agents Chemother.* 45:3287-3292.
 25. Sugino, Y., Y. Hinuma, T. Nada, Y. Tawada, H. Amano, T. Nakamura, Y. Hasegawa, K. Shimokata, N. Shibata, and Y. Arakawa. 2001. Antimicrobial activities and mechanisms of carbapenem resistance in clinical isolates of carbapenem-resistant *Pseudomonas aeruginosa* and *Acinetobacter* spp. *Kansenshogaku Zasshi* 75:662-670. (In Japanese.)
 26. Teran, F. J., J. E. Suarez, and M. C. Mendoza. 1991. Cloning, sequencing, and use as a molecular probe of a gene encoding an aminoglycoside 6'-N-acetyltransferase of broad substrate profile. *Antimicrob. Agents Chemother.* 35:714-719.
 27. Theaker, C., B. Azadian, and N. Soni. 2003. The impact of *Acinetobacter baumannii* in the intensive care unit. *Anaesthesia* 58:271-274.

Inhibitor-Sensitive AmpC β -Lactamase Variant Produced by an *Escherichia coli* Clinical Isolate Resistant to Oxyiminocephalosporins and Cephameycins

Yohei Doi,¹ Jun-ichi Wachino,¹ Masaji Ishiguro,² Hiroshi Kurokawa,¹ Kunikazu Yamane,¹ Naohiro Shibata,¹ Keigo Shibayama,¹ Keiko Yokoyama,¹ Haru Kato,¹ Tetsuya Yagi,¹ and Yoshichika Arakawa^{1*}

Department of Bacterial Pathogenesis and Infection Control, National Institute of Infectious Diseases, Tokyo,¹ and Suntory Institute for Bioorganic Research, Osaka,² Japan

Received 28 June 2003/Returned for modification 8 November 2003/Accepted 3 March 2004

Escherichia coli HKY28, a ceftazidime-resistant strain isolated from a urine specimen in Japan, produced an inhibitor-sensitive AmpC β -lactamase variant. The deduced amino acid sequence of the enzyme contained a number of substitutions and a tripeptide deletion (Gly286-Ser287-Asp288) compared with the sequence of native AmpC of *E. coli*. When the deletion was reverted by a 9-base insertion at the relevant site of *ampC* in the clone, the typical inhibitor-resistant phenotype of AmpC was restored, while at the same time the levels of resistance to ceftazidime, cefpirome, and cefepime were reduced eightfold or more. Molecular modeling studies indicated that a structural change took place in the H-10 helix as a result of the deletion, and this change caused an alteration of the substrate binding site, leading to a unique phenotype analogous to that of inhibitor-sensitive class A extended-spectrum β -lactamases. The degree of inhibition was greater with sulbactam and tazobactam than with clavulanic acid. To our knowledge, this is the first report to have characterized an *E. coli ampC* that encodes chromosomal AmpC β -lactamase sensitive to the available β -lactamase inhibitors.

The principal and most prevalent mechanism of resistance to β -lactam agents among pathogenic gram-negative bacteria is the production of β -lactamases (3, 17). One approach to overcoming the problem has been the development of β -lactams resistant to the hydrolytic activities of these enzymes. The other has been the development of β -lactamase inhibitors, which protect β -lactams from hydrolysis by β -lactamases when the inhibitors are used in combination with β -lactams (28). At present, three β -lactamase inhibitors, clavulanic acid, sulbactam, and tazobactam, are available for clinical use in combination with a number of penicillins. These inhibitors mainly target Ambler class A β -lactamases and inactivate their active-site serines, thus potentiating the actions of β -lactamase-sensitive compounds. Clavulanic acid and sulbactam are generally not effective in inhibiting the activities of AmpC β -lactamases, although some are known to be moderately inhibited by tazobactam (4, 14).

In 1994, we isolated an *Escherichia coli* clinical strain, HKY28, which produced a chromosomal AmpC β -lactamase that had an inhibitor-sensitive and extended-spectrum activity profile similar to those of class A extended-spectrum β -lactamases (ESBLs). However, the results of PCR experiments with representative TEM- and SHV-derived ESBLs and CTX-M-type β -lactamases were negative. In the present study we con-

ducted genetic, biochemical, and molecular modeling analyses of this unique AmpC β -lactamase variant.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. *E. coli* HKY28 was isolated from a culture of urine from an inpatient in Japan in 1994. *E. coli* XLI-Blue (Stratagene, La Jolla, Calif.) was used as the recipient strain for plasmids. *E. coli* BMH71-18mutS and *E. coli* MV1184 (Takara Bio Inc., Ohtsu, Japan) were used as the hosts in a site-directed mutagenesis experiment. Plasmid vectors pBCKS+ (Stratagene) and pKF18k (Takara Bio) were used for the cloning and site-directed mutagenesis experiments, respectively. For enzyme purification, *ampC*-deficient *E. coli* CS14-2 (7) was used as the host to avoid background AmpC production. Bacteria were grown in Luria-Bertani (LB) broth supplemented with the appropriate antibiotics, unless specified otherwise.

Antibiotics and susceptibility testing. The following β -lactam antibiotics and β -lactamase inhibitors were obtained from the indicated sources: aztreonam, Eisai Co., Ltd., Tokyo, Japan; ampicillin, amoxicillin, and cefminox, Meiji Seika Kaisha, Ltd., Tokyo, Japan; cefepime, Bristol Pharmaceuticals K. K., Tokyo, Japan; cefmetazole and chloramphenicol, Sankyo Co., Ltd., Tokyo, Japan; cefotaxime and cefpirome, Aventis Pharma, Ltd., Tokyo, Japan; cefoxitin and imipenem, Banyu Pharmaceutical Co., Ltd., Tokyo, Japan; ceftazidime and clavulanic acid, GlaxoSmithKline K. K., Tokyo, Japan; cephaloridine and moxalactam, Shionogi & Co., Ltd., Osaka, Japan; sulbactam, Pfizer Pharmaceuticals Inc., Tokyo, Japan; and tazobactam, Taiho Pharmaceutical Co., Ltd., Tokyo, Japan.

MICs were determined by the agar dilution method by the protocol recommended by the National Committee for Clinical Laboratory Standards (18).

PCR amplification. To amplify broad-spectrum β -lactamase genes from HKY28, PCR analysis was performed with sets of primers for various β -lactamases, including TEM- and SHV-derived ESBLs as well as CTX-M-1-, CTX-M-2-, and CTX-M-9-type β -lactamases, as described previously (27).

Transfer of ceftazidime resistance. Conjugation experiments were conducted with *E. coli* CSH2 as the recipient by broth mating and filter mating methods (7). Transconjugants were selected on LB agar supplemented with rifampin (50 μ g/ml), nalidixic acid (50 μ g/ml), and ceftazidime (4 μ g/ml).

Cloning and sequencing of β -lactamase gene. The basic recombinant DNA manipulations were carried out as described by Sambrook et al. (24). The

* Corresponding author. Mailing address: Department of Bacterial Pathogenesis and Infection Control, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-Murayama, Tokyo 208-0011, Japan. Phone: 81-42-561-0771, ext. 500. Fax: 81-42-561-7173. E-mail: yarakawa@nih.go.jp.

genomic DNA of HKY28 was prepared and digested with EcoRI. The resultant fragments were ligated with plasmid vector pBCKS+, and electrocompetent *E. coli* XL1-Blue was transformed with these recombinant plasmids. Transformants were selected for resistance to chloramphenicol (30 µg/ml) and ceftazidime (4 µg/ml). For determination of the MICs and use of the transformants for site-directed mutagenesis, the *ampC* gene of HKY28 was amplified with oligonucleotide primers *ampC*-U (5'-CGG AAT TCG GTT TTC TAC GGT CTG GC-3') and *ampC*-L (5'-CGG GAT CCG ATG ACA GCA AGG AAA AG-3'), which contained EcoRI and BamHI cleavage sites (indicated in boldface), respectively, at their 5' ends, by using Pyrobest DNA polymerase (Takara Bio). The EcoRI-BamHI fragment containing the *ampC* gene of *E. coli* HKY28 was ligated with pBCKS+ to yield pBE28W, which was then used to transform *E. coli* XL1-Blue and *E. coli* CS14-2. The coding sequences of the cloned fragments were determined by using custom sequencing primers as well as a BigDye Terminator Cycle Sequencing Ready Reaction kits and an ABI 3100 DNA sequencer (Applied Biosystems, Foster City, Calif.). The enzymes used for gene manipulations were purchased from Nippon Gene Co. Ltd. (Tokyo, Japan) or New England Biolabs, Inc. (Beverly, Mass.).

Reversion of AmpC deletion. Site-directed mutagenesis was performed to revert the 9-nucleotide deletion in the cloned *ampC* gene of *E. coli* HKY28 corresponding to a tripeptide deletion at positions 286 to 288 in AmpC. The reagents and strains contained in the Mutan-Express Km mutagenesis kit (Takara Bio) were used according to the procedures based on the oligonucleotide-directed dual Amber method (9) provided by the manufacturer. The following mutagenic primer containing the 9-nucleotide insertion (in boldface) was used: 5'-CCA GTG CAA TTT TAT TGT CAC TGC CGT TAA TGA TGA TGT CAG G-3'. After mutagenesis, the EcoRI-BamHI fragment containing the revertant *ampC* was ligated with pBCKS+ to yield pBE28R, which was then used to transform *E. coli* XL1-Blue and *E. coli* CS14-2.

Enzyme purification. *E. coli* CS14-2 harboring pBE28W or pBE28R was cultured overnight in 2 liters of LB broth supplemented with 30 µg of chloramphenicol per ml. Cells were harvested by centrifugation and washed with and then suspended in 3 ml of 50 mM 3-(*N*-morpholino)propanesulfonic acid (MOPS) buffer (pH 6.0). The cells were frozen and thawed twice and were then ultracentrifuged at 100,000 × g for 4 h at 4°C. For gel filtration, the supernatant containing β-lactamase was chromatographed through a HiLoad 16/60 Superdex 200 prepgrade (Pharmacia Biotech, Uppsala, Sweden) column preequilibrated with 50 mM MOPS buffer (pH 6.0). For cation-exchange chromatography, fractions with activity were then applied to a HiTrap SP HP column (Pharmacia Biotech) preequilibrated with the same buffer. The enzymes were eluted with a linear gradient of 0 to 0.5 M NaCl in the same buffer. The purity of the enzymes was checked by a sodium dodecyl sulfate-polyacrylamide gel electrophoresis.

Enzyme assays. Purified AmpC enzymes were assayed against various β-lactam substrates at 37°C in 50 mM phosphate buffer (pH 7.0) by using an auto-spectrophotometer (V-550; Nihon Bunko Ltd., Tokyo, Japan). The specific activity of the enzymes was defined as the activity that hydrolyzed 1 µmol of cephaloridine per min. K_m and k_{cat} values were obtained by a direct-weight fit to the Michaelis-Menten equation by using KaleidaGraph software (Hulinks, Tokyo, Japan). The concentrations of inhibitors giving a 50% reduction in hydrolysis of cephaloridine (IC_{50}) were measured after 10 min of preincubation of the enzymes with the inhibitors at 37°C and cephaloridine as the substrate at 1 mM. The affinities of the enzymes for the inhibitors (K_i s) were measured by competition procedures with cephaloridine in the same buffer with no preincubation of the enzyme or the inhibitor. To determine the isoelectric points, 10 µl of enzyme solution was loaded onto an Immobiline DryStrip (pH 3 to 10 and 6 to 11; Pharmacia Biotech), and electrophoresis was carried out with an IPGphor electrophoresis system (Pharmacia Biotech).

Modeling of substrate-enzyme complex structures. The crystal structure of the AmpC β-lactamase (Protein Data Bank accession number 2BLS) was used as the reference to build a model of the AmpC enzyme of *E. coli* HKY28. The tripeptide at the H-10 helix was deleted by the loop search method of the Homology module installed in Insight II software (version 2000; Molecular Simulations Inc., San Diego, Calif.). An initial structure of the enzyme was optimized by use of molecular dynamics calculations at 298 K by the cell multipole method, a distance-dependent dielectric constant, and a time step of 1 fs for 100 ps by sampling the conformation every 1 ps by use of Discover 3 software (version 98.0; Molecular Simulations Inc.). One hundred conformations were minimized until the final root-mean square deviation became less than 0.1 kcal/mol/Å, and the lowest energy conformation was selected for the substrate-docking study. The substrates were roughly docked into the ligand-binding cleft with the guidance of a hydrogen bond of a β-lactam carbonyl oxygen at the oxyanion hole as well as a hydrogen bond of the carboxylate oxygen with Tyr150 (12). The initial complex model was minimized, and then the substrate-binding site was covered by water

molecules (sphere thickness, 20 Å). The structure consisted of the substrate and the residues within 10 Å from the substrate, which were energy optimized in the presence of the water molecules by the molecular dynamics and minimization procedure described above. The lowest-energy structures were selected as energy-refined complex models.

Nucleotide sequence accession number. The nucleotide sequence encoding AmpC characterized in this study appears in the EMBL/GenBank/DBJ databases under accession number AB108683.

RESULTS

Susceptibility of parental strain. The MICs of β-lactams for parental strain *E. coli* HKY28 are shown in Table 1. Strain HKY28 was resistant to ampicillin, amoxicillin, cephaloridine, cefminox, and cefoxitin. It was also resistant to ceftazidime (MIC, 32 µg/ml) but remained susceptible to aztreonam and imipenem. Interestingly, the MIC of ampicillin was reduced by at least 8-fold when it was combined with sulbactam, and the MICs of cefotaxime were reduced by 16- and 8-fold when it was combined with sulbactam and tazobactam, respectively. Addition of sulbactam reduced the MIC of ceftazidime by eightfold. Overall, the reductions in the MICs were the greatest with sulbactam, followed by tazobactam and clavulanic acid.

PCR analysis of β-lactamase genes. By PCR *E. coli* HKY28 was negative for the genes for the TEM-, SHV-, CTX-M-1-, CTX-M-2- and CTX-M-9-type β-lactamases, which are the prevalent types of ESBLs in Japan.

Transfer of ceftazidime resistance. The ceftazidime resistance of *E. coli* HKY28 could not be transferred to recipient *E. coli* strain CSH2 by conjugation, despite repeated attempts.

Cloning and sequencing of resistance gene. A 6-kb EcoRI fragment containing a ceftazidime resistance determinant was cloned into the vector pBCKS+ and was termed pE753. Nucleotide sequencing analysis revealed a chromosomal locus of *E. coli* containing *ampC* flanked by *frdD* and *ble* but without any other β-lactamase gene. PCR-generated recombinant plasmid pBE28W containing *ampC* of *E. coli* HKY28 was found to possess an *ampC* gene identical to that of pE753 and conferred resistance to ceftazidime. The deduced amino acid sequence contained seven amino acid substitutions and three amino acid deletions (Gly286, Ser287, and Asp288) of the AmpC product compared with the sequence of *E. coli* K-12 (10) (Fig. 1). The promoter region of the *ampC* gene contained three mutations (a C-to-T change at position -73, a C-to-T change at position +6, and a G-to-A change at position +34) and a T insertion between positions -14 and -13 compared with the sequence of the corresponding region of the *E. coli* K-12 genome. Recombinant plasmid pBE28R, generated by site-directed mutagenesis, was confirmed to possess *ampC* of *E. coli* HKY28, except for the insertion of the 9-nucleotide sequence designed to restore the tripeptide deleted from *ampC* of *E. coli* HKY28.

Susceptibilities of clones to β-lactams. Both *E. coli* XL1-Blue harboring pBE28W (the HKY28 clone) and that harboring pBE28R (the revertant clone) displayed resistance or reduced susceptibilities to all β-lactams except cefpirome, cefepime, and imipenem; but the degree of resistance varied significantly between the two clones. The cefotaxime and ceftazidime MICs were fourfold or more higher for the HKY28 clone than for the revertant clone. The cefpirome and cefepime MICs were 64-fold higher for the HKY28 clone than

TABLE 1. Results of antibiotic susceptibility testing

β -Lactam	MIC ($\mu\text{g/ml}$)			
	<i>E. coli</i> HKY28	<i>E. coli</i> XLI-Blue(pBE28 W)	<i>E. coli</i> XLI-Blue(pBE28 R)	<i>E. coli</i> XLI-Blue
Amoxicillin	128	>128	>128	4
Amoxicillin-clavulanate ^a	128	>128	>128	4
Ampicillin	>128	>128	>128	2
Ampicillin-sulbactam ^b	32	64	>128	2
Piperacillin	8	8	8	0.5
Piperacillin-tazobactam ^c	4	2	4	0.5
Cefotaxime	16	32	8	0.06
Cefotaxime-clavulanate ^a	8	8	2	0.06
Cefotaxime-sulbactam ^b	1	1	2	0.06
Cefotaxime-tazobactam ^c	2	4	4	0.06
Ceftazidime	32	128	16	0.06
Ceftazidime-clavulanate ^a	16	32	8	0.13
Ceftazidime-sulbactam ^b	4	8	8	0.06
Ceftazidime-tazobactam ^c	16	8	8	0.13
Cephaloridine	64	128	128	4
Cefminox	32	32	32	0.5
Cefoxitin	16	32	>128	8
Cefmetazole	16	32	128	1
Moxalactam	8	4	8	0.25
Cefpirome	2	4	0.03	0.015
Cefepime	2	4	0.03	0.015
Aztreonam	8	16	16	0.06
Imipenem	0.13	0.13	0.13	0.13

^a Fixed concentration of clavulanate, 4 $\mu\text{g/ml}$.

^b Fixed concentration of sulbactam, 4 $\mu\text{g/ml}$.

^c Fixed concentration of tazobactam, 4 $\mu\text{g/ml}$.

for the revertant clone. On the other hand, the degree of resistance to cefoxitin and cefmetazole conferred by the revertant clone was significantly higher than that conferred by the HKY28 clone. When various β -lactam- β -lactamase inhibitor combinations were tested, the piperacillin, cefotaxime, and ceftazidime MICs for the HKY28 clone were reduced by up to 16-fold. The degree of reduction was the greatest when sulbactam was used as the inhibitor. The reductions in the MICs of the three inhibitors for the revertant clone were fourfold or less.

Isoelectric focusing. The isoelectric points were estimated to be 9.9 for the HKY28 AmpC (AmpC^D) and 9.8 for the revertant AmpC (AmpC^R). When the crude extract of *E. coli* HKY28 was subjected to analytical isoelectric focusing, only one band corresponding to AmpC^D was visualized with nitrocefin, confirming that AmpC^D is the only β -lactamase produced by *E. coli* HKY28 (data not shown).

Enzyme assays. The specific activities of AmpC^D and AmpC^R were 88 and 220 U/mg of protein, respectively. The

kinetic parameters (K_m and k_{cat}) and hydrolytic efficiencies (k_{cat}/K_m) of AmpC^D and AmpC^R against various β -lactams are given in Table 2. The k_{cat} values of AmpC^D were greater than those of AmpC^R for cefpirome and cefepime but lower for the rest of the substrates tested. However, for all substrates with the exception of cefotaxime, AmpC^D exhibited lower K_m values than AmpC^R. This difference was approximately 100-fold for ceftazidime, and overall, AmpC^D showed a 2.5-fold greater hydrolytic efficiency for ceftazidime than AmpC^R, despite the much poorer k_{cat} . AmpC^D exhibited much lower K_m values and higher k_{cat} values for cefpirome and cefepime than AmpC^R, resulting in approximately 40- and 20-fold greater hydrolytic efficiencies, respectively.

The IC₅₀s of the β -lactamase inhibitors for AmpC^D and AmpC^R and the K_i values of the enzymes against the inhibitors are listed in Table 3. AmpC^D exhibited approximately 5- to 10-fold lower K_i values than AmpC^R against all three inhibitors. Tazobactam was the best inhibitor and had the lowest IC₅₀ for AmpC^D.

TABLE 2. Kinetic activity of AmpC^D and AmpC^R

Substrate	AmpC ^D			AmpC ^R		
	K_m (μM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1} \text{s}^{-1}$)	K_m (μM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1} \text{s}^{-1}$)
Cephaloridine	100 \pm 10	64 \pm 4	6.4 \times 10 ⁵	780 \pm 40	300 \pm 10	3.9 \times 10 ⁵
Ampicillin	6.2 \pm 1.6	0.43 \pm 0.06	7.0 \times 10 ⁴	13 \pm 3	4.0 \pm 0.3	3.2 \times 10 ⁵
Cefoxitin	1.2 \pm 0.3	0.043 \pm 0.006	3.8 \times 10 ⁴	3.9 \pm 0.1	0.35 \pm 0.01	9.1 \times 10 ⁴
Ceftazidime	5.7 \pm 0.8	0.084 \pm 0.006	1.5 \times 10 ⁴	550 \pm 10	3.5 \pm 0.1	6.4 \times 10 ³
Cefotaxime	31 \pm 8	0.37 \pm 0.02	1.2 \times 10 ⁴	13 \pm 2	1.2 \pm 0.1	9.7 \times 10 ⁴
Cefpirome	21 \pm 1	1.5 \pm 0.1	7.1 \times 10 ⁴	120 \pm 20	0.21 \pm 0.03	1.8 \times 10 ³
Cefepime	49 \pm 5	1.0 \pm 0.1	2.1 \times 10 ⁴	200 \pm 40	0.21 \pm 0.02	1.1 \times 10 ³