

## Acknowledgments

We thank Drs. H. Kato and H. Nagatomo for their valuable comments and suggestions, T. Kazama and K. Tamaru for technical assistance with the analyses, Drs. K. Sakai, K. Shinohara, and E. Takahashi for kindly providing SHIV-C2/1 KS661, and Dr. J. Raymond for editing the manuscript. This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education and Science, Japan, a Health Sciences Research Grant from the Ministry of Health, Labour and Welfare, Japan, and a Research Grant on Health Sciences focusing on Drug innovation from the Japan Health Science Foundation.

## References

- Ahlers, J.D., Dunlop, N., Alling, D.W., Nara, P.L., Berzofsky, J.A., 1997. Cytokine-in-adjuvant steering of the immune response phenotype to HIV-1 vaccine constructs: granulocyte-macrophage colony-stimulating factor and TNF-alpha synergize with IL-12 to enhance induction of cytotoxic T lymphocytes. *J. Immunol.* 158, 3947–3958.
- Ahlers, J.D., Belyakov, I.M., Berzofsky, J.A., 2003. Cytokine, chemokine, and costimulatory molecule modulation to enhance efficacy of HIV vaccines. *Curr. Mol. Med.* 3, 285–301.
- Ahmed, R.K., Makitalo, B., Karlen, K., Nilsson, C., Biberfeld, G., Thorstensson, R., 2002. Spontaneous production of RANTES and antigen-specific IFN-gamma production in macaques vaccinated with SHIV-4 correlates with protection against SIVsm challenge. *Clin. Exp. Immunol.* 129, 11–18.
- Alkhatib, G., Combadiere, C., Broder, C.C., Feng, Y., Kennedy, P.E., Murphy, P.M., Berger, E.A., 1996. CC CKR5: a RANTES, MIP-1alpha, MIP-1beta receptor as a fusion cofactor for macrophage-tropic HIV-1. *Science* 272, 1955–1958.
- Aukrust, P., Liabakk, N.B., Muller, F., Lien, E., Espevik, T., Froland, S.S., 1994. Serum levels of tumor necrosis factor-alpha (TNF alpha) and soluble TNF receptors in human immunodeficiency virus type 1 infection—Correlations to clinical, immunologic, and virologic parameters. *J. Infect. Dis.* 169, 420–424.
- Aukrust, P., Muller, F., Froland, S.S., 1998. Circulating levels of RANTES in human immunodeficiency virus type 1 infection: effect of potent anti-retroviral therapy. *J. Infect. Dis.* 177, 1091–1096.
- Baba, T.W., Liska, V., Khimani, A.H., Ray, N.B., Dailey, P.J., Penninck, D., Bronson, R., Greene, M.F., McClure, H.M., Martin, L.N., Ruprecht, R.M., 1999. Live attenuated, multiply deleted simian immunodeficiency virus causes AIDS in infant and adult macaques. *Nat. Med.* 5, 194–203.
- Badley, A.D., Dockrell, D., Simpson, M., Schut, R., Lynch, D.H., Leibson, P., Paya, C.V., 1997. Macrophage-dependent apoptosis of CD4<sup>+</sup> T lymphocytes from HIV-infected individuals is mediated by FasL and tumor necrosis factor. *J. Exp. Med.* 185, 55–64.
- Cocchi, F., DeVico, A.L., Garzino-Demo, A., Arya, S.K., Gallo, R.C., Lusso, P., 1995. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8<sup>+</sup> T cells. *Science* 270, 1811–1815.
- Cooper, D.A., Tindall, B., Wilson, E.J., Imrie, A.A., Penny, R., 1988. Characterization of T lymphocyte responses during primary infection with human immunodeficiency virus. *J. Infect. Dis.* 157, 889–896.
- Enose, Y., Ui, M., Miyake, A., Suzuki, H., Uesaka, H., Kuwata, T., Kunisawa, J., Kiyono, H., Takahashi, H., Miura, T., Hayami, M., 2002. Protection by intranasal immunization of a nef-deleted, nonpathogenic SHIV against intravaginal challenge with a heterologous pathogenic SHIV. *Virology* 298, 306–316.
- Enose, Y., Kita, M., Yamamoto, T., Suzuki, H., Miyake, A., Horiuchi, R., Ibuki, K., Kaneyasu, K., Kuwata, T., Takahashi, E., Sakai, K., Shinohara, K., Miura, T., Hayami, M., 2004. Protective effects of nef-deleted SHIV or that having IFN-gamma against disease induced with a pathogenic virus early after vaccination. *Arch. Virol.* 149, 1705–1720.
- Fauci, A.S., 1993. Multifactorial nature of human immunodeficiency virus disease: implications for therapy. *Science* 262, 1011–1018.
- Fauci, A.S., 1996. Host factors and the pathogenesis of HIV-induced disease. *Nature* 384, 529–534.
- Folks, T.M., Clouse, K.A., Justement, J., Rabson, A., Duh, E., Kehrl, J.H., Fauci, A.S., 1989. Tumor necrosis factor alpha induces expression of human immunodeficiency virus in a chronically infected T-cell clone. *Proc. Natl. Acad. Sci. U.S.A.* 86, 2365–2368.
- Garzino-Demo, A., Moss, R.B., Margolick, J.B., Cleghorn, F., Sill, A., Blattner, W.A., Cocchi, F., Carlo, D.J., DeVico, A.L., Gallo, R.C., 1999. Spontaneous and antigen-induced production of HIV-inhibitory beta-chemokines are associated with AIDS-free status. *Proc. Natl. Acad. Sci. U.S.A.* 96, 11986–11991.
- Giavedoni, L., Ahmad, S., Jones, L., Yilma, T., 1997. Expression of gamma interferon by simian immunodeficiency virus increases attenuation and reduces postchallenge virus load in vaccinated rhesus macaques. *J. Virol.* 71, 866–872.
- Godard, C.M., Chernmann, J.C., 1998. Stimulation of bystander T-cell proliferation by tumor necrosis factor produced by HIV-1-infected macrophages. *J. Hum. Virol.* 1, 257–266.
- Gougeon, M.L., Lecoer, H., Dulioust, A., Enouf, M.G., Crouvoiser, M., Goujard, C., Debord, T., Montagnier, L., 1996. Programmed cell death in peripheral lymphocytes from HIV-infected persons: increased susceptibility to apoptosis of CD4 and CD8 T cells correlates with lymphocyte activation and with disease progression. *J. Immunol.* 156, 3509–3520.
- Griffin, D.E., Ward, B.J., Esolen, L.M., 1994. Pathogenesis of measles virus infection: an hypothesis for altered immune responses. *J. Infect. Dis.* 170, S24–S31.
- Gundlach, B.R., Linhart, H., Dittmer, U., Sopper, S., Reiprich, S., Fuchs, D., Fleckenstein, B., Hunsmann, G., Stahl-Hennig, C., Uberla, K., 1997. Construction, replication, and immunogenic properties of a simian immunodeficiency virus expressing interleukin-2. *J. Virol.* 71, 2225–2232.
- Gundlach, B.R., Lewis, M.G., Sopper, S., Schnell, T., Sodroski, J., Stahl-Hennig, C., Uberla, K., 2000. Evidence for recombination of live, attenuated immunodeficiency virus vaccine with challenge virus to a more virulent strain. *J. Virol.* 74, 3537–3542.
- Haga, T., Kuwata, T., Kozyrev, I., Kwofie, T.B., Hayami, M., Miura, T., 2000. Construction of an SIV/HIV type 1 chimeric virus with the human interleukin 6 gene and its production of interleukin 6 in monkey and human cells. *AIDS Res. Hum. Retrovir.* 16, 577–582.
- Haga, T., Shimizu, Y., Okoba, M., Kumabe, S., Goto, Y., Shinjo, T., Ichimura, H., Kuwata, T., Hayami, M., Miura, T., 2002. Construction and in vitro properties of chimeric simian and human immunodeficiency virus with the human TNF-alpha gene. *Microbiol. Immunol.* 46, 849–855.
- Herbein, G., Gordon, S., 1997. 55- and 75-kilodalton tumor necrosis factor receptors mediate distinct actions in regard to human immunodeficiency virus type 1 replication in primary human macrophages. *J. Virol.* 71, 4150–4156.
- Herbein, G., O'Brien, W.A., 2000. Tumor necrosis factor (TNF)-alpha and TNF receptors in viral pathogenesis. *Proc. Soc. Exp. Biol. Med.* 223, 241–257.
- Herbein, G., Mahlknecht, U., Batliwalla, F., Gregersen, P., Pappas, T., Butler, J., O'Brien, W.A., Verdin, E., 1998. Apoptosis of CD8<sup>+</sup> T cells is mediated by macrophages through interaction of HIV gp120 with chemokine receptor CXCR4. *Nature* 395, 189–194.
- Iida, T., Ichimura, H., Shimada, T., Ibuki, K., Ui, M., Tamaru, K., Kuwata, T., Yonehara, S., Imanishi, J., Hayami, M., 2000. Role of apoptosis induction in both peripheral lymph nodes and thymus in progressive loss of CD4<sup>+</sup> cells in SHIV-infected macaques. *AIDS Res. Hum. Retrovir.* 16, 9–18.
- Johnson, R.P., Desrosiers, R.C., 1998. Protective immunity induced by live attenuated simian immunodeficiency virus. *Curr. Opin. Immunol.* 10, 436–443.
- Kasahara, S., Ando, K., Saito, K., Sekikawa, K., Ito, H., Ishikawa, T., Ohnishi, H., Seishima, M., Kakumu, S., Moriwaki, H., 2003. Lack of tumor necrosis factor alpha induces impaired proliferation of hepatitis B virus-specific cytotoxic T lymphocytes. *J. Virol.* 77, 2469–2476.
- Kozyrev, I.L., Miura, T., Takemura, T., Kuwata, T., Ui, M., Ibuki, K., Iida, T., Hayami, M., 2002. Co-expression of interleukin-5 influences

- replication of simian/human immunodeficiency viruses in vivo. *J. Gen. Virol.* 83, 1183–1188.
- Kuwata, T., Igarashi, T., Ido, E., Jin, M., Mizuno, A., Chen, J., Hayami, M., 1995. Construction of human immunodeficiency virus 1/simian immunodeficiency virus strain mac chimeric viruses having vpr and/or nef of different parental origins and their in vitro and in vivo replication. *J. Gen. Virol.* 76, 2181–2191.
- Kwofie, T.B., Haga, T., Iida, T., Hayami, M., Miura, T., 2000. Plasma levels of the chemokine RANTES in macaque monkeys infected with pathogenic and non-pathogenic SIV/HIV-1 chimeric viruses at an early stage of infection. *J. Vet. Med. Sci.* 62, 1311–1312.
- Lane, B.R., Markovitz, D.M., Woodford, N.L., Rochford, R., Strieter, R.M., Coffey, M.J., 1999. TNF- $\alpha$  inhibits HIV-1 replication in peripheral blood monocytes and alveolar macrophages by inducing the production of RANTES and decreasing C-C chemokine receptor 5 (CCR5) expression. *J. Immunol.* 163, 3653–3661.
- Lazdins, J.K., Grell, M., Walker, M.R., Woods-Cook, K., Scheurich, P., Pfizenmaier, K., 1997. Membrane tumor necrosis factor (TNF) induced cooperative signaling of TNFR60 and TNFR80 favors induction of cell death rather than virus production in HIV-infected T cells. *J. Exp. Med.* 185, 81–90.
- Matsuyama, T., Kobayashi, N., Yamamoto, N., 1991. Cytokines and HIV infection: is AIDS a tumor necrosis factor disease? *AIDS* 5, 1405–1417.
- Navikas, V., Link, J., Persson, C., Olsson, T., Hojeberg, B., Ljungdahl, A., Link, H., Wahren, B., 1995. Increased mRNA expression of IL-6, IL-10, TNF- $\alpha$ , and perforin in blood mononuclear cells in human HIV infection. *J. Acquired Immune Defic. Syndr. Hum. Retrovir.* 9, 484–489.
- Pedersen, C., Dickmeiss, E., Gaub, J., Ryder, L.P., Platz, P., Lindhardt, B.O., Lundgren, J.D., 1990. T-cell subset alterations and lymphocyte responsiveness to mitogens and antigen during severe primary infection with HIV: a case series of seven consecutive HIV seroconverters. *AIDS* 4, 523–526.
- Poli, G., Kinter, A., Justement, J.S., Kehrl, J.H., Bressler, P., Stanley, S., Fauci, A.S., 1990. Tumor necrosis factor alpha functions in an autocrine manner in the induction of human immunodeficiency virus expression. *Proc. Natl. Acad. Sci. U.S.A.* 87, 782–785.
- Reinberger, S., Spring, M., Nisslein, T., Stahl-Hennig, C., Hunsmann, G., Dittmer, U., 1999. Kinetics of lymphocyte apoptosis in macaques infected with different simian immunodeficiency viruses or simian/human immunodeficiency hybrid viruses. *Clin. Immunol.* 90, 141–146.
- Rosenberg, E.S., Billingsley, J.M., Caliendo, A.M., Boswell, S.L., Sax, P.E., Kalams, S.A., Walker, B.D., 1997. Vigorous HIV-1-specific CD4<sup>+</sup> T cell responses associated with control of viremia. *Science* 278, 1447–1450.
- Shinohara, K., Sakai, K., Ando, S., Ami, Y., Yoshino, N., Takahashi, E., Someya, K., Suzaki, Y., Nakasone, T., Sasaki, Y., Kaizu, M., Lu, Y., Honda, M., 1999. A highly pathogenic simian/human immunodeficiency virus with genetic changes in cynomolgus monkey. *J. Gen. Virol.* 80, 1231–1240.
- Stahl-Hennig, C., Gundlach, B.R., Dittmer, U., ten Haaf, P., Heeney, J., Zou, W., Emilie, D., Sopper, S., Uberla, K., 2003. Replication, immunogenicity, and protective properties of live-attenuated simian immunodeficiency viruses expressing interleukin-4 or interferon-gamma. *Virology* 305, 473–485.
- Suryanarayana, K., Wiltrout, T.A., Vasquez, G.M., Hirsch, V.M., Lifson, J.D., 1998. Plasma SIV RNA viral load determination by real-time quantification of product generation in reverse transcriptase-polymerase chain reaction. *AIDS Res. Hum. Retrovir.* 14, 183–189.
- Tartaglia, L.A., Goeddel, D.V., Reynolds, C., Figari, I.S., Weber, R.F., Fendly, B.M., Palladino Jr., M.A., 1993. Stimulation of human T-cell proliferation by specific activation of the 75-kDa tumor necrosis factor receptor. *J. Immunol.* 151, 4637–4641.
- Ui, M., Kuwata, T., Igarashi, T., Ibuki, K., Miyazaki, Y., Kozyrev, I.L., Enose, Y., Shimada, T., Uesaka, H., Yamamoto, H., Miura, T., Hayami, M., 1999. Protection of macaques against a SHIV with a homologous HIV-1 Env and a pathogenic SHIV-89.6P with a heterologous Env by vaccination with multiple gene-deleted SHIVs. *Virology* 265, 252–263.
- von Sydow, M., Sonnerborg, A., Gaines, H., Strannegard, O., 1991. Interferon-alpha and tumor necrosis factor-alpha in serum of patients in various stages of HIV-1 infection. *AIDS Res. Hum. Retrovir.* 7, 375–380.
- Yokota, S., Geppert, T.D., Lipsky, P.E., 1988. Enhancement of antigen- and mitogen-induced human T lymphocyte proliferation by tumor necrosis factor-alpha. *J. Immunol.* 140, 531–536.
- Zheng, L., Fisher, G., Miller, R.E., Peschon, J., Lynch, D.H., Lenardo, M.J., 1995. Induction of apoptosis in mature T cells by tumour necrosis factor. *Nature* 377, 348–351.