

precipitation method, and the medium was changed every 2 days. At 14 days after transfection, the cells were stained with Giemsa solution and the foci that had formed were counted.

Rat 3Y1 cells were transfected with 5  $\mu$ g each of the above plasmids by the calcium phosphate precipitation method and cultured in the presence of 300  $\mu$ g/ml of G418. At 14 days after transfection, the cells were stained with Giemsa solution and the colonies that had formed were counted.

### Indirect immunofluorescence

H1299 cells were transfected with FLAG-DJ-1 or its mutant as described above. At 48 h after transfection, cells were fixed with a solution containing 4% paraformaldehyde and reacted with a combination of a mouse anti-FLAG monoclonal antibody (M2, Sigma), goat anti-GRP78 polyclonal antibody (N-20, Santa Cruz), and MitoTracker (Molecular Probe). The cells were then reacted with an FITC-conjugated anti-mouse IgG or rhodamine-conjugated anti-rabbit IgG and observed under a confocal laser fluorescent microscope.

### Isoelectric focusing

H1299 cells were irradiated with UV using a UV cross linker, and proteins were extracted from the cells at various times after UV-irradiation. Proteins were then separated in the pH 5–8 range of isoelectric focusing phoresis gel or 12.5% polyacrylamide gel containing SDS, transferred onto nitrocellulose membranes, and blotted with an anti-DJ-1 polyclonal antibody or antiactin-antibody (MAB1501R, CHEMICON) as described previously.<sup>41</sup>

### Establishment of a DJ-1-knockdown cell line

The nucleotide sequences of the upper strand of the oligonucleotide used for construction of an siRNA vector targeting mouse DJ-1 gene is 5'-GGATCCCGCCTTGCTAGTAGAATAAACTTCAAGAGAGTTTATTCTACTA GCAAGGTTTTTCCAAAAGCTT-3'. After annealing oligonucleotides corresponding to the upper and lower strands of DNA, these were inserted into *Bam*HI–*Hind*III sites of pRNA-U6.1/Neo (GenScript Corp.). These plasmids were transfected into mouse Flp-In™3T3 cells (Invitrogen) by the calcium phosphate precipitation method, and the cells were cultured in the medium in the presence of 200  $\mu$ g/ml G418 for 14 days. The cells that were resistant to the drug were then selected, and expression of DJ-1 was examined by Western blotting with an anti-mouse DJ-1 antibody.<sup>8</sup>

### Separation of soluble and insoluble forms of DJ-1

Human 293T cells were transfected with FLAG-DJ-1 or its mutant as described above. At 48 h after transfection, cells were lysed with a buffer containing 0.1% NP-40 and phosphate-buffered saline and by sonication. The supernatant fraction after centrifugation was used as soluble proteins, and the precipitates were further dissolved with a sample buffer containing SDS as described above and used as insoluble proteins. These proteins were blotted with an anti-FLAG antibody.

### Acknowledgements

We thank Yoko Misawa and Kiyomi Takaya for their technical assistance. This work was supported by grants-in-aid from the Ministry of Education, Science, Culture, Sports and Technology of Japan and the Ministry of Health, Labor and Welfare of Japan.

### References

- Nagakubo D, Taira T, Kitaura H, Ikeda M, Tamai K, Iguchi-Ariga SMM and Ariga H (1997) DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem. Biophys. Res. Commun.* 31: 509–513
- Taira T, Takahashi K, Kitagawa R, Iguchi-Ariga SMM and Ariga H (2001) Molecular cloning of human and mouse DJ-1 genes and identification of Sp1-dependent activation of the human DJ-1 promoter. *Gene* 263: 285–292
- Hod Y, Pentylala SN, Whyard TC and El-Maghrabi MR (1999) Identification and characterization of a novel protein that regulates RNA-protein interaction. *J. Cell. Biochem.* 72: 435–444
- Wagenfeld A, Yeung CH, Strupat K and Cooper TG (1998) Shedding of a rat epididymal sperm protein associated with infertility induced by ornidazole and alpha-chlorohydrin. *Biol. Reprod.* 8: 1257–1265
- Welch JE, Barbee RR, Roberts NL, Suarez JD and Klinefelter GR (1998) SP22: a novel fertility protein from a highly conserved gene family. *J. Androl.* 19: 385–393
- Wagenfeld A, Yeung CH, Shivaji S, Sundareswaran VR, Ariga H and Cooper TG (2000) Expression and cellular localization of contraception-associated protein. *J. Androl.* 21: 954–963
- Klinefelter GR, Welch JE, Perreault SD, Moore HD, Zucker RM, Suarez JD, Roberts NL, Bobseine K and Jeffay S (2002) Localization of the sperm protein SP22 and inhibition of fertility *in vivo* and *in vitro*. *J. Androl.* 23: 48–63
- Okada M, Matsumoto K, Niki T, Taira T, Iguchi-Ariga SMM and Ariga H (2002) DJ-1, a target protein for an endocrine disrupter, participates in the fertilization in mice. *Biol. Pharm. Bull.* 25: 853–856
- Yoshida K, Sato Y, Yoshiike M, Nozawa S, Ariga H and Iwamoto T (2003) Immunocytochemical localization of DJ-1 in human male reproductive tissue. *Mol. Reprod. Dev.* 66: 391–397
- Le Naour F, Misek DE, Krause MC, Deneux L, Giordano TJ, Scholl S and Hanash SM (2001) Proteomics-based identification of RS/DJ-1 as a novel circulating tumor antigen in breast cancer. *Clin. Cancer Res.* 7: 3328–3335
- Miura K, Bowman ED, Simon R, Peng AC, Robles AI, Jones RT, Katagiri T, He P, Mizukami H, Charboneau L, Kikuchi T, Liotta LA, Nakamura Y and Harris CC (2002) Laser capture microdissection and microarray expression analysis of lung adenocarcinoma reveals tobacco smoking- and prognosis-related molecular profiles. *Cancer Res.* 62: 3244–3250
- Takahashi K, Taira T, Niki T, Seino C, Iguchi-Ariga SMM and Ariga H (2001) DJ1 positively regulates the androgen receptor by impairing the binding of PIASx alpha to the receptor. *J. Biol. Chem.* 276: 37556–37563
- Niki T, Takahashi-Niki K, Taira T, Iguchi-Ariga SMM and Ariga H (2003) DJBP: A novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol. Cancer Res.* 1: 247–261
- Taira T, Iguchi-Ariga SMM and Ariga H (2004) Co-localization with DJ-1 is essential for the androgen receptor to exert its transcription activity that has been impaired by androgen-antagonists. *Biol. Pharm. Bull.* 27: 574–577
- Shinbo Y, Taira T, Niki T, Iguchi-Ariga SMM and Ariga H (2005) DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int. J. Oncol.* 26: 641–648
- Sekito A, Taira T, Niki T, Iguchi-Ariga SMM and Ariga H (2005) Stimulation of transforming activity of DJ-1 by Abstrakt, a DJ-1-binding protein. *Int. J. Oncol.* 26: 685–689
- Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC, Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A, Meco G, van Duijn CM, Oostra BA and Heutink P (2003) Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299: 256–259
- Abou-Sleiman PM, Healy DG, Quinn N, Lees AJ and Wood NW (2003) The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann. Neurol.* 54: 283–286
- Hague S, Rogaeva E, Hernandez D, Gulick C, Singleton A, Hanson M, Johnson J, Weiser R, Gallardo M, Ravina B, Gwinn-Hardy K, Crawley A, St George-Hyslop PH, Lang AE, Heutink P, Bonifati V, Hardy J and Singleton A (2003) Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann. Neurol.* 54: 271–274
- Hedrich K, Djarmati A, Schafer N, Hering R, Wellenbrock C, Weiss PH, Hilker R, Vieregge P, Ozelius LJ, Heutink P, Bonifati V, Schwinger E, Lang AE, Noth J, Bressman SB, Pramstaller PP, Riess O and Klein C (2004) DJ-1 (PARK7)

- mutations are less frequent than Parkin (PARK2) mutations in early-onset Parkinson disease. *Neurology* 62: 389–394
21. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R, Stenroos ES, Chandrasekharappa S, Athanassiadou A, Papapetropoulos T, Johnson WG, Lazzarini AM, Duvoisin RC, Di Iorio G, Golbe LI and Nussbaum RL (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276: 2045–2047
  22. Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, Yokochi M, Mizuno Y and Shimizu N (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392: 605–608
  23. Leroy E, Boyer R, Auburger G, Leube B, Ulm G, Mezey E, Harta G, Brownstein MJ, Jonnalagada S, Chernova T, Dehejia A, Lavedan C, Gasser T, Steinbach PJ, Wilkinson KD and Polymeropoulos MH (1998) The ubiquitin pathway in Parkinson's disease. *Nature* 395: 451–452
  24. Macedo MG, Anar B, Bronner IF, Cannella M, Squitieri F, Bonifati V, Hoogeveen A, Heutink P and Rizzo P (2003) The DJ-1L166P mutant protein associated with early onset Parkinson's disease is unstable and forms higher order protein complexes. *Hum. Mol. Genet.* 12: 2807–2816
  25. Miller DW, Ahmad R, Hague S, Baptista MJ, Canet-Aviles R, McLendon C, Carter DM, Zhu PP, Stadler J, Chandran J, Klinefelter GR, Blackstone C and Cookson MR (2003) L166P Mutant DJ-1, causative for recessive Parkinson's disease, is degraded through the ubiquitin-proteasome system. *J. Biol. Chem.* 278: 36588–36595
  26. Moore DJ, Zhang L, Dawson TM and Dawson VL (2003) A missense mutation (L166P) in DJ-1, linked to familial Parkinson's disease, confers reduced protein stability and impairs homo-oligomerization. *J. Neurochem.* 87: 1558–1567
  27. Gerner K, Holtorf E, Odoy S, Nuscher B, Yamamoto A, Regula JT, Beyer K, Haass C and Kahle PJ (2004) Differential effects of Parkinson's disease-associated mutations on stability and folding of DJ-1. *J. Biol. Chem.* 279: 6943–6951
  28. Olzmann JA, Brown K, Wilkinson KD, Rees HD, Huai Q, Ke H, Levey AI, Li L and Chin LS (2004) Familial Parkinson's disease-associated L166P mutation disrupts DJ-1 protein folding and function. *J. Biol. Chem.* 279: 8506–8515
  29. Bandyopadhyay R, Kingsbury AE, Cookson MR, Reid AR, Evans IM, Hope AD, Pittman AM, Lashley T, Canet-Aviles R, Miller DW, McLendon C, Strand C, Leonard AJ, Abou-Sleiman PM, Healy DG, Ariga H, Wood NW, de Silva R, Revesz T, Hardy JA and Lees AJ (2004) The expression of DJ-1 (PARK7) in normal human CNS and idiopathic Parkinson's disease. *Brain* 127: 420–430
  30. Rizzo P, Hinkle DA, Zhukareva V, Bonifati V, Severijnen LA, Martinez D, Ravid R, Kamphorst W, Eberwine JH, Lee VM, Trojanowski JQ and Heutink P (2004) DJ-1 colocalizes with tau inclusions: a link between parkinsonism and dementia. *Ann. Neurol.* 55: 113–118
  31. Neumann M, Muller V, Gerner K, Kretzschmar HA, Haass C and Kahle PJ (2004) Pathological properties of the Parkinson's disease-associated protein DJ-1 in alpha-synucleinopathies and tauopathies: relevance for multiple system atrophy and Pick's disease. *Acta Neuropathol.* 107: 489–496
  32. Heikkila R and Cohen G (1971) Inhibition of biogenic amine uptake by hydrogen peroxide: a mechanism for toxic effects of 6-hydroxydopamine. *Science* 172: 1257–1258
  33. Nicklas WJ, Youngster SK, Kindt MV and Heikkila RE (1987) MPTP, MPP+ and mitochondrial function. *Life Sci.* 40: 721–729
  34. Lotharius J and O'Malley KL (2000) The parkinsonism-inducing drug 1-methyl-4-phenylpyridinium triggers intracellular dopamine oxidation. A novel mechanism of toxicity. *J. Biol. Chem.* 275: 38581–38588
  35. Jenner P (2003) Oxidative stress in Parkinson's disease. *Ann. Neurol.* 53 (Suppl. 3): S26–S36
  36. Mitsumoto A, Nakagawa Y, Takeuchi A, Okawa K, Iwamatsu A and Takanezawa Y (2001) Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radical Res.* 35: 301–310
  37. Mitsumoto A and Nakagawa Y (2001) DJ-1 is an indicator for endogenous reactive oxygen species elicited by endotoxin. *Free Radical Res.* 35: 885–893
  38. Srisomsap C, Subhasitanont P, Otto A, Mueller EC, Punyart P, Wittmann-Liebold B and Svasti J (2002) Detection of cathepsin B up-regulation in neoplastic thyroid tissues by proteomic analysis. *Proteomics* 2: 706–712
  39. Kinumi T, Kimata J, Taira T, Ariga H and Niki E (2004) Cysteine-106 of DJ-1 is the most sensitive cysteine residue to hydrogen peroxide mediated oxidation *in vivo* in human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* 317: 722–728
  40. Yokota T, Sugawara K, Ito K, Takahashi R, Ariga H and Mizusawa H (2003) Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition. *Biochem. Biophys. Res. Commun.* 312: 1342–1348
  41. Taira T, Saito Y, Niki T, Iguchi-Ariga SMM, Takahashi K and Ariga H (2004) DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep.* 5: 213–218
  42. Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, Ali Z, Del Turco D, Bentivoglio AR, Healy DG, Albanese A, Nussbaum R, Gonzalez-Maldonado R, Deller T, Salvi S, Cortelli P, Gilks WP, Latchman DS, Harvey RJ, Dallapiccola B, Auburger G and Wood NW (2004) Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 304: 1158–1160
  43. Schmidt D and Muller S (2003) PIAS/SUMO: new partners in transcriptional regulation. *Cell Mol. Life Sci.* 60: 2561–2574
  44. Seeler JS and Dejean A (2003) Nuclear and unclear functions of SUMO. *Nat. Rev. Mol. Cell. Biol.* 4: 690–699
  45. Kahyo T, Nishida T and Yasuda H (2001) Involvement of PIAS1 in the sumoylation of tumor suppressor p53. *Mol. Cell* 8: 713–718
  46. Schmidt D and Muller S (2002) Members of the PIAS family act as SUMO ligases for c-Jun and p53 and repress p53 activity. *Proc. Natl. Acad. Sci. USA* 99: 2872–2877
  47. Nishida T and Yasuda H (2002) PIAS1 and PIASxalpha function as SUMO-E3 ligases toward androgen receptor and repress androgen receptor-dependent transcription. *J. Biol. Chem.* 277: 41311–41317
  48. Sachdev S, Bruhn L, Sieber H, Pichler A, Melchior F and Grosschedl R (2001) PIASy, a nuclear matrix-associated SUMO E3 ligase, represses LEF1 activity by sequestration into nuclear bodies. *Genes Dev.* 15: 3088–3103
  49. Takahashi-Niki K, Niki T, Taira T, Iguchi-Ariga SMM and Ariga H (2004) Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients. *Biochem. Biophys. Res Commun.* 320: 389–397
  50. Canet-Aviles RM, Wilson MA, Miller DW, Ahmad R, McLendon C, Bandyopadhyay S, Baptista MJ, Ringe D, Petsko GA and Cookson MR (2004) The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization. *Proc. Natl. Acad. Sci. USA* 101: 9103–9108
  51. Shendelman S, Jonason A, Martinat C, Leete T and Abeliovich A (2004) DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol.* 2: e362
  52. Martinat C, Shendelman S, Jonason A, Leete T, Beal MF, Yang L, Floss T and Abeliovich A (2004) Sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: an ES-derived cell model of primary parkinsonism. *PLoS Biol.* 2: e327
  53. Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS and Mak TW (2005) DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* 7: 263–273
  54. Xu J, Zhong N, Wang H, Elias JE, Kim CY, Woldman I, Pfl C, Gygi SP, Geula C and Yankner BA (2005) The Parkinson's disease-associated DJ-1 protein is a transcriptional co-activator that protects against neuronal apoptosis. *Hum. Mol. Genet.* 14: 1231–1241
  55. Pichler A, Gast A, Seeler JS, Dejean A and Melchior F (2002) The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* 108: 109–120
  56. Li Y, Wang H, Wang S, Quon D, Liu YW and Cordell B (2003) Positive and negative regulation of APP amyloidogenesis by sumoylation. *Proc. Natl. Acad. Sci. USA* 100: 259–264
  57. Honbou K, Suzuki NN, Horiuchi M, Niki T, Taira T, Ariga H and Inagaki F (2003) The crystal structure of DJ-1, a protein related to male fertility and Parkinson's disease. *J. Biol. Chem.* 278: 31380–31384
  58. Tao X and Tong L (2003) Crystal structure of human DJ-1, a protein associated with early-onset Parkinson's disease. *J. Biol. Chem.* 278: 31372–31379
  59. Wilson MA, Collins JL, Hod Y, Ringe D and Petsko GA (2003) The 1.1-A resolution crystal structure of DJ-1, the protein mutated in autosomal recessive early onset Parkinson's disease. *Proc. Natl. Acad. Sci. USA* 100: 9256–9261
  60. Huai Q, Sun Y, Wang H, Chin LS, Li L, Robinson H and Ke H (2003) Crystal structure of DJ-1/R5 and implication on familial Parkinson's disease. *FEBS Lett.* 549: 171–175
  61. Lee SJ, Kim SJ, Kim IK, Ko J, Jeong CS, Kim GH, Park C, Kang SO, Suh PG, Lee HS and Cha SS (2003) Crystal structures of human DJ-1 and *Escherichia*

- coli* Hsp31 that share an evolutionarily conserved domain. *J. Biol. Chem.* 278: 44552–44559
62. Orth M and Schapira AH (2002) Mitochondrial involvement in Parkinson's disease. *Neurochem. Int.* 40: 533–541
63. Chan HY, Warrick JM, Andriola I, Merry D and Bonini NM (2002) Genetic modulation of polyglutamine toxicity by protein conjugation pathways in *Drosophila*. *Hum. Mol. Genet.* 11: 2895–2904
64. Ueda H, Goto J, Hashida H, Lin X, Oyanagi K, Kawano H, Zoghbi HY, Kanazawa I and Okazawa H (2002) Enhanced SUMOylation in polyglutamine diseases. *Biochem. Biophys. Res. Commun.* 293: 307–313
65. Steffan JS, Agrawal N, Pallos J, Rockabrand E, Trotman LC, Slepko N, Illes K, Lukacovich T, Zhu YZ, Cattaneo E, Pandolfi PP, Thompson LM and Marsh JL (2004) SUMO modification of Huntingtin and Huntington's disease pathology. *Science* 304: 100–104

## DJ-1 interacts with HIPK1 and affects H<sub>2</sub>O<sub>2</sub>-induced cell death

AYA SEKITO<sup>1</sup>, SHIZUYO KOIDE-YOSHIDA<sup>1</sup>, TAKESHI NIKI<sup>2</sup>, TAKAHIRO TAIRA<sup>1,2</sup>,  
SANAE M.M. IGUCHI-ARIGA<sup>2,3</sup>, & HIROYOSHI ARIGA<sup>1,2</sup>

<sup>1</sup>Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan, <sup>2</sup>CREST, Japan Science and Technology Corporation, Saitama 332-0012, Japan, and <sup>3</sup>School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

Accepted by Professor E. Niki

(Received 11 August 2005; in revised form 26 October 2005)

### Abstract

DJ-1 is a novel oncogene and causative gene for the familial form of Parkinson's disease (PD). DJ-1 has multiple functions, including anti-oxidative stress by eliminating reactive oxygen species (ROS) and transcriptional regulation as a coactivator, and loss of these functions are thought to trigger the onset of PD. The mechanism underlying the prevention of cell death by DJ-1 is, however, not clear. In this study, we found that DJ-1 directly bound to homeodomain-interacting protein kinase 1 (HIPK1) *in vitro* and *in vivo* and that these proteins were colocalized in the nucleus. HIPK1 was then found to be degraded in human H1299 cells transfected with wild-type DJ-1 but not with a C106S DJ-1 mutant, a DJ-1 protein disrupting a catalytic domain of the putative protease, in a dose-dependent manner. Furthermore, although knockdown of either DJ-1 or HIPK1 rendered H1299 cells susceptible to H<sub>2</sub>O<sub>2</sub>-induced cell death, double-knockdown of DJ-1 and HIPK1 rendered H1299 cells resistant to H<sub>2</sub>O<sub>2</sub>-induced cell death, suggesting that the elevated level of HIPK1 induced by a low level of DJ-1 inhibits oxidative stress-induced cell death.

**Keywords:** *DJ-1, HIPK1, cell death, oxidative stress, Parkinson's disease*

### Introduction

DJ-1 was first identified by us as a novel oncogene that transforms mouse NIH3T3 cells in cooperation with activated ras [1]. Elevated expression of DJ-1 has been reported in cancer cells, including breast cancer, smoker-derived lung adenocarcinoma and prostate cancer [2–4]. DJ-1 was later found to participate in fertilization for sperm to penetrate into the zona pellucida of eggs [5–9]. DJ-1 was found to play a role in transcriptional regulation: a positive regulator of the androgen receptor [10–12] and p53 [13] and a negative regulator of PTEN tumor suppressor [14] and pyrimidine tract-binding protein-associated splicing factor (PSF) [15]. It has recently been shown that DJ-1 is responsible for onset of familial Parkinson's disease (PD), PARK7 [16], and 11 mutations in familial and sporadic forms of PD have been reported

[17–19]. DJ-1 has been shown to be expressed in almost all brain tissues in healthy men and patients with neurodegenerative diseases, including PD, Pick's disease and multiple system atrophy [20–22].

The pathology of PD involves an irreversible degeneration of the dopaminergic neuron. Various lines of evidence suggest that oxidative stresses lead to dopaminergic cell degeneration in PD [23–26]. Expression of DJ-1 has been shown to be induced by various reagents that induce reactive oxygen species (ROS) [27–30], and abnormal oxidative forms of DJ-1 have been found in some patients with sporadic forms of PD [20]. We previously reported that DJ-1 plays a role in the anti-oxidative stress reaction, in which ROS were scavenged by oxidizing DJ-1 itself, and that mutants of DJ-1, including various mutants found in PD patients, rendered cells susceptible to oxidative stress-induced cell death [31,32]. Other

Correspondence: H. Ariga, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita 12, Nishi 6, Kita-ku, Sapporo 060-0812, Japan. Tel: 81 11 707 3745. Fax: 81 11 706 4988. E-mail: hiro@pharm.hokudai.ac.jp

groups have also reported anti-oxidative activity of DJ-1 [33–35]. It has been shown that of three cysteines at amino acid numbers 46, 53 and 106 in DJ-1, C106 was first oxidized as a form of addition of SO<sub>3</sub>H or SO<sub>2</sub>H and then C46 and C53 were oxidized with increased amounts of H<sub>2</sub>O<sub>2</sub> added to cultured cells [30] and that C106 was indispensable for DJ-1 to exert its activity against oxidative stress [32–36]. In addition to oxidation of DJ-1, DJ-1 has been reported to be conjugated with SUMO-1 on a lysine residue at amino-acid number 130 (K130) [37]. It has also been reported that sumoylation of DJ-1 at K130 was essential for DJ-1 to exert its full activities, including activities for ras-dependent transformation, cell growth stimulation and anti-oxidative stress, and that improper sumoylation of L166P DJ-1, a DJ-1 mutant found in PD patients, occurred and resulted in loss of its function and formation of aggregation [37].

Oxidative stress is caused by ROS in cells in which large amounts of ROS are produced by alternative activities of scavenger proteins or by dysfunction of the mitochondrial respiratory chain pathway with reduction of complex I activity. We and other groups have shown that some DJ-1 is located in mitochondria in addition to the cytoplasm and nucleus [33,37,38] and that translocation of DJ-1 to mitochondria was stimulated by oxidative stress [39–42]. The functions of DJ-1 in mitochondria, however, are not clear. Recently, DJ-1 has been shown to interact with Daxx to inhibit apoptosis signal-regulating kinase 1 (ASK-1) activity and cell death by sequestering the death protein Daxx to the nucleus [43]. ASK-1 has been reported to be activated by Daxx that was translocated from the nucleus to cytoplasm after glucose deprivation or oxidative stress to cells, resulting in activation of the apoptosis pathway through Jun-N-terminal kinase (JNK) [44]. These findings suggest that there are at least two mechanisms by which DJ-1 acts against oxidative stress: one is elimination of ROS by oxidization of DJ-1 itself, and the other is activation of oxidative stress-induced apoptosis machinery.

In this study, we identified homeodomain-interacting protein kinase 1 (HIPK1) as a DJ-1-binding protein. HIPK1 has been shown to be involved in the apoptosis pathways of both Daxx-ASK-1 and p53 [45–49], and we have shown that DJ-1 bound to p53 to modulate its transcriptional activity [13]. We also showed that DJ-1 reduced the expression level of HIPK1 and affected the apoptosis activity of HIPK1.

## Materials and methods

### Cell culture

The human cell lines H1299 and 293T were cultured in Dulbecco's modified Eagle's medium supplemented with 10% calf serum.

### Construction of plasmids

Nucleotide sequences of the oligonucleotide used for PCR primers were as follows: hHIPK1 ATG, 5'-GAATTCATGGCATCACAGCTGCAA-3'; hHIPK1 sc, 5'-GGGAATCCCCACAGCAATGATTC-3'; hHIPK1 $\beta$  stop, 5'-GTCTGACTCAGCATCCAGCC-AGAAG-3'; hHIPK1 $\alpha$  stop, 5'-GGCTCGAGCTAT-AAGTAGGAATAC-3'; hHIPK2 ATG, 5'-GAATTC-ATGGCCTCACATGTGCAA-3'; hHIPK2 Bgl II antisense, 5'-CGACGCTTGACAGATCTCCATG TTC-3'; hHIPK2 Bgl II sense, 5'-CCAGAA CATGGAGATCTGCAAGCG-3'; hHIPK2 stop, 5'-CTCGAGTTATATGTAAGGGTACTG-3'.

Combinations of plasmids constructed, primers, templates and restriction enzyme sites and plasmids inserted are shown in Table I. PCR was carried out using the respective template and primers, and the resultant PCR product was digested with and inserted into the respective restriction enzyme site of the plasmid. Other plasmids used in this study were described previously [1,10,32].

### In vivo binding assay

Ten microgram of pcDNA3-Flag-HIPK1 $\alpha$  was transfected into human H1299 cells that were 60% confluent in a 10-cm dish by the calcium phosphate precipitation technique [49]. Forty-eight hours after transfection, the whole cell extract was prepared by the procedure described previously [10,32]. Approximately 2 mg of the H1299 cell proteins was first immunoprecipitated with a mouse anti-Flag antibody (M2, Sigma) under the same conditions as those described previously [10,32]. The precipitates were then separated in a 12% polyacrylamide gel containing SDS, blotted onto a nitrocellulose filter, and reacted with a rabbit anti-DJ-1 antibody [1] and with the mouse anti-Flag antibody. Immunoprecipitated proteins were then reacted with an HRP or IRDye800-conjugated second antibody and visualized by an ECL system or an infrared imaging system (Odyssey, LI-COR), respectively.

### In vitro binding assay

GST-DJ-1 or GST was expressed in and purified from *E. coli*. Purified GST-DJ-1 or GST was then incubated with 35S-HIPK1 $\alpha$  or 35S-HIPK1 $\beta$  synthesized *in vitro* using a reticulocyte lysate of a TnT-transcription-translation coupled system (Promega) and subjected to a pull-down assay using glutathione Sepharose as described previously [11,13]. The precipitates were separated in a 12% of polyacrylamide gel containing SDS and visualized by fluorography.

Table I. Plasmid construction.

Plasmid	Template	5'-primer	3'-primer	Restriction enzyme site	Plasmid inserted
pGADGHx-HIPK1 $\beta$	EST clone (Ac. BC033012)	hHIPK1 ATG	HIPK1 $\beta$ stop	EcoRI, XhoI	pGADGHx
pGADGHx-HIPK1 $\beta$ (sc)	EST clone (Ac. BC033012)	hHIPK1sc	HIPK1 $\beta$ stop	EcoRI, XhoI	pGADGHx
pGADGHx-HIPK1 $\alpha$ (sc)	EST clone (Ac. BC036057)	hHIPK1sc	HIPK1 $\alpha$ stop	EcoRI, XhoI	pGAD-GHx
pGADGHx-HIPK2	pcDNA-F-HIPK2	hHIPK2 ATG	hHIPK2 BgII anti	EcoRI, XhoI	pGAD-GHx
pGADGHx-HIPK1 $\alpha$	EST clone (Ac. BC033012)	hHIPK1 ATG	HIPK1 $\beta$ stop	EcoRI, XhoI	pGAD-GHx
pCMV-F-HIPK1 $\alpha$ (SC)	EST clone (Ac. BC036057)	hHIPK1sc	HIPK1 $\alpha$ stop	EcoRI, XhoI	PCMV-Flag
pCMV-F-HIPK1 $\alpha$	EST clone (Ac. BC033012)	hHIPK1 ATG	HIPK1 $\beta$ stop	EcoRI, XhoI	pCMV-Flag

Combinations of plasmids constructed, primers, templates and restriction enzyme sites and plasmids inserted were shown.

### RT-PCR

Total testis RNAs were prepared from human tissues (OriGene) or cultured cells by the acid guanidine thiocyanate-phenol-chloroform method, and cDNA was synthesized using the oligo dT primer and BcaBEST polymerase (Takara Co., Ltd). The first strand of cDNA products of HIPK1 $\alpha$  and HIPK1 $\beta$  was amplified with specific primers for the first 5 min at 94°C and then for 34–38 cycles of 1 min at 94°C, 1 min at 62°C and 1.5 min at 72°C. The first strand of cDNA products of DJ-1 was amplified with specific primers for the first 5 min at 94°C and then for 28 cycles of 1 min at 94°C, 2 min at 58°C and 3 min at 72°C.

Nucleotide sequences of the oligonucleotide used for PCR primers were as follows:

GAPDH-sense, 5'-TGAAGGTCCGAGTCAACG-GATTTGGT-3'; GAPDH-antisense, 5'-CATGTGGCCATGAGGTCCACCAC-3'; HIPK1-RT-sense, 5'-TCTTTGAGCAGCCCTTATTCCACT-3'; HIPK1-1-RT-antisense, 5'-GGGAGAAAAGATGAG-CAATGGAGC-3'; HIPK1-2-RT-antisense, 5'-GGG-GCTTCATTTGGCTTTATTTCAG-3'; Human DJ-1-sense, 5'-GGTGCAGGCTTGTAACATATAAC-3'; Human DJ-1-antisense, 5'-CTCTAAGTGATCGTCGACAGTTCGC-3'. The amplified products were separated on a 2% agarose gel and stained with ethidium bromide.

### Indirect immunofluorescence

H1299 cells were transfected with 5  $\mu$ g each of pcDNA3-DJ-1 and pEGFP-HIPK1 $\alpha$  by the calcium phosphate precipitation method [50]. Forty-eight hours after transfection, the cells were fixed with a solution containing 4% paraformaldehyde and reacted with a combination of a mouse anti-Flag-monoclonal antibody (M2, Sigma) and an anti-DJ-1 polyclonal antibody [1]. The cells were then reacted with a fluorescein isothiocyanate (FITC)-conjugated anti-rabbit IgG or rhodamine-conjugated anti-mouse IgG and observed under a confocal laser fluorescent microscope.

### Knockdown of genes by siRNA and cell viability assay

H1299 cells were cultured in a 96-well plate and transfected with 80 pmole/1.2  $\times$  10<sup>6</sup> cells of siRNAs targeting DJ-1 and HIPK1 genes using Lipofectamine 2000 (Invitrogen) according to the supplier's manual. At 24 h after transfection, the cells were treated with various amounts of hydrogen peroxide for 24 h and cell viability was then measured by an MTT assay using a cell counting kit-8 (DOJINDO). For RT-PCR and Western blot analysis, H1299 cells in a 6-cm dish were transfected with 80 pmole of siRNAs, and total RNAs and proteins were prepared at 72 and 48 h after transfection, respectively. The nucleotide

sequences for siRNAs targeting human DJ-1, human HIPK1 and luciferase genes were DJ-1: 5'-UGGAG-ACGGUCAUCCUGUdTdT-3' (upper strand) and 3'-dTdTACCUCUGCCAGUAGGGACA-5' (lower strand) HIPK1: 5'-GUUUCAACUGGACUACAA-GCAACA-AG-3' (upper strand) and 3'-UACAAA-GUUUGACCUGAUGUUCGUUGU-5' (lower strand), and luciferase: 5'-CGUACGCGGAAUA-CUUCGAdTdT-3' (upper strand) and 3'-dTdTGC-AUGCGCCUUAUGAAGCU-5' (lower strand). The siRNAs were synthesized by Greiner, Japan.

*Two-hybrid assay*

For a two-hybrid assay in yeast, wild-type DJ-1 and mutants of DJ-1 were inserted into the pGLex or pGAD-GHx vector and subjected to a two-hybrid assay as described previously [10].

**Results and discussion**

*Identification of HIPK1 as a Dj-1-binding protein*

We have reported that DJ-1 binds to PIASx $\alpha$  and DJBP and that it regulates AR transcription activity [10,11]. In addition to PIASx $\alpha$  and DJBP, cDNA spanning amino

acids 776–1210 of human homeodomain-interacting protein kinase 1 (HIPK1) was obtained by a yeast twohybrid screening. Mouse HIPK1 has been identified to be a protein interacting with the homeoproteins NKx-1.2 and NK-1 to act as a transcriptional cofactor (repressor) [51]. After a search of genomic and cDNA databases, we found two splicing isoforms of human HIPK1 and tentatively named them HIPK1 $\alpha$  and HIPK1 $\beta$ . The human HIPK1 gene comprises 16 exons, and HIPK1 $\alpha$  and HIPK1 $\beta$  are comprised of 1210 and 1075 amino acids corresponding to 1–16 and 1–15 exons, respectively (Figure 1A). After a nucleotide database search, EST clones of accession Nos. BC036057 and BC033012 were found to have the entire sequences of human HIPK1 $\alpha$  and HIPK1 $\beta$ , respectively. HIPK1 $\alpha$  and HIPK1 $\beta$  are composed of several domains, including a protein kinase domain, homeoprotein-interaction domain and PEST domain (Figure 1A) [51]. Since there is no report on human HIPK1 $\beta$ , the expressions of HIPK1 mRNAs were first examined by RT-PCR using various RNAs extracted from human tissues (Figure 1B and C). The results showed that human HIPK1 $\alpha$  mRNA, like human DJ-1 mRNA, was expressed ubiquitously in all of the tissues tested. HIPK1 $\beta$  was, on the other hand, found to be strongly expressed in the testis (Figure 1C).




Figure 1. Expression of HIPK1 mRNA in human tissues. (A) Structures of HIPK1 $\alpha$  and HIPK1 $\beta$  are shown. "YH" indicates the domain rich in tyrosine and histidine. (B) Positions of primers used for RT-PCR to amplify DNAs of HIPK1 $\alpha$  and HIPK1 $\beta$  were shown. Numbers written in Greece letters indicate the numbers of corresponding exons. Sizes of PCR products corresponding to HIPK1 $\alpha$  and HIPK1 $\beta$  to be amplified are 562 and 507 base pairs, respectively. (C) Human total RNAs were obtained from OriGene Technologies, and the expressions of HIPK1 $\alpha$  and HIPK1 $\beta$  were analyzed by RT-PCR with specific primers for each HIPK1 isoform, DJ-1 and GAPDH on total RNA as substrates. The amplified DNAs were separated in 2% agarose gels and visualized under UV light.




Figure 2. Associations of DJ-1 with HIPK1 $\alpha$  and HIPK1 $\beta$ . (A) Interactions between DJ-1 with HIPK1 $\alpha$  and HIPK1 $\beta$  were analyzed in a yeast twohybrid assay using a DJ-1 fusion protein with the LexA DNA-binding domain as bait and HIPK1-fusion proteins with the GAL4-activation domain as prey as described in "Materials and methods". Results obtained by using fusion proteins of the full-sized and C-terminal region of Daxx, Daxx and Daxx (c), respectively, with the LexA DNA-binding domain and the LexA DNA-binding domain alone as bait are also shown. HIPK1 $\alpha$ (sc) and HIPK1 $\beta$ (sc) indicate the regions of HIPK1 $\alpha$  and HIPK1 $\beta$  obtained by the screening. (B) Structures of HIPK1 $\alpha$  and HIPK2 are schematically shown, and identity of amino acids in each domain is also shown. (C) Interactions of DJ-1 with HIPK1 $\alpha$  and HIPK2 were analyzed in a yeast two-hybrid assay using a DJ-1 fusion protein with the LexA DNA-binding domain as bait and HIPK-fusion proteins with the GAL4-activation domain as prey as described in A. Results obtained by using fusion proteins of DJ-1 with the GAL4-activation domain and with the LexA DNA-binding domain as bait are also shown.

#### Associations of Dj-1 with HIPK1s

Associations of DJ-1 with HIPK1s were then examined by using a yeast two-hybrid assay, where DJ-1 and HIPK1s were fused to the LexA DNA-binding domain and GAL4 activation domain, respectively (Figure 2A). Since the C-terminal domain of a death domain-associated protein (Daxx) has been reported to associate with HIPK1 [45,49] and DJ-1 made a homodimer, the full-sized and C-terminal region of Daxx and DJ-1 were used as positive controls. While the C-terminal fragments of HIPK1 $\alpha$  and HIPK1 $\beta$  (HIPK1 $\alpha$ (sc) and HIPK1 $\beta$ (sc)), which were obtained in a screening of DJ-1-binding proteins, were found to strongly bind to both DJ-1 and the C-terminal region of Daxx (Daxx(C)), full-sized HIPK1 $\alpha$  but not HIPK1 $\beta$  was found to bind to DJ-1. Binding activities of DJ-1,

full-sized HIPK1 $\alpha$  and HIPK1 $\beta$  to Daxx were not clear in this two-hybrid assay. HIPK1 is a member of the HIPK family of proteins comprised of HIPK1, HIPK2 and HIPK3. Since HIPK2, whose amino acid composition is 62.9% homologous to that of HIPK1 $\alpha$  (Figure 2B), has been well characterized, interaction of DJ-1 with HIPK2 was also examined by the yeast two-hybrid assay (Figure 2C). The results showed that DJ-1 bound to full-sized HIPK1 $\alpha$  but not HIPK2. We therefore concentrated on HIPK1 $\alpha$  in this study.

Association of DJ-1 with HIPK1 $\alpha$  in cells was then examined. To do this, Flag-tagged HIPK1 $\alpha$  was transfected into human H1299 cells. Forty-eight hours after transfection, cell extracts were prepared from transfected cells, proteins in the extracts were immunoprecipitated with an anti-DJ-1 antibody or nonspecific IgG, and the precipitates were analyzed by Western



blotting with anti-Flag or anti-DJ-1 antibody (Figure 3A). The anti-DJ-1 antibody, but not IgG, was first confirmed to precipitate endogenously expressed DJ-1 (Figure 3A, lower part). The results also showed that Flag-HIPK1 $\alpha$  was co-immunoprecipitated with DJ-1 (Figure 3A, upper part), indicating

that DJ-1 is associated with HIPK1 $\alpha$  in H1299 cells. Association of DJ-1 with Flag-HIPK1 $\alpha$  was also observed in human 293T cells (data not shown). *In vitro* binding assays were then carried out using <sup>35</sup>S-labeled HIPK1 $\alpha$ (sc) that had been synthesized in a reticulocyte lysate and GST-DJ-1 or GST that had been




Figure 3. Interaction and colocalization of DJ-1 with HIPK1 $\alpha$ , a splicing variant of HIPK1. (A) Human H1299 cells were transfected with an expression vector for Flag-HIPK1 $\alpha$ . Forty-eight hours after transfection, whole cell extracts were prepared and then immunoprecipitated with an anti-DJ-1 antibody or non-specific IgG. Immunoprecipitates were then analyzed by Western blotting with an anti-Flag or anti-DJ-1 antibody. (B) GST or GST-DJ-1 was expressed in *E. coli* BL21(DE3) and applied to glutathione-Sepharose 4B. <sup>35</sup>S-HIPK1 $\alpha$ (sc) synthesized *in vitro* in a coupled transcription/translation system was then applied to the column. The labeled proteins that had bound to the column were separated in a gel and visualized by fluorography. 1/50 volumes of the labeled S-HIPK1 $\alpha$ (sc) used for the binding reaction were applied to the same gel (lane 2 in the upper part of figure). GST and GST-DJ-1 used in this experiment were separated on the gel and stained with coomassie brilliant blue (lower part). (C) H1299 cells were transfected with expression vectors for Flag-DJ-1 with or without GFP-HIPK1 $\alpha$  by the calcium phosphate precipitation technique. Forty-eight hours after transfection, the cells were fixed, reacted with an anti-Flag monoclonal antibody (M2, Sigma), and visualized with a rhodamine-conjugated anti-mouse IgG. GFP-HIPK1 $\alpha$  was visualized by self-fluorescence of GFP. The two figures have been merged.




Figure 4. Degradation of HIPK1 by DJ-1. H1299 cells in a 10-cm dish were transfected with 15  $\mu$ g of pcDNA3-Flag-HIPK1 $\alpha$ , 2.5  $\mu$ g of pEGFP-C1 and various amounts of pcDNA3-HA-DJ-1 or pcDNA3-HA-C106S DJ-1 by the calcium phosphate precipitation technique. Forty-eight hours after transfection, proteins in the cell extracts were subjected to Western blotting with anti-HA (MBL), anti-Flag antibodies (M2, Sigma), anti-GFP (MBL) and anti-actin (Chemicon) antibodies. The amounts of transfected pcDNA3-HA-DJ-1 or pcDNA3-HA-C106S DJ-1 were 1.25  $\mu$ g (lanes 4 and 7), 2.5  $\mu$ g (lanes 5 and 8) and 6.25  $\mu$ g (lanes 6 and 9).

expressed in and purified from *E. coli* (Figure 3B). The results showed that GST-DJ-1 bound to HIPK1 $\alpha$ (sc), suggesting that DJ-1 directly binds to HIPK1 $\alpha$ .

DJ-1 has been shown to be localized both in the cytoplasm and nucleus in human HeLa cells and is translocated from the cytoplasm to nucleus during the S-phase of the cell cycle upon induction by mitogen [1]. Recent findings have also shown that some DJ-1 is localized in mitochondria [33,37,40] and that mitochondrial localization of DJ-1 is stimulated by oxidative stress [39,41,42]. To determine the cellular localization of HIPK1 $\alpha$  and DJ-1, an expression vector for GFP-tagged HIPK1 $\alpha$  was cotransfected with Flag-tagged DJ-1 into human H1299 cells. Two days after transfection, the cells were stained with anti-Flag antibody, and Flag-DJ-1 and GFP- HIPK1 $\alpha$  were detected by a rhodamine-conjugated second antibody and self-fluorescence of GFP, respectively, and then visualized under a confocal laser microscope (Figure 2C). As previously described [47,49], HIPK1 $\alpha$  (green) was localized in the nucleus as speckles, and HIPK1 $\alpha$  and DJ-1 (red) were found to be colocalized in the nucleus as shown by the yellow color (Figure 3C, Merge).

#### Degradation of HIPK1 by Dj-1

During this study, we often observed that the expression of ectopically transfected HIPK1 $\alpha$  was not detected by Western blotting when DJ-1 was also transfected into various cell types. Since structural and biochemical analyses of DJ-1 suggest that DJ-1 is a protease [52,53], though controversial results have also been reported [54–56], various amounts of an expression vector for HA-tagged wild-type DJ-1 or C106S mutant DJ-1 were cotransfected with a constant amount of Flag-HIPK1 $\alpha$  into H1299 cells. Forty-eight hours after transfection, the expression levels of Flag-HIPK1 $\alpha$  and HA-DJ-1 were examined by Western blotting with anti-Flag and anti-HA antibodies (Figure 4). Cysteine at amino acid number

106 of DJ-1 (C106) is not only a putative catalytic center for protease activity of DJ-1 [52] but also an amino acid to be oxidized after oxidative stress [30,33]. C106S mutant DJ-1, a mutant in which there is substitution from cysteine to serine, has been shown to lose activity of DJ-1 against oxidative stress [32–35]. pCMV-EGFP was also cotransfected with DJ-1 and HIPK1 $\alpha$  to normalize transfection efficiencies. The results showed that the expression level of Flag-HIPK1 $\alpha$  was decreased by transfected HA-wild-type DJ-1, but not by C106S mutant DJ-1, in a dose-dependent manner. Since RT-PCR analysis showed that the expression level of mRNA of transfected Flag-HIPK1 $\alpha$  was not changed after cotransfection of HA-DJ-1 (data not shown), these results suggest that DJ-1 directly or indirectly degrades HIPK1 $\alpha$ .

#### Effect of reduced expression levels of Dj-1 and HIPK1 on H<sub>2</sub>O<sub>2</sub>-induced cell death

We and others have shown that DJ-1 protects cells against oxidative stress-induced cell death [31–35]. Pro- and anti-apoptotic functions of HIPK1 have, on the other hand, been reported. When cells were exposed to reagents that damage DNA, nuclear-localized HIPK1 phosphorylated p53 tumor suppressor, leading to activation of p53 transcription activity to trigger apoptosis [45]. When cells were deprived of glucose or receive oxidative stress, HIPK1 phosphorylated nucleus-localized Daxx, and phosphorylated Daxx was then translocated to the cytoplasm to associate with and activate an (ASK-1), resulting in activation of an apoptosis signal [45,49]. Alternatively, after stimulation of apoptosis by tumor necrosis factor (TNF) alpha, sumoylated HIPK1 in the nucleus was desumoylated and translocated to the cytoplasm to associate with ASK1-interacting protein (AIP1), a signal transducer in TNFalpha-induced ASK1 activation, resulting in an apoptosis signal [48]. HIPK1 (–/–) cells that had been established from

HIPK1-knockout mice were, however, more susceptible than HIPK1 (+/+) cells to apoptosis induced by DNA damage, suggesting that HIPK1 has an anti-apoptotic function [46].

To investigate roles of DJ-1 and HIPK1 in H<sub>2</sub>O<sub>2</sub>-induced apoptosis, H1299 cells were transfected with siRNAs targeting DJ-1 and HIPK1 genes (siDJ-1 RNA and siHIPK1 RNA). Twenty-four hours after transfection, the cells were treated with various concentrations of H<sub>2</sub>O<sub>2</sub> for 24 h and cell viability was then measured by an MTT assay (Figure 5). The expression levels of HIPK1 and DJ-1 were first

examined by RT-PCR, and less than 10% and about 30% of expression levels of DJ-1 and HIPK1 mRNAs, respectively, compared to the levels in cells transfected with siRNA targeting the luciferase gene, were confirmed (Figure 5A). As previously described in human SH-SY5Y and mouse NIH3T3 cells, in which the p53 gene is expressed [32], H1299 cells, in which the p53 gene is not expressed, transfected with siDJ-1 RNA were found to be rendered more susceptible to H<sub>2</sub>O<sub>2</sub>-induced cell death (Figure 5B), suggesting that DJ-1 protects cells from H<sub>2</sub>O<sub>2</sub>-induced cell death in a p53-independent manner. H1299 cells




Figure 5. Effect of HIPK1 on H<sub>2</sub>O<sub>2</sub>-induced cell death. (A) H1299 cells were transfected with siRNAs targeting DJ-1 and HIPK1 genes alone or together using Lipofectamine 2000. At 72 h after transfection, total RNAs were extracted and the expression levels of mRNAs of DJ-1, HIPK1 and GAPDH were examined by RTPCR as described in Materials and methods. (B) H1299 cells were transfected with siRNAs targeting DJ-1 and HIPK1 genes alone or together using Lipofectamine 2000. At 24 h after transfection, cells were treated with various amounts of H<sub>2</sub>O<sub>2</sub> for 24 h and cell viability was then measured by an MTT assay using a cell counting kit-8. The amounts of H<sub>2</sub>O<sub>2</sub> used were 5, 10, 25, 50, 100, 250, 500, 750 and 1000 μM. C H1299 cells in a 6-cm dish were transfected with 3 μg each of pcDNA3-Flag-HIPK1α, pcDNA3-HA-DJ-1 and pEGFP-C1 together with 80 pmol of siRNAs targeting DJ-1 and HIPK1 genes alone or together using Lipofectamine 2000. At 48 h after transfection, proteins in the cell extracts were subjected to Western blotting with anti-HA (MBL), anti-Flag antibodies (M2, Sigma) and anti-GFP (MBL) antibodies.

transfected with siHIPK1 RNA were found to be susceptible to H<sub>2</sub>O<sub>2</sub>-induced cell death, but the extent of susceptibility was less than that after transfection of siDJ-1 RNA into cells, suggesting that HIPK1 also plays a role in anti-oxidative stress and that weak susceptibility of siHIPK1-transfected cells to H<sub>2</sub>O<sub>2</sub>-induced cell death is affected by low knock-down efficiency of siHIPK1 RNA. H1299 cotransfected with siDJ-1 and siHIPK1 RNAs were, however, found to be more resistant to H<sub>2</sub>O<sub>2</sub>-induced cell death. Since DJ-1 degrades HIPK1 $\alpha$  as shown in Figure 4, it is thought that the elevated level of HIPK1 induced by DJ-1-knockdown protected H1299 cells from H<sub>2</sub>O<sub>2</sub>-induced cell death. We tried several times to make antibodies against HIPK1 using GST-HIPK1 or a synthetic peptide corresponding to the amino acids of HIPK1 as immunogens, but we did not succeed in establishing an anti-HIPK1 antibody that detects endogenous HIPK1. Commercially available anti-HIPK1 antibodies also did not work. To evaluate the effect of DJ-1-knockdown on the amount of HIPK1, therefore, Flag-tagged HIPK1 $\alpha$ , HA-tagged DJ-1 and GFP were cotransfected with various combinations of siRNAs targeting HIPK1 and DJ-1. Forty-eight hours after transfection, cell extracts were prepared and proteins in the extracts were analyzed by Western blotting with anti-Flag and HA antibodies (Figure 5C). GFP was used as an internal control. As shown in the levels of endogenous mRNAs, the expression levels of Flag-HIPK1 $\alpha$  and DJ-1-HA were reduced by transfection of respective siRNA alone (Figure 5C, lanes 2 and 3). When both siHIPK1 and siDJ-1 RNAs were transfected together, the expression level of DJ-1 was decreased but the extent of decrease was more than that after transfection of siDJ-1 RNA alone into cells, and the expression level of HIPK1 $\alpha$  was increased, suggesting that degradation of HIPK1 $\alpha$  was inhibited by DJ-1-knockdown (Figure 5C, lane 4). The results also suggest that HIPK1 protects H1299 cells against H<sub>2</sub>O<sub>2</sub>-induced cell death when the expression level of DJ-1 is extremely low stronger than when the expression level of DJ-1 is high. It is thought that both DJ-1 and HIPK1 play a role in anti-oxidative stress in H1299 cells under this condition and that the balance of the expression levels of DJ-1 and HIPK1 determine susceptibility to H<sub>2</sub>O<sub>2</sub>-induced cells death.

In this study, we identified HIPK1 as a DJ-1-binding-protein and found that HIPK1 was degraded by wild-type DJ-1 but not by C106S mutant DJ-1, a mutant lacking a putative catalytic amino acid for protease. Since protease activity of DJ-1 with C106 as a catalytic amino acid has been shown by structural and biochemical analyses [52,53], though controversial results have also been reported [54–56], it is likely that HIPK1 was degraded by DJ-1 directly or indirectly via other proteases that had been activated by DJ-1. We further found that HIPK1, like DJ-1,

prevented H1299 cells from H<sub>2</sub>O<sub>2</sub>-induced cells death and that knockdown of DJ-1 expression by siRNA resulted in enhancement of its protective activity, due to the elevated expression level of HIPK1. As described above, pro- and anti-apoptotic functions of HIPK1 have been reported [45–51]. These activities have been examined in various cell types that had been deprived of glucose or treated with TNF $\alpha$  and DNAdamaging reagents [45–51]. Addition of H<sub>2</sub>O<sub>2</sub> to cells has not been reported. These results suggest that pro- and anti-apoptotic functions of HIPK1 are determined in cell type- and inducer of cell death stimuli-dependent manners.

A number of studies have shown activity of DJ-1 against oxidative stress-induced cell death using various cell types, and at least three mechanisms underlying the anti-oxidative stress function of DJ-1 have been proposed: DJ-1 scavenges oxidative stress-induced ROS [32], DJ-1 regulates expressions of genes encoding proteins that play a role in preventing cell death [15,57], and interaction of DJ-1 with Daxx sequesters Daxx to the nucleus in order to leave ASK-1 in an inactive form [43]. On the other hand, it has been reported that Daxx phosphorylated by HIPK1 was translocated to the cytoplasm to activate ASK-1, leading to activation of the apoptosis pathway [49]. It is thought that when the high expression level of DJ-1 degrades HIPK1, Daxx is not phosphorylated to be sequestered to the nucleus, resulting in inactivation of the apoptosis pathway. DJ-1, therefore, plays a role in anti-oxidative stress by using various pathways.

#### Acknowledgements

We thank Yoko Misawa and Kiyomi Takaya for their technical assistance. This work was supported by grants-in-aid from the Ministry of Education, Science, Culture, Sports and Technology of Japan and the Ministry of Health, Labor and Welfare of Japan.

#### References

- [1] Nagakubo D, Taira T, Kitaura H, Ikeda M, Tamai K, Iguchi-Arigo SMM, Ariga H. DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem Biophys Res Commun* 1997;231:509–513.
- [2] Le Naour F, Misek DE, Krause MC, Deneux L, Giordano TJ, Scholl S, Hanash SM. Proteomics-based identification of RS/DJ-1 as a novel circulating tumor antigen in breast cancer. *Clin Cancer Res* 2001;7:3328–3335.
- [3] Miura K, Bowman ED, Simon R, Peng AC, Robles AI, Jones RT, Katagiri T, He P, Mizukami H, Charboneau L, Kikuchi T, Liotta LA, Nakamura Y, Harris CC. Laser capture microdissection and microarray expression analysis of lung adenocarcinoma reveals tobacco smoking- and prognosis-related molecular profiles. *Cancer Res* 2002;62:3244–3250.
- [4] Grzmil M, Voigt S, Thelen P, Hemmerlein B, Helmke K, Burfeind P. Up-regulated expression of the MAT-8 gene in prostate cancer and its siRNA-mediated inhibition of

- expression induces a decrease in proliferation of human prostate carcinoma cells. *Int J Oncol* 2004;24:97–105.
- [5] Wagenfeld A, Yeung CH, Strupat K, Cooper TG. Shedding of a rat epididymal sperm protein associated with infertility induced by ornidazole and alpha-chlorohydrin. *Biol Reprod* 1998;8:1257–1265.
  - [6] Welch JE, Barbee RR, Roberts NL, Suarez JD, Klinefelter GR. SP22: A novel fertility protein from a highly conserved gene family. *J Androl* 1998;19:385–393.
  - [7] Wagenfeld A, Yeung CH, Shivaji S, Sundareswaran VR, Ariga H, Cooper TG. Expression and cellular localization of contraception-associated protein. *J Androl* 2000;21:954–963.
  - [8] Klinefelter GR, Welch JE, Perreault SD, Moore HD, Zucker RM, Suarez JD, Roberts NL, Bobseine K, Jeffay S. Localization of the sperm protein SP22 and inhibition of fertility *in vivo* and *in vitro*. *J Androl* 2002;23:48–63.
  - [9] Okada M, Matsumoto K, Niki T, Taira T, Iguchi-Ariga SMM, Ariga H. DJ-1, a target protein for an endocrine disrupter, participates in the fertilization in mice. *Biol Pharm Bull* 2002;25:853–856.
  - [10] Takahashi K, Taira T, Niki T, Seino C, Iguchi-Ariga SMM, Ariga H. DJ-1 positively regulates the androgen receptor by impairing the binding of PIAS $\alpha$  to the receptor. *J Biol Chem* 2001;276:37556–37563.
  - [11] Niki T, Takahashi-Niki K, Taira T, Iguchi-Ariga SMM, Ariga H. DJBP: A Novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol Cancer Res* 2003;1:247–261.
  - [12] Taira T, Iguchi-Ariga SMM, Ariga H. Co-localization with DJ-1 is essential for the androgen receptor to exert its transcription activity that has been impaired by androgen antagonists. *Biol Pharm Bull* 2004;27:574–577.
  - [13] Shinbo Y, Taira T, Niki T, Iguchi-Ariga SMM, Ariga H. DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int J Oncol* 2005;26:641–648.
  - [14] Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS, Mak TW. DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* 2005;7:263–273.
  - [15] Xu J, Zhong N, Wang H, Elias JE, Kim CY, Woldman I, Pifl C, Gygi SP, Geula C, Yankner BA. The Parkinson's disease-associated DJ-1 protein is a transcriptional coactivator that protects against neuronal apoptosis. *Hum Mol Genet* 2005;14:1231–1241.
  - [16] Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC, Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A, Meco G, van Duijn CM, Oostra BA, Heutink P. Mutations in the DJ-1 gene associated with autosomal recessive early-onset Parkinsonism. *Science* 2003;299:256–259.
  - [17] Abou-Sleiman PM, Healy DG, Quinn N, Lees AJ, Wood NW. The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann Neurol* 2003;54:283–286.
  - [18] Hague S, Rogaeva E, Hernandez D, Gulick C, Singleton A, Hanson M, Johnson J, Weiser R, Gallardo M, Ravina B, Gwinn-Hardy K, Crawley A, St George-Hyslop PH, Lang AE, Heutink P, Bonifati V, Hardy J, Singleton A. Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann Neurol* 2003;54:271–274.
  - [19] Hedrich K, Djarmati A, Schafer N, Hering R, Wellenbrock C, Weiss PH, Hilker R, Vieregge P, Ozelius LJ, Heutink P, Bonifati V, Schwinger E, Lang AE, Noth J, Bressman SB, Pramstaller PP, Riess O, Klein C. DJ-1 (PARK7) mutations are less frequent than Parkin (PARK2) mutations in early-onset Parkinson disease. *Neurology* 2004;62:389–394.
  - [20] Bandopadhyay R, Kingsbury AE, Cookson MR, Reid AR, Evans IM, Hope AD, Pittman AM, Lashley T, Canet-Aviles R, Miller DW, McLendon C, Strand C, Leonard AJ, Abou-Sleiman PM, Healy DG, Ariga H, Wood NW, de Silva R, Revesz T, Hardy JA, Lees AJ. The expression of DJ-1 (PARK7) in normal human CNS and idiopathic Parkinson's disease. *Brain* 2004;127:420–430.
  - [21] Rizzu P, Hinkle DA, Zhukareva V, Bonifati V, Severijnen LA, Martinez D, Ravid R, Kamphorst W, Eberwine JH, Lee VM, Trojanowski JQ, Heutink P. DJ-1 colocalizes with tau inclusions: A link between Parkinsonism and dementia. *Ann Neurol* 2004;55:113–118.
  - [22] Neumann M, Muller V, Gorner K, Kretschmar HA, Haass C, Kahle PJ. Pathological properties of the Parkinson's disease-associated protein DJ-1 in alpha-synucleinopathies and tauopathies: Relevance for multiple system atrophy and Pick's disease. *Acta Neuropathol* 2004;107:489–496.
  - [23] Heikkila R, Cohen G. Inhibition of biogenic amine uptake by hydrogen peroxide: A mechanism for toxic effects of 6-hydroxydopamine. *Science* 1971;172:1257–1258.
  - [24] Nicklas WJ, Youngster SK, Kindt MV, Heikkila RE. MPTP, MPP+ and mitochondrial function. *Life Sci* 1987;40:721–729.
  - [25] Lotharius J, O'Malley KL. The Parkinsonism-inducing drug 1-methyl-4-phenylpyridinium triggers intracellular dopamine oxidation. A novel mechanism of toxicity. *J Biol Chem* 2000;275:38581–38588.
  - [26] Jenner P. Oxidative stress in Parkinson's disease. *Ann Neurol* 2003;53(3):S26–S36.
  - [27] Mitsumoto A, Nakagawa Y, Takeuchi A, Okawa K, Iwamatsu A, Takanezawa Y. Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radic Res* 2001;35:301–310.
  - [28] Mitsumoto A, Nakagawa Y. DJ-1 is an indicator for endogenous reactive oxygen species elicited by endotoxin. *Free Radic Res* 2001;35:885–893.
  - [29] Srisomsap C, Subhasitanont P, Otto A, Mueller EC, Punyarit P, Wittmann-Liebold B, Svasti J. Detection of cathepsin B up-regulation in neoplastic thyroid tissues by proteomic analysis. *Proteomics* 2002;2:706–712.
  - [30] Kinumi T, Kimata J, Taira T, Ariga H, Niki E. Cysteine-106 of DJ-1 is the most sensitive cysteine residue to hydrogen peroxide mediated oxidation *in vivo* in human umbilical vein endothelial cells. *Biochem Biophys Res Commun* 2004;317:722–728.
  - [31] Yokota T, Sugawara K, Ito K, Takahashi R, Ariga H, Mizusawa H. Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition. *Biochem Biophys Res Commun* 2003;312:1342–1348.
  - [32] Taira T, Saito Y, Niki T, Iguchi-Ariga SMM, Takahashi K, Ariga H. DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep* 2004;5:213–218.
  - [33] Canet-Aviles RM, Wilson MA, Miller DW, Ahmad R, McLendon C, Bandyopadhyay S, Baptista MJ, Ringe D, Petsko GA, Cookson MR. The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization. *Proc Natl Acad Sci USA* 2004;101:9103–9108.
  - [34] Shendelman S, Jonason A, Martinat C, Leete T, Abeliovich A. DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol* 2004;2:e362.
  - [35] Martinat C, Shendelman S, Jonason A, Leete T, Beal MF, Yang L, Floss T, Abeliovich A. Sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: An ES-derived cell model of primary Parkinsonism. *PLoS Biol* 2004;2:e327.
  - [36] Takahashi-Niki K, Niki T, Taira T, Iguchi-Ariga SMM, Ariga H. Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients. *Biochem Biophys Res Commun* 2004;320:389–397.
  - [37] Shinbo Y, Niki T, Taira T, Ooe H, Takahashi-Niki K, Maita C, Seino C, Iguchi-Ariga SMM, Ariga H. Proper SUMO-1

- conjugation is essential to DJ-1 to exert its full activities *Cell Death Diff* 2005., in press, available on line.
- [38] Zhang L, Shimoji M, Thomas B, Moore DJ, Yu S-W, Marupudi NI, Torp R, Torgner IA, Ottersen OP, Dawson TM, Dawson VL. Mitochondrial localization of the Parkinson's disease related protein DJ-1: Implications for pathogenesis. *Hum Mol Genet* 2005;14:2063-2073.
- [39] Jin J, Meredith GE, Chen L, Zhou Y, Xu J, Shie FS, Lockhart P, Zhang J. Quantitative proteomic analysis of mitochondrial proteins: Relevance to Lewy body formation and Parkinson's disease. *Mol Brain Res* 2005;134:119-138.
- [40] Blackinton J, Ahmad R, Miller DW, van der Brug MP, Canet-Aviles RM, Hague SM, Kaleem M, Cookson MR. Effects of DJ-1 mutations and polymorphisms on protein stability and subcellular localization. *Mol Brain Res* 2005;134:76-83.
- [41] Li HM, Niki T, Taira T, Iguchi-Ariga SMM, Ariga H. Association of DJ-1 with chaperones and enhanced association and colocalization with mitochondrial Hsp70 by oxidative stress. *Free Radic Res* 2005;39:1091-1099.
- [42] Ooc H, Taira T, Iguchi-Ariga SMM, Ariga H. Induction of reactive oxygen species by bisphenol A and abrogation of bisphenol A-induced cell injury by DJ-1. *Toxicol Sci* 2005;88:114-126.
- [43] Junn E, Taniguchi H, Jeong BS, Zhao X, Ichijo H, Mouradian MM. Interaction of DJ-1 with Daxx inhibits apoptosis signal-regulating kinase 1 activity and cell death. *Proc Natl Acad Sci USA* 2005;102:9691-9696.
- [44] Tobiume K, Inage T, Takeda K, Enomoto S, Miyazono K, Ichijo H. Molecular cloning and characterization of the mouse apoptosis signal-regulating kinase 1. *Biochem Biophys Res Commun* 1997;239:905-910.
- [45] Song JJ, Lee YJ. Role of the ASK1-SEK1-JNK1-HIPK1 signal in Daxx trafficking and ASK1 oligomerization. *J Biol Chem* 2003;278:47245-47252.
- [46] Kondo S, Lu Y, Debbas M, Lin AW, Sarosi I, Itie A, Wakeham A, Tuan J, Saris C, Elliott G, Ma W, Benchimol S, Lowe SW, Mak TW, Thukral SK. Characterization of cells and gene-targeted mice deficient for the p53-binding kinase homeodomain-interacting protein kinase 1 (HIPK1). *Proc Natl Acad Sci USA* 2003;100:5431-5436.
- [47] Li X, Zhang R, Luo D, Park SJ, Wang Q, Kim Y, Min W. Tumor necrosis factor alpha-induced desumoylation and cytoplasmic translocation of homeodomain-interacting protein kinase 1 are critical for apoptosis signal-regulating kinase 1-JNK/p38 activation. *J Biol Chem* 2005;280:15061-15070.
- [48] Song JJ, Lee YJ. Dissociation of Akt1 from its negative regulator JIP1 is mediated through the ASK1-MEK-JNK signal transduction pathway during metabolic oxidative stress: A negative feedback loop. *J Cell Biol* 2005;170:61-72.
- [49] Ecsedy JA, Michaelson JS, Leder P. Homeodomain-interacting protein kinase 1 modulates Daxx localization, phosphorylation, and transcriptional activity. *Mol Cell Biol* 2003;23:950-960.
- [50] Graham FJ, Van der Eb AJ. A new technique for the assay of infectivity of human adenovirus 5 DNA. *Virology* 1973;52:456-467.
- [51] Kim YH, Choi CY, Lee SJ, Conti MA, Kim Y. Homeodomain-interacting protein kinases, a novel family of co-repressors for homeodomain transcription factors. *J Biol Chem* 1998;273:25875-25879.
- [52] Honbou K, Suzuki NN, Horiuchi M, Niki T, Taira T, Ariga H, Inagaki F. The crystal structure of DJ-1, a protein related to male fertility and Parkinson's disease. *J Biol Chem* 2003;278:31380-31384.
- [53] Gorner K, Holtorf E, Odoy S, Nuscher B, Yamamoto A, Regula JT, Beyer K, Haass C, Kahle PJ. Differential effects of Parkinson's disease-associated mutations on stability and folding of DJ-1. *J Biol Chem* 2004;279:6943-6951.
- [54] Olzmann JA, Brown K, Wilkinson KD, Rees HD, Huai Q, Ke H, Levey AI, Li L, Chin LS. Familial Parkinson's disease-associated L166P mutation disrupts DJ-1 protein folding and function. *J Biol Chem* 2004;279:8506-8515.
- [55] Tao X, Tong L. Crystal structure of human DJ-1, a protein associated with early-onset Parkinson's disease. *J Biol Chem* 2003;278:31372-31379.
- [56] Wilson MA, Collins JL, Hod Y, Ringe D, Petsko GA. The 1.1-Å resolution crystal structure of DJ-1, the protein mutated in autosomal recessive early onset Parkinson's disease. *Proc Natl Acad Sci USA* 2003;100:9256-9261.
- [57] Nishinaga H, Takahashi-Niki K, Taira T, Andreadis A, Iguchi-Ariga SMM, Ariga H. Expression profiles of genes in DJ-1-knockdown and L166P DJ-1 mutant cells. *Neurosci Lett* 2005;390:54-59.



## Roles of distinct cysteine residues in *S*-nitrosylation and dimerization of DJ-1

Genta Ito <sup>a</sup>, Hiroyoshi Ariga <sup>b</sup>, Yasuhito Nakagawa <sup>c</sup>, Takeshi Iwatsubo <sup>a,\*</sup>

<sup>a</sup> *The Department of Neuropathology and Neuroscience, Graduate School of Pharmaceutical Sciences, University of Tokyo, Tokyo, Japan*

<sup>b</sup> *Department of Molecular Biology, Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan*

<sup>c</sup> *School of Pharmaceutical Sciences, Kitasato University, Tokyo, Japan*

Received 17 October 2005

Available online 18 November 2005

### Abstract

A significant proportion of early onset parkinsonism is inherited as an autosomal-recessive trait (AR-EP). DJ-1 was identified as one of the causative genes for AR-EP (PARK7), and DJ-1 protein has been implicated in oxidative stress response through oxidation of one of the three cysteine residues (i.e., Cys106). However, the individual roles of these cysteine residues remained unclear. We show by a systematic mutagenesis analysis that Cys46 and Cys53 of DJ-1, but not Cys106, are susceptible to *S*-nitrosylation in vitro as well as in cultured cells. Furthermore, alanine substitution of Cys46 diminished dimerization of DJ-1, a fundamental feature of this protein. These results indicate that distinct cysteine residues of DJ-1 harbor differential roles in relation to its structure and function.

© 2005 Elsevier Inc. All rights reserved.

**Keywords:** DJ-1; Parkinson's disease; *S*-nitrosylation; Dimerization; Neurodegeneration

Parkinson's disease (PD) is one of the most common neurodegenerative diseases affecting substantia nigra and presenting extrapyramidal symptoms including rigidity, tremor, and akinesia. A subset of PD patients manifest the symptoms early in their life (early onset PD: EP), and a significant proportion of EP is inherited as an autosomal-recessive trait (AR-EP). Subsequent to the discovery of parkin as the initial and major responsible gene for AR-EP (PARK2) [1], DJ-1 was identified as a causative gene for another type of AR-EP (PARK7) [2]. A number of studies have so far suggested that DJ-1 is a multi-functional protein, which is involved in transforming process of tumor cells [3], positive modulation of the transcriptional activity of androgen receptor [4,5], molecular chaperoning for  $\alpha$ -synuclein [6], or cell survival by antagonizing with PTEN [7]. Notably, the demonstration of an increase in the protein level of DJ-1 upon oxidative stress [8,9], as well as the anti-oxidative stress activity of DJ-1 shown in cultured

cells that overexpress DJ-1 [10,11], treated by RNAi for DJ-1 [12], or in DJ-1-deficient mice [13] or flies [14–17], strongly suggested that loss of anti-oxidative activity in the nigral neurons in PARK7 patients may cause the premature degeneration of these neurons. These findings also strongly supported the notion that oxidative stress may play a major role in the pathogenesis of PD and parkinsonism, based on genetics of familial PD.

X-ray crystallographic analyses have shown that DJ-1 forms a homodimer and that cysteine (Cys) 106 is selectively oxidized under an oxidative stress condition [18,19]. The specific oxidation of Cys106 upon oxidative stress was also confirmed by a rigorous protein chemical analysis [20]. Crystallographic analyses also showed that YDR533Cp and YajL/ThiJ, homologues of DJ-1 in *Saccharomyces cerevisiae* and *Escherichia coli*, respectively [21,22], form dimers, and the amino acid sequences located at the predicted dimer interface of DJ-1 and its homologues are highly conserved [23]. It has also been shown that L166P mutant DJ-1 linked to PARK7 fails to form a dimer and does not confer resistance to oxidative stress [10,24]. Based

\* Corresponding author. Fax: +81 3 5841 4708.

E-mail address: [iwatsubo@mol.f.u-tokyo.ac.jp](mailto:iwatsubo@mol.f.u-tokyo.ac.jp) (T. Iwatsubo).

on these findings, it is widely believed that dimerization is an important feature of DJ-1 that underlies its proper structure and functions.

Besides oxidation, protein *S*-nitrosylation is known as one of the reversible post-translational modifications that target the thiol group of Cys residues in proteins by the attack of nitrogen oxide (NO) [25], the latter being physiologically generated by NO synthase. It has been shown that the biological activities of a wide variety of proteins, e.g., thioredoxin [26], matrix metalloprotease-9 [27], and GAPDH [28], are modulated by *S*-nitrosylation. Protein *S*-nitrosylation is shown to be facilitated under an oxidative stress condition [29,30]. Notably, it has recently been shown that parkin is *S*-nitrosylated in the brains of patients with sporadic PD, leading to a decrease in its ubiquitin ligase activities [29,30].

These findings led us to examine the individual roles of the three distinct Cys residues in human DJ-1. In this report, we show that DJ-1 also is a target protein of *S*-nitrosylation *in vitro* and in cultured cells, and that Cys46 and Cys53, but not Cys106, are susceptible to this modification, using the biotin-switch assay, an efficient method to specifically detect protein *S*-nitrosylation [31]. We also show that Cys46 is involved in the dimerization of DJ-1.

## Materials and methods

**Construction of expression plasmids.** A cDNA encoding full-length human DJ-1 fused with an N-terminal FLAG tag in pcDNA3 vector (Invitrogen) was cloned as described (FLAG-DJ-1) [4]. cDNAs encoding mutant DJ-1 harboring Leu166Pro mutation (DJ-1/L166P), DJ-1/C46A, DJ-1/C53A, DJ-1/C106S, and C46A/C53A double mutation (DJ-1/C46/53A), DJ-1/C46/106A, DJ-1/C53/106A, or C46A/C53A/C106A triple mutation (Cys (-)) with N-terminal FLAG tags were generated by the long-PCR protocol using cDNAs encoding FLAG-DJ-1 as a template, and the following oligonucleotides were used as PCR primers: 5'-GGACCAGCTTCGAATTCGCGCCTGCAATTG-3' for DJ-1/L166P, 5'-CCCA GTACAGGCTAGTCGGATGTGGTCATTTG-3' for DJ-1/C46A, 5'-GTGGTCATTGCTCCTGATGCTAGCCTTGAA-3' for DJ-1/C53A, 5'-CTGATAGCCGCGATCGCTGCAGGTCCTAC-3' for DJ-1/C106A, 5'-CTGATAGCCGCGATATCTGCAGGTCCTAC-3' for DJ-1/C106S as forward primers, 5'-CAATTGCAGGCGCGAATTCGAAGCTGGTC C-3' for DJ-1/L166P, 5'-CAAATGACCACATCGCGACTAGCCTGTA CTGGG-3' for DJ-1/C46A, 5'-TTCAAGGCTAGCATCAGGAGC AATGACCAC-3' for DJ-1/C53A, 5'-GTAGGACCTGCAGCGATCGC GGCTATCAG-3' for and DJ-1/C106A, 5'-GTAGGACCTGCAGATA TCGCGGCTATCAG-3' for DJ-1/C106S as reverse primers. A cDNA encoding N-terminally myc-tagged DJ-1 (myc-DJ-1) was generated by PCR using *Platinum* Pfx (Invitrogen), and the following oligonucleotides were used as PCR primers: 5'-CCCAAGCTTCCACCATGGAACAAAA ACTCATCTCAGAAGAGGATCTGATGGCTTCCAAAAGAGCTC-3' as a forward primer and 5'-CCGCTCGAGC TAGTCTTTAAGAACAAG-3' as a reverse primer. The amplified cDNA was subcloned into pcDNA3 vector. cDNAs encoding DJ-1/L166P, DJ-1/C46A, DJ-1/C53A, and DJ-1/C106S fused with N-terminal myc tags were generated by the long-PCR protocol using a cDNA encoding myc-DJ-1 in pcDNA3 as the template, using primer pairs described above.

**Cell culture and transfection.** Monkey COS-1 cells and human SH-SY5Y neuroblastoma cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum and penicillin/streptomycin at 37 °C in 5% CO<sub>2</sub> atmosphere. Transient expression in COS-1 cells was performed by the DEAE-dextran method, and stable SH-SY5Y cell lines were generated by transfecting the cDNAs

in pcDNA3 vector using LipofectAMINE (Invitrogen) and selection in DMEM containing G418 (Calbiochem) at 500 µg/ml.

**Antibodies and immunochemical analyses.** The rabbit polyclonal antibody against the C-terminal region of human DJ-1 was generated by Nakagawa as described [8]. Other antibodies were purchased from Sigma (anti-FLAG M2 monoclonal, anti-FLAG polyclonal), Cell Signaling Technology (anti-myc 9B11 monoclonal), and Santa Cruz Biotechnology (anti-myc polyclonal), respectively. Sample preparations and analyses, including immunoblot analysis and co-immunoprecipitation, were performed as previously described [32,33], except for using 0.1% Triton X-100 as a detergent.

**Biotin-switch assay.** *S*-Nitrosylated proteins were detected using the biotin-switch assay as described by Jaffrey et al. [31] with some modifications. Briefly, samples that were prepared in HEN buffer (250 mM Hepes-NaOH, 1 mM EDTA, and 0.1 mM neocuproine, pH 7.7) containing 1% Triton X-100 were first incubated with *S*-nitrosoglutathione (GSNO; Sigma) or glutathione (GSH; WAKO) in darkness for 1 h at room temperature. The samples were then subjected to acetone precipitation to remove the NO donor and re-solubilized with 200 µl HENS buffer (HEN buffer containing 1% SDS) followed by addition of 2 µl of 2 M methyl methane thiosulfonate (MMTS; Pierce) to block the free thiol group. The samples were incubated at 50 °C for 20 min with frequent agitation by vortexing. MMTS was then removed by acetone precipitation, and the resultant precipitates were washed once with ice-cold acetone and then re-solubilized in 200 µl HENS buffer. The samples were then incubated with 2 mM ascorbic acid (Sigma) to release the NO from the *S*-nitrosylated thiol group, which was subsequently biotinylated by incubation with 400 µM biotin-HPDP (Pierce). The biotinylated proteins were then precipitated by incubating samples with 50 µl streptavidin-Sepharose beads High Performance (Amersham Biosciences). The streptavidin-Sepharose was then pelleted and washed 3 times with HEN buffer containing 0.5% SDS and 0.5% Triton X-100. The biotinylated proteins bound to the streptavidin beads were eluted by SDS-PAGE sample buffer and subjected to immunoblot analysis.

**Crosslinking assays.** Cells were rinsed with PBS once and harvested in PBS containing 0.1% Triton X-100. The samples were subsequently treated with 2.5 mg/ml of disuccinimidyl suberate (DSS) and incubated for 1 h at room temperature followed by dialysis against PBS for 8 h twice according to the manufacturer's instructions. The samples were then immunoprecipitated with anti-tag antibodies as indicated and subjected to an immunoblot analysis.

## Results

### *DJ-1 is S-nitrosylated at Cys46 and Cys53 in vitro and in cultured cells*

We first examined *in vitro S*-nitrosylation of DJ-1 using *S*-nitrosoglutathione (GSNO) as an NO donor by the biotin-switch assay. Basically, this method enables detection of *S*-nitrosylated thiol group of a Cys residue, by blocking the free thiol moiety by MMTS and then selectively biotinylating the originally *S*-nitrosylated moiety after reduction by ascorbate (covalently oxidized thiol group is not affected throughout the reactions). In the presence of 0.25 or 1 mM GSNO, we observed an extensive *S*-nitrosylation of FLAG-tagged (arrows) and endogenous (arrowheads) DJ-1 in lysates of stably transfected SH-SY5Y cells as revealed by a robust recovery of FLAG- or free DJ-1 by the biotin-switch assay (Fig. 1A, lanes 3 and 4). We next explored whether DJ-1 can be *S*-nitrosylated *in vivo* in cultured cells. For this purpose, we treated SH-SY5Y cells stably over-expressing FLAG-DJ-1 with GSNO and assessed the extent






Fig. 1. DJ-1 is *S*-nitrosylated in vitro and in cultured cells in vivo. (A) Lysates from SH-SY5Y cells stably overexpressing wild-type DJ-1 were subjected to in vitro *S*-nitrosylation and analyzed by the biotin-switch assay. Bound fractions to avidin beads were analyzed by immunoblotting with anti-FLAG (upper row) or anti-DJ-1 (lower row) antibodies. (B) SH-SY5Y cells stably overexpressing wild-type DJ-1 were treated with GSNO for 6 h at the indicated concentrations above the panels (mM). Bound ("ppt," lower row) and unbound ("sup," upper row) fractions to streptavidin beads were analyzed by immunoblotting with anti-FLAG antibody. Glutathione (GSH, lanes 1 and 2) was used as a negative control for the NO donor, *S*-nitrosoglutathione (GSNO, lanes 3 and 4). The numbers above each lane for GSNO and GSH indicate the concentration of reagents (mM). Note that lack of DJ-1 pulled down by streptavidin beads by MMTS treatment only (lane 2) indicates complete blocking of free thiol groups by this treatment, ensuring the specificity of detection of *S*-nitrosylated proteins. Arrow, FLAG-DJ-1; arrowhead, endogenous DJ-1.

of *S*-nitrosylation of DJ-1 by the biotin-switch assay. FLAG-tagged DJ-1 was biotinylated and pulled down by streptavidin beads (Fig. 1B, ppt, lanes 3 and 4, arrow) upon prior exposure to 1 or 3 mM GSNO, suggesting that DJ-1 can also be *S*-nitrosylated in vivo in cultured cells.

Previous reports have shown that DJ-1 is preferentially oxidized at Cys106 [18–20]. To examine which of the three Cys residues in DJ-1 is targeted to *S*-nitrosylation, we systematically mutated Cys residues of DJ-1 in various combinations, and overexpressed the Cys mutant (mt) DJ-1 in COS cells. We then treated the cell lysates by GSNO to induce *S*-nitrosylation of DJ-1 in vitro and detected *S*-nitrosylation of Cys mt DJ-1 by the biotin-switch assay. All the DJ-1 proteins harboring a single Cys substitution (i.e., at C46A, C53A or C106S) were *S*-nitrosylated, the C53A mt being least *S*-nitrosylated, followed by the C46A mt (Fig. 2A, lane 6). We then examined *S*-nitrosylation of the double Cys mt DJ-1. DJ-1 substituted at both Cys46 and Cys53 (C46/53A) was not *S*-nitrosylated (Fig. 2B, lane 6), whereas DJ-1 harboring either Cys46 or Cys53 (C46/106A, C53/106A) was *S*-nitrosylated, the C46/106A mt being less *S*-nitrosylated compared to C53/106A (Fig. 2B, lane 6). Taken together, we concluded that DJ-1 is *S*-nitrosylated at Cys46 and Cys53, but not at Cys106.




Fig. 2. DJ-1 is *S*-nitrosylated at Cys46 and Cys53. COS cells were transiently transfected with FLAG-DJ-1 harboring Cys substitutions at the indicated residues (A, single mutants; B, double mutants; Cys (-), triple mutant lacking all Cys residues), and the cell lysates were treated with GSNO or GSH in vitro as indicated above each lane. Bound ("ppt," lanes 4–6) and unbound ("sup," lanes 1–3) fractions to streptavidin beads were analyzed by immunoblotting with anti-FLAG antibody. Note that lane 6 represents the results of the biotin-switch assay. A faint band in lane 5 of Cys (-) is a non-specific one.

#### Dimer formation of DJ-1 is compromised by the C46A mutation

Based on the X-ray crystallographic analysis, wild-type (wt) DJ-1 has been shown to form a dimer: Cys53 residue was found to be located at the interface of the two DJ-1 proteins and Cys46 also resided close to the interface [18,19,23]. To examine the individual roles of these Cys residues on dimer formation of DJ-1, we generated wt FLAG- or myc-DJ-1 as well as C46A, C53A or C106S mt FLAG- or myc-DJ-1, and subjected these mutants to a co-immunoprecipitation analysis. Wt FLAG-DJ-1 and myc-DJ-1 doubly expressed in COS-1 cells were co-immunoprecipitated, whereas L166P mt FLAG-DJ-1 and myc-DJ-1 linked to PARK7 were not co-immunoprecipitated with each other, as previously described [34] (Fig. 3). C53A mt FLAG-DJ-1 was immunoprecipitated with C53A mt myc-DJ-1, as did C106S mt FLAG-DJ-1 and myc-DJ-1. In sharp contrast, C46A mt FLAG-DJ-1 and myc-DJ-1 showed only a trace amount of co-immunoprecipitation compared to wt or other Cys mt DJ-1, implicating Cys46 in the dimer formation of DJ-1 (Fig. 3). We further investigated the association of C46A mt DJ-1 by a chemical crosslinking analysis, using disuccinimidyl suberate (DSS), a primary amine-specific crosslinker that covalently crosslinks proteins with its interactors. We treated wt DJ-1 with DSS and then immunoprecipitated the covalently linked protein complexes, and found a clear band shift of DJ-1 to a position of ~50 kDa on immunoblots, which was equivalent to the size of a DJ-1 dimer, consistent with the results of a simple immunoprecipitation analysis (Fig. 4, DSS (+), arrow). In contrast to the results of simple co-immunoprecipitation, the intensity of the shifted band of C46A mt DJ-1 upon DSS treatment was




Fig. 3. Cys46 is required for the proper dimerization of DJ-1. COS-1 cells were transiently co-transfected with FLAG-DJ-1 and myc-DJ-1 of wild-type (wt), or those harboring corresponding Cys (C46A, C53A, and C106S) or L166P mutations. The cell lysates were immunoprecipitated with anti-myc or anti-FLAG antibodies (3rd and 4th rows from top, respectively), and then analyzed by immunoblotting with anti-FLAG (left panel) or anti-myc (right panel) antibodies, respectively. Control, immunoprecipitation using an irrelevant mouse IgG (LB509). Input shows the immunoblotting of fractions subjected to immunoprecipitation.




Fig. 4. Dimerization of DJ-1 assessed by co-immunoprecipitation followed by chemical crosslinking with DSS. Lysates of COS-1 cells co-transfected with wt or mt FLAG- and myc-DJ-1 (as in Fig. 3) were immunoprecipitated by anti-myc antibody, and analyzed by immunoblotting with anti-FLAG antibody, with (right panel) or without (left panel) crosslinking with DSS. The upper panel (input) shows the immunoblotting of fractions subjected to immunoprecipitation. Arrow, crosslinked DJ-1 dimer; arrowhead, DJ-1 monomer. Molecular mass standards are shown in kilodaltons at the left of the panel.

similar to that of wt DJ-1 (Fig. 4, DSS (+), arrow). These results suggest that substitution of Cys46 disrupts the physical interaction of two DJ-1 polypeptides, although the positioning of the two polypeptides may be relatively preserved, thus allowing the chemical crosslinking of the dimer in the presence of an excess amount of a crosslinker.

## Discussion

Human DJ-1 protein harbors three Cys residues, two of which (i.e., Cys46 and Cys106) are evolutionarily highly conserved [23], suggesting the functional and structural importance of these residues. In the present study, we show that (i) Cys46 and Cys53 are susceptible to *S*-nitrosylation,

and that (ii) Cys46 is required for the proper dimerization of DJ-1.

Protein chemical and crystallographic analyses have shown that Cys106 of DJ-1 is preferentially oxidized under an oxidative condition [18–20]. In this regard, the possibility that the exclusive detection of *S*-nitrosylation at Cys46 and Cys53 may reflect the complete and irreversible masking of the thiol group of Cys106 by an extensive oxidation upon overexpression in cultured cells, resulting in insusceptibility to *S*-nitrosylation of this residue. However, our observation that a significant amount of C46/53A double mt DJ-1 was biotinylated and pulled down upon omission of MMTS treatment that blocks the free thiol group (Fig. 2B, lane 5) strongly indicates that sufficient amount of free (i.e., non-oxidated) Cys106 was retained in the C46/53A mutant DJ-1 proteins, arguing against the liability of Cys106 to be *S*-nitrosylated. *S*-nitrosylation has been known to preferentially occur at Cys residues flanked by basic and acidic residues [35], although the amino acid sequences around Cys46 or Cys53 of DJ-1 do not fit with these characteristics. Therefore, the propensity of DJ-1 to be *S*-nitrosylated at Cys46 and Cys53 might be due to the three-dimensional dispositions of the Cys residues in DJ-1. It remains to be determined which of these two Cys residues is the major potential *S*-nitrosylation site: the extent of *S*-nitrosylation of these two Cys residues was apparently different between the single or double mutants, which also may be due to the minute differences in three-dimensional arrangements of the Cys residues. The effects of *S*-nitrosylation on the structure (e.g., dimerization) and function (e.g., anti-oxidative activity) of DJ-1 should further be clarified. Since protein *S*-nitrosylation may be facilitated under oxidative stress conditions [29,30], the latter being implicated in the pathophysiology of PD and related disorders [36], it is tempting to speculate that the function and structure of DJ-1 may be altered by *S*-nitrosylation in the brains of patients with sporadic or familial PD, like in the case of *S*-nitrosylation of parkin [29,30].

Cys46 has been predicted to be located close to the dimer interface [18,19,23]. Taken together with our observation that C46A substitution significantly affected the stable dimerization of DJ-1, we speculate that Cys46 plays a critical role in the maintenance of the structure of the DJ-1, especially at the dimer interface. However, the successful crosslinking of C46A mt DJ-1 as a dimer may suggest that the structural alteration of the dimer interface caused by this substitution may be a subtle one. Nonetheless, the fact that such a slight alteration led to a significant reduction in co-immunoprecipitation of dimer implicates Cys46 in the function and structure of DJ-1, which is in agreement with the phylogenetical conservation of Cys46. In contrast, Cys53, which is located at the dimer interface close to each other at a distance of 3.1 Å but not as highly conserved as Cys46 (e.g., unconserved in invertebrates), was not required for the dimerization of DJ-1.

Our observation that distinct Cys residues in DJ-1 harbor different functional and structural roles underscores the need for further investigations into the changes in DJ-1 in relation to oxidative and nitrosative stress conditions, especially in PD and related neurodegenerative conditions.

### Acknowledgments

The authors thank Y. Takanezawa, A. Mitsumoto, M. Hasegawa, T. Kuwahara, Y. Hanno, S. Takatori, and T. Uehara for helpful discussions. This study was supported by the Ministry of Education, Science, Culture and Sports on Priority Areas-Advanced Brain Science Project and -Research on Pathomechanisms of Brain Disorders, and for the 21st Century Center of Excellence Program, and by the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO), Japan.

### References

- [1] T. Kitada, S. Asakawa, N. Hattori, H. Matsumine, Y. Yamamura, S. Minoshima, M. Yokochi, Y. Mizuno, N. Shimizu, Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism, *Nature* 392 (1998) 605–608.
- [2] V. Bonifati, P. Rizzu, M.J. van Baren, O. Schaap, G.J. Breedveld, E. Krieger, M.C. Dekker, F. Squitieri, P. Ibanez, M. Joosse, J.W. van Dongen, N. Vanacore, J.C. van Swieten, A. Brice, G. Meco, C.M. van Duijn, B.A. Oostra, P. Heutink, Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism, *Science* 299 (2003) 256–259.
- [3] D. Nagakubo, T. Taira, H. Kitaura, M. Ikeda, K. Tamai, S.M. Iguchi-Ariga, H. Ariga, DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras, *Biochem. Biophys. Res. Commun.* 231 (1997) 509–513.
- [4] K. Takahashi, T. Taira, T. Niki, C. Seino, S.M. Iguchi-Ariga, H. Ariga, DJ-1 positively regulates the androgen receptor by impairing the binding of PIAS $\alpha$  to the receptor, *J. Biol. Chem.* 276 (2001) 37556–37563.
- [5] T. Niki, K. Takahashi-Niki, T. Taira, S.M. Iguchi-Ariga, H. Ariga, DJBP: a novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex, *Mol. Cancer Res.* 1 (2003) 247–261.
- [6] S. Shendelman, A. Jonason, C. Martinat, T. Leete, A. Abeliovich, DJ-1 is a redox-dependent molecular chaperone that inhibits  $\alpha$ -synuclein aggregate formation, *PLoS Biol.* 2 (2004) e362.
- [7] R.H. Kim, M. Peters, Y. Jang, W. Shi, M. Pintilie, G.C. Fletcher, C. DeLuca, J. Liepa, L. Zhou, B. Snow, R.C. Binari, A.S. Manoukian, M.R. Bray, F.F. Liu, M.S. Tsao, T.W. Mak, DJ-1, a novel regulator of the tumor suppressor PTEN, *Cancer Cell* 7 (2005) 263–273.
- [8] A. Mitsumoto, Y. Nakagawa, A. Takeuchi, K. Okawa, A. Iwamatsu, Y. Takanezawa, Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat, *Free Radic Res.* 35 (2001) 301–310.
- [9] J. Jin, G.E. Meredith, L. Chen, Y. Zhou, J. Xu, F.S. Shie, P. Lockhart, J. Zhang, Quantitative proteomic analysis of mitochondrial proteins: relevance to Lewy body formation and Parkinson's disease, *Brain Res. Mol. Brain Res.* 134 (2005) 119–138.
- [10] T. Taira, Y. Saito, T. Niki, S.M. Iguchi-Ariga, K. Takahashi, H. Ariga, DJ-1 has a role in antioxidative stress to prevent cell death, *EMBO Rep.* 5 (2004) 213–218.
- [11] R.M. Canet-Aviles, M.A. Wilson, D.W. Miller, R. Ahmad, C. McLendon, S. Bandyopadhyay, M.J. Baptista, D. Ringe, G.A. Petsko, M.R. Cookson, The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization, *Proc. Natl. Acad. Sci. USA* 101 (2004) 9103–9108.
- [12] T. Yokota, K. Sugawara, K. Ito, R. Takahashi, H. Ariga, H. Mizusawa, Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition, *Biochem. Biophys. Res. Commun.* 312 (2003) 1342–1348.
- [13] R.H. Kim, P.D. Smith, H. Aleyasin, S. Hayley, M.P. Mount, S. Pownall, A. Wakeham, A.J. You-Ten, S.K. Kalia, P. Horne, D. Westaway, A.M. Lozano, H. Anisman, D.S. Park, T.W. Mak, Hypersensitivity of DJ-1-deficient mice to 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) and oxidative stress, *Proc. Natl. Acad. Sci. USA* 102 (2005) 5215–5220.
- [14] M. Meulener, A.J. Whitworth, C.E. Armstrong-Gold, P. Rizzu, P. Heutink, P.D. Wes, L.J. Pallanck, N.M. Bonini, *Drosophila* DJ-1 mutants are selectively sensitive to environmental toxins associated with Parkinson's disease, *Curr. Biol.* 15 (2005) 1572–1577.
- [15] F.M. Menzies, S.C. Yenisseti, K.T. Min, Roles of *Drosophila* DJ-1 in survival of dopaminergic neurons and oxidative stress, *Curr. Biol.* 15 (2005) 1578–1582.
- [16] Y. Yang, S. Gehrke, M.E. Haque, Y. Imai, J. Kosek, L. Yang, M.F. Beal, I. Nishimura, K. Wakamatsu, S. Ito, R. Takahashi, B. Lu, Inactivation of *Drosophila* DJ-1 leads to impairments of oxidative stress response and phosphatidylinositol3-kinase/Akt signaling, *Proc. Natl. Acad. Sci. USA* 102 (2005) 13670–13675.
- [17] J. Park, S.Y. Kim, G.H. Cha, S.B. Lee, S. Kim, J. Chung, *Drosophila* DJ-1 mutants show oxidative stress-sensitive locomotive dysfunction, *Gene* 361C (2005) 133–139.
- [18] M.A. Wilson, J.L. Collins, Y. Hod, D. Ringe, G.A. Petsko, The 1.1-Å resolution crystal structure of DJ-1, the protein mutated in autosomal recessive early onset Parkinson's disease, *Proc. Natl. Acad. Sci. USA* 100 (2003) 9256–9261.
- [19] S.J. Lee, S.J. Kim, I.K. Kim, J. Ko, C.S. Jeong, G.H. Kim, C. Park, S.O. Kang, P.G. Suh, H.S. Lee, S.S. Cha, Crystal structures of human DJ-1 and *Escherichia coli* Hsp31, which share an evolutionarily conserved domain, *J. Biol. Chem.* 278 (2003) 44552–44559.
- [20] T. Kinumi, J. Kimata, T. Taira, H. Ariga, E. Niki, Cysteine-106 of DJ-1 is the most sensitive cysteine residue to hydrogen peroxide-mediated oxidation in vivo in human umbilical vein endothelial cells, *Biochem. Biophys. Res. Commun.* 317 (2004) 722–728.
- [21] M.A. Wilson, C.V. St. Amour, J.L. Collins, D. Ringe, G.A. Petsko, The 1.8-Å resolution crystal structure of YDR533Cp from *Saccharomyces cerevisiae*: a member of the DJ-1/ThiJ/PfpI superfamily, *Proc. Natl. Acad. Sci. USA* 101 (2004) 1531–1536.
- [22] M.A. Wilson, D. Ringe, G.A. Petsko, The atomic resolution crystal structure of the YajL (ThiJ) protein from *Escherichia coli*: a close prokaryotic homologue of the Parkinsonism-associated protein DJ-1, *J. Mol. Biol.* 353 (2005) 678–691.
- [23] K. Honbou, N.N. Suzuki, M. Horiuchi, T. Niki, T. Taira, H. Ariga, F. Inagaki, The crystal structure of DJ-1, a protein related to male fertility and Parkinson's disease, *J. Biol. Chem.* 278 (2003) 31380–31384.
- [24] D.W. Miller, R. Ahmad, S. Hague, M.J. Baptista, R. Canet-Aviles, C. McLendon, D.M. Carter, P.P. Zhu, J. Stadler, J. Chandran, G.R. Klinefelter, C. Blackstone, M.R. Cookson, L166P mutant DJ-1, causative for recessive Parkinson's disease, is degraded through the ubiquitin-proteasome system, *J. Biol. Chem.* 278 (2003) 36588–36595.
- [25] J.S. Stamler, D.I. Simon, J.A. Osborne, M.E. Mullins, O. Jaraki, T. Michel, D.J. Singel, J. Loscalzo, S-nitrosylation of proteins with nitric oxide: synthesis and characterization of biologically active compounds, *Proc. Natl. Acad. Sci. USA* 89 (1992) 444–448.
- [26] J. Haendeler, J. Hoffmann, V. Tischler, B.C. Berk, A.M. Zeiher, S. Dimmeler, Redox regulatory and anti-apoptotic functions of thioredoxin depend on S-nitrosylation at cysteine69, *Nat. Cell Biol.* 4 (2002) 743–749.

- [27] Z. Gu, M. Kaul, B. Yan, S.J. Kridel, J. Cui, A. Strongin, J.W. Smith, R.C. Liddington, S.A. Lipton, S-nitrosylation of matrix metalloproteinases: signaling pathway to neuronal cell death, *Science* 297 (2002) 1186–1190.
- [28] M.R. Hara, N. Agrawal, S.F. Kim, M.B. Cascio, M. Fujimuro, Y. Ozeki, M. Takahashi, J.H. Cheah, S.K. Tankou, L.D. Hester, C.D. Ferris, S.D. Hayward, S.H. Snyder, A. Sawa, S-nitrosylated GAPDH initiates apoptotic cell death by nuclear translocation following Siah1 binding, *Nat. Cell Biol.* 7 (2005) 665–674.
- [29] K.K. Chung, B. Thomas, X. Li, O. Pletnikova, J.C. Troncoso, L. Marsh, V.L. Dawson, T.M. Dawson, S-nitrosylation of parkin regulates ubiquitination and compromises parkin's protective function, *Science* 304 (2004) 1328–1331.
- [30] D. Yao, Z. Gu, T. Nakamura, Z.Q. Shi, Y. Ma, B. Gaston, L.A. Palmer, E.M. Rockenstein, Z. Zhang, E. Masliah, T. Uehara, S.A. Lipton, Nitrosative stress linked to sporadic Parkinson's disease: S-nitrosylation of parkin regulates its E3 ubiquitin ligase activity, *Proc. Natl. Acad. Sci. USA* 101 (2004) 10810–10814.
- [31] S.R. Jaffrey, H. Erdjument-Bromage, C.D. Ferris, P. Tempst, S.H. Snyder, Protein S-nitrosylation: a physiological signal for neuronal nitric oxide, *Nat. Cell Biol.* 3 (2001) 193–197.
- [32] T. Tomita, K. Maruyama, T.C. Saido, H. Kume, K. Shinozaki, S. Tokuhira, A. Capell, J. Walter, J. Grunberg, C. Haass, T. Iwatsubo, K. Obata, The presenilin 2 mutation (N141I) linked to familial Alzheimer disease (Volga German families) increases the secretion of amyloid beta protein ending at the 42nd (or 43rd) residue, *Proc. Natl. Acad. Sci. USA* 94 (1997) 2025–2030.
- [33] Y. Morohashi, N. Hatano, S. Ohya, R. Takikawa, T. Watabiki, N. Takasugi, Y. Imaizumi, T. Tomita, T. Iwatsubo, Molecular cloning and characterization of CALP/KChIP4, a novel EF-hand protein interacting with presenilin 2 and voltage-gated potassium channel subunit Kv4, *J. Biol. Chem.* 277 (2002) 14965–14975.
- [34] J.A. Olzmann, K. Brown, K.D. Wilkinson, H.D. Rees, Q. Huai, H. Ke, A.I. Levey, L. Li, L.S. Chin, Familial Parkinson's disease-associated L166P mutation disrupts DJ-1 protein folding and function, *J. Biol. Chem.* 279 (2004) 8506–8515.
- [35] J.S. Stamler, E.J. Toone, S.A. Lipton, N.J. Sucher, (S)NO signals: translocation, regulation, and a consensus motif, *Neuron* 18 (1997) 691–696.
- [36] W. Dauer, S. Przedborski, Parkinson's disease: mechanisms and models, *Neuron* 39 (2003) 889–909.