

FIG. 7. Sensitization of BPA-induced cell death in DJ-1-knockdown cells. Neuro2a (A) and GC1 cells (B) were transfected with siRNA against *DJ-1* or *luciferase* by lipofectamine plus (Invirogen). At 24 h after transfection, cells were administered various concentrations of BPA. At 48 h after administration, viabilities of cells were measured by an MTT assay. (A) Neuro2a ( $n = 5$ ). (B) GC1 ( $n = 5$ ). Asterisks indicate significant difference from transfection of siRNA targeting the *luciferase* gene; \* $p < 0.05$ ; \*\* $p < 0.01$ . Symbols of "untreated, vehicle, siLuc and siDJ-1" in figures indicate cells not transfected, transfected with lipofectamine plus alone, transfected with siRNA targeting the *luciferase* gene, and transfected with siRNA targeting the *DJ-1* gene, respectively.

to restore the activity of complex 1. Since DJ-1 was located in the cytoplasm, nucleus, and mitochondria in various types of cells (Bandopadhyay *et al.*, 2004; Nagakubo *et al.*, 1997; Shendelman *et al.*, 2004; Shinbo *et al.*, 2005a, in press; Wagenfeld *et al.*, 2000; Zhang *et al.*, 2005), and some DJ-1 was translocated to mitochondria after oxidative stress (Blackinton *et al.*, 2005; Jin *et al.*, 2005; Li *et al.*, 2005), the localization of DJ-1 in mitochondria in the cells that had been administered BPA was thought to affect BPA-induced ROS production.


The activity of DJ-1 to sustain the activity of complex 1 was, however, observed at 50 and 100  $\mu\text{M}$  BPA in Neuro2a and GC1 cells, respectively, and this activity was lost over these concentrations. Since, at high concentrations of BPA, cells began to die and the amount of DJ-1 also decreased, it is thought that the amount of DJ-1 is not sufficient to sustain the activity of complex 1. Alternatively, although DJ-1 contains three cysteines, and oxidation of a cysteine at amino acid number 106 (C106) is essential for DJ-1 to exert its anti-oxidative stress

activity (Taira *et al.*, 2004b; Takahashi-Niki *et al.*, 2004), oxidation of all three cysteines by excess ROS, which is produced by high doses of BPA, for instance, may result in loss of its activity (Kinumi *et al.*, 2004; Taira *et al.*, 2004b).

Although DJ-1 is an abundant protein and approximately  $5 \times 10^5$  molecules/cell of DJ-1 is present in cells, it is likely that, in addition to DJ-1, other ROS-scavenging proteins such as superoxide dismutase, glutathione peroxidase and catalase, and proteins belonging to other redox systems participate in protecting cells from BPA-induced cell injury. We have found that drastic changes in expressions of genes, including genes related to stress, apoptosis, oxidative stress, and neurotoxicity, occurred in DJ-1 knockdown and that expressions of some genes were regulated by DJ-1 (Nishinaga *et al.*, in press), suggesting that in addition to anti-oxidative activity of DJ-1, transcriptional function of DJ-1 toward genes related to stress functions is important.

In this study, levels of ROS production peaked at 12 h after BPA administration in GC1 cells and then decreased at 24 and

FIG. 6. Dysfunction of mitochondria in cultured cell induced by BPA administration. Neuro2a (A) and GC1 cells (B) were administered various concentrations of BPA. At 48 h after administration, mitochondria-rich fractions were prepared from cells, and complex 1 activities were measured using 70  $\mu\text{g}$  proteins as described in Materials and Methods. Twenty  $\mu\text{g}$  of proteins were analyzed by Western blotting to detect 39-, 30-, and 20-kDa subunits of complex 1. Intensities of bands were quantified as described in Materials and Methods, and relative expression of each subunit to that of untreated cells (untreated) at 0 h is also shown (C, Neuro2a; D, GC1). DJ-1 expression in Neuro2a (E) and GC1 cells (F) that had been administered BPA was analyzed by Western blotting with an anti-DJ-1 antibody. Twenty nM siRNAs against *luciferase* or *DJ-1* were transfected into cells. At 72 h after transfection, complex 1 activity and expressions of subunits of complex 1 and DJ-1 were analyzed as described above. (A, C, and E) Neuro2a ( $n = 6$ ). (B, D, and F) GC1 ( $n = 7$ ). Asterisks indicate significant difference from the control and transfection of siRNA targeting the *luciferase* gene; \* $p < 0.05$ ; \*\* $p < 0.01$ .



**FIG. 8.** Abrogation of BPA-induced cell death by DJ-1 in Neuro2a and GC1 cells. (A, B) Neuro2a and GC1 cells in 6-well plates were transfected with 650 and 400 ng of pcDNA3-F-DJ-1 or pcDNA3-F, respectively, using Lipofectamine 2000 (Invitrogen) according to the supplier's manual. At 72 h after transfection, proteins extracted from Neuro2a (A) and GC1 (B) cells were analyzed by Western blotting with anti-DJ-1 (ab4150, Abcam) and anti-actin (Chemicon) antibodies. Proteins were then reacted with an IRDye800-conjugated second antibody and visualized by an infrared imaging system (Odyssey, LI-COR). (C, D) Neuro2a and GC1 cells in 96-well plates were transfected with 16 and 10 ng of pcDNA3-F-DJ-1 or pcDNA3-F, respectively, using Lipofectamine 2000. At 24 h after transfection, Neuro2a (C) and GC1 (D) cells were administered various concentrations of BPA. At 48 h after administration, viabilities of cells were measured by an MTT assay. (C) Neuro2a ( $n = 5$ ). (D) GC1 ( $n = 5$ ). Asterisks indicate significant difference from transfection of pcDNA3, \* $p < 0.05$ ; \*\* $p < 0.01$ . Symbols of "untreated, vehicle, vector and DJ-1" in figures indicate cells not transfected, transfected with lipofectamine plus alone, transfected with pcDNA3-F, and transfected with pcDNA3-F-DJ-1, respectively.

48 h. ROS production in Neuro2A cells, however, continued to occur during these periods. In the kidney, liver, and testis, an isoform of UDP-glucuronosyltransferase (UGTs) has been shown to metabolize BPA to BPA-glucuronide (Reinhechel *et al.*, 1995). Although the presence of UGTs in GC1 cells has not been examined, BPA produced at 12 h after BPA administration might be metabolized in GC1 cells.

The present study showed that BPA induced production of various ROS, including highly reactive hydroxy radicals, and ROS have been shown to attack mitochondrial complex I (Yokota *et al.*, 1999). Nakagawa and Toyama reported reduction of complex I activity after incubation of a mitochondria-rich fraction with BPA *in vitro*, but degradation of mitochondria and production of ROS were not examined (Nakagawa and Toyama, 2000). Since the present study showed that BPA induces ROS production and inhibits complex I activity by disrupting complex I, it would be interesting to examine whether degradation of complex I by BPA also occurs *in vitro* using

an electron microscopy and whether this degradation is restored by DJ-1.

Environmental factors are thought to trigger the onset of PD. Several neurotoxins, including 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTT), rotenone, 6-hydroxydopamine (6-OHDA), and a metabolite of dopamine itself, have been reported to injure mitochondria, resulting in dopaminergic neuron death (Jenner, 2003). Recently, BPA has been reported to affect movement and to induce dopaminergic neuron death when BPA was administered to embryonal mice and into the substantia nigra of adult rats, respectively, though ROS levels were not examined (Kabuto *et al.*, 2004; Mizuo *et al.*, 2004; Suzuki *et al.*, 2003). BPA has been detected in several tissues and the blood serum of humans (Ikezuki *et al.*, 2002; Schonfelder *et al.*, 2002) and accumulates in adipose tissue and membranes, creating the potential for long-term exposure of humans to a low dose of BPA. Human exposure to PBA, and our present results, suggest that BPA could be a risk factor for the

onset of familial and sporadic PD. Furthermore, there are many factors in the environment that cause oxidative stress to humans, including smoking, UV light, alcohol, and environmental contaminants. These factors, including BPA, might act by themselves or in combination as risk factors for diseases such as cancer, infertility, and neurodegenerative diseases.

Since the activity of mitochondrial complex I was significantly diminished by siRNA targeting DJ-1 in the cells of this study, it is possible that DJ-1 has a novel function to maintain complex I activity. In DJ-1-knockdown cells, complex I was still intact. Although the precise molecular basis of this reaction is not clear at present, it is important to clarify its mechanism.

#### ACKNOWLEDGMENTS

We thank Yoko Misawa and Kiyomi Takaya for their technical assistance. This work was supported by grants-in-aid from the Ministry of Education, Science, Culture, Sports and Technology of Japan and the Ministry of Health, Labor and Welfare of Japan. Conflict of interest: none declared.

#### REFERENCES

- Abou-Sleiman, P. M., Healy, D. G., Quinn, N., Lees, A. J., and Wood, N. W. (2003). The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann. Neurol.* **54**, 283–286.
- Agarwal, A., Saleh, R. A., and Bedaiwy, M. A. (2003). Role of reactive oxygen species in the pathophysiology of human reproduction. *Fertil. Steril.* **79**, 829–843.
- Al-Hiyasat, A. S., Darmani, H., and Elbetieha, A. M. (2004). Leached components from dental composites and their effects on fertility of female mice. *Eur. J. Oral. Sci.* **112**, 267–272.
- Bandopadhyay, R., Kingsbury, A. E., Cookson, M. R., Reid, A. R., Evans, I. M., Hope, A. D., Pittman, A. M., Lashley, T., Canet-Aviles, R., Miller, D. W., et al. (2004). The expression of DJ-1 (PARK7) in normal human CNS and idiopathic Parkinson's disease. *Brain* **127**, 420–430.
- Benhar, M., Engelberg, D., and Levitzki, A. (2002). ROS, stress-activated kinases and stress signaling in cancer. *EMBO Rep.* **3**, 420–425.
- Biles, J. E., White, K. D., McNeal, T. P., and Begley, T. H. (1999). Determination of the diglycidyl ether of bisphenol A and its derivatives in canned foods. *J. Agric. Food Chem.* **47**, 1965–1969.
- Bindhumol, V., Chitra, K. C., and Mathur, P. P. (2003). Bisphenol A induces reactive oxygen species generation in the liver of male rats. *Toxicology* **188**, 117–124.
- Blackinton, J., Ahmad, R., Miller, D. W., van der Brug, M. P., Canet-Aviles, R. M., Hague, S. M., Kaleem, M., and Cookson, M. R. (2005). Effects of DJ-1 mutations and polymorphisms on protein stability and subcellular localization. *Mol. Brain Res.* **134**, 76–83.
- Bonifati, V., Rizzu, P., van Baren, M. J., Schaap, O., Breedveld, G. J., Krieger, E., Dekker, M. C., Squitieri, F., Ibanez, P., Joosse, M., et al. (2003). Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* **299**, 256–259.
- Canet-Aviles, R. M., Wilson, M. A., Miller, D. W., Ahmad, R., McLendon, C., Bandyopadhyay, S., Baptista, M. J., Ringe, D., Petsko, G. A., and Cookson, M. R. (2004). The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 9103–9108.
- Chitra, K. C., Latchoumycandane, C., and Mathur, P. P. (2003). Induction of oxidative stress by bisphenol A in the epididymal sperm of rats. *Toxicology* **14**, 119–127.
- Corti, O., Hampe, C., Darios, F., Ibanez, P., Ruberg, M., and Brice, A. (2005). Parkinson's disease: From causes to mechanisms. *C. R. Biol.* **328**, 131–142.
- Golden, T. R., Hinerfeld, D. A., and Melov, S. (2002). Oxidative stress and aging: Beyond correlation. *Aging Cell* **1**, 117–123.
- Hague, S., Rogaeva, E., Hernandez, D., Gulick, C., Singleton, A., Hanson, M., Johnson, J., Weiser, R., Gallardo, M., Ravina, B., et al. (2003). Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann. Neurol.* **54**, 271–274.
- Ikezuki, Y., Tsutsumi, O., Takai, Y., Kamei, Y., Taketani, Y. (2002). Determination of bisphenol A concentrations in human biological fluids reveals significant early prenatal exposure. *Hum. Reprod.* **17**, 2839–2841.
- Jenner, P. (2003). Oxidative stress in Parkinson's disease. *Ann. Neurol.* **53** (Suppl. 3), S26–S36.
- Jin, J., Meredith, G. E., Chen, L., Zhou, Y., Xu, J., Shie, F. S., Lockhart, P., and Zhang, J. (2005). Quantitative proteomic analysis of mitochondrial proteins: relevance to Lewy body formation and Parkinson's disease. *Mol. Brain Res.* **134**, 119–138.
- Junqueira, V. B., Simizu, K., Van Halsema, L., Koch, O. R., Barro, S. B., and Videla, L. A. (1988). Lindane-induced oxidative stress. I. Time course of changes in hepatic microsomal parameters, antioxidant enzymes, lipid peroxidative indices and morphological characteristics. *Xenobiotica* **18**, 1297–1304.
- Kabuto, H., Amakawa, M., and Shishibori, T. (2004). Exposure to bisphenol A during embryonic/fetal life and infancy increases oxidative injury and causes underdevelopment of the brain and testis in mice. *Life Sci.* **74**, 2931–2940.
- Kabuto, H., Hasuike, S., Minagawa, N., and Shishibori, T. (2003). Effects of bisphenol A on the metabolisms of active oxygen species in mouse tissues. *Environ. Res.* **93**, 31–35.
- Kim, R. H., Peters, M., Jang, Y., Shi, W., Pintilie, M., Fletcher, G. C., DeLuca, C., Liepa, J., Zhou, L., Snow, B., et al. (2005). DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* **7**, 263–273.
- Kinumi, T., Kimata, J., Taira, T., Ariga, H., and Niki, E. (2004). Cysteine-106 of DJ-1 is the most sensitive cysteine residue to hydrogen peroxide mediated oxidation *in vivo* in human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* **317**, 722–728.
- Klinefelter, G. R., Welch, J. E., Perreault, S. D., Moore, H. D., Zucker, R. M., Suarez, J. D., Roberts, N. L., Bobssine, K., and Jeffay, S. J. (2002). Localization of the sperm protein SP22 and inhibition of fertility *in vivo* and *in vitro*. *Andrology* **23**, 48–63.
- Kobzik, L., Godleski, J. J., and Brain, J. D. (1990). Oxidative metabolism in the alveolar macrophage: Analysis by flow cytometry. *J. Leukoc. Biol.* **47**, 295–303.
- Latchoumycandane, C., Chitra, K. C., and Mathur, P. P. (2002). The effect of 2,3,7,8-tetrachlorodibenzo-p-dioxin on the antioxidant system in mitochondrial and microsomal fractions of rat testis. *Toxicology* **171**, 127–135.
- Law, P., Campbell, S. D., Lepock, J. R., and Kruuv, J. (1986). Effects of butylated hydroxytoluene on membrane lipid fluidity and freezethaw survival in mammalian cell. *Cryobiology* **23**, 317–322.
- Li, H. M., Niki, T., Taira, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2005). Association of DJ-1 with chaperones and enhanced association and colocalization with mitochondrial Hsp70 by oxidative stress. *Free Radic. Res.* (in press).
- Martinat, C., Shendelman, S., Jonason, A., Leete, T., Beal, M. F., Yang, L., Floss, T., and Abeliovich, A. (2004). Sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: An ES-derived cell model of primary parkinsonism. *PLoS Biol.* **2**, e327.
- Mitsumoto, A., and Nakagawa, Y. (2001). DJ-1 is an indicator for endogenous reactive oxygen species elicited by endotoxin. *Free Radic. Res.* **35**, 885–893.

- Mitsumoto, A., Nakagawa, Y., Takeuchi, A., Okawa, K., Iwamatsu, A., and Takanezawa, Y. (2001). Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radic. Res.* **35**, 301–310.
- Mizuo, K., Narita, M., Miyagawa, K., Narita, M., Okuno, E., and Suzuki, T. (2004). Prenatal and neonatal exposure to bisphenol-A affects the morphine-induced rewarding effect and hyperlocomotion in mice. *Neurosci. Lett.* **356**, 95–98.
- Nagakubo, D., Taira, T., Kitaura, H., Ikeda, M., Tamai, K., Iguchi-Ariga, S. M. M., and Ariga, H. (1997). DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem. Biophys. Res. Commun.* **31**, 509–513.
- Nakagawa, Y., and Toyama, S. (2000). Metabolism and cytotoxicity of bisphenol A and other bisphenols in isolated rat hepatocytes. *Arch. Toxicol.* **74**, 99–105.
- Nikaido, Y., Yoshizawa, K., Danbara, N., Tsujita-Kyutoku, M., Yuri, T., Uehara, N., and Tsubura, A. (2004). Effects of maternal xenoestrogen exposure on development of the reproductive tract and mammary gland in female CD-1 mouse offspring. *Reprod. Toxicol.* **18**, 803–811.
- Niki, T., Takahashi-Niki, K., Taira, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2003). DJBP: A novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol. Cancer Res.* **1**, 247–261.
- Nishinaga, H., Takahashi-Niki, K., Taira, T., Andreadis, A., Iguchi-Ariga, S. M. M., and Ariga, H. (2005). Expression profiles of genes in DJ-1-knockdown and L166P DJ-1 mutant cells. *Neurosci. Lett.* (in press).
- Nunez, A. A., Kannan, K., Giesy, J. P., Fang, J., and Clemens, L. G. (2001). Effects of bisphenol A on energy balance and accumulation in brown adipose tissue in rats. *Chemosphere* **42**, 917–922.
- Okada, M., Matsumoto, K., Niki, T., Taira, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2002). DJ-1, a target protein for an endocrine disrupter, participates in the fertilization in mice. *Biol. Pharm. Bull.* **25**, 853–856.
- Olea, N., Pulgar, R., Perez, P., Olea-Serrano, F., Rivas, A., Novillo-Fertrell, A., Pedraza, V., Soto, A. M., and Sonnenschein, C. (1996). Estrogenicity of resin-based composites and sealants used in dentistry. *Environ. Health Perspect.* **104**, 298–305.
- Rego, A. C., and Oliveira, C. R. (2003). Mitochondrial dysfunction and reactive oxygen species in excitotoxicity and apoptosis: Implications for the pathogenesis of neurodegenerative diseases. *Neurochem. Res.* **28**, 1563–1574.
- Reinhechel, T., Wiswedel, I., Noack, H., and Augustin, W. (1995). Electrophoretic evidence for the impairment of complexes of the respiratory chain during iron/ascorbate induced peroxidation in isolated rat liver mitochondria. *Biochim. Biophys. Acta* **1239**, 45–50.
- Schonfelder, G., Wittfoht, W., Hopp, H., Talsness, C. E., Paulk, M., and Chahoud, I. (2002). Parent bisphenol A accumulation in the human maternal-fetal-placental unit. *Environ. Health Perspect.* **110**, A703–A707.
- Setsukinai, K., Urano, Y., Kakinuma, K., Majima, H. J., and Nagano, T. (2003). Development of Novel Fluorescence Probes That Can Reliably Detect Reactive Oxygen Species and Distinguish Specific Species. *J. Biol. Chem.* **278**, 3170–3175.
- Shen, J., and Cookson, M. (2004). Mitochondria and dopamine. New insights into recessive parkinsonism. *Neuron* **43**, 301–304.
- Shendelman, S., Jonason, A., Martinat, C., Leete, T., and Abeliovich, A. (2004). DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol.* **2**, e362.
- Shinbo, Y., Taira, T., Niki, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2005a). DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int. J. Oncol.* **26**, 641–648.
- Shinbo, Y., Niki, T., Taira, T., Ooe, H., Takahashi-Niki, K., Maita, C., Seino, C., Iguchi-Ariga, S. M. M., and Ariga, H. (2005b). Proper SUMO-1 conjugation is essential to DJ-1 to exert its full activities. *Cell Death Diff.* (in press).
- Stohs, S. J., Alsharif, N. Z., Shara, M. A., al-Bayati, Z. A., and Wahba, Z. Z. (1991). Evidence for the induction of an oxidative stress in rat hepatic mitochondria by 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Adv. Exp. Med. Biol.* **283**, 827–831.
- Suzuki, T., Mizuo, K., Nakazawa, H., Funae, Y., Fushiki, S., Fukushima, S., Shirai, T., and Narita, M. (2003). Prenatal and neonatal exposure to bisphenol-A enhances the central dopamine D1 receptor-mediated action in mice: Enhancement of the methamphetamine-induced abuse state. *Neuroscience* **117**, 639–644.
- Taira, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2004a). Co-localization with DJ-1 is essential for the androgen receptor to exert its transcription activity that has been impaired by androgen-antagonists. *Biol. Pharm. Bull.* **27**, 574–577.
- Taira, T., Saito, Y., Niki, T., Iguchi-Ariga, S. M. M., Takahashi, K., and Ariga, H. (2004b). DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep.* **5**, 213–218.
- Takahashi, K., Taira, T., Niki, T., Seino, C., Iguchi-Ariga, S. M. M., and Ariga, H. (2001). DJ-1 positively regulates the androgen receptor by impairing the binding of PIASx alpha to the receptor. *J. Biol. Chem.* **276**, 37556–37563.
- Takahashi-Niki, K., Niki, T., Taira, T., Iguchi-Ariga, S. M. M., and Ariga, H. (2004). Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients. *Biochem. Biophys. Res Commun.* **320**, 389–397.
- Toyama, Y., Suzuki-Toyota, F., Maekawa, M., Ito, C., and Toshimori, K. (2004). Adverse effects of bisphenol A to spermiogenesis in mice and rats. *Arch. Histol. Cytol.* **67**, 373–381.
- Tretter, L., Sipos, I., and Adam-Vizi, V. (2004). Initiation of neuronal damage by complex I deficiency and oxidative stress in Parkinson's disease. *Neurochem. Res.* **29**, 569–577.
- Wagenfeld, A., Yeung, C. H., Shivaji, S., Sundareswaran, V. R., Ariga, H., and Cooper, T. G. (2000). Expression and cellular localization of contraception-associated protein. *J. Androl.* **21**, 954–963.
- Wagenfeld, A., Yeung, C. H., Strupat, K., and Cooper, T. G. (1998). Shedding of a rat epididymal sperm protein associated with infertility induced by ornidazole and alpha-chlorohydrin. *Biol. Reprod.* **8**, 1257–1265.
- Welch, J. E., Barbee, R. R., Roberts, N. L., Suarez, J. D., and Klinefelter, G. R. (1998). SP22: A novel fertility protein from a highly conserved gene family. *J. Androl.* **19**, 385–393.
- Yokota, H., Iwano, H., Endo, M., Kobayashi, T., Inoue, H., Ikushiro, S., and Yuasa, A. (1999). Glucuronidation of the environmental oestrogen bisphenol A by an isoform of UDP-glucuronosyltransferase, UGT2B1, in rat liver. *Biochem. J.* **340**, 405–409.
- Yokota, T., Sugawara, K., Ito, K., Takahashi, R., Ariga, H., and Mizusawa, H. (2003). Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition. *Biochem. Biophys. Res. Commun.* **312**, 1342–1348.
- Yoshida, K., Sato, Y., Yoshiike, M., Nozawa, S., Ariga, H., and Iwamoto, T. (2003). Immunocytochemical localization of DJ-1 in human male reproductive tissue. *Mol. Reprod. Dev.* **66**, 391–397.
- Zhang, L., Shimoji, M., Thomas, B., Moore, D. I., Yu, S.-W., Marupudi, N. I., Torp, R., Torgner, I. A., Ottersen, O. P., Dawson, T. M., et al. (2005). Mitochondrial localization of the Parkinson's disease related protein DJ-1: Implications for pathogenesis. *Hum. Mol. Genet.* **14**, 2063–2073.

# Proper SUMO-1 conjugation is essential to DJ-1 to exert its full activities

Y Shinbo<sup>1</sup>, T Niki<sup>2</sup>, T Taira<sup>1,2</sup>, H Ooe<sup>1,2</sup>, K Takahashi-Niki<sup>2</sup>, C Maita<sup>2</sup>, C Seino<sup>1</sup>, SMM Iguchi-Ariga<sup>2,3</sup> and H Ariga<sup>\*,1,2</sup>

<sup>1</sup> Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo, Japan

<sup>2</sup> CREST, Japan Science and Technology Corporation, Kawaguchi, Saitama, Japan

<sup>3</sup> Graduate School of Agriculture, Hokkaido University, Sapporo, Japan

\* Corresponding author: H Ariga, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-ku, Sapporo 060-0812, Japan.

Tel: +81-11-706-3745; Fax: +81-11-706-4988;

E-mail: hiro@pharm.hokudai.ac.jp

Received 21.2.05; revised 19.4.05; accepted 03.5.05; published online 24.6.05

Edited by RA Knight

## Abstract

DJ-1 is a multifunctional protein that plays roles in transcriptional regulation and antioxidative stress, and loss of its function is thought to result in the onset of Parkinson's disease (PD). Here, we report that DJ-1 was sumoylated on a lysine residue at amino-acid number 130 (K130) by PIASx $\alpha$  or PIASy. The K130 mutation abrogated all of the functions of DJ-1, including *ras*-dependent transformation, cell growth promotion and anti-UV-induced apoptosis activities. Sumoylation of DJ-1 was increased after UV irradiation concomitant with a pI shift to an acidic point of DJ-1. Furthermore, L166P, a mutant DJ-1 found in PD patients, and K130RX, an artificial mutant containing four mutations in DJ-1, were improperly sumoylated, and they became insoluble, partly localized in the mitochondria and degraded by the proteasome system. Both L166P-expressing cells and DJ-1-knockdown cells were found to be highly susceptible to UV-induced cell apoptosis.

*Cell Death and Differentiation* (2006) 13, 96–108.

doi:10.1038/sj.cdd.4401704; published online 24 June 2005

**Keywords:** DJ-1; SUMO-1 conjugation; apoptosis; oxidative stress; cell transformation; cell growth

**Abbreviations:** AR, androgen receptor; PD, Parkinson's disease

## Introduction

DJ-1 was first identified by our group as a novel candidate of the oncogene product that transformed mouse NIH3T3 cells in cooperation with activated *ras*.<sup>1</sup> The human *DJ-1* gene is mapped at chromosome 1p36.2–p36.3, where a hot spot of chromosome abnormalities has been reported in several tumors.<sup>2</sup> However, the mechanism by which cells are transformed has not been clarified. Another group has also identified RS, another name for human DJ-1, as a regulatory component of an RNA-binding protein complex.<sup>3</sup> Further-

more, CAP-1 or SP22, a rat homologue of human DJ-1, has been identified by other laboratories as a key protein related to infertility of male rats that were exposed to sperm toxicants such as ornidazole and epichlorohydrin. DJ-1 in the sperm and epididymis decreased in parallel with the subsequent infertility of the rats and was later found to be related to infertility.<sup>4–9</sup> It has also been shown that DJ-1 is a circulating tumor antigen in breast cancer, in which DJ-1 is secreted from cells to serum,<sup>10</sup> and that DJ-1 is overexpressed in smoker-derived lung adenocarcinoma.<sup>11</sup> DJ-1 is therefore thought to play roles in both somatic cells and sperm.

We have shown that DJ-1 acts as a positive regulator for the androgen receptor (AR) by sequestering PIASx $\alpha$  or DJBP.<sup>12–14</sup> PIASx $\alpha$ , a member of the PIAS family of proteins, and DJBP, a novel DJ-1-binding protein, function as negative regulators for AR by preventing AR DNA-binding activity and by recruiting histone deacetylase complex, respectively.<sup>12,13</sup> We have also showed that DJ-1 bound to p53 to restore p53 transcription activity inhibited by Topors/p53BP3 and that transforming activity of DJ-1 was stimulated by Abstrakt, another DJ-1-binding protein.<sup>15,16</sup>

Deletion and point (L166P) mutations of *DJ-1* have recently been shown to be responsible for the onset of familial Parkinson's disease (PD), PARK7,<sup>17</sup> and other homozygous and heterozygous mutations of *DJ-1* have been identified in patients with familial or sporadic PD.<sup>18–20</sup> PD involves an irreversible degeneration of the dopaminergic nigrostriatal pathway. Genes responsible for rare familial early-onset PD, including  $\alpha$ -synuclein,<sup>21</sup> Parkin<sup>22</sup> and UCH-L1,<sup>23</sup> have been identified, and they are thought to play a role in ubiquitin–proteasome dysfunction in PD. L166P DJ-1 has been reported to be degraded, in part, by the ubiquitin–proteasome system.<sup>24–28</sup> DJ-1 was shown to be expressed in almost all brain tissues in healthy men and patients with neurodegenerative diseases, including PD, Pick's disease and multiple system atrophy.<sup>29–31</sup> Various lines of evidence also suggest that oxidative stresses contribute to the cascade leading to dopaminergic cell degeneration in PD,<sup>32–35</sup> and expression of DJ-1 was shown to be induced by oxidative stresses.<sup>36–39</sup> A pI shift of DJ-1 towards a more acidic isoform has been observed in PD patients.<sup>29</sup> We previously reported that DJ-1 plays a role in the antioxidative stress reaction, in which reactive oxygen species (ROS) were eliminated *in vitro* and *in vivo* by oxidizing DJ-1 itself, and that mutations of DJ-1, including L166P, lead to cell death.<sup>40,41</sup> Recently, PINK1, a causative gene product of *PARK6*, has also been suggested to play a role in an antioxidative stress response,<sup>42</sup> but precise roles of PINK1 are not clarified.

SUMO-1 conjugation to lysine residues of proteins is catalyzed in three steps by E1, E2 and E3 enzymes and modulates interactions, localization, activity and stability of proteins to be sumoylated (see recent reviews and references therein).<sup>43,44</sup> E3 SUMO-1 ligases promote attachment of the SUMO modifier to target proteins, and some SUMO-1 ligases have been identified. PIAS family proteins, which include

PIAS1, PIAS3, PIAS $\alpha$ , PIAS $\beta$  and PIAS $\gamma$ , were initially identified as negative regulators of STAT-transcription factors and later found to function as transcriptional coregulators in various other cellular pathways, including Wnt signaling, the p53 pathway and steroid hormone signaling. PIAS proteins were then found to act as E3 SUMO-1 ligases that stimulate SUMO-1 conjugation to proteins, including p53,<sup>45-47</sup> AR,<sup>47</sup> Jun<sup>46</sup> and Lef-1.<sup>48</sup> In most cases, the PIAS/SUMO pathway appears to be an important mechanism of transcriptional regulation.

In this study, we found that mutation of lysine at amino-acid number 130 of DJ-1 abrogated all of the functions of DJ-1 and that improperly sumoylation to L166P mutant DJ-1 makes DJ-1 insoluble, which may trigger the onset of PD.

## Results


### SUMO-1 conjugation of DJ-1 in cells

We have reported that DJ-1 binds to PIAS $\alpha$  and regulates the AR transcription activity.<sup>12</sup> In addition to PIAS $\alpha$ , both Ubc9 and SUMO-1 were identified as DJ-1-binding proteins by a yeast two-hybrid screening. Ubc9 is an E2 enzyme in the SUMO-1 conjugation system. These findings indicate the possibility of sumoylation of DJ-1. To test this possibility, human H1299 cells were lysed in SDS-containing buffer, boiled, and immunoprecipitated with an anti-DJ-1 antibody or nonspecific IgG. Precipitates were analyzed by Western blotting with an anti-DJ-1 antibody or an anti-SUMO-1 antibody (Figure 1). In addition to a 26-kDa band corresponding to DJ-1, a 44 kDa band appeared in the immunoprecipitate with the anti-DJ-1 antibody but not with IgG (Figure 1, lanes 2 and 1, respectively). Since SUMO-1 conjugation to proteins is known to give an addition of approximately 20 kDa to proteins in SDS-polyacrylamide gel and since the 44-kDa band reacted with the anti-SUMO-1 antibody, the results clearly showed that DJ-1 was sumoylated *in vivo* in H1299 cells (Figure 1, lane 4).

Then EST clones for cDNAs encoding DJ-1 homologs of chicken, fish (Atlantic salmon) and *Drosophila* were obtained,

and monkey DJ-1 cDNA was constructed by RT-PCR using total RNA from monkey COS1 cells as a template. A search for amino-acid sequences for DJ-1 homologs of *Xenopus*, nematode and *Drosophila* were carried out in the NIH-NCI Database. Interestingly, two homologs of DJ-1 were found to be present both in nematode and *Drosophila*. Amino-acid sequences of DJ-1s were found to be highly conserved among species, and the identities of the sequences of monkey, mouse, rat, chicken, *Xenopus*, fish and *Drosophila* to that of human DJ-1 are 98, 91, 91, 89, 78, 80, 52 (nematode 1), 44 (nematode 2), 52 (*Drosophila* 1) and 58% (*Drosophila* 2), respectively (Figure 2a). Of 16 lysines, candidates for the sumoylated amino acid in human DJ-1, the lysine at amino-acid number 130 was found to be conserved among all of the species (Figure 2a). We therefore converted all of the lysines of human DJ-1 to arginine and tested the SUMO-1 conjugation to these mutants of DJ1s in H1299 cells into which FLAG-DJ-1, HA-Ubc9 and T7-SUMO-1 were cotransfected. At 48 h after transfection, the cells were lysed in SDS-containing buffer, boiled, and immunoprecipitated with an anti-FLAG antibody. Precipitates were analyzed by Western blotting with the anti-FLAG antibody (Figure 2b). The results showed that wild-type DJ-1 and all of the arginine-substituted mutants except for K130R were sumoylated, indicating that K130 is a major sumoylation site of DJ-1 (Figure 2, lane 12). It was noted that both the K99R mutant and its sumoylated form of DJ-1 run slowly in gel, suggesting that this mutant possesses a conformation different from that of wild-type DJ-1 (Figure 2, lane 10).

We previously reported that DJ-1 binds to both PIAS $\alpha$  and PIAS $\gamma$ ,<sup>12</sup> both of which are known to function as an E3 SUMO-1 ligase. We therefore tested whether PIAS $\alpha$  and PIAS $\gamma$  are E3 SUMO-1 ligases for DJ-1. H1299 cells were cotransfected with FLAG-DJ-1, HA-Ubc9, T7-SUMO-1 and various amounts of HA-PIAS $\alpha$  or -PIAS $\gamma$  and SUMO-1 conjugation to DJ-1 was examined. Expressions of HA-PIAS $\alpha$  and -PIAS $\gamma$  were first confirmed by blotting the precipitates with an anti-HA antibody (Figure 3a, lanes 4-9, 11 and 12 in lower part). The results showed that while the HA-tag alone did not stimulate sumoylation of DJ-1, both HA-PIAS $\alpha$  and -PIAS $\gamma$  stimulated



**Figure 1** SUMO-1 conjugation to DJ-1 *in vivo*. Proteins were extracted from H1299 cells, immunoprecipitated with an anti-DJ-1 polyclonal antibody, and blotted with an anti-DJ-1 monoclonal antibody or anti-SUMO-1 polyclonal antibody as described in Materials and Methods



**Figure 2** Alignment of deduced amino-acid sequences of DJ-1s from human to *Drosophila* and identification of the sumoylation site on DJ-1. (a) Accession numbers for amino-acid sequences are D61380 (human), AB073863(monkey), AB015652 (mouse), AJ007291 and AF157511 (rat), AB076264 (chicken), AF394958 (*Xenopus*), AB079598 (Atlantic salmon), AAB37889 (nematode 1), AAK18904 (nematode 2), AB079599 (*Drosophila* 1) and AAF58316 (*Drosophila* 2). The identical amino acids are drawn in gray, and the lysines are indicated by dots. Lysines corresponding to lysine at amino-acid number 130 of human DJ-1 are boxed. (b) H1299 cells were transfected with FLAG-DJ-1 and lysine-substituted mutants of DJ-1. At 48 h after transfection, proteins extracted from transfected cells were immunoprecipitated with an anti-FLAG antibody and blotted with the anti-FLAG antibody as described in Materials and Methods

sumoylation of wild-type DJ-1 but not K130R in a dose-dependent manner (Figure 3a), indicating that both PIAS $\alpha$  and PIAS $\gamma$  work as E3 SUMO-1 ligases for DJ-1. To examine


the effect of PIAS $\alpha$  on sumoylation to endogenous DJ-1, H1299 cells were cotransfected with PIAS $\alpha$ -HA, T7-SUMO-1 and Ubc9-HA, and SUMO-1 conjugation to DJ-1 was



examined. As in the case of the ectopic expression system shown in Figure 3a, endogenous DJ-1 was sumoylated by PIASxα in a dose-dependent manner (Figure 3b, lanes 2–5), and similar results were obtained when PIASy was cotransfected into H1299 cells (data not shown). These results clearly indicate that both PIASxα and PIASy are E3 SUMO-1 ligases for DJ-1.

### Abrogation of DJ-1 activities by K130 mutation


To know the importance of K130 for DJ-1 activities, transformation activities of several lysine-substitution mutants of DJ-1 were tested in rat 3Y1 cells by cotransfection of DJ-1 mutants with activated *ras* (Figure 4a). As previously reported in mouse NIH3T3 cells,<sup>1</sup> wild-type DJ-1 gave a number of transformed foci only in the case of cotransfection with activated *ras*. While DJ-1 mutants of both K62, 63R, in which lysines at #62 and 63 were changed to arginine, and K93R still gave numbers of foci similar to that given by wild-type DJ-1, DJ-1 mutants of K130R and K130RX, in which three other amino acids, S57R, E96G and H126Y, had been changed,<sup>12</sup> as well as K130 gave smaller numbers of foci than that



**Figure 3** Stimulation of SUMO-1 conjugation to DJ-1 by PIASxα and PIASy. (a) H1299 cells were cotransfected with FLAG-DJ-1, T7-SUMO-1, Ubc9-HA and various amounts of PIASxα-HA or PIASy. At 48 h after transfection, proteins extracted from transfected cells were immunoprecipitated with an anti-FLAG antibody and blotted with the anti-FLAG antibody or an anti-HA antibody as described in Materials and Methods. The amounts of expression vectors for PIASxα-HA and PIASy were 2.5, 5.0 and 7.5 μg in lanes 4 and 7, 5 and 8, and 6 and 9, respectively. (b) H1299 cells were cotransfected with T7-SUMO-1, Ubc9-HA and various amounts of PIASxα-HA. At 48 h after transfection, proteins extracted from transfected cells were immunoprecipitated with an anti-DJ-1 antibody and analyzed by Western blotting with the anti-DJ-1 antibody or an anti-SUMO-1 antibody as described in Materials and Methods. The amounts of expression vectors for PIASxα-HA were 2.5, 5.0 and 7.5 μg in lanes 3, 4 and 5, respectively

without activated *ras*. These results clearly indicate that K130 in DJ-1 is of functional importance for the transforming activity of DJ-1.

Many proteins possessing transforming activity often have cell growth-stimulating activity. To test this activity of DJ-1, rat 3Y1 cells were transfected with the expression vectors for wild-type and mutant forms of DJ-1 that were cloned in pcDNA3 containing a neomycin-resistant gene, and they were cultured in a medium containing G418. G418-resistant colonies were then counted 14 days after transfection (Figure 4b). Transfection with c-Myc as a positive control gave a large number of colonies, while transfection with the vector alone or no-DNA control (none) gave few or no colonies. Wild-type or non-K130 mutants of DJ-1, K62, 63R and K93R, gave similar numbers of colonies to those of c-Myc. K130R and K130RX mutants of DJ-1, on the other hand, abolished the colony-forming activity to the level of the vector alone. These results indicate that DJ-1 possesses cell growth-promoting activity that is dependent upon K130.



**Figure 4** Effects of lysine mutations of DJ-1 on the transforming and cell growth promoting activities of DJ-1. (a) Rat 3Y1 cells were transfected with expression vectors for activated wild-type DJ-1 or mutants of DJ-1 with or without activated H-ras. In total, 14 days after transfection, the transformed cell foci were counted. (b) Rat 3Y1 cells were transfected with neomycin-resistant gene-containing expression vectors for wild-type DJ-1 or mutants of DJ-1, and they were cultured in the presence of G418. At 14 days after transfection, the G418-resistant cell colonies were counted. Pictures of the colonies are shown

### Induction of DJ-1 expression and stimulation of SUMO-1 conjugation to DJ-1 by UV irradiation

Expression of DJ-1 was induced in various cells that had been subjected to treatment with paraquat, endotoxin or iron, which yielded ROS.<sup>36–39</sup> We also showed that expression of DJ-1 was induced in human neuroblastoma SH-SY5Y cells treated with hydrogen peroxide.<sup>41</sup> Since ultraviolet (UV) irradiation is known also to induce ROS, we first examined whether DJ-1 expression is induced after 20 J/m<sup>2</sup> UV irradiation by using p53-positive and p53-negative cell lines, ME180 and H1299 cells, respectively, in which p53 is a positive control as an induced protein against UV irradiation (Figure 5A). In ME180 cells, while expression of p53 was induced at 2 h after UV irradiation, expression of DJ-1 was induced at 15 min after UV irradiation and the level peaked at 30–45 min and then gradually decreased up to 3 h after UV irradiation (Figure 5A-a). In H1299 cells, induction of DJ-1 expression started at 15 min after UV irradiation and the level peaked at 30–45 min and then gradually decreased until 6 h after UV irradiation (Figure 5A-b). These results suggest that expression of DJ-1 is induced by UV irradiation in a p53-independent manner.

Shifts of the isoelectric point of DJ-1 in cells caused by oxidative stresses have been reported.<sup>36–41</sup> H1299 cells were irradiated with 20 and 50 J/m<sup>2</sup> UV, and proteins prepared from cells were separated on isoelectric focusing gel over the pI range of 5–8 (Figure 5B). After blotting with an anti-DJ-1 antibody, pI of DJ-1 was found to be shifted to a more acidic point, from 6.3 to 5.2, at 1 h after UV irradiation (Figure 5B). SUMO-1 conjugation to DJ-1 was also found to be stimulated at 25 min, peak at 30 min and then gradually decreased after UV irradiation (Figure 5C). These results suggest that DJ-1 is first sumoylated and then oxidized after UV irradiation.

ME180 cells after UV irradiation were stained with a rabbit anti-DJ-1 polyclonal antibody, reacted with an FITC-conjugated anti-rabbit IgG, and visualized under a confocal laser microscope (Figure 5D). Nuclei and the mitochondria in the cytoplasm were also stained with DAPI and MitoTracker, respectively. Endogenous DJ-1 was found to be localized mainly in the cytoplasm and sparsely in the nucleus without UV irradiation (Figure 5D, time 0 h). After UV irradiation, almost all the DJ-1 were translocated to the nucleus, which was shown in the merged figures as light blue color (Figure 5D, time 0.25–0.75 h). At 1 h after UV irradiation, some DJ-1 were relocalized in the cytoplasm and almost all the DJ-1 were located in the cytoplasm at 24 h after UV irradiation. These results indicate that a portion of DJ-1 is shuttled in between the cytoplasm and nucleus of UV-irradiated H1299 cells.


### Improper sumoylation and insoluble form of L166P mutant of DJ-1 found in PD patients

Homozygous point mutations of the *DJ-1* gene have been found in Italian kindred of PD patients,<sup>17</sup> and its gene product L166P mutant form of DJ-1 has been reported to be unstable and to be degraded, in part, by the ubiquitin–proteasome system.<sup>24–28,49</sup> We therefore examined the sumoylation status of L166P DJ-1 (Figure 6). H1299 cells were transfected with FLAG-tagged wild type and three mutants of DJ-1


together with SUMO-1 and Ubc9, and protein extracts from transfected cells were immunoprecipitated with an anti-FLAG antibody. Precipitates were then separated on two gels and analyzed by Western blotting with the anti-FLAG antibody or an anti-SUMO-1 antibody (Figure 6A). Sumoylation of wild-type DJ-1 but not K130R DJ-1 was first confirmed (Figure 6A-a, lanes 1, 2, 6 and 7). Characteristics of the band of 62 kDa observed above the DJ-1-SUMO-1 bands are not clear at present. It is interesting that L166P and K130RX showed ladder bands over the range of SUMO-1-DJ-1 in a blot with the anti-FLAG antibody and that these bands also reacted with the anti-SUMO-1 antibody (Figure 6A-a, lanes 8, 9, 3 and 4), indicating that L166P and K130RX mutants of DJ-1 are conjugated with multi- or poly-SUMO-1, or alternatively that these mutant proteins became misfolded, resulting in the formation of the abnormal aggregates. Since a lysine residue at amino-acid number 130 (K130) in K130RX DJ-1 was changed to arginine, it is possible that sumoylation occurs on lysine residue(s) other than K130 in K130RX DJ-1 and that this is also true for sumoylation of L166P DJ-1. To examine this possibility, K130 in L166P DJ-1 was changed to arginine and sumoylation assay of this mutant (K130R, L166P DJ-1) was carried out. This mutant showed ladder bands as did K130RX and L166P (Figure 6A-a, lanes 5 and 10), indicating that sumoylation occurred some lysine residues, including L130, in L166P DJ-1. This improper sumoylation of L166P and K130RX mutants of DJ-1 was also observed in an *in vitro* reconstituted system using recombinant proteins comprised of SAE1/SAE2 complex, Ubc9 and SUMO-1 (Figure 6A-b, lanes 3 and 4), suggesting that the structure of L166P and K130RX mutants of DJ-1 possess easier accessibility to be sumoylated than does wild-type DJ-1 and was changed to be improperly sumoylated.

We then examined the solubility of wild-type DJ-1 and mutants of DJ-1 in cells. Human 293T cells were transfected with FLAG-tagged wild-type DJ-1 and four mutants of DJ-1, L166P, K130R, K130RX and C106S. An amino acid at C106 is a putative active center for protease activity of DJ-1, and a C106S mutant had no protease and antioxidative stress activities.<sup>28,41,50–52</sup> At 48 h after transfection, the soluble proteins were extracted with 0.1% NP-40 and insoluble proteins were further dissolved in 2.0% SDS, and these proteins were analyzed by Western blotting with an anti-FLAG antibody (Figure 6B). While wild-type DJ-1, C106S DJ-1 and K130R DJ-1 were in soluble forms, both L166P and K130RX were found to be in insoluble forms. Since insoluble forms of DJ-1 have been observed in patients with degenerative neuronal diseases, it is interesting that improperly sumoylated DJ-1 becomes insoluble (see Discussion).

We then examined the cellular localization of these mutants of DJ-1. H1299 cells were transfected with FLAG-tagged wild-type DJ-1 and four mutants of DJ-1. At 48 h after transfection, cells were stained with an anti-FLAG antibody and visualized with an FITC-conjugated anti-mouse antibody (Figure 7). Since L166P DJ-1 has been reported to be localized, in part, in the mitochondria<sup>17</sup> and we have reported that DJ-1 also played a role in antiendoplasmic reticulum (ER) stress-induced cell death,<sup>40</sup> cells were also reacted with MitoTracker and an anti-GRP78 to see the mitochondria and ER, respectively (Figures 7a and data not shown, respectively).



**Figure 5** Induction of expression, pI shift and stimulation of sumoylation of DJ-1. (A) ME180 cells (a) and H1299 cells (b) were irradiated with 20 J/m<sup>2</sup> of UV, and at various times after irradiation, proteins extracted from cells were blotted with an anti-p53 (FL-393, Santa Cruz), anti-DJ-1 polyclonal and anti-actin antibodies (upper panel). Bands that reacted with the antibody were quantitated using a densitometer (lower panel). (B) H1299 cells were irradiated with 20 and 50 J/m<sup>2</sup> of UV, and proteins in the extracts were then analyzed by isoelectric focusing phoresis gel as described in Materials and Methods. (C) H1299 cells were irradiated with 20 J/m<sup>2</sup> of UV. At various times after irradiation, proteins extracted from cells were immunoprecipitated with an anti-DJ-1 polyclonal antibody and blotted with an anti-DJ-1 monoclonal antibody or anti-actin antibody as described in Materials and Methods. (D) ME180 cells were irradiated with 20 J/m<sup>2</sup> of UV. At various times after irradiation, cells were stained with an anti-DJ-1 polyclonal antibody. The cells were then reacted with a FITC-conjugated anti-rabbit IgG and observed under a confocal laser fluorescent microscope. The cells were also stained with DAPI and MitoTracker-red (Molecular probe)




**Figure 6** Improper sumoylation of L166P mutant of DJ-1. (A) H1299 cells were cotransfected with FLAG-DJ-1 or mutants, T7-SUMO-1 and Ubc9-HA. At 48 h after transfection, proteins extracted from transfected cells were immunoprecipitated with an anti-FLAG antibody and analyzed by Western blotting with the anti-FLAG antibody or an anti-SUMO-1 antibody as described in Materials and Methods. (b) *In vitro* sumoylation of DJ-1 and its mutant were carried out using the reaction mixture containing <sup>35</sup>S-DJ-1 and its mutated DJ-1 and the recombinant proteins of E1 enzyme, Ubc9 and SUMO-1 as described in Materials and Methods. Labeled proteins in the mixture were separated on polyacrylamide gel and visualized by fluorography. (B) 293T cells were transfected with FLAG-DJ-1 or its mutants. At 48 h after transfection, soluble proteins were first extracted from transfected cells with 0.1% SDS-containing buffer and insoluble proteins were then dissolved with 2% SDS. These were analyzed by Western blotting with an anti-FLAG antibody

Nuclei were also stained with DAPI. Wild-type DJ-1 and K130R DJ-1 were localized both in the cytoplasm and nucleus, and some portion of DJ-1 were also localized in the mitochondria as shown by merged figures in which the green and red have become yellow (Figure 7a, k and l). K130RX, L166P DJ-1 and a double mutant (K130R, L166P) were, on the other hand, found to be predominantly localized in the cytoplasm and almost all of them were localized in the mitochondria (Figure 7a, m, n and o). Neither wild-type DJ-1 nor any of the four mutants of DJ-1, on the other hand, was found to be localized in the ER in cells without ER stress (data not shown). Wild-type DJ-1 and the four mutants of DJ-1 were also not found to be localized in the Golgi apparatus (data not shown).

Since L166P DJ-1 has been reported to be degraded, in part, by the ubiquitin-proteasome system<sup>24-28,49</sup> and DJ-1 possess cysteine protease activity,<sup>28</sup> we then examined the

stability of mutants of DJ-1 against degradation by proteasome and cysteine protease using their specific inhibitors, MG132 and E64, respectively. Proteins were extracted from NIH3T3 cells transfected with FLAG-tagged wild-type DJ-1 and mutants of DJ-1 in the presence or absence of MG132 or E64 and blotted with an anti-FLAG antibody or an antiactin antibody (Figure 8). The results first showed that the amounts of L166P and K130RX DJ-1 were smaller than those of wild-type DJ-1 and K130R DJ-1 in the absence of MG132 (Figure 8, lanes 1, 4, 7, 10, 13 and 16) and that E64 had no effect on the stability of wild-type DJ-1 or any of the mutants of DJ-1 (Figure 8, lanes 3, 6, 9, 12, 15 and 18). Although wild-type DJ-1 and K130R DJ-1 were not affected by MG132, the amounts of L166P and K130RX increased partially in the presence of MG132 (Figure 8, lanes 14 and 17 in the middle part), suggesting that K130RX and L166P DJ-1 were degraded, in part, by the proteasome system. It was noted that



**Figure 7** Localization of DJ-1 and its mutants in cells. H1299 cells were transfected with FLAG-wild-type (Wt) or its mutants. At 48 h after transfection, cells were fixed, reacted with a combination of an anti-FLAG antibody (a–e), DAPI and MitoTracker (f–j), and visualized with a fluorescein isothiocyanate-conjugated anti-mouse antibody (a–e) and self-fluorescences of DAPI and MitoTracker (f–j). The two figures have been merged (k–o)

slow-migrating bands of K130RX DJ-1 appeared in the presence of MG132 (Figure 8, lane 17 in the upper part). Although it is possible that these bands correspond to the ubiquitinated K130RX DJ-1, it is not certain at present due to the inability of identification of these bands against an antiubiquitin antibody (data not shown). These results suggest that mutants of DJ-1 localized in the mitochondria are improperly sumoylated, insoluble and unstable.

### Loss of antiapoptosis activity of DJ-1 mutants

We then examined the effects of K130R and L166P mutations of DJ-1 on UV-induced apoptosis. Mouse NIH3T3-Wt, -K130R and -L166P cells, which are stable cell lines expressing wild-type, K130R and L166P DJ-1 as described previously,<sup>41</sup> and NIH3T3-D2 cells, which are stable DJ-1-knockdown cells expressing siRNA toward DJ-1,<sup>49</sup> were irradiated with two doses of UV. At 22 h after irradiation, DNA from cells was stained with propidium iodide and the distribution of cells in the cell cycle was analyzed by flow cytometry (Figure 9). DJ-1 expression in D2 cells decreased to about 10% of that of parental NIH3T3 cells (Figure 9a) and the characteristics of D2 cells has been described.<sup>49</sup> Apoptotic cells that appeared at the position below the G1/G0 phase were counted using the 'ModiFit LT' program (Figure 9b). As in the case of hydrogen peroxide-induced cell death,<sup>41</sup> cells harboring exogenously added wild-type DJ-1 were much more resistant to UV than were parental nontransfected cells (host). Of the cell lines harboring mutants, on the other hand, the cell line harboring L166P was found to be most sensitive to UV and the K130R line followed in terms of sensitivity to UV. DJ-1-knockdown cells were found to have similar sensitivity to that of L166P cells.


These results clearly indicate that DJ-1 plays a role in protection against UV-induced apoptosis and that mutations or knockdown of DJ-1 lead to cell death.

## Discussion


### SUMO-1 conjugation to DJ-1

In this study, we first found that a lysine at amino-acid number 130 of DJ-1 (K130) was sumoylated by PIAS $\alpha$  and PIAS $\gamma$  in cells and that this sumoylation of DJ-1 was essential for DJ-1 to exert its full activities, including *ras*-dependent transforming and cell growth-stimulating activities. PIAS family proteins, which include PIAS1, PIAS3, PIAS $\alpha$ , PIAS $\beta$  and PIAS $\gamma$ , are known to function as E3 SUMO-1 ligases and the proteins to be sumoylated by each PIAS family protein were identified.<sup>43,44</sup> In most case, transcription factors have been identified as target proteins for PIAS family proteins and DJ-1 is also a transcription factor as a coactivator to regulate expressions of genes for AR, p53 and PTEN.<sup>12,13,15,53,54</sup> Since identifications of almost all the proteins that had been sumoylated were performed by overexpression system using expression vectors for SUMO-1 and Ubc9, the level of sumoylation of DJ-1, which was observed without overexpression of SUMO-1 or Ubc9, is thought to be high.

DJ-1 expression and its sumoylation were induced by UV irradiation and DJ-1 was translocated from the cytoplasm to nucleus. Since induction of DJ-1 expression after irradiation of UV occurred both in p53 (–) and p53 (+) cells and this occurred faster than did induction of p53 expression in p53 (+) cells, induction of DJ-1 expression by UV irradiation is p53-independent. UV irradiation is known to induce ROS and



**Figure 8** Stability of DJ-1 and its mutants. Mouse NIH3T3 cell lines harboring FLAG-wild-type DJ-1 and its mutants were cultured in the presence of 10 mM MG132 or 8  $\mu$ g/ml of E64 for 12 h. Proteins extracted from cells were analyzed by Western blotting with anti-FLAG and antiactin antibodies. Host indicates parental NIH3T3 cells



**Figure 9** Abrogation of antiapoptosis activity of DJ-1 by mutations. (a) Proteins extracted from NIH3T3 cells (host) and DJ-1-knockdown cells (D2) were analyzed by Western blotting with anti-mouse DJ-1 and antiactin antibodies. The proteins that reacted with primary antibodies were visualized with IRDye800-conjugated or Alexa Fluor680-conjugated secondary antibodies using an infrared imaging system (Odyssey, LI-COR). (b) Mouse NIH3T3 cell lines harboring FLAG-wild-type DJ-1 and its mutants were irradiated with 100 and 2000 J/m<sup>2</sup> UV. At 22 h after irradiation, DNA in cells was stained with propidium iodide and the distribution of cells in the cell cycle was analyzed by flow cytometry. Distribution of cells in the cell cycle by analysis of flow cytometry was quantitated by using a 'ModiFit LT' program. Results ( $n=3$ ) were statistically processed using Student's *t*-test. Significance: \* $P < 0.01$  versus apoptotic cells in wt-DJ-1 cells

to give the oxidative stress to cells, and DJ-1 expression has been reported to be induced by several chemicals that had also been known to induce ROS.<sup>36-39</sup> UV irradiation of cells also shifted the pI of DJ-1 to acidic points, as in the case of direct addition of hydrogen peroxide, one of the ROS, to human SH-SY5Y and mouse NIH3T3 cells, in which ROS were scavenged by DJ-1, as described previously.<sup>41</sup> The amount of pI-shifted DJ-1, however, was again reduced 4 h after UV irradiation concomitant with a reduction of the amount of the sumoylated form of DJ-1 and relocalization of DJ-1 in the cytoplasm, suggesting that self-oxidation of DJ-1 following SUMO-1 conjugation to DJ-1 occurs depending on

the amounts of ROS in cells and that the sumoylated form of DJ-1 is an active form to eliminate ROS. High-dose of UV irradiation of cells induces apoptosis, and an NIH3T3 cell line expressing exogenously added wild-type DJ-1 became resistant to UV-induced apoptosis compared to the parental cells. This antiapoptotic activity of DJ-1 was abrogated by the mutation into an amino acid K130, a sumoylation site of DJ-1, as well as those of *ras*-dependent transforming and cell growth-stimulating activities, indicating that SUMO-1 conjugation to K130 of DJ-1 is essential for DJ-1 to function.

#### Improper SUMO-1 conjugation to L166P DJ-1, a mutant of DJ-1 found in PD patients

In a latter part of this paper, we described the properties of L166P DJ-1, a mutant of DJ-1 found in PD patients,<sup>17</sup> and K130RX DJ-1, an artificial mutant possessing four mutations including K130R, in terms of sumoylation and loss of function. Both L166LP and K130RX were improperly sumoylated *in vivo* compared to wild-type DJ-1, to which one SUMO-1 molecule was conjugated. It is possible that this improper SUMO-1 conjugation to L166P and K130RX occurs on multiple lysine residues (multisumoylation) or on a specific lysine residue with poly-SUMO-1 (polysumoylation). It is also possible that L166P and K130RX are misfolded, resulting in sumoylation of multiple lysine residues (multisumoylation) and then abnormal aggregation. Although polysumoylation, which had been thought not to occur, has recently been reported,<sup>55,56</sup> it is not clear whether there was excess sumoylation of DJ-1 mutants in either case. Since K130RX lost lysine at 130, a lysine(s) other than K130 was sumoylated. Furthermore, both L166LP and K130RX were also improperly sumoylated in an *in vitro* reconstituted system using recombinant proteins. The crystal structure of DJ-1 has been determined by five groups, including us, and it has been shown that L166 is located in the  $\alpha$ -helix near the C-terminal region of DJ-1 and that L166P mutation might disrupt dimer formation.<sup>57-61</sup> Structure of K130RX was also thought to be

drastically changed by the computer-based analysis. The protective activity of DJ-1 against hydrogen peroxide-induced cell death has been reported to be abrogated by L166P mutation.<sup>41,49</sup> It is therefore thought that the structures of these mutants themselves changed to allow for improper sumoylation or abnormal aggregation. Both L166P and K130RX DJ-1 were found to become insoluble. Recently, insoluble forms of DJ-1 have been reported to be colocalized with tau or  $\alpha$ -synuclein in fibrillar inclusions in patients with Pick's disease and multiple system atrophy, suggesting that DJ-1 is related to various neurodegenerative diseases, including PD.<sup>30,31</sup> It would therefore be interesting to examine whether such insoluble forms of DJ-1 in patients with neurodegenerative diseases are sumoylated.

Localization of K130R DJ-1 in the cytoplasm and nucleus was found to be the same as that of wild-type DJ-1, suggesting that SUMO-1 conjugation is not sufficient to determine the localization of DJ-1 in cells. Some of wild-type DJ-1 and K130R DJ-1 were also found to be localized in the mitochondria. Both L166P and K130RX DJ-1 were then found to be localized in the mitochondria. Since the mitochondrion is a major ROS-production organelle and since low activity levels of the complex 1 that reside in mitochondria have been found in PD patients (see review and references therein),<sup>62</sup> it is interesting that improperly sumoylated DJ-1 was localized in the mitochondria. Since DJ-1 has no mitochondria-import/export sequences, wild-type DJ-1 and mutants of DJ-1, especially L166P and L130RX, may associate with other protein(s) that are targeted in mitochondria. To determine the importance of different localizations of DJ-1 in cells, localization of DJ-1 in specific organelles such as mitochondria in patients with various neurodegenerative diseases, including PD, must be investigated.

K130RX is unstable and degraded, in part, by the ubiquitin-proteasome system, like L166P as described previously.<sup>24-28,49</sup> In the presence of the proteasome inhibitor MG132, a ladder above the position of the K130RX monomer was observed in polyacrylamide gel. This is likely to be a ubiquitinated form of K130RX DJ-1, and further study is needed to confirm this. Wild-type DJ-1 and its mutants were not affected by the presence of E64, an inhibitor of cysteine protease, indicating that DJ-1 is not regulated by its protease activity.

We established DJ-1-knockdown cells (D2 cells) in which siRNA targeting to the mouse DJ-1 gene was expressed under the control of the U6 promoter,<sup>49</sup> and we found that D2 cells were highly susceptible to UV-induced apoptosis, indicating that DJ-1 plays a role in antiapoptotic activity. A cell line that harbors exogenously added L166P DJ-1 had sensitivity against UV-induced apoptosis similar to that of D2 cells. It is therefore thought that L166P DJ-1 acts in a dominant-negative fashion at least in this cell line, in which wild-type DJ-1 is present. In PD patients harboring L166P mutation of the *DJ-1* gene, loss of function of DJ-1 triggers onset of disease.

It has been reported that improper sumoylation was related to the onset of polyglutamine diseases, Alzheimer's disease and Huntington's disease.<sup>63-65</sup> Together with the results of those studies, the results of this study suggest that proper sumoylation is essential for DJ-1 to exert its full activities and

that improper sumoylation makes DJ-1 insoluble, mislocalized and unstable, which trigger the onset of PD.

## Materials and Methods

### Cells

Human H1299, ME180, 293T, HeLa and rat 3Y1 cells were cultured in Dulbecco's modified Eagle's medium with 10% calf serum.

### Construction of lysine mutants of DJ-1

Nucleotide sequences of the oligonucleotide used for PCR primers were as follows: DJ-1-K62R (+), 5'-CTGAAGATGCAAGAAAAGAG-3'; DJ-1-K62R (-), 5'-CATATGGTCCCTCTTTTCTTGC-3'; DJ-1-K62, 63R (+), 5'-CTGAAGATGCAAGAAGAGAG-3'; DJ-1-K62, 63R (-), 5'-CATATGGTCCCTCTTCTTGC-3'; DJ-1-K63R (+), 5'-CTGAAGATGCAAAAAAGAGAG-3'; DJ-1-K63R (-), 5'-CATATGGTCCCTCTTTTCTTGC-3'; DJ-1-K130R (+), 5'-CAACACACCCTCTTGCTAGAGAC-3'; DJ-1-K130R (-), 5'-CATCATTTTGTCTCTAGCAAG-3'; T7, 5'-CCCTATAGTGAGTCGTATTA-3'; Sp6, 5'-TTTAGGTGACACTATAGAAT-3'. The first PCR was carried out on pcDNA3-FLAG-DJ-1 as a template either with DJ-1-K (+) and T7 or with DJ-1K (-) and Sp6 as primers. The products were mixed and used as templates for the second PCR with T7 and Sp6 as primers. The resultant PCR product was digested with *Hind*III and *Xho*I and inserted into the respective site of pcDNA3.

### SUMO-1 conjugation to DJ-1 *in vivo*

Proteins were extracted from human H1299 cells with a buffer containing 0.1% NP-40, 50 mM Tris-HCl (pH 7.5) and 150 mM NaCl and by sonication. Proteins were then immunoprecipitated with an anti-DJ-1 polyclonal antibody-conjugated agarose,<sup>1</sup> and the precipitates were dissolved with a sample buffer containing 30 mM Tris-HCl (pH 6.8), 6% SDS, 0.006% bromophenol blue, 12% 2-mercaptoethanol and 50% glycerol, boiled for 15 min, and blotted with an anti-DJ-1 monoclonal antibody (3E8, specificity described by Yoshida *et al.*<sup>9</sup>) or an anti-SUMO-1 polyclonal antibody (FL-101, Santa Cruz). H1299 cells were transfected with 5  $\mu$ g of FLAG-DJ-1 or its mutants, 2  $\mu$ g of Ubc9-HA and 2  $\mu$ g of T7-SUMO-1 by the calcium phosphate precipitation method. At 48 h after transfection, the proteins extracted from cells were immunoprecipitated with an anti-FLAG monoclonal antibody (M2, Roche), blotted with an anti-T7 monoclonal antibody (Novagen) or the anti-FLAG monoclonal antibody, and visualized by ECL (Amersham BioScience).

### SUMO-1 conjugation to DJ-1 *in vitro*

Recombinant proteins of human SUMO-1, E1 enzyme (SAE1 and SAE2) and Ubc9 were purchased from Alexis Biochemicals. <sup>35</sup>S-FLAG-DJ-1 and its mutant were synthesized *in vitro* using a TnT-Coupled reticulocyte lysate system (Promega). Reaction mixture containing 100 ng of E1 enzyme, 900 ng of Ubc9, 10  $\mu$ g of SUMO-1 and <sup>35</sup>S-FLAG-DJ-1 and its mutant was incubated at 30°C for 2 h, mixed with the sample buffer, and boiled for 5 min. The proteins were then separated on SDS-containing polyacrylamide gel and visualized by fluorography.

### Focus forming and growth assays

Rat 3Y1 cells cultured in a 10-cm dish were transfected with 1  $\mu$ g each of pEJ6.6, pCMV-F-DJ-1 or pCMV-F-DJ-1-mutant by the calcium phosphate

precipitation method, and the medium was changed every 2 days. At 14 days after transfection, the cells were stained with Giemsa solution and the foci that had formed were counted.

Rat 3Y1 cells were transfected with 5  $\mu$ g each of the above plasmids by the calcium phosphate precipitation method and cultured in the presence of 300  $\mu$ g/ml of G418. At 14 days after transfection, the cells were stained with Giemsa solution and the colonies that had formed were counted.

### Indirect immunofluorescence

H1299 cells were transfected with FLAG-DJ-1 or its mutant as described above. At 48 h after transfection, cells were fixed with a solution containing 4% paraformaldehyde and reacted with a combination of a mouse anti-FLAG monoclonal antibody (M2, Sigma), goat anti-GRP78 polyclonal antibody (N-20, Santa Cruz), and MitoTracker (Molecular Probe). The cells were then reacted with an FITC-conjugated anti-mouse IgG or rhodamine-conjugated anti-rabbit IgG and observed under a confocal laser fluorescent microscope.

### Isoelectric focusing

H1299 cells were irradiated with UV using a UV cross linker, and proteins were extracted from the cells at various times after UV-irradiation. Proteins were then separated in the pH 5–8 range of isoelectric focusing phoresis gel or 12.5% polyacrylamide gel containing SDS, transferred onto nitrocellulose membranes, and blotted with an anti-DJ-1 polyclonal antibody or antiactin-antibody (MAB1501R, CHEMICON) as described previously.<sup>41</sup>

### Establishment of a DJ-1-knockdown cell line

The nucleotide sequences of the upper strand of the oligonucleotide used for construction of an siRNA vector targeting mouse DJ-1 gene is 5'-GGATCCGCTTGTCTAGTAGAATAAAGCTTCAAGAGAGTTTATTCTACTA GCAAGGTTTTTCCAAAAGCTT-3'. After annealing oligonucleotides corresponding to the upper and lower strands of DNA, these were inserted into *Bam*HI–*Hind*III sites of pRNA-U6.1/Neo (GenScript Corp.). These plasmids were transfected into mouse Flp-In™3T3 cells (Invitrogen) by the calcium phosphate precipitation method, and the cells were cultured in the medium in the presence of 200  $\mu$ g/ml G418 for 14 days. The cells that were resistant to the drug were then selected, and expression of DJ-1 was examined by Western blotting with an anti-mouse DJ-1 antibody.<sup>8</sup>

### Separation of soluble and insoluble forms of DJ-1

Human 293T cells were transfected with FLAG-DJ-1 or its mutant as described above. At 48 h after transfection, cells were lysed with a buffer containing 0.1% NP-40 and phosphate-buffered saline and by sonication. The supernatant fraction after centrifugation was used as soluble proteins, and the precipitates were further dissolved with a sample buffer containing SDS as described above and used as insoluble proteins. These proteins were blotted with an anti-FLAG antibody.

### Acknowledgements

We thank Yoko Misawa and Kiyomi Takaya for their technical assistance. This work was supported by grants-in-aid from the Ministry of Education, Science, Culture, Sports and Technology of Japan and the Ministry of Health, Labor and Welfare of Japan.

### References

- Nagakubo D, Taira T, Kitaura H, Ikeda M, Tamai K, Iguchi-Arigo SMM and Ariga H (1997) DJ-1, a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. *Biochem. Biophys. Res. Commun.* 31: 509–513
- Taira T, Takahashi K, Kitagawa R, Iguchi-Arigo SMM and Ariga H (2001) Molecular cloning of human and mouse DJ-1 genes and identification of Sp1-dependent activation of the human DJ-1 promoter. *Gene* 263: 285–292
- Hod Y, Pentylala SN, Whyard TC and El-Maghrabi MR (1999) Identification and characterization of a novel protein that regulates RNA-protein interaction. *J. Cell. Biochem.* 72: 435–444
- Wagenfeld A, Yeung CH, Strupat K and Cooper TG (1998) Shedding of a rat epididymal sperm protein associated with infertility induced by ornidazole and alpha-chlorohydrin. *Biol. Reprod.* 8: 1257–1265
- Welch JE, Barbee RR, Roberts NL, Suarez JD and Klinefelter GR (1998) SP22: a novel fertility protein from a highly conserved gene family. *J. Androl.* 19: 385–393
- Wagenfeld A, Yeung CH, Shivaji S, Sundareswaran VR, Ariga H and Cooper TG (2000) Expression and cellular localization of contraception-associated protein. *J. Androl.* 21: 954–963
- Klinefelter GR, Welch JE, Perreault SD, Moore HD, Zucker RM, Suarez JD, Roberts NL, Bobseine K and Jeffay S (2002) Localization of the sperm protein SP22 and inhibition of fertility *in vivo* and *in vitro*. *J. Androl.* 23: 48–63
- Okada M, Matsumoto K, Niki T, Taira T, Iguchi-Arigo SMM and Ariga H (2002) DJ-1, a target protein for an endocrine disrupter, participates in the fertilization in mice. *Biol. Pharm. Bull.* 25: 853–856
- Yoshida K, Sato Y, Yoshiike M, Nozawa S, Ariga H and Iwamoto T (2003) Immunocytochemical localization of DJ-1 in human male reproductive tissue. *Mol. Reprod. Dev.* 66: 391–397
- Le Naour F, Misk DE, Krause MC, Deneux L, Giordano TJ, Scholl S and Hanash SM (2001) Proteomics-based identification of RS/DJ-1 as a novel circulating tumor antigen in breast cancer. *Clin. Cancer Res.* 7: 3328–3335
- Miura K, Bowman ED, Simon R, Peng AC, Robles AI, Jones RT, Katagiri T, He P, Mizukami H, Charboneau L, Kikuchi T, Liotta LA, Nakamura Y and Harris CC (2002) Laser capture microdissection and microarray expression analysis of lung adenocarcinoma reveals tobacco smoking- and prognosis-related molecular profiles. *Cancer Res.* 62: 3244–3250
- Takahashi K, Taira T, Niki T, Seino C, Iguchi-Arigo SMM and Ariga H (2001) DJ1 positively regulates the androgen receptor by impairing the binding of PIAS $\alpha$  to the receptor. *J. Biol. Chem.* 276: 37556–37563
- Niki T, Takahashi-Niki K, Taira T, Iguchi-Arigo SMM and Ariga H (2003) DJBP: A novel DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol. Cancer Res.* 1: 247–261
- Taira T, Iguchi-Arigo SMM and Ariga H (2004) Co-localization with DJ-1 is essential for the androgen receptor to exert its transcription activity that has been impaired by androgen-antagonists. *Biol. Pharm. Bull.* 27: 574–577
- Shinbo Y, Taira T, Niki T, Iguchi-Arigo SMM and Ariga H (2005) DJ-1 restores p53 transcription activity inhibited by Topors/p53BP3. *Int. J. Oncol.* 26: 641–648
- Sekito A, Taira T, Niki T, Iguchi-Arigo SMM and Ariga H (2005) Stimulation of transforming activity of DJ-1 by Abstrakt, a DJ-1-binding protein. *Int. J. Oncol.* 26: 685–689
- Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC, Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A, Meco G, van Duijn CM, Oostra BA and Heutink P (2003) Mutations in the DJ-1 gene associated with autosomal recessive early-onset parkinsonism. *Science* 299: 256–259
- Abou-Sleiman PM, Healy DG, Quinn N, Lees AJ and Wood NW (2003) The role of pathogenic DJ-1 mutations in Parkinson's disease. *Ann. Neurol.* 54: 283–286
- Hague S, Rogaeva E, Hernandez D, Gulick C, Singleton A, Hanson M, Johnson J, Weiser R, Gallardo M, Ravina B, Gwinn-Hardy K, Crawley A, St George-Hyslop PH, Lang AE, Heutink P, Bonifati V, Hardy J and Singleton A (2003) Early-onset Parkinson's disease caused by a compound heterozygous DJ-1 mutation. *Ann. Neurol.* 54: 271–274
- Hedrich K, Djarmati A, Schafer N, Hering R, Wellenbrock C, Weiss PH, Hilker R, Vieregge P, Ozellus LJ, Heutink P, Bonifati V, Schwinger E, Lang AE, Noth J, Bressman SB, Pramstaller PP, Riess O and Klein C (2004) DJ-1 (PARK7)



- mutations are less frequent than Parkin (PARK2) mutations in early-onset Parkinson disease. *Neurology* 62: 389–394
21. Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H, Rubenstein J, Boyer R, Stenroos ES, Chandrasekharappa S, Athanassiadou A, Papapetropoulos T, Johnson WG, Lazzarini AM, Duvoisin RC, Di Iorio G, Golbe LI and Nussbaum RL (1997) Mutation in the alpha-synuclein gene identified in families with Parkinson's disease. *Science* 276: 2045–2047
  22. Kitada T, Asakawa S, Hattori N, Matsumine H, Yamamura Y, Minoshima S, Yokochi M, Mizuno Y and Shimizu N (1998) Mutations in the parkin gene cause autosomal recessive juvenile parkinsonism. *Nature* 392: 605–608
  23. Leroy E, Boyer R, Auburger G, Leube B, Ulm G, Mezey E, Harta G, Brownstein MJ, Jonnalagada S, Chernova T, Dehejia A, Lavedan C, Gasser T, Steinbach PJ, Wilkinson KD and Polymeropoulos MH (1998) The ubiquitin pathway in Parkinson's disease. *Nature* 395: 451–452
  24. Macedo MG, Anar B, Bronner IF, Cannella M, Squitieri F, Bonifati V, Hoogeveen A, Heutink P and Rizzo P (2003) The DJ-1L166P mutant protein associated with early onset Parkinson's disease is unstable and forms higher order protein complexes. *Hum. Mol. Genet.* 12: 2807–2816
  25. Miller DW, Ahmad R, Hague S, Baptista MJ, Canet-Aviles R, McLendon C, Carter DM, Zhu PP, Stadler J, Chandran J, Klinefelter GR, Blackstone C and Cookson MR (2003) L166P Mutant DJ-1, causative for recessive Parkinson's disease, is degraded through the ubiquitin-proteasome system. *J. Biol. Chem.* 278: 36588–36595
  26. Moore DJ, Zhang L, Dawson TM and Dawson VL (2003) A missense mutation (L166P) in DJ-1, linked to familial Parkinson's disease, confers reduced protein stability and impairs homo-oligomerization. *J. Neurochem.* 87: 1558–1567
  27. Gerner K, Holtorf E, Odoj S, Nuscher B, Yamamoto A, Regula JT, Beyer K, Haass C and Kahle PJ (2004) Differential effects of Parkinson's disease-associated mutations on stability and folding of DJ-1. *J. Biol. Chem.* 279: 6943–6951
  28. Oltmann JA, Brown K, Wilkinson KD, Rees HD, Huai Q, Ke H, Levey AI, Li L and Chin LS (2004) Familial Parkinson's disease-associated L166P mutation disrupts DJ-1 protein folding and function. *J. Biol. Chem.* 279: 8506–8515
  29. Bandopadhyay R, Kingsbury AE, Cookson MR, Reid AR, Evans IM, Hope AD, Pittman AM, Lashley T, Canet-Aviles R, Miller DW, McLendon C, Strand C, Leonard AJ, Abou-Sleiman PM, Healy DG, Ariga H, Wood NW, de Silva R, Revesz T, Hardy JA and Lees AJ (2004) The expression of DJ-1 (PARK7) in normal human CNS and idiopathic Parkinson's disease. *Brain* 127: 420–430
  30. Rizzo P, Hinkle DA, Zhukareva V, Bonifati V, Severijnen LA, Martinez D, Ravid R, Kamphorst W, Eberwine JH, Lee VM, Trojanowski JQ and Heutink P (2004) DJ-1 colocalizes with tau inclusions: a link between parkinsonism and dementia. *Ann. Neurol.* 55: 113–118
  31. Neumann M, Muller V, Gerner K, Kretschmar HA, Haass C and Kahle PJ (2004) Pathological properties of the Parkinson's disease-associated protein DJ-1 in alpha-synucleinopathies and tauopathies: relevance for multiple system atrophy and Pick's disease. *Acta Neuropathol.* 107: 489–496
  32. Heikkila R and Cohen G (1971) Inhibition of biogenic amine uptake by hydrogen peroxide: a mechanism for toxic effects of 6-hydroxydopamine. *Science* 172: 1257–1258
  33. Nicklas WJ, Youngster SK, Kindt MV and Heikkila RE (1987) MPTP, MPP+ and mitochondrial function. *Life Sci.* 40: 721–729
  34. Lotharius J and O'Malley KL (2000) The parkinsonism-inducing drug 1-methyl-4-phenylpyridinium triggers intracellular dopamine oxidation. A novel mechanism of toxicity. *J. Biol. Chem.* 275: 38581–38588
  35. Jenner P (2003) Oxidative stress in Parkinson's disease. *Ann. Neurol.* 53 (Suppl. 3): S26–S36
  36. Mitsumoto A, Nakagawa Y, Takeuchi A, Okawa K, Iwamatsu A and Takanezawa Y (2001) Oxidized forms of peroxiredoxins and DJ-1 on two-dimensional gels increased in response to sublethal levels of paraquat. *Free Radical Res.* 35: 301–310
  37. Mitsumoto A and Nakagawa Y (2001) DJ-1 is an indicator for endogenous reactive oxygen species elicited by endotoxin. *Free Radical Res.* 35: 885–893
  38. Srisomsap C, Subhasitanont P, Otto A, Mueller EC, Panyarit P, Wittmann-Liebold B and Svasti J (2002) Detection of cathepsin B up-regulation in neoplastic thyroid tissues by proteomic analysis. *Proteomics* 2: 706–712
  39. Kinumi T, Kimata J, Taira T, Ariga H and Niki E (2004) Cysteine-106 of DJ-1 is the most sensitive cysteine residue to hydrogen peroxide mediated oxidation *in vivo* in human umbilical vein endothelial cells. *Biochem. Biophys. Res. Commun.* 317: 722–728
  40. Yokota T, Sugawara K, Ito K, Takahashi R, Ariga H and Mizusawa H (2003) Down regulation of DJ-1 enhances cell death by oxidative stress, ER stress, and proteasome inhibition. *Biochem. Biophys. Res. Commun.* 312: 1342–1348
  41. Taira T, Saito Y, Niki T, Iguchi-Ariga SMM, Takahashi K and Ariga H (2004) DJ-1 has a role in antioxidative stress to prevent cell death. *EMBO Rep.* 5: 213–218
  42. Valente EM, Abou-Sleiman PM, Caputo V, Muqit MM, Harvey K, Gispert S, Ali Z, Del Turco D, Bentivoglio AR, Healy DG, Albanese A, Nussbaum R, Gonzalez-Maldonado R, Deller T, Salvi S, Cortelli P, Gilks WP, Latchman DS, Harvey RJ, Dallapiccola B, Auburger G and Wood NW (2004) Hereditary early-onset Parkinson's disease caused by mutations in PINK1. *Science* 304: 1158–1160
  43. Schmidt D and Muller S (2003) PIAS/SUMO: new partners in transcriptional regulation. *Cell Mol. Life Sci.* 60: 2561–2574
  44. Seeler JS and Dejean A (2003) Nuclear and unclear functions of SUMO. *Nat. Rev. Mol. Cell. Biol.* 4: 690–699
  45. Kahyo T, Nishida T and Yasuda H (2001) Involvement of PIAS1 in the sumoylation of tumor suppressor p53. *Mol. Cell* 8: 713–718
  46. Schmidt D and Muller S (2002) Members of the PIAS family act as SUMO ligases for c-Jun and p53 and repress p53 activity. *Proc. Natl. Acad. Sci. USA* 99: 2872–2877
  47. Nishida T and Yasuda H (2002) PIAS1 and PIASxalpha function as SUMO-E3 ligases toward androgen receptor and repress androgen receptor-dependent transcription. *J. Biol. Chem.* 277: 41311–41317
  48. Sachdev S, Bruhn L, Sieber H, Pichler A, Melchior F and Grosschedl R (2001) PIASy, a nuclear matrix-associated SUMO E3 ligase, represses LEF1 activity by sequestration into nuclear bodies. *Genes Dev.* 15: 3088–3103
  49. Takahashi-Niki K, Niki T, Taira T, Iguchi-Ariga SMM and Ariga H (2004) Reduced anti-oxidative stress activities of DJ-1 mutants found in Parkinson's disease patients. *Biochem. Biophys. Res Commun.* 320: 389–397
  50. Canet-Aviles RM, Wilson MA, Miller DW, Ahmad R, McLendon C, Bandopadhyay S, Baptista MJ, Ringe D, Petsko GA and Cookson MR (2004) The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization. *Proc. Natl. Acad. Sci. USA* 101: 9103–9108
  51. Shendelman S, Jonason A, Martinat C, Leete T and Abeliovich A (2004) DJ-1 is a redox-dependent molecular chaperone that inhibits alpha-synuclein aggregate formation. *PLoS Biol.* 2: e362
  52. Martinat C, Shendelman S, Jonason A, Leete T, Beal MF, Yang L, Floss T and Abeliovich A (2004) Sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: an ES-derived cell model of primary parkinsonism. *PLoS Biol.* 2: e327
  53. Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L, Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS and Mak TW (2005) DJ-1, a novel regulator of the tumor suppressor PTEN. *Cancer Cell* 7: 263–273
  54. Xu J, Zhong N, Wang H, Elias JE, Kim CY, Woldman I, Pifl C, Gygi SP, Geula C and Yankner BA (2005) The Parkinson's disease-associated DJ-1 protein is a transcriptional co-activator that protects against neuronal apoptosis. *Hum. Mol. Genet.* 14: 1231–1241
  55. Pichler A, Gast A, Seeler JS, Dejean A and Melchior F (2002) The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* 108: 109–120
  56. Li Y, Wang H, Wang S, Quon D, Liu YW and Cordell B (2003) Positive and negative regulation of APP amyloidogenesis by sumoylation. *Proc. Natl. Acad. Sci. USA* 100: 259–264
  57. Honbou K, Suzuki NN, Horiuchi M, Niki T, Taira T, Ariga H and Inagaki F (2003) The crystal structure of DJ-1, a protein related to male fertility and Parkinson's disease. *J. Biol. Chem.* 278: 31380–31384
  58. Tao X and Tong L (2003) Crystal structure of human DJ-1, a protein associated with early-onset Parkinson's disease. *J. Biol. Chem.* 278: 31372–31379
  59. Wilson MA, Collins JL, Hod Y, Ringe D and Petsko GA (2003) The 1.1-A resolution crystal structure of DJ-1, the protein mutated in autosomal recessive early onset Parkinson's disease. *Proc. Natl. Acad. Sci. USA* 100: 9256–9261
  60. Huai Q, Sun Y, Wang H, Chin LS, Li L, Robinson H and Ke H (2003) Crystal structure of DJ-1/RS and implication on familial Parkinson's disease. *FEBS Lett.* 549: 171–175
  61. Lee SJ, Kim SJ, Kim IK, Ko J, Jeong CS, Kim GH, Park C, Kang SO, Suh PG, Lee HS and Cha SS (2003) Crystal structures of human DJ-1 and *Escherichia*

- coli* Hsp31 that share an evolutionarily conserved domain. *J. Biol. Chem.* 278: 44552–44559
62. Orth M and Schapira AH (2002) Mitochondrial involvement in Parkinson's disease. *Neurochem. Int.* 40: 533–541
  63. Chan HY, Warrick JM, Andriola I, Merry D and Bonini NM (2002) Genetic modulation of polyglutamine toxicity by protein conjugation pathways in *Drosophila*. *Hum. Mol. Genet.* 11: 2895–2904
  64. Ueda H, Goto J, Hashida H, Lin X, Oyanagi K, Kawano H, Zoghbi HY, Kanazawa I and Okazawa H (2002) Enhanced SUMOylation in polyglutamine diseases. *Biochem. Biophys. Res. Commun.* 293: 307–313
  65. Steffan JS, Agrawal N, Pallos J, Rockabrand E, Trotman LC, Slepko N, Illes K, Lukacsovich T, Zhu YZ, Cattaneo E, Pandolfi PP, Thompson LM and Marsh JL (2004) SUMO modification of Huntingtin and Huntington's disease pathology. *Science* 304: 100–104

## DJ-1 interacts with HIPK1 and affects H<sub>2</sub>O<sub>2</sub>-induced cell death

AYA SEKITO<sup>1</sup>, SHIZUYO KOIDE-YOSHIDA<sup>1</sup>, TAKESHI NIKI<sup>2</sup>, TAKAHIRO TAIRA<sup>1,2</sup>,  
SANAE M.M. IGUCHI-ARIGA<sup>2,3</sup>, & HIROYOSHI ARIGA<sup>1,2</sup>

<sup>1</sup>Graduate School of Pharmaceutical Sciences, Hokkaido University, Sapporo 060-0812, Japan, <sup>2</sup>CREST, Japan Science and Technology Corporation, Saitama 332-0012, Japan, and <sup>3</sup>School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan

Accepted by Professor E. Niki

(Received 11 August 2005; in revised form 26 October 2005)

### Abstract

DJ-1 is a novel oncogene and causative gene for the familial form of Parkinson's disease (PD). DJ-1 has multiple functions, including anti-oxidative stress by eliminating reactive oxygen species (ROS) and transcriptional regulation as a coactivator, and loss of these functions are thought to trigger the onset of PD. The mechanism underlying the prevention of cell death by DJ-1 is, however, not clear. In this study, we found that DJ-1 directly bound to homeodomain-interacting protein kinase 1 (HIPK1) *in vitro* and *in vivo* and that these proteins were colocalized in the nucleus. HIPK1 was then found to be degraded in human H1299 cells transfected with wild-type DJ-1 but not with a C106S DJ-1 mutant, a DJ-1 protein disrupting a catalytic domain of the putative protease, in a dose-dependent manner. Furthermore, although knockdown of either DJ-1 or HIPK1 rendered H1299 cells susceptible to H<sub>2</sub>O<sub>2</sub>-induced cell death, double-knockdown of DJ-1 and HIPK1 rendered H1299 cells resistant to H<sub>2</sub>O<sub>2</sub>-induced cell death, suggesting that the elevated level of HIPK1 induced by a low level of DJ-1 inhibits oxidative stress-induced cell death.

**Keywords:** DJ-1, HIPK1, cell death, oxidative stress, Parkinson's disease

### Introduction

DJ-1 was first identified by us as a novel oncogene that transforms mouse NIH3T3 cells in cooperation with activated ras [1]. Elevated expression of DJ-1 has been reported in cancer cells, including breast cancer, smoker-derived lung adenocarcinoma and prostate cancer [2–4]. DJ-1 was later found to participate in fertilization for sperm to penetrate into the zona pellucida of eggs [5–9]. DJ-1 was found to play a role in transcriptional regulation: a positive regulator of the androgen receptor [10–12] and p53 [13] and a negative regulator of PTEN tumor suppressor [14] and pyrimidine tract-binding protein-associated splicing factor (PSF) [15]. It has recently been shown that DJ-1 is responsible for onset of familial Parkinson's disease (PD), PARK7 [16], and 11 mutations in familial and sporadic forms of PD have been reported

[17–19]. DJ-1 has been shown to be expressed in almost all brain tissues in healthy men and patients with neurodegenerative diseases, including PD, Pick's disease and multiple system atrophy [20–22].

The pathology of PD involves an irreversible degeneration of the dopaminergic neuron. Various lines of evidence suggest that oxidative stresses leads to dopaminergic cell degeneration in PD [23–26]. Expression of DJ-1 has been shown to be induced by various reagents that induce reactive oxygen species (ROS) [27–30], and abnormal oxidative forms of DJ-1 have been found in some patients with sporadic forms of PD [20]. We previously reported that DJ-1 plays a role in the anti-oxidative stress reaction, in which ROS were scavenged by oxidizing DJ-1 itself, and that mutants of DJ-1, including various mutants found in PD patients, rendered cells susceptible to oxidative stress-induced to cell death [31,32]. Other

Correspondence: H. Ariga, Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita 12, Nishi 6, Kita-ku, Sapporo 060-0812, Japan. Tel: 81 11 707 3745. Fax: 81 11 706 4988. E-mail: hiro@pharm.hokudai.ac.jp

groups have also reported anti-oxidative activity of DJ-1 [33–35]. It has been shown that of three cysteines at amino acid numbers 46, 53 and 106 in DJ-1, C106 was first oxidized as a form of addition of SO<sub>3</sub>H or SO<sub>2</sub>H and then C46 and C53 were oxidized with increased amounts of H<sub>2</sub>O<sub>2</sub> added to cultured cells [30] and that C106 was indispensable for DJ-1 to exert its activity against oxidative stress [32–36]. In addition to oxidation of DJ-1, DJ-1 has been reported to be conjugated with SUMO-1 on a lysine residue at amino-acid number 130 (K130) [37]. It has also been reported that sumoylation of DJ-1 at K130 was essential for DJ-1 to exert its full activities, including activities for ras-dependent transformation, cell growth stimulation and anti-oxidative stress, and that improper sumoylation of L166P DJ-1, a DJ-1 mutant found in PD patients, occurred and resulted in loss of its function and formation of aggregation [37].

Oxidative stress is caused by ROS in cells in which large amounts of ROS are produced by alternative activities of scavenger proteins or by dysfunction of the mitochondrial respiratory chain pathway with reduction of complex I activity. We and other groups have shown that some DJ-1 is located in mitochondria in addition to the cytoplasm and nucleus [33,37,38] and that translocation of DJ-1 to mitochondria was stimulated by oxidative stress [39–42]. The functions of DJ-1 in mitochondria, however, are not clear. Recently, DJ-1 has been shown to interact with Daxx to inhibit apoptosis signal-regulating kinase 1 (ASK-1) activity and cell death by sequestering the death protein Daxx to the nucleus [43]. ASK-1 has been reported to be activated by Daxx that was translocated from the nucleus to cytoplasm after glucose deprivation or oxidative stress to cells, resulting in activation of the apoptosis pathway through Jun-N-terminal kinase (JNK) [44]. These findings suggest that there are at least two mechanisms by which DJ-1 acts against oxidative stress: one is elimination of ROS by oxidization of DJ-1 itself, and the other is activation of oxidative stress-induced apoptosis machinery.

In this study, we identified homeodomain-interacting protein kinase 1 (HIPK1) as a DJ-1-binding protein. HIPK1 has been shown to be involved in the apoptosis pathways of both Daxx-ASK-1 and p53 [45–49], and we have shown that DJ-1 bound to p53 to modulate its transcriptional activity [13]. We also showed that DJ-1 reduced the expression level of HIPK1 and affected the apoptosis activity of HIPK1.

## Materials and methods

### Cell culture

The human cell lines H1299 and 293T were cultured in Dulbecco's modified Eagle's medium supplemented with 10% calf serum.

### Construction of plasmids

Nucleotide sequences of the oligonucleotide used for PCR primers were as follows: hHIPK1 ATG, 5'-GAATTCATGGCATCACAGCTGCAA-3'; hHIPK1 sc, 5'-GGGAATCCCCACAGCAATGATTC-3'; hHIPK1 $\beta$  stop, 5'-GTCTGACTCAGCATCCAGCCAGAAG-3'; hHIPK1 $\alpha$  stop, 5'-GGCTCGAGCTATAGTAGGAATAC-3'; hHIPK2 ATG, 5'-GAATTCATGGCCTCACATGTGCAA-3'; hHIPK2 Bgl II antisense, 5'-CGACGCTTGAGATCTCCATG TTC-3'; hHIPK2 Bgl II sense, 5'-CCAGAA CATGGAGATCTGCAAGCG-3'; hHIPK2 stop, 5'-CTCGAGTTATATGTAAGGGTACTG-3'.

Combinations of plasmids constructed, primers, templates and restriction enzyme sites and plasmids inserted are shown in Table I. PCR was carried out using the respective template and primers, and the resultant PCR product was digested with and inserted into the respective restriction enzyme site of the plasmid. Other plasmids used in this study were described previously [1,10,32].

### In vivo binding assay

Ten microgram of pcDNA3-Flag-HIPK1 $\alpha$  was transfected into human H1299 cells that were 60% confluent in a 10-cm dish by the calcium phosphate precipitation technique [49]. Forty-eight hours after transfection, the whole cell extract was prepared by the procedure described previously [10,32]. Approximately 2 mg of the H1299 cell proteins was first immunoprecipitated with a mouse anti-Flag antibody (M2, Sigma) under the same conditions as those described previously [10,32]. The precipitates were then separated in a 12% polyacrylamide gel containing SDS, blotted onto a nitrocellulose filter, and reacted with a rabbit anti-DJ-1 antibody [1] and with the mouse anti-Flag antibody. Immunoprecipitated proteins were then reacted with an HRP or IRDye800-conjugated second antibody and visualized by an ECL system or an infrared imaging system (Odyssey, LI-COR), respectively.

### In vitro binding assay

GST-DJ-1 or GST was expressed in and purified from *E. coli*. Purified GST-DJ-1 or GST was then incubated with 35S-HIPK1 $\alpha$  or 35S-HIPK1 $\beta$  synthesized *in vitro* using a reticulocyte lysate of a TnT-transcription-translation coupled system (Promega) and subjected to a pull-down assay using glutathione Sepharose as described previously [11,13]. The precipitates were separated in a 12% of polyacrylamide gel containing SDS and visualized by fluorography.