

Fig. 6 Western blot analysis of cytochrome *c* release. (a) Time course of mitochondrial cytochrome *c* release into the cytosol. (b) Densitometric analysis of cytochrome *c* release expressed as a ratio to COX. The cells with Mito-G93A and Mito-G85R exhibited significantly more cytochrome *c* release than those with Cyto-G93A and Cyto-G85R. Dorfin significantly reduced the amount of mitochondrial cytochrome *c* released into the cytosol induced by Cyto-G93A and Cyto-G85R, whereas it did not affect that induced by Mito-SOD1. Values are mean \pm SD ($n = 4$). * $p < 0.05$ (two-way ANOVA with Tukey–Kramer post-hoc test).

the activation of caspase-9 and caspase-3 induced by Cyto-G93A and Cyto-G85R (Figs 7 and 8). However, Dorfin did not reduce the activation of caspase-9 and caspase-3 induced by Mito-G93A and Mito-G85R (Figs 7 and 8), as it did not reduce the release of cytochrome *c* induced by Mito-G93A and Mito-G85R (Fig. 6). These findings combined with the aforementioned observations suggest that the reduction in the amount of mitochondrial mutant SOD1 due to Dorfin results in attenuated activation of the mitochondrial PCD pathway and prevents eventual cell death.

Discussion

In the present study, we first demonstrated that Dorfin, an E3 for mutant SOD1s, attenuated the activation of the mitochondrial PCD pathway and prevented eventual cell death in a neuronal cell model of FALS by reducing the amount of mutant SOD1 in the mitochondria. Dorfin reduced the levels of both cytosolic and mitochondrial mutant SOD1-EGFP fusion proteins that were expressed by Cyto-G93A and Cyto-G85R without organelle-oriented signals, whereas Dorfin did not affect the level of mutant SOD1-EGFP fusion protein that was expressed by Mito-G93A and Mito-G85R with mitochondrial localizing signals. The reduction in the level of




Fig. 7 Western blot analysis of caspase-9 activation. (a) Time course of the activation of caspase-9. (b) Densitometric analysis of caspase-9 activation. The cells with Mito-G93A and Mito-G85R exhibited significantly more activation of caspase-9 than those with Cyto-G93A and Cyto-G85R. Dorfin significantly reduced the activation of caspase-9 induced by Cyto-G93A and Cyto-G85R, whereas it did not reduce that induced by Mito-SOD1. Values are mean \pm SD ($n = 4$). * $p < 0.05$ (two-way ANOVA with Tukey–Kramer post-hoc test).




Fig. 8 Western blot analysis of caspase-3 activation. (a) Time course of activation of caspase-3. (b) Densitometric analysis of caspase-3 activation. The cells with Mito-G93A and Mito-G85R exhibited significantly more activation of caspase-3 than those with Cyto-G93A and Cyto-G85R. Dorfin significantly reduced the activation of caspase-3 induced by Cyto-G93A and Cyto-G85R, whereas it did not reduce that induced by Mito-SOD1. Values are mean \pm SD ($n = 4$). * $p < 0.05$ (two-way ANOVA with Tukey–Kramer post-hoc test).

mitochondrial SOD1-EGFP was observed earlier than that of total or cytosolic SOD1-EGFP. Moreover, Dorfin was present in the cytosol, not in the mitochondria. These findings indicated that the mitochondrial mutant SOD1 without organelle-oriented signals (Cyto-G93A and Cyto-G85R) might be translocated from the cytosol, and we suggest that Dorfin reduces the mitochondrial accumulation of mutant SOD1 by enhancing the degradation of mutant SOD1 in the cytosol through the ubiquitin–proteasomal pathway, thereby reducing the uptake of mutant SOD1 into the mitochondria.

Many reports have documented mitochondrial involvement in ALS and FALS. Mitochondrial degeneration with vacuolization or membrane disintegration in motor neurons is one of the earliest pathological findings in FALS Tg mice (Dal Canto and Gurney 1994; Wong *et al.* 1995; Hirano 1996; Kong and Xu 1998; Jaarsma *et al.* 2000; Higgins *et al.* 2003). Moreover, mitochondrial dysfunction such as altered calcium homeostasis (Carri *et al.* 1997; Menzies *et al.* 2002b), decreased respiratory chain complex activity (Mattiuzzi *et al.* 2002; Menzies *et al.* 2002a), alteration of mitochondria-related gene expression (Yoshihara *et al.* 2002) and an increase in reactive oxygen species (Beretta *et al.* 2003) have been reported in *in vitro* and *in vivo* models of FALS. Several studies have documented that SOD1, which has been considered a cytosolic enzyme, also exists in the mitochondrial intermembrane space (Okado-Matsumoto and Fridovich 2001; Sturtz *et al.* 2001; Higgins *et al.* 2002) and that the mitochondrial vacuoles are lined with mutant SOD1 in a FALS Tg mice model (Jaarsma *et al.* 2001; Higgins *et al.* 2003). Although the mitochondria-oriented vector we used here is designed to localize proteins to the mitochondrial matrix, we predict that SOD1-EGFP also exists in the mitochondrial intermembrane space through the process of its uptake into the mitochondrial matrix in our model, although we were not able to confirm this. Recent studies also revealed that SOD1 in the mitochondria originates from the uptake of SOD1 in the cytosol (Sturtz *et al.* 2001; Okado-Matsumoto and Fridovich 2002; Field *et al.* 2003). At least our result provided enough evidence that Dorfin interacts with mutant SOD1 in the cytosol, not in the mitochondria. Thus we suggest that Dorfin indirectly reduces the mitochondrial accumulation of mutant SOD1 by reducing the uptake of mutant SOD1 into the mitochondria.

Previous studies demonstrated that the mitochondrial PCD pathway, cytochrome *c* release and subsequent caspase activation, might contribute to the motor neuron cell death in FALS (Durham *et al.* 1997; Martin 1999; Li *et al.* 2000; Pasinelli *et al.* 2000; Guégan *et al.* 2001; Kriz *et al.* 2002; Zhu *et al.* 2002). Thus, inhibiting the activation of the mitochondrial PCD pathway is potentially useful in the treatment of FALS. Methods for this include inhibition of cytochrome *c* release by minocycline (Zhu *et al.* 2002; Kriz *et al.* 2002), co-expression of bcl-2 (Lee *et al.* 2001) or X-chromosome-linked inhibitor of apoptosis protein

(Ishigaki *et al.* 2002), and treatment with a broad caspase inhibitor zVAD-fmk (Pasinelli *et al.* 2000; Takeuchi *et al.* 2002a) or a caspase-9 specific inhibitor zLEHD-fmk (Takeuchi *et al.* 2002a). In this study, we demonstrated that Dorfin reduces the amount of mitochondrial mutant SOD1, attenuates the activation of the mitochondrial PCD pathway and prevents eventual neuronal cell death. It is therefore possible that reducing the amount of mutant SOD1 in the mitochondria may be adopted as a new therapeutic strategy for mutant SOD1-associated FALS.

Recent studies have suggested that some E3s, including Dorfin, act in a quality-control system to degrade cytosolic or transmembranous unfolded abnormal proteins (Moynihan *et al.* 1999; Fang *et al.* 2001; Meacham *et al.* 2001; Murata *et al.* 2001; Yoshida *et al.* 2002). The mitochondria also have a quality-control system that depends on mitochondria-specific molecular chaperones and ATPases associated with diverse cellular activities (AAA) proteases such as chaperonin 60 (Gottesman *et al.* 1997), mitochondrial heat-shock protein 70 (Savel'ev *et al.* 1998), and homologs of Lon, Yme1p, ClpP and ClpX (Wang *et al.* 1993; Suzuki *et al.* 1997; Langer 2000; Shah *et al.* 2000; Kang *et al.* 2002; Röttgers *et al.* 2003). A recent study documented that the accumulation of unfolded abnormal proteins in the mitochondria itself up-regulated the nuclear gene expression encoding mitochondrial-specific molecular chaperones (Zhao *et al.* 2002). Even though the mitochondria are able to dispose of abnormal proteins, they appear to have limited capacity to do this. They also seem to release death signals when abnormal proteins overflow their disposing capacity. Combination therapy such as Dorfin and mitochondria-specific molecular chaperones or AAA proteases thus seems more effective. Further investigations are needed to develop this therapeutic avenue.

There remains the problem of how the mutant SOD1 induces the mitochondrial PCD pathway. One of our previous studies revealed that bcl-2 family pro-apoptotic proteins, such as Bax, Bak, Bid, Bad and Bim, and other mitochondrial death signals such as apoptosis-inducing factor (AIF) and second mitochondria-derived activator of caspase (Smac) were not involved in the neuronal cell death in our model (Takeuchi *et al.* 2002a). Other studies have reported that translocation of Bax and cleavage of Bid were associated with neuronal cell death in the FALS Tg mouse model (Guégan *et al.* 2001; 2002), but there is a possibility that the surrounding environment of motor neurons such as astrocytes, microglia or dying neurons might have been affected in these models. Moreover, we have indicated that a non-apoptotic form of PCD might contribute to neuronal cell death through the mitochondrial PCD pathway in our model (Takeuchi *et al.* 2002a). Another report also mentioned that a non-apoptotic type of PCD acting through the mitochondrial PCD pathway might underlie mutant SOD1-related neurotoxicity (Guégan and Przedborski 2003). Further *in vivo*

investigations are needed to shed light on the mechanism of mutant SOD1-mediated neuronal cell death.

In this study we demonstrated that Dorfin, an E3 for mutant SOD1s, significantly reduced the level of mutant SOD1 in the mitochondria, attenuated the subsequent activation of the mitochondrial PCD pathway and prevented eventual neuronal cell death in a neuronal cell model of FALS. Reducing the accumulation of mutant SOD1 in the mitochondria may have an important place in the therapeutic strategy for mutant SOD1-associated FALS, and Dorfin may play a key role in this.

Acknowledgements

We are grateful to Dr Keiji Tanaka (Department of Molecular Oncology, The Tokyo Metropolitan Institute of Medical Science) for his helpful comments. This work was supported by grants from the Ministry of Health, Labor and Welfare of Japan, and a Center of Excellence grant from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Ardley H. C., Tan N. G. S., Rose S. A., Markham A. F. and Robinson P. A. (2001) Features of the Parkin/Ariadne-like ubiquitin ligase, HHARI, that regulate its interaction with the ubiquitin-conjugating enzyme, UbCH7. *J. Biol. Chem.* **276**, 19640–19647.
- Beretta S., Sala G., Mattavelli L., Ceresa C., Casciati A., Ferri A., Carr M. T. and Ferrarese C. (2003) Mitochondrial dysfunction due to mutant copper/zinc superoxide dismutase associated with amyotrophic lateral sclerosis is reversed by *N*-acetylcysteine. *Neurobiol. Dis.* **13**, 213–221.
- Carri M. T., Ferri A., Battistoni A., Famhy L., Gabbianelli R., Poccia F. and Rotilio G. (1997) Expression of a Cu,Zn superoxide dismutase typical of familial amyotrophic lateral sclerosis induces mitochondrial alteration and increase of cytosolic Ca²⁺ concentration in transfected neuroblastoma SH-SY5Y cells. *FEBS Lett.* **414**, 365–368.
- Dal Canto M. C. and Gurney M. E. (1994) Development of central nervous system pathology in a murine transgenic model of human amyotrophic lateral sclerosis. *Am. J. Pathol.* **145**, 1271–1279.
- Durham H. D., Roy J., Dong L. and Figlewicz Z. D. A. (1997) Aggregation of mutant Cu/Zn superoxide dismutase proteins in a culture model of ALS. *J. Neuropathol. Exp. Neurol.* **56**, 523–530.
- Fang S., Ferrone M., Yang C., Jensen J. P., Tiwari S. and Weissman A. M. (2001) The tumor autocrine motility factor receptor, gp78, is a ubiquitin protein ligase implicated in degradation from the endoplasmic reticulum. *Proc. Natl Acad. Sci. USA* **98**, 14422–14427.
- Field L. S., Furukawa Y., O'Halloran T. V. and Culotta V. C. (2003) Factors controlling the uptake of yeast Cu/Zn superoxide dismutase into mitochondria. *J. Biol. Chem.* **278**, 28052–28059.
- Gottesman S., Wickner S. and Maurizi M. R. (1997) Protein quality control: triage by chaperones and proteases. *Genes Dev.* **11**, 815–823.
- Guégan C. and Przedborski S. (2003) Programmed cell death in amyotrophic lateral sclerosis. *J. Clin. Invest.* **111**, 153–161.
- Guégan C., Vila M., Rosoklija G., Hays A. P. and Przedborski S. (2001) Recruitment of the mitochondrial-dependent apoptotic pathway in amyotrophic lateral sclerosis. *J. Neurosci.* **21**, 6569–6576.
- Guégan C., Vila M., Teissman P., Chen C., Onténiente B., Li M., Friedlander R. M. and Przedborski S. (2002) Instrumental activation of Bid by caspase-1 in a transgenic mouse model of ALS. *Mol. Cell. Neurosci.* **20**, 553–562.
- Higgins C. M., Jung C., Ding H. and Xu Z. (2002) Mutant Cu,Zn superoxide dismutase that causes motoneuron degeneration is present in mitochondria in the CNS. *J. Neurosci.* **22** RC215.
- Higgins C. M. J., Jung C. and Xu Z. (2003) ALS-associated mutant SOD1G93A causes mitochondrial vacuolation by expansion of the intermembrane space and by involvement of SOD1 aggregation and peroxisomes. *BMC Neurosci.* **4**, 16–29.
- Hirano A. (1996) Neuropathology of ALS: an overview. *Neurology* **47**, S63–S66.
- Ishigaki S., Liang Y., Yamamoto M., Niwa J., Ando Y., Yoshihara T., Takeuchi H., Doyu M. and Sobue G. (2002) X-linked inhibitor of apoptosis protein is involved in mutant SOD1-mediated neuronal degeneration. *J. Neurochem.* **82**, 576–584.
- Jaarsma D., Haasdijk E. D., Grashorn J. A. C., Hawkins R., van Duijn W., Verspaget H. W., London J. and Holstege J. C. (2000) Human Cu/Zn superoxide dismutase (SOD1) overexpression in mice causes mitochondrial vacuolization, axonal degeneration, and premature motoneuron death and accelerates motoneuron disease in mice expressing a familial amyotrophic lateral sclerosis mutant SOD1. *Neurobiol. Dis.* **7**, 623–643.
- Jaarsma D., Rognoni F., van Duijn W., Verspaget H. W., Haasdijk E. D. and Holstege J. C. (2001) CuZn superoxide dismutase (SOD1) accumulates in vacuolated mitochondria in transgenic mice expressing amyotrophic lateral sclerosis-linked SOD1 mutations. *Acta Neuropathol.* **102**, 293–305.
- Kang S. G., Ortega J., Singh S. K., Wang N., Huang N., Steven A. C. and Maurizi M. R. (2002) Functional proteolytic complexes of the human mitochondrial ATP-dependent protease, hClpXP. *J. Biol. Chem.* **277**, 21095–21102.
- Kong J. and Xu Z. (1998) Massive mitochondrial degeneration in motor neurons triggers the onset of amyotrophic lateral sclerosis in mice expressing a mutant SOD1. *J. Neurosci.* **18**, 3241–3250.
- Kriz J., Nguyen M. D. and Julien J. P. (2002) Minocycline slows disease progression in a mouse model of amyotrophic lateral sclerosis. *Neurobiol. Dis.* **10**, 268–278.
- Langer T. (2000) AAA proteases: cellular machines for degrading membrane proteins. *Trends Biochem. Sci.* **25**, 247–251.
- Lee M. H., Hyun D.-H., Halliwell B. and Jenner P. (2001) Effect of overexpression of wild-type and mutant Cu/Zn-superoxide dismutases on oxidative stress and cell death induced by hydrogen peroxide, 4-hydroxynonenal or serum deprivation: potentiation of injury by ALS-related mutant superoxide dismutases and protection by Bcl-2. *J. Neurochem.* **78**, 209–220.
- Li M., Ona V. O., Guégan C. et al. (2000) Functional role of caspase-1 and caspase-3 in an ALS transgenic mouse model. *Science* **288**, 335–339.
- Martin L. J. (1999) Neuronal death in amyotrophic lateral sclerosis is apoptosis: possible contribution of a programmed cell death mechanism. *J. Neuropathol. Exp. Neurol.* **58**, 459–471.
- Mattiazzi M., D'Aurelio M., Gajewski C. D., Martushova K., Kiaei M., Beal M. F. and Manfredi G. (2002) Mutated human SOD1 causes dysfunction of oxidative phosphorylation in mitochondria of transgenic mice. *J. Biol. Chem.* **277**, 29626–29633.
- Meacham G. C., Patterson C., Zhang W., Younger J. M. and Cyr D. M. (2001) The Hsc70 co-chaperone CHIP targets immature CFTR for proteasomal degradation. *Nat. Cell Biol.* **3**, 100–105.
- Menzies F. M., Cookson M. R., Taylor R. W., Turnbull D. M., Chrzanoska-Lightowlers Z. M., Dong L., Figlewicz D. A. and Shaw P. J. (2002a) Mitochondrial dysfunction in a cell culture model of familial amyotrophic lateral sclerosis. *Brain* **125**, 1522–1533.

- Menzies F. M., Ince P. G. and Shaw P. J. (2002b) Mitochondrial involvement in amyotrophic lateral sclerosis. *Neurochem. Int.* **40**, 543–551.
- Moynihn T. P., Ardley H. C., Nuber U., Rose S. A., Jones P. F., Markham A. F., Scheffner M. and Robinson P. A. (1999) The ubiquitin-conjugating enzymes UbcH7 and UbcH8 interact with RING finger/IBR motif-containing domains of HHARI and H7-AP1. *J. Biol. Chem.* **274**, 30963–30968.
- Murata S., Minami Y., Minami M., Chiba T. and Tanaka K. (2001) CHIP is a chaperone-dependent E3 ligase that ubiquitylates unfolded protein. *EMBO Report* **2**, 1133–1138.
- Niwa J., Ishigaki S., Doyu M., Suzuki T., Tanaka K. and Sobue G. (2001) A novel centrisomal RING-finger protein, Dorfin, mediates ubiquitin ligase activity. *Biochem. Biophys. Res. Commun.* **281**, 706–713.
- Niwa J., Ishigaki S., Hishikawa N., Yamamoto M., Doyu M., Murata S., Tanaka K., Taniguchi N. and Sobue G. (2002) Dorfin ubiquitylates mutant SOD1 and prevents mutant SOD1-mediated neurotoxicity. *J. Biol. Chem.* **277**, 36793–36798.
- Okado-Matsumoto A. and Fridovich I. (2001) Subcellular distribution of superoxide dismutase (SOD) in rat liver. *J. Biol. Chem.* **276**, 38388–38393.
- Okado-Matsumoto A. and Fridovich I. (2002) Amyotrophic lateral sclerosis: a proposed mechanism. *Proc. Natl Acad. Sci. USA* **99**, 9010–9014.
- Pasinelli P., Houseweart M. K., Brown R. H. Jr and Cleveland D. W. (2000) Caspase-1 and -3 are sequentially activated in motor neuron death in Cu,Zn superoxide dismutase-mediated familial amyotrophic lateral sclerosis. *Proc. Natl Acad. Sci. USA* **97**, 13901–13906.
- Raoul C., Estévez A. G., Nishimune H., Cleveland D. W., deLapeyrière O., Henderson C. E., Haase G. and Pettmann B. (2002) Motoneuron death triggered by a specific pathway downstream of Fas; potentiation by ALS-linked SOD1 mutations. *Neuron* **35**, 1067–1083.
- Rosen D. R., Siddique T., Patterson D. *et al.* (1993) Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. *Nature* **362**, 59–62.
- Röttgers K., Zufall N., Guiard B. and Voos W. (2003) The ClpB homolog Hsp78 is required for the efficient degradation of proteins in the mitochondrial matrix. *J. Biol. Chem.* **277**, 45829–45837.
- Savel'ev A. S., Novikova L. A., Kovaleva I. E., Luzikov V. N., Neupert W. and Langer T. (1998) ATP-dependent proteolysis in mitochondria. *J. Biol. Chem.* **273**, 20596–20602.
- Shah Z. H., Hakkaart G. A. J., Arku B., de Jong L., van der Spek H., Grivell L. A. and Jacobs H. T. (2000) The human homologue of the yeast mitochondrial AAA metalloprotease Yme1p complements a yeast *yme1* disruptant. *FEBS Lett.* **478**, 267–270.
- Sturtz L. A., Diekert K., Jensen L. T., Lill R. and Culotta V. C. (2001) A fraction of yeast Cu,Zn-superoxide dismutase and its metallochaperone, CCS, localize to the intermembrane space of mitochondria. *J. Biol. Chem.* **276**, 38084–38089.
- Suzuki C. K., Rep M., van Dijn J. M., Suda K., Grivell L. A. and Schatz G. (1997) ATP-dependent proteases that also chaperone protein biogenesis. *Trends Biochem. Sci.* **22**, 118–123.
- Takeuchi H., Kobayashi Y., Ishigaki S., Doyu N. and Sobue G. (2002a) Mitochondrial localization of mutant superoxide dismutase I triggers caspase-dependent cell death in a cellular model of familial amyotrophic lateral sclerosis. *J. Biol. Chem.* **277**, 50966–50972.
- Takeuchi H., Kobayashi Y., Yoshihara T., Niwa J., Doyu M., Ohtsuka K. and Sobue G. (2002b) Hsp70 and Hsp40 improve neurite outgrowth and suppress intracytoplasmic aggregate formation in cultured neuronal cells expressing mutant SOD1. *Brain Res.* **949**, 11–22.
- Wang N., Gottesman S., Willingham M. C., Gottesman M. M. and Maurizi M. R. (1993) A human mitochondrial ATP-dependent protease that is highly homologous to bacterial Lon protease. *Proc. Natl Acad. Sci. USA* **90**, 11247–11251.
- Wong P. C., Pardo C. A., Borchelt D. R., Lee M. K., Copeland N. G., Jenkins N. A., Sisodia S. S., Cleveland D. W. and Price D. L. (1995) An adverse property of a familial ALS-linked SOD1 mutation causes motor neuron disease characterized by vacuolar degeneration of mitochondria. *Neuron* **14**, 1105–1116.
- Yim M. B., Kang J. H., Yim H. S., Kwak H. S., Chock P. B. and Stadtman E. R. (1996) A gain-of-function of an amyotrophic lateral sclerosis-associated Cu,Zn-superoxide dismutase mutant: an enhancement of free radical formation due to a decrease in K_m for hydrogen peroxide. *Proc. Natl Acad. Sci. USA* **93**, 5709–5714.
- Yoshida Y., Chiba T., Tokunaga F. *et al.* (2002) E3 ubiquitin ligase that recognizes sugar chains. *Nature* **418**, 438–442.
- Yoshihara T., Ishigaki S., Yamamoto M., Liang Y., Niwa J., Takeuchi H., Doyu M. and Sobue G. (2002) Differential expression of inflammation- and apoptosis-related genes in spinal cords of a mutant SOD1 transgenic mouse model of familial lateral sclerosis. *J. Neurochem.* **80**, 158–167.
- Zhao Q., Wang J., Levichkin I. V., Stasinopoulos S., Ryan M. T. and Hoogenraad N. J. (2002) A mitochondrial specific stress response in mammalian cells. *EMBO J.* **21**, 4411–4419.
- Zhu S., Stavrovskaya I. G., Drozda M. *et al.* (2002) Minocycline inhibits cytochrome *c* release and delays progression of amyotrophic lateral sclerosis. *Nature* **417**, 74–78.



DDB2, the xeroderma pigmentosum group E gene product, is directly ubiquitylated by Cullin 4A-based ubiquitin ligase complex

Noriyuki Matsuda^a, Keiko Azuma^{a,b}, Masafumi Saijo^c, Shun-ichiro Iemura^d,
Yusaku Hioki^d, Tohru Natsume^d, Tomoki Chiba^a, Kiyoji Tanaka^c, Keiji Tanaka^{a,*}

^a Department of Molecular Oncology, Tokyo Metropolitan Institute of Medical Science, 3-18-22 Honkomagome, Bunkyo-ku, Tokyo 113-8613, Japan

^b Department of Biology, Ochanomizu University, 2-1-1 Ohtsuka, Tokyo 112-8610, Japan

^c Graduate School of Frontier Biosciences, Osaka University and Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Corporation, 1-3 Yamada-oka, Suita, Osaka 565-0871, Japan

^d National Institutes of Advanced Industrial Science and Technology, Biological Information Research Center (JBIRC), Kohhoh-ku, Tokyo 135-0064, Japan

Accepted 17 December 2004

DDB2, the xeroderma pigmentosum group E gene product, is directly ubiquitylated by Cullin 4A-based ubiquitin ligase complex

Noriyuki Matsuda^a, Keiko Azuma^{a,b}, Masafumi Saijo^c, Shun-ichiro Iemura^d,
Yusaku Hioki^d, Tohru Natsume^d, Tomoki Chiba^a, Kiyoji Tanaka^c, Keiji Tanaka^{a,*}

^a Department of Molecular Oncology, Tokyo Metropolitan Institute of Medical Science, 3-18-22 Honkomagome, Bunkyo-ku, Tokyo 113-8613, Japan

^b Department of Biology, Ochanomizu University, 2-1-1 Ohtsuka, Tokyo 112-8610, Japan

^c Graduate School of Frontier Biosciences, Osaka University and Core Research for Evolutional Science and Technology (CREST), Japan Science and Technology Corporation, 1-3 Yamada-oka, Suita, Osaka 565-0871, Japan

^d National Institutes of Advanced Industrial Science and Technology, Biological Information Research Center (JBIRC), Kohtoh-ku, Tokyo 135-0064, Japan

Accepted 17 December 2004

Abstract

Xeroderma pigmentosum (XP) is a genetic disease characterized by hypersensitivity to UV irradiation and high incidence of skin cancer caused by inherited defects in DNA repair. Mutational malfunction of damaged-DNA binding protein 2 (DDB2) causes the XP complementation group E (XP-E). DDB2 together with DDB1 comprises a heterodimer called DDB complex, which is involved in damaged-DNA binding and nucleotide excision repair. Interestingly, by screening for a cellular protein(s) that interacts with Cullin 4A (Cul4A), a key component of the ubiquitin ligase complex, we identified DDB1. Immunoprecipitation confirmed that Cul4A interacts with DDB1 and also associates with DDB2. To date, it has been reported that DDB2 is rapidly degraded after UV irradiation and that overproduction of Cul4A stimulates the ubiquitylation of DDB2 in the cells. However, as biochemical analysis using pure Cul4A-containing E3 is missing, it is still unknown whether the Cul4A complex directly ubiquitylates DDB2 or not. We thus purified the Cul4A-containing E3 complex to near homogeneity and attempted to ubiquitylate DDB2 in vitro. The ubiquitylation of DDB2 was reconstituted using this pure E3 complex, indicating that DDB–Cul4A E3 complex in itself can ubiquitylate DDB2 directly. We also showed that an amino acid substitution, K244E, in DDB2 derived from a XP-E patient did not affect its ubiquitylation.

© 2005 Elsevier B.V. All rights reserved.

Keywords: Nucleotide excision repair; E3; Ubiquitin; DDB1; DDB2; Cullin 4A

1. Introduction

Several proteins that bind specifically to ultraviolet (UV) irradiation damaged-DNA have been discovered by electrophoretic mobility shift assay or filter-binding assay since 1970s [1]. Previous studies that have characterized the damaged-DNA binding (DDB) protein indicated that the minimal DDB complex is a heterodimer comprised of a 127 kDa DDB1 subunit and 48 kDa DDB2 subunit. The binding activ-

ity of damaged DNA is thought to reside in this heterodimeric complex (for reviews, see [2,3]).

Xeroderma pigmentosum (XP) is a rare genetic disease characterized by clinical and cellular hypersensitivity to UV radiation and high incidence of skin cancer [4]. Cells from XP patients show defective repair of DNA damage that had been induced by UV or chemical agents, and tendency for skin carcinogenesis. In 1988, Chu and Chang [5] reported that cells from XP complementation group E (XP-E) individuals (GM02415/XP2RO) lacked this damaged-DNA binding activity, suggesting that DDB is functionally involved in the XP-E disease. This is also true for some other alleles of XP-E

* Corresponding author. Tel.: +81 3 3823 2237; fax: +81 3 3823 2237.
E-mail address: tanakak@rinshoken.or.jp (K. Tanaka).

patients [6]. Further evidence for the involvement came from microinjection experiments indicating that the purified DDB complex complements the XP-E cells' defect [7,8]. Other studies demonstrated that ectopic expression of human *DDB2* enhanced DNA repair in Chinese hamster V79 cells, which rarely express endogenous rodent *DDB2* [9]. Soon after the identification of *DDB1* and *DDB2* genes, Nichols et al. [10] revealed that *DDB2* was in fact mutated in XP-E cells lacking DDB activity. However, the molecular basis of the XP-E phenotype was ambiguous, because several groups found that cells from other patients with XP-E had normal levels of DDB activity (DDB+) and possessed no mutation in *DDB2* gene (reviewed in [2,3]). This discrepancy was puzzling until recently. Based on a thorough analysis, however, it was found that some DDB+ cell lines were mistakenly assigned to XP-E, and now it appears that all known authentic cases of XP-E are caused by *DDB2* mutations [11,12].

In eukaryotic cells, selective protein degradation is largely mediated by the ubiquitin/proteasome system. When ubiquitin is attached to the target protein by the ubiquitylation machineries, the proteasome recognizes the poly-ubiquitylated substrate to be degraded. This ubiquitin conjugating system requires the cascade reaction of three enzymes, namely E1, a ubiquitin-activating enzyme, E2, a ubiquitin-conjugating enzyme, and E3, a ubiquitin ligase. In 1999, Shiyanov et al. [13] reported that Cullin 4A (Cul4A) associates with the DDB complex. The cullin family of proteins compose a multimeric E3 complex. Cullin 1, which is the most well characterized cullin, serves as a rigid scaffold of its E3 complex and catalyses ubiquitylation through appropriate positioning of E2 and the substrate [14]. Other cullin family proteins including Cul4A are believed to function as well. The interaction between Cul4A and DDB1 was also demonstrated by several other groups recently ([15–20] and this work). These results, together with the rapid degradation of DDB2 after UV irradiation [21,22], suggest the involvement of Cullin 4A in DDB2 ubiquitylation and degradation. Strikingly, over-production of Cul4A stimulates the ubiquitylation of DDB2 [15,16]. However, since the latter studies did not show biochemical evidence of Cul4A involvement in the ubiquitylation of DDB2, it is still unknown whether the Cul4A-containing E3 complex in itself directly ubiquitylates DDB2 or not. To further investigate the mode of this ubiquitylation, an *in vitro* reconstitution by biochemical approach is obviously required. Here, we show that DDB2 can be ubiquitylated directly by the purified DDB–Cul4A E3 complex in a reconstitution *in vitro* experiment.

2. Materials and methods

2.1. Protein identification by LC–MS/MS analysis

The Cullin 4A-associated complexes were digested with *Achromobacter* protease-I and the resulting peptides were analyzed using a nanoscale LC–MS/MS system as described

previously [23,24]. The peptide mixture was applied to a Mightysil-PR-18 (1 μ m particle, Kanto Chemicals, Tokyo, Japan) frit-less column (45 mm \times 0.150 mm i.d.) and separated using a 0–40% gradient of acetonitrile containing 0.1% formic acid over 30 min at a flow rate of 50 nl/min. Eluted peptides were sprayed directly into a quadruple time-of-flight hybrid mass spectrometer (Q-T of *Ultima*, Micromass, Manchester, UK). MS and MS/MS spectra were obtained in data-dependent mode. Up to four precursor ions above an intensity threshold of 10 counts/s were selected for MSMS analyses from each survey scan. All MS/MS spectra were searched for protein sequences of Swiss Prot and RefSeq (NCBI) using batch processes of Mascot software package (Matrix Science, London, UK).

2.2. Cell culture condition

High-Five insect cells were maintained as an adherent culture in Grace insect media (Invitrogen, Carlsbad, CA, USA) supplemented with 8% fetal bovine serum (Sigma, St. Louis, MO, USA) and 1% penicillin-streptomycin (Invitrogen). ts41 cells established from Chinese hamster [25] were maintained in Dulbecco's modified Eagle's medium (Sigma) containing 10% fetal bovine serum and 1% penicillin-streptomycin under 5% CO₂ condition at 34 °C.

2.3. Immunoprecipitation experiment

To express DDB1, DDB2 and cullin family proteins, all plasmids were constructed from pcDNA3 or pcDNA3.1 plasmid (Invitrogen). Additional details of the plasmid construction processes will be provided upon request. Mammalian ts41 cells at 48 h after DNA transfection were harvested, washed by phosphate-buffer saline (PBS) and lysed with buffer A containing 20 mM Tris–HCl, pH 7.5, 150 mM NaCl, 0.5% Nonidet P-40 and 10% glycerol. After removal of the debris by centrifugation, anti-Flag antibody (M2)-conjugated agarose (Sigma) was added to the lysate and the mixture was incubated at 4 °C for 2 h under constant rotation. After extensive washing of immunoprecipitates with buffer A, binding proteins were eluted with sodium dodecyl sulphate (SDS)-containing buffer and boiled at 95 °C for 5 min. The eluate was subjected to immunoblotting using anti-Flag (M2; Sigma), anti-Myc (Santa Cruz, Delaware, CA, USA), anti-Cul4A (our laboratory collection) and anti-DDB1 antibodies (Zymed, San Francisco, CA, USA).

2.4. Protein purification

To overproduce His-DDB1, Flag-DDB2, Cullin 4A-HA and T7-Rbx1 proteins in insect cells, the tagged full-length cDNAs were inserted into pFastBac donor plasmid (Invitrogen). Additional details of the plasmid construction processes can be provided upon request. Subsequent production of baculovirus particles was carried out according to the protocol provided by the manufacturer. Baculovirus particles for His-

DDB1 and Flag-DDB2 were used to simultaneously infect High-Five cells, as well as viruses for Cullin4A-HA and T7-Rbx1. Insect cells were incubated for 48 h after infection, washed using PBS at 4 °C and then harvested by centrifugation. The cell extract was collected using buffer B containing 20 mM Tris-HCl, pH 7.5, 0.5% Nonidet P-40, 150 mM NaCl, 100 μ M ZnSO₄, 10 mM 2-mercaptoethanol, 6% glycerol and a protease inhibitor mixture without ethylenediaminetetraacetic acid (EDTA) (Roche, Mannheim, Germany). After centrifugation, the cell lysates were mixed together and incubated at 4 °C for 5 h with occasional gentle mixing. For initial purification, the cell lysate was loaded on a single-stranded DNA cellulose (Sigma) column equilibrated with buffer B. The column was then washed with buffer B containing 0.3 M NaCl followed by elution with buffer B containing 0.7 M NaCl. The eluted fraction was subsequently purified with nickel-chelating agarose (Qiagen, Stanford, CA, USA) pre-equilibrated with buffer B and eluted by 120 mM imidazole. This purified complex was further separated on a glycerol gradient sedimentation, which was carried out through a 10–40% glycerol gradient in 25 mM Tris-HCl, pH 7.5, 1 mM dithiothreitol (DTT) and 2 mM ATP for 22 h at 25,000 rpm ultracentrifugation. Fractions of 1 ml were collected from the top of the gradient and subjected to silver staining and immunoblotting.

To purify DDB2 (K244E)-containing complex, cell lysates containing His-DDB1, Flag-DDB2 (K244E), Cullin 4A-HA and T7-Rbx1 proteins in buffer B were collected as mentioned above. The DDB2 (K244E) complex was roughly purified with nickel-chelating agarose (Qiagen) pre-equilibrated with buffer B and eluted by 100 mM imidazole. Obtained fractions were then loaded onto HiTrap Heparin HP column (Amersham Biosciences, Piscataway, NJ, USA), washed with buffer C [20 mM Tris-HCl, pH 7.5, 150 mM NaCl, 100 μ M ZnSO₄, 1 mM DTT, 4.5% glycerol and protease inhibitor mixture without EDTA (Roche)], and eluted with a 0.15–0.75 M NaCl gradient in buffer C. The DDB2 (K244E)-containing complex was eluted around 0.5 M NaCl and was subjected to dialysis with buffer D containing 20 mM Tris-HCl, pH 8.0, 20 mM NaCl, 100 μ M ZnSO₄, and 1 mM DTT. A protein complex containing wild-type DDB2 was simultaneously isolated by the same method and used as a control.

To collect the authentic DDB-Cul4A complex from mammalian cells, HeLa cells stably expressing N-terminally FLAG-HA-tagged DDB2 were used. The genuine DDB-Cul4A complex was immunoprecipitated with anti-FLAG antibody followed by anti-HA antibody as described previously [17]. The eluates were further purified by Mini Q (Amersham Biosciences) column chromatography instead of glycerol density gradient centrifugation.

2.5. *In vitro* ubiquitylation assay

The ubiquitylation assay was essentially performed as described previously [26,27]. Briefly, the purified DDB-Cul4A

complex was incubated in 25 mM Tris-HCl, pH 7.5, 1 mM DTT, 25 μ M MG132 (Peptide Inc., Osaka, Japan), 5 mM MgCl₂, 100 μ M ZnSO₄, 2 mM ATP, 50 μ g of ubiquitin (Sigma)/ml, 2 μ g of E1/ml and 70 μ g of various E2-expressing *Escherichia coli* lysate/ml at 32 °C for 2 h and subjected to immunoblotting with anti-His (penta-His antibody; Qiagen, Stanford, CA, USA), anti-HA (HA.11, Berkeley Antibody Company, Berkeley, CA, USA), anti-Flag (M2; Sigma) and anti-T7 (Novagen, Madison, WI, USA) antibodies. In some cases, GST-ubiquitin was used instead of native ubiquitin.

3. Results

3.1. DDB complex physically interacts with Cullin 4A

To explore the molecular function of Cullin 4A, we examined the cellular partner(s) that interact with Cul4A in cells. A thorough analysis of human EST and genome sequences showed that the registered human Cul4A sequence (659 amino acid protein [28]) lacks its N-terminal 100 amino acid residues and thus the full-length Cul4A was obtained by PCR-assisted cDNA cloning and used hereafter. The complete nucleotide sequence of full-length Cul4A has been registered under accession number AB178950.

Flag-tagged Cul4A was expressed in HEK293 cells followed by immunoprecipitation by anti-Flag antibody. The immunoprecipitates were eluted with a Flag peptide and then digested with Lys-C endopeptidase (A. protease I) and the cleaved fragments were directly analyzed using a highly sensitive “direct nano-flow LC-MS/MS” system (for detail, see Section 2). Following database search, a dozen of peptides were assigned to MS/MS spectra obtained from four nano-LC-MS/MS analyses for the Flag-Cul4A-associated complexes and DDB1 was identified as one of the Cul4A-interacting proteins.

To confirm the interaction between Cul4A and DDB1, we performed immunoprecipitation experiment. Plasmids carrying Flag-tagged cullin family proteins (Cul1, 2, 3, 4A, 4B and 5) and myc-tagged DDB1 were concurrently transfected into ts41 cells. Extracts of the transfected or mock-transfected cells were subjected to immunoprecipitation using anti-Flag antibody followed by immunoblotting with anti-DDB1 antibody. As shown in Fig. 1A, Cul4A significantly interacted with DDB1. Cul1 also bound DDB1 weakly, whereas the other Cullins tested did not interact with DDB1. We next examined whether DDB2 also associates with Cul4A, because DDB1 and DDB2 are part of the DDB complex. Plasmids carrying 6myc-tagged cullin family proteins were transfected into ts41 cells along with a plasmid harboring Flag-tagged DDB2. Each extract was then subjected to immunoprecipitation using anti-Flag antibody and immunoblotting with anti-myc antibody. Consistent with the above results, DDB2 also interacted strongly with Cul4A and weakly with Cul1 and Cul4B (Fig. 1B). DDB2 did not bind with other cullin fam-




Fig. 1. DDB complex interacts with Cullin 4A. (A) Cul4A interacts with DDB1. Flag-tagged cullin family proteins and Myc-tagged DDB1 were simultaneously transfected into ts41 cells. After immunoprecipitation (IP) by anti-Flag antibody, the resulting immunoprecipitates were subjected to immunoblotting using anti-DDB1 antibody. (B) Cul4A also associates with DDB2. IP was similarly performed using FLAG-tagged DDB2 and Myc-tagged cullin family protein concurrently transfected into ts41 cells. After IP by anti-FLAG antibody, the resulting immunoprecipitates were analyzed using anti-Myc antibody.

ily proteins (Cul2, 3 and 5) examined. These results showed that DDB complex preferentially interacts with Cul4A, as reported previously [16].

3.2. Purification of DDB–Cul4A complex

We next attempted to purify DDB–Cul4A E3 complex using baculovirus expression system to perform biochemical experiments. Flag tag was fused to DDB2 at its N-terminus to facilitate its detection. This Flag-tagged DDB2 is thought to be functional because recent studies showed that ectopic expression of Flag-DDB2 enhanced DNA repair in Chinese hamster V79 cells [9], and purified Flag-DDB2 protein could restore damaged-DNA binding activity in extracts of XP-E patient cells [12]. DDB complex has been purified previously using DNA affinity column [13] and we also used DNA cellulose for initial purification of this complex. His6-tagged DDB1 and Flag-tagged DDB2 were simultaneously expressed in High-Five insect cells by the baculovirus induction system. Cul4A-HA and T7-Rbx1 were expressed concurrently as well. Each cell lysate was mixed and the resulting protein complex was purified by sequential column chromatography on single-stranded DNA cellulose, nickel-chelating agarose and subsequent 10–40% glycerol gradient by ultracentrifugation. The E3 complex comprised of DDB1, DDB2, Cul4A and Rbx1 was collected to near homogeneity as a peak fraction of glycerol gradient as shown in Fig. 2. Note that several other proteins were also detected in the final preparation (for example, a typical protein is shown by an asterisk in Fig. 2). However, since the peak fraction of such protein was inconsistent with that of the E3 complex in the glycerol gradient (data not shown), we think the protein is a contaminant derived from insect cells or a degradation

product of the expressed protein, rather than a protein physiologically associated with the E3 complex.

3.3. DDB2 is ubiquitylated by purified Cul4A complex

Using this purified complex, we next tried to reconstitute the ubiquitylation of DDB2 to check whether DDB–Cul4A complex per se can ubiquitylate DDB2. Since E3 generally requires specific E2 to mediate ubiquitylation, we tested eight different E2 enzymes (E2-20k, E2-25k, Ubc3, Ubc4, UbcH5a, UbcH5c, Ubc7 and Ubc8). Slower-migrating ladders derived from auto-ubiquitylation of Cul4A (see below) were observed only from the reaction with Ubc4, UbcH5a and UbcH5c, whereas the other E2 enzymes tested did not support this modification (Fig. 3A). We thus used UbcH5 family as a source of E2 in the following experiments. Purified DDB–Cul4A complex was incubated with ATP, ubiquitin, E1 and UbcH5a, and subjected to immunoblotting with the antibody for each component. As expected, ladders derived from the auto-ubiquitylation of Cul4A were observed (Fig. 3B, single asterisk in the middle panel). Moreover, apparent high molecular-mass ladders were evident when DDB2 was detected using the anti-Flag antibody (Fig. 3B, single asterisk in the left panel). In order to demonstrate that this modification was due to ubiquitylation, we repeated the ubiquitylation assay in the presence or absence of ubiquitin. The slower migrating ladders were not detected without ubiquitin, and the addition of GST-ubiquitin instead of native ubiquitin resulted in the appearance of larger molecular-mass bands (Fig. 3B, double asterisks), indicating that this modification indeed is ubiquitylation. In the case of DDB1, a single high-molecular band also emerged after *in vitro* ubiquitylation (Fig. 3B, right panel). However, this ubiquitylation sig-




Fig. 2. Purification of the baculovirus-expressed DDB–Cul4A complex. The DDB–Cul4A complex was purified by sequential column chromatography and subsequently separated onto a 10–40% glycerol gradient by ultracentrifugation. The peak fraction of DDB–Cul4A complex was resolved by SDS-PAGE and visualized by silver staining. Asterisk shows the contaminant protein (see Section 3).

nal of DDB1 was fainter than that of Cul4A and DDB2 (see Section 4).

To further investigate the biochemical characteristics of DDB–Cul4A complex, we next purified it under more physiological conditions. HeLa cells stably expressing FLAG-HA-tagged DDB2 [17] were used to collect E3 complex. The DDB2-containing complex was immunoprecipitated with

anti-FLAG antibody followed by anti-HA antibody as described previously [17] and the eluates were further purified by Mini Q column chromatography. The authentic E3 complex, comprised of DDB1, DDB2, Cul4A and Rbx1, was purified to almost homogeneity (Fig. 4A). When this complex was incubated with ATP, ubiquitin, E1 and UbcH5a, apparent high molecular-mass ladders derived from the ubiquity-




Fig. 3. In vitro reconstitution of DDB2 ubiquitylation. (A) The DDB–Cul4A E3 complex cooperates with Ubc4 and UbcH5 subfamily of E2 enzymes. Purified DDB–Cul4A E3 was incubated with the indicated E2 enzymes and subjected to immunoblotting with anti-HA antibody to identify the auto-ubiquitylation. (B) DDB2 was directly ubiquitylated by the DDB–Cul4A complex. Pure DDB1–DDB2–Cul4A complex was subjected to in vitro ubiquitylation assay in the absence (no) or presence of ubiquitin (Ub) or GST-ubiquitin (GST-Ub) and analyzed by immunoblotting with each antibody. Single asterisks show the ubiquitin conjugation and double asterisks indicate GST-ubiquitin conjugation.




Fig. 4. In vitro ubiquitylation of DDB2 using authentic DDB–Cul4A complex. (A) Purification of the genuine DDB–Cul4A complex. Purified complex was resolved by SDS-PAGE and visualized by silver staining. (B) In vitro ubiquitylation of DDB2 and auto-ubiquitylation of Cul4A. The authentic DDB–Cul4A complex was subjected to in vitro ubiquitylation assay in the absence (no) or presence of ubiquitin (Ub) or GST-ubiquitin (GST-Ub) and analyzed by immunoblotting with anti-HA (DDB2), anti-Cul4A and anti-DDB1 antibodies. Single asterisks show the ubiquitin conjugation and double asterisks indicate GST-ubiquitin conjugation.

lation of DDB2 and Cul4A were again observed (Fig. 4B, single asterisk). The exclusion of ubiquitin from the assay quenched these bands and replacement of native ubiquitin with GST-ubiquitin retarded their mobility (Fig. 4B, double asterisks). In contrast, DDB1 was rarely ubiquitylated, although a faint ubiquitylation signal was observed after long exposure (Fig. 4B, right panel). Because the DDB–Cul4A complex derived from both insect (Fig. 3) and mammalian (Fig. 4) cells directly ubiquitylated DDB2, we concluded that DDB2 was ubiquitylated by genuine DDB–Cul4A complex.

3.4. XP-E mutation does not affect ubiquitylation of DDB2 in vitro

Two cell lines established from XP-E patients, XP2RO and XP82TO, have been characterized in detail. XP2RO and XP82TO cells harbor naturally occurring single amino acid substitutions, R273H and K244E, in DDB2 protein, respectively. It has been reported that the XP82TO mutant protein (DDB2-K244E) interacts normally with DDB1 and Cullin 4A. Conversely, XP2RO mutant protein (DDB2-R273H) interacts with neither of them [13,29]. We also confirmed by immunoprecipitation experiments that DDB2-K244E interacts with DDB1 and Cullin 4A normally, but DDB2-R273H did not associate with either of them (Fig. 5A). Intriguingly, Ropic-Otrin et al. [21] reported that UV-induced rapid degradation of DDB2 protein did not occur in XP82TO cell line. This information prompted us to test whether K244E mutation affects the in vitro ubiquitylation of DDB2. Because DDB1–DDB2 (K244E)–Cul4A complex did not interact effectively with DNA cellulose, we were unable to purify it

compared with the wild-type complex (data not shown). We thus purified this mutant protein complex by affinity chromatography on nickel-chelating column and subsequent heparin column, and the bound DDB2 (K244E) complex was eluted around 0.5 M NaCl (Fig. 5B). As a control, the wild type DDB2-containing complex was simultaneously isolated by the same method. The DDB1–DDB2 (K244E)–Cul4A complex was incubated with ATP, ubiquitin, E1 and UbcH5a, and subjected to immunoblotting with the anti-Flag antibody. The mutant DDB2 protein was ubiquitylated in a manner equivalent to that of the wild-type control (Fig. 5C), indicating that XP82TO mutation (K244E) did not affect the ubiquitylation of DDB2 in vitro. This result also suggests that the mutated site of DDB2 (244th K) per se is not the unique ubiquitylation site.

4. Discussion

The DDB complex is regulated through several processes when cells are exposed to UV irradiation, namely very rapid translocation into the nucleus and binding to chromatin [17,29–31], hasty degradation of DDB2 protein [21,22] and final transcriptional induction of *DDB2* mRNA [32,33]. Chemical inhibition of proteasomes prevents rapid degradation of DDB2 protein, suggesting that this process is mediated by the ubiquitin/proteasome system. Among these regulation processes of DDB2, proteolytic degradation is the most intriguing because several recent reports [13,15–17] and our present results have shown a tight relationship between the DDB complex and proteins involved in ubiquitylation.




Fig. 5. XP82TO mutation (K244E) does not affect the *in vitro* ubiquitylation of DDB2. (A) DDB2-K244E interacts with DDB1 and Cul4A, but DDB2-R273H associates with neither. Flag-DDB2 (WT, K244E or R273H) was co-transfected with Myc-Cul4A and HA-DDB1 into ts41 cells. After immunoprecipitation (IP) by anti-Flag antibody, the resulting immunoprecipitates were subjected to immunoblotting using anti-Cul4A and anti-DDB1 antibodies. (B) The DDB2 (K244E) complex was resolved by SDS-PAGE and visualized by silver staining. (C) Ubiquitylation of DDB2 protein with XP-E mutation (DDB2-K244E) was comparable with that of wild-type DDB2 *in vitro*. The DDB–Cul4A complex containing mutant or wild type DDB2 was subjected to *in vitro* ubiquitylation in the presence (Ub) or absence (-) of ubiquitin. Asterisk indicates the ubiquitin-conjugated DDB2.

Interestingly, ectopic over-production of Cullin 4A accelerates the degradation of DDB2, suggesting that Cul4A ubiquitylates DDB2 [15,16]. However, no reconstitution experiments were performed and thus this information did not exclude the possible involvement of other E3(s) downstream of Cul4A in the ubiquitylation of DDB2, rather than directly by Cul4A. This situation prompted us to reconstitute the *in vitro* ubiquitylation of DDB2 and we presented in this study biochemical evidence for the ubiquitylation of DDB2 directly by the DDB–Cullin 4A complex.

4.1. *In vitro* ubiquitylation of each subunit of the DDB complex

It is well established that a significant fraction of DDB2 is degraded promptly after UV irradiation [21,22] and is also degraded in a cell cycle-dependent manner [16]. Conversely, it is still controversial whether another component of the DDB complex, DDB1, is a target of ubiquitylation and subsequent

degradation. Zhou's group reported that overproduction of Cul4A in cells stimulates the ubiquitylation of DDB1 [15]. In contrast, neither ectopically expressed Cul4A nor UV irradiation accelerates degradation of DDB1 was reported by other groups [12,16]. In our reconstitution experiment, DDB1 was very weakly ubiquitylated in the DDB–Cul4A complex from insect cells (Fig. 3B) and seldom ubiquitylated in the complex from HeLa cells (Fig. 4B). Because the HeLa cell-derived complex is purer and was considered to be isolated under more physiological conditions, this result supports the notion that DDB1 is not ubiquitylated by the Cul4A E3 complex. Even though DDB1 was faintly ubiquitylated, such mono- or di-ubiquitylation is insufficient for the proteasomal degradation. Therefore, we favor the scenario that not DDB1 but DDB2 is the target of ubiquitylation by Cul4A E3 complex *in vivo* [12,16].

4.2. XP-E mutation did not affect the ubiquitylation of p48 *in vitro*

Rapic-Otrin et al. [21] reported that UV-induced rapid degradation of DDB2 did not occur in the XP-E cell line (XP82TO) whose DDB2 harbors a K244E mutation. Because this mutant protein (DDB2-K244E) can interact with DDB1 and Cullin 4A (Fig. 5A [13]) but not with damaged DNA [8,12,34], this result suggests that the binding activity to damaged-DNA is necessary for the degradation of DDB2. Another possibility is that the mutated site of DDB2 (244th K) per se is the main ubiquitylation site, as suggested previously [21]. However, the latter is unlikely because we showed that this mutant protein was still ubiquitylated in a manner similar to the wild-type DDB2 protein *in vitro* (Fig. 5C). Perhaps binding to damaged-DNA renders the conformation of DDB complex more acquiescent for ubiquitylation and/or UV recruits DDB to some specialized chromatin place where the other ubiquitylation machinery is easy to access *in vivo*.

4.3. Biochemical role of DDB in nucleotide excision repair

In XP-E cells lacking the DDB activity, the nucleotide excision repair (NER) of cyclobutane pyrimidine dimer (CPD) is significantly impaired [32], suggesting the importance of DDB complex in NER *in vivo*. However, this DDB complex is not essential for the reconstitution of the cell-free NER *in vitro*. The NER reaction was successfully reconstituted in the absence of DDB [35–37], although it may exhibit some stimulatory or inhibitory effects under certain conditions [38–40]. One interpretation of these results is that some partner protein(s) of DDB complex may be missing in such NER assay *in vitro*. Recent studies [13,15–20,41–43] and the present work emphasize the role of the DDB1 complex in the ubiquitin ligation. We can thus speculate that the effect and requirement of DDB could change if other ubiquitylation machinery was added to the *in vitro* NER assay.

4.4. Biological function of DDB2 ubiquitylation

As mentioned above, DDB2 protein is rapidly degraded after UV-irradiation in vivo [21,22] and we showed here that the DDB–Cul4A complex could directly ubiquitylate DDB2. What is the function of DDB2 ubiquitylation and subsequent degradation? After damaged-DNA recognition, DDB is thought to hand over the DNA lesion to the following NER component(s) including XPC [2]. An appealing hypothesis is that clearance of DDB2 by ubiquitylation and succeeding degradation facilitates accession of the following NER factor(s) to the DNA lesion. However, it is not clear at present whether the ubiquitylation of DDB2 is only required for its UV-induced degradation, or is essential to change some biological character of DDB2 preceding degradation. Because various non-proteolytic functions of ubiquitylation have been identified recently [44], it is still conceivable that the ubiquitylation of DDB2 might have an additional role besides degradation. Moreover, we still do not know whether DDB2 ubiquitylation is a pertinent event for DDB function in DNA repair, or is the only side effect accompanied by ubiquitylation of authentic, relevant substrate. To define the precise role of DDB–Cullin 4A complex-mediated ubiquitylation, further studies are obviously required; especially the identification of the physiological substrate. Such experiments are currently underway in our laboratories.

Acknowledgements

We are grateful to Dr. Yoshihiro Nakatani of Harvard Medical School for providing HeLa cells expressing epitope-tagged DDB2 and to Dr. Kaoru Sugawara of RIKEN for the critical reading of the manuscript. We also thank all members of Prof. K. Tanaka laboratory for the helpful discussions. This work was supported by CREST of Japan Science and Technology (JST), Takeda Science Foundation (to M.S.) and Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan (to K.T.).

References

- [1] R.S. Feldberg, L. Grossman, A DNA binding protein from human placenta specific for ultraviolet damaged DNA, *Biochemistry* 15 (1976) 2402–2408.
- [2] J. Tang, G. Chu, Xeroderma pigmentosum complementation group E and UV-damaged DNA-binding protein, *DNA Repair* 1 (2002) 601–616.
- [3] B.B.O. Wittschieben, R.D. Wood, DDB complexities, *DNA Repair* 2 (2003) 1065–1069.
- [4] D. Bootsma, K.H. Kremer, J.E. Cleaver, J.H.J. Hoeijmakers, Nucleotide excision repair syndromes: xeroderma pigmentosum, Cockayne syndrome and trichothiodystrophy, in: B. Vogelstein, K.W. Kinzler (Eds.), *The Genetic Basis of Human Cancer*, McGraw Hill, New York, 1997, pp. 245–274.
- [5] G. Chu, E. Chang, Xeroderma pigmentosum group E cells lack a nuclear factor that binds to damaged DNA, *Science* 242 (1988) 564–567.
- [6] B.J. Hwang, S. Toering, U. Francke, G. Chu, p48 activates a UV-damaged-DNA binding factor and is defective in xeroderma pigmentosum group E cells that lack binding activity, *Mol. Cell. Biol.* 18 (1998) 4391–4399.
- [7] S. Keeney, A.P. Eker, T. Brody, W. Vermeulen, D. Bootsma, J.H. Hoeijmakers, S. Linn, Correction of the DNA repair defect in xeroderma pigmentosum group E by injection of a DNA damage-binding protein, *Proc. Natl. Acad. Sci. U.S.A.* 91 (1994) 4053–4056.
- [8] V. Ropic-Otrin, I. Kuraoka, T. Nardo, M. McLenigan, A.P. Eker, M. Stefanini, A.S. Levine, R.D. Wood, Relationship of the xeroderma pigmentosum group E DNA repair defect to the chromatin and DNA binding proteins UV-DDB and replication protein A, *Mol. Cell. Biol.* 18 (1998) 3182–3190.
- [9] J.Y. Tang, B.J. Hwang, J.M. Ford, P.C. Hanawalt, G. Chu, Xeroderma pigmentosum p48 gene enhances global genomic repair and suppresses UV-induced mutagenesis, *Mol. Cells* 5 (2000) 737–744.
- [10] A.F. Nichols, P. Ong, S. Linn, Mutations specific to the xeroderma pigmentosum group E Ddb-phenotype, *J. Biol. Chem.* 271 (1996) 24317–24320.
- [11] T. Itoh, S. Linn, T. Ono, M. Yamaizumi, Reinvestigation of the classification of five cell strains of xeroderma pigmentosum group E with reclassification of three of them, *J. Invest. Dermatol.* 114 (2000) 1022–1029.
- [12] V. Ropic-Otrin, V. Navazza, T. Nardo, E. Botta, M. McLenigan, D.C. Bisi, A.S. Levine, M. Stefanini, True XP group E patients have a defective UV-damaged DNA binding protein complex and mutations in DDB2 which reveal the functional domains of its p48 product, *Hum. Mol. Genet.* 12 (2003) 1507–1522.
- [13] P. Shiyanov, A. Nag, P. Raychaudhuri, Cullin 4A associates with the UV-damaged DNA-binding protein DDB, *J. Biol. Chem.* 274 (1999) 35309–35312.
- [14] N. Zheng, B.A. Schulman, L. Song, J.J. Miller, P.D. Jeffrey, P. Wang, C. Chu, D.M. Koepf, S.J. Elledge, M. Pagano, R.C. Conaway, J.W. Conaway, J.W. Harper, N.P. Pavletich, Structure of the Cul1-Rbx1-Skp1-F boxSkp2 SCF ubiquitin ligase complex, *Nature* 416 (2002) 703–709.
- [15] X. Chen, Y. Zhang, L. Douglas, P. Zhou, UV-damaged DNA-binding proteins are targets of CUL-4A-mediated ubiquitination and degradation, *J. Biol. Chem.* 276 (2001) 48175–48182.
- [16] A. Nag, T. Bondar, S. Shiv, P. Raychaudhuri, The xeroderma pigmentosum group E gene product DDB2 is a specific target of cullin 4A in mammalian cells, *Mol. Cell. Biol.* 21 (2001) 6738–6747.
- [17] R. Groisman, J. Polanowska, I. Kuraoka, J. Sawada, M. Saijo, R. Drapkin, A.F. Kisselev, K. Tanaka, Y. Nakatani, The ubiquitin ligase activity in the DDB2 and CSA complexes is differentially regulated by the COP9 signalosome in response to DNA damage, *Cell* 113 (2003) 357–367.
- [18] C. Liu, K.A. Powell, K. Mundt, L. Wu, A.M. Carr, T. Caspari, Cop9/signalosome subunits and Pcu4 regulate ribonucleotide reductase by both checkpoint-dependent and -independent mechanisms, *Genes Dev.* 17 (2003) 1130–1140.
- [19] T. Bondar, A. Ponomarev, P. Raychaudhuri, Ddb1 is required for the proteolysis of the *Schizosaccharomyces pombe* replication inhibitor Spd1 during S phase and after DNA damage, *J. Biol. Chem.* 279 (2004) 9937–9943.
- [20] I.E. Wertz, K.M. O'Rourke, Z. Zhang, D. Dornan, D. Arnott, R.J. Deshaies, V.M. Dixit, Human De-etiolated-1 regulates c-Jun by assembling a CUL4A ubiquitin ligase, *Science* 303 (2004) 1371–1374.
- [21] V. Ropic-Otrin, M.P. McLenigan, D.C. Bisi, M. Gonzalez, A.S. Levine, Sequential binding of UV DNA damage binding factor and degradation of the p48 subunit as early events after UV irradiation, *Nucleic Acids Res.* 30 (2002) 2588–2598.
- [22] M.E. Fitch, I.V. Cross, S.J. Turner, S. Adimoolam, C.X. Lin, K.G. Williams, J.M. Ford, The DDB2 nucleotide excision repair gene product p48 enhances global genomic repair in p53 deficient human fibroblasts, *DNA Repair* 2 (2003) 819–826.

- [23] T. Natsume, Y. Yamauchi, H. Nakayama, T. Shinkawa, M. Yanagida, N. Takahashi, T. Isobe, A direct nanoflow liquid chromatography-tandem mass spectrometry system for interaction proteomics, *Anal. Chem.* 74 (2002) 4725–4733.
- [24] M. Komatsu, T. Chiba, K. Tatsumi, S.-I. Iemura, I. Tanida, N. Okazaki, T. Ueno, E. Kominami, T. Natsume, K. Tanaka, A novel protein-conjugating system for Ufm1, a ubiquitin-fold modifier, *EMBO J.* 23 (2004) 1977–1986.
- [25] S. Handeli, H. Weintraub, The ts41 mutation in Chinese hamster cells leads to successive S phases in the absence of intervening G2, M, and G1, *Cell* 71 (1992) 599–611.
- [26] N. Matsuda, T. Suzuki, K. Tanaka, A. Nakano, Rma1, a novel type of RING finger protein conserved from Arabidopsis to human, is a membrane-bound ubiquitin ligase, *J. Cell Sci.* 114 (2001) 1949–1957.
- [27] N. Imai, N. Matsuda, K. Tanaka, A. Nakano, S. Matsumoto, W. Kang, Ubiquitin ligase activities of Bombyx mori nucleopolydiovirus RING finger proteins, *J. Virol.* 77 (2003) 923–930.
- [28] L.C. Chen, S. Manjeshwar, Y. Lu, D. Moore, B.M. Ljung, W.L. Kuo, S.H. Dairkee, M. Wernick, C. Collins, H.S. Smith, The human homologue for the *Caenorhabditis elegans* cul-4 gene is amplified and overexpressed in primary breast cancers, *Cancer Res.* 58 (1998) 3677–3683.
- [29] P. Shivanov, S.A. Hayes, M. Donepudi, A.F. Nichols, S. Linn, B.L. Slagle, P. Raychaudhuri, The naturally occurring mutants of DDB are impaired in stimulating nuclear import of the p125 subunit and E2F1-activated transcription, *Mol. Cell. Biol.* 19 (1999) 4935–4943.
- [30] V. Ropic-Otrin, M. McLenigan, M. Takao, A.S. Levine, M. Protic, Translocation of a UV-damaged DNA binding protein into a tight association with chromatin after treatment of mammalian cells with UV light, *J. Cell Sci.* 110 (1997) 1159–1168.
- [31] W. Liu, A.F. Nichols, J.A. Graham, R. Dualan, A. Abbas, S. Linn, Nuclear transport of human DDB protein induced by ultraviolet light, *J. Biol. Chem.* 275 (2000) 21429–21434.
- [32] B.J. Hwang, J.M. Ford, P.C. Hanawalt, G. Chu, Expression of the p48 xeroderma pigmentosum gene is p53-dependent and is involved in global genomic repair, *Proc. Natl. Acad. Sci. U.S.A.* 96 (1999) 424–428.
- [33] T. Itoh, C. O’Shea, S. Linn, Impaired regulation of tumor suppressor p53 caused by mutations in the xeroderma pigmentosum DDB2 gene: mutual regulatory interactions between p48(DDB2) and p53, *Mol. Cell. Biol.* 23 (2003) 7540–7553.
- [34] S. Keeney, H. Wein, S. Linn, Biochemical heterogeneity in xeroderma pigmentosum complementation group E, *Mutat. Res.* 273 (1992) 49–56.
- [35] M. Araki, C. Masutani, T. Maekawa, Y. Watanabe, A. Yamada, R. Kusumoto, D. Sakai, K. Sugawara, Y. Ohkuma, F. Hanaoka, Reconstitution of damage DNA excision reaction from SV40 minichromosomes with purified nucleotide excision repair proteins, *Mutat. Res.* 459 (2000) 147–160.
- [36] S.J. Araujo, F. Tirode, F. Coin, H. Pospiech, J.E. Syvaaja, M. Stucki, U. Hubscher, J.M. Egly, R.D. Wood, Nucleotide excision repair of DNA with recombinant human proteins: definition of the minimal set of factors, active forms of TFIIH, and modulation by CAK, *Genes Dev.* 14 (2000) 349–359.
- [37] D. Mu, C.H. Park, T. Matsunaga, D.S. Hsu, J.T. Reardon, A. Sancar, Reconstitution of human DNA repair excision nuclease in a highly defined system, *J. Biol. Chem.* 270 (1995) 2415–2418.
- [38] A. Aboussekhra, M. Biggerstaff, M.K. Shivji, J.A. Vilpo, V. Moncollin, V.N. Podust, M. Protic, U. Hubscher, J.M. Egly, R.D. Wood, Mammalian DNA nucleotide excision repair reconstituted with purified protein components, *Cell* 80 (1995) 859–868.
- [39] M. Wakasugi, M. Shimizu, H. Morioka, S. Linn, O. Nikaido, T. Matsunaga, Damaged DNA-binding protein DDB stimulates the excision of cyclobutane pyrimidine dimers in vitro in concert with XPA and replication protein A, *J. Biol. Chem.* 276 (2001) 15434–15440.
- [40] M. Wakasugi, A. Kawashima, H. Morioka, S. Linn, A. Sancar, T. Mori, O. Nikaido, T. Matsunaga, DDB accumulates at DNA damage sites immediately after UV irradiation and directly stimulates nucleotide excision repair, *J. Biol. Chem.* 277 (2002) 1637–1640.
- [41] Y. Yanagawa, J.A. Sullivan, S. Komatsu, G. Gusmaroli, G. Suzuki, J. Yin, T. Ishibashi, Y. Saijo, V. Rubio, S. Kimura, J. Wang, X.W. Deng, Arabidopsis COP10 forms a complex with DDB1 and DET1 in vivo and enhances the activity of ubiquitin conjugating enzymes, *Genes Dev.* 18 (2004) 2172–2181.
- [42] L.A. Higa, I.S. Mihaylov, D.P. Banks, J. Zheng, H. Zhang, Radiation-mediated proteolysis of CDT1 by CUL4-ROC1 and CSN complexes constitutes a new checkpoint, *Nat. Cell Biol.* 5 (2003) 1008–1015.
- [43] J. Hu, C.M. McCall, T. Ohta, Y. Xiong, Targeted ubiquitination of CDT1 by the DDB1–CUL4A-ROC1 ligase in response to DNA damage, *Nat. Cell Biol.* 6 (2004) 1003–1009.
- [44] J.D. Schnell, L. Hicke, Non-traditional functions of ubiquitin and ubiquitin-binding proteins, *J. Biol. Chem.* 278 (2003) 35857–35860.

Glycoprotein-specific ubiquitin ligases recognize N-glycans in unfolded substrates

Yukiko Yoshida^{1,2+}, Eru Adachi^{1,2}, Kanako Fukiya^{1,2}, Kazuhiro Iwai^{2,3} & Keiji Tanaka¹

¹Tokyo Metropolitan Institute of Medical Science, Bunkyo-ku, Tokyo, Japan, ²CREST, Japan Science and Technology Corporation (JST), Saitama, Japan, and ³Department of Molecular Cell Biology, Graduate School of Medicine, Osaka City University, Osaka, Japan

Misfolded or unassembled polypeptides in the endoplasmic reticulum (ER) are retro-translocated into the cytosol and degraded by the ubiquitin-proteasome system. We reported previously that the SCF^{Fbs1,2} ubiquitin-ligase complexes that contribute to ubiquitination of glycoproteins are involved in the ER-associated degradation pathway. Here we investigated how the SCF^{Fbs1,2} complexes interact with unfolded glycoproteins. The SCF^{Fbs1} complex was associated with p97/VCP AAA ATPase and bound to integrin- β 1, one of the SCF^{Fbs1} substrates, in the cytosol in a manner dependent on p97 ATPase activity. Both Fbs1 and Fbs2 proteins interacted with denatured glycoproteins, which were modified with not only high-mannose but also complex-type oligosaccharides, more efficiently than native proteins. Given that Fbs proteins interact with innermost chitobiose in N-glycans, we propose that Fbs proteins distinguish native from unfolded glycoproteins by sensing the exposed chitobiose structure.

Keywords: ubiquitin ligase; glycoprotein; unfold; N-glycan

EMBO reports (2005) 6, 239–244. doi:10.1038/sj.embor.7400351

INTRODUCTION

Most secretory and membrane proteins are co-translationally translocated into the lumen of the endoplasmic reticulum (ER). In the ER, these proteins are modified by N-linked oligosaccharides and subjected to 'quality control' in which aberrant proteins are distinguished from properly folded proteins (Ellgaard & Helenius, 2003). When the improperly folded or incompletely assembled proteins fail to restore their functional states, they are degraded by the ER-associated degradation (ERAD) system, which involves retrograde transfer of proteins from the ER to the cytosol followed by degradation by the proteasome. p97/VCP, a cytosolic ATPase member of the AAA ATPase family, may have several roles in the

ERAD pathway. It has been suggested that the p97-Ufd1-Npl4 complex is required for the extraction of misfolded proteins from the ER into the cytosol (Tsai *et al*, 2002). Such a retro-translocation step would most probably be mediated by dual recognition of the substrates by p97, with the complex binding both the nonubiquitinated segment of a substrate and the attached polyubiquitin chain (Ye *et al*, 2003). The p97 complex, which associates with the ER membrane proteins VIMP and Derlin-1 through the amino-terminal domain of p97, is thought to be required for the extraction of misfolded proteins from the ER (Lilley & Ploegh, 2004; Ye *et al*, 2004). However, p97 has also been proposed to be important for the release of ERAD substrates after their export from the ER, and thereby for their accessibility to the proteasome (Elkabetz *et al*, 2004). The role of each of these functions remains to be established.

In the ubiquitin system, the ubiquitin ligase 'E3' has an important role in the selection of target proteins for ubiquitination. At present, several E3s have been identified in the ERAD pathway, such as Hrd1 (Bays *et al*, 2001) and Doa10 (Swanson *et al*, 2001) in yeast, and gp78 (Fang *et al*, 2001), CHIP (Meacham *et al*, 2001) and Parkin (Imai *et al*, 2001) in mammals. In addition, we have recently identified a new member of the ERAD-linked E3 family, SCF^{Fbs}, which participates in ERAD for selective elimination of glycoproteins (Yoshida *et al*, 2002, 2003). Whereas Hrd1, Doa10 and gp78 are localized in the ER, SCF^{Fbs} complexes are localized in the cytosol similar to CHIP and Parkin.

The SCF is composed of Cullin1/Cdc53, Skp1, Roc1/Rbx1, and one member of the large family of F-box proteins, which are involved in trapping target proteins (Deshaies, 1999). Fbs1 and Fbs2 (F-box protein that recognizes sugar chains) interact with glycoproteins containing high-mannose oligosaccharides, protein modification of which occurs in the ER. Our recent X-ray crystallographic and nuclear magnetic resonance (NMR) studies of the substrate-binding domain of Fbs1 have shown that Fbs1 recognizes the inner chitobiose of high-mannose oligosaccharides by a small hydrophobic pocket located at the top of the β -barrel (Mizushima *et al*, 2004).

In this study, we examined whether Fbs proteins discriminate between folded and unfolded glycoproteins because ERAD substrates are thought to be unfolded. Both Fbs1 and Fbs2 preferably bind to denatured proteins that contain not only

¹Tokyo Metropolitan Institute of Medical Science, 3-18-22, Hon-komagome, Bunkyo-ku, Tokyo 113-8613, Japan

²CREST, Japan Science and Technology Corporation (JST), Saitama 332-0012, Japan

³Department of Molecular Cell Biology, Graduate School of Medicine, Osaka City University, Osaka 545-8585, Japan

*Corresponding author. Tel: +81 3 3823 2105; Fax: +81 3 3823 2965;

E-mail: yyosida@rinshoken.or.jp

Received 16 August 2004; revised 11 January 2005; accepted 13 January 2005; published online 18 February 2005




Fig 1 | Fbs1 binds to integrin-β1 dependent on p97 ATPase activity. (A) Fbs1 associates with p97 in microsomal fraction. Endogenous Fbs1 was immunoprecipitated from 100,000g supernatant (S) and precipitate (P) fractions of brains of adult mice. Lysate (15 μg each) and immunoprecipitates were analysed by immunoblotting with antibodies against Fbs1, p97, calreticulin and GRP78. (B) Interaction of Fbs proteins with p97. Lysates of 293T cells transiently expressing Flag-tagged F-box proteins (–, empty vector) and HA-tagged p97 were subjected to immunoprecipitation, and the resulting precipitates were analysed by immunoblotting. The asterisk shows immunoglobulin heavy chains. (C) Fbs1 binding to integrin-β1 in the cytosol depends on p97 ATPase activity. 293T cells were transfected with Flag-tagged Fbs1, V5-tagged integrin-β1 and HA-tagged p97 (–, empty vector; W, wild-type p97; M, mutant p97 (K524A)). Fbs1 was immunoprecipitated from supernatant (S) and precipitate (P) fractions. Expressions of p97, Fbs1 and integrin-β1 in fractionated lysates (5 μg each) and the amount of integrin-β1 associated with Fbs1 were analysed by immunoblotting using anti-V5 antibody. The membranous fraction (Mem.) was prepared from 24,000g precipitate. Asterisks show immunoglobulin heavy and light chains.

high-mannose but also complex-type oligosaccharides over native counterparts. The results showed that these F-box proteins probably interact with the innermost chitobiose in N-glycans in only unfolded glycoprotein in the ERAD pathway, considering that chitobiose moieties are usually masked by the folded polypeptide.

RESULTS
SCF^{Fbs} associates with the p97 complex

We have isolated Fbs1 from mouse brain cytosol as a novel sugar-binding protein that functions as a substrate-binding subunit in SCF-type E3 for ERAD (Yoshida *et al*, 2002), but the ubiquitination machinery for ERAD is probably associated with the ER membrane. To test the localization of Fbs1 proteins in mouse brain, we prepared anti-Fbs1 polyclonal antibodies. Lysates from adult mouse brain were fractionated into 100,000g supernatant (S)

and precipitate (P) fractions excluding 24,000g precipitate, and the presence of Fbs1 was analysed by immunoblotting (Fig 1A). Fbs1 was detected in the P as well as the S fractions, suggesting that Fbs1 interacts with proteins that associate with the ER membrane. As p97/VCP is thought to be involved in the retro-transport of ERAD substrates (Tsai *et al*, 2002), we examined the interaction of Fbs1 with p97. As shown in Fig 1A, Fbs1 was co-immunoprecipitated with p97 from the P but not from the S fraction. To determine whether other F-box proteins interact with p97, the Flag-tagged F-box proteins were expressed, together with haemagglutinin (HA)-tagged p97, in 293T cells, immunoprecipitated and analysed by immunoblotting (Fig 1B). Both Fbs1 and Fbs2 but not βTrCP1, an F-box protein with WD repeats for substrate recognition, were co-immunoprecipitated with p97. These results suggest that a part of Fbs proteins binds specifically to the p97-containing complex at the ER.




Fig 2 | Pull-down analysis of the interactions of Fbs1, Fbs1W and Fbs2 with native and denatured *N*-glycoproteins. Extracts of cells expressing Flag-tagged Fbs1, the W280A mutant of Fbs1 (Fbs1W) and Fbs2 were incubated with native or guanidine-HCl-treated (denatured) glycoproteins (lanes 1–5) or deglycosylated fetuin (lane 6)-immobilized beads. The beads were washed and then boiled with sample buffer. Lysates (7.5 μ g) and bound proteins were analysed by immunoblotting using anti-Flag antibody (top) and the structures of *N*-glycans in the glycoproteins tested are shown at the bottom.

Fbs1 binds to integrin- β 1 dependent on p97 activity

We identified pre-integrin- β 1, which was modified with high-mannose oligosaccharides, as one of the Fbs1 substrates (Yoshida *et al*, 2002). As the Fbs1–pre-integrin- β 1 interaction occurs in the cytosol, retro-translocation of integrin- β 1 from the ER into the cytosol is required for Fbs1 binding. To analyse the involvement of p97 in the interaction between Fbs1 and pre-integrin- β 1, 293T cells were transfected with expression plasmids encoding Flag-tagged Fbs1, V5-tagged integrin- β 1 together with HA-tagged p97 or its mutant, and treated with the proteasome inhibitor MG132 for 1.5 h before collecting cells. Immunoprecipitates with anti-Flag antibody of S and P fractions of transfected 293T cells were immunoblotted with anti-V5 antibody to detect co-immunoprecipitated integrin- β 1 (Fig 1C). Most of pre-integrin- β 1 was localized in the P fractions, and the expression of the ATPase-defective p97 mutant (K524A) increased the amount of total pre-integrin- β 1 in the P fraction (lanes 4–6). The amount of pre-integrin- β 1 associated with Fbs1 was greater in wild-type p97-expressing cells than in mutant p97-expressing cells in both S and P fractions (lanes 7–12). These results suggest that Fbs proteins recognize and ubiquitinate pre-integrin- β 1 retro-translocated by p97, and this modification may facilitate pre-integrin- β 1 binding to the p97–Ufd1–Npl4 complex, as well as its extraction from the ER.

Fbs interacts with denatured *N*-glycoproteins

Glycoproteins retro-translocated from the ER are not native proteins. Therefore, to examine whether Fbs proteins recognize

denatured glycoproteins better than native proteins, first we carried out a pull-down assay using several *N*-glycoproteins (Fig 2). We have reported that both Fbs1 and Fbs2 recognize the innermost chitobiose structure in high-mannose oligosaccharides (Yoshida *et al*, 2003). Although both ovalbumin and ribonuclease B (RNaseB) contain high-mannose oligosaccharides, Fbs1 effectively bound to RNaseB alone (see Discussion). Fbs1 could bind to asialofetuin and ovomucoid, but could hardly bind to fetuin and ovalbumin. The ability of Fbs2 to bind to asialofetuin, fetuin and ovomucoid was weaker than that to proteins attached to the high-mannose oligosaccharides. Interestingly, both Fbs proteins could bind to all the denatured *N*-glycoproteins tested but not to denatured deglycosylated proteins (lane 6), whereas the W280A mutant of Fbs1 (Fbs1W) that fails to interact with the innermost GlcNAc moiety in *N*-glycan (Mizushima *et al*, 2004) could not bind to any native or denatured glycoproteins, suggesting that the denaturation of glycoproteins increases the accessibility to the innermost chitobiose of *N*-glycans by Fbs proteins.

We next examined whether the binding potency and substrate specificity of Fbs proteins are influenced by denaturation of cellular glycoproteins. Lysates from the mouse brain or Neuro2a cells were treated with or without 6 M guanidine-HCl, diluted ten times with the lysis buffer and incubated with the His-tagged Fbs proteins produced by *Escherichia coli* (Fig 3; supplementary information 1 online). Guanidine-HCl at 0.6 M had no influence on Fbs binding to glycoproteins (supplementary information 2 online). The glycoproteins bound to Fbs were isolated using Ni-NTA affinity chromatography and detected by lectin blotting (Fig 3). Blotting with GNA, a lectin that binds to high-mannose oligosaccharide, showed that denaturation markedly increased the number of proteins bound to Fbs. The spectrum of Fbs1-bound protein bands in the brain detected by WGA, a lectin specific for terminal GlcNAc or sialic acids, was similar to those detected by GNA, suggesting that these proteins are modified by both high-mannose and complex-type oligosaccharides. Conversely, the proteins detected by RCA120, a lectin that binds to terminal galactose- β 1-4GlcNAc, were different to those detected by GNA. Both the quantities and species of RCA120-reactive proteins recognized by Fbs1 were also considerably increased by denaturation. Treatment of denatured proteins with peptide: *N*-glycanase (PNGase F) almost diminished their binding to Fbs. Furthermore, Fbs1W could hardly bind these glycoproteins. These results strongly suggest that both Fbs1 and Fbs2 bind to the innermost GlcNAc moiety irrespective of the terminal sugar moieties, and that the accessibility of Fbs proteins to the innermost GlcNAc moiety is enhanced by denaturation of the substrate glycoproteins. As all *N*-linked oligosaccharides contain innermost chitobiose structure, Fbs proteins seem to be capable of binding most *N*-glycoproteins when denatured.

SCF^{Fbs1} ubiquitinates denatured glycoproteins

To see whether SCF^{Fbs1} ubiquitinates denatured glycoproteins more efficiently than native counterparts, we performed an *in vitro* ubiquitination assay using purified components including recombinant SCF^{Fbs1} proteins. Efficient ubiquitination of GlcNAc-terminated fetuin (GTF), which is an *in vitro* substrate for SCF^{Fbs1} (Yoshida *et al*, 2002), was detected by immunoblotting using an anti-fetuin antibody. When an excess amount of substrates existed, denatured asialofetuin was efficiently ubiquitinated,




Fig 3 | Interactions of Fbs1 and Fbs2 with cellular glycoproteins containing *N*-linked oligosaccharides in native and denatured states. Native (N), denatured (D) or PNGase-treated proteins prepared from mouse brain (A) or Neuro2a cells (B) were incubated with recombinant Fbs1-, Fbs1W- and Fbs2-immobilized beads. Native and PNGase-treated lysates (15 µg each; lanes 1 and 2) and proteins bound to Fbs1, Fbs1W or Fbs2 were analysed by lectin blotting using HRP-labelled GNA, WGA and RCA.

whereas ubiquitination of native asialofetuin was marginal (Fig 4; supplementary information 3 online). No ubiquitination of GTF or denatured asialofetuin was detected in the absence of E1, E2, ATP or substrate, and SCF^{Fbs1W} (Mizushima *et al*, 2004) failed to ubiquitinate these substrates (supplementary information 4 online). These results demonstrate that the higher affinity of Fbs1 for denatured *N*-glycoproteins results in a more efficient ubiquitination of the denatured substrates than native counterparts.

DISCUSSION

In the early secretory pathway, *N*-glycosylation facilitates conformational maturation by promoting the glycoprotein-folding machinery, and functions as tags for ER retention and targeting to the ERAD pathway. The calnexin-calreticulin cycle, consisting of two homologous lectins, calnexin and calreticulin, which interact with monoglucosylated *N*-glycans, in concert with UDP-glucose:glycoprotein glucosyltransferase (GT) and glucosidase II, has a central role in folding and ER retention. Conversely, it is shown that α -mannosidase I and EDEM have a pivotal role in selective disposal of misfolded glycoproteins (Ellgaard & Helenius, 2003; Yoshida, 2003). Among these oligosaccharide-related

molecules in the ER, only GT has been shown to recognize incompletely folded proteins (Parodi, 2000). In the cytosol, the *N*-glycans in proteins extracted from the ER are removed before proteolysis by PNGase. In addition, PNGase can discriminate between non-native and folded glycoproteins, favouring the former (Hirsch *et al*, 2004). In this study, we showed that the Fbs proteins preferentially bind to denatured glycoproteins over properly folded proteins. As the retro-translocated proteins in the cytosol from the ER are misfolded, it is conceivable that *N*-glycan recognition proteins in the cytosol can sense misfolded states.

Although GT and PNGase can distinguish the folding states of substrates, the structural elements required for identification of their targets are not fully understood. *In vitro* studies have shown that GT also preferentially re-glucosylates glycoproteins in partially folded, molten globule conformations (Caramelo *et al*, 2003), and that an important feature for recognition is the exposure of hydrophobic clusters and innermost GlcNAc residue (Sousa & Parodi, 1995). Fbs1 interacts with the inner chitobiose in *N*-glycans of glycoproteins by a specific binding surface located at one tip of the β -sandwich of its substrate-binding domain (Mizushima *et al*, 2004). The intramolecular interactions of




Fig 4 *In vitro* ubiquitination of native GlcNAc-terminated fetuin (GTF), asialofetuin (ASF) and denatured ASF by SCF^{Fbs1} ligase. The high-molecular-mass ubiquitinated fetuin ((GST-Ub)*n*-fetuin) was detected by immunoblotting with anti-fetuin antibody.

innermost GlcNAc residue and the polypeptide moiety generally hamper the binding of Fbs1 to the chitobiose portions of glycoproteins as a result of steric hindrance in their native states. Therefore, Fbs1 recognizes the innermost position of *N*-glycans as a signal for unfolded glycoproteins. Conversely, as RNaseB contains an oligosaccharide that does not contact the polypeptide chain except at the covalent attachment point (Williams *et al*, 1987), it is likely that RNaseB interacts with Fbs1 even in the native form probably due to the exceptional freedom of the innermost chitobiose portion (Fig 2). The present results confirmed that Fbs proteins bind to denatured glycoproteins.

Considering that ubiquitination of ERAD substrates is linked to retro-translocation and rapid degradation, ubiquitin ligases for ERAD might be associated with the ER membrane. Moreover, as many glycoproteins are efficiently deglycosylated by PNGase after retrograde transfer into the cytosol (Blom *et al*, 2004), the rapid recognition of substrate by Fbs proteins before deglycosylation is critical. We found a part of Fbs proteins in association with p97 in the microsomal fractions, and extraction of the substrate of SCF^{Fbs1} was dependent on the ATPase activity of p97 followed by association with Fbs1. Thus, SCF^{Fbs1} seems to be positioned in the ER membrane in such a way that it can ubiquitinate substrates immediately after retro-translocation to the cytosol.

Fbs proteins can bind not only high-mannose oligosaccharides but also various types of *N*-glycans in glycoproteins. These modified glycoproteins other than the high-mannose oligosaccharides are not ERAD substrates. Therefore, our finding suggests that SCF^{Fbs1} mediates ubiquitination of exogenous or membrane proteins endocytosed into the cells. This is not unusual, because it is well known that extracellular proteins incorporated by phagocytosis into dendritic cells are presented to MHC class I molecules after proteasomal degradation (Castellino *et al*, 2000). Other studies demonstrated the transfer of endocytosed proteins into the cytosol by unknown mechanisms before their proteasomal processing and/or destruction (Kovacsovic-Bankowski & Rock, 1995). Further functional analysis of Fbs family proteins can shed light on the degradation of endocytosed proteins.

METHODS

Transfection, plasmids, antibodies, immunoprecipitation and immunoblotting. 293T cells were transfected as described previously (Yoshida *et al*, 2002). HA-p97-expressing plasmid was a kind gift from S. Khochbin (INSERM, France). HA-p97 (K524A)-expressing plasmid was constructed by site-directed mutagenesis. Human integrin- β 1 complementary DNA was cloned from cDNA clone (ATCC 988953) in pTracer-EF-V5His vector (Invitrogen, Carlsbad, CA, USA). The anti-mouse Fbs1 serum was generated in rabbits by standard procedures using a synthetic peptide corresponding to residues 1–14 (MDGDGDPFVSHP) of the predicted protein coupled to keyhole limpet haemocyanin. Monoclonal antibodies to p97 and V5 epitope were purchased from Progen (Heidelberg, Germany) and Invitrogen, respectively, and polyclonal antibodies to calreticulin and GRP78 were from Affinity Bioreagents (Exeter, Devon, UK). Antibodies to Flag, HA and fetuin have been described previously (Yoshida *et al*, 2003). Immunoprecipitation from whole-cell extracts or subcellular fractionation of cells and immunoblotting were performed in TBS-T (1% Triton X-100, 50 mM Tris-HCl pH 7.5, 150 mM NaCl and protease inhibitors) as described previously (Yoshida *et al*, 2002).

Pull-down assay. Fetuin, asialofetuin, ovalbumin, RNaseB and ovomucoid were purchased from Sigma-Aldrich (St Louis, MO, USA). For preparation of deglycosylated fetuin (DGF), 10 mg of asialofetuin was incubated with 200 U of PNGase F (Roche, Mannheim, Germany) in 50 mM phosphate buffer, pH 7.2, at 37 °C for 24 h. The enzyme-treated proteins were loaded onto successive WGA and RCA-lectin agarose columns. The flow-through fraction from both columns was used as DGF. Each 10 mg glycoprotein was immobilized to 0.5 ml of Affi-gel 10 or 15 (Bio-Rad, Richmond, CA, USA). For preparation of denatured-glycoprotein-immobilized beads, after each half of glycoprotein-immobilized beads was incubated in 6 M guanidine-HCl for 2 h, the beads were washed five times with ten volumes of 20 mM Tris-HCl (pH 7.5)/150 mM NaCl (TBS) containing 0.5% NP-40 (TBS-N). Each cell extract prepared with TBS-N from Flag-tagged Fbs1, Fbs1 W280A mutant or Fbs2-expressing 293T cells (30 μ g) was incubated with 15 μ l of various glycoprotein-immobilized beads. Bound proteins were eluted by boiling with SDS sample buffer and were analysed by immunoblotting.

Binding assay and lectin blotting. The substrate-binding domain of mouse Fbs1 (117–297) and its W280A mutant were cloned into pET15b (Mizushima *et al*, 2004), that of mouse Fbs2 (46–295) was cloned into pET33b, and expressed in *E. coli*. The His-tagged Fbs proteins were bound to Ni-NTA agarose beads (Qiagen, Hilden, Germany). Mouse brains and Neuro2a cells were homogenized in TBS-N and protease inhibitors. After centrifugation of the homogenate at 15,000g for 30 min, guanidine-HCl was dissolved with one-third of the supernatant (protein concentration 5 mg ml⁻¹) up to 6 M. Guanidine-HCl-treated and untreated lysates were diluted ten times with TBS-N. Another aliquot was treated with PNGase F subsequent to denaturation by heating for 5 min at 100 °C in the presence of 1% SDS and was then diluted ten times with TBS-N. The dilutes and PNGase-treated lysates were precleared with Ni-NTA agarose and then the flow-through fractions were incubated with the Fbs-protein-bound beads for 18 h at 4 °C. The beads were washed with TBS-N containing 20 mM imidazole. The adsorbed proteins were eluted by 0.2 M

imidazole in TBS-N. Eluted proteins were separated by SDS-PAGE, and blotted onto a membrane (Immobilon). After the blotted membranes were blocked with 3% bovine serum albumin in PBS, lectin blotting was performed using horseradish peroxidase (HRP)-labelled GNA (EY Laboratories), RCA120 and WGA (Seikagaku-kogyo, Japan).

In vitro ubiquitination assays. Preparation of GTF and *in vitro* ubiquitination assays were performed as described previously (Yoshida et al, 2002). Denatured asialofetuin was prepared by 200 times dilution of 20 mg ml⁻¹ asialofetuin treated with 6 M guanidine-HCl. Details of the assay condition are described in supplementary information 5 online.

Supplementary information is available at *EMBO reports* online (<http://www.emboports.org>).

ACKNOWLEDGEMENTS

We thank F. Tokunaga for helpful comments on the manuscript. This work was supported in part by Grants-in-Aid from the Ministry of Education, Science, Culture of Japan and the Hayashi Memorial Foundation for Female Natural Scientists, and the Seki Memorial Foundation for Science, Tokyo, Japan.

REFERENCES

- Bays NW, Gardner RG, Seelig LP, Joazeiro CA, Hampton RY (2001) Hrd1p/Der3p is a membrane-anchored ubiquitin ligase required for ER-associated degradation. *Nat Cell Biol* **3**: 24–29
- Blom D, Hirsch C, Stern P, Tortorella D, Ploegh HL (2004) A glycosylated type I membrane protein becomes cytosolic when peptide: N-glycanase is compromised. *EMBO J* **23**: 650–658
- Caramelo JJ, Castro OA, Alonso LG, De Prat-Gay G, Parodi AJ (2003) UDP-Glc:glycoprotein glucosyltransferase recognizes structured and solvent accessible hydrophobic patches in molten globule-like folding intermediates. *Proc Natl Acad Sci USA* **100**: 86–91
- Castellino F et al (2000) Receptor-mediated uptake of antigen/heat shock protein complexes results in major histocompatibility complex class I antigen presentation via two distinct processing pathways. *J Exp Med* **191**: 1957–1964
- Deshai RJ (1999) SCF and Cullin/Ring H2-based ubiquitin ligases. *Annu Rev Cell Dev Biol* **15**: 435–467
- Elkabetz Y, Shapira I, Rabinovich E, Bar-Nun S (2004) Distinct steps in dislocation of luminal endoplasmic reticulum-associated degradation substrates: roles of endoplasmic reticulum-bound p97/Cdc48p and proteasome. *J Biol Chem* **279**: 3980–3989
- Ellgaard L, Helenius A (2003) Quality control in the endoplasmic reticulum. *Nat Rev Mol Cell Biol* **4**: 181–191
- Fang S, Ferrone M, Yang C, Jensen JP, Tiwari S, Weissman AM (2001) The tumour autocrine motility factor receptor, gp78, is a ubiquitin protein ligase implicated in degradation from the endoplasmic reticulum. *Proc Natl Acad Sci USA* **98**: 14422–14427
- Hirsch C, Misaghi S, Blom D, Pacold ME, Ploegh HL (2004) Yeast N-glycanase distinguishes between native and non-native glycoproteins. *EMBO Rep* **5**: 201–206
- Imai Y, Soda M, Inoue H, Hattori N, Mizuno Y, Takahashi R (2001) An unfolded putative transmembrane polypeptide, which can lead to endoplasmic reticulum stress, is a substrate of Parkin. *Cell* **105**: 891–902
- Kovacs-Kovacs-Bankowski M, Rock KL (1995) A phagosome-to-cytosol pathway for exogenous antigens presented on MHC class I molecules. *Science* **267**: 243–246
- Lilley BN, Ploegh HL (2004) A membrane protein required for dislocation of misfolded proteins from the ER. *Nature* **429**: 834–840
- Meacham GC, Patterson C, Zhang W, Younger JM, Cyr DM (2001) The Hsc70 co-chaperone CHIP targets immature CFTR for proteasomal degradation. *Nat Cell Biol* **3**: 100–105
- Mizushima T et al (2004) Structural basis of sugar-recognizing ubiquitin ligase. *Nat Struct Mol Biol* **11**: 365–370
- Parodi AJ (2000) Protein glucosylation and its role in protein folding. *Annu Rev Biochem* **69**: 69–93
- Sousa M, Parodi AJ (1995) The molecular basis for the recognition of misfolded glycoproteins by the UDP-Glc:glycoprotein glucosyltransferase. *EMBO J* **14**: 4196–4203
- Swanson R, Locher M, Hochstrasser M (2001) A conserved ubiquitin ligase of the nuclear envelope/endoplasmic reticulum that functions in both ER-associated and Mat α 2 repressor degradation. *Genes Dev* **15**: 2660–2674
- Tsai B, Ye Y, Rapoport TA (2002) Retro-translocation of proteins from the endoplasmic reticulum into the cytosol. *Nat Rev Mol Cell Biol* **3**: 246–255
- Williams RL, Greene SM, McPherson A (1987) The crystal structure of ribonuclease B at 2.5-Å resolution. *J Biol Chem* **262**: 16020–16031
- Ye Y, Meyer HH, Rapoport TA (2003) Function of the p97-Ufd1-Npl4 complex in retrotranslocation from the ER to the cytosol: dual recognition of nonubiquitinated polypeptide segments and polyubiquitin chains. *J Cell Biol* **162**: 71–84
- Ye Y, Shibata Y, Yun C, Ron D, Rapoport TA (2004) A membrane protein complex mediates retro-translocation from the ER lumen into the cytosol. *Nature* **429**: 841–847
- Yoshida Y (2003) A novel role for N-glycans in the ERAD system. *J Biochem (Tokyo)* **134**: 183–190
- Yoshida Y et al (2002) E3 ubiquitin ligase that recognizes sugar chains. *Nature* **418**: 438–442
- Yoshida Y, Tokunaga F, Chiba T, Iwai K, Tanaka K, Tai T (2003) Fbs2 is a new member of the E3 ubiquitin ligase family that recognizes sugar chains. *J Biol Chem* **278**: 43877–43884