

図5 統合失調症における白質の異常（拡散テンソル画像） (Mori et al, submitted)




図6 Dysbindin 変異型マウスにおける social interaction の低下




図7 統合失調症患者と健常者における dysbindin の遺伝子型と記憶機能 (言語性記憶指数) (上折れ線: 健常者; 下折れ線: 統合失調症患者) (A: protective haplotype; O: others)




図8 BDNF 遺伝子のマイクロサテライト多型の構造


AGCACTAGCTGCCTATTCCAAAAATGTGTAAAACACCACTCAGCTTTTTAAAAGTAG  
GATAAACTCAGAGCGCGCGCACA  
<CGCGCGCGCG>①  
<CACACACACACACACACACACACA>②  
<GAGAGA>③  
ACATCTCTAGTAAAAAGAAAAGTTGAGCTTTCTTAGCTAGATGTGTGTATTAGCCAGA  
AAAAGCCAAGGAGTGAAGGGTTTTAGAGAACTGGAGGAGATAAAGTGGA

①(CG)n: n=0, 4, or 5

②(CA)n: n=10 ~ 15

③(GA)n: n=2 or 3

図9 未服薬統合失調症患者と健常者の抹消血遺伝子発現の網羅的解析



## 研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Okada T, Hashimoto R, Numakawa T, Iijima Y, Kosuga A, Tatsumi M, Kamijima K, Kato T, Kunugi H	A complex polymorphic region in the brain-derived neurotrophic factor (BDNF) gene confers susceptibility to bipolar disorder and affects transcriptional activity.	Mol Psychiatry		In press	2006
Ohnishi T, Hashimoto R, Mori T, Nemoto K, Moriguchi Y, Iida H, Noguchi H, Nakabayashi T, Hori H, Ohmori M, Tsukue R, Anami K, Hirabayashi N, Harada S, Arima K, Saitoh O, Kunugi H:	The association between the Val158Met polymorphism of the catechol-O-methyl transferase gene and morphological abnormalities of the brain in chronic schizophrenia.	Brain	129 (Pt 2)	399-410	2006
Nemoto K, Ohnishi T, Mori T, Moriguchi Y, Hashimoto R, Asada T, Kunugi H.	The Val66Met polymorphism of the brain-derived neurotrophic factor gene affects age-related brain morphology.	Neurosci Lett.	397(1-2)	25-29	2006
Chiba S, Hashimoto R, Hattori S, Yohda M, Lipska B, Weinberger DR, Kunugi H	Effect of antipsychotic drugs on DISC1 and dysbindin expression in mouse frontal cortex and hippocampus.	J Neural Transm		In press	2006
Kunugi H, Hashimoto R, Okada T, Hori H, Nakabayashi T, Baba A, Kudo K, Omori M, Takahashi S, Tsukue R, Anami K, Hirabayashi N, Kosuga A, Tatsumi M, Kamijima K, Asada T, Harada S,	Arima K, Saitoh O: Possible association between nonsynonymous polymorphisms of the anaplastic lymphoma kinase (ALK) gene and schizophrenia in a Japanese population.	J Neural Transm		In press	2006

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Hashimoto R, Hattori S, Chiba S, Yagasaki Y, Okada T, Kumamaru E, Mori T, Nemoto K, Tanii H, Hori H, Noguchi H, Numakawa T, Ohnishi T, Kunugi H.	Susceptibility genes for schizophrenia.	Psychiatry Clin Neurosci	Suppl 1	S4-S10	2006

## ORIGINAL ARTICLE

# A complex polymorphic region in the brain-derived neurotrophic factor (BDNF) gene confers susceptibility to bipolar disorder and affects transcriptional activity

T Okada<sup>1</sup>, R Hashimoto<sup>1</sup>, T Numakawa<sup>1</sup>, Y Iijima<sup>1</sup>, A Kosuga<sup>2</sup>, M Tatsumi<sup>2,3</sup>, K Kamijima<sup>2</sup>, T Kato<sup>4</sup> and H Kunugi<sup>1</sup>

<sup>1</sup>Department of Mental Disorder Research, National Institute of Neuroscience, National Center of Neurology and Psychiatry, Tokyo, Japan; <sup>2</sup>Department of Psychiatry, Showa University School of Medicine, Tokyo, Japan; <sup>3</sup>Yokohama Shinryo Clinic, Kanagawa, Japan and <sup>4</sup>Laboratory for Molecular Dynamics of Mental Disorders, Brain Science Institute, RIKEN, Saitama, Japan

Previous studies have suggested that genetic variations in the brain-derived neurotrophic factor (BDNF) gene may be associated with several neuropsychiatric diseases including bipolar disorder. The present study examined a microsatellite polymorphism located approximately 1.0 kb upstream of the translation initiation site of the BDNF gene for novel sequence variations, association with bipolar disorder, and effects on transcriptional activity. Detailed sequencing analysis revealed that this polymorphism is not a simple dinucleotide repeat, but it is highly polymorphic with a complex structure containing three types of dinucleotide repeats, insertion/deletion, and nucleotide substitutions that gives rise to a total of 23 novel allelic variants. We obtained evidence supporting the association between this polymorphic region (designated as BDNF-linked complex polymorphic region (BDNF-LCPR)) and bipolar disorder. One of the major alleles ('A1' allele) was significantly more common in patients than in controls (odds ratio 2.8, 95% confidential interval 1.5–5.3,  $P=0.001$ ). Furthermore, a luciferase reporter gene assay in rat primary cultured neurons suggests that this risk allele (A1) has a lower-transcription activity, compared to the other alleles. Our results suggest that the BDNF-LCPR is a functional variation that confers susceptibility to bipolar disorder and affects transcriptional activity of the BDNF gene.

*Molecular Psychiatry* advance online publication, 28 March 2006; doi:10.1038/sj.mp.4001822

**Keywords:** association study; brain-derived neurotrophic factor (BDNF); bipolar disorder; polymorphism; susceptibility; transcriptional activity

## Introduction

Brain-derived neurotrophic factor (BDNF) belongs to the neurotrophic factor family and promotes the development, regeneration, survival and maintenance of function of neurons.<sup>1</sup> It modulates synaptic plasticity and neurotransmitter release across multiple neurotransmitter systems, as well as the intracellular signal-transduction pathway.<sup>2</sup> BDNF has been implicated in the pathogenesis of mood disorders and in the mechanism of action of therapeutic agents such as mood stabilizers and antidepressants.<sup>3</sup> BDNF protein was reduced in postmortem brains of patients with bipolar disorder, compared to controls.<sup>4</sup> Chronic electroconvulsive seizure and antidepressant drug

treatments increase mRNA of BDNF and its receptor trkB.<sup>5</sup> Lithium may also exert its neuroprotective effect through enhancing expression of BDNF and trkB.<sup>6</sup>

The BDNF gene is, therefore, an attractive candidate gene which may give susceptibility to bipolar disorder.<sup>7</sup> In accordance with this, at least three previous studies reported a significant association between the Val66Met polymorphism (NCBI dbSNP rs6265) of the BDNF gene and bipolar disorder in Caucasian populations.<sup>8–10</sup> In these studies, the Val66 allele was consistently found to have a risk-increasing effect on the development of bipolar disorder. However, this association was not replicated in other Caucasian<sup>11–13</sup> or Asian populations including ours.<sup>14–16</sup>

Another polymorphism of the BDNF gene that has been well studied as to the possible association with neuropsychiatric diseases is the 'GT repeat' located approximately 1.0 kb upstream of the translation initiation site of the gene.<sup>17</sup> With respect to the possible effect on mood disorders, a significant

Correspondence: Dr H Kunugi, Department of Mental Disorder Research, National Institute of Neuroscience, National Center of Neurology and Psychiatry, 4-1-1, Ogawahigashi, Kodaira, Tokyo 187-8502, Japan.

E-mail: hkunugi@ncnp.go.jp

Received 8 November 2005; revised 19 January 2006; accepted 21 February 2006

linkage disequilibrium with bipolar disorder<sup>8</sup> and a significant association with childhood onset mood disorder<sup>18</sup> have been reported in Caucasian populations, although one study failed to find such an association with bipolar disorder.<sup>13</sup> Furthermore, there is some evidence suggesting that this polymorphism plays a role in the pathogenesis of schizophrenia.<sup>13,19,20</sup> However, there is no study that examined whether this polymorphism has functional effects. Since micro- and minisatellite polymorphisms even located in intron have been shown to play a role in the expression of many genes,<sup>21</sup> it might be intriguing to examine whether this microsatellite polymorphism of the BDNF gene is associated with transcriptional activity in an allele-dependent manner.

The aim of the present study was to examine this microsatellite polymorphism (designated here as BDNF-linked complex polymorphic region (BDNF-LCPR) due to its complex structure) for novel sequence variations, association with bipolar disorder, and effects on transcriptional activity.

## Materials and methods

### Subjects

Subjects were 153 patients with bipolar disorder (71 males) and the same number of controls (71 males), matched for age, sex, ethnicity, and geographical area. These subjects, who were recruited from Showa University Hospital and Shiga University of Medical Science Hospital, Japan, were previously genotyped for the Val66Met polymorphism of the BDNF gene, yielding a result of no significant association.<sup>16</sup> Mean age (standard deviation (s.d.)) in the patients was 47.8 (s.d. 15.3) years and that in the controls 47.1 (11.0). All the patients and controls were biologically unrelated Japanese. Consensus diagnosis of bipolar disorder was made for each patient by at least two experienced psychiatrists according to the Diagnostic and Statistical Manual of Mental Disorders, 4th ed. (DSM-IV),<sup>22</sup> based on unstructured interviews and medical records. Among the patients, 94 individuals (61%) were diagnosed with bipolar I and the remain-

ing 59 with bipolar II disorder. Patients who had one or more comorbid axis I disorders were excluded. The mean age of onset and number of episodes were 37.8 (s.d. 15.2) years and 3.9 (1.4) times, respectively. Thirty-four patients (22.2%) had at least one episode with psychotic features. Sixty-seven patients (43.8%) had a family history of major psychiatric illness (mood disorders or schizophrenia spectrum disorders) within their second-degree relatives. The controls were screened with a semi-structured interview and those individuals who had current or past contact to psychiatric services were excluded. In addition, those individuals who had a family history of major psychiatric illness or those who had a current or past history of regular use of psychotropic medication, including hypnotics, were excluded from the control group. After description of the study, written informed consent for the participation of the study was obtained from every subject. The study protocol was approved by ethics committee of each institution.

### Sequence analysis

Venous blood was drawn and genomic DNA was extracted according to standard procedures. To determine accurate DNA sequences for the BDNF-LCPR, we cloned this polymorphic region and performed direct sequencing. An approximately 400 base-pair (bp) DNA fragment encompassing the polymorphic region was amplified by polymerase chain reaction (PCR) with primers of *Hind*III-tagged BDNF-LCPR-F1 and *Hind*III-tagged BDNF-LCPR-R1 (see Table 1 and Figure 1a). The purified PCR products were ligated into the *Hind*III site of the pBluescriptII SK (+) vector (Toyobo, Tokyo, Japan). The vector was transformed into *Escherichia coli*, DH5 $\alpha$  and incubated. For sequencing, PCR amplification was performed with primers of GTTGTAACGACGGCCA GTG (Universal primer) and GGAAACAGCTATGAC CATG (Reverse primer). At least four clones were examined for each individual. Direct sequencing was performed with the CEQ8000 Genetic Analysis System (Beckman Coulter, Fullerton, CA, USA).

Cloning and sequencing analysis described above suggested that the polymorphic region is not a simple

**Table 1** Primer sequences for sequencing the BDNF-linked complex polymorphic region (BDNF-LCPR)

Primer No.	Primer Name	UCSC Chromosome11 5' → 3'	Numbers in Fig. 1 5' → 3'	Primer Sequence 5' → 3'
1	BDNF-LCPR-F1	27637949 → 27637930	139 → 158	TAGAGCAACCCTCTGGCAA
2	BDNF-LCPR-R1	27637545 → 27637567	543 → 521	TGTCATGAAAACAATGTGTCTGG
3	BDNF-LCPR-F2	27637844 → 27637822	244 → 266	CCAAAATGTGTAAAACACCACTC
4	B-BDNF-LCPR-F2	27637844 → 27637822	244 → 266	<b>Biotin</b> -CCAAAATGTGTAAAACACCACTC
5	BDNF-LCPR-R2	27637715 → 27637741	373 → 347	GAAAGCTCAACTTTTCTTTTACTAGA
6	B-BDNF-LCPR-R2	27637715 → 27637741	373 → 347	<b>Biotin</b> -GAAAGCTCAACTTTTCTTTTACTAGA
7	BDNF-LCPR-F3	27637810 → 27637791	278 → 297	AGTAGGATAAACTCAGAGCG
8	BDNF-LCPR-R3	27637730 → 27637749	358 → 339	CTTTTACTAGAGATGTTCT
9	Reverse		1 → 19	GGAAACAGCTATGACCATG
10	Universal		641 → 621	GTTGTAAAACGACGGCCAGTG





experiments and determined genotype for every subject. Genotypes were read blind to affection status.

#### Association analysis with bipolar disorder

The presence of Hardy–Weinberg equilibrium in genotype distribution was examined by using the  $\chi^2$ -test for goodness of fit. Allele frequencies of the BDNF-LCPR were compared between patients and controls by using the  $\chi^2$ -test for independence. Then linkage disequilibrium and haplotype-based association analysis for the BDNF-LCPR and the Val66Met polymorphisms were carried out. These statistical analyses were performed by using the SPSS v11 (SPSS Japan Inc., Tokyo, Japan) and the COCAPHASE v2.403 program (<http://www.hgmp.mrc.ac.uk/~fdudbrid/software/unphased/>). All *P*-values reported are two-tailed.

#### Luciferase reporter gene assay in primary cultured neurons


Primary cultures were prepared from the cortex of postnatal 2 days old rats (SLC, Shizuoka, Japan) as described previously.<sup>23</sup> To generate plasmids for the luciferase gene reporter assay (Figure 2a), the BDNF-LCPR was amplified by PCR with primers of *Sma*I-tagged BDNF-LCPR-F1 and *Sma*I-tagged BDNF-LCPR-R1 (Table 1 and Figure 1). The PCR products were inserted into the *Sma*I site upstream of the SV40 promoter in the pGL3-Promoter vector (Promega, Tokyo, Japan). The four major alleles were subject to the assay. Plasmid constructs were transfected at 5 days *in vitro*. Cells on 24-well plates were co-transfected with 800 ng of pGL3-Promoter firefly luciferase vectors that included major alleles of the BDNF-LCPR and 25 ng of phRL-TK renilla luciferase vector (Promega, Tokyo, Japan) as an internal control by using Lipofectamine 2000 reagent (Invitrogen, Tokyo, Japan). Empty pGL3-Promoter vector was transfected simultaneously.

At 24 h after transfection, luciferase activity was measured by using Dual-Luciferase Reporter Assay System (Promega, Tokyo, Japan) and a Lumat LB9507 luminometer (Berthold Technologies, Bad Wildbad, Germany), as described previously.<sup>24</sup> Firefly and renilla luciferase activities were quantified sequentially as relative light units (RLU) by addition of their respective substrates. The ratio of firefly RLU to renilla RLU of each sample was automatically computed. Then the activity of each construct was expressed as the relative value compared to that of empty pGL3-Promoter vector (relative luciferase expression, RLE). Primary cultured cells were prepared three times and transfection was performed triplicate for each cell culture. Comparisons in RLE were carried out by analysis of variance (ANOVA) or *t*-test.

## Results

#### Detection of novel variants

The structure of the BDNF gene<sup>25,26</sup> and DNA sequence of the cloned fragment according to the



**Figure 2** Luciferase reporter gene assay on the four major alleles of the BDNF-linked complex polymorphic region (BDNF-LCPR). (a) Schematic illustration of the luciferase assay construct for the BDNF-LCPR. (b) Relative luciferase expression (RLE) for pGL3-Promoter vector with insertion of each allele (A1, A2, A3, or A4) of the BDNF-LCPR in comparison with pGL3-Promoter vector without insertion of BDNF-LCPR (con). Error bars represent standard deviations (s.d.). \*\*: RLE for the A1 allele was significantly lower than the remaining three alleles combined ( $t = -3.4$ ,  $df = 34$ ,  $P = 0.002$ ).

University of California, Santa Cruz (UCSC) genome database are illustrated in Figure 1a. We detected a total of 23 allelic variants in the BDNF-LCPR (registered to the DDBJ/EMBL/GenBank database, accession numbers AB212736 to AB212758). Sequences and allele frequencies in patients with bipolar disorder and controls are shown in Table 2. Allelic variants of the BDNF-LCPR consisted of three components of dinucleotide repeat of (CA)<sub>del/2</sub>(CG)<sub>del/4/5</sub>, (CA)<sub>9–15</sub>, and (GA)<sub>2/3</sub>, which were combined in succession (Figure 1b). In addition, there were four exceptional rare variants that contained a single nucleotide substitution (variants 2 and 4) or insertion of two nucleotides of cg (variants 1 and 3) immediately 5' side of the repeats. The 'GT repeat' due to the original report<sup>17</sup> was CA, but not GT, repeat when the sequence was read in the forward direction of the BDNF gene. There were four major alleles of Del-12-3 (allele 1; A1), 4-12-3 (A2), 5-12-2 (A3), and 5-13-3 (A4). To perform statistical analyses, the remaining rare alleles were combined and considered to be 'allele 5 (A5)'. Supplementary figures S1 and S2 show images of direct sequencing of the major alleles, which were cloned and an example of pyrosequencing depicted in 'pyrogram'.

**Table 2** Detected alleles and their frequencies in patients with bipolar disorder and controls for the BDNF-linked complex polymorphic region (BDNF-LCPR)

Allele name	Sequence 5' → 3'	Fragment Size (bp) <sup>a</sup>	Allele counts (%)		
			Patients	Controls	Total
Del-11-3	agagcgcgcg(del)(ca) <sub>11</sub> (ga) <sub>3</sub> acat	393	2 (0.7)	1 (0.3)	3 (0.5)
Del-12-2	agagcgcgcg(del)(ca) <sub>12</sub> (ga) <sub>2</sub> acat	393	5 (1.6)	8 (2.6)	13 (2.1)
Del-12-3 (A1)	agagcgcgcg(del)(ca) <sub>12</sub> (ga) <sub>3</sub> acat	395	36 (11.8)	14 (4.6)	50 (8.2)
Del-15-3	agagcgcgcg(del)(ca) <sub>15</sub> (ga) <sub>3</sub> acat	401	1 (0.3)	0 (0.0)	1 (0.2)
4-11-3	agagcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>11</sub> (ga) <sub>3</sub> acat	405	0 (0.0)	1 (0.3)	1 (0.2)
4-12-2	agagcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>12</sub> (ga) <sub>2</sub> acat	405	0 (0.0)	1 (0.3)	1 (0.2)
4-12-3 (A2)	agagcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>12</sub> (ga) <sub>3</sub> acat	407	32 (10.5)	48 (15.7)	80 (13.1)
4-13-2	agagcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>13</sub> (ga) <sub>2</sub> acat	407	3 (1.0)	3 (1.0)	6 (1.0)
4-13-3	agagcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>13</sub> (ga) <sub>3</sub> acat	409	1 (0.3)	3 (1.0)	4 (0.7)
5-9-2	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>9</sub> (ga) <sub>2</sub> acat	401	1 (0.3)	0 (0.0)	1 (0.2)
5-10-3	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>10</sub> (ga) <sub>3</sub> acat	405	6 (2.0)	6 (2.0)	12 (2.0)
5-11-2	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>11</sub> (ga) <sub>2</sub> acat	405	4 (1.3)	2 (0.7)	6 (1.0)
5-11-3	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>11</sub> (ga) <sub>3</sub> acat	407	1 (0.3)	0 (0.0)	1 (0.2)
5-12-2 (A3)	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>12</sub> (ga) <sub>2</sub> acat	407	82 (26.8)	89 (29.1)	171 (27.9)
5-12-3	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>12</sub> (ga) <sub>3</sub> acat	409	5 (1.6)	6 (2.0)	11 (1.8)
5-13-2	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>13</sub> (ga) <sub>2</sub> acat	409	12 (3.9)	11 (3.6)	23 (3.8)
5-13-3 (A4)	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>13</sub> (ga) <sub>3</sub> acat	411	110 (35.9)	105 (34.3)	215 (35.1)
5-14-2	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>14</sub> (ga) <sub>2</sub> acat	411	0 (0.0)	1 (0.3)	1 (0.2)
5-14-3	agagcgcgcgcaca(cg) <sub>5</sub> (ca) <sub>14</sub> (ga) <sub>3</sub> acat	413	3 (1.0)	5 (1.6)	8 (1.3)
<i>Exceptional variants</i>					
Variant 1	agagcgcgcgcg(del)(ca) <sub>13</sub> (ga) <sub>3</sub> acat	399	1 (0.3)	0 (0.0)	1 (0.2)
Variant 2	agagcgcgcgcacatg(cg) <sub>4</sub> (ca) <sub>12</sub> (ga) <sub>2</sub> acat	405	0 (0.0)	1 (0.3)	1 (0.2)
Variant 3	agagcgcgcgcgcaca(cg) <sub>4</sub> (ca) <sub>13</sub> (ga) <sub>3</sub> acat	411	1 (0.3)	0 (0.0)	1 (0.2)
Variant 4	agagtgcgcgcaca(cg) <sub>5</sub> (ca) <sub>12</sub> (ga) <sub>2</sub> acat	407	0 (0.0)	1 (0.3)	1 (0.2)
Total chromosomes			306 (100.0)	306 (100.0)	612 (100.0)

<sup>a</sup>Fragment size of PCR product amplified by primers of BDNF-LCPR-F1 and BDNF-LCPR-R1 (see Table 1).

#### Association analysis with bipolar disorder

Genotype and allele distributions in patients and controls are shown in Table 3. The genotype distributions were in Hardy–Weinberg equilibrium (for the patients:  $\chi^2=4.5$ ,  $df=13$ ,  $P=0.98$ ; for the controls:  $\chi^2=8.2$ ,  $df=13$ ,  $P=0.83$ ). The overall allele frequencies differed significantly between patients and controls ( $\chi^2=13.4$ ,  $df=4$ ,  $P=0.0093$ ). The global- $P$ -value estimated by the permutation test of 10 000 simulations, correcting for multiple testing, yielded a similar result ( $P=0.010$ ). The A1 allele was clearly more common in patients than in controls (11.8 vs 4.6%, odds ratio (OR) 2.8, 95% confidential interval (CI) 1.5–5.3,  $\chi^2=10.5$ ,  $df=1$ ,  $P=0.001$ ). When the three components of the BDNF-LCPR, that is, (CA)<sub>del/2</sub> (CG)<sub>del/4/5</sub>, (CA)<sub>9–15</sub>, and (GA)<sub>2/3</sub>, were examined separately, only the first component showed a significant association with bipolar disorder (Table 4). Deletion of the first component, as seen in the A1 allele, was significantly more common in patients than in controls (14.6 vs 8.3%, OR 1.9, 95% CI 1.1–3.2,  $\chi^2=5.9$ ,  $df=1$ ,  $P=0.015$ ).

Then we examined linkage disequilibrium and haplotype-based association for the BDNF-LCPR and

the Val66Met polymorphism. As reported previously,<sup>16</sup> there was no significant association between the Val66Met polymorphism and bipolar disorder in the current sample; the frequencies of the Val66 allele were 0.60 and 0.62 in patients and comparison groups, respectively ( $\chi^2=0.25$ ,  $df=1$ ,  $P=0.62$ ). Results of haplotype-based analysis for these two polymorphisms are shown in Table 5. There was a very tight linkage disequilibrium between the BDNF-LCPR and Val66Met polymorphism ( $D'=0.91$  for patients and  $D'=0.90$  for controls;  $\chi^2=512$ ,  $df=28$ ,  $P=1.6 \times 10^{-90}$  in total subjects). The Val66 allele was linked to the A1, A2, or A3 allele, while the Met66 allele was to the A4 allele. The haplotype-based association analysis yielded a significant result (global  $P=0.0069$ ) estimated by the permutation test, correcting for multiple testing. Since the A1 allele was completely linked to the Val66 allele, the most significant individual  $P$ -value of 0.001 was obtained when the A1-Val66 was assumed to be the risk. When pairwise linkage disequilibrium across three components of the BDNF-LCPR and Val66Met was examined individually, there was a tight linkage disequilibrium between

**Table 3** Genotype and allele distributions in patients with bipolar disorder and controls for the BDNF-linked complex polymorphic region (BDNF-LCPR)

Genotype/allele	Counts (%)		
	Patients	Controls	Total
<b>Genotype</b>			
A1/A1	2 (1.3)	0 (0.0)	2 (0.7)
A1/A2	5 (3.2)	1 (0.7)	6 (2.0)
A1/A3	12 (7.8)	3 (2.0)	15 (4.9)
A1/A4	9 (5.9)	9 (5.9)	18 (5.9)
A1/A5	6 (3.9)	1 (0.6)	7 (2.3)
A2/A2	3 (2.0)	4 (2.6)	7 (2.3)
A2/A3	7 (4.6)	18 (11.8)	25 (8.2)
A2/A4	9 (5.9)	13 (8.5)	22 (7.2)
A2/A5	5 (3.3)	8 (5.2)	13 (4.2)
A3/A3	11 (7.2)	11 (7.2)	22 (7.2)
A3/A4	29 (19.0)	32 (20.9)	61 (19.9)
A3/A5	12 (7.8)	14 (9.1)	26 (8.5)
A4/A4	23 (15.0)	17 (11.1)	40 (13.1)
A4/A5	17 (11.1)	17 (11.1)	34 (11.1)
A5/A5	3 (2.0)	5 (3.2)	8 (2.6)
Total subjects	153 (100.0)	153 (100.0)	306 (100.0)
<b>Allele</b>			
A1	36 (11.8)	14 (4.6)	50 (8.2)
A2	32 (10.5)	48 (15.7)	80 (13.1)
A3	82 (26.8)	89 (29.1)	171 (27.9)
A4	110 (35.9)	105 (34.3)	215 (35.1)
A5	46 (15.0)	50 (16.3)	96 (15.7)
Total chromosomes	306 (100.0)	306 (100.0)	612 (100.0)

each component of the BDNF-LCPR and the Val66-Met, while linkage disequilibrium within the three components were much weaker (Supplementary Table S1). The deletion of the first component of the BDNF-LCPR was completely linked to the Val66 allele; the (CA)<sub>del</sub>(CG)<sub>del</sub> allele was completely linked to the Val 66 allele, while the Val66 allele was linked to any of the (CA)<sub>del/2</sub>(CG)<sub>del/4/5</sub> alleles.

#### Luciferase reporter gene assay in primary cultured neurons

Figure 2b shows observed RLEs for the major four alleles of the BDNF-LCPR, compared to RLE without insertion of such alleles (empty pGL3-Promoter vector). RLE decreased due to insertion of the polymorphic region for all the alleles compared to the empty pGL3-Promoter vector, suggesting that the BDNF-LCPR and its flanking region may have a silencer-like effect on transcriptional activity. When RLE was compared among the four alleles, there was a significant difference ( $F=5.9$ ,  $df=3$ ,  $32$ ,  $P=0.003$ , ANOVA). RLE for the A1 allele was the smallest among the four alleles. When RLE for the A1 allele was compared to that for the remaining three alleles combined, the difference was significant ( $t=-3.4$ ,  $df=34$ ,  $P=0.002$ ), providing evidence suggesting that

the A1 allele is associated with lower transcriptional activity.

#### Discussion

The present study demonstrated that a microsatellite polymorphism of the BDNF gene originally reported as a 'GT repeat'<sup>17</sup> is not a simple dinucleotide repeat, but a very complex structure of polymorphism, containing three types of dinucleotide repeats, insertion/deletion, and nucleotide substitutions, which is consistent in part with a recent report.<sup>27</sup> We therefore designated this region as BDNF-linked complex polymorphic region (BDNF-LCPR). The nucleotide sequences were determined by combination of pyrosequencing together with direct sequencing after cloning. Thus sequencing errors are unlikely. As a result, a total of 23 novel allelic variants were detected, although only five alleles had been identified in the original report.<sup>17</sup> We obtained evidence suggesting an association between the BDNF-LCPR and bipolar disorder. This is in accordance with a previous study<sup>8</sup> that reported a significant association between this polymorphism and bipolar disorder. However, detected alleles and their distribution considerably differ between this previous study<sup>8</sup> and the current study since the former genotyped the polymorphism by fragment-size analysis. We detected multiple alleles for each fragment size; for example, the A2 and A3 alleles had the same fragment size (407 bp, see Table 2). Therefore, fragment size analysis is not enough to perform an association study on the BDNF-LCPR.

Of note, the microsatellite corresponding to the BDNF-LCPR and its flanking region are conserved in rodents at similar location relative to the translation initiation site of the BDNF gene (1065 bp upstream in humans, 921 bp in rats, and 963 bp in mice). The nucleotide sequences flanking the microsatellite were highly homologous between humans and rodents (rat: 68% and mouse 66%, according to our calculation based on sequences from GenBank accession number AABR03134358.1 for rat and AY057907 for mouse). We then examined whether the BDNF-LCPR is associated with transcriptional activity in an allele-dependent manner, using luciferase reporter gene assay on primary cultured neurons from the rat brain cortex. The results provided evidence that the A1 allele is associated with lower transcriptional activity, compared to the other major alleles. This is interesting because the A1 allele, which is 12 or 16 bp shorter than the other major alleles (see Table 2), were found to be increased in patients with bipolar disorder, compared to controls. These results suggest that the A1 allele plays a role in giving susceptibility to bipolar disorder by reducing transcriptional activity of the BDNF gene. Since the A1 allele has deletion of the first component of the BDNF-LCPR, and this deletion was significantly more common in patients than in controls, it is possible that such deletion might be responsible for altering transcriptional

**Table 4** Allelic association analysis of each of the three components of the BDNF-LCPR with bipolar disorder

Allele	Counts (%)		
	Patients	Controls	Significance (P-value)
1. (CA) <sub>del/2</sub> (CG) <sub>del/4/5</sub>			
Del-del	44 (14.6)	25 (8.3)	0.015
2-4	38 (12.6)	54 (17.9)	0.07
2-5	220 (72.8)	223 (73.8)	0.78
Total chromosomes <sup>a</sup>	302 (100)	302 (100)	0.044 <sup>b</sup>
2. (CA) <sub>9-15</sub>			
9	1 (0.3)	0 (0.0)	0.24
10	6 (2.0)	6 (1.9)	1.00
11	7 (2.3)	4 (1.3)	0.36
12	160 (52.3)	168 (54.9)	0.52
13	128 (41.8)	122 (39.9)	0.62
14	3 (1.0)	6 (2.0)	0.31
15	1 (0.3)	0 (0.0)	0.24
Total chromosomes	306 (100)	306 (100)	1.00 <sup>b</sup>
3. (GA) <sub>2/3</sub>			
2	107 (35.0)	117 (38.2)	0.40
3	199 (65.0)	189 (61.8)	0.40
Total chromosomes	306 (100)	306 (100)	0.44 <sup>b</sup>

Individual alleles were tested for association by grouping all others together and applying the  $\chi^2$ -test (df = 1).

<sup>a</sup>For the (CA)<sub>del/2</sub>(CG)<sub>del/4/5</sub>, four individuals who carried an exceptionally rare variant (see Table 2) were excluded from the analysis.

<sup>b</sup>Global P-values were estimated by the permutation test with 10 000 simulations, correcting for multiple testing.

**Table 5** Haplotype-based association analysis for the BDNF-linked complex polymorphic region (BDNF-LCPR) and Val66Met polymorphism in patients with bipolar disorder and controls

Haplotype	Counts (%)		
	Patients	Controls	Significance (P-value)
A1-Val	36 (11.8)	14 (4.6)	0.001
A2-Val	32 (10.4)	48 (15.7)	0.054
A3-Val	82 (26.8)	89 (29.1)	0.53
A4-Val	0 (0.0)	1 (0.3)	0.23
A4-Met	110 (36.0)	104 (34.0)	0.61
A5-Val	35 (11.4)	39 (12.7)	0.62
A5-Met	11 (3.6)	11 (3.6)	1.0
Total chromosomes	306 (100)	306 (100)	0.0069 <sup>a</sup>

Individual haplotypes were tested for association by grouping all others together and applying the  $\chi^2$  test (df = 1).

<sup>a</sup>Global P-value was estimated by the permutation test with 10 000 simulations.

activity and conferring the susceptibility. Our result is in line with a recent finding that BDNF protein was reduced in postmortem brains of patients with bipolar disorder, compared to controls.<sup>4</sup>

In previous studies<sup>9-10</sup> that reported a positive association between the Val66met polymorphism of the BDNF gene and bipolar disorder, the Val66 allele

was consistently found to be the risk allele. However, other studies<sup>11-16</sup> failed to find such an association. The Val66Met polymorphism has been found to have functional effects. The Met66 allele was associated with poorer episodic memory, abnormal hippocampal activation, and lower hippocampal *n*-acetyl aspartate in humans and that the Met66 allele showed lower

depolarization-induced secretion and failed to localize to secretory granules or synapses in neurons.<sup>28</sup> The relationship between the Met66 allele and poorer episodic memory has been further demonstrated.<sup>29</sup> Since impairment in verbal episodic memory is one of the most consistently reported cognitive problems in individuals with bipolar disorder,<sup>30,31</sup> it is not feasible that the Val66 allele, but not the Met66 one, has consistently been reported to be the risk allele for bipolar disorder.<sup>8-10</sup> In our linkage disequilibrium analysis between the BDNF-LCPR and the Val66Met polymorphisms, the A1 allele was completely linked to the Val66 allele, which may explain, at least in part, the inconsistent results in the previous studies. That is, the A1 allele might be a responsible allele; however, its linkage to the Val66 allele have made the Val66 allele over-represented in some samples but not in the other samples, since the Val66 allele is linked to not only the A1 allele but also A2, A3, and A5 alleles. To demonstrate this hypothesis, the association between the BDNF-LCPR and bipolar disorder should be reevaluated based on the current findings. In addition, studies examining the possible association of the BDNF-LCPR with brain structure and functions are warranted.

Several studies have performed an association study between the 'GT repeat' and schizophrenia, which have also yielded conflicting results.<sup>13,19,20,32-35</sup> To resolve the inconsistent findings, further studies based on the current information are required.

In conclusion, we demonstrated that a microsatellite of the BDNF gene, which was originally reported as a 'GT repeat'<sup>17</sup> is not a simple dinucleotide repeat, but has a complex structure of polymorphism. Association analysis and luciferase reporter gene assay suggest that the BDNF-LCPR is a functional polymorphism that confers susceptibility to bipolar disorder and affects transcriptional activity.

### Acknowledgments

We thank Shizuno T and Okada K for their technical assistance in laboratory. This study was supported by the Health and Labor Sciences Research Grants (Research on Psychiatric and Neurological Diseases and Mental Health), the Japan Health Sciences Foundation (Research on Health Sciences focusing on Drug innovation) and Mitsubishi Pharma Research Foundation (HK).

### References

- Maisonpierre PC, Belluscio L, Friedman B, Alderson RF, Wiegand SJ, Furth ME et al. NT-3, BDNF, and NGF in the developing rat nervous system: parallel as well as reciprocal patterns of expression. *Neuron* 1990; **5**: 501-509.
- Thoenen H. Neurotrophins and neuronal plasticity. *Science* 1995; **270**: 593-598.
- Duman RS. Synaptic plasticity and mood disorders. *Mol Psychiatry* 2002; **7**(Suppl 1): S29-S34.
- Knable MB, Barci BM, Webster MJ, Meador-Woodruff J, Torrey EF. Molecular abnormalities of the hippocampus in severe psychiatric illness: postmortem findings from the Stanley Neuropathology Consortium. *Mol Psychiatry* 2004; **9**: 609-620.
- Nibuya M, Morinobu S, Duman RS. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci* 1995; **15**: 7539-7547.
- Hashimoto R, Takei N, Shimazu K, Christ L, Lu B, Chuang DM. Lithium induces brain-derived neurotrophic factor and activates TrkB in rodent cortical neurons: an essential step for neuroprotection against glutamate excitotoxicity. *Neuropharmacology* 2002; **43**: 1173-1179.
- Green E, Craddock N. Brain-derived neurotrophic factor as a potential risk locus for bipolar disorder: evidence, limitations, and implications. *Curr Psychiatry Rep* 2003; **5**: 469-476.
- Neves-Pereira M, Mundo E, Muglia P, King N, Macciardi F, Kennedy JL. The brain-derived neurotrophic factor gene confers susceptibility to bipolar disorder: evidence from a family-based association study. *Am J Hum Genet* 2002; **71**: 651-655.
- Sklar P, Gabriel SB, McInnis MG, Bennett P, Lim YM, Tsan G et al. Family-based association study of 76 candidate genes in bipolar disorder: BDNF is a potential risk locus. *Mol Psychiatry* 2002; **7**: 579-593.
- Geller B, Badner JA, Tillman R, Christian SL, Bolhofner K, Cook Jr EH. Linkage disequilibrium of the brain-derived neurotrophic factor Val66Met polymorphism in children with a prepubertal and early adolescent bipolar disorder phenotype. *Am J Psychiatry* 2004; **161**: 1698-1700.
- Oswald P, Del-Favero J, Massat I, Souery D, Claes S, Van Broeckhoven C et al. Non-replication of the brain-derived neurotrophic factor (BDNF) association in bipolar affective disorder: a Belgian patient-control study. *Am J Med Genet* 2004; **129B**: 34-35.
- Skibinska M, Hauser J, Czernski PM, Leszczynska-Rodziewicz A, Kosmowska M, Kapelski P et al. Association analysis of brain-derived neurotrophic factor (BDNF) gene Val66Met polymorphism in schizophrenia and bipolar affective disorder. *World J Biol Psychiatry* 2004; **5**: 215-220.
- Neves-Pereira M, Cheung JK, Pasdar A, Zhang F, Breen G, Yates P et al. BDNF gene is a risk factor for schizophrenia in a Scottish population. *Mol Psychiatry* 2005; **10**: 208-212.
- Hong CJ, Huo SJ, Yen FC, Tung CL, Pan GM, Tsai SJ. Association study of a brain-derived neurotrophic-factor genetic polymorphism and mood disorders, age of onset and suicidal behavior. *Neuropsychobiology* 2003; **48**: 186-189.
- Nakata K, Ujike H, Sakai A, Uchida N, Nomura A, Imamura T et al. Association study of the brain-derived neurotrophic factor (BDNF) gene with bipolar disorder. *Neurosci Lett* 2003; **337**: 17-20.
- Kunugi H, Iijima Y, Tatsumi M, Yoshida M, Hashimoto R, Kato T et al. No association between the Val66Met polymorphism of the brain-derived neurotrophic factor gene and bipolar disorder in a Japanese population: a multicenter study. *Biol Psychiatry* 2004; **56**: 376-378.
- Pröschel M, Saunders A, Roses AD, Muller CR. Dinucleotide repeat polymorphism at the human gene for the brain-derived neurotrophic factor (BDNF). *Hum Mol Genet* 1992; **1**: 353.
- Strauss J, Barr CL, George CJ, King N, Shaikh S, Devlin B et al. Association study of brain-derived neurotrophic factor in adults with a history of childhood onset mood disorder. *Am J Med Genet* 2004; **131B**: 16-19.
- Krebs MO, Guillin O, Bourdell MC, Schwartz JC, Olie JP, Poirier MF et al. Brain derived neurotrophic factor (BDNF) gene variants association with age at onset and therapeutic response in schizophrenia. *Mol Psychiatry* 2000; **5**: 558-562.
- Muglia P, Vicente AM, Verga M, King N, Macciardi F, Kennedy JL. Association between the BDNF gene and schizophrenia. *Mol Psychiatry* 2003; **8**: 146-147.
- Comings DE. Polygenic inheritance and micro/minisatellites. *Mol Psychiatry* 1998; **3**: 21-31.
- American Psychiatric Association. *American Psychiatric Association: Diagnostic and Statistical Manual of Mental Disorders*, 4th edn. American Psychiatric Association: Washington DC, 1994.
- Numakawa T, Yamagishi S, Adachi N, Matsumoto T, Yokomaku D, Yamada M et al. Brain-derived neurotrophic factor-induced potentiation of Ca(2+) oscillations in developing cortical neurons. *J Biol Chem* 2002; **277**: 6520-6529.

- 24 Tadokoro K, Hashimoto R, Tatsumi M, Kamijima K, Kunugi H. Analysis of enhancer activity of a dinucleotide repeat polymorphism in the neurotrophin-3 gene and its association with bipolar disorder. *Neuropsychobiology* 2004; **50**: 206–210.
- 25 Aoyama M, Asai K, Shishikura T, Kawamoto T, Miyachi T, Yokoi T *et al*. Human neuroblastomas with unfavorable biologies express high levels of brain-derived neurotrophic factor mRNA and a variety of its variants. *Cancer Lett* 2001; **164**: 51–60.
- 26 Garzon D, Yu G, Fahnestock M. A new brain-derived neurotrophic factor transcript and decrease in brain-derived neurotrophic factor transcripts 1, 2 and 3 in Alzheimer's disease parietal cortex. *J Neurochem* 2002; **82**: 1058–1064.
- 27 Koizumi H, Hashimoto K, Shimizu E, Iyo M, Mashimo Y, Hata A. Further analysis of microsatellite marker in the BDNF gene. *Am J Med Genet* 2005; **135**: 103.
- 28 Egan MF, Kojima M, Callicott JH, Goldberg TE, Kolachana BS, Bertolino A *et al*. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell* 2003; **112**: 257–269.
- 29 Dempster E, Touloupoulou T, McDonald C, Bramon E, Walshe M, Filbey F *et al*. Association between BDNF val66 met genotype and episodic memory. *Am J Med Genet* 2005; **134**: 73–75.
- 30 Sweeney JA, Kmiec JA, Kupfer DJ. Neuropsychologic impairments in bipolar and unipolar mood disorders on the CANTAB neurocognitive battery. *Biol Psychiatry* 2000; **48**: 674–684.
- 31 Deckersbach T, Savage CR, Reilly-Harrington N, Clark L, Sachs G, Rauch SL. Episodic memory impairment in bipolar disorder and obsessive-compulsive disorder: the role of memory strategies. *Bipolar Disord* 2004; **6**: 233–244.
- 32 Sasaki T, Dai XY, Kuwata S, Fukuda R, Kunugi H, Hattori M *et al*. Brain-derived neurotrophic factor gene and schizophrenia in Japanese subjects. *Am J Med Genet* 1997; **74**: 443–444.
- 33 Hawi Z, Straub RE, O'Neill A, Kendler KS, Walsh D, Gill M. No linkage or linkage disequilibrium between brain-derived neurotrophic factor (BDNF) dinucleotide repeat polymorphism and schizophrenia in Irish families. *Psychiatry Res* 1998; **81**: 111–116.
- 34 Wassink TH, Nelson JJ, Crowe RR, Andreasen NC. Heritability of BDNF alleles and their effect on brain morphology in schizophrenia. *Am J Med Genet* 1999; **88**: 724–728.
- 35 Virgos C, Martorell L, Valero J, Figuera L, Civeira F, Joven J *et al*. Association study of schizophrenia with polymorphisms at six candidate genes. *Schizophr Res* 2001; **49**: 65–71.

Supplementary Information accompanies the paper on the Molecular Psychiatry website (<http://www.nature.com/mp>).

# The association between the Val158Met polymorphism of the catechol-O-methyl transferase gene and morphological abnormalities of the brain in chronic schizophrenia

Takashi Ohnishi,<sup>1,2,4</sup> Ryota Hashimoto,<sup>2</sup> Takeyuki Mori,<sup>1,2</sup> Kiyotaka Nemoto,<sup>1</sup> Yoshiya Moriguchi,<sup>1</sup> Hidehiro Iida,<sup>4</sup> Hiroko Noguchi,<sup>2</sup> Tetsuo Nakabayashi,<sup>2,3</sup> Hiroaki Hori,<sup>2,3</sup> Mayu Ohmori,<sup>3</sup> Ryoutaro Tsukue,<sup>3</sup> Kimitaka Anami,<sup>3</sup> Naotugu Hirabayashi,<sup>3</sup> Seiichi Harada,<sup>3</sup> Kunimasa Arima,<sup>3</sup> Osamu Saitoh<sup>3</sup> and Hiroshi Kunugi<sup>2</sup>

<sup>1</sup>Department of Radiology, National Center Hospital of Mental, Nervous and Muscular Disorders, National Center of Neurology and Psychiatry, <sup>2</sup>Department of Mental Disorder Research, National Institute of Neuroscience, National Center of Neurology and Psychiatry, <sup>3</sup>Department of Psychiatry, National Center Hospital of Mental, Nervous, and Muscular Disorders, National Center of Neurology and Psychiatry, Tokyo and <sup>4</sup>Department of Investigative Radiology, Research Institute, National Cardiovascular Center, Osaka, Japan

Correspondence to: Takashi Ohnishi, Department of Radiology, National Center Hospital of Mental, Nervous, and Muscular Disorders, National Center of Neurology and Psychiatry 4-1-1 Ogawa Higashi, Kodaira City,

Tokyo, Japan 187-0031

E-mail: tohnishi@hotmail.com

**The catechol-O-methyl transferase (COMT) gene is considered to be a promising schizophrenia susceptibility gene. A common functional polymorphism (Val158Met) in the COMT gene affects dopamine regulation in the prefrontal cortex (PFC). Recent studies suggest that this polymorphism contributes to poor prefrontal functions, particularly working memory, in both normal individuals and patients with schizophrenia. However, possible morphological changes underlying such functional impairments remain to be clarified. The aim of this study was to examine whether the Val158Met polymorphism of the COMT gene has an impact on brain morphology in normal individuals and patients with schizophrenia. The Val158Met COMT genotype was obtained for 76 healthy controls and 47 schizophrenics. The diagnostic effects, the effects of COMT genotype and the genotype-diagnosis interaction on brain morphology were evaluated by using a voxel-by-voxel statistical analysis for high-resolution MRI, a tensor-based morphometry. Patients with schizophrenia demonstrated a significant reduction of volumes in the limbic and paralimbic systems, neocortical areas and the subcortical regions. Individuals homozygous for the Val-COMT allele demonstrated significant reduction of volumes in the left anterior cingulate cortex (ACC) and the right middle temporal gyrus (MTG) compared to Met-COMT carriers. Significant genotype-diagnosis interaction effects on brain morphology were noted in the left ACC, the left parahippocampal gyrus and the left amygdala-uncus. No significant genotype effects or genotype-diagnosis interaction effects on morphology in the dorsolateral PFC (DLPFC) were found. In the control group, no significant genotype effects on brain morphology were found. Schizophrenics homozygous for the Val-COMT showed a significant reduction of volumes in the bilateral ACC, left amygdala-uncus, right MTG and left thalamus compared to Met-COMT schizophrenics. Our findings suggest that the Val158Met polymorphism of the COMT gene might contribute to morphological abnormalities in schizophrenia.**

**Keywords:** schizophrenia; polymorphism; COMT; ACC; DLPFC



**Abbreviations:** ACC = anterior cingulate cortex; COMT = catechol-*O*-methyl transferase; DLPFC = dorsolateral prefrontal cortex; FDR = false discovery rate; IQ = intelligence quotient; JART = Japanese version of National Adult Reading Test; ROI = region of interest; SPM = statistical parametric mapping; TBM = tensor-based morphometry

Received July 15, 2005. Revised September 21, 2005. Accepted October 27, 2005. Advance Access publication December 5, 2005

## Introduction

Schizophrenia is a severe neuropsychiatric disorder with deficits of multiple domains of cognitive functions, volition and emotion. Family and twin studies have provided cumulative evidence for a genetic basis of schizophrenia (Kendler, 1983; McGue *et al.*, 1983; Sullivan *et al.*, 2003); however, identification of the underlying susceptibility loci has been limited. Collective data have suggested that the aetiology of schizophrenia involves the interplay of complex polygenic influences and environmental risk factors operating on brain maturational processes (Harrison *et al.*, 2005).

*In vivo* neuroimaging studies have demonstrated that brain abnormalities should play an important role in the pathophysiology of schizophrenia. Structural MRI studies have demonstrated relatively consistent brain abnormalities in patients with schizophrenia, such as enlargement of the ventricular system and regional volume decrease in the temporal lobe structures (Gaser *et al.*, 2001; Okubo *et al.*, 2001; Shenton *et al.*, 2001; Davidson and Heinrichs, 2003). Studies with schizophrenics and their healthy siblings demonstrate that even healthy siblings share some of morphological abnormalities observed in schizophrenia (Steel *et al.*, 2002; Gogtay *et al.*, 2003). A recent morphological MR study revealed that a common polymorphism of the brain-derived neurotrophic factor, one of the well-known schizophrenia susceptibility genes, affected the anatomy of the hippocampus and prefrontal cortex (PFC) in healthy individuals (Pezawas *et al.*, 2004). Furthermore, some studies have suggested that environmental factors interact with genetic factors (Cannon *et al.*, 1993; Nelson *et al.*, 2004). For example, obstetric complications are well known non-genetic risk factors of schizophrenia. However, a previous study suggested that obstetric complications might induce brain morphological abnormalities in schizophrenics and their siblings, but not in comparison with subjects at low genetic risk for schizophrenia (Cannon *et al.*, 1993). These facts suggest that genetic factors should have considerable impact on brain morphology in patients with schizophrenia.

Catechol-*O*-methyl transferase (COMT) is a promising schizophrenia susceptibility gene because of its role in monoamine metabolism (Goldberg *et al.*, 2003; Stefanis *et al.*, 2004; Harrison *et al.*, 2005). A common single nucleotide polymorphism (SNP) of the COMT gene producing an amino acid substitution of methionine (met) to valine (val) at position 108/158 (Val158Met) affects dopamine regulation in the PFC (Palmatier *et al.*, 1999). This polymorphism impacts on the stability of the enzyme, such that the Val-COMT allele has significantly lower enzyme activity than the Met-COMT allele (Weinberger *et al.*, 2001; Chen *et al.*, 2004). Several

studies have revealed that the Val-COMT allele is associated with poorer performances, compared to the Met-COMT allele, in cognitive tasks of frontal function such as the Wisconsin Card Sorting Test (WCST) and N-back task (Egan *et al.*, 2001; Weinberger *et al.*, 2001; Goldberg *et al.*, 2003). The underlying mechanism of such behavioural differences may be related to lower prefrontal dopamine levels arising from higher dopamine catabolism mediated by the Val-COMT allele (Chen *et al.*, 2004; Tunbridge *et al.*, 2004).

The results of studies on the association between the Val158Met polymorphism and schizophrenia have, however, been controversial (Daniels *et al.*, 1996; Kunugi *et al.*, 1997; Ohmori *et al.*, 1998; Norton *et al.*, 2002; Galderisi *et al.*, 2005; Ho *et al.*, 2005). The result of a meta-analysis was even more inconclusive (Fan *et al.*, 2005). Such inconsistency was also found in associations between frontal functions and the Val158Met polymorphism (Egan *et al.*, 2001; Weinberger *et al.*, 2001; Goldberg *et al.*, 2003; Ho *et al.*, 2005). The possible morphological changes due to the COMT gene might be present and play a role in susceptibility to schizophrenia and in giving rise to impaired frontal functions. However, morphological changes underlying functional impairments remain to be clarified.

A recent advancement of methods for MR volumetry, such as voxel-based morphometry and deformation-based morphometry [or tensor-based morphometry (TBM)], allows us to explore and analyse brain structures of schizophrenics (Wright *et al.*, 1995; Gaser *et al.*, 2001). Using TBM techniques, we investigated the association between the Val158-Met polymorphism of the COMT gene and brain morphology in normal individuals and patients with schizophrenia. The aim of this study was to clarify whether there are significant genotype and/or genotype-disease interaction effects on brain morphology.

## Methods

### Subjects

Seventy-six healthy subjects and forty-seven patients with schizophrenia participated in the study. All the subjects were biologically unrelated Japanese. Written informed consent was obtained from all the subjects in accordance with ethical guidelines set by a local ethical committee. All normal subjects were screened using a questionnaire on medical history and excluded if they had neurological, psychiatric or medical conditions that could potentially affect the CNS, such as substance abuse or dependence, atypical headache, head trauma with loss of consciousness, asymptomatic or symptomatic cerebral infarctions detected by T<sub>2</sub>-weighted MRI, hypertension, chronic lung

disease, kidney disease, chronic hepatic disease, cancer, or diabetes mellitus. The patients were diagnosed on the basis of DSM-IV criteria, information from medical records and a clinical interview. All patients were stable and/or partially remitted at the time of MR measurement and neuropsychological tests.

According to genotypes, each group (control and schizophrenia) was categorized into three groups; the homozygous Val-COMT group (control:  $n = 38$ , two were left-handed, schizophrenia:  $n = 19$ , one was left-handed), the Val/Met-COMT group (control:  $n = 25$ , three were left-handed, schizophrenia:  $n = 22$ , all were right-handed) and the remaining homozygous Met-COMT group (control:  $n = 13$ , all were right-handed, schizophrenia:  $n = 6$ , all were right-handed). Because of the small number of subjects with homozygous Met-COMT, the Val/Met-COMT and homozygous Met-COMT groups were combined and treated as one group, the Met-COMT carriers. Table 1 shows the characteristics of each group. All groups were of comparable age, gender ( $\chi^2$  test,  $df = 3$ ,  $P = 0.38$ ) and handedness ( $\chi^2$ -test,  $df = 3$ ,  $P = 0.53$ ). No genotype effects and genotype-diagnosis interaction effects were found in years of education, scores of full scale Intelligence Quotient (IQ) and scores of premorbid IQ [Japanese version of National Adult Reading Test (JART) score], however, patients who had fewer years of education ( $P < 0.0001$ ), had lower scores of both full scale IQ and JART ( $P < 0.001$ ). The duration of illness, medication and hospitalization, the age at disease onset and drug dose (chlorpromazine equivalent) of those homozygous for the Val-COMT did not differ from the Met-COMT carriers.

### SNP genotyping

Venous blood was drawn from subjects and genomic DNA was extracted from whole blood according to the standard procedures. The Val158Met polymorphism of the COMT gene (dbSNP accession: rs4680) was genotyped using the TaqMan 5'-exonuclease allelic discrimination assay, described previously (Hashimoto *et al.*, 2004, 2005). Briefly, primers and probes for detection of the SNP are: forward primer 5'-GACTGTGCCGACATCAC-3', reverse primer 5'-CAGGCATGCACACCTTGTC-3', probe 1 5'-VIC-TTTCGCTG-GCGTGAAG-MGB-3' and probe 2 5'-FAM-CGCTGGCATGAAG-MGB-3'. PCR cycling conditions were: at 95°C for 10 min, 50 cycles of 92°C for 15 s and 60°C for 1 min.

### MRI procedures

All MR studies were performed on a 1.5 tesla Siemens Magnetom Vision plus system. A three dimensional (3D) volumetric acquisition of a T<sub>1</sub>-weighted gradient echo sequence produced a gapless series of thin sagittal sections using an MPRage sequence (TE/TR, 4.4/11.4 ms; flip angle, 15°; acquisition matrix, 256 × 256; 1 NEX, field of view, 31.5 cm; slice thickness, 1.23 mm).

### Image analysis (TBM)


The basic principle of TBM is to analyse the local deformations of an image and to infer local differences in brain structure. In TBM, MRI scans of individual subjects are mapped to a template image with three-dimensional (3D) non-linear normalization routines. Local deformations were estimated by a univariate Jacobian approach. The basic principle of TBM is the same as a method used in a previous report described as deformation-based morphometry (Gaser *et al.*, 2001). Firstly, inhomogeneities in MR images were corrected using a bias correction function in statistical parametric mapping (SPM2),

then the corrected image was scalp-edited by masking with a probability image of brain tissue obtained from each image using a segmentation function in SPM2. Using a linear normalization algorithm in SPM2, all brains were resized to a voxel size of 1.5 mm and adjusted for orientation and overall width, length and height (Fig. 1A). Therefore, brains were transformed to the anatomical space of a template brain whose space is based on Talairach space (Talairach and Tournoux, 1988). Subsequent non-linear normalization introduced local deformations to each brain to match it to the same scalp-edited template brain (Fig. 1C). The non-linear transformation was done using the high-dimension-warping algorithm (Ashburner and Friston, 2004). After the high dimensional warping, each image (Fig. 1B) looks similar to the template (Fig. 1C). Figure 2 demonstrated a mean MR image of 76 controls (left) and a mean MR image of 47 schizophrenics after high dimensional warping (Fig. 2). We obtained 3D deformation fields for every brain (Fig. 1D). Each of these 3D deformation fields consists of displacement vectors for every voxel, which describe the 3D displacement needed to locally deform the brain to match it to the template. We calculated the Jacobian determinants to obtain voxel by voxel parametric maps of local volume change relative to the template brain (Fig. 1E). The local Jacobian determinant is a parameter commonly used in continuum mechanics (Gurtin, 1987), which characterizes volume changes, such as local shrinkage or enlargement caused by warping. The parametric maps of Jacobian determinants were analysed using SPM2, which implements a 'general linear model'. To test hypotheses about regional population effects and interaction, data were analysed by an analysis of covariance (ANCOVA) without global normalization. There was no significant difference in age among the four groups, however, patients with schizophrenia, particularly those homozygous for the Val-COMT allele, were older than controls. Therefore, we treated age and years of education and scores of JART as nuisance variables. Since TBM explores the entire brain (grey matter, CSF space and white matter) at once, the search volume of TBM has a large number of voxels and since our interest was in morphological changes in the grey matter and CSF space, we excluded white matter tissue from analyses by using an explicit mask (Fig. 1F). We used  $P < 0.001$ , corrected for multiple comparisons with false discovery rate (FDR)  $< 0.05$  as a statistical threshold. The resulting sets of  $t$  values constituted the statistical parametric maps {SPM ( $t$ )}. Firstly, we estimated the main effects, the genotype effect in total subjects (the Val/Val-COMT versus the Met-COMT carriers) and the diagnostic effect (schizophrenia versus controls) and then the genotype-diagnosis interaction effect was estimated. Furthermore, the effects of genotypes in each group (controls carrying the Val/Val-COMT gene versus controls carrying the Met-COMT gene and schizophrenics carrying the Val/Val-COMT gene versus schizophrenics carrying the Met-COMT gene) were estimated within the ANCOVA design matrix. Anatomical localization accorded both to MNI coordinates and Talairach coordinates obtained from M. Brett's transformations ([www.mrc-cbu.cam.ac.uk/Imaging/mninspace.html](http://www.mrc-cbu.cam.ac.uk/Imaging/mninspace.html)) and are presented as Talairach coordinates (Talairach and Tournoux, 1988). Since previous studies have demonstrated the association between the Val158Met polymorphism and the dorsolateral PFC (DLPFC), we applied an additional hypothesis-driven region of interest (ROI) method to test regional population effects in the DLPFC. For this ROI analysis, we used the Wake Forest University PickAtlas (Maldjian *et al.*, 2003) within the ANCOVA design matrix for SPM analysis. We set  $P < 0.05$  (uncorrected) with a small volume correction ( $P < 0.05$  within the ROI) to assess grey matter volume changes in the DLPFC (Brodmann area 46, 9 and 8).


Table 1 Subject characteristics

	Control Val/Val	Met carriers	Schizophrenia Val/Val	Met carriers	Diagnosis F (P)	Genotype F (P)*	Genotype by diagnosis F (P)
Number of subjects	38	38	19	28			
Gender (M/F)	16 out of 22	14 out of 24	11 out of 8	13 out of 15			
Handedness (R/L)	36 out of 2	35 out of 3	18 out of 1	28 out of 0			
Age (years)	41.47 (13.42)	39.26 (10.6)	45.98 (15.29)	43.05 (10.57)	3.633 (0.059)	1.7 (0.195)	0.21 (0.647)
Education (years)	17 (3.16)	16.06 (2.57)	12.67 (2.43)	13.33 (3.31)	30.855 (<0.0001)	0.047 (0.828)	1.61 (0.208)
Full scale IQ (WAIS-R)	113.42 (12.05)	108.93 (13.58)	80.69 (17.68)	88.958 (22.08)	57.9 (<0.001)	0.29 (0.59)	3.41 (0.068)
JART	78.8 (10.45)	75.42 (13.65)	54.69 (20.74)	62.25 (27.06)	23.366 (<0.001)	0.292 (0.59)	2.014 (0.159)
Wechsler Memory Scale—Revised							
Verbal memory	111.78 (15.001)	111.061 (12.89)	78.0 (21.623)	81.33 (18.57)	86.93 (<0.001)	0.147 (0.702)	0.354 (0.553)
Visual memory	112.1 (8.51)	106.55 (11.99)	74.78 (24.32)	83.29 (20.613)	85.51 (<0.001)	0.204 (0.65)	4.605 (0.03)
General memory	113.31 (13.92)	110.85 (12.22)	74.43 (21.3)	79.33 (19.14)	111.93 (<0.001)	0.135 (0.715)	1.226 (0.27)
Attention/concentration	104.47 (13.25)	102.94 (16.51)	87.79 (19.09)	92.54 (17.38)	16.08 (0.001)	0.228 (0.634)	0.866 (0.14)
Delayed recall	111.88 (15.46)	112.48 (10.08)	77.07 (20.92)	81.21 (19.19)	99.74 (<0.001)	0.52 (0.475)	0.284 (0.59)
WCST (preservative error)	2.5 (3.89)	3.14 (3.90)	12.08 (11.54)	8.52 (10.63)	24.5 (<0.0001)	0.93 (0.34)	1.93 (0.17)
Digit span	11.12 (3.25)	10.77 (3.34)	7.83 (3.93)	9.09 (2.74)	12.165 (0.0007)	0.415 (0.52)	1.28 (0.261)
Onset age			25.38 (10.34)	23.74 (7.992)		0.52	
Duration of illness (years)			19.86 (14.93)	18.84 (9.8)		0.77	
Duration of hospitalization (months)			66 (153.41)	59.59 (91.18)		0.86	
Duration of medication (years)			12.86 (14.21)	16.4 (9.89)		0.29	
Drug dose of typical antipsychotic drugs (mg/day, chlorpromazine equivalent)			617.9 (720.18)	700.38 (752.67)		0.69	
Drug dose of atypical antipsychotic drugs (mg/day, chlorpromazine equivalent)							
Mean (standard deviation)			282.3 (428.29)	340.23 (482.19)		0.66	

Mean (standard deviation); WAIS-R = Wechsler Adult Intelligence Scale—Revised; JART = Japanese version of National Adult Reading Test; WCST = Wisconsin Card Sorting Test.



**Fig. 1** Steps of analysis for tensor-based morphometry. An example is shown for a single subject in one axial slice. The single object brain (A) has been corrected for orientation and overall size to the template brain (C). Non-linear spatial normalization removes most of the anatomical differences between the two brains by introducing local deformations to the object brain, which then (B) looks as similar as possible to the template. Image (D) shows the deformations applied to the object brain by a deformed grid. Statistical analysis can be done univariate using the local Jacobian determinant as a derivative of the field (E). An explicit mask image (F) was used to explore morphology in the grey matter and CSF space.



**Fig. 2** Mean images after high dimensional warping control subjects and schizophrenics. *Left:* The mean image of warped MR images obtained from 76 controls. Even after averaging, the mean image is not blurred. *Right:* The mean image of warped MR images obtained from 47 schizophrenics. The mean image of schizophrenic looks similar to that of controls.

## Results

### Behavioural data

Patients had a lower full scale IQ, measured by the Wechsler Adult Intelligence Scale—Revised, than controls. They also had a lower expected premorbid IQ measured by a JART,

lower scores of Wechsler Memory Scale—Revised and demonstrated poorer performance of working memory measures such as the number of preservative errors in the WCST and digit span (Table 1). No genotype or genotype-diagnosis interaction effects were found in working memory measures