

32. Benson MA, Newey SE, Martin-Rendon E, Hawkes R, Blake DJ. Dysbindin, a novel coiled-coil-containing protein that interacts with the dystrobrevins in muscle and brain. *J Biol Chem* 2001; **276**: 24232-24241.
33. Blake DJ, Weir A, Newey SE, Davies KE. Function and genetics of dystrophin and dystrophin-related proteins in muscle. *Physiol Rev* 2002; **82**: 291-329.
34. Blake DJ, Kroger S. The neurobiology of duchenne muscular dystrophy: learning lessons from muscle? *Trends Neurosci* 2000; **23**: 92-9.
35. Li W, Zhang Q, Oiso N et al. Hermansky-Pudlak syndrome type 7 (HPS-7) results from mutant dysbindin, a member of the biogenesis of lysosome-related organelles complex 1 (BLOC-1). *Nat Genet* 2003; **35**: 84-89.
36. Huizing M, Boissy RE, Gahl WA. Hermansky-Pudlak syndrome: vesicle formation from yeast to man. *Pigment Cell Res* 2002; **15**: 405-19.
37. Stefansson H, Sarginson J, Kong A et al. Association of neuregulin 1 with schizophrenia confirmed in a Scottish population. *Am J Hum Genet* 2003; **72**: 83-7.
38. Williams NM, Preece A, Spurlock G et al. Support for genetic variation in neuregulin 1 and susceptibility to schizophrenia. *Mol Psychiatry* 2003; **8**: 485-7.
39. Yang JZ, Si TM, Ruan Y et al. Association study of neuregulin 1 gene with schizophrenia. *Mol Psychiatry* 2003; **8**: 706-9.
40. Tang JX, Chen WY, He G et al. Polymorphisms within 5' end of the Neuregulin 1 gene are genetically associated with schizophrenia in the Chinese population. *Mol Psychiatry* 2004; **9**: 11-2.

41. Zhao X, Shi Y, Tang J et al. A case control and family based association study of the neuregulin1 gene and schizophrenia. *J Med Genet* 2004; **41**: 31-4.
42. Li T, Stefansson H, Gudfinnsson E et al. Identification of a novel neuregulin 1 at-risk haplotype in Han schizophrenia Chinese patients, but no association with the Icelandic/Scottish risk haplotype. *Mol Psychiatry* 2004; **9**: 698-704.
43. Corvin AP, Morris DW, McGhee K et al. Confirmation and refinement of an 'at-risk' haplotype for schizophrenia suggests the EST cluster, Hs.97362, as a potential susceptibility gene at the Neuregulin-1 locus. *Mol Psychiatry* 2004; **9**: 208-13.
44. Bakker SC, Hoogendoorn ML, Selten JP et al. Neuregulin 1: genetic support for schizophrenia subtypes. *Mol Psychiatry* 2004; **9**: 1061-3.
45. Hong CJ, Huo SJ, Liao DL, Lee K, Wu JY, Tsai SJ. Case-control and family-based association studies between the neuregulin 1 (Arg38Gln) polymorphism and schizophrenia. *Neurosci Lett* 2004; **366**: 158-61.
46. Petryshen TL, Middleton FA, Kirby A et al. Support for involvement of neuregulin 1 in schizophrenia pathophysiology. *Mol Psychiatry* 2005; **10**: 366-74, 328.
47. Kampman O, Anttila S, Illi A et al. Neuregulin genotype and medication response in Finnish patients with schizophrenia. *Neuroreport* 2004; **15**: 2517-20.
48. Thiselton DL, Webb BT, Neale BM et al. No evidence for linkage or association of neuregulin-1 (NRG1) with disease in the Irish study of high-density schizophrenia families (ISHDSF). *Mol Psychiatry* 2004; **9**: 777-83; image 729.

49. Iwata N, Suzuki T, Ikeda M et al. No association with the neuregulin 1 haplotype to Japanese schizophrenia. *Mol Psychiatry* 2004; **9**: 126-7.
50. Hashimoto R, Straub RE, Weickert CS, Hyde TM, Kleinman JE, Weinberger DR. Expression analysis of neuregulin-1 in the dorsolateral prefrontal cortex in schizophrenia. *Mol Psychiatry* 2004; **9**: 299-307.
51. Law AJ, Lipska BK, Shannon Weickert C et al. Splice variant-specific alterations of Neuregulin-1 gene expression in hippocampus in schizophrenia. *Program No. 109.7. 2004 Abstract Viewer/Itinerary Planner. Washington, DC: Society for Neuroscience, 2004. Online 2003.*
52. Fischbach GD, Rosen KM. ARIA: a neuromuscular junction neuregulin. *Annu Rev Neurosci* 1997; **20**: 429-58.
53. Ozaki M, Sasner M, Yano R, Lu HS, Buonanno A. Neuregulin-beta induces expression of an NMDA-receptor subunit. *Nature* 1997; **390**: 691-4.
54. Yang X, Kuo Y, Devay P, Yu C, Role L. A cysteine-rich isoform of neuregulin controls the level of expression of neuronal nicotinic receptor channels during synaptogenesis. *Neuron* 1998; **20**: 255-70.
55. Rieff HI, Raetzman LT, Sapp DW, Yeh HH, Siegel RE, Corfas G. Neuregulin induces GABA(A) receptor subunit expression and neurite outgrowth in cerebellar granule cells. *J Neurosci* 1999; **19**: 10757-66.
56. Liu Y, Ford B, Mann MA, Fischbach GD. Neuregulins increase alpha7 nicotinic acetylcholine receptors and enhance excitatory synaptic transmission in GABAergic interneurons of the hippocampus. *J Neurosci* 2001; **21**: 5660-9.

57. Gerlai R, Pisacane P, Erickson S. Heregulin, but not ErbB2 or ErbB3, heterozygous mutant mice exhibit hyperactivity in multiple behavioral tasks. *Behav Brain Res* 2000; **109**: 219-27.
58. Buonanno A, Fischbach GD. Neuregulin and ErbB receptor signaling pathways in the nervous system. *Curr Opin Neurobiol* 2001; **11**: 287-96.
59. Steinthorsdottir V, Stefansson H, Ghosh S et al. Multiple novel transcription initiation sites for NRG1. *Gene* 2004; **342**: 97-105.
60. Zhang X, Tochigi M, Ohashi J et al. Association study of the DISC1/TRAX locus with schizophrenia in a Japanese population. *Schizophr Res* 2005.
61. Callicott JH, Straub RE, Pezawas L et al. Variation in DISC1 affects hippocampal structure and function and increases risk for schizophrenia. *Proc Natl Acad Sci U S A* 2005; **102**: 8627-32.
62. Hodgkinson CA, Goldman D, Jaeger J et al. Disrupted in schizophrenia 1 (DISC1): association with schizophrenia, schizoaffective disorder, and bipolar disorder. *Am J Hum Genet* 2004; **75**: 862-72.
63. Hennah W, Varilo T, Kestila M et al. Haplotype transmission analysis provides evidence of association for DISC1 to schizophrenia and suggests sex-dependent effects. *Hum Mol Genet* 2003; **12**: 3151-9.
64. Thomson PA, Wray NR, Millar JK et al. Association between the TRAX/DISC1 locus and both bipolar disorder and schizophrenia in the Scottish population. *Mol Psychiatry* 2005; **10**: 657-68, 616.

65. Kockelkorn TT, Arai M, Matsumoto H et al. Association study of polymorphisms in the 5' upstream region of human DISC1 gene with schizophrenia. *Neurosci Lett* 2004; **368**: 41-5.
66. Devon RS, Anderson S, Teague PW et al. Identification of polymorphisms within Disrupted in Schizophrenia 1 and Disrupted in Schizophrenia 2, and an investigation of their association with schizophrenia and bipolar affective disorder. *Psychiatr Genet* 2001; **11**: 71-8.
67. Sachs NA, Sawa A, Holmes SE, Ross CA, DeLisi LE, Margolis RL. A frameshift mutation in Disrupted in Schizophrenia 1 in an American family with schizophrenia and schizoaffective disorder. *Mol Psychiatry* 2005; **10**: 758-64.
68. Ozeki Y, Tomoda T, Kleiderlein J et al. Disrupted-in-Schizophrenia-1 (DISC-1): mutant truncation prevents binding to NudeE-like (NUDEL) and inhibits neurite outgrowth. *Proc Natl Acad Sci U S A* 2003; **100**: 289-94.
69. Morris JA, Kandpal G, Ma L, Austin CP. DISC1 (Disrupted-In-Schizophrenia 1) is a centrosome-associated protein that interacts with MAP1A, MIPT3, ATF4/5 and NUDEL: regulation and loss of interaction with mutation. *Hum Mol Genet* 2003; **12**: 1591-608.
70. Miyoshi K, Honda A, Baba K et al. Disrupted-In-Schizophrenia 1, a candidate gene for schizophrenia, participates in neurite outgrowth. *Mol Psychiatry* 2003; **8**: 685-94.
71. Millar JK, Christie S, Porteous DJ. Yeast two-hybrid screens implicate DISC1 in brain development and function. *Biochem Biophys Res Commun* 2003; **311**: 1019-25.

72. Sawamura N, Sawamura-Yamamoto T, Ozeki Y, Ross CA, Sawa A. A form of DISC1 enriched in nucleus: Altered subcellular distribution in orbitofrontal cortex in psychosis and substance/alcohol abuse. *Proc Natl Acad Sci U S A* 2005; **102**: 1187-92.
73. Brandon NJ, Schurov I, Camargo LM et al. Subcellular targeting of DISC1 is dependent on a domain independent from the Nudel binding site. *Mol Cell Neurosci* 2005; **28**: 613-24.
74. James R, Adams RR, Christie S, Buchanan SR, Porteous DJ, Millar JK. Disrupted in Schizophrenia 1 (DISC1) is a multicompartimentalized protein that predominantly localizes to mitochondria. *Mol Cell Neurosci* 2004; **26**: 112-22.
75. Murphy KC. Schizophrenia and velo-cardio-facial syndrome. *Lancet* 2002; **359**: 426-30.
76. Goldberg TE, Egan MF, Gscheidle T et al. Executive subprocesses in working memory: relationship to catechol-O-methyltransferase Val158Met genotype and schizophrenia. *Arch Gen Psychiatry* 2003; **60**: 889-96.
77. Gallinat J, Bajbouj M, Sander T et al. Association of the G1947A COMT (Val(108/158)Met) gene polymorphism with prefrontal P300 during information processing. *Biol Psychiatry* 2003; **54**: 40-8.
78. Matsumoto M, Weickert CS, Beltaifa S et al. Catechol O-methyltransferase (COMT) mRNA expression in the dorsolateral prefrontal cortex of patients with schizophrenia. *Neuropsychopharmacology* 2003; **28**: 1521-30.
79. Tunbridge E, Burnet PW, Sodhi MS, Harrison PJ. Catechol-o-methyltransferase (COMT) and proline dehydrogenase (PRODH) mRNAs in the dorsolateral

prefrontal cortex in schizophrenia, bipolar disorder, and major depression.

Synapse 2004; **51**: 112-8.

80. Glatt SJ, Faraone SV, Tsuang MT. Association between a functional catechol O-methyltransferase gene polymorphism and schizophrenia: meta-analysis of case-control and family-based studies. *Am J Psychiatry* 2003; **160**: 469-76.
81. Zou F, Li C, Duan S et al. A family-based study of the association between the G72/G30 genes and schizophrenia in the Chinese population. *Schizophr Res* 2005; **73**: 257-61.
82. Korostishevsky M, Kremer I, Kaganovich M et al. Transmission disequilibrium and haplotype analyses of the G72/G30 locus: Suggestive linkage to schizophrenia in Palestinian Arabs living in the North of Israel. *Am J Med Genet B Neuropsychiatr Genet* 2005.
83. Schumacher J, Jamra RA, Freudenberg J et al. Examination of G72 and D-amino-acid oxidase as genetic risk factors for schizophrenia and bipolar affective disorder. *Mol Psychiatry* 2004; **9**: 203-7.
84. Wang X, He G, Gu N et al. Association of G72/G30 with schizophrenia in the Chinese population. *Biochem Biophys Res Commun* 2004; **319**: 1281-6.
85. Korostishevsky M, Kaganovich M, Cholostoy A et al. Is the G72/G30 locus associated with schizophrenia? single nucleotide polymorphisms, haplotypes, and gene expression analysis. *Biol Psychiatry* 2004; **56**: 169-76.
86. Mulle JG, Chowdari KV, Nimgaonkar V, Chakravarti A. No evidence for association to the G72/G30 locus in an independent sample of schizophrenia families. *Mol Psychiatry* 2005; **10**: 431-3.

87. Chen YS, Akula N, Detera-Wadleigh SD et al. Findings in an independent sample support an association between bipolar affective disorder and the G72/G30 locus on chromosome 13q33. *Mol Psychiatry* 2004; **9**: 87-92; image 5.
88. Hattori E, Liu C, Badner JA et al. Polymorphisms at the G72/G30 gene locus, on 13q33, are associated with bipolar disorder in two independent pedigree series. *Am J Hum Genet* 2003; **72**: 1131-40.
89. Zhang F, St Clair D, Liu X et al. Association analysis of the RGS4 gene in Han Chinese and Scottish populations with schizophrenia. *Genes Brain Behav* 2005; **4**: 444-8.
90. Williams NM, Preece A, Spurlock G et al. Support for RGS4 as a susceptibility gene for schizophrenia. *Biol Psychiatry* 2004; **55**: 192-5.
91. Morris DW, Rodgers A, McGhee KA et al. Confirming RGS4 as a susceptibility gene for schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 2004; **125**: 50-3.
92. Chen X, Dunham C, Kendler S et al. Regulator of G-protein signaling 4 (RGS4) gene is associated with schizophrenia in Irish high density families. *Am J Med Genet B Neuropsychiatr Genet* 2004; **129**: 23-6.
93. Chowdari KV, Mirnics K, Semwal P et al. Association and linkage analyses of RGS4 polymorphisms in schizophrenia. *Hum Mol Genet* 2002; **11**: 1373-80.
94. Cordeiro Q, Talkowski ME, Chowdari KV, Wood J, Nimgaonkar V, Vallada H. Association and linkage analysis of RGS4 polymorphisms with schizophrenia and bipolar disorder in Brazil. *Genes Brain Behav* 2005; **4**: 45-50.

95. Sobell JL, Richard C, Wirshing DA, Heston LL. Failure to confirm association between RGS4 haplotypes and schizophrenia in Caucasians. *Am J Med Genet B Neuropsychiatr Genet* 2005.
96. Grillet N, Pattyn A, Contet C, Kieffer BL, Goriadis C, Brunet JF. Generation and characterization of Rgs4 mutant mice. *Mol Cell Biol* 2005; **25**: 4221-8.
97. Emamian ES, Hall D, Birnbaum MJ, Karayiorgou M, Gogos JA. Convergent evidence for impaired AKT1-GSK3beta signaling in schizophrenia. *Nat Genet* 2004; **36**: 131-137.
98. Schwab SG, Hoefgen B, Hanses C et al. Further Evidence for Association of Variants in the AKT1 Gene with Schizophrenia in a Sample of European Sib-Pair Families. *Biol Psychiatry* 2005.
99. Ikeda M, Iwata N, Suzuki T et al. Association of AKT1 with schizophrenia confirmed in a Japanese population. *Biol Psychiatry* 2004; **56**: 698-700.
100. Ohtsuki T, Inada T, Arinami T. Failure to confirm association between AKT1 haplotype and schizophrenia in a Japanese case-control population. *Mol Psychiatry* 2004; **9**: 981-3.
101. Brazil DP, Park J, Hemmings BA. PKB binding proteins. Getting in on the Akt. *Cell* 2002; **111**: 293-303.
102. Rodgers EE, Theibert AB. Functions of PI 3-kinase in development of the nervous system. *Int J Dev Neurosci* 2002; **20**: 187-97.
103. Brunet A, Datta SR, Greenberg ME. Transcription-dependent and -independent control of neuronal survival by the PI3K-Akt signaling pathway. *Curr Opin Neurobiol* 2001; **11**: 297-305.