

**Figure 4.** Results of the whole-brain voxel-based statistical parametric mapping analysis of the *trans*-1,2,3,5,6,10-beta-hexahydro-6-[4-(methylthio)phenyl]pyrrolo-[2,1-a]isoquinoline ( $[^{11}\text{C}](+)\text{McN-5652}$ ) distribution volumes (DVs). A, Locations of methamphetamine abuser and control differences in  $[^{11}\text{C}](+)\text{McN-5652}$  DVs. Areas with significantly reduced  $[^{11}\text{C}](+)\text{McN-5652}$  DVs in methamphetamine abusers compared with those in controls ( $P < .001$ , corrected for cluster level) are given in Table 2. B, Locations of clusters with significant negative correlations between Aggression Questionnaire scores and  $[^{11}\text{C}](+)\text{McN-5652}$  DVs in methamphetamine abusers ( $P < .05$ , corrected for voxel level) (Table 3). Each top row shows 3-dimensional glass brain views; each bottom row, detected area superimposed onto normal template magnetic resonance images.

amphetamine, and their histories were retrospectively confirmed by the abusers and their family members through

detailed Structured Clinical Interview for *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*–

**Table 2. Voxel-Based Analysis of Regional Brain [<sup>11</sup>C](+)McN-5652 Distribution Volume Reductions in 12 Methamphetamine Abusers Compared With 12 Control Subjects\***

Location	Cluster-Level Analysis		Voxel-Level Analysis		Talairach Coordinates		
	Corrected P Value	Voxels, No.	Corrected P Value	z Score	x	y	z
Right insular cortex	<.001	45 315	.009	5.33	34	13	-4
Left caudate nucleus	NA	NA	.02	5.12	-10	19	-4
Right claustrum	NA	NA	.03	5.02	32	0	-3

Abbreviations: [<sup>11</sup>C](+)McN-5652, *trans*-1,2,3,5,6,10-*beta*-hexahydro-6-[4-(methylthio)phenyl]pyrrolo-[2,1-*a*]isoquinoline; NA, not available.

\*The significance threshold was  $P < .05$  at the corrected voxel level and  $P < .05$  at the corrected cluster level. Coordinates are given in millimeters from the origin at the midpoint of the anterior commissure for voxels of peak significance.

**Table 3. Voxel-Based Analysis of Regional Brain [<sup>11</sup>C](+)McN-5652 Distribution Volumes Negatively Associated With Aggression Questionnaire Scores in 12 Methamphetamine Abusers\***

Location	Cluster-Level Analysis		Voxel-Level Analysis		Talairach Coordinates		
	Corrected P Value	Voxels, No.	Corrected P Value	z Score	x	y	z
Right orbitofrontal cortex	<.001	20	.007	5.49	6	26	-21
Left inferior temporal cortex	<.001	38	.007	5.48	-57	-30	-19
Left orbitofrontal cortex	.001	10	.02	5.17	-10	34	-24
Right anterior cingulate cortex	<.001	12	.03	5.13	10	49	10

Abbreviation: [<sup>11</sup>C](+)McN-5652, *trans*-1,2,3,5,6,10-*beta*-hexahydro-6-[4-(methylthio)phenyl]pyrrolo-[2,1-*a*]isoquinoline.

\*The significance threshold was  $P < .05$  at the corrected voxel level and  $P < .05$  at the corrected cluster level. Coordinates are given in millimeters from the origin at the midpoint of the anterior commissure for voxels of peak significance.

based interviews. Furthermore, in this study, the severity of aggression clearly paralleled the decreases in serotonin transporter density in the brain, which in turn were found to be associated with the duration of methamphetamine use. Therefore, it seems unlikely that the increased aggression observed in these methamphetamine abusers reflected a preexisting disposition or personality trait.

Except for the scores on the AQ, none of the scores on the clinical rating scales for psychiatric symptoms were correlated with the decrease in serotonin transporter density. Methamphetamine has been reported to affect not only serotonergic neurons but also several other types of neurons, such as the dopaminergic, glutamatergic, and  $\gamma$ -aminobutyric acid (GABA)-ergic neurons, all of which have been implicated in the presence of a variety of psychiatric symptoms (eg, delusions, hallucinations, and anxiety).<sup>63</sup> It is possible that changes in various types of neurons might have affected or modified the clinical symptoms evaluated herein. Another plausible interpretation for the negative results is that, as seen in Table 1, the severity of most of the residual symptoms assessed in this study ranged from mild to moderate, and the variances of their distributions were relatively small; together, these factors may have biased the results toward the null hypothesis.

Herein, we recruited methamphetamine abusers from the community; they were recreational abusers of methamphetamine only, and none of them had used other illicit drugs or had taken toxic or high doses of methamphetamine. Although our strategy allowed us to evaluate

the pure effects of methamphetamine on the human brain, the findings may not be generalized to the broad population of methamphetamine abusers. However, the combined use of methamphetamine with other illicit drugs is infrequent in Japan, as indicated by Japanese National Police Agency records in 2002.<sup>64</sup> One reason for this is that cannabis, cocaine, and major illicit drugs other than methamphetamine are not widely distributed in Japan.<sup>64</sup> Furthermore, a national survey of 233 methamphetamine abusers reported that only 2.6% of the abusers had undergone methamphetamine intoxication,<sup>65</sup> suggesting that abusers of an overdose of methamphetamine are rare in Japan. Consequently, our findings are considered to be fairly generalizable to the population of methamphetamine abusers, at least in Japan.

In this study, all the methamphetamine abusers exhibited some psychopathologic symptoms, even in an abstinent state. To our knowledge, no previous studies have examined the incidence of psychopathologic abnormalities in abstinent methamphetamine abusers recruited from the general community. In a study by Wada and Fukui,<sup>65</sup> who investigated the psychopathologic characteristics of 233 abstinent methamphetamine abusers recruited from hospitals in Japan (the period of abstinence exceeded 1 month; the mean  $\pm$  SD duration of methamphetamine use was  $11.1 \pm 7.9$  years), almost all the abusers exhibited some psychopathologic symptoms, such as auditory hallucinations, delusions of reference/persecution, mood disturbances, anxiety, insomnia, irritability, impulsivity, and personality changes, including the antisocial personality type. Such observations cannot be applied to absti-

ment abusers in the community as a whole but may provide some support for the high occurrence of psychopathologic symptoms observed in this study. In Japan, most methamphetamine abusers take the substance intravenously,<sup>8</sup> whereas in a study<sup>66</sup> from the United States, approximately 90% of methamphetamine abusers had no history of intravenous or intramuscular injection of methamphetamine. Furthermore, a study by Domier and colleagues<sup>66</sup> revealed that among recently abstinent methamphetamine abusers who had discontinued its use for several months, the injecting abusers had a significantly higher incidence of psychopathologic symptoms than the noninjecting abusers. These results suggest that in Japan, the intravenous intake of methamphetamine could predispose its abusers to persistent psychiatric problems, even after the cessation of methamphetamine use. Nevertheless, it remains an important and unresolved issue whether a reduction in serotonin transporter could be expected to occur in abusers with no psychopathologic signs or symptoms. To verify our findings that methamphetamine abuse is linked to a reduction in brain serotonin transporters, which in turn underlies persistent psychopathologic symptoms, additional studies that also incorporate a group of methamphetamine abusers with no apparent psychopathologic problems are required.

Wilson and colleagues<sup>67</sup> examined serotonin concentrations in postmortem tissue samples from human brains with a history of long-term methamphetamine abuse, although they did not study serotonin transporters per se. They concluded that there were no substantial alterations in serotonin concentrations in the global brain except in the medial prefrontal cortex (Brodmann area 11: a reduction of 56% compared with controls) and in the orbitofrontal cortex (Brodmann area 12: a reduction of 61% compared with controls). These results seem to contradict our observation of reductions in serotonin transporters in widely distributed brain regions. The discrepancy between the results of that postmortem study and those of present study is puzzling. However, one possible explanation for this discrepancy could be related to differences in the pattern and amount of drug use between the samples.<sup>1,16,17</sup> In Western countries, methamphetamine abusers often use other drugs, mainly cocaine or cannabis<sup>68-70</sup>; however, no information is provided with respect to this issue in the study by Wilson and colleagues.<sup>67</sup> Because methamphetamine is more likely to produce neurotoxic effects in serotonergic neurons than either cocaine or cannabis,<sup>71,72</sup> methamphetamine abusers who use this drug only could have experienced more severe damage to serotonergic neurons than abusers who simultaneously use other drugs, such as cocaine or cannabis. Furthermore, similar to most methamphetamine abusers in Japan, those in this study intravenously injected the substance. The intravenous intake may further potentiate the neurotoxic effects of methamphetamine.

To our knowledge, this is the first study to demonstrate a severe and long-lasting reduction in the density of the serotonin transporter in the living brains of methamphetamine abusers. The observed decrease in serotonin transporter density was also found to be associated with elevated levels of aggression. The present findings,

combined with the results of previous animal studies, suggest that those who abuse methamphetamine may be at substantial risk for severe serotonin neuronal damage in the brain, potentially leading to persistently elevated aggression, even in those in a currently abstinent state.

Submitted for Publication: September 21, 2004; final revision received April 30, 2005; accepted July 7, 2005.

Correspondence: Yoshimoto Sekine, MD, PhD, Department of Psychiatry and Neurology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Hamamatsu, Shizuoka 431-3192 Japan (ysekine@hama-med.ac.jp).

Acknowledgment: This work was supported by a Grant-in-Aid for the Center of Excellence from the Ministry of Education, Culture, Sports, Science, and Technology, Tokyo, Japan; research grant 16A for nervous and mental disorders from the Ministry of Health, Labor, and Welfare, Tokyo; a grant for the research on serotonin from the Public Health Research Foundation/New Frontiers of Neurotransmitter Research, Tokyo; and the Stanley Foundation, Bethesda, Md.

## REFERENCES

1. McCann UD, Wong DF, Yokoi F, Villemagne V, Dannals RF, Ricaurte GA. Reduced striatal dopamine transporter density in abstinent methamphetamine and methcathinone users: evidence from positron emission tomography studies with [<sup>11</sup>C]WIN-35,428. *J Neurosci*. 1998;18:8417-8422.
2. Volkow ND, Chang L, Wang GJ, Fowler JS, Leonido-Yee M, Franceschi D, Sedler MJ, Gatley SJ, Hitzemann R, Ding YS, Logan J, Wong C, Miller EN. Association of dopamine transporter reduction with psychomotor impairment in methamphetamine abusers. *Am J Psychiatry*. 2001;158:377-382.
3. Sekine Y, Iyo M, Ouchi Y, Matsunaga T, Tsukada H, Okada H, Yoshikawa E, Futatsubashi M, Takei N, Mori N. Methamphetamine-related psychiatric symptoms and reduced brain dopamine transporters studied with PET. *Am J Psychiatry*. 2001;158:1206-1214.
4. Farrell M, Marsden J, Ali R, Ling W. Methamphetamine: drug use and psychoses becomes a major public health issue in the Asia Pacific region. *Addiction*. 2002;97:771-772.
5. Hall W, Hando J, Darke S, Ross J. Psychological morbidity and route of administration among amphetamine users in Sydney, Australia. *Addiction*. 1996;91:81-87.
6. London ED, Simon SL, Berman SM, Mandelkern MA, Lichtman AM, Bramen J, Shinn AK, Miotto K, Learn J, Dong Y, Matochik JA, Kurian V, Newton T, Woods R, Rawson R, Ling W. Mood disturbances and regional cerebral metabolic abnormalities in recently abstinent methamphetamine abusers. *Arch Gen Psychiatry*. 2004;61:73-84.
7. Seivewright N. Disorders relating to the use of amphetamine and cocaine. In: Gelder MG, Lopez-Ibor JJ, Andreasen N, eds. *New Oxford Textbook of Psychiatry*. New York, NY: Oxford University Press; 2000:531-534.
8. Konuma K. Use and abuse of amphetamines in Japan. In: Cho AK, Segal DS, eds. *Amphetamine and Its Analogs*. San Diego, Calif: Academic Press; 1994:415-435.
9. Sekine Y, Minabe Y, Kawai M, Suzuki K, Iyo M, Isoda H, Sakahara H, Ashby CR Jr, Takei N, Mori N. Metabolite alterations in basal ganglia associated with methamphetamine-related psychiatric symptoms: a proton MRS study. *Neuropsychopharmacology*. 2002;27:453-461.
10. Sato M, Chen CC, Akiyama K, Otsuki S. Acute exacerbation of paranoid psychotic state after long-term abstinence in patients with previous methamphetamine psychosis. *Biol Psychiatry*. 1983;18:429-440.
11. Ricaurte GA, Schuster CR, Seiden LS. Long-term effects of repeated methylamphetamine administration on dopamine and serotonin neurons in the rat brain: a regional study. *Brain Res*. 1980;193:153-163.
12. Davidson C, Gow AJ, Lee TH, Ellinwood EH. Methamphetamine neurotoxicity: necrotic and apoptotic mechanisms and relevance to human abuse and treatment. *Brain Res Brain Res Rev*. 2001;36:1-22.
13. Matsuzaki H, Namikawa K, Kiyama H, Mori N, Sato K. Brain-derived neurotrophic factor rescues neuronal death induced by methamphetamine. *Biol Psychiatry*. 2004;55:52-60.

14. Ricaurte GA, Sabol KE, Seiden LS. Functional consequences of neurotoxic amphetamine exposure. In: Cho AK, Segal DS, eds. *Amphetamine and Its Analogs*. San Diego, Calif: Academic Press; 1994:297-313.
15. Kita T, Wagner GC, Nakashima T. Current research on methamphetamine-induced neurotoxicity: animal models of monoamine disruption. *J Pharmacol Sci*. 2003;92:178-195.
16. Villemagne V, Yuan J, Wong DF, Dannals RF, Hatzidimitriou G, Mathews WB, Ravert HT, Musachio J, McCann UD, Ricaurte GA. Brain dopamine neurotoxicity in baboons treated with doses of methamphetamine comparable to those recreationally abused by humans: evidence from [<sup>11</sup>C]WIN-35,428 positron emission tomography studies and direct in vitro determinations. *J Neurosci*. 1998;18:419-427.
17. Volkow ND, Chang L, Wang GJ, Fowler JS, Franceschi D, Sedler M, Gatley SJ, Miller E, Hitzemann R, Ding YS, Logan J. Loss of dopamine transporters in methamphetamine abusers recovers with protracted abstinence. *J Neurosci*. 2001;21:9414-9418.
18. Sekine Y, Minabe Y, Ouchi Y, Takei N, Iyo M, Nakamura K, Suzuki K, Tsukada H, Okada H, Yoshikawa E, Futatsubashi M, Mori N. Association of dopamine transporter loss in the orbitofrontal and dorsolateral prefrontal cortices with methamphetamine-related psychiatric symptoms. *Am J Psychiatry*. 2003;160:1699-1701.
19. McCann UD, Szabo Z, Scheffel U, Dannals RF, Ricaurte GA. Positron emission tomographic evidence of toxic effect of MDMA ("Ecstasy") on brain serotonin neurons in human beings. *Lancet*. 1998;352:1433-1437.
20. Reneman L, Lavalaye J, Schmand B, de Wolff FA, van den Brink W, den Heeten GJ, Booij J. Cortical serotonin transporter density and verbal memory in individuals who stopped using 3,4-methylenedioxymethamphetamine (MDMA or "ecstasy"): preliminary findings. *Arch Gen Psychiatry*. 2001;58:901-906.
21. Szabo Z, McCann UD, Wilson AA, Scheffel U, Owonikoko T, Mathews WB, Ravert HT, Hilton J, Dannals RF, Ricaurte GA. Comparison of (+)-(11)C-McN5652 and (11)C-DASB as serotonin transporter radioligands under various experimental conditions. *J Nucl Med*. 2002;43:678-692.
22. Semple DM, Ebmeier KP, Glabus MF, O'Carroll RE, Johnstone EC. Reduced in vivo binding to the serotonin transporter in the cerebral cortex of MDMA ("ecstasy") users. *Br J Psychiatry*. 1999;175:63-69.
23. Scheffel U, Szabo Z, Mathews WB, Finley PA, Yuan J, Callahan B, Hatzidimitriou G, Dannals RF, Ravert HT, Ricaurte GA. Fenfluramine-induced loss of serotonin transporters in baboon brain visualized with PET. *Synapse*. 1996;24:395-398.
24. Bengel D, Isaacs KR, Heils A, Lesch KP, Murphy DL. The appetite suppressant d-fenfluramine induces apoptosis in human serotonergic cells. *Neuroreport*. 1998;9:2989-2993.
25. McCann UD, Yuan J, Ricaurte GA. Neurotoxic effects of +/-fenfluramine and phentermine, alone and in combination, on monoamine neurons in the mouse brain. *Synapse*. 1998;30:239-246.
26. Poole R, Brabbins C. Drug induced psychosis. *Br J Psychiatry*. 1996;168:135-138.
27. Takebayashi K, Sekine Y, Takei N, Minabe Y, Isoda H, Takeda H, Nishimura K, Nakamura K, Suzuki K, Iwata Y, Sakahara H, Mori N. Metabolite alterations in basal ganglia associated with psychiatric symptoms of abstinent toluene users: a proton MRS study. *Neuropsychopharmacology*. 2004;29:1019-1026.
28. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*. Washington, DC: American Psychiatric Press; 1994.
29. First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Patient Version*. Washington, DC: American Psychiatric Press; 1997.
30. Al-Dirbashi OY, Kuroda N, Wada M, Takahashi M, Nakashima K. Quantification of methamphetamine, amphetamine and enantiomers by semi-micro column HPLC with fluorescence detection; applications on abusers' single hair analyses. *Biomed Chromatogr*. 2000;14:293-300.
31. Buss AH, Perry M. The aggression questionnaire. *J Pers Soc Psychol*. 1992;63:452-459.
32. Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol*. 1959;32:50-55.
33. Hamilton M. Development of a rating scale for primary depressive illness. *Br J Soc Clin Psychol*. 1967;6:278-296.
34. Mohr P, Horacek J, Motlova L, Libiger J, Czobor P. Prolactin response to d-fenfluramine challenge test as a predictor of treatment response to haloperidol in acute schizophrenia. *Schizophr Res*. 1998;30:91-99.
35. Overall JE, Gorham DR. The Brief Psychiatric Rating Scale. *Psychol Rep*. 1962;10:799-812.
36. Volkow ND, Wang GJ, Fischman MW, Foltin RW, Fowler JS, Abumrad NN, Vitkun S, Logan J, Gatley SJ, Pappas N, Hitzemann R, Shea CE. Relationship between subjective effects of cocaine and dopamine transporter occupancy. *Nature*. 1997;386:827-830.
37. Ouchi Y, Nobeza S, Okada H, Yoshikawa E, Futatsubashi M, Kaneko M. Altered glucose metabolism in the hippocampal head in memory impairment. *Neurology*. 1998;51:136-142.
38. Watanabe M, Shimizu K, Omura T, Takahashi M, Kosugi T, Yoshikawa E, Sato N, Okada H, Yamashita T. A new high resolution PET scanner dedicated to brain research. *IEEE Trans Nucl Sci*. 2002;49:634-639.
39. Parsey RV, Kegeles LS, Hwang DR, Simpson N, Abi-Dargham A, Mawlawi O, Slifstein M, Van Heertum RL, Mann JJ, Laruelle M. In vivo quantification of brain serotonin transporters in humans using [<sup>11</sup>C]McN 5652. *J Nucl Med*. 2000;41:1465-1477.
40. Ouchi Y, Yoshikawa E, Okada H, Futatsubashi M, Sekine Y, Iyo M, Sakamoto M. Alterations in binding site density of dopamine transporter in the striatum, orbitofrontal cortex, and amygdala in early Parkinson's disease: compartment analysis for β-CFT binding with positron emission tomography. *Ann Neurol*. 1999;45:601-610.
41. Brody AL, Mandelkern MA, London ED, Childress AR, Lee GS, Bota RG, Ho ML, Saxena S, Baxter LR Jr, Madsen D, Jarvik ME. Brain metabolic changes during cigarette craving. *Arch Gen Psychiatry*. 2002;59:1162-1172.
42. Ouchi Y, Nobeza S, Yoshikawa E, Futatsubashi M, Kanno T, Okada H, Torizuka T, Nakayama T, Tanaka K. Postural effects on brain hemodynamics in unilateral cerebral artery occlusive disease: a positron emission tomography study. *J Cereb Blood Flow Metab*. 2001;21:1058-1066.
43. Ouchi Y, Okada H, Yoshikawa E, Futatsubashi M, Nobeza S. Absolute changes in regional cerebral blood flow in association with upright posture in humans: an orthostatic PET study. *J Nucl Med*. 2001;42:707-712.
44. Simpson HB, Lombardo I, Slifstein M, Huang HY, Hwang DR, Abi-Dargham A, Liebowitz MR, Laruelle M. Serotonin transporters in obsessive-compulsive disorder: a positron emission tomography study with [<sup>11</sup>C]McN 5652. *Biol Psychiatry*. 2003;54:1414-1421.
45. Mikolajczyk K, Szabatin M, Rudnicki P, Grodzki M, Burger CA. JAVA environment for medical image data analysis: initial application for brain PET quantitation. *Med Inform (Lond)*. 1998;23:207-214.
46. Tauscher J, Kapur S, Verhoeff NP, Hussey DF, Daskalakis ZJ, Tauscher-Wisniewski S, Wilson AA, Houle S, Kasper S, Zipsursky RB. Brain serotonin 5-HT (1A) receptor binding in schizophrenia measured by positron emission tomography and [<sup>11</sup>C]WAY-100635. *Arch Gen Psychiatry*. 2002;59:514-520.
47. Talairach J, Tournoux P. *Co-planer Stereotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*. Stuttgart, Germany: Georg Thieme; 1988.
48. Ito K, Morrish PK, Rakshi JS, Uema T, Ashburner J, Bailey DL, Friston KJ, Brooks DJ. Statistical parametric mapping with <sup>18</sup>F-dopa PET shows bilaterally reduced striatal and nigral dopaminergic function in early Parkinson's disease. *J Neurosurg Psychiatry*. 1999;66:754-758.
49. Ouchi Y, Kanno T, Okada H, Yoshikawa E, Futatsubashi M, Nobeza S, Torizuka T, Tanaka K. Changes in dopamine availability in the nigrostriatal and mesocortical dopaminergic systems by gait in Parkinson's disease. *Brain*. 2001;124:784-792.
50. Wang GJ, Volkow ND, Chang L, Miller E, Sedler M, Hitzemann R, Zhu W, Logan J, Ma Y, Fowler JS. Partial recovery of brain metabolism in methamphetamine abusers after protracted abstinence. *Am J Psychiatry*. 2004;161:242-248.
51. Kovachich GB, Aronson CE, Brunswick DJ. Effects of high-dose methamphetamine administration on serotonin uptake sites in rat brain measured using [<sup>3</sup>H]cynonimpipramine autoradiography. *Brain Res*. 1989;505:123-129.
52. Davidson RJ, Putnam KM, Larson CL. Dysfunction in the neural circuitry of emotion regulation: a possible prelude to violence. *Science*. 2000;289:591-594.
53. Coccaro EF, Kavoussi RJ, Hauger RL, Cooper TB, Ferris CF. Cerebrospinal fluid vasopressin levels: correlates with aggression and serotonin function in personality-disordered subjects. *Arch Gen Psychiatry*. 1998;55:708-714.
54. Coccaro EF. Central serotonin and impulsive aggression. *Br J Psychiatry Suppl*. 1989;8:52-62.
55. Virkkunen M, Rawlings R, Tokola R, Poland RE, Guidotti A, Nemeroff C, Bissette G, Kalogeris K, Karonen SL, Linnoila M. CSF biochemistry, glucose metabolism, and diurnal activity rhythms in alcoholic, violent offenders, fire setters, and healthy volunteers. *Arch Gen Psychiatry*. 1994;51:20-27.
56. Linnoila M, DeJong J, Virkkunen M. Monoamines, glucose metabolism, and impulse control. *Psychopharmacol Bull*. 1989;25:404-406.
57. Roy A, Adinolf B, Linnoila M. Acting out hostility in normal volunteers: negative correlation with levels of 5HIAA in cerebrospinal fluid. *Psychiatry Res*. 1988;24:187-194.
58. Linnoila M, Virkkunen M, Scheinin M, Nuutila A, Rimon R, Goodwin FK. Low cerebrospinal fluid 5-hydroxyindoleacetic acid concentration differentiates impulsive from nonimpulsive violent behavior. *Life Sci*. 1983;33:2609-2614.
59. Grafman J, Schwab K, Warden D, Pridgen A, Brown HR, Salazar AM. Frontal lobe injuries, violence, and aggression: a report of the Vietnam Head Injury Study. *Neurology*. 1996;46:1231-1238.
60. Parsey RV, Oquendo MA, Simpson NR, Ogden RT, Van Heertum R, Arango V, Mann JJ. Effects of sex, age, and aggressive traits in man on brain serotonin

- 5-HT<sub>1A</sub> receptor binding potential measured by PET using [C-11]WAY-100635. *Brain Res.* 2002;954:173-182.
61. Lai MK, Tsang SW, Francis PT, Esiri MM, Keene J, Hope T, Chen CP. Reduced serotonin 5-HT<sub>1A</sub> receptor binding in the temporal cortex correlates with aggressive behavior in Alzheimer disease. *Brain Res.* 2003;974:82-87.
  62. van den Bree MB, Svikis DS, Pickens RW. Genetic influences in antisocial personality and drug use disorders. *Drug Alcohol Depend.* 1998;49:177-187.
  63. Wrona MZ, Yang Z, Zhang F, Dryhurst G. Potential new insights into the molecular mechanisms of methamphetamine-induced neurodegeneration. *NIDA Res Monogr.* 1997;173:146-174.
  64. National Police Agency (Japan). *Criminal White Paper* [in Japanese]. Tokyo, Japan: Printing Bureau, Ministry of Finance of Japan; 2004.
  65. Wada K, Fukui S. Relationship between years of methamphetamine use and symptoms of methamphetamine psychosis [in Japanese]. *Arukoro Kenkyuto Yakubutsu Ison.* 1990;25:143-158.
  66. Domier CP, Simon SL, Rawson RA, Huber A, Ling W. A comparison of injecting and noninjecting methamphetamine users. *J Psychoactive Drugs.* 2000;32:229-232.
  67. Wilson JM, Kalasinsky KS, Levey AI, Bergeron C, Reiber G, Anthony RM, Schmunk GA, Shannak K, Haycock JW, Kish SJ. Striatal dopamine nerve terminal markers in human, chronic methamphetamine users. *Nat Med.* 1996;2:699-703.
  68. Richter KP, Ahluwalia HK, Mosier MC, Nazir N, Ahluwalia JS. A population-based study of cigarette smoking among illicit drug users in the United States. *Addiction.* 2002;97:861-869.
  69. Sumnall HR, Wagstaff GF, Cole JC. Self-reported psychopathology in polydrug users. *J Psychopharmacol.* 2004;18:75-82.
  70. Smart RG, Mann RE, Tyson LA. Drugs and violence among Ontario students. *J Psychoactive Drugs.* 1997;29:369-373.
  71. Jacobsen LK, Staley JK, Malison RT, Zoghbi SS, Seibyl JP, Kosten TR, Innis RB. Elevated central serotonin transporter binding availability in acutely abstinent cocaine-dependent patients. *Am J Psychiatry.* 2000;157:1134-1140.
  72. Croft RJ, Klugman A, Baldeweg T, Gruzeller JH. Electrophysiological evidence of serotonergic impairment in long-term MDMA ("ecstasy") users. *Am J Psychiatry.* 2001;158:1687-1692.

#### Clinical Trials Registration

In concert with the International Committee of Medical Journal Editors, *Archives of General Psychiatry* will require, as a condition of consideration for publication, registration of clinical trials in a public trials registry (such as <http://ClinicalTrials.gov> or <http://controlled-trials.com>). Trials must be registered at or before the onset of patient enrollment. This policy applies to any clinical trial starting enrollment after March 1, 2006. For trials that began enrollment before this date, registration will be required by June 1, 2006. The trial registration number should be supplied at the time of submission.

For details about this new policy see the editorials by DeAngelis et al in the September 8, 2004 (2004;292:1363-1364) and June 15, 2005 (2005;293:2927-2929) issues of *JAMA*.



## Advanced paternal age associated with an elevated risk for schizophrenia in offspring in a Japanese population

Kenji J. Tsuchiya<sup>a</sup>, Shu Takagai<sup>a</sup>, Masayoshi Kawai<sup>a</sup>, Hideo Matsumoto<sup>b</sup>,  
Kazuhiko Nakamura<sup>a</sup>, Yoshio Minabe<sup>a</sup>, Norio Mori<sup>a</sup>, Nori Takei<sup>a,c,\*</sup>

<sup>a</sup>Department of Psychiatry and Neurology, Hamamatsu University School of Medicine, Hamamatsu, Japan

<sup>b</sup>Department of Psychiatry, Tokai University School of Medicine, Isehara, Japan

<sup>c</sup>Division of Psychological Medicine, Institute of Psychiatry, London, UK

Received 14 December 2004; received in revised form 2 March 2005; accepted 4 March 2005

Available online 21 April 2005

### Abstract

**Objective:** Advanced paternal age at birth as a risk for schizophrenia in the adult offspring has been reported in previous studies exclusively conducted in Western countries and Israel. The question has arisen whether this finding could be replicated in countries with socially and culturally different attitudes toward marriage, including factors such as age at marriage. To address this question, we conducted a case-control study of a Japanese population.

**Methods:** The subjects were representative inpatients with a DSM-IV diagnosis of schizophrenia. Unrelated healthy volunteers were recruited as control subjects. This study was conducted as one of a series of the projects by use of “The Mother and Child Health Handbooks (MCHHs),” from which information on parental characteristics around the time of birth, including parental ages at birth, had been extracted and recorded on computer.

**Results:** Ninety-nine subjects with schizophrenia and 381 healthy control subjects enrolled for the study. Advanced paternal, but not maternal, age was associated with an elevated risk for schizophrenia. Reproducibility of the association across different cultures is suggestive of a causal link.

© 2005 Elsevier B.V. All rights reserved.

**Keywords:** Schizophrenia; Paternal age; Risk factor; Case-control study; Japan

### 1. Introduction

Recent studies have consistently reported that advanced paternal age at birth is a risk factor for schizophrenia in the offspring (Brown et al., 2002; Byrne et al., 2003; Dalman and Allebeck, 2002; El Saadi et al., 2004; Malaspina et al., 2001; Zammit et al., 2003). Paternal germ line mutation may be a likely

\* Corresponding author. Department of Psychiatry and Neurology, Hamamatsu University School of Medicine, 1-20-1 Handayama, Hamamatsu 431-3192, Japan. Tel.: +81 53 435 2295; fax: +81 53 435 3621.

E-mail address: [ntakei@hama-med.ac.jp](mailto:ntakei@hama-med.ac.jp) (N. Takei).

explanation (Malaspina, 2001), but the mechanism underlying this association is not yet fully understood.

Previous studies examining this association have been conducted exclusively in European countries, Israel, USA, and Australia. Attitudes toward marriage, in particular, age at the time of marriage, may vary according to cultural backgrounds. If the association between advanced paternal age and the risk for schizophrenia were reproduced among populations of different ethnic origin, the relationship addressed here would be more indicative of causal significance, and genetic explanations including de novo mutation on schizophrenia-susceptible genes would become realistic (Malaspina, 2001).

Therefore, we investigated whether or not the claimed association between advanced paternal age at birth and the risk for schizophrenia among offspring would be reproducible in a Japanese population.

## 2. Methods

This study was conducted as a series of our projects which were launched to investigate pregnancy- and birth-related events as risk factors for schizophrenia in Japan. The projects were set out to exclusively utilize the Mother and Child Health Handbook (MCHH), a set of notes covering a wide range of information, including pregnancy and delivery events, as well as newborns' characteristics and parental data. In Japan, every woman receives the MCHH from the local municipal office when pregnant

and, in general, keeps it long after the birth of the offspring as a memorial. Ample information derived from the MCHHs has been systematically recorded on computer and, thus far, used for a series of studies (Kawai et al., 2004; Matsumoto et al., 1999, 2001).

We sought the participation of subjects with schizophrenia who were admitted to the University Hospital of the Hamamatsu University School of Medicine during the period extending from January 1, 1999 to December 31, 2002. A structured interview was conducted by trained psychiatrists (K.J.T., K.M., N.M. and N.T.) using the Schedule for Clinical Interview for Axis-I Diagnosis of DSM-IV (SCID-I) (First et al., 1996). Those individuals, who met the criteria for a DSM-IV diagnosis of schizophrenia (295.xx) and whose biological mothers were identified, were recruited. We also included unrelated healthy volunteers who were residents in the community from which the patient group was derived, and who were confirmed to have had no psychiatric history.

As regards the parental age and parity, we relied upon computerized MCHH data. As these data have become available since the year 1967, only individuals born in and after 1968 were included in the present study. Maternal and paternal ages were divided into tertiles according to the distribution of age (see Table 1). The family history of psychiatric diagnoses was examined via interviews with each subject and one close relative, most often the mother; subjects were designated as "positive" for a psychiatric family history if they had at least one first-degree relative with psychosis. Odds ratios (ORs) and

Table 1  
Odds ratios for schizophrenia among offspring in relation to parental age

	Cases ( <i>n</i> = 99)	Controls ( <i>n</i> = 381)	Crude estimates		Adjusted estimates <sup>a</sup>	
			ORs	95% CI	ORs	95% CI
<i>Maternal age (year)</i>						
≤25 (reference)	31	128	1		1	
26–28	40	118	1.40	(0.82, 2.38)	1.08	(0.60, 1.93)
≥29	28	135	0.86	(0.49, 1.51)	0.59	(0.29, 1.20)
Test for trend			<i>p</i> = 0.60		<i>p</i> = 0.14	
<i>Paternal age (year)</i>						
≤28 (reference)	22	138	1		1	
29–31	36	120	1.88	(1.05, 3.37)	2.08	(1.12, 3.86)
≥32	41	123	2.09	(1.18, 3.71)	3.00	(1.49, 6.04)
Test for trend			<i>p</i> = 0.013		<i>p</i> = 0.002	

<sup>a</sup> Adjusted for age and gender of the subject, parity, family history, and age of the other parent.

95% confidence intervals (CIs) for schizophrenia in relation to parental age categories were estimated using logistic regression. Along with the crude OR, we computed the OR adjusted for relevant covariates in order to eliminate the confounding effects. In light of the moderate sample size of the present study, we opted for a parsimonious model to increase precision rather than a model in which all available variables were adjusted for. We adopted a logistic regression model with a forward procedure (Kleinbaum et al., 1998). Following the recommendation by Kleinbaum et al. (1998), we chose a loosened alpha level of .20 to include potential confounders in the model. For the statistical analyses, we used STATA, version 8.1 (Stata Corporation, 2003).

Written informed consent was obtained from each participant. This study design was approved by the University Hospital Ethics Committee.

### 3. Results

#### 3.1. Characteristics in patients and controls

Ninety-nine patients with schizophrenia and 381 control subjects participated in this study. The mean age differed significantly between the patients and controls (25.4 years [S.D. 5.5] for the patients and 23.0 years [S.D. 4.3] for the controls;  $t=4.53$ ,  $df=478$ ,  $p<.001$ ). More male subjects were included among the patients ( $n=56$ , 57%) than among the controls ( $n=179$ , 47%);  $\chi^2=2.89$ ,  $df=1$ ,  $p=.089$ . A higher proportion of a family history of psychosis was noted among the patients (8/99, 8%) than among the controls (14/381, 4%);  $\chi^2=3.49$ ,  $df=1$ ,  $p=.062$ . There was no marked or significant difference in the distribution of first-born subjects between the patients (40/99, 40%) and controls (157/381, 41%);  $\chi^2=0.02$ ,  $df=1$ ,  $p=.89$ .

#### 3.2. Parental age and risk for schizophrenia in the offspring

Maternal age at birth did not differ significantly between the two groups (27.3 years [S.D. 3.5] for the patients and 27.3 years [S.D. 3.5] for the controls;  $t=0.12$ ,  $df=478$ ,  $p=.91$ ); however, there was a trend toward greater paternal age at birth among the patients with schizophrenia than among the controls (30.9

years [S.D. 3.7] for the patients and 30.1 years [S.D. 4.0] for the controls;  $t=1.87$ ,  $df=478$ ,  $p=.062$ ).

Differences in risk were then examined across three classes of maternal age at birth using logistic regression with no adjustment for covariates; the test for a linear trend revealed no association (likelihood ratio test (LRT)=0.28,  $df=1$ ,  $p=.60$ ) (Table 1). However, as regards paternal age, the test for a linear trend did reveal a significant association (LRT=6.28,  $df=1$ ,  $p=.013$ ). Those subjects whose paternal age was 29 to 31 years at birth had a 1.9-fold increased risk, and those with a paternal age of 32 years or older had a 2.1-fold increased risk, compared to those with a paternal age of 28 years or younger.

Subsequently, we incorporated potential confounding factors into the logistic regression model with a forward procedure, as described above. Accordingly, age, sex, family history, and age of the other parent were entered into the model as covariates. The linear trend that had been observed in the crude analysis of paternal age remained significant, but became even more significant after adjustment for these covariates (LRT=9.37,  $df=1$ ,  $p=.002$ ). In effect, the estimated risk among the offspring of the highest paternal age class was three times greater than that of the offspring of the lowest paternal age class. However, as for maternal age, no association emerged, even after the covariates were entered into the model (LRT=2.21,  $df=1$ ,  $p=.14$ ).

A third factor in the context of the relationship between two factors (paternal age and a risk for the disorder) may account for the implication of advanced paternal age as a risk factor, i.e., deviation in paternal behavior. Namely, men with a predisposition to the disorder may tend to get married and to father children at a relatively high age. As information on the behavioral traits of the fathers of the subjects included in the present study was not available, we chose to examine the interaction of “family history” and “paternal age” in the same regression model. In this analysis, we found no significant interaction effect (LRT=1.40,  $df=1$ ,  $p=.24$ ).

### 4. Discussion

In the Japanese population, we found an association between advanced paternal age at birth and the



risk for schizophrenia in the offspring. This finding is consistent with that of previous studies (Brown et al., 2002; Byrne et al., 2003; Dalman and Allebeck, 2002; El Saadi et al., 2004; Malaspina et al., 2001; Zammit et al., 2003). However, these prior studies did not find support for an association between maternal age and the risk of offspring developing the disorder. The results of the present study also showed no such association, and are hence once again in agreement with previous reports.

It is highly likely that marital habits, including preferred age at marriage among men, as well as fathering age, vary across countries and cultures. In fact, the mean paternal age at birth in a Swedish study was 32.2 years for control offspring (Zammit et al., 2003), which was slightly older than that (31.5 years) in an Israeli study (Malaspina et al., 2001) (information on mean paternal age is not available in the other studies cited above). In contrast, our Japanese study showed a comparatively younger paternal age at birth (30.1 years) in the control subjects. These data indicate some degree of cultural influence on fathering age. Nevertheless, all of the previous studies cited here, as well as our study, showed the same finding of an association between paternal age at birth and the risk for schizophrenia, strongly suggesting the presence of non-culturally determined factor(s).

Apart from cultural factors, other technical issues, in particular the diagnosis, need to be considered. Previous studies have employed several different definitions of schizophrenia, including broadly defined psychoses as well as schizophrenia within a “narrow” framework; each of the definitions employed may exert a different impact on the association of interest. As regards studies utilizing a broad concept of psychosis that includes schizophrenia, four studies incorporated subsamples with psychoses, namely affective psychosis as well as non-affective psychosis (three parts of the study by El Saadi et al., 2004; Brown et al., 2002). Among these studies, two studies showed a significant association between advanced paternal age and risk (the Swedish and Danish part of the study by El Saadi et al., 2004), and one study showed a trend toward the association (Brown et al., 2002), although the remaining study did not find support for the association (the Australian part of the study by El Saadi et al., 2004). Therefore, the reported findings remain inconsistent. On the other hand, four studies investigated subjects

with “narrow” schizophrenia (e.g., 295 in ICD-9 or F20 in ICD-10), and all showed a significant association (Brown et al., 2002; Dalman and Allebeck, 2002; Zammit et al., 2003; Byrne et al., 2003).

In this context, of particular interest is a study by Brown et al. (2002), who directly compared the strength of the association according to the diagnoses opted for (i.e., psychoses vs. schizophrenia). They demonstrated that the relative risk for schizophrenia among individuals with oldest fathers at birth (45 years or over) was 3.6 (95% CI: 1.0 to 15.5), whereas the relative risk for psychoses among those with fathers belonging to the same age class was 2.7 (95% CI: 0.8 to 7.2), compared with those with the youngest fathers of 15–24 years. These findings indicate that the use of narrowly defined schizophrenia strengthens the association. Strictly speaking, as we adopted a DSM-IV diagnosis of schizophrenia, our findings can only be compared with those studies which adopted a similar (i.e., “narrow”) definition (Brown et al., 2002; Byrne et al., 2003; Dalman and Allebeck, 2002; Zammit et al., 2003). All such studies, including ours, provide supportive evidence for the conclusion that advanced paternal age is associated with an increased risk for schizophrenia among the offspring. In addition, two studies have shown that the paternal age effect is stronger for schizophrenia than for psychoses other than schizophrenia (Malaspina et al., 2001; Zammit et al., 2003). Therefore, it is likely that advanced paternal age is more specifically related to schizophrenia than to other types of psychosis.

Another important factor that may have influenced our findings is birth order. Naturally, later-born children are likely to have an older father compared to first-born children. A study has reported that later birth order is associated with an increased risk for schizophrenia (Sham et al., 1993). Thus, some of the present findings might have been partly due to the effect of birth order. Among all the subjects in the present study, the mean paternal age at the birth of the first-born children was significantly younger, i.e., 28.5 years (S.D. 3.1), compared to that of the fathers of the later-born children, i.e., 31.5 years (S.D. 4.0) ( $t=8.80$ ,  $df=478$ ,  $p<.001$ ). However, parity (or birth order) was not associated with an increased risk for the disorder (first-born vs. later-born,  $LRT=0.02$ ,  $df=1$ ,  $p=0.89$ ;  $OR=0.97$  [95% CI: 0.62 to 1.52]). Furthermore, in order to rule out any residual

confounding effect of birth order, we adjusted for this factor in our logistic regression model.

On the other hand, the behavioral characteristics of the fathers might account for the association we found between advanced paternal age and a risk for schizophrenia in the offspring. Many studies have indicated that “schizotypy”, or schizotypal personality traits reflect a genetic background of schizophrenia and run in the pedigree of patients with the disorder (Kendler et al., 1995; Mata et al., 2000). The behavior of fathers with such traits may be characterized by impaired sociality, which may lead to relatively late marriage. Therefore, our finding of advanced paternal age at birth in the pre-schizophrenic offspring could be accounted for by their father’s behavioral characteristics. However, such a scenario of genetic influence should equally pertain to the maternal side, that is, maternal behavioral characteristics. We found no association of maternal age with the risk, which is opposed to the genetic account of the finding in this study. In addition, the ORs representing the strength of the association between paternal age and the risk remained unchanged after adjustment for a family history of psychosis. Moreover, no interaction effect of family history and the paternal age was evident. In accord with these findings, a study by Byrne et al. (2003), who examined two risk sets (all subjects and subjects without a family history of psychiatric diagnosis), has shown that the strength of the association between advanced paternal age and the risk for schizophrenia does not differ between the two sets of the sample. Therefore, it is unlikely that the association between advanced paternal age and the risk for schizophrenia is explained exclusively by family history, or by paternal “schizotypy”.

A different approach has also been applied to address the relation of family history to the paternal age effect. Malaspina et al. (2002) examined the paternal age separately for schizophrenia probands with at least one first- or second-degree relative suffering from non-affective psychosis (i.e., familial schizophrenia) and those without such relatives (i.e., sporadic schizophrenia). This was conducted on the basis that if paternal age were older in sporadic cases than in familial cases, then new germ line mutations would be a possible explanation for the paternal age effect. Indeed, the results provided evidence supporting their prediction (Malaspina et al., 2002). However,

this finding was not supported by a recent study of Pulver et al. (2004), who also investigated parental age at birth in familial vs. sporadic cases, and found no group difference. Although the findings of the two studies conflict with each other, they, at the very least, do not give support to a genetic link between the paternal age at birth and the risk of the disorder.

It should be noted that Dalman and Allebeck (2002) reported that those individuals with fathers belonging to the oldest age class or to the youngest age class at birth may be associated with an increased risk, compared to those with fathers belonging to the middle age class. We failed to replicate such a ‘U’-shaped relationship of paternal age to the risk for the disorder; instead, our findings indicated a linear relationship between advanced paternal age and the risk for schizophrenia.

#### *4.1. Limitations and strengths*

We only used three age classes for categorizing paternal age. In order to scrutinize the possible U-shaped relationship discussed above, more age classes would be optimal. This was, however, hampered due to the relatively small sample sizes in our study. In this regard, large studies are clearly of benefit. As regards the evaluation of family history, we did not conduct a structured interview for the evaluation of each of the subject’s relatives. We therefore might have misclassified the status of family history, and this misclassification may have in turn obscured the true genetic influence on the risk association explored here.

On the other hand, the ethnic composition of the present sample was homogenous; that is, all of the subjects recruited for this study were Japanese and had been born in Japan. This is the first replication study on this topic conducted in an Asian country, and such reproducibility implies that the risk-increasing effect of advanced paternal age at birth is observable across different cultures, which further enhances the likelihood of a true association.

#### **Acknowledgments**

A part of this study was presented at the 12th Biennial Winter Workshop for Schizophrenia, Davos,

Switzerland, on February 10, 2004. This study was financially supported by a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan: Dr. Takei is a recipient of a Grant-in-Aid for Science Research (B) (2) No. 14370288. The authors would also like to thank Christina Dalman, MD, PhD, for her helpful comments.

## References

- Brown, A.S., Schaefer, C.A., Wyatt, R.J., Begg, M.D., Goetz, R., Bresnahan, M.A., Harkavy-Friedman, J., Gorman, J.M., Malaspina, D., Susser, E.S., 2002. Paternal age and risk of schizophrenia in adult offspring. *Am. J. Psychiatry* 159, 1528–1533.
- Byrne, M., Agerbo, E., Ewald, H., Eaton, W.W., Mortensen, P.B., 2003. Parental age and risk of schizophrenia: a case-control study. *Arch. Gen. Psychiatry* 60, 673–678.
- Dalman, C., Allebeck, P., 2002. Paternal age and schizophrenia: further support for an association. *Am. J. Psychiatry* 159, 1591–1592.
- El Saadi, O., Pedersen, C.B., McNeil, T.F., Saha, S., Welham, J., O'Callaghan, E., Cantor-Graae, E., Chant, D., Mortensen, P.B., McGrath, J., 2004. Paternal and maternal age as risk factors for psychosis: findings from Denmark, Sweden and Australia. *Schizophr. Res.* 67, 227–236.
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B.W., 1996. Structured Clinical Interview for DSM-IV Axis I Disorders (Version 2.0). American Psychiatric Publishing Inc., Arlington, VA. (Japanese translation by S. Takahashi, Japan UNI Agency Inc. Tokyo).
- Kawai, M., Minabe, Y., Takagai, S., Ogai, M., Matsumoto, H., Mori, N., Takei, N., 2004. Poor maternal care and high maternal body mass index in pregnancy as a risk factor for schizophrenia in offspring. *Acta Psychiatr. Scand.* 110, 257–263.
- Kendler, K.S., McGuire, M., Gruenberg, A.M., Walsh, D., 1995. Schizotypal symptoms and signs in the Roscommon family study. Their factor structure and familial relationship with psychotic and affective disorders. *Arch. Gen. Psychiatry* 52, 296–303.
- Kleinbaum, D.G., Kupper, L.L., Muller, K.E., Nizam, A., 1998. *Applied Regression Analysis and Other Multivariable Methods*, 3rd ed. Duxbury Press, Pacific Grove, CA.
- Malaspina, D., 2001. Paternal factors and schizophrenia risk: de novo mutations and imprinting. *Schizophr. Bull.* 27, 379–393.
- Malaspina, D., Harlap, S., Fennig, S., Heimann, D., Nahon, D., Feldman, D., Susser, E.S., 2001. Advancing paternal age and the risk of schizophrenia. *Arch. Gen. Psychiatry* 58, 361–367.
- Malaspina, D., Corcoran, C., Fahim, C., Berman, A., Harkavy-Friedman, J., Yale, S., Goetz, D., Goetz, R., Harlap, S., Gorman, J., 2002. Paternal age and sporadic schizophrenia: evidence for de novo mutations. *Am. J. Med. Genet.* 114, 299–303.
- Mata, I., Sham, P.C., Gilvary, C.M., Jones, P.B., Lewis, S.W., Murray, R.M., 2000. Childhood schizotypy and positive symptoms in schizophrenic patients predict schizotypy in relatives. *Schizophr. Res.* 44, 129–136.
- Matsumoto, H., Takei, N., Saito, H., Kachi, K., Mori, N., 1999. Childhood-onset schizophrenia and obstetric complications: a case-control study. *Schizophr. Res.* 38, 93–99.
- Matsumoto, H., Takei, N., Saito, H., Kachi, K., Mori, N., 2001. The association between obstetric complications and childhood-onset schizophrenia: a replication study. *Psychol. Med.* 31, 907–914.
- Pulver, A.E., McGrath, J.A., Liang, K.Y., Lasseter, V.K., Nestadt, G., Wolyniec, P.S., 2004. An indirect test of the new mutation hypothesis associating advanced paternal age with the etiology of schizophrenia. *Am. J. Med. Genet.* 124B, 6–9.
- Sham, P.C., MacLean, C.J., Kendler, K.S., 1993. Risk of schizophrenia and age difference with older siblings. Evidence for a maternal viral infection hypothesis? *Br. J. Psychiatry* 163, 627–633.
- Stata Corporation., 2003. *Stata Special Edition Version 8.1 for Windows*. Stata Corporation, College Station, TX.
- Zammit, S., Allebeck, P., Dalman, C., Lundberg, I., Henningson, T., Owen, M.J., Lewis, G., 2003. Paternal age and risk for schizophrenia. *Br. J. Psychiatry* 183, 405–408.

Research report

## Neural substrates in judgment process while playing go: a comparison of amateurs with professionals

Yasuomi Ouchi<sup>a,\*</sup>, Toshihiko Kanno<sup>a</sup>, Etsuji Yoshikawa<sup>b</sup>, Masami Futatsubashi<sup>b</sup>,  
Hiroyuki Okada<sup>b</sup>, Tatsuo Torizuka<sup>a</sup>, Mitsuo Kaneko<sup>a</sup>

<sup>a</sup>Positron Medical Center, Hamamatsu Medical Center, 5000 Hirakuchi, Hamakita, 434-0041, Japan

<sup>b</sup>Central Research Laboratory, Hamamatsu Photonics K.K. Hamakita, 434-8601, Japan

Accepted 15 October 2004

Available online 10 November 2004

### Abstract

A professional go player shows incomparable ability in judgment during go game. Positron emission tomography (PET) was used to investigate the neural substrates of professional go player's judgment process. Eight professional go players and six amateur players were instructed to think over silently in the opening-stage game (*fuseki*, territorial planning) problems and the life-or-death (*tsume*, checkmate judgment) problems presented on the monitor in front of them for 60 s of H<sub>2</sub><sup>15</sup>O PET scans and to state the answer afterwards. We found that in the territorial planning problems the parietal activation was equally observed in both groups with the additional prefrontal activation in the amateur group, and in the checkmate-decision problems the precuneus and cerebellum were activated in professionals while the premotor and parietooccipital cortices (visuospatial processing region) were extensively activated in amateurs. The comparison of the two groups showed stronger activations in the precuneus and cerebellum in the professionals in contrast to the premotor activation in amateurs during checkmate judgment. In addition, the cerebellum was remarkably activated in the higher ranking professional players. These findings suggested the cerebellum and precuneus play important roles in processing of accurate judgment by visual imagery and nonmotor learning memory processes in professional go players.

© 2004 Elsevier B.V. All rights reserved.

*Theme:* Neural basis of behavior

*Topic:* Cognition

*Keywords:* Go; Professional; Judgment; Precuneus; Cerebellum; Positron emission tomography

### 1. Introduction

There are several neuroimaging studies concerning the neural substrates of cognitive processes involved in professional thinking [13,20,25]. In these studies, the contribution of the prefrontal and parietal cortices was reported to be of great importance. Specifically, the precuneus was remarkably activated during the execution of precise topographical memory recall for professional taxi drivers [20] and in visuospatial processing for abacus

experts [13]. In addition to these cerebral activations in cognitive processes, recent findings on the cerebellum in nonmotor cognitive processes are worth noticing because the cerebellum is activated in many other cognitive tasks such as memory, error detection, attention, sensory discrimination and timing [1,2,10,15,18]. Furthermore, since the cerebellum also functions under the condition of the theory of mind [5], it is easy to speculate that an expert would exploit this “cognitive organ” more efficiently in cognitive tasks. However, this issue remains to be investigated.

The game go involves similar thought processes as the Western-style board game of chess. A previous activation study during a chess game revealed neural substrates of

\* Corresponding author. Fax: +81 53 585 0367.

E-mail address: [ouchi@pmc.hmedc.or.jp](mailto:ouchi@pmc.hmedc.or.jp) (Y. Ouchi).

problem-solving processes in the brain, except for the cerebellum, in nonprofessional players [21]. Playing go is different in nature from the chess game. Chessmen have different roles in chess, but go stones all have the same value. Thus, brain activity highly depends on the contemplation of the position of the stones, not on the selection of chessman. This may result because the perception of the objects determines the actions that can be made toward them [12]. The go game, which does not involve selection of objects, may provide simpler but likely more profound cognitive processing in terms of mental imagination [7]. Obtaining a professional title depends exclusively on the period of experience, knowledge, and ability and the ability to perform mental operations that amateurs cannot attain [7]. Visuospatial processing may be greater in go experts than in novices [14]. Thus, the professional rank may be used as an index of player's ability and utilized as a covariate for examining the neural correlates of the professional go player's judgment process. One activation study of the go game was recently reported that used fMRI [6], but the study design was different from ours in subjects (amateur vs. professional and amateur) and tasks (determining next move vs. judging in different situations the correct or incorrect responses).

The purpose of the current study was to investigate neural substrates of professional strategic thinking by comparing brain activations of certificated professional go players recruited from the Japan Go Association with the neural substrates of amateurs during deliberate life-or-death (checkmate judgment) and territorial planning situations of the game using positron emission tomography (PET) with  $H_2^{15}O$ .

## 2. Materials and methods

### 2.1. Subjects

Eight male right-handed professional go players (mean age  $\pm$  S.D.  $41.1 \pm 11.3$  years) from Japan Go Association and six amateur go players (four male and two female,  $59.5 \pm 5.4$  years) gave their informed consent to participate in the present experiment, which was approved by a local ethics committee at Hamamatsu Medical Center. Professionals had experienced for more than 15 years and practiced go almost everyday for various competitions; in contrast, amateurs had experienced for more than 10 years but practiced go on a recreational basis. Each participant underwent magnetic resonance imaging (MRI) using a 0.3 T static magnet system (MRP7000AD, Hitachi, Tokyo, Japan) before PET measurement, showing no morphological abnormality. All professional players were classified by the certificated professional rank ("dan" or grade ranging from first to ninth grade), consisting of one professional player with 1-dan, one with 3-dan, one with 4-dan, one with 7-dan, one with 8-dan, and three with 9-dan.

### 2.2. Tasks

In the real situation of a go game, two players, sitting face to face and holding black or white stones, take turns placing stones onto one of many vacant line intersections on the board. The goal is to occupy as large an area as possible with given stones. In the present study, three situations were given to all participants. In the first task, each subject performed one baseline task; (Base) staring at the center of the go board without any thought. Then each subject performed two nonmotor cognitive tasks; (Task 1) thinking about moving stones to expand their territory in the given situation on the screen (territorial planning), (Task 2) determining the final move in a life-or-death situation in order to occupy the territory in competition (checkmate judgment) (Fig. 1A). In the two nonmotor tasks, no motor performance was allowed during the scans. In Task 1, placing stones during this stage of the go game was unlimited and there was no correct place to put the stones.




Fig. 1. Go task and task performance. (A) Examples of stones arrangement for Task 1 (territorial planning) and Task 2 (checkmate judgment), (B) visual analog scales (VAS) for task difficulty (black bar) and rates of correct answer (CR, white bar) in professional and amateur groups, which was the result of Task 2. The asterisk indicates the presence of significant difference ( $p < 0.05$ ,  $\chi^2$  test).

However, in Task 2, there was only one way to solve the problem. In Task 2, the participants were required to choose one correct answer out of three options. Three different versions of each problem and two control tasks were alternatively presented in a counterbalanced manner. Each subject was instructed to refrain from moving the eyes during tasks and the ocular movement was monitored through the video focusing on the participant's face. The amateurs performed similar tasks with easier contents designed for nonprofessionals. Each task was presented on a screen three times for 90 s. The participants were required to state their answer for each Task 2 problem and assess the difficulty for each Task 1 and Task 2 problem as a visual analog scale (VAS, from the easiest 1 to the most difficult 10) every time after completion of each scan for Task 2. The VAS value for Task 2 was defined as follows: VAS value for Task 2=(individual VAS score in Task 2)/(mean VAS score averaged between Task 1 and Task 2)×100.

### 2.3. Apparatus and procedure

The experiment was conducted with a brain-purpose high-resolution Hamamatsu Photonics PET scanner [30] capable of yielding 47-slice images simultaneously with a spatial resolution of 2.9 mm (full width at half maximum) transaxially and 3.0 mm axially and with a 163-mm axial field of view. After a 10-min transmission scan for attenuation correction using a  $^{68}\text{Ge}/^{68}\text{Ga}$  source with the subject's head fixed by a radiosurgery-purpose thermo-plastic facemask, a 60-s emission scan was acquired from when the radiotracer first entered the cerebral circulation after intravenous bolus injection of 300 MBq of  $\text{H}_2^{15}\text{O}$  [26]. The timing of the PET start to collect the rising phase of the head curve radioactivity was described previously [16]. After back projection and filtering with a Hanning filter of a cut-off frequency of 0.2 cycles per pixel, image resolution of reconstructed regional cerebral blood flow was  $6.0\times 6.0\times$

3.6 mm full width at half maximum and the voxel size was measured to be  $1.3\times 1.3\times 3.4$  mm.

### 2.4. Data analysis

The whole-brain CBF data were analyzed using SPM99 software (Wellcome Department of Cognitive Neurology, London, UK, <http://www.fil.ion.ucl.ac/spm/spm99.html> [9]). The analytic procedure of SPM was basically the same as the methods in our previous reports [16,22]. Briefly, spatially normalized data based on the standard stereotaxic brain atlas [28] after being realigned to the first image data were smoothed by an isotropic Gaussian kernel of 8 mm. The effect of variance from global CBF was excluded by proportional scaling with the global CBF normalized to 50 ml/100g/min. So, the individual rCBF response was regarded as centered adjusted rCBF value. The resultant Z-maps were displayed on the three-dimensional MRI data obtained from all participants prior to each PET session. Between-group comparison was performed using VAS as a covariate because the difficulty of go problems presented was different from each group. In correlation analyses, the ranks in professional go players were used as covariates for testing the judgment process at different stages (both Task 1 and Task 2). Significant differences in CBF between conditions were estimated with a statistical threshold set at  $p<0.05$  corrected for multiple comparisons at voxel levels. In the SPM correlation analysis, statistical significance was given as  $p<0.001$ , uncorrected.

## 3. Results

### 3.1. Performance

There was no significant difference in VAS for task difficulty between groups. However, amateurs less frequently




Fig. 2. Brain activation during opening stage thinking (green) (Task 1 vs. Base) and the life-or-death judgment (red) (Task 2 vs. Base) irrespective of answer content in the amateur and professional go groups ( $p<0.05$ , corrected). Top row: amateur, bottom row: professional. Yellow denotes an overlap of green and red areas.

Table 1  
Activated brain regions in checkmate decision with and without correct answers relative to the baseline condition in professional and amateur go players

Group	Activated area	BA	Coordinates (x y z)	Z score
<i>Successful response</i>				
Professional	L precuneus	7	-12 -78 50	4.62
	R precuneus	7	18 -78 54	4.22
	R cerebellum (posterior lobe)	-	26 -44 -46	4.01
Amateur	Precuneus	7	0 -76 52	5.32
	R intraparietal sulcus	40/7	50 -44 42	4.98
	L precuneus	7	-16 -76 50	4.66
<i>Failed response</i>				
Professional	L supramarginal cortex	40	-60 -38 46	5.71
	R supramarginal cortex	40	60 -42 42	5.23
Amateur	R intraparietal sulcus	40/7	42 -52 44	6.60
	R precuneus	7	2 -76 48	6.27
	L intraparietal sulcus	40/7	-30 -60 48	5.94
	R superior precentral sulcus	6	30 -2 58	5.73
	L superior precentral sulcus	6	-24 6 58	5.39
	R middle temporal gyrus	19	42 -80 20	5.37
	L precuneus	7	-16 -72 50	5.32
	L supramarginal cortex	40	-50 -50 40	5.18

BA: Brodmann area, R: right, L: left.

made correct answers than professionals ( $p < 0.05$ ,  $\chi^2$  test) (Fig. 1B). The video showed various degrees of the eye blink and minimal eye movement during each scan in all participants, indicating that the present finding could eliminate the effect of saccadic eye movement. No motor behavior was found during scans in all subjects.

Table 2  
Brain regions significantly activated during judging correctly in two groups

Group	Activated area	BA	Coordinates (x y z)	Z score
Professional	R cerebellum (anterior lobe)	-	20 -74 -18	4.73
	L cerebellum (anterior lobe)	-	-24 -76 -14	4.43
	L precuneus	7	-18 -66 60	4.36
	R cerebellum (posterior lobe)	-	36 -48 -36	4.29
Amateur	L superior precentral sulcus	6	-22 0 60	4.87
	R cuneus	17	14 -102 0	4.60
	L supramarginal cortex	40	-60 -28 40	4.53
	L superior precentral sulcus	6	26 0 64	4.51

BA: Brodmann area, R: right, L: left.

### 3.2. PET results

Within-subject subtraction analysis irrespective of outcome (correct or false answer) showed extensive activations in the parietal and prefrontal cortices and cerebellum bilaterally in amateur go players in either type of task (upper images, Fig. 2), whereas more focal activated regions were observed in the parietal and temporooccipital cortices and right cerebellum in the professional group (bottom images, Fig. 2). When the players made correct answers in Task 2 (checkmate judgment), the bilateral superior parietal cortices (precuneus) and the right cerebellum was activated in the professional group, and the precuneus and intraparietal sulcus region were significantly activated in amateurs (Table 1, Fig. 3, green). When thinking incorrectly, the bilateral supramarginal cortices were activated in the professional group, and the broader cortical regions covering the premotor and parietal cortices bilaterally were activated in the amateur group (Table 1, Fig. 3, red).

In the checkmate judgment process with correct response, the precuneus and cerebellar cortex were more




Fig. 3. Brain activation during the life-or-death deliberation with (green) and without (red) correct answers in the amateur and professional go groups ( $p < 0.05$ , corrected). Top row: amateur, bottom row: professional. Yellow denotes an overlap of green and red areas.




Fig. 4. Brain regions more activated during the life-or-death deliberation with correct reply in the professional than amateur group (green), and more activated in the amateur than professional group (red) ( $p < 0.05$ , corrected).

strongly activated in the professional group, while the premotor and parietooccipital activation were stronger in the amateur group (Table 2, Fig. 4). Scattered plotting revealed that the CBF increase in the left premotor cortex correlated significantly with the task difficulty (VAS) in the amateur group (Fig. 5A,  $p < 0.05$ ,  $r = 0.62$ ), whereas the right cerebellar CBF increase correlated with the score in the professional group (Fig. 5B,  $p < 0.05$ ,  $r = 0.61$ ).

Correlation analysis in the professional group showed that the bilateral cerebellum in territorial planning (Task 1) and the right cerebellum in checkmate-decision (Task 2) correlated positively with professional ranks (Fig. 6,  $p < 0.001$ , uncorrected).

#### 4. Discussion

The present study revealed different neural substrates involved in cognitive processes between professional and amateur go players (irrespective of successful or failed outcome during the go game). The precuneus and cerebellum were engaged more in the checkmate judgment process in professional players, while the premotor and occipitoparietal cortices were remarkably activated in amateurs.

Although unsuccessful performance caused broader brain activation during the checkmate judgment in the amateur group (Fig. 3), only focal regions were detected in the professional group, suggesting more economical use of the brain energy in professional go players. As suggested in a recent study of professional musicians [17], the different cortical representations in judgment process could be interpreted as a result of cortical plasticity in highly skilled go players.

In view of correct or incorrect responses during the game, it was found that the precuneus was significantly activated during the checkmate judgment with correct answers in both professional and amateur players. This region connects to the anterior cingulate, prefrontal, lateral parietal and temporal cortices [29]. This anatomical connection suggests that the precuneus plays a role in orchestrating multimodal associative functions. Hence, almost all the participants, when performing successfully, would have utilized every part region of the brain related to the precuneus activity. It has been reported that the precuneus is activated more during silent tasks for motor imagery [4,24] and during motor imagery of complete finger movement than during explicit execution [11]. Thus, it is suggested that the precuneus might allow complete mental reproduction of the configuration of go stones on the imagined board in the players with correct responses. The left precuneus was engaged in this strategic thinking in professional players. Therefore, one can assume that professional go players exploit this ability efficiently and make the most of the motor imagery skills through highly vigilant or conscious retrieval of acquired memory [2]. In contrast, unsuccessful deliberation caused supramarginal activation in professionals and the prefrontal-temporoparietal activation in amateurs. A recent fMRI study showed that the supramarginal cortex might be involved in an enactment effect that improves performance encoding ability [27]. This suggests that trial and error of different




Fig. 5. Correlation analyses between VAS scores (%) and premotor CBF responses (%) in the amateur group (A) and between VAS scores and cerebellar CBF responses in the professional group (B).






Fig. 6. Activated brain regions significantly correlated with professional “dan” ranks during the territorial planning (red) and the life-or-death judgment (green) in the professional group ( $p < 0.001$ , uncorrected).

moves might have activated the supramarginal cortex in professional players. In contrast with the professional go players, amateur players lacking in strategic or enactment ability required incorporation of the prefrontal (premotor) and parietooccipital cortex, including the precuneus, to solve the problem (Fig. 4). This premotor activity was positively associated with the difficulty of the go problem (Fig. 5A). This pattern of activation in amateurs is consistent with the activation results in a recent fMRI study showing that the premotor and precuneus regions were activated conjointly during visuomotor operation [13] and in a chess study revealing activations of the superior frontal and occipitoparietal regions during checkmate judgment [21]. The difference of these patterns in judgment processes between professional and amateur go players may reflect a distinctive professional way of thinking leading to greater efficiency, which requires a smaller number of active neurons in specific regions incorporated in the critical situations during the go game.

Another interesting result from the comparison of the professional and amateur group in the present study was that the cerebellum was more engaged in the endgame strategy in professional players. The cerebellum has been implicated in motor imagery [8,19] and problem solving [18] in nonmotor cognitive operations. As the rank of professional go players became higher, this cerebellar cortex was more activated as shown in Fig. 6. In addition, this cerebellar activity correlated positively with VAS for task difficulty (Fig. 5B). This seemingly contradictory finding suggests that VAS of professionals in the present study might indicate uneasy emotion, because the cerebellar activation was correlated with gradual increase in task-demand during mental rehearsal [3]. Actually, some higher ranking professionals exhibited high VAS scores and excellent achievement. Thus, the cerebellum in the professional players might be associated with predictive control that guides online imagery motor performance during tense situations of the go game.

There are a few methodological caveats that have to be taken into account for the present study. The task contents in life-or-death problems given to amateurs were different from those given to professionals because amateurs would

not have been able to solve the difficult tasks given to the professionals. Although the VAS for task difficulty was found similar between the groups, the strategy to solve the problem might be different. The present study did not attempt to elucidate each step of problem solving or strategic cognitive processes. This issue may be important for unraveling the mystery of the expert’s mind. The questionnaire after each session of PET scans revealed that some professionals had solved the problems early within the 60-s period and spent the remaining times verifying the answer. This might cause not only weaker activation of brain region responsible for the execution of solving tasks, but also incorporation of brain regions irrelevant to the judgment process. Thus, the present result might reflect the summation of neural substrates for judgment and imagery retrieval processes occurring in the professional mind.

In summary, our results have shown that the professional judgment requires the precuneus and cerebellar activations during the go game. This suggests that visual imagery and motor imagination may be important for the highly skilled tactics of professionals. In contrast, it seems that the extensive frontoparietal regions functioning partly in visuomotor processing operate dominantly in execution of solving problems in amateur go players. In view of brain rehabilitation for the elderly and patients with dementia, playing a game of go may cause premotor activation, which may be of great value for stimulating the brain, because incorporation of the premotor region activity is necessary for executing arithmetic cognitive tasks in Alzheimer’s disease patients [23]. The different pattern of cognitive processes between expert and amateur go players may reflect the brain functional plasticity or functional specialization that is acquired later in domain-specific experts.

#### Acknowledgments

We would like to thank Mr. Zenta Takekawa (Japan Go Association) for his task selection, Mr. Fumitoshi Nakamura and the staff of the Japan Go Association for their constant support in the present study.

## References

- [1] G. Allen, R.B. Buxton, E.C. Wong, et al., Attentional activation of the cerebellum independent of motor involvement, *Science* 275 (1997) 1940–1943.
- [2] N.C. Andreasen, D.S. O'Leary, S. Paradiso, et al., The cerebellum plays a role in conscious episodic memory retrieval, *Hum. Brain Mapp.* 8 (1999) 226–234.
- [3] H. Boecker, A.O. Ceballos-Baumann, P. Bartenstein, et al., A H<sub>2</sub>(15)O positron emission tomography study on mental imagery of movement sequences—the effect of modulating sequence length and direction, *Neuroimage* 17 (2002) 999–1009.
- [4] E. Bonda, M. Petrides, S. Frey, Neural correlates of mental transformations of the body-in-space, *Proc. Natl. Acad. Sci. U. S. A.* 92 (1995) 11180–11184.
- [5] C. Calarge, N.C. Andreasen, D.S. O'Leary, Visualizing how one brain understands another: a PET study of theory of mind, *Am. J. Psychiatry* 160 (2003) 1954–1964.
- [6] X. Chen, D. Zhang, X. Zhang, et al., A functional MRI study of high-level cognition: II. The game of GO, *Brain Res. Cogn. Brain Res.* 16 (2003) 32–37.
- [7] C. Cho, *Go: A Complete Introduction of the Game*, Kiseido Publishing, Tokyo, 1997, 1–138 pp.
- [8] J. Decety, H. Sjöholm, E. Ryding, et al., Cerebellum participates in mental activity: tomographic measurements of regional cerebral blood flow, *Brain Res.* 535 (1990) 313–317.
- [9] K.J. Friston, A.P. Holmes, K.J. Worsley, et al., Statistical parametric mapping in functional imaging: a general linear approach, *Hum. Brain Mapp.* 2 (1995) 189–210.
- [10] J.H. Gao, L.M. Parsons, J.M. Bower, et al., Cerebellum implicated in sensory acquisition and discrimination rather than motor control, *Science* 272 (1996) 545–547.
- [11] E. Gerardin, A. Sirigu, S. Lehericy, et al., Partially overlapping neural networks for real and imagined hand movements, *Cereb. Cortex* 10 (2000) 1093–1104.
- [12] J. Grezes, J. Decety, Does visual perception of object afford action? Evidence from a neuroimaging study, *Neuropsychologia* 40 (2002) 212–222.
- [13] T. Hanakawa, M. Honda, T. Okada, et al., Neural correlates underlying mental calculation in abacus experts: a functional magnetic resonance imaging study, *Neuroimage* 19 (2003) 296–307.
- [14] T. Hatta, T. Kogure, A. Kawakami, Hemisphere specialization of Go experts in visuospatial processing, *Am. J. Psychol.* 112 (1999) 571–584.
- [15] M. Ito, Movement and thought: identical control mechanisms by the cerebellum, *Trends Neurosci.* 16 (1993) 448–450.
- [16] M. Iwase, Y. Ouchi, H. Okada, et al., Neural substrates of human facial expression of pleasant emotion induced by comic films: a PET Study, *Neuroimage* 17 (2002) 758–768.
- [17] L. Jancke, N.J. Shah, M. Peters, Cortical activations in primary and secondary motor areas for complex bimanual movements in professional pianists, *Brain Res. Cogn. Brain Res.* 10 (2000) 177–183.
- [18] S.G. Kim, K. Ugurbil, P.L. Strick, Activation of a cerebellar output nucleus during cognitive processing, *Science* 265 (1994) 949–951.
- [19] A.R. Luft, M. Skalej, A. Stefanou, et al., Comparing motion- and imagery-related activation in the human cerebellum: a functional MRI study, *Hum. Brain Mapp.* 6 (1998) 105–113.
- [20] E.A. Maguire, R.S. Frackowiak, C.D. Frith, Recalling routes around London: activation of the right hippocampus in taxi drivers, *J. Neurosci.* 17 (1997) 7103–7110.
- [21] P. Nichelli, J. Grafman, P. Pietrini, et al., Brain activity in chess playing, *Nature* 369 (1994) 191.
- [22] Y. Ouchi, H. Okada, E. Yoshikawa, et al., Brain activation during maintenance of standing postures in humans, *Brain* 122 (1999) 329–338.
- [23] Y. Ouchi, E. Yoshikawa, M. Futatsubashi, et al., Activation in the premotor cortex during mental calculation in patients with Alzheimer's disease: relevance of reduction in posterior cingulate metabolism, *Neuroimage* 22 (2004) 155–163.
- [24] L.M. Parsons, P.T. Fox, J.H. Downs, et al., Use of implicit motor imagery for visual shape discrimination as revealed by PET, *Nature* 375 (1995) 54–58.
- [25] M. Pesenti, L. Zago, F. Crivello, et al., Mental calculation in a prodigy is sustained by right prefrontal and medial temporal areas, *Nat. Neurosci.* 4 (2001) 103–107.
- [26] M.E. Raichle, W.E.W. Martin, P. Herscovitch, et al., Brain blood flow measured with intravenous H<sub>2</sub>(15)O: II. Implementation and validation, *J. Nucl. Med.* 24 (1983) 790–798.
- [27] M.O. Russ, W. Mack, C.R. Grama, et al., Enactment effect in memory: evidence concerning the function of the supramarginal gyrus, *Exp. Brain Res.* 149 (2003) 497–504.
- [28] J. Talairach, P. Tournoux, *Co-planer Stereotaxic Atlas of the Human Brain: 3-Dimensional Proportional System: An Approach to Cerebral Imaging*, Georg Thieme, Stuttgart, 1988.
- [29] E.P.M. Vianna, G. Van Hoesen, J. Parvizi, Efferent connections of the primate posterior cingulate, retrosplenial and mesial parietal cortices, *J. Neurosci.* 587 (2002) 12.
- [30] M. Watanabe, K. Shimizu, T. Omura, et al., A new high resolution PET scanner dedicated to brain research, *IEEE Trans. Nucl. Sci.* 49 (2002) 634–639.

## Cerebral hemodynamics evaluation by near-infrared time-resolved spectroscopy: Correlation with simultaneous positron emission tomography measurements

Etsuko Ohmae,<sup>a,\*</sup> Yasuomi Ouchi,<sup>b</sup> Motoki Oda,<sup>a</sup> Toshihiko Suzuki,<sup>a</sup> Shuji Nobesawa,<sup>b</sup> Toshihiko Kanno,<sup>b</sup> Etsuji Yoshikawa,<sup>a</sup> Masami Futatsubashi,<sup>a</sup> Yukio Ueda,<sup>a</sup> Hiroyuki Okada,<sup>a</sup> and Yutaka Yamashita<sup>a</sup>

<sup>a</sup>Central Research Laboratory, Hamamatsu Photonics K.K., 5000 Hirakuchi, Hamamatsu, Shizuoka 434-8601, Japan

<sup>b</sup>Positron Medical Center, Hamamatsu Medical Center, 5000 Hirakuchi, Hamamatsu, Shizuoka 434-0041, Japan

Received 18 August 2004; revised 3 March 2005; accepted 4 August 2005  
Available online 13 September 2005

We compared pharmacologically-perturbed hemodynamic parameters (cerebral blood volume; CBV, and flow; CBF) by acetazolamide administration in six healthy human subjects studied with positron emission tomography (PET) and near-infrared (NIR) time-resolved spectroscopy (TRS) simultaneously to investigate whether NIR-TRS could measure *in vivo* hemodynamics in the brain tissue quantitatively. Simultaneously with the PET measurements, TRS measurements were performed at the forehead with four different optode spacing from 2 cm to 5 cm. Total hemoglobin and oxygen saturation (SO<sub>2</sub>) measured by TRS significantly increased after administration of acetazolamide at any optode spacing in all subjects. In PET study, CBV and CBF were estimated in the following three volumes of interest (VOIs) determined on magnetic resonance images, VOI<sub>1</sub>: scalp and skull, VOI<sub>2</sub>: gray matter region, VOI<sub>3</sub>: gray and white matter regions. Acetazolamide treatment elevated CBF and CBV significantly in VOI<sub>2</sub> and VOI<sub>3</sub> but VOI<sub>1</sub>. TRS-derived CBV was more strongly correlated with PET-derived counterpart in VOI<sub>2</sub> and VOI<sub>3</sub> when the optode spacing was above 4 cm, although optical signal from cerebral tissue could be caught with any optode spacing. As to increase of the CBV, 4 cm of optode spacing correlated best with VOI<sub>2</sub>. To support the result of TRS-PET experiment, we also estimated the contribution ratios of intracerebral tissue to observed absorption change based on diffusion theory. The contribution ratios at 4 cm were estimated as follows: 761 nm: 50%, 791 nm: 72%, 836 nm: 70%. These results demonstrated that NIR-TRS with 4 cm of optode spacing could measure cerebral hemodynamic responses optimally and quantitatively.

© 2005 Elsevier Inc. All rights reserved.

### Introduction

Near-infrared spectroscopy (NIRS) allows simple, non-invasive measurement of the oxygenation state and hemodynamics in living tissue by utilizing the differential in absorption spectrum between oxygenated and deoxygenated hemoglobin. This field got its start from the finding by Jobsis (1977) that when a cat's head is irradiated with near-infrared (NIR) light, the intensity of the transmitted light shows changes according to the oxygen metabolic state in the tissues. Since then, there has been growing study and technique of NIRS measurement.

Making that advantage, this method has been expected for use in surgical operations (Kakahana et al., 1996; De Blasi et al., 1997) and neonate respiration care (Meek et al., 1999; Isobe et al., 2000). Besides the clinical field, topographical imaging by multi-channels measurement is being performed to observe brain activity on the cortex (Watanabe et al., 2000; Tanosaki et al., 2001).

NIRS encompasses some different techniques and analysis, and we are adopting approaches of time-resolved spectroscopy (TRS; Oda et al., 1996; Yamashita et al., 1998), phase modulated spectroscopy (PMS; Tuchiya and Urakami, 1996; Iwai et al., 2001) or spatially resolves spectroscopy (SRS; Suzuki et al., 1999) method, etc. to quantification.

In contrast to the wide applicability of NIRS to brain monitoring, fundamental and critical questions still remain to be clarified, one of which is light propagation in the human head. The effect of the various external tissues of the head such as skin, skull and cerebrospinal fluid (CSF) on photon propagation in the internal cerebral tissue has not yet been fully examined *in vivo*.

Several researchers (Firbank et al., 1995; Okada et al., 1997) expressed doubts about the use of NIRS on adult human heads due to problems from the multi-layered structure of the scalp, skull, CSF and the cerebral tissue. Firbank first showed that the presence of CSF had a significant effect on the light distribution. It was

---

\* Corresponding author. Fax: +81 53 586 6180.

E-mail address: etuko-o@crl.hpk.co.jp (E. Ohmae).

Available online on ScienceDirect (www.sciencedirect.com).

reported that the NIRS signal from the adult human head was only 10–20% of the total signal due to the effect of CSF from the Monte Carlo simulation. These doubts are related to the essential question of where to measure using the photon, and research studies are being conducted using both simulations (Okada and Delpy, 2000, 2003a,b; Misonoo and Okada, 2001) and experimental measurements. On the contrary, several researchers (McCormick et al., 1992; Harris et al., 1994; Germon et al., 1995; Kohri et al., 2001) have reported that NIRS can detect the brain signals more specifically by increasing the optode spacing from experimental measurements.

In this study, in order to investigate the relation between the optode spacing and light sampling depth, we observed change in the cerebral blood volume (CBV) of six adult subjects by administration of a drug with simultaneous measurement of the TRS system which can measure the blood volume and the oxygen saturation ( $\text{SO}_2$ ) quantitatively and positron emission tomography (PET), and we compared the CBV by TRS (TRS CBV) with CBV by PET (PET CBV) and estimated the contribution ratios of intracerebral tissue to the observed absorption change at three different wavelengths.

## Materials and methods

### Subjects

Six healthy male subjects (mean age,  $42.6 \pm 5.08$ ; range, 37 to 51 years) were studied. Informed consent was obtained from all subjects before experiment. It was confirmed that they had no previous history of intracranial disorders and also that there were no anatomical abnormalities by making a check with a magnetic resonance imaging (MRI; 0.3 T MRP7000AD, Hitachi Ltd., Japan).

### Three-wavelength TRS system

We used TRS-10 system (Hamamatsu Photonics K.K., Japan) (Oda et al., 2000) to obtain TRS-CBV in our experiment. This system uses time-correlate single photon counting (TCPC) method for

measuring the temporal function of the sample. The system measures the intensity of light in a time domain and enables analysis of the data with the time domain photo diffusion equation (Patterson et al., 1989). The block diagram of this system is shown in Fig. 1.

As the light source, this system uses semiconductor lasers called “Picosecond Light Pulser (PLP, Hamamatsu Photonics K.K., Japan)” emitting light pulses at three different wavelengths (761 nm, 791 nm, 836 nm) with a peak power of 60 mW, average power of 30  $\mu\text{W}$ , the full width at half maximum (FMHM) of 100 ps and repetition frequency of 5 MHz. The detector section of this system consists of a photomultiplier tube (PMT, H6279-MOD, Hamamatsu Photonics K.K., Japan) followed by constant fraction discriminators (CFD), time-to-amplitude converters (TAC), A/D converters and histogram memories.

The system instrumental function is about 160 ps FMWH. The three PLPs emit light pulses on a time series, and the 3-wavelength optical pulses (761, 791, 836 nm) are guided into one optical fiber via a fiber coupler (CH20G-D3-CF, Mitsubishi Gas Chemical Company Inc., Japan). An optical switch (SC SERIES, JDS FITEL Inc., Canada) in this experiment selected the light irradiation point. A neutral density filter installed between the optical switch and each irradiation fiber maintained the light entering the PMT at a correct level. Each single optical fiber (GC200/250L, FUJIKURA Ltd., Japan) used for light irradiation has a numerical aperture (N.A.) of 0.25 and a core diameter of 200  $\mu\text{m}$ . The optical bundle fiber (LB21E, HOYA Corp., Japan) used to collect the light has an N.A. of 0.21 and a bundle diameter of 3 mm.

### TRS data analysis

The observed temporal profiles were fitted into the photon diffusion equation (Patterson et al., 1989) using the non-linear least square fitting method. The reduced scattering ( $\mu_s'$ ) and absorption coefficients ( $\mu_a$ ) for three wavelengths were calculated (Appendix A). Then oxyhemoglobin (TRS  $\text{HbO}_2$ ), deoxyhemoglobin (TRS Hb), total hemoglobin (TRS tHb) and oxygen saturation ( $\text{SO}_2$ ) were calculated with least square method (Appendix B). We then converted the TRS tHb into the TRS CBV for comparison with the PET CBV (Appendix C). Additionally, we calculated the partial mean pathlength of extracerebral tissue ( $L_{\text{ext}}$ ) at each wavelength




Fig. 1. Block diagram of time-resolved spectroscopy system (TRS-10: Hamamatsu Photonics K.K.).