

REFERENCES

- Fukuyama Y, Osawa M, Suzuki H. Congenital progressive muscular dystrophy of the Fukuyama type—Clinical, genetic and pathological considerations. *Brain Dev* 1981;3:1–29
- Kamoshita S, Konishi Y, Segawa M, Fukuyama Y. Congenital muscular dystrophy as a disease of the central nervous system. *Arch Neurol* 1976; 33:513–16
- Kobayashi K, Nakahori Y, Miyake M, et al. An ancient retrotransposal insertion causes Fukuyama-type congenital muscular dystrophy. *Nature* 1998;394:388–92
- Barkovich AJ, Kuzniecky RI, Dobyns WB, Jackson GD, Becker LE, Evrard P. A classification scheme for malformations of cortical development. *Neuropediatrics* 1996;27:59–63
- Takada K, Nakamura H, Tanaka J. Cortical dysplasia in congenital muscular dystrophy with central nervous system involvement (Fukuyama type). *J Neuropathol Exp Neurol* 1984;43:395–407
- Nakano I, Funahashi M, Takada K, Toda T. Are breaches in the glia limitans the primary cause of the micropolygyria in Fukuyama-type congenital muscular dystrophy (FCMD)? Pathological study of the cerebral cortex of an FCMD fetus. *Acta Neuropathol (Berl)* 1996;9: 313–21
- Yamamoto T, Shibata N, Kanazawa M, et al. Early ultrastructural changes in the central nervous system in Fukuyama congenital muscular dystrophy. *Ultrastruct Pathol* 1997;2:355–60
- Saito Y, Murayama S, Kawai M, Nakano I. Breached cerebral glia limitans—basal lamina complex in Fukuyama-type congenital muscular dystrophy. *Acta Neuropathol (Berl)* 1999;98:330–36
- Ishii H, Hayashi YK, Nonaka I, Arahata K. Electron microscopic examination of basal lamina in Fukuyama congenital muscular dystrophy. *Neuromuscul Disord* 1997;7:191–97
- Takada K, Rin YS, Kasagi S, Sato K, Nakamura H, Tanaka J. Long survival in Fukuyama congenital muscular dystrophy: Occurrence of neurofibrillary tangles in the nucleus basalis of Meynert and locus ceruleus. *Acta Neuropathol* 1986;71:228–32
- Oka A, Itoh M, Takashima S. The early induction of cyclooxygenase 2 associated with neurofibrillary degeneration in brains of patients with Fukuyama-type congenital muscular dystrophy. *Neuropediatrics* 1999;30: 34–37
- Wisniewski K, Jervis GA, Moretz RC, Wisniewski HM. Alzheimer neurofibrillary tangles in diseases other than senile and presenile dementia. *Ann Neurol* 1979;5:288–94
- Mandybur TI, Nagpaul AS, Pappas Z, Niklowitz WJ. Alzheimer neurofibrillary change in subacute sclerosing panencephalitis. *Ann Neurol* 1977;1:103–7
- Suzuki K, Parker CC, Pentchev PG, et al. Neurofibrillary tangles in Niemann-Pick disease type C. *Acta Neuropathol* 1995;89:227–38
- Autio-Harmanen H, Oldfors A, Sourander P, Renlund M, Dammert K, Simila S. Neuropathology of Salla disease. *Acta Neuropathol (Berl)* 1988; 75:481–90
- Auer IA, Schmidt ML, Lee VM, et al. Paired helical filament tau (PHFtau) in Niemann-Pick type C disease is similar to PHFtau in Alzheimer's disease. *Acta Neuropathol (Berl)* 1995;90:547–51
- Vermersch P, Sergeant N, Ruchoux MM, et al. Specific tau variants in the brains of patients with myotonic dystrophy. *Neurology* 1996;47:711–17
- The National Institute on Aging, and Reagan Institute Working Group on Diagnostic Criteria for the Neuropathological Assessment of Alzheimer's Disease. *Neurobiol Aging* 1997;18(suppl):S1–2
- Toda T, Miyake M, Kobayashi K, et al. Linkage-disequilibrium mapping narrows the Fukuyama-type congenital muscular dystrophy (FCMD) candidate region to 100 kb. *Am J Hum Genet* 1996;59:1313–20
- Toda T, Ikegawa S, Okui K, et al. Refined mapping of a gene responsible for Fukuyama-type congenital muscular dystrophy: Evidence for strong linkage disequilibrium. *Am J Hum Genet* 1994;55:946–50
- Gallyas F. Silver staining of Alzheimer's neurofibrillary changes by means of physical development. *Acta Morphol Acad Sci Hung* 1971;19:1–8
- Grogan TM, Casey TT, Miller PC, Rangel CS, Nunnery DW, Nagle RB. Automation of immunohistochemistry. *Adv Pathol Lab Med* 1993;6:253–83
- Grogan TM, Rangel C, Rimsza L, et al. Kinetic-mode, automated double-labeled immunohistochemistry and in situ hybridization in diagnostic pathology. *Adv Pathol Lab Med* 1995;8:79–100
- Hasegawa M, Jakes R, Crowther RA, Lee VM, Ihara Y, Goedert M. Characterization of mAb AP422, a novel phosphorylation-dependent monoclonal antibody against tau protein. *FEBS Lett* 1996;384:25–30
- Umeda Y, Taniguchi S, Arima K, et al. Alterations in human tau transcripts correlate with those of neurofilaments in sporadic tauopathies. *Neurosci Lett* 2004;359:151–54
- Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol (Berl)* 1991;82:239–59
- Saito Y, Mizuguchi M, Oka A, Takashima S. Fukutin protein is expressed in neurons of the normal developing human brain but is reduced in Fukuyama-type congenital muscular dystrophy brain. *Ann Neurol* 2000; 47:756–64
- Sasaki J, Ishikawa K, Kobayashi K, et al. Neuronal expression of the fukutin gene. *Hum Mol Genet* 2000;9:3083–90
- Yamamoto T, Kato Y, Karita M, et al. Fukutin expression in glial cells and neurons: Implication in the brain lesions of Fukuyama congenital muscular dystrophy. *Acta Neuropathol (Berl)* 2002;104:217–24
- Michele DE, Barresi R, Kanagawa M, et al. Post-translational disruption of dystroglycan-ligand interactions in congenital muscular dystrophies. *Nature* 2002;418:417–22
- Sugita S, Saito F, Tang J, Satz J, Campbell K, Sudhof TC. A stoichiometric complex of neuexins and dystroglycan in brain. *J Cell Biol* 2001;154: 435–45
- Liu F, Iqbal K, Grundke-Iqbal I, Hart GW, Gong CX. O-GlcNAcylation regulates phosphorylation of tau: A mechanism involved in Alzheimer's disease. *Proc Natl Acad Sci U S A* 2004;101:10804–9
- Maroun LL, Graem N. Autopsy standards of body parameters and fresh organ weights in nonmacerated and macerated human fetuses. *Pediatr Dev Pathol* 2005;8:204–17