

It has been reported that various types of bacteria are detected in oral maxillofacial infections and periodontal diseases, and many efforts have been made to determine the pathogenicity of these bacteria. Nevertheless, the relationship between bacteria and POMC symptoms has not been widely investigated. Our results suggest that bacteria, frequently detected in the oral cavity, were present in the fluids of POMC, and that these bacteria were partially responsible for the progression of POMC symptoms, although further studies on the presence of bacterial species other than the 20 species selected in this study and their etiologic roles are required. PCR methods are effective in detecting fastidious bacteria in maxillofacial infections and these methods will assist in conventional bacteriological examination, including culturing methods.

Acknowledgments

This study was supported in part by Grants-in-Aid for Scientific Research (17659659, 17591985, 16390601) from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References

- Ashimoto A, Chen C, Bakker I, Slots J. Polymerase chain reaction detection of 8 putative periodontal pathogens in subgingival plaque of gingivitis and advanced periodontitis lesions. *Oral Microbiol Immunol* 1996; 11: 266–273.
- Brook I, Frazier EH. Infections caused by *Propionibacterium* species. *Rev Infect Dis* 1991; 13: 819–822.
- Carranza FA, Bulkacz J. Defense mechanisms of the gingival. In: Carranza FA, Newman MG, eds. *Clinical Periodontology, 8th edn*. Philadelphia: W.B. Saunders, 1996: 103–107.
- Clarridge JE III, Attorri S, Musher DM, Hebert J, Dunbar S. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* ('*Streptococcus milleri* Group') are of different clinical importance and are not equally associated with abscess. *Clin Infect Dis* 2001; 32: 1511–1515.
- Csukas Z, Banizs B, Rozgonyi F. Studies on the cytotoxic effects of *Propionibacterium acnes* strains isolated from cornea. *Microb Pathog* 2004; 36: 171–174.
- Doan N, Contreras A, Flynn J, Slots J, Chen C. Molecular identification of *Dialister pneumosintes* in subgingival plaque of humans. *J Clin Microbiol* 2000; 38: 3043–3047.
- Dymock D, Weighan AI, Scully C, Wade WG. Molecular analysis of microflora associated with dentoalveolar abscesses. *J Clin Microbiol* 1996; 34: 537–542.
- Eady EA, Ingham E. *Propionibacterium acnes* – friend or foe. *Rev Med Microbiol* 1994; 5: 163–173.
- Furcht C, Eschrich K, Merte K. Detection of *Eikenella corrodens* and *Actionobacillus actionomycetemcomitans* by use of the polymerase chain reaction (PCR) *in vitro* and in subgingival plaque. *J Clin Periodontol* 1996; 23: 891–897.
- Gardner DG, Gullane PJ. Mucocoeles of the maxillary sinus. *Oral Surg Oral Med Oral Pathol* 1986; 62: 538–543.
- Gothberg KAT, Little JW, King DR, Bean RL. A clinical study of cysts arising from mucosa of the maxillary sinus. *Oral Surg Oral Med Oral Pathol* 1976; 41: 52–58.
- Hadar T, Shvero J, Nogevis BI, Yaniv E. Mucus retention cyst of the maxillary sinus: the endoscopic approach. *Br J Oral Maxillofac Surg* 2000; 38: 227–229.
- Hashimura T, Sato M, Hoshino E. *Mogibacterium timidum* and *Eubacterium saphenum* from pulpal and periradicular samples using the polymerase chain reaction (PCR) method. *Int Endod J* 2001; 34: 463–470.
- Heim Dahl A, von Konow L, Satoh T, Nord CE. Clinical appearance of orofacial infections of odontogenic origin in relation to microbiological findings. *J Clin Microbiol* 1985; 22: 299–302.
- Hykin PG, Tobal K, McIntyre G, Matheson MM, Towler HM, Lighan SL. The diagnosis of delayed post-operative endophthalmitis by polymerase chain reaction of bacterial DNA in vitreous samples. *J Med Microbiol* 1994; 40: 408–415.
- Imai T. Mucocoele of the maxillary sinus after radical operation for sinusitis. *Z Otol Tokyo* 1933; 39: 723–735.
- Ito K, Miyata H, Watanabe K, Ueno K. Bacteriology of infectious disease in otorhinolaryngology (1). Bacteriological study of paranasal cyst. *Nippon Jibiinkouka Gakkaikai Kaiho* 1992; 95: 1229–1238.
- Jacobs JA, Schot CS, Bunschoten AE, Schouls LM. Rapid species identification of '*Streptococcus milleri*' strains by line blot hybridization: identification of a distinct 16S rRNA population closely related to *Streptococcus constellatus*. *J Clin Microbiol* 1996; 34: 1717–1721.
- Jappe U, Ingham E, Henwood J, Holland KT. *Propionibacterium acnes* and inflammation in acne; *P. acnes* has T-cell mitogenic activity. *Br J Dermatol* 2002; 146: 202–209.
- Kanamori S, Kusano N, Shinzato T, Saito A. The role of the capsule of the *Streptococcus milleri* group in the pathogenicity. *J Infect Chemother* 2004; 10: 105–109.
- Kazor CE, Mitchell PM, Lee AM, Stokes LN, Loesche WJ, Dewhirst FE, Paster BJ. Diversity of bacterial populations on the tongue dorsa of patients with halitosis and healthy patients. *J Clin Microbiol* 2003; 41: 558–563.
- Kubo I. A buccal cyst occurred after radical operation of the maxillary sinus. *Z Otol Tokyo* 1927; 33: 896–897.
- Kuriyama T, Karasawa T, Nakagawa K, Saiki Y, Yamamoto E, Nakamura S. Bacteriologic features and antimicrobial susceptibility in isolates from orofacial odontogenic infections. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2000; 90: 600–608.
- Kuriyama T, Karasawa T, Nakagawa K, Yamamoto E, Nakamura S. Bacteriology and antimicrobial susceptibility of gram-positive cocci isolated from pus specimens of orofacial odontogenic infections. *Oral Microbiol Immunol* 2002; 17: 132–135.
- Kuriyama T, Nakagawa K, Kawashiri S, Yamamoto E, Nakamura S, Karasawa T. The virulence of mixed infection with *Streptococcus constellatus* and *Fusobacterium nucleatum* in a murine orofacial infection model. *Microbes Infect* 2000; 2: 1425–1430.
- Kuriyama T, Nakagawa K, Kawashiri S, Yamamoto E, Saiki Y. Study of pathogens of odontogenic infections using the oral floor abscess model in mouse. – potential of *Streptococcus constellatus* and *Fusobacterium nucleatum* for abscess production. – *Jpn J Oral Maxillofac Surg* 1998; 44: 382–387.
- Lane DJ. 16S/23S rRNA sequencing. In: Stackebrandt E, Goodfellow M, eds. *Nucleic Acid Techniques in Bacterial Systematics*. Chichester: John Wiley & Sons, 1991: 115–175.
- Linton AH, Hinton MH. The normal microbiota of the body. In: Linton AH, Heather MD, eds. *Topley and Wilson's Principles of Bacteriology and Immunity, Vol. 2, 8th edn*. London: Edward Arnold, 1990: 312–329.
- Mandell GL, Bennett JE, Mandell DR. *Douglas and Benett's Principles and Practice of Infectious Diseases, 5th edn*. London: A. Harcourt Health Sciences Company, Churchill Livingstone, 2000.
- Marks SC, Latoni JD, Mathog RH. Mucocoeles of the maxillary sinus. *Otolaryngol Head Neck Surg* 1997; 117: 18–21.
- Moore WE, Moore LV. The bacteria of periodontal diseases. *Periodontol* 2000; 1994; 5: 66–77.
- Niederman R, Zhang J, Kashket S. Short-chain carboxylic-acid-stimulated, PMN-mediated gingival inflammation. *Crit Rev Oral Biol Med* 1997; 8: 269–290.
- Okamoto M, Maeda N, Kondo K, Leung KP. Hemolytic and hemagglutinating activities of *Prevotella intermedia* and *Prevotella nigrescens*. *FEMS Microbiol Lett* 1999; 178: 299–304.
- Paju S, Bernstein JM, Hasse EM, Scannapieco FA. Molecular analysis of bacterial flora associated with chronically inflamed maxillary sinuses. *J Med Microbiol* 2003; 52: 591–597.
- Paster BJ, Boches SK, Galvin JL, Ericson RE, Lau CN, Levanos VA, et al. Bacterial diversity in human subgingival plaque. *J Bacteriol* 2001; 183: 3770–3783.
- Potempa J, Banbula A, Travis J. Role of bacterial proteinases in matrix destruction and modulation of host responses. *Periodontol* 2000; 24: 153–192.
- Roche Y, Yoshimori RN. *In-vitro* activity of spiramycin and metronidazole alone in combination against clinical isolates from odontogenic abscesses. *J Antimicrob Chemother* 1997; 41: 353–357.
- Ruoff KL. *Streptococcus anginosus* ('*Streptococcus milleri*') the unrecognized pathogen. *Clin Microbiol Rev* 1998; 1: 102–108.
- Rupf S, Merte K, Eschrich K. Quantification of bacteria in oral samples by

- competitive polymerase chain reaction. *J Dent Res* 1999; 78: 850–856.
40. Socransky SS, Haffajee AD. Evidence of bacterial etiology: a historical perspective. *Periodontol* 2000 1994; 5: 7–25.
 41. Song Y, Liu C, McTeague M, Vu A, Liu JY, Finegold SM. Rapid identification of Gram-positive anaerobic coccal species originally classified in the genus *Peptostreptococcus* by multiplex PCR assays using genus- and species-specific primers. *Microbiology* 2003; 149: 1719–1727.
 42. Tachikawa J. Clinico-pathological study of postoperative maxillary cysts. *Shikwa Gakuho* 1975; 75: 1117–1142.
 43. Takahashi N, Saito K, Schachtele CF, Yamada T. Acid tolerance and acid-neutralizing activity of *Porphyromonas gingivalis*. *Prevotella intermedia* and *Fusobacterium nucleatum*. *Oral Microbiol Immunol* 1997; 12: 323–328.
 44. Takahashi N, Sato T, Yamada T. Metabolic pathways for cytotoxic end product formation from glutamate- and aspartate-containing peptides by *Porphyromonas gingivalis*. *J Bacteriol* 2000; 182: 4704–4710.
 45. Travis J, Pike R, Imamura T, Potempa J. *Porphyromonas gingivalis* proteinases as virulence factors in the development of periodontitis. *J Periodontal Res* 1997; 32: 120–125.
 46. Wahlfors J, Meurman JH, Vaisanen P, Alakuijala P, Korhnen A, Torkko H, Janne J. Simultaneous detection of *Actionobacillus actinomycetemcomitans* and *Porphyromonas gingivalis* by a rapid PCR method. *J Dent Res* 1995; 74: 1796–1801.
 47. Wanahita A, Goldsmith EA, Musher DM, Clarridge JE III, Rubio J, Krishnan B, et al. Interaction between human polymorphonuclear leukocytes and *Streptococcus milleri* group bacteria. *J Infect Dis* 2002; 185: 85–90.
 48. Whiley RA, Beighton D, Winstanley TG, Fraser HY, Hardie JM. *Streptococcus intermedius*, *Streptococcus constellatus*, and *Streptococcus anginosus* (the *Streptococcus milleri* group): association with different body sites and clinical infections. *J Clin Microbiol* 1992; 30: 243–244.
 49. Wilson MJ, Weighan AJ, Wade WG. Applications of molecular ecology in the characterization of uncultured microorganism associated with human disease. *Rev Med Microbiol* 1997; 8: 91–101.

Synergistic Inhibition by Combination of Fluoride and Xylitol on Glycolysis by Mutans Streptococci and Its Biochemical Mechanism

H. Maehara^a Y. Iwami^b H. Mayanagi^a N. Takahashi^b

^aDepartment of Lifelong Oral Health Sciences, Division of Pediatric Dentistry, and ^bDepartment of Oral Biology, Division of Oral Ecology and Biochemistry, Tohoku University Graduate School of Dentistry, Sendai, Japan

Key Words

Fluoride · Glycolysis · Mutans streptococcus · Xylitol

Abstract

The purpose of this study was to evaluate the combined inhibitory effect of fluoride and xylitol on acid production by mutans streptococci, *Streptococcus mutans* NCTC10449 and *Streptococcus sobrinus* 6715, from glucose under strictly anaerobic conditions at fixed pH 5.5 and 7.0. The bacteria were grown in a tryptone-yeast extract broth under strictly anaerobic conditions (N₂: 80%; H₂: 10%; CO₂: 10%). Reaction mixtures for acid production from glucose contained bacterial cells with fluoride (0–6.4 mM) and/or xylitol (60 mM). Acidic end products of glucose fermentation and intracellular glycolytic intermediates were assayed. The combination of fluoride and xylitol inhibited acid production more effectively than fluoride or xylitol alone. In the presence of fluoride and xylitol, the proportion of lactic acid in the total amount of acidic end products decreased, while the proportion of formic and acetic acids increased. Analyses of intracellular glycolytic intermediates revealed that xylitol inhibited the upper part of the glycolytic pathway, while fluoride inhibited the lower part. This study indicates that fluoride and xylitol together have synergistic inhibitory effects on the acid production of mutans streptococci and suggests that xylitol has the potential to enhance inhibitory effects of low concentrations of fluoride.

Copyright © 2005 S. Karger AG, Basel

Both fluoride and xylitol have been widely used as cariostatic agents. Fluoride inhibits the acid production and growth of oral bacteria such as the cariogenic bacteria mutans streptococci [Kashket et al., 1977; Hamilton and Ellwood, 1978; van Houte, 1980; Okuda and Frostell, 1982; Marsh et al., 1985; Zameck and Tinanoff, 1987], particularly under anaerobic conditions [Hata et al., 1990]. The effects on streptococcal cells of fluoride are partly ascribed to the inhibition of enolase, one of the series of glycolytic enzymes [Jenkins, 1999]. This inhibition decreases the intracellular level of phosphoenolpyruvate (PEP), and thus decreases bacterial sugar uptake via PEP-dependent phosphotransferase system (PEP-PTS) [Jenkins, 1999]. In addition, fluoride can directly inhibit bacterial proton-translocating ATPase (H⁺-ATPase) that is considered to partly contribute to the excretion of proton out of the cells, leading to the acidification of intracellular pH. The dissociation of unionized hydrofluoric acid into H⁺ and F⁻ in the cells also promotes intracellular acidification [Jenkins, 1999]. This intracellular acidification can further reduce bacterial metabolic activity.

Xylitol is a valuable sugar alcohol for caries prevention [Scheinin et al., 1976; Trahan, 1995] because the majority of oral bacteria, including mutans streptococci, do not ferment it [Birkhed et al., 1985]. Xylitol is taken into the cells and phosphorylated to xylitol-5-phosphate (X5P) by PEP-PTS [Assev and Rölla, 1984; Trahan et al., 1985; Trahan, 1995]. Since X5P cannot be metabolized further [Hausman et al., 1984; Assev and Rölla, 1986], the intracellularly accumulated X5P inhibits glycolytic enzymes

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2005 S. Karger AG, Basel
0008-6568/05/0396-0521\$22.00/0

Accessible online at:
www.karger.com/crc

Nobuhiro Takahashi
Department of Oral Biology, Division of Oral Ecology and Biochemistry
Tohoku University Graduate School of Dentistry
4-1 Seiryō-machi, Aoba-ku, Sendai, 980-8575 (Japan)
Tel. +81 22 717 8294. Fax +81 22 717 8297. E-Mail nobu-t@mail.tains.tohoku.ac.jp

such as phosphoglucose isomerase (PGI) and phosphofructokinase (PFK) and subsequently represses sugar metabolism [Assev et al., 1983; Vadeboncoeur et al., 1983; Assev and Rölla, 1986; Bradshaw and Marsh, 1994]. After that, the X5P can be dephosphorylated into xylitol and excreted out of the cells [Trahan et al., 1991]. In this 'futile cycle' [Assev and Rölla, 1986], the bacterial cells are forced to waste energy and their growth is obstructed as a result [Mäkinen and Scheinin, 1976; Assev et al., 1980, 1983; Vadeboncoeur et al., 1983; Loesche, 1984; Assev and Rölla, 1986; Bradshaw and Marsh, 1994].

Since both fluoride and xylitol have inhibitory effects on sugar metabolism of mutans streptococci as stated above, using them together in clinical situations is expected to inhibit acid production of mutans streptococci effectively. Scheie et al. [1988] demonstrated the combined inhibitory effect of fluoride and xylitol on the growth of mutans streptococci and suggested the inhibitory step of glucose metabolism. Moreover, Rogers and Bert [1992] showed the possibility that xylitol enhances the inhibitory effect of fluoride on mutans streptococci growing in continuous culture. However, the biochemical mechanism of the fluoride and xylitol combination on dental plaque bacteria has not been elucidated yet.

Therefore, this study aimed to evaluate the combined inhibitory effect of fluoride and xylitol on the acid production of mutans streptococci from glucose under strictly anaerobic conditions like those of dental plaque with a pH-stat that can exactly keep the constant pH of reaction mixtures. Furthermore, we attempted to elucidate the biochemical mechanism by analyzing the profiles of intracellular glycolytic intermediates.

Materials and Methods

Bacterial Strain

Streptococcus mutans NCTC10449 and *Streptococcus sobrinus* 6715 were used in this study.

Anaerobic Procedure

All the experiments for bacterial cultivation and metabolism were conducted under strictly anaerobic conditions [Yamada and Carlsson, 1975] like those of the deep layers of mature dental plaque [Ritz, 1967; Kenney and Ash, 1969; Katayama et al., 1975], which are closely associated with causation of dental caries. The storage and cultivation of bacteria were performed in an anaerobic chamber (type NHC, gas phase: N₂, 80%; H₂, 10%; CO₂, 10%, Hirasawa Works, Tokyo, Japan). Preparation of bacterial cell suspensions and the bacterial metabolism experiments were carried out in another anaerobic chamber (type NH, gas phase: N₂, 90%; H₂, 10%, Hirasawa Works). During centrifugation outside the anaerobic chamber for harvesting and washing, the bacterial cells were pro-

ected from exposure to air by double sealed centrifuge tubes (Kubota Commercial Affairs, Tokyo, Japan) [Miyasawa et al., 2003]. To exclude oxygen, growth media and solutions were placed in the anaerobic chambers for at least 3 days prior to use, while powdered reagents and experimental instruments were placed in the anaerobic chamber for at least 1 day prior to use.

Bacterial Growth Conditions

The bacterial strains, cultured on blood agar plates and stored at 4°C in the anaerobic chamber, were inoculated into 5 ml of tryptone-yeast extract (TYE; pH 7) broth containing 1.7% tryptone (Difco, Detroit, Mich., USA), 0.3% yeast extract (Difco), 85.5 mM NaCl, 14.4 mM dipotassium phosphate and 11.1 mM glucose, and incubated anaerobically at 37°C overnight. The bacterial suspension was subcultured into another 100 ml of TYE broth (inoculum size 5%) for 12 h, and then further cultured into 800 ml of TYE broth (inoculum size 5%). Bacterial growth was monitored by optical density at 660 nm of aliquots from the culture using a spectrophotometer (model UV-160 Shimadzu, Ltd., Kyoto, Japan). The cells were harvested at the early exponential phase of growth by centrifugation (10,000 g for 10 min at 4°C). The cells were washed twice with a solution containing 150 mM KCl and 5 mM MgCl₂ (4°C) by centrifugation (13,000 g for 8 min at 4°C) and the washed cells were finally resuspended in the same solution. The cell suspension was adjusted to 2.7 mg dry weight cells per milliliter and stored at 4°C under anaerobic conditions until the following experiments.

Acid Production from Glucose in the Presence of Fluoride and/or Xylitol

The reaction mixture contained bacterial cells (0.9 mg dry weight cells), 150 mM KCl and 5 mM MgCl in 2 mM potassium phosphate buffer (pH 5.5 or 7.0). Potassium fluoride (0–6.4 mM), xylitol (60 mM) or both were added into the reaction mixtures. The control reaction mixture contained no fluoride or xylitol. The reaction mixture was maintained at pH 5.5 or 7.0 by titration with 60 mM KOH using a pH stat (AUTO pH STAT; model AUT-211S, TOA Electronics, Tokyo, Japan) with agitation by a magnetic stirrer, and then preincubated at 35°C for 4 min. The reaction was started by the addition of 10 mM glucose.


Analysis of Carboxylic Acids from Glucose Fermentation

Lactic, formic and acetic acids were assayed following the method of Takahashi et al. [1987]. At 10 min after the addition of glucose, aliquots of the reaction mixture of *S. mutans* NCTC 10449 were sampled and immediately mixed with 0.6 N perchloric acid. The resultant mixture was brought out from the anaerobic chamber, filtered to remove cell debris through a membrane filter (pore size 0.20 µm; Advantec, polypropylene, Toyo Roshi Ltd., Tokyo, Japan), diluted with 0.2 N HCl and assayed with a carboxylic acid analyzer (model Eyela S-3000, Tokyo Rika Co. Ltd., Tokyo, Japan).

Determination of Glycolytic Intermediates

Glycolytic intermediates were extracted from cells following the method of Iwami et al. [2001]. The cells of *S. mutans* NCTC 10449 were collected by filtering 3.0 ml of reaction mixture through a membrane filter (pore size 0.45 µm; Acrodisc, polyethersulfone, Pall Gelman Laboratory, Ann Arbor, Mich., USA) at 0, 2, 4 and 6 min after the addition of glucose. The glycolytic intermediates

Fig. 1. Effect of fluoride on the acid production for 10 min from 10 mM glucose in the presence and absence of 60 mM xylitol by the cells of *S. mutans* NCTC 10449 and *S. sobrinus* 6715 at pH 7.0 and 5.5. F = Fluoride alone; F + X = fluoride plus xylitol. The data are the means of three independent experiments. Vertical bars indicate standard deviations from three independent experiments.



were immediately extracted by passing 2.0 ml of 0.6 *N* perchloric acid through the filter. The filter was filtered twice besides with the same perchloric acid solution to extract glycolytic intermediates fully. The extracts were neutralized to pH of 6.0 using 1 *M* triethanolamine-HCl and 5 *M* K₂CO₃ with chilling at 4°C, and filtered through another membrane filter (pore size 0.20 μm; Advantec, polypropylene, Toyo Roshi Ltd., Tokyo, Japan) to remove sediment. The neutralized extracts were stored at 4°C until assayed.

Glycolytic intermediates in the cell extracts were determined at 35°C by the fluorometric methods of Maitra and Estabrook [1964] and Takahashi et al. [1991] with some modifications. A fluorescence spectrophotometer (model CAF-110, Jasco Corp., Tokyo, Japan) was used with a slit width of 10 nm and excitation/emission wavelengths of 340/470 nm. The assay mixture for glucose 6-phosphate (G6P) and fructose 6-phosphate (F6P) contained 20 mM triethanolamine hydrochloride-NaOH buffer (TA buffer, pH 7.5), 0.06 mM NADP and the cell extract. The amounts of G6P were determined from the increase of NADPH after the addition of 0.6 U/ml glucose 6-phosphate dehydrogenase (EC 1.1.1.49). After NADPH increase reached a plateau, 3.3 U/ml phosphoglucose isomerase (EC 5.3.1.9) was added, and the amount of F6P was then determined from the subsequent increase of NADPH. The assay mixture for dihydroxyacetone phosphate (DHAP), glyceraldehyde 3-phosphate (G3P) and fructose 1,6-bisphosphate (FBP) contained 20 mM TA buffer, 0.5 mM ethylenediamine tetraacetate, 7.5 mM NADH and the cell extract. Glycerol 3-phosphate dehydrogenase (EC 1.1.1.8; 0.85 U/ml), triose phosphate isomerase (EC 5.3.1.1; 25 U/ml) and aldolase (EC 4.1.2.13; 0.23 U/ml) were added sequentially to determine the amounts of DHAP, G3P and FBP from the respective decrease of NADH. The assay mixture for 3-phospho-

glycerate (3PG) and 2-phosphoglycerate (2PG) contained 20 mM TA buffer, 2 mM cysteine, 1.5 mM MgCl₂, 1.5 mM ATP, 0.01 mM NADH and the cell extract. Glyceraldehyde 3-phosphate dehydrogenase (EC 1.2.1.12; 2.5 U/ml), 3-phosphoglycerate kinase (EC 2.7.2.3; 8.4 U/ml) and phosphoglycerate mutase (EC 5.4.2.1; 4.5 U/ml) were added sequentially to determine the amounts of 3PG and 2PG from the respective decrease of NADH. The assay mixture for PEP and pyruvate (PYR) contained 20 mM TA buffer, 10 mM KCl, 5 mM MgCl₂, 0.5 mM ADP, 5 μM NADH and the cell extract. Lactate dehydrogenase (EC 2.7.1.40; 3.5 U/ml) and pyruvate kinase (EC 1.1.1.27; 2.4 U/ml) were added sequentially to determine the amounts of PYR and PEP from the respective decrease of NADH.

Statistical Analysis

All numerical data were given as means with standard deviations. Statistical significance was assessed using ANOVA and Dunnett test. Differences were considered significant at the level *p* < 0.05.

Results

Acid Production from Glucose in the Presence of Fluoride and/or Xylitol

In the presence of fluoride, the acid production by *S. mutans* NCTC10449 from glucose for 10 min decreased at both pH 7.0 and 5.5. The decrease of acid production




Fig. 2. Effect of fluoride and/or xylitol on the acid production curves of the cells of *S. mutans* NCTC 10449 and *S. sobrinus* 6715 after addition of 10 mM glucose at pH 7.0. F = 1.6 and 2.4 mM fluoride alone for *S. mutans* and *S. sobrinus*, respectively; X = 60 mM xylitol alone; F + X = fluoride plus xylitol. The data represent three independent experiments. The results were similar among three independent experiments.


was dose-dependent (fig. 1). At pH 7.0, 2.4 mM fluoride was required to reduce acid production by 50% or more, while only 0.35 mM fluoride had a similar effect at pH 5.5. Similar results were obtained in *S. sobrinus* 6715 (fig. 1).

The inhibition of acid production by fluoride was enhanced in the presence of xylitol. At pH 7.0, the acid production by *S. mutans* decreased by 21.7% in the presence of 1.6 mM fluoride alone and by 8.9% in the presence of 60 mM xylitol alone, while the acid production was decreased by 58.4% in the presence of a combination of 1.6 mM fluoride and 60 mM xylitol. The enhanced inhibition caused by the addition of xylitol was also observed at pH 5.5, with 16.3% inhibition in the presence of

0.3 mM fluoride alone and 80.1% inhibition by the combination of 0.3 mM fluoride plus 60 mM xylitol. A similar inhibition by the combination of fluoride and xylitol was observed in *S. sobrinus* 6715.

In the absence of fluoride or xylitol, the acid production by *S. mutans* NCTC 10449 started immediately after the addition of glucose and gradually slowed down (fig. 2, control). In the presence of xylitol alone (fig. 2, X), no acid production was detected for about 1 min after the addition of glucose. Once it started, the acid production pattern was similar to the control, though the acid production rate was slower. In the presence of fluoride (fig. 2, F) and the combination of fluoride and xylitol (fig. 2, F + X), the acid production was very low right after the addition

Fig. 3. Effect of fluoride and/or xylitol on the production of acidic end products from 10 mM glucose for 10 min by the cells of *S. mutans* NCTC 10449 at pH 7.0. Left = Amounts of acidic end products; right = profiles of acidic end products. F = 1.6 mM fluoride alone; X = 60 mM xylitol alone; F + X = fluoride plus xylitol. The data are the means of three independent experiments. Error bars indicate standard deviations from three independent experiments. * Significant difference from control ($p < 0.05$).



of glucose and increased gradually and became nearly constant by 10 min. Similar inhibitory effects of fluoride and xylitol were observed in *S. sobrinus* 6715.

Production of Carboxylic Acids from Glucose in the Presence of Fluoride and/or Xylitol

The effects of fluoride and xylitol on the formation of acidic end products from glucose and the profile of intracellular glycolytic intermediates were evaluated at a fluoride concentration of 1.6 mM and pH 7.0 using *S. mutans* NCTC 10449, where the combined effect of fluoride and xylitol was evident. The total amount of carboxylic acids (sum of lactic, formic and acetic acids) produced by *S. mutans* NCTC10449 at pH 7.0 for 10 min was 4.49 nmol/mg of cells in the control, while those in the presence of fluoride alone and xylitol alone were 3.75 and 4.27 nmol/mg of cells, respectively. In the presence of fluoride and xylitol combined, the total amount decreased to 2.23 nmol/mg of cells (fig. 3). These results showed that the combination of fluoride and xylitol was more effective in inhibiting the acid production than fluoride or xylitol alone, similar to the results of acid production rate obtained by pH stat (fig. 1). In addition, lactic acid as a proportion of the total amount of carboxylic acids decreased, while the proportion of formic and acetic acids increased in the presence of xylitol alone. A similar result was observed in the presence of a combination of fluoride and xylitol (fig. 3, right).

Profile of Intracellular Glycolytic Intermediates in the Presence of Fluoride and/or Xylitol

In the presence of fluoride alone, the levels of FBP and PYR were lower and the levels of 3PG and 2PG were

higher than those in the control during glucose fermentation (fig. 4). In contrast, in the presence of xylitol alone, the levels of glycolytic intermediates except PYR were mostly lower. Moreover, in the presence of the combination of fluoride and xylitol, the levels of G6P through G3P were similar to those in the presence of xylitol alone, while the levels of 3PG through PYR were similar to those in the presence of fluoride alone (fig. 4).

At 0 min (just before glucose addition), the control cells had low levels of glycolytic intermediates, except 3PG, 2PG and PEP (the intermediates which comprise the 'PEP potential' for sugar uptake into cells [Thompson, 1979]). The levels of these 'PEP potential' intermediates (3PG, 2PG and PEP) at 0 min in the presence of fluoride alone were twice as high as those of the control, while the levels of those were negligible in the presence of xylitol alone. The level of 'PEP potential' in the combination of fluoride and xylitol was close to the control at 0 min, though PEP was not detected (fig. 4).

Discussion

Acid production from glucose by mutans streptococci was inhibited in the presence of fluoride or xylitol as previously reported [Kashket et al., 1977; Hamilton and Ellwood, 1978; Assev et al., 1983; Vadeboncoeur et al., 1983; Marsh et al., 1985; Bradshaw and Marsh, 1994]. In addition, this study has demonstrated that the inhibitory effect was enhanced by a combination of fluoride and xylitol. This is consistent with previous results for combined effects of fluoride and xylitol on the growth of mutans streptococci [Scheie et al., 1988; Rogers and Bert, 1992].




Fig. 4. Effect of fluoride and/or xylitol on the levels of glycolytic intermediates of the cells of *S. mutans* NCTC 10449 during glucose fermentation. The levels of glycolytic intermediates at 0, 2, 4, and 6 min after addition of 10 mM glucose to the cells *S. mutans* NCTC 10449 in the presence of 1.6 mM fluoride and/or 60 mM xylitol. F = 1.6 mM fluoride alone; X = 60 mM xylitol alone; F + X = fluoride plus xylitol. Error bars indicate standard deviations from three independent experiments. Significant difference from control (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$). # = Not detected. The data are the means of three independent experiments.

Moreover, percent inhibition by combination of fluoride and xylitol was larger than the sum of percent inhibition by fluoride alone and percent inhibition by xylitol alone, indicating that the combined effect was synergistic (fig. 1, 2).

The observation that fluoride did not change the profile of end products while xylitol decreased the proportion of lactic acid implies a difference in inhibitory mechanism between xylitol and fluoride (fig. 3). This finding led us to the analysis of glycolytic intermediates to understand the biochemical mechanism for the synergistic effects of xylitol and fluoride.

The presence of fluoride alone increased the level of 2PG and 3PG in the lower part of glycolysis as reported previously [Hata et al., 1990]. This indicates that fluoride inhibited enolase (a glycolytic enzyme catalyzing the conversion of 2PG to PEP) and resulted in the accumulation of 3PG and 2PG and the decrease of PEP (fig. 4). In bacterial cells, PEP is a phosphoryl donor for sugar uptake by PEP-PTS [Schachtele and Mayo, 1973; Dills et al., 1980] and builds 'PEP potential' together with 3PG and 2PG, precursors of PEP. Although 3PG and 2PG levels were high in the presence of fluoride, the slow supply of PEP from 3PG and 2PG caused by the fluoride inhibition could decrease glucose uptake via PEP-PTS and slow

down the entire glycolytic process [Kanapka and Hamilton, 1971; Bender et al., 1985] (fig. 2, 4).

In the presence of xylitol alone, in contrast, most glycolytic intermediates decreased as acid production was inhibited, suggesting the repression of sugar uptake by xylitol. In addition, the marked decrease of FBP as previously reported [Miyasawa et al., 2003] may be due to the inhibition by X5P of PGI and PFK catalyzing the conversion of G6P to FBP [Trahan, 1995]. Since FBP is an activator of lactate dehydrogenase, the decrease in FBP caused inactivation of lactate dehydrogenase and consequent reduction of lactic acid production (fig. 3) as previously reported [Miyasawa et al., 2003].

The combination of fluoride and xylitol increased the intracellular levels of 3PG and 2PG as observed in the presence of fluoride alone, while the other glycolytic intermediates decreased similarly to the presence of xylitol alone. These observations indicate that the simultaneous inhibition, e.g. the inhibition of enolase by fluoride and the inhibition of PGI, PFK and glucose uptake by xylitol, repressed the entire glycolytic pathway more efficiently.

These inhibitory effects seemed more efficient in the initial stage of acid production. During preincubation in the presence of fluoride alone, PEP could not be consumed and 3PG and 2PG were accumulated through flu-

oride inhibition of enolase (fig. 4). After addition of glucose, the 'PEP potential' could be used for glucose uptake, however, the enolase inhibition by fluoride continued to interfere with the conversion of 2PG into PEP and consequently slowed down the entire glycolytic flow (fig. 2). In the presence of xylitol alone, the level of 'PEP potential' was low due to the consumption of PEP (fig. 4) through the phosphorylation of xylitol to X5P by PEP-PTS [Assev and Rölla, 1984; Trahan et al., 1985; Trahan, 1995; Miyasawa et al., 2003]. As a result, the lack of 'PEP potential' delayed acid production from glucose for about 1 min after glucose addition (fig. 2). The combination of fluoride and xylitol was more effective. The combination caused both enolase inhibition and low 'PEP potential' (fig. 4), resulting in the prolonged delay of initiation of acid production (fig. 2).

The inhibitory effect of fluoride on dental plaque acidogenicity in vivo is still unclear since there are numerous clinical observations but very few appropriate methods for evaluating antimicrobial activity of fluoride in vivo [van Loveren, 2001]. Reduction of plaque acidogenicity after sucrose challenge has been reported in plaque from subjects who used fluoridated water or rinsed their mouths daily with 48 mM (912 ppm) sodium fluoride [Edgar et al., 1970; Geddes and McNee, 1982]. Moreover, Neff [1967] reported that when applied simultaneously with sucrose, 0.52 mM (10 ppm) sodium fluoride was sufficient to inhibit acid production of dental plaque. However, Giertsen et al. [1999] argued that mouth rinsing with 12 mM (228 ppm) sodium fluoride had no significant effect on the acidogenic potential of dental plaque. In experiments using a continuous culture of mixed oral bac-

teria [Rogers et al., 1991; Bradshaw and Marsh, 1994] and an artificial biofilm [Bradshaw et al., 2002] that mimic some aspects of the dental plaque micro-ecosystem, fluoride and xylitol as well as a fluoride-xylitol combination were able to reduce the acid production of the micro-ecosystem with a reduction of the number of mutans streptococci. These sophisticated studies support the possibility that the combination of fluoride and xylitol could reduce dental plaque acidogenicity in vivo, though a proper method for clinical application of fluoride and xylitol should be developed as soon as possible.

In conclusion, direct inhibition of glycolytic enzymes, enolase, PGI and PFK and the consequent inhibition on the supply of 'PEP potential' for sugar uptake appears to be the cause of the synergistic inhibition by combination of fluoride and xylitol on glycolysis by mutans streptococci. Attempts have been made to use both fluoride and xylitol together to enhance remineralization of tooth surfaces [Smits and Arends, 1985; Arends et al., 1990; Gaffar et al., 1998; Amaechi et al., 1998]. Besides remineralization, this study suggests that using fluoride and xylitol together can effectively inhibit the sugar metabolism of mutans streptococci and also that xylitol has the potential to enhance the inhibitory effects of low concentrations of fluoride.

Acknowledgments

This work was supported by Grants-in-Aid for Science Research (B) No. 13470446, No. 14370687 and No. 16390601 from the Ministry of Education, Science, Sports and Culture, Japan.

References

- ▶ Amaechi BT, Higham SM, Edgar WM: The influence of xylitol and fluoride on dental erosion in vitro. *Arch Oral Biol* 1998;43:157-161.
- ▶ Arends J, Smits M, Ruben JL, Christoffersen J: Combined effect of xylitol and fluoride on enamel demineralization in vitro. *Caries Res* 1990;24:256-257.
- Assev S, Rölla G: Evidence for presence of a xylitol phosphotransferase system in *Streptococcus mutans* OMZ 176. *Acta Pathol Microbiol Immunol Scand [B]* 1984;92:89-92.
- Assev S, Rölla G: Further studies on the growth inhibition of *Streptococcus mutans* OMZ 176 by xylitol. *Acta Pathol Microbiol Immunol Scand [B]* 1986;94:97-102.
- Assev S, Vegarud G, Rölla G: Growth inhibition of *Streptococcus mutans* strain OMZ 176 by xylitol. *Acta Pathol Microbiol Scand [B]* 1980; 88:61-63.
- Assev S, Wåler SM, Rölla G: Further studies on the growth inhibition of some oral bacteria by xylitol. *Acta Pathol Microbiol Immunol Scand [B]* 1983;91:261-265.
- ▶ Bender GR, Thibodeau EA, Marquis RE: Reduction of aciduranc of streptococcal growth and glycolysis by fluoride and gramicidin. *J Dent Res* 1985;64:90-95.
- ▶ Birkhed D, Kalfas S, Svensäter G, Edwardsson S: Microbiological aspects of some caloric sugar substitutes. *Int Dent J* 1985;35:9-17.
- ▶ Bradshaw DJ, Marsh PD: Effect of sugar alcohols on the composition and metabolism of a mixed culture of oral bacteria grown in a chemostat. *Caries Res* 1994;28:251-256.
- ▶ Bradshaw DJ, Marsh PD, Hodgson RJ, Visser JM: Effects of glucose and fluoride on competition and metabolism within in vitro dental bacterial communities and biofilms. *Caries Res* 2002;36:81-86.
- ▶ Dills SS, Apperson A, Schmidt MR, Saier MH: Carbohydrate transport in bacteria. *Microbiol Rev* 1980;44:385-418.
- ▶ Edgar WM, Jenkins GN, Tatevossian A: The inhibitory action of fluoride on plaque bacteria: Further evidence. *Br Dent J* 1970;128:129-132.
- ▶ Gaffar A, Blake-Haskins JC, Sullivan R, Simone A, Schmidt R, Saunders F: Cariostatic effects of a xylitol/NaF dentifrice in vivo. *Int Dent J* 1998;48:32-39.

- Geddes DA, McNee SG: The effect of 0.2 per cent (48 mM) NaF rinses daily on human plaque acidogenicity in situ (Stephan curve) and fluoride content. *Arch Oral Biol* 1982;27:765-769.
- Giertsen E, Emberland H, Scheie AA: Effects of mouth rinses with xylitol and fluoride on dental plaque and saliva. *Caries Res* 1999;33:23-31.
- Hamilton IR, Ellwood DC: Effects of fluoride on carbohydrate metabolism by washed cells of *Streptococcus mutans* grown at various pH values in a chemostat. *Infect Immun* 1978;19:434-442.
- Hata S, Iwami Y, Kamiyama K, Yamada T: Biochemical mechanisms of enhanced inhibition of fluoride on the anaerobic sugar metabolism by *Streptococcus sanguis*. *J Dent Res* 1990;69:1244-1247.
- Hausman SZ, Thompson J, London J: Futile xylitol cycle in *Lactobacillus casei*. *J Bacteriol* 1984;160:211-215.
- van Houte J: Bacterial specificity in the etiology of dental caries. *Int Dent J* 1980;30:305-326.
- Iwami Y, Takahashi-Abbe S, Takahashi N, Yamada T, Kano N, Mayanagi H: The time-course of acid excretion, levels of fluorescence dependent on cellular nicotinamide adenine nucleotide and glycolytic intermediates of *Streptococcus mutans* cells exposed and not exposed to air in the presence of glucose and sorbitol. *Oral Microbiol Immunol* 2001;16:34-39.
- Jenkins GN: Review of fluoride research since 1959. *Arch Oral Biol* 1999;44:985-992.
- Kanapka JA, Hamilton IR: Fluoride inhibition of enolase activity in vivo and its relationship to the inhibition of glucose-6-P formation in *Streptococcus salivarius*. *Arch Biochem Biophys* 1971;146:167-174.
- Kashket S, Rodriguez VM, Bunick FJ: Inhibition of glucose utilization in oral streptococci by low concentrations of fluoride. *Caries Res* 1977;11:301-307.
- Katayama T, Suzuki T, Okada S: Clinical observation of dental plaque maturation: Application of oxidation-reduction indicator dyes. *J Periodontol* 1975;46:610-613.
- Kenney EB, Ash MM: Oxidation reduction potential of developing plaque, periodontal pockets and gingival sulci. *J Periodontol* 1969;40:630-633.
- Loesche WJ: The effect of sugar alcohols on plaque and saliva level of *Streptococcus mutans*. *Swed Dent J* 1984;8:125-135.
- van Loveren C: Antimicrobial activity of fluoride and its in vivo importance: Identification of research questions. *Caries Res* 2001;35(suppl 1):65-70.
- Maitra PK, Estabrook RW: A fluorometric method for the enzymic determination of glycolytic intermediates. *Anal Biochem* 1964;41:472-484.
- Mäkinen KK, Scheinin A: Turku sugar studies. VII. Principal biochemical findings on whole saliva and plaque. *Acta Odontol Scand* 1976;34:241-283.
- Marsh PD, McDermid AS, Keevil CW, Ellwood DC: Effect of environmental conditions on the fluoride sensitivity of acid production by *S. sanguis* NCTC 7865. *J Dent Res* 1985;64:85-89.
- Miyasawa H, Iwami Y, Mayanagi H, Takahashi N: Xylitol inhibition of anaerobic acid production by *Streptococcus mutans* at various pH levels. *Oral Microbiol Immunol* 2003;18:215-219.
- Neff D: Acid production from different carbohydrate sources in human plaque in situ. *Caries Res* 1967;1:78-87.
- Okuda K, Frostell G: The effect of fluoride on the acid production of *Streptococcus mutans* and other oral streptococci. *Swed Dent J* 1982;6:29-36.
- Ritz HL: Microbial population shifts in developing human dental plaque. *Arch Oral Biol* 1967;12:1561-1568.
- Rogers AH, Bert AG: Effects of xylitol and fluoride on the response to glucose pulses of *Streptococcus mutans* T8 growing in continuous culture. *Oral Microbiol Immunol* 1992;7:124-126.
- Rogers AH, Pilowsky KA, Zilm PS, Gully NJ: Effects of pulsing with xylitol on mixed continuous cultures of oral streptococci. *Aust Dent J* 1991;36:231-235.
- Schachtele CF, Mayo JA: Phosphoenolpyruvate-dependent glucose transport in oral streptococci. *J Dent Res* 1973;52:1209-1215.
- Scheie AA, Assev S, Rölla G: Combined effect of xylitol, NaF and ZnCl₂ on growth and metabolism of *Streptococcus sobrinus* OMZ 176. *APMIS* 1988;96:761-767.
- Scheinin A, Mäkinen KK, Ylitalo K: Turku sugar studies. V. Final report on the effect of sucrose, fructose and xylitol diets on the caries incidence in man. *Acta Odontol Scand* 1976;34:179-216.
- Smits MT, Arends J: Influence of xylitol- and/or fluoride-containing toothpastes on the remineralization of surface softened enamel defects in vivo. *Caries Res* 1985;19:528-535.
- Takahashi N, Abbe K, Takahashi-Abbe S, Yamada T: Oxygen sensitivity of sugar metabolism and interconversion of pyruvate formate-lyase in intact cells of *Streptococcus mutans* and *Streptococcus sanguis*. *Infect Immun* 1987;55:652-656.
- Takahashi N, Iwami Y, Yamada T: Metabolism of intracellular polysaccharide in the cells of *Streptococcus mutans* under strictly anaerobic conditions. *Oral Microbiol Immunol* 1991;6:299-304.
- Thompson J: Lactose metabolism in *Streptococcus lactis*: Phosphorylation of galactose and glucose moieties in vivo. *J Bacteriol* 1979;140:774-785.
- Trahan L: Xylitol: A review of its action on mutans streptococci and dental plaque - its clinical significance. *Int Dent J* 1995;45:77-92.
- Trahan L, Bareil M, Gauthier L, Vadeboncoeur C: Transport and phosphorylation of xylitol by a fructose phosphotransferase system in *Streptococcus mutans*. *Caries Res* 1985;19:53-63.
- Trahan L, Néron S, Bareil M: Intracellular xylitol-phosphate hydrolysis and efflux of xylitol in *Streptococcus sobrinus*. *Oral Microbiol Immunol* 1991;6:41-50.
- Vadeboncoeur C, Trahan L, Mouton C, Mayrand D: Effect of xylitol on the growth and glycolysis of acidogenic oral bacteria. *J Dent Res* 1983;62:882-884.
- Yamada T, Carlsson J: Regulation of lactate dehydrogenase and change of fermentation products in streptococci. *J Bacteriol* 1975;124:55-61.
- Zameck RL, Tinanoff N: Effects of NaF and SnF₂ on growth, acid and glucan production of several oral streptococci. *Arch Oral Biol* 1987;32:807-810.

Difference in the xylitol sensitivity of acid production among *Streptococcus mutans* strains and the biochemical mechanism

H. Miyasawa-Hori¹, S. Aizawa²,
N. Takahashi¹

¹Division of Oral Ecology and Biochemistry, Department of Oral Biology and ²Division of Pediatric Dentistry, Department of Lifelong Oral Health Sciences, Tohoku University Graduate School of Dentistry, Sendai, Japan

Miyasawa-Hori H, Aizawa S, Takahashi N. Difference in the xylitol sensitivity of acid production among *Streptococcus mutans* strains and the biochemical mechanism. *Oral Microbiol Immunol* 2006; 21: 1–6. © Blackwell Munksgaard, 2006.

Xylitol inhibits the glycolysis and growth of *Streptococcus mutans*, but to different degrees among strains. Thus, we studied the biochemical mechanism through which the inhibition varies, using *S. mutans* strains ATCC 31989, NCTN 10449, and NCIB 11723, which are highly sensitive, moderately sensitive, and resistant to xylitol, respectively, under strictly anaerobic conditions such as those found in deep layers of dental plaque. Xylitol (30 mM) decreased the rate of acid production from glucose (10 mM) in ATCC 31989, NCTC 10449, and NCIB 11723 by 86, 26, and 0%, respectively. The activities of the xylitol : phosphoenolpyruvate phosphotransferase system (PEP-PTS) relative to those of glucose : PEP-PTS were 120, 16, and 3%, respectively. These findings indicated that the higher the xylitol : PEP-PTS activity was and the more effectively xylitol decreased glucose : PEP-PTS activity, the more sensitive the strain was to xylitol. In ATCC 31989 and NCTC 10449, intracellular accumulation of xylitol 5-phosphate and decreases of fructose 1,6-bisphosphate and glucose 6-phosphate were observed. Furthermore, in the presence of xylitol (30 mM), glucose : PEP-PTS activities decreased by 34, 17, and 0%, respectively. These results suggest that the following inhibitory mechanisms are active in the xylitol-sensitive *mutans* streptococci: direct inhibition of glycolytic enzymes by xylitol 5-phosphate derived from xylitol : PEP-PTS and, possibly, indirect inhibition through competition for the phosphoryl donor, HPr-P, between glucose and xylitol : PEP-PTS.

Key words: acid production; phosphoenolpyruvate-sugar phosphotransferase system; *Streptococcus mutans*; xylitol



Nobuhiro Takahashi, Division of Oral Ecology and Biochemistry, Department of Oral Biology, Tohoku University Graduate School of Dentistry, 4–1 Seiryomachi, Aoba-ku, Sendai, 980–8575, Japan
Tel.: +81 22 7178294;
fax: +81 22 7178297;
e-mail: nobu-t@mail.tains.tohoku.ac.jp
Accepted for publication October 26, 2005

Xylitol is widely used as a noncariogenic sugar substitute because it is not fermented by oral bacteria (6). Xylitol has been reported to inhibit the growth of *mutans* streptococci in the presence of glucose, galactose, mannose, lactose, maltose, sucrose, sorbitol or mannitol as a carbon source *in vitro* (1, 5, 7, 8, 16, 24, 25), and the acid production from glucose by resting cells of *S. mutans* (7, 13, 25). Xylitol is also known to selectively inhibit the growth of *S. mutans* in mixed culture using a chemostat (3, 15).

The major route of sugar transport by microorganisms is via the phosphoenolpyruvate phosphotransferase system (PEP-PTS). Two sugar-nonspecific proteins, enzyme-I and histidine-containing phosphocarrier protein (HPr), and a sugar-specific protein, enzyme-II are required for PEP-PTSs. PEP phosphorylates enzyme-I to phospho-enzyme-I, which in turn transfers the phosphoryl group to HPr. In many cases, phospho-HPr (HPr-P) generated from phospho-enzyme-I, transfers the phosphoryl group directly to enzyme-

II, which in turn phosphorylates incoming sugar (10, 14).

Bacterial cells are thought to incorporate xylitol as xylitol 5-phosphate through xylitol : PEP-PTS and the xylitol 5-phosphate inhibits the enzyme activity of sugar metabolism, resulting in the inhibition of both bacterial growth and acid production (18). In addition, the futile cycle, in which xylitol 5-phosphate is dephosphorylated to xylitol with waste of PEP potential, can also retard the growth of *S. mutans* (18).

	Journal Name	OMI	273		Dispatch: 3.1.06	Journal: OMI	CE: Blackwell
	Manuscript No.				Author Received:	No. of pages: 6	PE: Sankar

However, some strains are xylitol sensitive whereas others are resistant (19), and the degree of inhibition varies among strains (25). Thus, we studied the biochemical mechanism of the variable inhibition, using three strains of *S. mutans* previously characterized as highly sensitive, moderately sensitive, and resistant to xylitol under strictly anaerobic conditions such as those found in deep layers of dental plaque.

Material and methods

Bacterial strains and growth conditions

We used the following strains of *S. mutans*: NCTC 10449, ATCC 31989, and NCIB 11723. *S. mutans* NCTC 10449 was a gift as a xylitol-sensitive strain from Prof. L. Trahan (Universite Laval, Québec, Canada) (20). Each strain was inoculated into a complex medium containing 1.7% tryptone (Difco Laboratories, Detroit, MI), 0.3% yeast extract (Difco), 85 mM NaCl, and 11 mM glucose as described (25) under strictly anaerobic conditions in an anaerobic chamber (N₂, 80%; H₂, 10%; CO₂, 10%, NHC-type, Hirasawa Works, Tokyo, Japan) and incubated at 35°C overnight. Cell cultures were transferred into the same complex medium and precultured overnight at 35°C. The cell cultures were again transferred into the same complex medium (5% inoculum size) and grown at 35°C. The bacterial cells were harvested by centrifugation (7000 × g for 15 min at 4°C) at an early logarithmic phase of growth (optical density at 660 nm [OD₆₆₀] ≈ 0.3) under anaerobic conditions as described previously (17). Bacterial purity was regularly confirmed by culturing on blood agar plates.

Acid production from glucose in the presence of xylitol

The following experiments were conducted in a different type of anaerobic chamber (N₂, 90%; H₂, 10%, NH-type, Hirasawa Works). Cells were washed twice with cold 2 mM potassium phosphate buffer (pH 7.0) containing 150 mM KCl and 5 mM MgCl₂, and suspended in the same buffer. The optical density of the cell suspension at 660 nm was adjusted to 3.5 (1.9 mg of cells [dry weight] per ml).

The cell suspensions were agitated with a magnetic stirrer at 35°C. The reaction was started by adding a mixture of 10 mM glucose and 0 or 30 mM xylitol to the cell suspensions. The rate of acid production

by the cells was monitored at pH 7.0 using an automatic pH titrator (model AUT-211S, Toa Electronics Ltd, Kobe, Japan) with 50 mM KOH. The rate of acid production at 2 min after adding glucose or the glucose-xylitol mixture was calculated as μmol of protons per min per mg dry weight of cells.

Before and after the incubation for 10 min, cell suspensions (0.9 ml) were sampled and mixed immediately with 0.1 ml of 6 N perchloric acid. The mixtures were filtered (pore size 0.20 μm, polypropylene; ADVANTEC, Toyo Roshi Ltd, Tokyo, Japan) and cell-free filtrates were diluted with 0.2 N hydrochloric acid and stored at 4°C for the assay of acidic end products.

Analysis of acidic end products

Amounts of acidic end products, lactic, acetic, formic and pyruvic acids were quantified using a carboxylic acid analyzer (model EYELA S-3000X; Tokyo Rikakikai Co., Ltd, Tokyo, Japan) in stored cell-free filtrates, as described previously (17).

PEP-PTS activities for glucose, xylitol and fructose (glucose, xylitol: and fructose: PEP-PTS activities)

The PEP-PTS activities were estimated by a modification of the method of Kornberg & Reeves (9) as described previously (13). Cells were harvested, washed twice as described above and stored at -20°C. After thawing, the cells were suspended in 2 mM potassium phosphate buffer (pH 7.0) containing 150 mM KCl and 5 mM MgCl₂ (OD₆₆₀ = 5.0). Toluene was added at a final concentration of 1% to the cell suspension, and mixed vigorously for 1 min. After centrifugation (1200 × g for 3 min), the cells were suspended in the same buffer (OD₆₆₀ ≈ 50). The PEP-PTS activities for glucose, xylitol or fructose at pH 7.0 were estimated as a decrease of reduced nicotinamide adenine dinucleotide (NADH) in reaction mixtures containing 0.1 mM NADH, 53 μg of cells (dry weight)/ml, 1 mM phosphoenolpyruvate, 11 U/ml lactate dehydrogenase (EC 1.1.1.27, rabbit muscle; Roche Diagnostics, Indianapolis, IN), 100 mM potassium phosphate buffer (pH 7.0) at 35°C. The reaction was started by adding 5, 10, 30, 60 or 120 mM glucose, xylitol or fructose. The decrease of NADH was monitored using a dual wavelength spectrophotometer (model 557; Hitachi Ltd, Tokyo, Japan) at 340 nm.

Inhibition of glucose : PEP-PTS activity in the presence of xylitol

Glucose : PEP-PTS activity in the presence of xylitol was also determined. Toluene-treated cells as described above were suspended in a reaction mixture containing 1 mM NADP, 53 μg of cells [dry weight]/ml, 1 mM phosphoenolpyruvate, 3.5 U/ml glucose 6-phosphate dehydrogenase (EC 1.1.1.49, yeast; Roche Diagnostics) and 100 mM potassium phosphate buffer (pH 7.0) at 35°C. The reaction was started by adding 10 mM glucose and 0, 10, 30, 60 or 120 mM xylitol to the cell suspensions. The increase of NADPH was monitored spectrophotometrically at 340 nm.

Assays of glycolytic intermediates and xylitol 5-phosphate

Cell suspensions were reacted with 10 mM glucose containing 0 or 30 mM xylitol as described above for the experiment of acid production. After the incubation for 2 min, the cells were collected by passing the reaction mixture through a membrane filter (pore size 0.45 μm, polyethersulfone; Acrodisc, Pall Gelman Laboratory, Ann Arbor, MI). Glycolytic intermediates and xylitol 5-phosphate in the cells were immediately extracted in 0.6 N perchloric acid, and neutralized with 5 M K₂CO₃ in air. The neutralized extracts were stored at 4°C for subsequent assays of the glycolytic intermediates and xylitol 5-phosphate, and at -20°C for 3-phosphoglycerate assays.

The glycolytic intermediates in the cell extracts were enzymatically determined at 35°C by a modification of the enzymatic method of Minakami et al. (12). Xylitol 5-phosphate was estimated as described previously (13). The assay mixture for xylitol 5-phosphate contained 1.1 mM NAD, 5 mM MgCl₂, 0.1 mM EDTA, and the extracts in 50 mM Tris-HCl buffer (pH 8.5) at 35°C. The reaction was started by the addition of 4.2 U/ml polyol dehydrogenase (EC 1.1.1.14, sorbitol dehydrogenase, sheep liver; Roche Diagnostics) and 50 U/ml alkaline phosphatase (EC 3.1.3.1, calf intestine; Roche Diagnostics). The increase of NADH was monitored spectrophotometrically at 340 nm.

Statistical methods

Differences in rates of acid production, rates of glucose : PEP-PTS activities with xylitol and in profiles of glycolytic intermediates were analyzed by the Mann-Whitney *U*-test. Differences in amounts of

Table 1. Relative rate of acid production and the formation of acidic end products from 10 mM glucose (G) and 10 mM glucose plus 30 mM xylitol (G +X).

<i>S. mutans</i> strain	Substrate	Relative rate of acid production	Acidic end products		
			Lactate	Acetate	Formate
ATCC 31989	G	100 ^a	1.44 ± 0.28 ^b	0.34 ± 0.09	0.34 ± 0.06
	G +X	14 ± 1 [#]	0.07 ± 0.01*	0.25 ± 0.04	0.27 ± 0.04
NCTC 10449	G	100	1.07 ± 0.07	0.92 ± 0.04	0.90 ± 0.05
	G +X	74 ± 9 [#]	0.42 ± 0.03	1.00 ± 0.07	1.11 ± 0.13
NCIB 11723	G	100	1.72 ± 0.21	0.67 ± 0.08	0.75 ± 0.10
	G +X	102 ± 6	1.62 ± 0.22	0.69 ± 0.09	0.78 ± 0.12

^a Relative rate of acid production (mean ± standard deviation, %) obtained from six independent experiments. Significant difference between relative rates of acid production in the presence and absence of xylitol: #*P* < 0.01.

^b Amounts of acidic end products (mean ± standard deviation, μmol/mg cells) obtained from three independent experiments. Significant difference between amounts of acidic end products in the presence and absence of xylitol: **P* < 0.05.

acidic end products were analyzed by the Dunn test.

Results

Inhibitory effect of xylitol on acid production

Xylitol significantly inhibited the acid production from glucose of *S. mutans* ATCC 31989 and NCTC 10449. In the presence of 30 mM xylitol, the acid production rates of ATCC 31989 and NCTC 10449 were decreased by 86 ± 1% (*P* < 0.01) and 26 ± 9% (*P* < 0.01), respectively, whereas that of NCIB 11723 was not inhibited (Table 1).

The total amounts of acidic end products generated by ATCC 31989 and NCTC 10449 cells decreased in the presence of xylitol. The reduction of lactic acid was

remarkable and it was significant in ATCC 31989 (*P* < 0.05), resulting in an end product shift to formate-acetate-dominant (Table 1). Xylitol had no effect on NCIB11723.

Glucose, xylitol and fructose : PEP-PTS activities

All *S. mutans* strains had PEP-PTS activities for glucose, xylitol, and fructose (Fig. 1). In ATCC 31989, fructose : PEP-PTS activity at 5–120 mM fructose and xylitol : PEP-PTS activity at 30–120 mM xylitol were higher than glucose : PEP-PTS activity. In particular, xylitol : PEP-PTS activity at 30 mM xylitol was 120 ± 14% of glucose : PEP-PTS activity at 10 mM glucose. In NCTC 10449, both fructose and xylitol : PEP-PTS activities

were lower than glucose : PEP-PTS activity. Xylitol : PEP-PTS activity at 30 mM xylitol was 16 ± 6% of glucose : PEP-PTS activity at 10 mM glucose. Both fructose and xylitol : PEP-PTS activities were low in NCIB 11723, and xylitol : PEP-PTS at 30 mM xylitol activity was only 3 ± 1% of glucose : PEP-PTS activity at 10 mM glucose.

Decrease in glucose : PEP-PTS activity in the presence of xylitol

In the presence of added xylitol, glucose : PEP-PTS activities of ATCC 31989 and NCTC 10449 were decreased (Fig. 2). The presence of 30 mM xylitol decreased glucose : PEP-PTS activity to 67 ± 6 and 83 ± 5%, respectively. As the xylitol concentration increased, the decrease became larger and statistically significant over 60 mM xylitol. However, little inhibition was observed in NCIB 11723.

Effect of xylitol on the profile of glycolytic intermediates during glucose metabolism

When metabolizing glucose only, all of the *S. mutans* strains had large amounts of fructose 1,6-bisphosphate, but the profiles of glycolytic intermediates downstream of




Fig. 1. PEP-PTS activities for glucose (○), xylitol (●) and fructose (shaded circle) of *Streptococcus mutans* ATCC 31989, NCTC 10449, and NCIB 11723. Vertical bars indicate standard deviations from three independent experiments. PEP-PTS activity for 10 mM glucose was regarded as 100%.




Fig. 2. PEP-PTS activities for 10 mM glucose of *Streptococcus mutans* in the presence of xylitol. ATCC 31989 (●), NCTC 10449 (□) and NCIB 11723 (◇). Significant difference between the PEP-PTS activities in the presence and absence of xylitol: **P* < 0.05, ***P* < 0.01. Vertical bars indicate standard deviations from three independent experiments.




Fig. 3. Glycolytic intermediates and xylitol 5-phosphate of *S. mutans* ATCC 31989, NCTC 10449, and NCIB 11723, 2 min after adding glucose (10 mM) or a mixture of glucose (10 mM) and xylitol (30 mM). Glycolytic intermediates (G, □) in the absence of xylitol. Glycolytic intermediate (G+X) and xylitol 5-phosphate (X) in the presence of xylitol. G6P, glucose 6-phosphate; F6P, fructose 6-phosphate; FBP, fructose 1,6-bisphosphate; DHAP, dihydroxyacetone phosphate; G3P, glyceraldehyde 3-phosphate; 3PG, 3-phosphoglycerate; 2PG, 2-phosphoglycerate; PEP, phosphoenolpyruvate; PYR, pyruvate. Significant difference between intermediate levels in the presence and absence of xylitol: * $P = 0.05$. Horizontal bars indicate standard deviations from four independent experiments.

dihydroxyacetone phosphate were slightly different among strains (Fig. 3).

When glucose was being metabolized by the ATCC 31989 and NCTC 10449 strains in the presence of xylitol, xylitol 5-phosphate accumulated and the amounts of glycolytic intermediates decreased, particularly those of fructose 1,6-bisphosphate ($P = 0.05$) in ATCC 31989 and NCTC 10449 and glucose 6-phosphate (G6P) ($P = 0.05$) in ATCC 31989. However, such inhibition was not evident in NCIB 11723.

Discussion

As previously reported (13, 25), xylitol inhibited the acid production of *S. mutans* strains and decreased the lactate production in strains ATCC 31989 and NCTC 10449 (Table 1), but the degree of these inhibitions varied between strains. On the other hand, in strain NCIB 11723, xylitol did not inhibit acid production or shift the end product profile.

Trahan (18) proposed an inhibitory mechanism of xylitol in which *S. mutans* transports xylitol as xylitol 5-phosphate through the activity of xylitol : PEP-PTS and, consequently, the xylitol 5-phosphate inhibits phosphoglucose isomerase and phosphofructokinase, the glycolytic enzymes for G6P conversion to fructose 1,6-bisphosphate, resulting in a decrease in intracellular levels of fructose 1,6-bisphosphate and the entire glycolytic rate. Miyasawa et al. (13) and Maehara et al. (11) then confirmed this notion, and found the decrease in lactate production was due to the decrease in fructose 1,6-bisphosphate,

an absolute activator of streptococcal lactate dehydrogenase. The present study found that higher xylitol : PEP-PTS activity indicated more inhibition of acid production by xylitol (Table 1 and Fig. 1). This observation supports the notion that xylitol 5-phosphate produced by xylitol : PEP-PTS activity is responsible for glycolytic inhibition and suggests that *S. mutans* strains with higher xylitol : PEP-PTS activity are more sensitive to xylitol inhibition. It has been proposed that xylitol is transported via a constitutive fructose : PEP-PTS and that xylitol : PEP-PTS activity appears as part of the constitutive fructose : PEP-PTS activity (18, 19, 22). We found here that high fructose : PEP-PTS activities in *S. mutans* were accompanied by high xylitol : PEP-PTS activities (Fig. 1), supporting this notion. Furthermore, it is reported that the *fxpC* gene of the constitutive fructose : PEP-PTS was located in the genomes of xylitol-sensitive streptococci and the *fxpC*-defective mutant was resistant to xylitol (2), although no information is available about *fxpC* gene in the strains used in our study.

Analyses of metabolic intermediates revealed that the xylitol-sensitive strains ATCC 31989 and NCTC 10449 accumulated xylitol 5-phosphate and decreased fructose 1,6-bisphosphate in the presence of xylitol (Fig. 3). These results support the notion that xylitol inhibits the glycolytic enzymes required for G6P conversion to fructose 1,6-bisphosphate. Despite the powerful inhibitory effect of xylitol (Table 1) and high xylitol : PEP-PTS activity (Fig. 1), the accumulation of xylitol 5-phosphate in ATCC 31989 seemed

to be smaller than that in NCTC 10449. It is suggested that the glycolytic enzymes of ATCC 31989 are more sensitive to xylitol and a small amount of xylitol 5-phosphate is enough to inhibit the entire glycolytic metabolism.

Not only fructose 1,6-bisphosphate but also G6P significantly decreased in a highly xylitol-sensitive strain ATCC 31989 (Fig. 3), indicating that xylitol itself can inhibit the glucose uptake system (e.g. glucose : PEP-PTS) and result in a decrease in intracellular G6P. This was confirmed by the observation that the presence of xylitol decreased the glucose : PEP-PTS activity in xylitol-sensitive strains, ATCC 31989 and NCTC 10449 (Fig. 2). This could be due to competition for the phosphoryl donor, HPr-P, between the glucose and the xylitol : PEP-PTSs. In the ATCC 31989 strain, with high xylitol : PEP-PTS activity and powerful xylitol inhibition of glucose : PEP-PTS activity, HPr-P could phosphorylate xylitol to xylitol 5-phosphate efficiently and result in a direct inhibition by xylitol 5-phosphate on glycolytic enzymes and an indirect inhibition on glucose phosphorylation by HPr-P.

NCIB 11723 isolated from human dental plaque by Carlsson (4) has natural xylitol resistance like other xylitol-resistant strains isolated from xylitol consumers (21, 23). It has been suggested that the xylitol resistance is due to the absence of a constitutive fructose : PEP-PTS by which xylitol is also incorporated, thus preventing xylitol-resistant strains from incorporating xylitol (19, 21). In the present study, NCIB 11723 had little xylitol : PEP-PTS

activity (Fig. 1) and did not accumulate xylitol 5-phosphate (Fig. 3) in the presence of a low concentration of xylitol, supporting this notion. As the xylitol concentration increased, however, xylitol : PEP-PTS activities of NCIB 11723 appeared (Fig. 1). In the presence of 120 mM xylitol, the xylitol : PEP-PTS activity compared with that of glucose : PEP-PTS reached $12 \pm 4\%$. However, 120 mM xylitol did not inhibit glucose : PEP-PTS activity (Fig. 2) and negligibly inhibited acid production from glucose (data not shown). These findings indicate that xylitol has less affinity for HPr-P than glucose in NCIB 11723. Thus, the strain cannot incorporate xylitol as xylitol 5-phosphate in the presence of both xylitol and glucose.

In conclusion, xylitol sensitivity varies among *S. mutans* strains: the higher the xylitol : PEP-PTS activity, and the more effectively xylitol decreases glucose : PEP-PTS activity, the more sensitive the strain is to xylitol. Xylitol has two inhibitory mechanisms:

- direct inhibition of glycolytic enzymes by xylitol 5-phosphate derived from xylitol : PEP-PTS;
- possibly, indirect inhibition of sugar uptake through competition for the phosphoryl donor, HPr-P, between the glucose and the xylitol : PEP-PTS.

Acknowledgments

This study was supported in part by a research fellowship (no. 163025 to HH) and Grants-in-Aid for Scientific Research (B) (no. 16390601 to NT) from the Japan Society for the Promotion of Science.

References

1. Assev S, Wåler SM, Rølla G. Further studies on the growth inhibition of some oral bacteria by xylitol. *Acta Pathol Microbiol Immunol Scand [B]* 1983; 91: 261–265.
2. Benchabane H, Lortie LA, Buckley ND, Trahan L, Frenette M. Inactivation of the *Streptococcus mutans fspC* gene confers resistance to xylitol, a caries-preventive natural carbohydrate sweetener. *J Dent Res* 2002; 81: 380–386.
3. Bradshaw DJ, Marsh PD. Effect of sugar alcohols on the composition and metabolism of a mixed culture of oral bacteria grown in a chemostat. *Caries Res* 1994; 28: 251–256.
4. Carlsson J. Presence of various types of non-haemolytic streptococci in dental plaque and in other sites of the oral cavity in man. *Odontol Revy* 1967; 18: 55–74.
5. Gauthier L, Vadeboncoeur C, Mayrand D. Loss of sensitivity to xylitol by *Streptococcus mutans* LG-1. *Caries Res* 1984; 18: 289–295.
6. Gehring F, Mäkinen KK, Larmas M, Scheinin A. Turku sugar studies. IV. An intermediate report on the differentiation of polysaccharide-forming streptococci (*S. mutans*). *Acta Odontol Scand* 1974; 32: 435–444.
7. Kakuta H, Iwami Y, Mayanagi H, Takahashi N. Xylitol inhibition of acid production and growth of mutans streptococci in the presence of various dietary sugars under strictly anaerobic conditions. *Caries Res* 2003; 37: 404–409.
8. Kenney EB, Ash MM. Oxidation reduction potential of developing plaque, periodontal pockets and gingival sulci. *J Periodontol* 1969; 40: 630–633.
9. Kornberg HL, Reeves RE. Inducible phosphoenolpyruvate-dependent hexose phosphotransferase activities in *Escherichia coli*. *Biochem J* 1972; 128: 1339–1344.
10. Lengeler JW, Jahreis K, Wehmeier UF. Enzymes II of the phosphoenolpyruvate-dependent phosphotransferase systems: their structure and function in carbohydrate transport. *Biochim Biophys Acta* 1994; 1188: 1–28.
11. Maehara H, Iwami Y, Mayanagi H, Takahashi N. Synergistic inhibition by combination of fluoride and xylitol on glycolysis by mutans streptococci and its biochemical mechanism. *Caries Res* 2005; 39: 521–528.
12. Minakami S, Suzuki C, Saito T, Yoshikawa H. Studies on erythrocyte glycolysis. I. Determination of the glycolytic intermediates in human erythrocytes. *J Biochem* 1965; 58: 543–550.
13. Miyasawa H, Iwami Y, Mayanagi H, Takahashi N. Xylitol inhibition of anaerobic acid production by *Streptococcus mutans* at various pH levels. *Oral Microbiol Immunol* 2003; 18: 215–219.
14. Postma PW, Lengeler JW, Jacobson GR. Phosphoenolpyruvate: carbohydrate phosphotransferase systems of bacteria. *Microbiol Rev* 1993; 57: 543–594.
15. Rogers AH, Pilowsky KA, Zilm PS, Gully NJ. Effects of pulsing with xylitol on mixed continuous cultures of oral streptococci. *Aust Dent J* 1991; 36: 231–235.
16. Rølla G, Oppermann RV, Waaler SM, Assev S. Effect of aqueous solutions of sorbitol-xylitol on plaque metabolism and on growth of *Streptococcus mutans*. *Scand J Dent Res* 1981; 89: 247–250.
17. Takahashi N, Abbe K, Takahashi-Abbe S, Yamada T. Oxygen sensitivity of sugar metabolism and interconversion of pyruvate formate-lyase in intact cells of *Streptococcus mutans* and *Streptococcus sanguis*. *Infect Immun* 1987; 55: 652–656.
18. Trahan L. Xylitol: a review of its action on mutans streptococci and dental plaque – its clinical significance. *Int Dent J* 1995; 45: 77–92.
19. Trahan L, Bareil M, Gauthier L, Vadeboncoeur C. Transport and phosphorylation of xylitol by a fructose phosphotransferase system in *Streptococcus mutans*. *Caries Res* 1985; 19: 53–63.
20. Trahan L, Bourgeau G, Breton R. Emergence of multiple xylitol-resistant (fructose PTS-) mutants from human isolates of mutans streptococci during growth on dietary sugars in the presence of xylitol. *J Dent Res* 1996; 75: 1892–1900.
21. Trahan L, Mouton C. Selection for *Streptococcus mutans* with an altered xylitol transport capacity in chronic xylitol consumers. *J Dent Res* 1987; 66: 982–988.
22. Trahan L, Neron S, Bareil M. Intracellular xylitol-phosphate hydrolysis and efflux of xylitol in *Streptococcus sobrinus*. *Oral Microbiol Immunol* 1991; 6: 41–50.
23. Trahan L, Soderling E, Drean MF, Chevrier MC, Isokangas P. Effect of xylitol consumption on the plaque-saliva distribution of mutans streptococci and the occurrence and long-term survival of xylitol-resistant strains. *J Dent Res* 1992; 71: 1785–1791.
24. Vadeboncoeur C, St Martin S, Brochu D, Hamilton IR. Effect of growth rate and pH on intracellular levels and activities of the components of the phosphoenolpyruvate: sugar phosphotransferase system in *Streptococcus mutans* Ingbritt. *Infect Immun* 1991; 59: 900–906.
25. Vadeboncoeur C, Trahan L, Mouton C, Mayrand D. Effect of xylitol on the growth and glycolysis of acidogenic oral bacteria. *J Dent Res* 1983; 62: 882–884.