

Nishikawa A, Hirose M.	gland carcinogenesis after initiation with diethylnitrosamine (DEN) and 1,2-dimethylhydrazine (DMH).				
Okazaki K, Ishii Y, Kitamura Y, Maruyama S, Umemura T, Miyauchi M, Yamagishi M, Imazawa T, Nishikawa A, Yoshimura Y, Nakazawa H, Hirose M.	Dose dependent promotion of forestomach carcinogenesis by combined treatment with sodium nitrite and ascorbic acid after initiation with N-methyl-N'-nitro-N-nitrosoguanidine - Possible contribution of nitric oxide- associated oxidative DNA damage.	Cancer Sci.	97	175-182	2006
Kitamura Y, Yamagishi M, Umemura T, Nishikawa A, Hirose M.	Lack of enhancing effects of sodium nitrite on 2-amino-1-methyl-6-phenylimidazo[4,5- <i>b</i>]pyridine (PhIP)-induced mammary carcinogenesis in female Sprague-Dawley rats.	Cancer Lett.	235	69-74	2006
Kitamura Y, Umemura T, Kanki K, Ishii Y, Kuroiwa Y, Masegi T, Nishikawa A, Hirose M.	Lung as a new target in rats of 2-amino-3-methylimidazo[4,5- <i>f</i>]quinoline carcinogenesis: Results of a two-stage model initiated with N-bis(2-hydroxypropyl)nitrosamine.	Cancer Sci.	97	368-373	2006
Sakano K, Mizutani M, Murata M, Oikawa S, Hiraku Y, Kawanishi S.	Procyanidin B2 has anti- and pro-oxidant effects on metal-mediated DNA damage.	Free Radic. Biol. Med.	39	1041-1049	2005
Oikawa S, Ito T, Iwayama M, Kawanishi S.	Radical production and DNA damage induced by carcinogenic 4-hydrazinobenzoic acid, an ingredient of mushroom <i>Agaricus bisporus</i> .	Free Radic. Res.	40	31-39	2005
Piao F, Ma N, Hiraku Y, Murata M, Oikawa S.	Oxidative DNA damage in relation to neurotoxicity in	J. Occup. Health	47	445-449	2005

S, Cheng F, Zhong L, Yamauchi T, Kawanishi S, Yokoyama K.	the brain of mice exposed to arsenic at environmentally relevant levels.				
Kawanishi S, Oikawa S, Murata M.	Evaluation for safety of antioxidant chemopreventive agents.	Antioxid Redox Signal	7	1728-1739	2005
Hongo T, Kajikawa M, Ishida S, Ozawa S, Ohno Y, Sawada J, Umezawa A, Ishikawa S, Kobayashi T, and Honda H	Three-dimensional high-density culture of HepG2 cells in a 5-ml radial-flow bioreactor for construction of artificial liver	J. Bioscience and Bioengineering	99	237-244	2005
Kubo T, Kim S-R, Sai K, Saito Y, Nakajima T, Matsumoto K, Saito H, Shirao K, Yamamoto N, Minami H, Ohtsu A, Yoshida T, Saijo N, Ohno Y, Shogo Ozawa S, Sawada J.	Functional Characterization of Three Naturally Occurring Single Nucleotide Polymorphisms in the CES2 gene encoding carboxylesterase 2 (hCE-2).	Drug Metab. Disp.	33	1482-1487	2005
Yawata A, Kim SR, Miyajima A, Kubo T, Ishida S, Saito Y, Nakajima Y, Katori N, Matsumoto Y, Fukuoka M, Ohno Y, Ozawa S, Sawada J	Polymorphic tandem repeat sequences of the thymidylate synthase gene correlates with cellular-based sensitivity to fluoropyrimidine antitumor agents.	Cancer Chemother. Pharmacol.	56	465-472	2005
Kuribayashi, M., Asamoto, M., Suzuki, S., Hokaiwado, N.,	Lack of modification of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) rat	Cancer Lett.	232	289-299	2005

Ogawa, K. and Shirai, T.	hepatocarcinogenesis by caffeine, a CYP1A2 inducer, points to complex counteracting influences.				
-----------------------------	----------------------------------------------------------------------------------------------------------	--	--	--	--