

Table 19
 Incidence of hepatocellular hypertrophy in centrilobular area in male rats
 - Carcinogenicity study -

	Dose (%)				
	0	0.02	0.3	1.0	3.0
No. of rats examined	50	47	44	49	50
Hepatocellular hypertrophy, slight	5	3	4	8	18*

*, significantly different from the control value at $p < 0.01$

Table 20-1

Incidence of neoplastic lesions in rats of the control and high dose groups
- Carcinogenicity study -

Organ	Findings	Male		Female	
		0% 50	3% 50	0% 50	3% 50
Liver	(M) Angiosarcoma	1	0	0	0
	(B) Adenoma	0	0	1	0
Kidney	(M) Liposarcoma	1	0	0	0
Heart	(M) Endocardial schwannoma, malignant	1	0	0	0
	(B) Endocardial schwannoma	0	2	0	0
Brain	(B) Astrocytoma	0	1	0	0
Pituitary	(M) Anterior adenocarcinoma	0	0	1	3
	(B) Anterior adenoma	13	12	22	16
Thymus	(M) Thymoma	1	3	3	3
Pancreas	(M) Islet cell adenocarcinoma	2	3	1	0
	(B) Islet cell adenoma	1	2	1	1
	(B) Acinar cell adenoma	0	1	0	0
Adrenal	(M) Malignant pheochromocytoma	1	0	0	0
	(B) Pheochromocytoma	2	0	1	1
	(B) Adenoma of cortex	0	0	0	1
	(B) Neuroganglioma	0	0	1	0
LN	(M) Malignant lymphoma, systemic	0	1	1	1
	(B) Hemangioma	1	1	0	0
Thyroid	(B) C-cell adenoma	3	6	3	3
	(B) Follicular cell adenoma	3	1	1	0
Parathyroid	(B) Adenoma	0	0	1	0

Table 20-2

Incidence of neoplastic lesions in rats of the control and high dose groups -Continued-
 - Carcinogenicity study -

Organ	Findings	Male		Female	
		0% 50	3% 50	0% 50	3% 50
<i>continued</i>					
Stomach	(M) Fibrosarcoma	1	0	0	0
Intestine	(B) Fibroma	0	0	1	0
Muscle	(B) Granular cell tumor	0	1	0	0
Mammary gland	(M) Adenocarcinoma	0	0	5	2
	(B) Adenoma	0	0	5	1
	(B) Fibroadenoma	1	0	9	4
Testis	(B) Leydig cell tumor	0	2		
Prostate	(B) Adenoma of ventral lobe	1	0		
Preputial gland	(B) Adenoma	1	0		
Ovary	(M) Granulosa-theca cell tumor, malignant			1	0
	(M) Cystadenocarcinoma			1	0
	(B) Granulosa-theca cell tumor			1	1
	(B) Luteoma			0	1
Uterus	(M) Stromal sarcoma			1	0
	(M) Endometrial adenocarcinoma			0	1
	(B) Endometrial stromal polyp			7	5
Nasal cavity	(M) Osteosarcoma	1	0	0	0
Spinal cord	(B) Osteoma	0	0	1	0
Abdominal cavity	(M) Angiosarcoma	1	0	0	0
Subcutis/skin	(B) Fibroma	3	0	0	0
	(B) Keratoacanthoma	0	1	0	0
	(M) SCC	0	0	1	0
	(B) Lipoma	0	0	1	0
	(M) Malignant Schwannoma	0	1	0	0
Scrotum	(M) Mesothelioma	1	1		
Parotid gland	(M) Adenocarcinoma	0	0	0	1

研究成果の刊行に関する一覧表

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Nishikawa, M., Wanibuchi, H., Morimura, K., Kinoshita, A., Nishikawa, T., Hayashi, S., Yano, Y. and Fukushima, S.:	N-acetylcysteine and S-methylcysteine inhibit MeIQx rat hepatocarcinogenesis in the post-initiation stage.	Carcinogenesis	27	982-988	2005
Wei, M., Morimura, K., Wanibuchi, H., Shen, J., Salim, E. I., Moku, M., Hakoi, K. and Fukushima, S.	JTE-522, a selective cyclooxygenase-2 inhibitor, inhibits induction but not growth and invasion of 1,2-dimethylhydrazine-induced tubular adenocarcinomas of colon in rats.	Int. J. Cancer	113	354-358	2005
Kuroda, K., Yoshida, K., Yoshimura, M., Endo, Y., Wanibuchi, H., Fukushima, S. and Endo, G.	Genotoxicity of dimethylarsinous acid: high induction of tetraploids.	Appl. Organometal. Chem.	19	221-225	2005
Kang, J.S., Morimura, K., Salim, E.I., Wanibuchi, H., Yamaguchi, S. and Fukushima, S.	Persistence of liver cirrhosis in association with proliferation of nonparenchymal cells and altered location of α -smooth muscle actin-positive cells.	Toxicol. Pathol.	33	329-335	2005
Kang, J.S., Wanibuchi, H., Morimura, K., Puatanachokchai, R., Salim, E.I., Hagihara, A., Seki, S. and Fukushima, S.	Enhancement by estradiol 3-benzoate in thioacetamide-induced liver cirrhosis of rats.	Toxicol. Sci.	85	720-726	2005
Fukushima, S., Wanibuchi, H., Morimura, K., Nakae, D., Tsuda, H., Imaida, K., Shirai, T., Tatematsu, M., Tsukamoto, T., Hirose, M., Furukawa, F.	Lack of potential of low dose N-nitrosodimethylamine to induce preneoplastic lesions, glutathione S-transferase placental form-positive foci, in rat liver.	Cancer Lett.	222	11-15	2005

Doi, K., Wanibuchi, H., Salim, E. I., Morimura, K., Kinoshita, A., Kudoh, S., Hirata, K., Yoshikawa, J. and Fukushima, S.	Lack of large intestinal carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5- <i>b</i>]pyridine at low doses in rats initiated with azoxymethane.	Int. J. Cancer.	115	870-878	2005
Fukushima, S., Morimura, K., Wanibuchi, H., Kinoshita, A. and Salim, E. I.	Current and emerging challenges in toxicopathology: carcinogenic threshold of phenobarbital and proof of arsenic carcinogenicity using rat medium-term bioassays for carcinogens.	Toxicol. Appl. Pharmacol.	207	S225-S229	2005
Fukushima, S., Kinoshita, A., Puatanachokchai, R., Kushida, M., Wanibuchi, H. and Morimura, K.	Hormesis and dose-response-mediated mechanisms in carcinogenesis: evidence for a threshold in carcinogenicity of non-genotoxic carcinogens.	Carcinogenesis	26	1835-1845	2005
Murai, T., Mori, Y., Tatematsu, K., Koide, A., Hagiwara, A., Makino, S., Mori, S., Wanibuchi, H. and Fukushima, S.	Differences in susceptibility to <i>N</i> -butyl- <i>N</i> -(4-hydroxybutyl)nitrosamine-induced urinary bladder carcinogenesis between SD/gShi rats with spontaneous hypospermatogenesis and SD/cShi rats with spontaneous hydronephrosis.	Cancer Sci.	96	637-644	2005
Kushida, M., Sukata, T., Uwagawa, S., Ozaki, K., Kinoshita, A., Wanibuchi, H., Morimura, K., Okuno, Y. and Fukushima, S.	Low dose DDT inhibition of hepatocarcinogenesis initiated by diethylnitrosamine in male rats: Possible mechanisms.	Toxicol. Appl. Pharmacol.	208	285-294	2005
Kushida, M., Wanibuchi, H., Morimura, K., Kinoshita, A., Kang, J. S., Puatanachokchai, R., Wei, M., Funae, Y and Fukushima, S.	Dose-dependence of Promotion of 2-amino-3,8-dimethylimidazo[4,5- <i>f</i>]quinoxaline-induced rat hepatocarcinogenesis by ethanol: Evidence for a threshold.	Cancer Sci.	96	747-757	2005
Morimura, K., Kang, J. S., Wei, M., Wanibuchi, H., Tsuda, H. and Fukushima, S.	Lack of urinary bladder carcinogenicity of sodium L-ascobate in human <i>c-Ha-ras</i> proto-oncogene transgenic rats.	Toxicol. Pathol.	33	764-767	2005

Nishikawa, M., Wanibuchi, H., Morimura, K., Kinoshita, A., Nishikawa, T., Hayashi, S., Yano, Y. and Fukushima, S.	N-acetylcysteine and S- methylcysteine inhibit MeIQx rat hepatocarcinogenesis in the post-initiation stage.	Carcinogene sis	27	982-988	2005
Hagiwara, A., Murai, T., Miyata E., Nabae, K., Doi, Y., Yoshino, H., Ichihara, T., Mori, S. and Fukushima, S.	Influence of strain and diet on hepatocarcinogenicity of N- butyl-N-(4- hydroxybutyl)nitrosamine (BBN) in rats: slight inhibition of preneoplastic liver lesion development by uracil.	J. Toxicol. Pathol.	18	105-110	2005
Wei, M., Hori, T., Wanibuchi, H., Morimura, K., Kang, J.S., Puatanachokchai, R. and Fukushima, S.	Existence of no-observed effect levels for 2-amino- 3,8-dimethylimidazo[4,5- f]quinoxaline on hepatic preneoplastic lesion development in BN rats.	Cancer Letters	231	304-308	2006
Romanenko, A., Morimura, K., Kinoshita, A., Wanibuchi, H., Vozianov, A. and Fukushima, S.	Aberrant expression of E- cadherin and β -catenin in association with transforming growth factor- β 1 in urinary bladder lesions in humans after the Chernobyl accident.	Cancer Sci.	97	45-50	2006
Shen, J., Wanibuchi, H., Waalkes, M. P., Salim, E. I. Kinoshita, A., Yoshida, K., Endo, G. and Fukushima, S.	A comparative study of the sub-chronic toxic effects of three organic arsenical compounds on the urothelium in F344 rats; gender-based differences in response.	Toxicol. Appl. Pharmacol.	210	171-180	2006
Kang, J. S., Morimura, K., Toda, C., Wanibuchi, H., Wei, M., Kojima, N. and Fukushima, S.	Testicular toxicity of DEHP, but not DEHA, is elevated under conditions of thioacetamide-induced liver damage.	Reproductiv e Toxicology	21	253-259	2006
Romanenko, A. M., Kinoshita, A., Wanibuchi, H., Wei, M., Zaparin, W. K., Vinnichenko, W. I., Vozianov, A. F. and Fukushima, S.	Involvement of ubiquitination and sumoylation in bladder lesions induced by persistent long-term low dose ionizing radiation in humans.	The Journal of Urology	175	739-743	2006

Hagihara, A., Wanibuchi, H., Puatanachokchai, R., Kang, J. S., Miyazi, N., Seki, S. and Fukushima, S.	Difference in sensitivity of F344 rats from different breeders to phenobarbital hepatocarcinogenicity.	J Toxicol Pathol.	19	29-36	2006
Kang, J. S., Waniuchi, H., Morimura, K., Totsuka, Y., Yoshimura, I. and Fukushima, S.	Existence of a no effect level for MeIQx hepatocarcinogenicity on background of thioacetamide-induced liver damage in rats.	Cancer Sci.	97	453-458	2006
Kushida, M., Aiso, S., Morimura, K., Wei, M., Wanibuchi, H., Nagano, K. and Fukushima, S.	Absence of β -Catenin Alteration in Hepatic Tumors Induced by p- Nitroanisole in Crj:BDF1 Mice.	Toxicologic Pathology	34	237-242	2006
Wanibuchi, H., Wei, M., Karim, M. R., Morimura, K., Doi, K., Kinoshita, A. and Fukushima, S.	Existence of no hepatocarcinogenic effect levels of 2-amino-3, 8- dimethylimidazo [4, 5-f] quinoxaline with or without coadministration with ethanol.	Toxicologic Pathology	34	232-236	2006
Waddell, W. J., Fukushima, S. and Williams, G. M.	Concordance of thresholds for carcinogenicity of N- nitrosodierhylamine.	Arch Toxicol	80	305-309	2006
Uda, N., Kashimto, N., Sumioka, I., Kyo, E., Sumi, S. and Fukushima, S.	Aged garlic extract inhibits development of putative preneoplastic lesions in rat hepatocarcinogenesis.	American Society for Nutrition,		855S- 860S	2006
Kitano, M., Chen, Tian-Xin., Morimura, K., Wei, M., Hidaka, T., Hosoe, K. and Fukushima, S.	Biphasic increase of epithelial DNA synthesis in the urinary bladder of rats treated with a tumor promoter, sodium L- ascorbate.	J. Toxicol. Pathol.	19	53-56	2006
Kitano, M., Wada, J., Ariki, Y., Kato, M., Wanibuchi, H., Morimura, K., Hidaka, T., Hosoe, T. and Fukushima, S.	Possible tumor development from double positive foci for TGF- α and GST-P observed in early stages on rat hepatocarcinogenesis.	Cancer Sci.	97	478-483	2006
Mori, S., Murai, T., Wanibuchi, H., Hagihara, A., Puatanachokchai, P. and Fukushima, S.	Susceptibility of four F ₁ hybrids of male rats to the promoting effects of sodium L-ascorbate in two-stage urinary bladder carcinogenesis.	J Toxicol Pathol.	19	87-91	2006

Puatanachokchai, R., Morimura, K., Wanibuchi, H., Oka, M., Kinoshita, A., Fukui, M., Yamaguchi, S., Funae, Y. and Fukushima, S.	Alpha-benzene hexachloride exerts hormesis in preneoplastic lesion formation of rat hepatocarcinogenesis with the possible role for hepatic detoxifying enzymes.	Cancer Lett.	240	102-113	2006
Nakajima, Y., Endo, Y., Inoue, Y., Yamanaka, K., Kato, K., Wanibuchi, H. and Endo, G.	Ingestion of Hijiki seaweed and risk of arsenic poisoning +.	Appl Organometal Chem.	20	557-564	2006
Kinoshita, A., Wanibuchi, H., Wei, M. and Fukushima, S.	Hormesis in carcinogenicity of non-genotoxic carcinogens.	J.Toxicol.P athol.	19	111-122	2006
Romanenko, A. M., Morimura, K., Kinoshita, A., Wanibuchi, H., Takahashi, S., Zaparin, W. K., Vinnichenko, W. K., Vozianov, A. F. and Fukushima S.	Upregulation of fibroblast growth factor receptor 3 and epidermal growth factor receptors, in association with Raf-1, in urothelial dysplasia and carcinoma in situ after the Chernobyl accident.	Cancer Sci.	97	1168- 1174	2006
Shirai T, Kawabe M, Ichihara T, Fujiwara O, Taki M, Watanabe S, Wake K, Yamanaka Y, Imaida K, Asamoto M, Tamano S.	Chronic exposure to a 1.439 GHz electromagnetic field used for cellular phones does not promote N- ethylnitrosourea induced central nervous system tumors in F344 rats.	Bioelectrom agnetics	26	59-68	2005
Ichihara T, Miyashita K, Kawabe M, Imaida K, Asamoto M, Ogiso T, Tamano S, Hirose M, Shirai T.	Lack of combination hepatocarcinogenicity of harman, norharman and amitrole when given with NaNO ₂ in the rat.	Journal of Toxicologic al Sciences	30	1-6	2005
Ichihara T, Tamano S, Yoshino H, Imaida K, Ishikawa H, Kakizoe T, Shirai T.	Lack of chemoprevention or promotion effects of docosahexaenoic acid on small intestine, colon, liver, lung, thyroid, esophagus, kidney, and forestomach carcinogenesis in a rat medium-term multi- organ carcinogenesis model.	Journal of Toxicologic Pathology	18	53-89	2005
Zeng Y, Yokohira M, Takeuchi H, Saoo K, Yamakawa K, Matsuda Y, Hosokawa, K, Imaida K.	Lack of significant modifying effect of arctiin on prostate carcinogeesis in probasin/SV40 T antigen transgenic rats.	Cancer Letters	222	145-15	2005

Zeng Y, Saoo K, Yokohira M, Takeuchi H, Yamakawa K, Matsuda Y, J.Q. Li, Imaida K.	Dietary D-psicose, a rare sugar, does not show any modifying effects in a medium-term liver carcinogenesis bioassay in F344 male rat.	J. Toxicol. Pathol.	18	85-88	2005
Zeng Y, Yokohira M, Saoo K, Takeuchi H, Chen K, Yamakawa K, Matsuda K, Kakehi Y, Imaida K.	Inhibition of prostate carcinogenesis in probasin/SV40 T antigen transgenic rats by raloxifene, an antiestrogen with anti-androgen action, but not nimesulide, a selective cyclooxygenase-2 inhibitor	Carcinogenesis	26	1109-1116	2005
Fukushima S, Wanibuchi H, Morimura K, Nakae D, Tsuda H, Imaida K, Shirai T, Tatematsu M, Tsukamoto T, Hirose M, Furukawa F.	Lack of potential of low dose N-nitrosodimethylamine to induce preneoplastic lesions, glutathione S-transferase placental form-positive foci, in rat liver.	Cancer Letters	222	11-15	2005
Yokohira M, Takeuchi H, Yamakawa K, Saoo K, Ikeda M, Matsuda Y, Zeng Y, Hosokawa K, Maeta H, Imaida K.	Establishment of a biological system for detection of lung toxicity due to fine particle instillation: Sequential histopathological changes with acute and subacute lung damage due to intratracheal instillation of quartz in F344 male rats.	Journal of Toxicologic Pathology	18	13-18	2005
Takeuchi H, Saoo K, Matsuda Y, Yokohira M, Yamakawa K, Zeng Y, Miyazaki M, Fujieda M, Kamataki T, Imaida K.	Dose dependent inhibitory effects of dietary 8-methoxypsoralen on NNK-induced lung tumorigenesis in female A/J mice.	Cancer Letters			2005
Matsuda Y, Saoo K, Hosokawa K, Yamakawa K, Yokohira M, Zeng Y, Takeuchi H, Iwai J, Shirai T, Obika K, Imaida, K.	Epithelioid malignant peripheral nerve sheath tumor. Report of a case with inflammatory infiltration.	Pathology - Research and Practice	201	355-360	2005
今井田克己, 玉野静光, 萩原昭裕, 福島昭治, 白井智之, 伊東信行	内分泌活性物質の発癌性作用および発癌修飾性検出のためのラット中期バイオアッセイの適用.	内分泌活性物質のヒトおよび野生生物に及ぼす影響		779-783	2005

中江 大	佐々木研究所における化学物質のリスク評価・管理に資する毒性病理学的研究(1) コリン欠乏アミノ酸食投与ラット肝発がんモデルによる知見	化学生物総合管理	1	331-352	2005
Uematsu F, Takahashi M, Yoshida M, Igarashi M, Watanabe N, Suzuki N, Abe M, Rusyn I, Floyd RA and Nakae D.	Distinct patterns of gene expression in hepatocellular carcinomas and adjacent non-cancerous, cirrhotic liver tissues in rats fed a choline-deficient, L-amino acid-defined diet.	Cancer Sci.	96	414-424	2005
Powell CL, Kosyk O, Bradford BU, Denda A, Uematsu F, Nakae D and Rusyn I.	Temporal correlation of pathology and DNA damage with gene expression in a choline deficient model of rat liver injury.	Hepatology	42	1137-1147	2005
Kotake Y, Kishida H, Nakae D and Floyd RA.	Nitric oxide production by primary liver cells isolated from amino acid-fed rats.	Methods Enzymol	396	535-541	2005
Fukushima S, Wanibuchi H, Morimura K, Nakae D, Tsuda H, Imaida K, Shirai T, Tatematsu M, Tsukamoto T, Hirose M and Furukawa F.	Lack of potential of low dose <i>N</i> -nitrosodimethylamine to induce preneoplastic lesions, glutathione <i>S</i> -transferase placental form-positive foci, in rat liver.	Cancer Lett.	222	11-15	2005
Shimomoto T, Yoshida M, Katsuda S, Takahashi M, Uematsu F, Kuniyasu H, Maekawa A and Nakae D	• -Smooth muscle actin-positive stromal cells reactive to estrogens surround endometrial glands in rats but not in mice.	J Toxicol Pathol.	18	47-52	2005
Sakurai T, Takei M, Ogasawara J, Watanabe N, Sanpei M, Yoshida M, Nakae D.	Sakurai T, Nakano N, Kizaki T, Ohno H and Izawa T, Exercise training enhances tumor necrosis factor-• -induced expressions of anti-apoptotic genes without alterations in caspase-3 activity in rat epididymal adipocytes.	Jpn J Physiol.	55	181-189	2005
Yamada D, Yoshida M, Williams YN, Fukami T, Kikuchi S, Masuda M, Maruyama T, Ohta T, Nakae D, Maekawa A, Kitamura T and Murakami Y.	Disruption of spermatogenic cell adhesion and male infertility in mice lacking TSLC1/IGSF4, an immunoglobulin superfamily cell adhesion molecule.	Mol Cell Biol	26	3610-3624	2005