






Table 2. Continued.

Compound	Structure and CAS name	CAS RN MW Formula	Use/origin	Environmental occurrence	Trade names, comments, and nontarget species effects
3,4,5,6-Tetrabromo- <i>o</i> -cresol	 3,4,5,6-Tetrabromo- <i>o</i> -cresol	576-55-6 423.72 C <sub>7</sub> H <sub>4</sub> Br <sub>4</sub> O	Antiseptic, fungicide	POTWs in Germany: tetrabromo- <i>o</i> -cresol found in both influents and effluents (<0.1 µg/L) (70)	
Timolol	 ( <i>S</i> )-1-[(1,1-Dimethylethyl)-amino]-3-[4-(4-morpholinyl)-1,2,5-thiadiazol-3-yl]oxy]-2-propanol	26839-75-8 316.42 C <sub>13</sub> H <sub>24</sub> N <sub>4</sub> O <sub>3</sub> S	Beta-blocker (antihypertensive)	POTW max. effluent: 0.07 µg/L; max. in surface waters: 0.01 µg/L (18)	e.g., hemihydrate: Betimol
Tolfenamic acid	 2-[(3-Chloro-2-methylphenyl)-amino]benzoic acid	13710-19-5 261.71 C <sub>14</sub> H <sub>12</sub> ClNO <sub>2</sub>	Analgesic/anti-inflammatory	Not detected in POTW effluent or surface waters (18); in Brazilian STW effluent 1.6 µg/L (69)	Veterinary NSAID e.g., Tolfedine
Triclosan	 5-Chloro-2-(2,4-dichlorophenoxy)phenol	3380-34-5 289.54 C <sub>12</sub> H <sub>7</sub> Cl <sub>3</sub> O <sub>2</sub>	Antiseptic	0.05-0.15 µg/L in water (113). Antibacterial widely used for 30 years in a vast array of consumer products. Its usage as a preservative and disinfectant continues to grow. Triclosan's use in commercial products spans footwear (in hosiery and insoles of shoes called "odor-eaters"), hospital handsoap, acne creams (e.g., Clearasil), and rather recently as a slow-release product called Microban, which is incorporated in a wide variety of plastic products (from children's toys to kitchen utensils, such as cutting boards)	e.g., Irgasan DP 300
Verapamil	 α-[3-[[2-(3,4-Dimethoxyphenyl)-ethyl]methylamino]propyl]-3,4-dimethoxy-α-(1-methyl-ethyl)benzeneacetonitrile; e.g., hydrochloride: Vasolan	52-53-9 454.61 C <sub>27</sub> H <sub>38</sub> N <sub>2</sub> O <sub>4</sub>	Cardiac drug (calcium ion influx inhibitor) (antihypertensive)	No occurrence data	µM concentrations and lower greatly increase toxicity of certain drugs for many aquatic organisms (32) because of inhibition of "multidrug transporters"; removal of other drugs is reduced, lengthening the exposure time; <i>Daphnia</i> immobilization EC <sub>50</sub> 0.11–0.67 mM (125)

Abbreviations: CAS RN, Chemical Abstracts Service Registry Number; max, maximum; MW, molecular weight; NA, not applicable. \*In the first survey of Canadian aquatic life (154), maximum musk lipid concentrations from populated areas in Canada showed values in distinct contrast with those from Europe: musk ketone (2.2–17.7 µg/g lipid; mussels, winter flounder, clams); 0.2–0.7 (trout, eel, lobster); Galaxolide: 0.01–3.0 µg/g in the same samples. For a given sample, the lipid concentrations of musk ketone (the dominant nitro musk) were several-fold to over an order of magnitude greater than the concentrations for the predominant polycyclic (Galaxolide); most samples contained lower concentrations of musk xylene and Tonalide. These data contrast with those from Europe—the concentrations of nitro and polycyclics are comparable in Canada, whereas the nitro musks are 1–3 orders of magnitude lower than the polycyclics in Europe (presumably reflecting different usage patterns and the fact that the nitro musks have not been restricted in the West). In comparison with Europe, insignificant quantities of amino metabolites of nitro musks were found in Canadian aquatic life, presumably a reflection of the lower degree of sewage treatment in Canada [e.g., (2)].


ratio of drug concentrations in bile to that in the surrounding water can increase many orders of magnitude as exposure duration increases (15). Detection of exposure of fish to many drugs can thereby be facilitated through the analysis of bile.

The introduction of drugs into the environment is partly a function of the quantity of drugs manufactured, the dosage frequency and amount [the 200 most frequently prescribed drugs, representing about two-thirds of all

prescriptions filled in the United States for the most recently documented year, are listed in RxList (16)], the excretion efficiency of the parent compound and metabolites, propensity of the drug to sorb to solids, and the metabolic transformation capability of subsequent sewage treatment (or landfill) microorganisms. Publicly owned wastewater treatment plants (POTWs) receive influent from both domestic, municipal, and industrial (including pharmaceutical manufacture) sewage systems. The processed

liquid effluents from primary and secondary treatments are then discharged to surface waters and the residual solids (sludge) to landfills/farms; land disposal, including manure from treated animals at CAFOs, creates the potential for introduction into groundwaters or surface waters (via wet weather run-off). Theoretically, PPCPs in sewage sludge applied to crop lands could be taken up by plants.

Compounds surviving the various phases of metabolism and other degradative or



**Figure 1.** Metabolic approach to increasing the polarity (and excreability) of drugs.

sequestering actions (i.e., display environmental persistence) can then pose an exposure risk for organisms in the environment. Even the less/nontoxic conjugates (glucuronides) can later be converted back to the original bioactive compounds via enzymatic ( $\beta$ -glucuronidases) or chemical hydrolysis (e.g., acetylsalicylic acid can be hydrolyzed to the free salicylic acid). Some degradation products can even be more bioactive than the parent compound. Therefore, conjugates can essentially act as storage reservoirs from which the free drugs can later be released into the environment. Up to 90% of certain drugs can become conjugated (17,18), conjugation varying as a function of chemical class. These pathways of introduction into the environment have been summarized by Velagaleti (10).

**Sewage treatment plants.** Treatment facilities, primarily POTWs or sewage treatment works (STWs), which include privately owned works as well, play a key role in the introduction of pharmaceuticals into the environment [see Rogers (19) for a review of the fate of synthetic chemicals in sewage treatment plants]. STWs were designed to handle human waste of mainly natural origin, primarily via the acclimated degradative action of microorganisms (the efficiency of metabolism of a given drug can increase with duration of treatment because of enzyme induction and cellular adaptation) and the coagulation/flocculation of suspended solids; sometimes, tertiary treatment (e.g., chemical/ultraviolet [UV] oxidation) is used. Most anthropogenic chemicals introduced along with this normal waste suffer unknown fates. Two primary mechanisms remove substances from the incoming waste stream: *a*) microbial degradation to lower molecular weight products, leading sometimes to complete mineralization— $\text{CO}_2$  and  $\text{H}_2\text{O}$ ; and *b*) sorption to filterable solids, which are later removed with the sludge.

Although the microbiota of sewage treatment systems may have been exposed to many PPCPs for a number of years, two factors work against the effective microbial removal of these substances from STWs. First, the concentrations of most drugs are probably so low that the lower limits for enzyme affinities may not be met. For example, the daily loadings of PPCPs into STWs are largely a function of the serviced human population, the dosages/duration of medications

consumed, and the metabolic/excretory half-lives, which are all large variables. As an example, the daily load of a subset of pharmaceuticals to a particular POTW near Frankfurt/Main, Germany, ranged from tens to hundreds of grams, with approximate individual removal efficiencies varying widely from 10 to 100% but trending to around 60% (18). This particular POTW serviced about a third of a million people at a flow rate of roughly 60,000  $\text{m}^3/\text{day}$ . Despite the number of studies on treatment efficiencies, a widespread investigation is still lacking for the differences in removal efficiencies for distinct types of STWs as well as for individual treatment techniques. The extent to which a particular plant uses primary, secondary, and tertiary technologies will greatly influence removal efficiencies; the technologies employed vary widely among cities. The biodegradative fate of most compounds in STWs is governed by nongrowth-limiting (enzyme-saturating) substrate concentrations (copiotrophic metabolism). In contrast, PPCPs are present in STWs at concentrations at enzyme-saturating levels, which necessitates oligotrophic metabolism. These micropollutants might be handled by only a small subset of specialist oligotrophic organisms whose occurrence is probably more prevalent in native environments characterized by low-carbon fluxes (e.g., sediments and associated pore waters, where desorption mass transfer is limiting) than in STWs. This means that degradation of PPCPs may occur more prevalently in the receiving waters/sediments than in STWs.

Second, many new drugs are introduced to the market each year; some of these drugs are from entirely new classes never seen before by the microbiota of an STW. Each of these presents a new challenge to biodegradation. A worst-case scenario may not be unusual—the concentration of a drug leaving an STW in the effluent could essentially be the same as that entering. Only the several-fold to multiple order of magnitude dilution when the effluent is mixed into the receiving water, assuming a sufficiently high natural flow, serves to reduce the concentration; obviously, smaller streams have increased potential for having higher concentrations of any PPCP that has been introduced. In general, most pharmaceuticals resist extensive microbial degradation (e.g., mineralization) (10). Although some parent drugs often show poor solubility in water (10), leading to preferential sorption to suspended particles, they can thereby sorb to colloids and therefore be discharged in the aqueous effluent. Metabolites, including breakdown products and conjugates, will partition mainly to the aqueous effluent. Some published data demonstrate that many parent drugs do make their way

into the environment (see references cited in Table 2 under "Environmental Occurrence").

The efficiency of removal of pharmaceuticals by STWs is largely unknown. Currently, the most extensive study of treatment efficiency (18) reports removal from German STWs of 14 drugs representing five broad physiologic categories. Removal of the parent compound (keep in mind that possible subsequent metabolites were not accounted for) ranged from 7% (carbamazepine, an anti-epileptic) to 96% (propranolol, a beta-blocker); most removal efficiencies averaged about 60%. Fenofibrate, acetaminophen, and salicylic acid, *o*-hydroxyhippuric acid, and gentisic acid (acetylsalicylic acid metabolites) could not be detected in effluent; salicylic acid was found in the influent at concentrations up to 54  $\mu\text{g}/\text{L}$ . It is important to understand that absent the stoichiometric accounting of metabolic products, one cannot distinguish between the three major fates of a substance: *a*) degradation to lower molecular weight compounds, *b*) physical sequestration by solids (and subsequent removal as sludge), and *c*) conjugates that can later be hydrolyzed to yield the parent compound (e.g., clofibric and fenofibric acid conjugates) (18). Therefore, by simply following disappearance (removal) of a substance, one cannot conclude that it was structurally altered or destroyed—it may simply reside in another state or form. Identifying metabolic products is difficult not only because of the number of metabolites (sometimes several per parent compound) but also because standard reference materials are difficult to obtain commercially and can be costly.

Despite high removal rates in STWs for some drugs, upsets in the homeostasis of a treatment plant can result in higher than normal discharges. For example, Ternes (18) found that wet weather runoff dramatically reduced the removal rates for certain drugs (e.g., several nonsteroidal anti-inflammatory drugs [NSAIDs] and lipid regulators) in a facility located close to Frankfurt/Main. During the increased period of influent flow, the removal rate dropped to below 5% from over 60% previously; several days were required for the removal rates to recover. Clearly, even for drugs efficiently removed, the operational state of the STW can have a dramatic effect on the removal efficiencies. Other transients that could affect removal include transitions between seasons and sporadic plug-flow influx of toxicants from various sources. Overflows from STW failure or overcapacity events (e.g., floods, excessive water use) lead to direct, untreated introduction of sewage into the environment. In efforts to improve tributary conditions (by increasing stream flow), some cities have considered increasing the percentage of annual overflow events (e.g., see the Portland,

Oregon, proposal (20). The highest concentration in an STW effluent reported by Ternes (18) was for bezafibrate (4.6 µg/L); the highest concentration in surface water also was for bezafibrate (3.1 µg/L ppb).

**Landfills.** PPCPs can be introduced to landfills both directly via domestic and industrial routes and indirectly via sewage sludge. Holm et al. (21) first reported leachates carrying pharmaceuticals from a landfill. Large amounts of numerous sulfonamides (antibiotics) and barbiturates from domestic waste and from a pharmaceutical manufacturer were disposed of at a Danish landfill over a 45-year period. High concentrations (ppm) of many of these drugs were found in leachates close to the landfill; these compounds even accounted for 5% of the total nonvolatile organic carbon found in the leachate. It was also found that the concentrations dropped off dramatically tens of meters down gradient, presumably a result of microbial attenuation.

**Drinking water.** Few pharmaceuticals have been identified in domestic drinking water, probably because of the dearth of monitoring efforts and because the required detection limits are too low for current routine analytical technology. In Germany, however, clofibric acid concentrations up to 165 ng/L (22) and 270 ng/L (23) have been measured in tap water; the presumed source was from recharged groundwaters that had been contaminated by sewage. Stumpf et al. (24) and Ternes et al. (9) found several pharmaceuticals in German drinking water in the lower nanograms-per-liter range, with a maximum of 70 ng/L for clofibric acid. Additionally, these investigators found that diclofenac, bezafibrate, phenazone, and carbamazepine were sometimes present. In the majority of the samples analyzed, however, no drugs were observed. The investigations performed to date therefore indicate that contamination of drinking water does not appear to be a general problem. Depending on the water source for drinking water production, however, certain facilities can experience contamination, especially if the source is polluted groundwater and if polishing technology does not remove the PPCP [e.g., see Heberer et al. (23) and Stumpf et al. (24)]. A major unaddressed issue regarding human health is the long-term effects of ingesting via potable waters very low, subtherapeutic doses of numerous pharmaceuticals multiple times a day for many decades. This concern especially relates to infants, fetuses, and people suffering from certain enzyme deficiencies (which can even be food-induced, e.g., microsomal oxidase inhibition by grapefruit juice).

**Drinking water regulations.** Regulations designed to safeguard receiving waters (from sewage treatment) and drinking water were historically designed to protect the consumer

from the obvious threats of pathogens, widely used industrial chemicals, and certain radionuclides. The treatment processes used by state-of-the-art POTWs evolved from the need to remove these limited sets of contaminants. In areas of water scarcity, the future will see more and more reuse of treated sewage to meet drinking water needs. This will impose a severe burden on water providers to ensure that all chemical contaminants have been removed to the greatest extent possible. It will also require the ability to identify as many of the plethora of potential chemicals in the upgraded water as possible.

According to the National Research Council (NRC) (25), more than two dozen major U.S. utilities release so much effluent to receiving waters that when the natural flows are low, the discharged waste composes 50% of the eventual flow. Any residual, unidentified contaminants therefore are diluted 2-fold at best. In more densely populated countries (e.g., United Kingdom), this figure can rise as high as 90% of flow during times of low rainfall (26).

**Domestic animals.** Whereas the concentration of many drugs is greatly attenuated through sewage treatment plants, larger quantities of many pharmaceuticals are used in various animal husbandry operations, especially CAFOs. With aquaculture in particular, which uses many anti-infectives and anesthetics, the chance for introduction into the immediate environment is greatly enhanced, and the possibility of direct human consumption of therapeutic quantities is correspondingly heightened. Even in the United States the extremely large populations of pet dogs and cats are recipients of numerous drugs (e.g., tranquilizers and antidepressants)—some prescribed by veterinarians and others intended for their owners' use as pet owners sometimes administer medications to their pets to test off-label uses for themselves. PPCPs (both veterinary drugs and OTC products) used with terrestrial domestic animals can be dispersed into the environment through the same routes as those PPCPs used for humans, with the added major route of run-off/leaching of on-ground fecal material.

### Significant Aspects of Ecotoxicology

**Shortcomings of effluent toxicologic screening: comprehensive chemical characterization cannot be replaced—chemical characterization and toxicity screening must be better integrated.** There are two debates in the realm of ecotoxicology, both of which have ramifications with respect to performing ecologic risk assessments (ERAs) for PPCPs. The first is the relevance of purposefully simplified, defined-species toxicity tests to predicting/extrapolating pollutant impacts

on the more highly organized and complex structural/functional levels of communities or ecosystems (processes) [see Boudou and Ribeyre (27)]; this is truer for PPCPs than for pesticides, as the former were generally never designed to have any intended effects on wildlife and therefore any knowledge as to what types of effects to look for is clearly more limited. Can changes in a complex system be predicted from knowledge of a small subset of the underlying components? The second is the question of whether it is necessary to know the spectrum of possible physiologic effects, given a multitude of organisms, or possible mechanisms (modes) of action before looking for and ascribing causation to changes at the population level and higher. Considering this, one can only pose at this time the rhetorical question as to whether the risk posed by the presence of pollutants in complex waste streams (e.g., PPCPs in STW effluents) can be detected/quantified by the use of current toxicity screening tests never designed to embrace the spectrum of end points (some exquisitely subtle) that may be involved. The most conservative approach would be one that captures the coordinated use of toxicity-directed screening and chemistry-directed characterization, feeding the results of each to the other, to better reveal the nature of any stressors.

Although most pharmaceuticals are designed to target specific metabolic pathways in humans and domestic animals, they can have numerous often unknown effects on metabolic systems of nontarget organisms, especially invertebrates. Although many nontarget organisms share certain receptors with humans, effects on nontarget organisms are usually unknown. It is important to recognize that for many drugs, their specific modes of action even in the target species are also unknown. For these drugs, it is impossible to predict what effects they might have on nontarget organisms. Without knowing the mode of action, coupled with not knowing the possible receptors, it is impossible to design rational toxicity testing procedures at the organism level. In the final analysis, given the vast array of mechanisms of drug action and side effects, the total number of different toxicity tests possibly required to screen the effluent from a typical STW could be impractically large. The current batteries of acute/chronic toxicity tests used for ecotoxicity screening merely supply gross indications of directly measurable acute effects. Even if the known mode of action is considered when selecting ecotoxicity tests [as recommended by Henschel et al. (28)], this falsely presupposes that other modes of action are nonexistent or nominal.

Regulatory agencies only in the last few years have recognized that pharmaceuticals should be screened to determine possible

effects on nontarget species. The world's first requirement for ecotoxicity testing as a prerequisite for registration of a pharmaceutical was established in 1995 and first implemented in Germany according to European Union (EU) guideline 92/18 EWG for veterinary pharmaceuticals. For a more in-depth discussion, see Henschel et al. (28), and for a general discussion of the issues in aquatic ecotoxicology, see Boudou and Ribeyre (27).

Screening waste effluent and receiving waters for toxicologic effects can at best be only partially effective because the range of physiologic effects is too broad and relevant to a vast array of aquatic and terrestrial organisms, spanning everything from acute toxicity to very subtle behavioral or genetic changes, of which the consequences are not immediately manifested and can be detected only over long periods of time. There are too many scenarios to discuss in an efficient, comprehensive manner. The complexity of accounting for a wide range of mechanisms of action was made clear in the National Research Council's recent report on endocrine disruptors (29). Although for this class of pollutants the number of modes of action is very large, they represent only a subset of those for PPCPs in general. Quite clearly, any successful toxicity-directed methodology for risk assessment of complex effluents or environmental samples should also make use of a well-developed knowledge of the chemical constituents and their modes of action; current approaches are not yet sufficiently comprehensive. The complexity of this task is further magnified when the effect and necessarily its mode of action have not even been elucidated.

A popular means of attempting to identify the toxic constituents, using toxicity identification and evaluation, in complex waste such as sewage effluent is that of bioassay-directed fractionation screening (30), in which chemical separation techniques yield distinct chemical-class fractions that are then subjected to toxicity testing. Those fractions showing activity against the selected end point are then subjected to chemical identification protocols. Even if one accepts the limitations of selecting appropriate end points (the number with environmental relevance would be enormous), this extremely time-consuming approach would miss any combined effects, whether antagonistic or synergistic, of multiple chemicals. Direct, rigorous chemical characterization of problematic samples clearly must play a role in the identification of toxicants that might elicit previously unrealized toxic effects in nontarget organisms.

**The trend toward optically pure pharmaceuticals: fewer side effects and lower concentrations.** Most pharmaceuticals are racemic mixtures. For a specific optically active drug, it is theorized that only one of its

optical isomers is responsible for the desired physiologic, therapeutic effects; the other isomers are at best inactive, or even worse, responsible for many of the untoward side-effects that most drugs display. A recent trend in the pharmaceutical industry, and now supported by the FDA, is to produce only the optically pure therapeutic isomer (31). This has the potential to not only lessen side effects, but for some drugs, the total dosage can be lessened by at least 50%. This could help in reducing the burden on sewage treatment plants. The significance of the industry's switch to optically pure isomers is that the number of metabolites and other SRSs entering the environment will be reduced at least by half, and the use of the active ingredient will also be reduced by at least 50% because the potency will effectively increase. At the same time, however, the trend of pharmaceuticals toward higher potency will increase the difficulty of environmental monitoring because the required detection levels will be lowered.

**Synergistic effects and potentiation: the potentially critical role of "multixenobiotic resistance."** The biochemical interactions of drugs, often leading to adverse effects, is well known in humans. Little is known, however, of this interplay in aquatic organisms. The following is provided as an example of the complex potential for adverse drug interactions (one actually leading to increased exposure), as it also illuminates the interwoven pathways that ultimately determine exposure. Mostly during this decade, a new mechanism for elimination of xenobiotics from organisms (first observed in tumor cells) has been elucidated—multidrug transporters. This excretory system, also called multixenobiotic transporters, comprises proteins that facilitate the active export of potentially toxic substances, primarily those of moderate lipophilicity, from inside cells. The best-known transporters are the *P*-glycoproteinlike (*Pgp*) transporters (*P* is for permeability altering), or P170 (because of their 170-kDa mass), which have been well characterized in mammals, especially tumor cells, and bacteria.

The toxicologic significance of these nonspecific transporters in maintaining a first line of defense against exposure to multiple xenobiotics in aquatic species has been largely pioneered and reviewed by Epel (32) and by Kurelec and co-workers (33–35); this system confers what has become known as multidrug or multixenobiotic resistance (MDR or MXR). Although these protective proteins have not been found in all aquatic organisms, they have been found in many, especially filter feeders and bottom dwellers (those having potentially high exposures to xenobiotics). This extrusion pump protein system, and possibly others as yet identified, facilitate the

removal and prevent the entrance of those compounds not metabolized or conjugated. They seem to have nonspecific recognition, working for many pesticides, drugs, and natural toxins alike. The action of this transporter system can be inhibited by certain substances such as verapamil ( $\alpha$ -[3-[[2-(3,4-dimethoxyphenyl)ethyl]methylamino]propyl]-3,4-dimethoxy- $\alpha$ -(1-methylethyl)benzene acetonitrile), a cardiac drug—calcium ion influx inhibitor—that directly binds to the active site of *Pgp*. Exposure to verapamil at micromolar concentrations and lower greatly increases the toxicity of a number of drugs or other xenobiotics for many aquatic organisms (32), as the toxicant cannot be readily removed from the exposed organism; exposure time is thereby lengthened by its intracellular accumulation. This elimination system does not function for highly hydrophobic substances (e.g., DDT, polychlorinated biphenyls [PCBs]) and as such might play a more critical role in eliciting effects from exposure to the less hydrophobic PPCPs. Xenobiotics may irreversibly inhibit (cyclosporine A inhibits ATPase), competitively inhibit (verapamil, quinidine, reserpine at low concentrations or high concentrations of general lipophilic compounds such as petroleum oil), or indirectly modulate (e.g., via phosphorylation) MXR regulation or expression (staurosporine inhibits protein kinase C *Pgp* regulator), resulting in its reversal.

The slow escalation, by induction or genetic enrichment, of MXR occurrence and activity among aquatic organisms can give the illusion that the toxicity potential in the aquatic environment is stable or even decreasing when in reality it may be increasing. The introduction of a new substance, at what would normally be a no-effect level, that disrupts the activity of MXR could thereby lead to a profound cascade of unanticipated and unaccounted-for toxic events—a phenomenon akin to what is being termed toxicant-induced loss of tolerance in humans. Organisms in an aquatic environment that have adapted via MXR to certain levels of a suite of toxicants could experience widespread interspecies toxic events should their MXR be inhibited by the addition of a single agent capable of inhibiting MXR, even one that ordinarily would elicit no effect on its own. The resulting effects would be inexplicable if considered solely on the basis of exposure to the new toxicant.

Little is known about which xenobiotics have activity within this relatively newly identified class of chemicals, referred to as chemosensitizers, or their frequency of occurrence in the environment. Smital and Kurelec (35), however, showed that unidentified agents in samples from polluted waters enhance the accumulation of aromatic amines

in clams, mussels, snails, and sponges. Some examples of known MXR inhibitors (34,35), other than verapamil include trifluoroperazine (Stelazine, which is a calmodulin antagonist and an antipsychotic tranquilizer), reserpine (antihypertensive), quinidine and amiodarone (anti-arrhythmics), cyclosporins (immunosuppressants), anthracyclines (non-cytotoxic cytotoxin analogs), and progesterone (steroid); some natural substances such as agent(s) in grapefruit juice are also known to inhibit the P-glycoprotein system (36).

### Environmental Studies on Pharmaceuticals

Given the numbers and quantities of pharmaceuticals manufactured and used throughout the world and that many of these chemicals are designed to have profound physiologic effects, comparatively little research has been published on their occurrence in the environment, effects on nontarget organisms, or assessment of environmental impact. Literally thousands of distinct drugs are approved for use throughout the world. Many of these are manufactured and used in very large quantities. The world's combined literature (the vast majority of these studies have originated in Europe, but the issue applies equally worldwide) has addressed only a very small percentage of these compounds, and the huge array of associated metabolites and other transformation products, many of which undoubtedly have strong physiologic activity, simply compounds the magnitude of the problem.

When drugs are detected in the environment (e.g., surface waters), their concentrations are generally in the ng/L– $\mu$ g/L (ppt–ppb) range. Although parts-per-billion concentrations may not pose much acute risk, it is completely unknown whether other receptors in nontarget organisms are sensitive. It must also be recognized that even though individual concentrations of any drug might be low, the combined concentrations from drugs sharing a common mechanism of action could be substantial. Exposures in the aquatic environment are of particular concern, since aquatic organisms (as opposed to those spending at least some time in terrestrial settings) are subject to continual, unabated life-cycle exposures. This is a highly significant consideration for pharmaceuticals (or bioactive metabolites) that are refractory to structural transformations and are continually introduced into surface waters from sewage treatment plants. Moreover, the polar, non-volatile nature of most drugs prevents their escape from the aquatic realm. Effectively, even PPCPs with relatively short environmental half-lives assume the qualities of highly persistent pollutants because they are continually replenished by infusion to the aquatic environment from STWs.

### Environmental Occurrences

(Note: The names, structures, Chemical Abstracts Service Registry Numbers, and some of the data for environmental occurrences cited in this paper are summarized in Table 2.) Probably the first report of a prescription drug in the environment (sewage treatment effluent) was made over 20 years ago by Garrison et al. (37), who reported clofibric acid (the active metabolite from the lipid regulators clofibrate, etofibrate, and theofibrate) concentrations of 0.8–2.0  $\mu$ g/L in raw sewage and activated sludge effluent. They also found the ubiquitous caffeine and nicotine to be the two most prevalent compounds in influent and effluent from activated sludge, but they did not find the parent clofibrate in any sample. In parallel, Hignite and Azarnoff (38) reported salicylic acid and clofibric acid in the influent and effluent from a Kansas City, Missouri, municipal sewage treatment plant [the history of clofibric acid identified in the environment has been summarized by Stan and Heberer (39)]. Clofibric acid was routinely detected in the effluent of this Missouri STW at an average effluent rate of 2.1 kg/day; over a 10-month period its loading remained in the tight range of 0.76–2.92 kg/day. Similarly, salicylic acid, a hydrolytic metabolite of aspirin, averaged 8.6 kg/day but ranged more widely from 0.55 to 28.7 kg/day. Stan and Heberer also observed that the influent concentrations of clofibric acid were only 20% higher than the effluent concentrations, showing that this chemical resisted removal by the STW. In contrast, for salicylic acid, the influent concentration was about an order of magnitude higher than the effluent, showing more efficient removal.

It therefore was clearly recognized over 20 years ago that the continual, daily introduction of kilogram quantities of drugs from a given STW into receiving waters could result in sustained concentrations with the potential to lead to exposures in aquatic organisms. Little more transpired in the literature, however, during the next 15 years, although clofibric acid continued to appear in a number of monitoring efforts that did not target PPCPs. The most complete investigation to date of the occurrence of pharmaceuticals in both the influent and effluent of POTWs (and also in various surface waters) has been published by Ternes (18).

The distribution of pharmaceuticals is a large function of their production volumes, which can rival those for many pesticides. There are thousands of registered drugs that are dispensed both as prescriptions and OTC; this makes it difficult to estimate usage rates for those pharmaceuticals sold via both routes (e.g., many analgesics). In Germany, roughly 2,900 drugs are permitted in human medicine alone (18). Many countries dispense

drugs in the absence of prescriptions. The two primary sources for release into the environment are from human and veterinary applications. Ternes (18) states that at least for lipid regulators and NSAIDs the source is almost entirely from human usage, as these drugs are infrequently (or never) used in veterinary medicine. In general, the literature shows that most pharmaceuticals, when detected, are present in surface waters in a concentration range of 1 ng/L–1  $\mu$ g/L. To put this in perspective, Richardson and Bowron (40) state that 1,000 kg of a chemical distributed evenly among the rivers in England and Wales would yield a concentration of about 0.1  $\mu$ g/L. Many pharmaceuticals are consumed in amounts far exceeding this; in fact, Richardson and Bowron report 170 pharmaceuticals used annually in excess of this amount.

### Terrestrial and Atmospheric Exposure

*Minor route for PPCPs in contrast to pesticides.* The majority of PPCPs introduced into the environment is undoubtedly into aquatic systems; the terrestrial environment receives only a secondary input. Although the primary source for terrestrial exposure is probably from disposal of biosolids from STWs and from animal wastes both applied to land and stored in open-air pits (waste lagoons), other possible sources for veterinary pharmaceuticals result from animal dips and direct deposition of dung from medicated animals. To date, most attention has been focused on the application of animal wastes to land, primarily because of the suspected introduction of antibiotics and nutrients, not because of PPCPs other than veterinary antibiotics, which are used in comparatively smaller amounts. It should be noted, however, that even though the introduction of veterinary antibiotics into the environment, both terrestrial and aquatic, via animal wastes is widely discussed, the topic has experienced little attention in the peer-reviewed literature (41,42). This topic also relates directly to the human health concern of introducing/promoting antibiotic resistance in bacteria, both native to and introduced into the environment [see section on "Antibiotics" and Williams and Heymann (43)].

The polar nature of the majority of drugs/metabolites leads to facile leaching from land disposal areas into groundwater or wet weather runoff into surface waters. The remainder (largely those designed to pass the blood–brain barrier) have lipophilic character, rendering them prone to bioconcentration from consumption of water or bioaccumulation from consumption of tissue.

Dung-feeding fauna such as birds, beetles, worms, flies, and microorganisms could experience immediate exposure to excreted terrestrial veterinary pharmaceuticals and metabolites.

These organisms in turn could suffer effects themselves from exposure or, alternatively, pass on accumulated residues further up the food chain. All other routes of dispersal to other environmental compartments also play roles, with the distinct exception of direct volatilization, because nearly all PPCPs, with the exception of medical gases and fragrances in contrast with many other anthropogenic compounds are polar or otherwise nonvolatile. The major volatile pharmaceuticals are the inhalable anesthetics (e.g., halothane); these hydrofluoroalkanes are known to oxidize in the atmosphere, like the conventional hydro[chloro]fluorocarbon refrigerants, to yield the highly persistent, toxic, and ubiquitous product trifluoroacetic acid (TFA). This source of TFA is believed to be minor (44).

## Drug Classes and Environmental Occurrences

### Hormones/Mimics

**Potential for receptor interaction may not be rare.** An excellent overview of hormone systems is given by the Endocrine Disruptor Screening and Testing Advisory Committee (EDSTAC) (45). Steroids were the first physiologic compounds to be reported in sewage effluent (46–49) and as such were the first pharmaceuticals to capture the attention of environmental scientists. Estrogenic drugs, primarily synthetic xenoestrogens, are used extensively in estrogen-replacement therapy and in oral contraceptives, in veterinary medicine for growth enhancement, and in athletic performance enhancement. A special issue of *The Science of the Total Environment* (50) is devoted to drugs (especially hormones) as pollutants in the aquatic environment.

Although the synthetic oral contraceptive (17 $\alpha$ -ethynylestradiol) occurs generally at low concentration (< 7 ng/L) in POTW effluent, it is still suspected, in combination with the steroidal estrogens 17 $\beta$ -estradiol and estrone [30], of causing vitellogenin production (feminization) in male fish. Feminization is a phenomenon first observed for fish in sewage treatment lagoons in the mid-1980s (26). An overview of pharmaceutical hormones in the environment is presented by Arcand-Hoy et al. (51). The estrogenic activity of various waters (from sewage to drinking water) has been shown to vary dramatically, spanning six orders of magnitude. Some other widely used synthetic hormone modulators include Proscar/Propecia (finasteride: an androgen hormone inhibitor) and various thyroxine analogs (thyroid hormones); nothing is known of the environmental fates of these compounds. In general, the lipophilicity of these hormones is sufficiently great that at least a large portion are removed via sorptive processes in sewage treatment (52,53) and

therefore partition to the sludge; but even the low concentrations that remain in the effluents may be capable of exerting physiologic effects in aquatic biota.

In addition to these synthetic steroids and xenoestrogens is a suite of naturally occurring estrogen hormones, for example, phytoestrogens such as the complex series of leguminous isoflavonoids, including genistein, daidzein, and glycitein in soy. Further complicating the picture are a host of newly suspected endocrine-disrupting compounds (EDCs), more recently referred to as hormonally active agents (HAAs) by the NRC (29), which have gained attention in the last few years, primarily as a result of the 1996 publication *Our Stolen Future* by Colburn et al. (54). These inadvertent EDCs include such commonly recognized industrial pollutants and products as halogenated dioxins/furans, PCBs, organohalogen pesticides, phthalates, and bisphenol A.

The issue of screening many of the major commercial chemicals (over 87,000 total) for endocrine disruption potential has been formalized with the creation of the EDSTAC, which had been charged by the U.S. EPA with the task of implementing a screening and testing program by August 1999 (45). The Chemical Manufacturers Association (CMA) also has launched an intensive health effects investigation for over 3,000 high-volume chemicals (called the Health and Environmental Research Initiative) (55). It is significant, however, that pharmaceuticals are not specifically targeted by the EDSTAC (or the CMA) in its tiered screening program that focuses on pesticides, commodity chemicals, naturally occurring nonsteroidal estrogens (phytoestrogens and mycotoxins), food additives, cosmetics, nutritional supplements, and representative mixtures (for possible synergistic effects). Even though the strategy gives top priority to “chemicals with widespread exposure at the national level” (55), PPCPs are not specially targeted. It is also significant that the screening strategy will initially focus on only the three primary hormone systems—estrogen, androgen, and thyroid—hormone systems of relatively unknown importance to invertebrates (45).

A controversial hypothesis regarding multiple toxicants (sharing a common mode of action), when each is present at a low level, is that of synergism. Evidence of synergism among estrogenic mimics (where the effect can be elicited at orders-of-magnitude lower concentration than predicted by additive action) was reported by Arnold et al. (56). This study created much controversy by purporting synergistic action of low-level chemical mixtures. Subsequent studies by Gaido et al. (57) and others rebutted this hypothesis. They did not find any evidence of synergism in mixtures of mild estrogenic pollutants.

McLachlan (58) later withdrew the article by Arnold et al. (56), but the issue has not been put to rest, especially given Arnold's other publications on this subject including Arnold et al. (59) and references cited therein. Another controversial issue is that of inverted (U-shaped) dose–response in which toxicity diminution tracks lower concentrations down to a certain level, at which point toxicity again increases. Consequently, higher dose effects might not be useful in predicting the type or magnitude of effects from lower doses (29). This unresolved issue, coupled with the controversy of whether toxicity thresholds necessarily exist, could severely impede EDSTAC's ability to reach its objective because the concentration ranges that must be investigated would be greatly expanded.

**Low molecular weight nonpeptidyl molecules can mimic hormones.** Another subclass of hormonelike substances includes those that are being purposefully designed to mimic the activity of therapeutically significant hormones. A long-sought objective has been to obviate the need for hormone-replacement therapy (e.g., insulin) by designing small synthetic (nonpeptidyl) molecules that mimic the hormone's effect yet can be ingested orally, taken up by the gut, and remain stable for a sufficiently long period of time in the blood. The first report of a “designer” hormone mimic (60,61), a polybenzimidazole that activates the receptor for a cytokine that regulates white blood cell production, perhaps portends the advent of many synthetic hormone mimics in therapeutic medicine. If the finding can be generalized, it could mean that the possible routes of hormone disruption by simple molecules could extend beyond that of the estrogen/androgen system.

With the exception of estrogenic mimics, the possibility of disrupting the activity of proteinaceous hormones by lower molecular weight anthropogenic chemicals has been held in low regard. This view has been based on the fact that a relatively large, complex proteinaceous molecule (the hormone) neatly “fits” within the complex three-dimensional domain of its target receptor, whereas in contrast a much smaller nonproteinaceous molecule would have little to offer in terms of recognition specificity. It has been believed that the complexity of larger proteins such as insulin was required to enable recognition by the corresponding receptors; smaller compounds simply did not convey enough three-dimensional information to have high-binding constants for one or multiple receptors.

The report by Tian et al. (60) demonstrates for the first time that a relatively small nonpeptide molecule can bind to a receptor normally dedicated to a proteinaceous hormone. While this has high therapeutic significance

(this research might catalyze concerted attempts to develop the first protein-mimicking and therefore perhaps hormone-mimicking low molecular weight drugs), it also alludes to the possibility that existing anthropogenic compounds might have a greater chance of interacting with hormone receptors than was previously believed. Although the synthetic substance was three to six orders of magnitude less potent, its ability to bind to the receptor was undisputed (in the mouse *in vitro* and, more importantly, *in vivo*).

### Antibiotics

**In addition to pathogen resistance, genotoxicity may be a concern.** A large body of literature exists on antibiotics in the environment. Veterinary and animal husbandry, especially aquaculture, usage plays a major role in their introduction into the environment. In one study of hospital effluent, fluoroquinolones was the chemical class contributing the major portion to overall DNA toxicity (62); ciprofloxacin, for example, was identified at 3–87 µg/L. Hirsch et al. (41) analyzed German STW effluents and groundwaters/surface waters for 18 antibiotics representing macrolides, sulfonamides, penicillins, and tetracyclines. Although the penicillins (susceptible to hydrolysis) and the tetracyclines (can precipitate with calcium and similar cations) were not found, the others were detected in the microgram per liter range. Indeed, the rampant, widespread (and sometimes indiscriminate) use of antibiotics, coupled with their subsequent release into the environment, is the leading proposed cause of accelerated/spreading resistance among bacterial pathogens, which is exacerbated by the fact that resistance is maintained even in the absence of continued selective pressure (an irreversible occurrence). Sufficiently high concentrations could also have acute effects on bacteria. Such exposures could easily lead to altered microbial community structures in nature and thereby affect the higher food chain. Their use in aquaculture results in eventual human consumption. For a discussion of promotion of antibiotic resistance, see the policy article by Witte (63). Hartmann et al. (62) propose that genotoxicity in hospital effluent may result more from antibiotics than from antineoplastics.

Recently, a number of stream surveys documented the significant prevalence of native bacteria that display resistance to a wide array of antibiotics including vancomycin (64). Isolates from wild geese near Chicago, Illinois, are reported to be resistant to ampicillin, tetracycline, penicillin, and erythromycin (65). All these reports could simply indicate that the natural occurrence of antibiotic resistance in native bacterial populations is much higher than expected or that these bacteria are being selected for by the

uncontrolled release of antibiotics into the environment. If the latter is true then, excluding the significance of antibiotics themselves in the environment, their occurrence can be viewed as marking or indicating the possible presence of other PPCPs.

### Blood Lipid Regulators

#### **Fibrates—high usage. Fibric acid metabolites—ubiquitous, persistent pollutants.**

Clofibrac acid was the first prescription drug (actually an SRS) reported in a sewage effluent (37,39), and it continues to be one of the most frequently reported PPCPs in monitoring studies. Clofibrac acid (2-[4]-chlorophenoxy-2-methyl propanoic acid), the active metabolite from a series of widely used blood lipid regulators, and which also happens to be structurally related to the phenylalkanoic acid herbicide mecoprop (the methylphenoxy structural analog), has captured much attention from investigators in Europe. Stan et al. (22) first reported clofibrac acid in Berlin tap water at concentrations between 10 and 165 ng/L. Heberer and Stan (66) found clofibrac acid at levels up to 4 µg/L in groundwater under a sewage treatment farm; they also found clofibrac acid concentrations up to 270 ng/L in drinking water samples. They concluded that it is not removed by sewage/water treatment processes.

Buser et al. (67) report finding clofibrac acid in various Swiss waters ranging from rural to urban lakes. Concentrations ranged from 1–9 ng/L (ppt), whereas the parallel concentrations for mecoprop were higher at 8–45 ng/L; little of either compound was found in a relatively remote mountain lake, indicating no atmospheric deposition. Because this drug is not manufactured in Switzerland, its route of introduction into the environment had to be through medical use and subsequent excretion/disposal. Although these concentrations are very low, they are significant in that they are similar to the concentrations found for any of the conventional ubiquitous and persistent pollutants, sometimes referred to as persistent organic pollutants (POPs) or persistent bioaccumulative toxicants (PBTs) such as lindane [see Jones and de Voogt (68) for an overview]. In one of the lakes studied, Buser et al. (67) calculated steady-state amounts of clofibrac acid to be roughly 19 kg (with export and import amounts balancing each other). Perhaps more significantly, they also found amounts of clofibrac acid up to 7.8 ng/L in the North Sea; the parallel concentrations of mecoprop in the same North Sea samples were lower, up to only 2.7 ng/L, indicating that mecoprop was less persistent than clofibrac acid.

Stumpf et al. (24) and Ternes (18) reported bezafibrate, gemfibrozil, and clofibrac/fenofibrac acids in river waters at the

nanogram per liter level. Stumpf et al. (69) reported that the removal efficiencies from Brazilian STWs for clofibrac/fenofibrac acids, bezafibrate, and gemfibrozil ranged from only 6–50%, verifying extremely limited degradation for these compounds. This chemical class is ubiquitous because the daily human dosages are generally high (grams per day). Buser et al. (67) concluded that the concentrations seen in urban Swiss and German rivers, coupled with essentially the same concentrations in the North Sea, lead to an annual input of 50–100 tons of clofibrac acid into the North Sea. The concentration of clofibrac acid in the environment is more a function of dilution than of degradation. Clofibrac acid is the most widely and routinely reported drug found in open waters. It would be expected that its occurrence in other parts of the world would parallel these studies.

### Nonopioid Analgesics/Nonsteroidal Anti-Inflammatory Drugs

Stumpf et al. (24) were the first to identify diclofenac, ibuprofen, acetylsalicylic acid, and ketoprofen in sewage and river water. Ternes (18) reported levels of diclofenac, indometacin, ibuprofen, naproxen, ketoprofen, and phenazone in POTW effluent exceeding 1 µg/L; all these except ketoprofen were also found in surface waters at concentrations severalfold lower. In another study, Ternes et al. (70) reported average concentrations of acetylsalicylic acid generally less than 1 µg/L in most POTW effluents as well as less than 0.14 µg/L in rivers. They also reported salicylic acid concentrations of 54 µg/L in POTW influents, with two other acetylsalicylic acid metabolites, gentisic acid (4.6 µg/L) and *o*-hydroxyhippuric acid (6.8 µg/L). While low levels (0.5 µg/L) of salicylic acid appeared in the effluents, no detectable amounts of the metabolites could be found. Ternes et al. (70) also found naproxen in all POTW effluents examined and in river waters (–0.05–0.4 µg/L); two veterinary NSAIDs, meclofenamic and tofenamic acids, were not detectable in any river sample. In their screening of waters in Berlin, Heberer et al. (23) found that the most prevalent drugs, other than clofibrac acid, were the NSAIDs diclofenac, ibuprofen, and propyphenazone. In groundwater from a drinking water plant, they found diclofenac, ibuprofen, and *N*-methylphenacetin (from phenacetin) (23). In the influent to Swiss STWs, Buser et al. (71) found diclofenac at concentrations of 0.5–1.8 µg/L, whereas the concentrations in the respective effluents were only moderately reduced (at most 50%). In the receiving water (Swiss lakes/rivers), they found 11–310 ng/L but only 1–12 ng/L in exiting waters. They concluded that photolysis was the major cause of the diminished concentrations of diclofenac in surface waters

(71). Buser et al. (72) showed that ibuprofen, while present in influents at 1–3.3 µg/L, was easily degraded to yield low effluent concentrations (nanograms/liter) in contrast to other NSAIDs, which were more refractory. This study is also one of the few that examined the enantiomeric selectivity in the degradation of the parent optical isomers as well as the production of metabolites.

### Beta-Blockers/ $\beta_2$ -Sympathomimetics

Hirsch et al. (73) and Ternes (18) identified the beta-blockers metoprolol and propranolol, with lesser amounts of betaxolol, bisoprolol, and nadolol, in POTW effluent. Only metoprolol and propranolol were found in surface waters at concentrations just above the limit of detection. The  $\beta_2$ -sympathomimetics (bronchodilators) terbutalin and salbutamol (albuterol in the United States), but rarely clenbuterol and fenoterol, were detected in POTW effluent and only at low concentrations, less than 0.2 µg/L. They were rarely seen in surface waters. It may be significant to note that medications delivered by inhalers could result in portions of the dose being deposited externally because of improper dosing technique.

Fenfluramine (*N*-ethyl- $\alpha$ -methyl-3-[trifluoromethyl] benzene ethanamine hydrochloride), known as Pondimin in addition to other brand names, is a sympathomimetic amine, which was used as a popular diet (anorectic) drug and was removed from the U.S. market in 1998 by the FDA because of heart valve damage. Although no one has looked for fenfluramine in sewage, it is known to enhance the release of serotonin (3-(2-aminoethyl)indol-5-ol or 5-hydroxytryptamine creatinine sulfate [5-HT]); in the crayfish, 5-HT in turn triggers release of ovary-stimulating hormone, resulting in larger oocytes with enhanced amounts of vitellin (consequences unknown) (74). Similarly, in fiddler crabs, fenfluramine at a dose of 125 nmol stimulates (through 5-HT) the production of gonad-stimulating hormone, which accelerates testicular maturation (75).

### Antidepressants/Obsessive-Compulsive Regulators

*Subtle but possibly profound effects on nontarget [aquatic] species.* Selective serotonin reuptake inhibitors (SSRIs) are a major class of widely prescribed antidepressants that includes Prozac, Zoloft, Luvox, and Paxil. These drugs enjoy widespread and heavy use. One of the few series of studies reported in the literature that addresses the effects of drugs on nontarget organisms (albeit not the intent of the studies) was performed in a quest for more effective spawning inducers for economically important bivalves (76). Fong's studies and those of other physiologists studying the

function of serotonin in a wide array of aquatic creatures could prove highly significant in any discussion of the importance of low levels of pharmaceuticals in the environment. Fong's work is perhaps the most significant to date for showing the potential for dramatic physiologic effects on nontarget species (in this case invertebrates) by low (ppb) concentrations of pharmaceuticals.

Serotonin is a biogenic amine common in both vertebrate and invertebrate nervous systems. SSRIs increase serotonin neurotransmission by inhibiting its reuptake at the synapses by inhibiting the transporter enzymes. In addition to playing a key role in mammalian neurotransmission, serotonin is involved in a wide array of physiologic regulatory roles in molluscs, among most other creatures. For bivalves, reproductive functions including spawning, oocyte maturation, and parturition are regulated by serotonin, (76). Serotonin controls a wide spectrum of additional behaviors and reflexes in molluscs, including heartbeat rhythm, feeding/biting, swimming motor patterns, beating of cilia, and induction of larval metamorphosis (77). It also stimulates release of various neurohormones in crustaceans (hyperglycemic hormone, red pigment-dispersing hormone, neurodepressing hormone, and molt-inhibiting hormone) and ovarian maturation (78).

It has long been known that serotonin at concentrations of  $10^{-4}$  to  $10^{-3}$  M (–0.18–1.8 g/L) induces spawning in bivalves. Some commercial farmers make use of this by adding serotonin to induce spawning. Fong (76) found that Prozac (fluoxetine) and Luvox (fluvoxamine) are the most potent inducers ever found, eliciting spawning behavior in zebra mussels at aqueous concentrations many orders of magnitude lower than serotonin. Fluoxetine elicited significant spawning in male mussels at concentrations of  $10^{-7}$  M (–150 µg/L); females were an order of magnitude less sensitive at  $10^{-6}$  M. Fluvoxamine was the most potent of the SSRIs, eliciting significant spawning in male mussels, at  $10^{-9}$  M (–0.318 µg/L); females were two orders of magnitude less sensitive, at  $10^{-7}$  M. In males, spawning was complete in the first hour, while females were slower (within 2 hr). Paxil (paroxetine) was the least potent of these three SSRIs, eliciting male spawning, but to a lesser degree, at  $10^{-6}$  M, and having no inducing effect on females at any concentration. It should be noted that Fong states that the evidence is not clear whether these compounds are indeed acting as SSRIs, or via some other mechanism. It is also unknown how these compounds are taken up by molluscs (76).

In another study, Fong et al. (79) showed that fluvoxamine induces significant parturition in fingernail clams at 1 nM; 1 nM fluvoxamine

also potentiated the effect of 10 µM 5-HT by almost 5-fold. Paroxetine was less potent, requiring a concentration of 10 µM to effect significant parturition. In contrast, even at concentrations of 100 µM, fluoxetine displayed no effect, although it was capable at 5 µM of potentiating 5-HT at concentrations that were otherwise subthreshold. It is interesting that the order of potency for inducing parturition in clams differs from the order for induction of spawning in mussels (above). This points to the complexity of considering any approach involving extrapolations from one species to another or from one drug to another within a given class.

In crustaceans, Kulkarni et al. (74) found that fluoxetine significantly potentiates the effect of 5-HT in crayfish, enhancing the release of ovary-stimulating hormone, which results in larger oocytes with enhanced amounts of vitellin; any ecologic consequences of higher vitellin protein levels are unknown. Similarly, in fiddler crabs, fluoxetine at a dose of 125 nmol stimulates (through 5-HT) the production of gonad-stimulating hormone, which accelerates testicular maturation (75).

It is clear that aquatic life can be exquisitely sensitive to at least some of this class of compounds. Although some SSRIs are extremely potent, others have almost no effect, which possibly makes the approach of assessing ecologic risk on a class-by-class basis infeasible.

Concentration of SSRIs plays a complicated role with respect to effects. For example, Couper and Leise (77) found that while injected fluoxetine induced significant metamorphosis in a gastropod,  $10^{-4}$  M induced less metamorphosis than  $10^{-6}$  M. Simple extrapolations of effects from higher concentrations do not necessarily have any relevance to effects at lower concentrations.

The potential for SSRIs to elicit subtle effects on aquatic life is further extended by serotonin reuptake mechanisms that also are a factor in snails and squids (76), particularly in the regulation of aggression (80). Yet another example of a subtle effect that would go unnoticed is the fighting behavior of lobsters, in which serotonin causes behavior reversal by stimulating subordinates to engage in fighting against dominants by reducing their propensity to retreat (80).

### Antiepileptics

*Antiepileptics are ubiquitous and prevalent due to poor STW removal.* Carbamazepine was the drug detected most frequently and in highest concentrations during a study by Ternes (18). This drug was detected in all POTWs and receiving waters, with a maximum concentration of 6.3 µg/L. Ternes hypothesized that the ubiquitous occurrence resulted from the very low removal efficiency from POTWs, which was calculated to be



only 7%. Sacher et al. (81) found carbamazepine levels in the river Rhine in Germany up to 0.90 µg/L and always above 0.1 µg/L.

### Antineoplastics

**Antineoplastics are highly [geno]toxic compounds, primarily from hospitals, with poor removal from STWs.** Antineoplastic agents, antitumor agents primarily used only within hospitals for chemotherapy, are found sporadically and in a range of concentrations, probably because only small amounts are introduced to STWs via domestic sewage because of their long-lived physiologic retention. These compounds act as nonspecific alkylating agents (i.e., specific receptors are not involved) and therefore have the potential to act as either acute or long-felt stressors (mutagens/carcinogens/teratogens/embryotoxins) in any organism. The fact that two oxazaphosphorines, ifosfamide and cyclophosphamide, were found in certain effluents in the low microgram-per-liter range indicates that these highly toxic compounds, which are probably refractory to microbial degradation at POTWs (82), can find their way into the environment. Indeed, Steger-Hartmann et al. (82) found levels of cyclophosphamide in sewage influent from servicing hospitals ranging from undetectable to 143 ng/L; the levels in the effluent reached 17 ng/L.

Additional evidence pointing to the refractory nature of ifosfamide is presented by Kümmerer et al. (83), who found that concentrations of ifosfamide in hospital effluent matched the predicted values of up to 1.91 µg/L; also the concentrations in the influent and effluent of POTWs that serviced chemotherapy hospitals were essentially unchanged (influent/effluent maximum, 43 ng/L; median, 6.5–9.3 ng/L). Kümmerer et al. (83) found ifosfamide to be totally refractory to removal by POTWs and to totally resist alteration during a 2-month bench-scale POTW simulation.

Another class of antineoplastics, the platinates, includes carboplatin and cisplatin. Although the stability of these compounds in sewage systems is unknown, Kümmerer et al. (84) calculated that if they were present in hospital sewage effluents as the intact parent compound, they could be present at daily average concentrations of up to 600 ng/L (on the basis of total platinum). Although the majority of the dose for these compounds is excreted in the urine in the first day, a large amount (~30%) resides in the body and is slowly excreted over a period of years and therefore could be excreted to residential sewage systems. Falter and Wilken (85) showed that while these compounds are difficult to determine analytically, their potential to remain in the aqueous phase after sewage treatment is high.

White and Rasmussen (86), in the most detailed overview to date on the genotoxicity of wastewaters, elaborate that while the genotoxic potency of industrial wastewaters is often the highest, the overall loading of genotoxic compounds to surface waters is far greater, up to several orders of magnitude, from municipal treatment plants. They present a striking correlation between the occurrence of direct-acting mutagens in surface waters and the human population served by the discharging STWs. This correlation points to the activities/metabolism of humans, not industrial activities, as the origin for these mutagens. A number of possible sources for the mutagens are discussed, an obvious one of which is antineoplastic drugs.

These data point to antineoplastics as a class of drugs of potential concern for environmental effects, not just for their acute toxicity but perhaps more for their ability to effect subtle genetic changes, the cumulative impact of which over time can lead to more profound ecologic change. Hospitals are the major source of genotoxic drugs. POTWs that service hospitals, especially multiple hospitals, are likely candidates for releasing these chemicals into surface waters.

### Impotence Drugs

**This class of drugs displays widespread use, new modes of action, and unknown effects on nontarget organisms.** Even though a number of drugs from various chemical classes have been used over the years for treating impotence, the emergence of Viagra (sildenafil citrate) has focused tremendous attention on this market. The significance of this therapeutic class of drugs, with new ones awaiting FDA approval, is that they all tend to have distinct modes of action, most of which differ from those of traditional drugs. While potential effects on wildlife are totally unknown, the fact that Viagra, for example, works by inhibiting a phosphodiesterase responsible for regulating the concentration of cyclic guanosine monophosphate, which indirectly relaxes muscles and increases blood flow (87), gives cause for concern regarding the disruption of this common phosphodiesterase in unintended target species. Impotence drugs will prove to have very high usage rates, especially since they are one of the most common drugs available without prescription over the Internet, yielding high potential for environmental exposure and possibly nontarget effects.

### Tranquilizers

**Little is known about possible occurrence of tranquilizers.** Ternes (18) reported diazepam in almost half of the POTWs but only in low concentrations of less than 0.04 µg/L; it could not be detected in surface waters. Genicola (88) reports diazepam in the groundwater

from a monitoring well at a Superfund site near Atlantic City, New Jersey. Concentrations were approximately 10–40 µg/L and probably originated in a landfill in which pharmaceutical manufacturers disposed of chemicals.

### Retinoids

**High usage rates and profound activity in amphibians lends cause for concern.** Retinoids, low molecular weight lipophilic derivatives of vitamin A, can have profound effects upon the development of various embryonic systems (89), especially amphibians in which retinoic acid receptors have been hypothesized to play a role in frog deformities. Although naturally occurring, retinoids have been used for a number of years for a wide array of medical conditions including skin disorders (e.g., Accutane [isotretinoin] for acne), antiaging treatments (e.g., Retin-A [tretinoin] for skin wrinkles), and cancer (e.g., Vesanoid [tretinoin] for leukemia). Isotretinoin (13-cis-retinoic acid) is related to both retinoic acid and retinol (vitamin A). Tretinoin is among the top 200 prescribed drugs in the United States. Methoprene, an insecticidal synthetic retinoic acid mimic, is photolabile and yields numerous photo-products, some of which also elicit strong retinoic acid activity (90). Although retinoic acids would also be expected to be photolabile (and therefore not persistent), their products may also still possess receptor activity.

### Diagnostic Contrast Media

**Diagnostic contrast media have very high usage rates, display considerable persistence, show no evidence for mineralization, and have low physiologic activity.** Detailed X-ray images of soft tissues are routinely captured by the use of contrast media. Some of the more widely used members of contrast media are highly substituted and sterically hindered amidated, iodinated aromatics such as diatrizoate and iopromide (91), which are used worldwide at annual rates exceeding 3,000 tons. Kalsch (91) found these compounds to be quite resistant to transformation in STWs and in river waters. When transformations were effected, they merely terminated with unidentified resistant metabolites. Ternes et al. (92) recently reported significant amounts of iopromide in rivers.

In municipal STW effluents, Ternes et al. (92) found concentrations as high as 15 µg/L (iopamidol) and 11 µg/L (iopromide). In an STW close to Frankfurt/Main, they found two other contrast agents, diatrizoate and iomeprol, at concentrations up to 8.7 µg/L, as well as iothalamic acid and ioxithalamic acid in the nanogram-per-liter range. In rivers and streams, five iodinated diagnostics were repeatedly detected, with median values up to 0.49 µg/L for iopamidol and up to 0.23 µg/L

for diatrizoate. Isolated maximum values above 100 µg/L for diatrizoate indicated that relatively high local concentrations can occur, especially in small streams containing a high percentage of STW discharges. Maximum groundwater concentrations for iodinated contrast agents ranged up to 2.4 µg/L and may well represent a worst case with respect to occurrence of pharmaceuticals in native waters. In Germany alone, individual contrast agents can experience annual usage rates of 100 tonnes. Such high usage, coupled with inefficient human metabolism (95% unmetabolized) and ineffective elimination of iodinated contrast agents by STWs, can lead to very high environmental accumulations and persistence. Despite these negative attributes, contrast agents have no bioaccumulation potential and low toxicity (93); Steger-Hartmann et al. (93) also found no acute toxicity for bacteria (*Vibrio fischeri*), algae (*Scenedesmus subspicatus*), crustaceans (*Daphnia*), and fish (*Danio rerio*, *Leuciscus idus melanotus*) exposed to no more than 10 g/L of iohexol, iotrolan, diatrizoate, or iopromide.

### Personal Care Products in the Environment

For the purposes of this review, personal care products are defined as chemicals marketed for direct use by the consumer (excluding OTC medication with documented physiologic effects) and having intended end uses primarily on the human body (products not intended for ingestion, with the exception of food supplements). In general, these chemicals are directed at altering odor, appearance, touch, or taste while not displaying significant biochemical activity. Most of these chemicals are used as the active ingredients or preservatives in cosmetics, toiletries, or fragrances. They are not used for treatment of disease, but some may be intended to prevent diseases (e.g., sunscreen agents). In contrast to drugs, almost no attention has been given to the environmental fate or effects of personal care products—the focus has traditionally been on the effects from intended use on human health. Many of these substances are used in very large quantities frequently more than recommended.

**Table 3.** Personal care products produced in Germany (1993).

Product category	Tons produced
Bath additives	162,300
Shampoos, hair tonic	103,900
Skin care products	75,500
Hair sprays, setting lotions, hair dyes	71,000
Oral hygiene products	69,300
Soaps	62,600
Sun screens	7,900
Perfumes, aftershaves	6,600
Total	559,100

Personal care products differ from pharmaceuticals in that large amounts can be directly introduced to the environment. For example, these products can be released directly into recreational waters or volatilized into the air (e.g., musks). Because of this direct release they can bypass possible degradation in POTWs. Also, in contrast to pharmaceuticals, less is known about the effects of this broad and diverse class of chemicals on nontarget organisms, especially aquatic organisms. Data are also limited on the unexpected effects on humans. For example, common sunscreen ingredients, 2-phenylbenzimidazole-5-sulfonic acid and 2-phenylbenzimidazole, can effect DNA breakage when exposed to UV-B (94).

The quantities of personal care products produced commercially can be very large. For example, in Germany alone the combined annual output for eight separate categories has been estimated (95) at 559,000 tons for 1993 (Table 3). A few examples are given below of common personal care products that are ubiquitous pollutants and that may possess substantial bioactivity.

*Fragrances (musks) are ubiquitous, persistent, bioaccumulative pollutants that are sometimes highly toxic; amino musk transformation products are toxicologically significant.* Synthetic musks comprise a series of structurally similar chemicals (which emulate the odor but not the structure of the expensive, natural product from the Asian musk deer) used in a broad spectrum of fragranced consumer items, both as fragrance and as fixative. Included are the older, synthetic nitro musks (e.g., ambrette, musk ketone, musk xylene, and the lesser known musks moskene and tibetene) and a variety of newer, synthetic polycyclic musks that are best known by their individual trade names or acronyms. The polycyclic musks (substituted indanes and tetralins) are the major musks used today, accounting for almost two-thirds of worldwide production) and especially the inexpensive nitro musks (nitrated aromatics accounting for about one-third of worldwide production) are used in nearly every commercial fragrance formulation (cosmetics, detergents, toiletries) and most other personal care products with fragrance; they are also used as food additives and in cigarettes and fish baits (96).

The nitro musks are under scrutiny in a number of countries because of their persistence and possible adverse environmental impacts and therefore are beginning to be phased out in some countries. Musk xylol has proved carcinogenic in a rodent bioassay and is significantly absorbed through human skin; from exposure to combined sources, a person could absorb 240 µg/day (97). The human lipid concentration of various musks parallels that of other bioaccumulative pollutants such as PCBs (98). Worldwide production of

synthetic musks in 1988 was 7000 tons (96); worldwide production for nitro musks in 1993 was 1,000 tons, two-thirds of which were musk xylene (99).

Synthetic musks first began to be identified in environmental samples almost 20 years ago (100,101). By 1981, Yamagishi et al. (100) had identified musk xylene and musk ketone in gold fish (*Carassius auratus langsdorfi*) in Japanese rivers and not much later (101) in river water, sewage, marine mussels (*Mytilus edulis*), and oysters (*Crassostrea gigas*). Yamagishi's studies comprised the first comprehensive monitoring efforts, identifying musk xylene and musk ketone in freshwater fish, marine shellfish, river water, and STW waters. Musk xylene was found in all samples, and musk ketone was found in 80% of the 74 samples analyzed. Concentrations in STW effluents ranged from 25 to 36 ng/L for musk xylene and from 140 to 410 ng/L for musk ketone. Concentrations of musk xylene in fish muscle were in the tens of parts per billion, whereas those for musk ketone were less than 10 µg/kg, with highest values occurring in fish downstream of STWs. In contrast, for shellfish the concentrations were lower, between 1 and 5.3 µg/kg, presumably because of their lower lipid contents. In river water, musk xylene occurred in all samples, whether upstream or downstream of STWs and ranged between 1 and 23 ng/L; those of musk ketone were generally in the same range, but in distinct contrast they were not detectable in upstream samples.

Geyer et al. (102) have published an excellent review on residues of nitro musk fragrances in fish and mussels as well as in breast milk and human lipids and the current ecotoxicologic and toxicologic knowledge for these personal care products. Residues of musk xylene and musk ketone found in the fillet of freshwater fish (e.g., pike, eel, bass, Zander, rainbow trout) from rivers of North Germany were between 10 and 350 µg/kg lipid and 10 and 380 µg/kg lipid for musk xylene and musk ketone, respectively. In mussels (*Mytilus edulis*) 10–30 µg/kg lipid of both fragrances were detected. In human breast milk from German women, musk xylene and musk ketone were detected between 10 and 240 mg/kg lipid (102). Recently, the literature has a number of additional publications from Europe, especially Germany and Switzerland. Rimkus et al. (103) give a brief overview of the occurrence of musks in the environment. Kafferlein et al. (99) and Geyer et al. (102) published the most thorough reviews to date on the occurrence (in the environment and in personal care products), transformation, and toxicology of the ubiquitous musk xylene; these reviews summarize many more occurrence studies (for musk xylene) than mentioned here.

Musks are refractory to biodegradation (other than reduction of nitro musks to amino derivatives), which explains why they have been measured in water bodies throughout the world (96). They also are very lipophilic [octanol-water partition coefficients are similar to those for DDT and hexachlorocyclohexane (104)] and therefore can bioconcentrate/bioaccumulate (102,103). Concern has been expressed regarding developmental toxicity in aquatic organisms. Musk ambrette (2,6-dinitro-3-methoxy-4-*tert*-butyl toluene) may play a role in damaging the nervous system (105).

Draisci et al. (106) examined freshwater fish in Italy and identified two of five targeted polycyclic musks in most fish samples; a hexahydro-hexamethylcyclopental-benzopyran (HHCB, trade name Galaxolide) and an acetylhexamethyltetralin (AHTN, trade name Tonalide) were identified at levels ranging from less than 4 ng/g (ppb) to 105 ng/g in fish muscle tissue. In the Swiss river Glatt, Müller et al. (98) identified Galaxolide, Tonalide, and Celestolide (ADBI, 4-acetyl-6-*tert*-butyl-1,1-dimethylindane) at concentrations of 136, 75, and 3.2 ng/L, respectively; they also found the nitro musks tibetene, ambrette, moskene, ketone, and xylene at concentrations of 0.04, <0.03, 0.08, 8.3, and 0.62 ng/L, respectively. Eschke et al. (cited in 107) identified Galaxolide, Tonalide, and Celestolide in the fatty tissue of bream and perch from the Ruhr River, Germany, at average concentrations between 2.5 and 4.6 mg/kg (ppm), illustrating the extreme bioaccumulation potential for these compounds. Recently, Heberer et al. (108) investigated the contamination of surface waters in Berlin, Germany, (and vicinity) receiving high percentages of treated sewage and found maximum concentrations above 10 µg/L for the polycyclic musks Galaxolide, Tonalide, and Celestolide.

Winkler et al. (104) measured musks in 31 particulate matter and water samples from the Elbe River, Germany. In all particulate matter samples, concentrations for musk ketone were 4–22 ng/g, for Galaxolide 148–736 ng/g, and for Tonalide 194–770 ng/g; Celestolide was found at concentrations of 4–43 ng/g in 23 of the particulate matter samples. The values for the three most prevalent musks were within the same order of magnitude as those for 15 polycyclic aromatic hydrocarbons (PAHs) and exceeded those for 14 common polychlorinated organic pollutants (only hexachlorobiphenyl [HCB] and *p,p'*-DDT were of similar concentration). Also found in all the 31 water samples were musk ketone (2–10 ng/L), Galaxolide (36–152 ng/L), and Tonalide (24–88 ng/L); Celestolide was found only at 2–8 ng/L. These higher values exceeded those for all the polychlorinated organics and the PAHs. The

occurrences of individual musks are sometimes correlated as a result of their use as mixtures in commercial products. In Germany, the nitro musks are being replaced by the polycyclic musks, therefore resulting in lower concentrations for musk ketone (104).

It is not surprising that musks have been detected in air. Kallenborn et al. (109) detected three polycyclic musks and two nitro musks in Norwegian outdoor air samples. The polycyclic musks were more prevalent. Concentrations of all these musks ranged from low picograms per cubic meter to hundreds of picograms per cubic meter. The most common was the polycyclic musk Galaxolide, but the relative ratios among the musks are a function of usage (which varies among countries) and photolability.

Although the significance of the aquatic toxicity of the nitro and polycyclic musks is debatable [genotoxicity from the polycyclics seems to not be a concern] (110), the aminobenzene (reduced) versions of the nitro musks can be highly toxic; these reduced derivatives are undoubtedly created under the anaerobic conditions of sewage sludge digestion. Behecti et al. (111) tested the acute toxicity of four reduced analogs of musk xylene on *Daphnia magna*. The *p*-aminodinitro compound exhibited the most toxicity of the four, with extremely low median effective concentration (EC<sub>50</sub>) values averaging 0.25 µg/L (0.25 ppb).

Recently, the amino transformation products of nitro musks were identified in sewage treatment effluent and in the Elbe River, Germany. Gatermann et al. (96) identified musk xylene and musk ketone together with their amino derivatives 4- and 2-amino musk xylenes and 2-amino musk ketone. In sewage treatment influent, the concentrations of musk xylene and musk ketone were 150 and 550 ng/L, respectively. In the effluent, their concentrations dropped to 10 and 6 ng/L, respectively. In contrast, although the amino derivatives could not be detected in the influent, their concentrations in the effluents dramatically increased, showing extensive transformation of the parent nitro musks: 2-amino musk xylene (10 ng/L), 4-amino musk xylene (34 ng/L), and 2-amino musk ketone (250 ng/L). It was concluded that the amino derivatives could be expected in sewage effluent at concentrations more than an order of magnitude higher than the parent nitro musks. In the Elbe, 4-amino musk xylene was found at higher concentrations (1–9 ng/L) than the parent compound.

Given that the amino nitro musk transformation products *a*) are more water soluble than the parent musks, *b*) still have significant octanol-water partition coefficients (high bioconcentration potential), and *c*) are more toxic than the parent nitro musks, more

attention should be focused on these compounds. Because synthetic musks are ubiquitous, used in large quantities, introduced into the environment almost exclusively via treated sewage effluent, and are persistent and bioconcentratable, they are prime candidates for monitoring in both water and biota as indicators for the presence of other PPCPs. Their analysis, especially in biota, has been thoroughly discussed by Gatermann et al. (96) and by Rimkus et al. (103).

### Preservatives

Parabens (alkyl-*p*-hydroxybenzoates) are one of the most widely and heavily used suites of antimicrobial preservatives in cosmetics (skin creams, tanning lotions, etc.), toiletries, pharmaceuticals, and even foodstuffs (up to 0.1% wt/wt). Although the acute toxicity of these compounds is very low, Routledge et al. (112) report that these compounds (methyl through butyl homologs) display weak estrogenic activity in several assays. Although the risk from dermal application in humans is unknown, the probable continual introduction of these benzoates into sewage treatment systems and directly to recreational waters from the skin leads to the question of risk to aquatic organisms. Butylparaben showed the most competitive binding to the rat estrogen receptor at concentrations one to two orders of magnitude higher than that of nonylphenol and showed estrogenic activity in a yeast estrogen screen at 10<sup>-6</sup> M.

### Disinfectants/Antiseptics

Triclosan (Irgasan DP 300, a chlorinated diphenyl ether: 2,4,4'-trichloro-2'-hydroxydiphenyl ether) is an antiseptic agent that has been widely used for almost 30 years in a vast array of consumer products. Its use as a preservative and disinfectant continues to grow; for example, it is incorporated at < 1% in Colgate's Total toothpaste, the first toothpaste approved by the FDA to fight gingivitis. While triclosan is registered with the U.S. EPA as a pesticide, it is freely available OTC. Triclosan's use in commercial products includes footwear (in hosiery and insoles of shoes called Odor-Eaters), hospital handsoap, acne creams (e.g., Clearasil), and rather recently as a slow-release product called Microban, which is incorporated into a wide variety of plastic products from children's toys to kitchen utensils such as cutting boards. Many of these uses can result in direct discharge of triclosan to sewage systems, and as such this compound can find its way into receiving waters depending on its resistance to microbial degradation. Okumura and Nishikawa (113) found traces of triclosan ranging from 0.05 to 0.15 µg/L in water. Although triclosan has long been regarded as a biocide, a toxicant having a wide-ranging,

nonspecific mechanism(s) of action—in this case gross membrane disruption, McMurry et al. (114) report that triclosan is rather an antibacterial having particular enzymatic targets (lipid synthesis). As such, bacteria could develop resistance to triclosan. As with all antibiotics in the environment, this could lead to development of resistance and change in microbial community structure (diversity).

A wide array of disinfectants are used in rather large amounts not just by hospitals, but also by households and livestock breeders. These compounds are often substituted phenolics as well as others such as triclosan. Biphenylol, 4-chlorocresol, chlorophene, bromophene, 4-chloroxylenol, and tetrabromo-*o*-cresol (70) are some of the active ingredients, at percentage volumes of < 1–20%. A survey of 49 STWs in Germany (70) routinely found biphenylol and chlorophene in both influents, up to 2.6 µg/L for biphenylol and up to 0.71 µg/L for chlorophene, and effluents. The removal of chlorophene from the effluent was less extensive than for biphenylol, with surface waters having concentrations similar to that of the effluents.

### Sunscreen Agents

The occurrence of sunscreen agents (UV filters) in the German lake Meerfelder Maar was investigated by Nagtegaal et al. (115). The combined concentrations of six sunscreen agents (SSAs) identified in perch (*Perca fluviatilis*) in the summer of 1991 were as high as 2.0 mg/kg lipid and in roach (*Rutilus rutilus* L) in the summer of 1993, as high as 0.50 mg/kg lipid. Methylbenzylidene camphor (MBC) was detected in roach from three other German lakes. These lipophilic SSAs seem to occur widely in fish from small lakes used for recreational swimming. Both fish species had body burdens of SSA on par with PCBs and DDT. The bioaccumulation factor, calculated as quotient of the MBC concentration in the whole fish (21 µg/kg) versus that in the water (0.004 µg/L), exceeded 5,200, indicating high lipophilicity. The fact that SSAs (e.g., 2-hydroxy-4-methoxybenzophenone [oxybenzone] and 2-ethylhexyl-4-methoxycinnamate) can be detected in human breast milk [16 and 417 ng/g lipid, respectively (116)] shows the potential for dermal absorption and bioconcentration in aquatic species. No data have been published on newer SSAs such as avobenzene (1-[4-(1,1-dimethylethyl)phenyl]-3(4-methoxyphenyl)-1,3-propanedione).

### Nutraceuticals/Herbal Remedies

During the last several years, the popularity of nutritional supplements was codified by the creation of a new term for the subclass of highly bioactive food supplements called nutraceuticals (117), also referred to as nutriceuticals. Nutraceuticals are a rapidly

growing commercial class of bioactive compounds, usually botanicals, intended as supplements to the diet. Nutraceuticals and many herbal remedies can have potent physiologic effects. These are a mainstay of alternative medicine and have enjoyed explosive growth in use in the United States and other countries during the last decade. Many are used as food supplements that have either proven or hypothesized biologic activity but are not classified as drugs by the FDA, primarily because a given botanical usually has not one but an array of distinct compounds whose assemblage elicits the putative effect and because these arrays cannot be easily standardized. As such they are not regulated and are available OTC (heavily promoted via the Internet). Even in those cases in which the natural product is identical to a prescription pharmaceutical (e.g., the Chinese red-yeast product Cholestin newly introduced to the United States contains lovastatin, an active ingredient in the approved prescription drug Mevacor used to lower cholesterol levels), a recent ruling (118) prevented the FDA from regulation.

With the accelerating inverted age structure of our society, coupled with the U.S. 1994 Dietary Supplement Health and Education Act (DSHEA) (119) (which eases regulations on the introduction and marketing of supplements), the use of nutraceuticals could greatly escalate. The significance of dietary supplements in the United States was epitomized by the creation of the Office of Dietary Supplements (ODS) via the DSHEA in 1995 under the National Institutes of Health (NIH) (119). The ODS maintains a searchable database (International Bibliographic Information on Dietary Supplements [IBIDS]) of published scientific literature on dietary supplements (120). The NIH was also mandated to create the National Center for Complementary and Alternative Medicine (NCCAM, 121) to “facilitate the evaluation of alternative medical treatment modalities to determine their effectiveness.”

Although these substances are readily available OTC, albeit in poorly characterized/standardized forms, an effort is underway to patent various nutraceuticals by standardizing the extracts and thereby making them available only by prescription. This effort is being pioneered by PharmaPrint, Inc. (Irvine, CA) (122), which has applied to FDA for various investigational new drug applications. The patenting of hundreds of multiple-molecule nutraceuticals for therapeutic purposes could lead to more widespread use of these substances.

As an example, a recent addition to this class is a substance called huperzine A, an alkaloid extracted from a Chinese moss,

which has been documented to improve memory. It is therefore experiencing strong demand for treating Alzheimer's disease and has captured the attention of those who follow the nutraceutical market because of its true pharmaceutical qualities. The significance of this particular compound is that it possesses acute biologic activity as a cholinesterase inhibitor identical to that of organophosphorus and carbamate insecticides. It is so effective that the medical community is concerned about its abuse/misuse, especially since it is legal. While huperzine A, and alkaloids in general (compounds with heterocyclic nitrogen, proton-accepting group, and strong bioactivity), are naturally occurring compounds, their susceptibility to biodegradation in STWs or in open waters is unknown. This is the case for almost all nutraceuticals.

Another example is Kava, which is prepared from the root of *Piper methysticum*, long used throughout the South Pacific because of its mild narcotic effect among a host of other effects. The active ingredients in Kava are believed to be a suite of lipophilic lactones comprising substituted  $\alpha$ -pyrones (methysticin, kavain, yangonin, and others) (123). These compounds display a host of effects in humans, but nothing is known about their effects on other organisms or fate in STWs.

There are countless other nutraceuticals, both new and ancient, experiencing vigorous consumption. These few examples illustrate the unknowns regarding whether these compounds are being excreted, surviving sewage treatment, and then eliciting effects on aquatic organisms. Nutraceuticals and herbal remedies would have the same potential fate in the environment as pharmaceuticals, with the added dimension that their usage rates could be much higher, as they are readily available and taken without the controls of prescription medication. Because these compounds are natural products, however, they would be expected to be more easily biodegraded.

Although the argument can be made that naturally occurring compounds would not pose an ecologic risk, this ignores that *a*) the concentrations of these compounds in effluents could be higher than they are in the environment in which they occur naturally, and *b*) many of these substances/mixtures come only from isolated parts of the world (e.g., Kava, huperzine), and their use/dispersal in other parts of the world would essentially make them anthropogenic. The use of these compounds serves to redistribute their normal occurrence in the environment, and even though they might be naturally occurring, this promotes exposure to organisms that normally would never occur.

## Nontarget Organism Exposure to and Effects from PPCPs

### Environmental Exposure

*Persistence is not critical if the source is constant, leading to perpetual aquatic exposure.* Many PPCP ingredients seem to have considerable persistence in the environment; blood lipid regulators and musks are examples. Although environmental persistence usually is a major determinant of exposure in the environment, for pollutants that are used on a continual basis and are introduced to the environment through STWs, the supply is continually replenished. In the absence of very short half-lives, exposures even to nonpersistent compounds could be significant. This is especially true for aquatic organisms, which are captives of their environment and perpetually exposed.

Seasonal variations in pharmaceutical infusion to surface waters from sewage treatment effluents may not be a factor. No seasonal variations were noted for musks; weekly variations of severalfold in concentration have been noted in the River Elbe in Germany (104).

### Effects on Nontarget Organisms

*Although acute data are lacking, subtle effects might be the major concern.* The potential effects of PPCPs on nontarget species, especially on aquatic organisms, are mostly unknown. No concerted research effort has focused on the ecotoxicology of PPCPs. Some isolated studies, however, have included pharmaceuticals in various toxicity assays relevant to aquatic life. One study in particular was begun in Scandinavia.

In 1989 under the direction of the Scandinavian Society of Cell Toxicology (SSCT), an international toxicologic evaluation study was initiated—Multicenter Evaluation of In Vitro Cytotoxicity Tests (MEIC). While the main purpose of the MEIC was to thoroughly investigate a select list of chemicals for evaluating human toxicity by employing and benchmarking a wide battery of *in vitro* and *in vivo* tests, the MEIC generated a range of ecotoxicologic data for various species including fish, amphibians, crustaceans, and single-cell eukaryotes. Of the 50 selected chemicals, at least 18 were drugs. Although limited and focusing on the more common, obvious end points, the MEIC data sets are some of the only available that catalog the effects of certain drugs on aquatic life. The MEIC was concluded in 1996, and the data are still being evaluated. A database of SSCT's MEIC cytotoxicity data can be found at the web site for the Cytotoxicology Laboratory, Uppsala, Sweden (CTLU) (124).

An example of one of the MEIC studies relevant to aquatic effects is the study by Lilius et al. (125) that presents effects data for

all 50 MEIC chemicals on two species of the crustacean *Daphnia*. Of 50 MEIC reference chemicals, 18 pharmaceuticals had immobilization EC<sub>50</sub> values ranging from below 0.01 mM (e.g., 0.0037 for amitriptyline and 0.0017 for thioridazine) to less than 10 mM (e.g., 6 mM for phenobarbital and 8.2 mM for aspirin); 9 of the 18 drugs had EC<sub>50</sub> values below 0.1 mM. In comparison, the EC<sub>50</sub> values for a range of common industrial chemicals and pesticides were generally in the same millimolar range (e.g., phenol, 0.078; 2,4-dichlorophenoxyacetic acid, 0.65; nicotine, 0.023; potassium cyanide, 0.0086; lindane, 0.005). Another MEIC study (126) reports data from a number of tests relevant to the aquatic environment (several crustaceans, rotifer, and *Microtox*). Using the OECD (Organisation for Economic Co-operation and Development) Test Guideline 202, Part II (chronic *Daphnia* reproduction), Kalbfus and Kopf (127) reported results for clofibrate and salicylic acid: for clofibrate, NOEC (no observed effect concentration) of 10 µg/L and EC<sub>50</sub> (24 hr) of 106 µg/L, whereas the NOEC for salicylic acid was three orders of magnitude higher (10 mg/L); the clofibrate NOEC concentrations for algae and luminescence bacteria ranged between 5 and 40 µg/L, whereas those for salicylic acid were between 15 and 60 mg/L.

Acute toxicity, the major type of end point investigated with nontarget species, is only one of many possible ecotoxicologic end points of concern. Investigation of multi-generational life-cycle effects is almost entirely lacking for any nontarget species. This is surprising, as in the aquatic environment—given that exposures to PPCPs would be more constant than episodic—organisms are exposed for their entire life cycles. Perhaps more important, however, subtle behavioral modifications or genetic alterations have the potential to lead to profound long-term ecologic effects for which it could prove extremely difficult to pinpoint the cause. Acute toxicity should not be a primary concern because it can be so easily detected, and mitigation measures can be designed in a timely manner. Rather, concern should be directed toward effects occurring undetected that can lead to long-term adverse impacts, which in turn are perceived simply as natural variation or evolution. This concern is amplified compared with that for pesticides, as nearly nothing is known about the effects of PPCPs on aquatic or terrestrial life. We do know, however, that these substances have the potential to be profoundly bioactive through a constellation of different modes of action. The toxicologic data that exist for nontarget species are almost exclusively focused on antibiotics and woefully lacking for fish [see Halling-Sørensen et al. (5) for a

tabulated summary of effects data]. The studies of Fong et al. (76,79), presented earlier under "Antidepressants," are the best examples to date of subtle effects resulting from low concentrations. Another overlooked issue regarding effects is that of organisms from lower trophic levels whose presence is critical to community homeostasis.

### Subtle Effects—Beyond Our Notice

*Acute toxicity is only a small part of a larger puzzle.* Many drugs that are used successfully to modify the behavior of humans could have unforeseen effects on nontarget organisms. These effects could be so subtle that their consequences may be imperceptible but nevertheless profound when elicited over long periods of time. For example, subtle behavioral effects could be responsible for changing any number of attributes for a particular species, resulting in changes over numerous generations that would otherwise be deemed part of normal adaption/evolution. Many drugs even have unpredicted or unanticipated effects in humans. Such unforeseen biologic effects could prove even more profound and unexpected with nontarget species.

The ability to elicit numerous subtle effects across a wide range of species is embodied by no single class of pharmaceuticals better than SSRIs. This class of drugs shows the potential for PPCPs to elicit a constellation of effects that would be hard to detect in natural settings, or to tease apart from what would otherwise be considered normal behavior.

Perhaps the most important concern regarding the exposure of aquatic and terrestrial organisms to PPCPs is that the effects could be so subtle that they would escape any effort to detect them, with the result that imperceptible changes could accumulate until they had a significant impact—not necessarily on the individual organisms but rather on the population or community, perhaps after generations of change. Subtle effects via regulation by any of the countless pathways/networks of signal transduction in aquatic species can range from modification or reversal of attraction and boldness to avoidance and shyness affecting all behavior characteristics spanning the range from feeding to mating to directional sensing (e.g., chemotaxis, gravitaxis). On the surface, the result would simply be attributed to natural adaptation or confounded by other natural changes.

Kurelec (128) has formalized a concept, genotoxic disease syndrome (GDS), that embodies the idea of nonobvious change as effected in the aquatic environment, especially as exacerbated by compromising the activity of MXR defense systems. As proposed, GDS is seen as the gradual accumulation of a wide spectrum of toxic events, none of which alone

results in an obvious outcome. The cumulative effects fall under the general, diffuse, ill-defined alteration/impairment/inhibition of enzyme systems, protein turnover, metabolism, and cytotoxic repair, leading to reduced fitness, gradual degeneration/atrophy of tissues/organs, reduced growth, accelerated aging, impaired immunologic systems, impaired reproduction, higher incidence of disease, and impaired adaptation/survival/succession.

Kurelec (33) reported another example of behavioral alteration. The MXR inhibitor verapamil when added to polluted river waters (at a level of ~1 ppm that normally did not affect fish) elicited dramatic avoidance attempts by the fish, escalating to the point of frantic escape attempts when 2-aminoanthracene was also added at a concentration 0.53  $\mu$ M, which would otherwise not elicit a response.

The significance of subtle and cumulative impacts is only beginning to be recognized by environmental toxicologists. Weiss (129) wrote that just as the predicted rise in "crack babies" whose pregnant mothers used cocaine was never observed—rather only a small but significant 3% reduction in IQ was observed—the same subtle effects are very likely occurring from environmental toxicants. Weiss points out that the effects on humans by lead (Pb), methyl mercury, PCBs, and endocrine disruptors in general probably are manifested in almost undetectable changes, and these may accumulate over time to yield truly profound changes that would not be distinguishable from natural events. The specter of subtle, cumulative effects could make current toxicity-directed screening largely useless in any effort to test waste effluents for toxicologic end points.

Abnormal behavior can masquerade as seemingly normal deviation within a natural statistical variation. Change can occur so slowly that it appears to result from natural events, with no reason to presume artificial causation. It is difficult to connect the issues of cause and ultimate effect, in part because of the ambiguous and subjective nature of subtle effects, but especially when these effects are confounded as aggregations of numerous, unrelated interactions. Weiss (130) points out that statistical tests based on means and *p* values are incapable of detecting the very subtle changes that low-level toxicants can impart. Slight shifts within the statistical distributions of any particular characteristic expressed among a population are not necessarily reflected by statistics of the mean/median; these changes may well be more obvious, however, when considering an individual organism rather than looking across a population.

Another aspect to gradual, undetected changes relates to community composition/organization, especially in microorganisms, in which the composition of species can be

affected over time (i.e., successional effects) as a result of the presence of anthropogenic chemicals (131). The pressure for succession can result from adverse effects (e.g., toxicity preferentially to one species over another) or from conferring an advantage to one species (e.g., use of the toxicant as a nutrient source). Succession of community structure tends to occur on such a long time line that cause and effect issues are rarely considered.

A number of toxicity testing approaches have been developed over the years, some of which employ less obvious end points for the nonspecific detection of the presence, but not the identities, of toxicants. One of the best known rapid approaches uses the reduction in light output of a bioluminescent bacterium (e.g., *Microtox*) (132). Another more recent example (133) automatically detects changes in the gravitaxis of a unicellular flagellate—indeed, a subtle end point in itself and one that would probably go unnoticed and undoubtedly lead to die off. Such approaches are badly needed for detecting changes in toxicant concentrations in wastewater effluents and for directing subsequent chemical characterization to identify the putative toxicants.

## Environmental Assessment

Although various levels of prospective ERA via standardized tests are required in the United States and Europe as part of the drug registration process (see "Approach of Regulatory Agencies" below), meaningful effort on this front is simply not possible with the currently limited state of knowledge on environmental fate, transport, and effects of pharmaceuticals; to date, retrospective studies (e.g., ERAs based on environmental survey) are rare. Examples of prospective ecologic assessments can be seen in Henschel et al. (28), who performed these assessments for four high-use pharmaceuticals in Germany: salicylic acid (the main metabolite of acetylsalicylic acid, aspirin), paracetamol (4-acetamidophenol, acetaminophen—analgesic/antipyretic), clofibrac acid (chlorophenoxyisobutyric acid; blood lipid regulator), and methotrexate (4-amino-10-methyl-folic acid; chemotherapy folic acid antagonist that disrupts nucleotide synthesis). Unmetabolized, the loading of these drugs into bodies of water in Germany could be hundreds of tons per year. Henschel et al. (28) found that although all four drugs would have passed traditional ecotoxicity screening, methotrexate would not have passed at least one nonstandard test. This showed that the current guidelines could be underestimating nontarget effects.

The OECD, an intergovernmental organization with representatives from 29 countries, publishes the *OECD Test Guidelines* (134), a collection of methods used to assess the hazards of chemicals and of chemical preparations

such as pesticides and pharmaceuticals. Assessment of methodology for aquatic toxicity of chemicals has been recently reviewed by the OECD (135). General information on ecologic risk assessment is available from the U.S. EPA (136).

The enormous array of pharmaceuticals will continue to diversify and grow as the human genome is mapped. Today, there are about 500 distinct biochemical receptors at which drugs are targeted; U.S. private R&D investment in new pharmaceuticals in 1998 was nearly \$18 billion. The number of targets is expected to increase up to 20-fold (yielding 3,000–10,000 drug targets) in the near future according to the Pharmaceutical Research & Manufacturers Association (137). In 1998, the FDA approved 30 new nonbiologic drugs, one of which was Viagra (138). The FDA Modernization Act of 1997 (139) will also help to accelerate this growth. Most of the new drugs have totally unpublished environmental transformation/fate/effects properties; two examples of highly prescribed new drugs are Viagra (sildenafil: 1-[[3-(6,7-dihydro-1-methyl-7-oxo-3-propyl-1*H*-pyrazolo[4,3-*d*]pyrimidin-5-yl)-4ethoxyphenyl]-sulfonyl]-4-methylpiperazine]) and Propecia/Proscar (finasteride: [5 $\alpha$ -17 $\beta$ -*N*-(1,1-dimethylethyl)-3-oxo-4-azaandrost-1-ene-17-carboxamide; used in treating male baldness and benign prostatic hyperplasia). This explosion in new drugs will severely exacerbate our limited knowledge of drugs in the environment and possibly increase the exposure/effects risks to nontarget organisms. Finally, the current proliferation of web sites offering prescription drugs by mail will only exacerbate the growing use/misuse of a wide array of drugs.

## Approach of Regulatory Agencies

There are only two major activities with respect to managing the release of pharmaceuticals in the environment. One results from the research that has occurred in various European/Scandinavian countries over the last few decades, culminating in guidelines from the EU. The other comes from the FDA. It is important to understand, however, that responsibility in the United States for monitoring drugs in the environment does not currently rest with either the FDA or the U.S. EPA. Few other alternative approaches for assessing ecologic risk posed by pharmaceuticals have been proposed. In one of the more comprehensive approaches, Roembke et al. (8) used the basic ecologic risk assessment approach, upon which the U.S. EPA's current approach (136) is based, to discuss alternatives; in particular, they noted that acute effects testing cannot be relied upon by itself—chronic effects testing is extremely important.

## European Union Activities

**Agency for the Evaluation of Medicinal Products—EMEA/CVMP/055/96.** In the early 1980s, government regulators [e.g., European Commission—Pharmaceuticals and Cosmetics (EEC) which sets rules governing medicinal products in the EU (140)] first showed concern over the release of veterinary pharmaceuticals and their metabolites into the environment and any untoward effects they might have on biota: “the potential risks for the environment resulting from the use of the medicinal product.” Veterinary medicinals were targeted presumably because they were perceived as having a more direct route of introduction to the environment (e.g., fish farms, parasite dips, farm runoff). Only much later has concern been expressed with respect to human drugs, but no regulations/guidelines have been established. An analogous directive has yet to be published by the EU for human pharmaceuticals. A good overview of the approach used by the EEC was published by Henschel et al. (28) and Montforts et al. (42).

In EMEA/CVMP/055/96-final (141), the CVMP (European Committee for Veterinary Medicinal Products) sets forth its final guidance for risk assessments for veterinary medicinal products, excluding biologics. The guidance elaborates on the phased-assessments set forth in EEC Directive 92/18/EEC, where Phase I assesses the potential for release to the environment [derivation of predicted environmental concentrations (PECs)]; for more information on establishing PECs for exposure assessment, see OECD (142). Phase II is broken into two tiers: tier A evaluates possible fate and effects, and tier B (should cause for concern regarding specific biologic species result from tier A findings) looks at effects on specific biota that might receive exposure. ERAs are required for new veterinary drugs. The report must include potential for environmental exposure (considering patterns of use and routes of administration [internal vs external], metabolism/excretion [metabolites representing less than 20% of the dose are excluded from concern], and disposal), fate and effects, and any needed risk management strategies. The guidance mandates the use of worst-case exposure scenarios. EMEA's guidelines are also being developed specifically for environmental impact assessments of veterinary medicinals by EMEA (143).

Effects testing includes algal growth inhibition, fish acute/chronic/bioaccumulation exposure, avian dietary and reproductive, earthworm toxicity, terrestrial plant growth, and activated sludge respiration inhibition. The guidance document seems to recognize the incredible diversity of stressor-receptor possibilities that could

result from pharmaceuticals and their metabolites entering the environment: “there may be considerable variation in receptor specificity/sensitivity between species” (143). This is complicated further by the fact that the mode(s) of action responsible for the desired therapeutic effect of many drugs is poorly understood (sometimes totally unknown). For example, although various modes of action for the disease-modifying agents used to treat rheumatoid arthritis (e.g., methotrexate and hydroxychloroquine, intended for use in chemotherapy and malaria, respectively) have been known for many years, the actual mechanisms by which the symptoms of this particular disease are alleviated are mostly unknown. It would therefore be impossible to forecast what type of effects could be anticipated.

**Biocides.** The new EU Biocide Directive (144) covers the commercialization of biocidal products (e.g., disinfectants), but few of these are used in personal care products. Significantly, however, the Directive emphasizes ecotoxicologic issues (on par with human health issues), including fate and ecologic effects.

## FDA

Concern regarding introduction of pharmaceuticals to the environment in the United States is addressed by the FDA, which requires Environmental Assessments (EAs), as required under National Environmental Policy Act of 1969 (NEPA), and the specifics of which are set forth in “Guidance for Industry: Environmental Assessment of Human Drug and Biologics Application” (145) for all drug applications/actions meeting minimum criteria. As with the EU's approach, concern rests primarily on acute and chronic effects as measured by traditional toxicity tests. Much less concern is expressed for behavioral effects, whether avoidance, breeding, etc. The FDA does, however, recognize “extraordinary circumstances” where there is the potential for serious harm to the environment or for an action to “significantly affect the quality of the human environment” (145). This notion includes not just toxicity to environmental organisms but also “environmental effects other than toxicity, such as lasting effects on ecological dynamics” (145). Clearly this could cover subtle behavioral modifications from which effects accumulate over time/generations, eventually leading to measurable change but unrecognized as such. NEPA [40 CFR 1508.27; also see Appendix C in the FDA document (145)] also defines “significantly” around two issues—“context” and “intensity” (severity of impact). Among the 10 issues with respect to “intensity,” one relates to:

Whether the action is related to other actions with individually insignificant but cumulatively significant impacts.

Significance exists if it is reasonable to anticipate a cumulatively significant impact on the environment.

The FDA's approach is very similar to that of the EU. The FDA requires an EA if the expected environmental concentration (EEC, analogous to the PEC—predicted environmental concentration) at the point of entry to the aquatic environment (the expected introduction concentration [EIC]) exceeds 1 ppb. The EIC is calculated assuming that all the drug product produced for 1 year enters POTWs, that the drug's usage is spread across the country in proportion to the population, and that none of the parent drug is metabolized or transformed; this can be altered if transformation data are available, but if metabolites or other SRSs are present at greater than 10% of the parent level, then toxicology must also be known. Its value is calculated as the product of *a*) kilogram per year of active product produced for use, *b*) reciprocal of the influent to POTWs (liters per day), *c*) reciprocal of 365 days/year, and *d*)  $10^9$   $\mu\text{g}/\text{kg}$ .

Like the EMEA, the FDA uses a tiered approach to determine if regulatory actions are required. This approach centers around assessment factors used to determine when ecotoxicity testing is not needed. An assessment factor is calculated by dividing the test end point (e.g., lethal dose or  $\text{EC}_{50}$ ) by the maximum EEC (which is equal to the greater of EEC or EIC). Lower factors necessitate more extensive testing. Further testing is also necessitated by drugs that bioaccumulate or with SRSs that are more toxic than the parent drug. The weakest aspect to this approach is that the toxicity of the SRSs is assessed from what is known for human toxicology rather than for potential nontarget organisms. Moreover, given the current knowledge of fate, transport, and ecotoxicity of anthropogenic chemicals, there are simply too many unknowns to be able to predict whether a pharmaceutical (or its transformation products) will find its way into the environment at a particular concentration. The FDA also requires EAs for drugs that also occur naturally in the environment if their usage and subsequent discharge (to POTWs, landfills, etc.) will alter the natural, ambient concentration.

A significant shortcoming of either of these two current regulatory approaches to determining ecologic risk results from not taking into account the cumulative (additive/synergistic/antagonistic) impacts of drugs affecting the same receptors. The EEC value for any given drug could easily be exceeded when the cumulative concentrations of like-mode-of-action drugs are considered, especially in those instances where numerous competing drugs are commercially available

in any class. Needless to say, this approach also ignores the possibility of synergistic effects from drugs of other classes.

## Conclusions and Recommendations

This review aims to catalyze a discussion in the environmental science community to determine the significance of PPCPs in the environment and to foster further research efforts, if warranted. The intent is not to outline a tiered approach for a research strategy, but rather to highlight where further research might be needed in each of various areas organized around the "risk paradigm," as set forth by the National Research Council (146) (Table 1). A step-wise strategy to determine if a major research effort needs to be launched would require efforts at establishing the incidence of PPCP occurrence in the environment coupled with parallel determinations of whether effects among a wide spectrum of aquatic organisms can occur at documented concentrations of PPCPs and whether cost-effective modifications of STW operation can dramatically improve removal efficiencies. The authors' personal recommendations and summary of significant conclusions are presented in Table 1. It is hoped that this broad overview presents a wide perspective and proper context for this emerging problem.

## REFERENCES AND NOTES

- Pressley SA. N. Carolina effort seeks to wipe out 999. *Washington Post*, p. A03, Sunday, 25 April 1999. Available: <http://search.washingtonpost.com> [cited 31 August 1999].
- Bonner M, Wristen KG. The National Sewage Report Card (Number Two): Rating the Treatment Methods and Discharges of 21 Canadian Cities. Sierra Legal Defence Fund Report. August 1999. Sierra Legal Defence Fund, Toronto, Ontario, Canada. Available: <http://www.sierralegal.org/reports.htm>
- Pfleger K. Mass Spectral and GC Data of Drugs, Poisons, Pesticides, Pollutants and Their Metabolites. 2nd rev ed. Vols 1-3 published in 1992. Weinheim, Germany:VCH, 1999.
- Pfleger K. Mass Spectral and GC Data of Drugs, Poisons, Pesticides, Pollutants and Their Metabolites. Vol 4. 2nd rev ed and expanded ed. Weinheim, Germany:VCH, 1999.
- Halling-Sørensen B, Nors Nielsen S, Lanzky PF, Ingerslev F, Holten Lützhøft HC, Jørgensen SE. Occurrence fate and effects of pharmaceutical substances in the environment - a review. *Chemosphere* 36(2):357-393 (1998).
- Montague P. Drugs in the Water. *Rachel's Environment & Health Weekly* #614. Available: [gopher://ftp.std.com/70/11/periodicals/rachel/](http://ftp.std.com/70/11/periodicals/rachel/) [cited 3 September 1998].
- Raloff J. Drugged waters—Does it matter that pharmaceuticals are turning up in water supplies? *Science News* 153:187-189 (1998).
- Roembke J, Knacker Th, Stahl-Schmidt-Allner P. Studie über Umweltprobleme im Zusammenhang mit Arzneimitteln. [Study about environmental problems in context with drugs.] F+E Vorhabens Nr. 106 04 121 Umweltbundesamt Berlin. German Report of the Research and Development Project no 106 04 121 of Federal Ministry of Research and Development, Berlin, Germany, 1996.
- Ternes TA, Hirsch R, Stumpf M, Eggert T, Schuppert B, Haberer K. Nachweis und Screening von Arzneimittelrückständen Diagnostika und Antiseptika in der aquatischen Umwelt. [Identification and screening of pharmaceuticals diagnostics and antiseptics in the aquatic environment.] Bundesministerium für Bildung Wissenschaft Forschung Technologie (BMBF)/53170. German Report of the Federal Ministry of Education, Science, Research and Technology, (Ref no 02WU9567/3). Bonn, Germany, 1999.
- Velagaleti R. Behavior of pharmaceutical drugs (human and animal health) in the environment. *Drug Inform J* 31:715-722 (1997).
- Bosch X. Household antibiotic storage [Letter]. *Science* 281:785 (1998).
- Associated Press. Wyoming officials considering Yellowstone pollution fines. *Las Vegas Review-Journal*, p. 9B, 31 October 1998.
- Milstein M. Park sewage systems on the verge of failure internal report states. Available: <http://www.billingsgazette.comain.htm> [cited 8 March 1999].
- James MO. Overview of in vivo metabolism of drugs by aquatic species. *Vet Hum Toxicol* 28(suppl 1):2-8 (1986).
- Guarino AM, Lech JJ. Metabolism disposition and toxicity of drugs and other xenobiotics in aquatic species. *Vet Hum Toxicol* 28(suppl 1):38-44 (1986).
- RxList. The Internet Drug Index (The Top 200 Prescriptions: 1998 US Prescriptions Based on More than 2.4 Billion U.S. Prescriptions). Available via: <http://www.rxlist.com/> [at: <http://www.rxlist.com/top200.htm>] [cited 30 August 1999].
- Forth W, Henschler D, Rummel W, Starke K. Allgemeine und spezielle Pharmakologie und Toxikologie. 6th rev ed. [General and special pharmacology and toxicology.] Mannheim/Leipzig/Wien/Zürich:Wissenschaftsverlag, 1996.
- Ternes TA. Occurrence of drugs in German sewage treatment plants and rivers. *Water Res* 32(11):3245-3260 (1998).
- Rogers HR. Sources behaviour and fate of organic contaminants during sewage treatment and in sewage sludges. *Sci Total Environ* 185:3-26 (1996).
- Learn S. Sewage tradeoff: add overflow, help tributaries. The Oregonian, Thursday 22 April 1999. Available: <http://www.oregonlive.com/news/99/04/st042215.html> [cited 31 August 1999].
- Holm JV, Rügge K, Bjerg PL, Christensen TH. Occurrence and distribution of pharmaceutical organic compounds in the groundwater downgradient of a landfill (Grindsted, Denmark). *Environ Sci Technol* 29(5):1415-1420 (1995).
- Stan H-J, Heberer T, Linkerhäger M. Occurrence of clofibric acid in the aquatic system—is the use in human medical care the source of the contamination of surface ground and drinking water? *Vom Wasser* 83:57-68 (1994).
- Heberer T, Schmidt-Bäumler K, Stan H-J. Occurrence and distribution of organic contaminants in the aquatic system in Berlin. Part I: Drug residues and other polar contaminants in Berlin surface and ground water. *Acta Hydrochim Hydrobiol* 26(5):272-278 (1998).
- Stumpf M, Ternes TA, Haberer K, Seel P, Baumann W. Nachweis von Arzneimittelrückständen in Kläranlagen und Fließgewässern [Determination of drugs in sewage treatment plants and river water]. *Vom Wasser* 86:291-303 (1996).
- J.R. When sewage is recycled for drinking. *Science News* 153:186 (1998).
- Routledge EJ, Sheahan D, Desbrow C, Brighty GC, Waldock M, Sumpter JP. Identification of estrogenic chemicals in STW effluent. 2: In vivo responses in trout and roach. *Environ Sci Technol* 32:1559-1565 (1998).
- Boudou A, Ribeyre F. Aquatic ecotoxicology: from the ecosystem to the cellular and molecular levels. *Environ Health Perspect* 105(suppl 1):21-35 (1997).
- Henschel K-P, Wenzel A, Diedrich M, Fliedner A. Environmental hazard assessment of pharmaceuticals. *Reg Toxicol Pharmacol* 25:220-225 (1997).
- National Research Council. *Hormonally Active Agents in the Environment*. Washington, DC:National Academy Press, 1999. Available: <http://books.nap.edu/books/0309064199/html> [cited 31 August 1999].
- Desbrow C, Routledge EJ, Brighty GC, Sumpter JP, Waldock M. Identification of estrogenic chemicals in STW effluent. 1: Chemical fractionation and in vitro biological screening. *Environ Sci Technol* 32(11):1549 (1998).
- Rogers RS. Sepracor: Skating on thin 'ice'. *Chem Eng News* 30:11-13 (1998).
- Epel D. Use of multidrug transporters as first lines of defense against toxins in aquatic organisms. *Comp Biochem Physiol A* 120:23-28 (1998).
- Kurelec B. The multixenobiotic resistance mechanism in aquatic organisms. *Crit Rev Toxicol* 22(1):23-43 (1992).
- Kurelec B. A new type of hazardous chemical: the chemosensitizers of multixenobiotic resistance. *Environ Health Perspect* 105 (suppl 4):855-60 (1997).
- Smital T, Kurelec B. The chemosensitizers of multixenobiotic resistance mechanism in aquatic invertebrates: a new class of pollutants. *Mutat Res* 399(1):43-53 (1999).
- Soldner A, Christians U, Susanto M, Wacher V, Silverman J, Benet LZ. Grapefruit juice activates P-glycoprotein-mediated drug transport. *Pharmaceut Res* 16(4):478 (1999).
- Garrison AW, Pope JD, Allen FR. GC/MS Analysis of organic compounds in domestic wastewaters. In: *Identification and Analysis of Organic Pollutants in Water* (Keith CH, ed). Ann Arbor, MI:Ann Arbor Science Publishers, 1976:517-556.
- Hignite C, Azamoff DL. Drugs and drug metabolites as environmental contaminants: chlorophenoxisobutyrate and salicylic acid in sewage water effluent. *Life Sci* 20:337-342 (1977).
- Stan HJ, Heberer T. Pharmaceuticals in the aquatic environment. *Analysis Mag* 25(7):20-23 (1997).
- Richardson ML, Bowron JM. The fate of pharmaceutical chemicals in the aquatic environment. *J Pharm Pharmacol* 37:1-12 (1985).
- Hirsch R, Ternes T, Haberer K, Kratz K-L. Occurrence of antibiotics in the aquatic environment. *Sci Total Environ* 225(1-2):109-118 (1999).
- Montforts MHMM, Kalf DF, van Vlaardingen PLA, Linders JBHJ. The exposure assessment for veterinary medicinal products. *Sci Total Environ* 225(1-2):119-133 (1999).
- Williams RJ, Heymann RL. Containment of antibiotic resistance. *Science* 279:1153-1154 (1998).
- Jordan A, Frank H. Trifluoroacetate in the environment. Evidence for sources other than HFC/HCFs. *Environ Sci Technol* 33:522-527 (1999).
- Endocrine Disruptor Screening and Testing Advisory Committee. Final Report. August 1998. Available: <http://www.epa.gov/opptintr/opptendo/finalrpt.htm>. Also EDSTAC Home Page. EPA's Endocrine Disruptor Screening and Testing Advisory Committee (EDSTAC) Home Page. Available: <http://www.epa.gov/opptintr/opptendo/index.htm> [cited 31 August 1999].
- Aheme GV, Briggs R. The relevance of the presence of certain synthetic steroids in the aquatic environment. *J Pharm Pharmacol* 41:735-736 (1989).
- Shore LS, Gurevita M, Shemesh M. Estrogen as an environmental pollutant. *Bull Environ Contam Toxicol* 51:361-66 (1993).
- Tabak HH, Bunch RL. Steroid hormones as water pollutants. I: Metabolism of natural and synthetic ovulation-inhibiting hormones by microorganisms of activated sludge and primary settled sewage. *Dev Ind Microbiol* 11:367-376 (1970).
- Tabak HH, Bloomfield RN, Bunch RL. Steroid hormones as water pollutants. II: Studies on the persistence and stability of natural urinary and synthetic ovulation-inhibiting hormones and treated wastewaters. *Dev Ind Microbiol* 22:497-519 (1981).
- Ternes T, Wilken R-D, eds. *Drugs and Hormones as Pollutants of the Aquatic Environment: Determination and Ecotoxicological Impacts*. *Sci Total Environ* 225(1-2):176 pp (1999).
- Arcand-Hoy LD, Nimrod AC, Benson WH. Endocrine-modulating substances in the environment: estrogenic effects of pharmaceutical products. *Int J Toxicology* 17(2):139-158 (1998).
- Johnson AC, Williams RJ, Ulahannan T. Comment on identification of estrogenic chemicals in STW effluent. 1: Chemical fractionation and in vitro biological screening. *Environ Sci Technol* 33(2):369-370 (1999).
- Routledge EJ, Waldock M, Sumpter JP. Response to Comment on identification of estrogenic chemicals in STW effluent. 1. Chemical fractionation and in vitro biological screening [Letter response]. *Environ Sci Technol* 33(2):371 (1999).
- Colburn T, Dumanoski D, Myers JP. *Our Stolen Future: Are We Threatening Our Fertility, Intelligence, and Survival? A Scientific Detective Story*. New York:Plume/Penguin, 1997. Also see <http://www.osf-facts.org> [cited 4 September 1999].
- Chemical Manufacturers Association. *The Chemical Industry's Health and Environmental Effects Research Initiative*. Available: <http://www.cmahq.com/CMAWebsite3.nsf/pages/healthresearch> [cited 31 August 1999].
- Arnold SF, Klotz DM, Collins BM, Vonier PM, Guillet L J Jr, McLachlan JA. Synergistic activation of estrogen receptor with combinations of environmental chemicals. *Science* 272:1489-1492 (1996).
- Gaido KW, McDonnell DP, Korach KS, Safe SH. Estrogenic activity of chemical mixtures: Is there synergism? *Chit Activities* 17(2):1-7 (1997).
- McLachlan JA. Synergistic effect of environmental estrogens: report withdrawn. *Science* 277:459d-463d (1997).
- Arnold SF, Bergeron JM, Tran DQ, Collins BM, Vonier PM, Crews D, Toscano WA Jr, McLachlan JA. Synergistic responses of steroidal estrogens in vitro (yeast) and in vivo (turtles). *Biochem Biophys Res Commun* 235(2):336-342 (1997).
- Tian S-S, Lamb P, King AG, Miller SG, Kessler L, Luengo JI, Averill L, Johnson RK, Gleason JG, Pelus LM, Dillon SB, Rosen J. A small nonpeptidyl mimic of granulocyte-colony-stimulating factor. *Science* 281:257-259 (1998).
- Barinaga M. Small molecule fills hormone's shoes. *Science* 281:149-151 (1998).
- Hartmann A, Alder AC, Koller T, Widmer RM. Identification of



- fluoroquinolone antibiotics as the main source of *umuC* genotoxicity in native hospital wastewater. *Environ Toxicol Chem* 17(3):377-382 (1998).
63. Witte W. Medical consequences of antibiotic use in agriculture. *Science* 279:996-997 (1998).
64. Ash RJ, Mauch B, Moulder W, Morgan M. Antibiotic-resistant bacteria in U.S. rivers. Abstract no Q-383. In: Proceedings of the Conference of the American Society for Microbiology 99th Annual Meeting, June 1999, Chicago, Illinois. Herndon, VA:ASM Press, <http://www.asmsa.org/mgsrc/absprgbk.htm>
65. Eichorst S, Pfeifer A, Magill NG, Tischler ML. Antibiotic resistance among bacteria isolated from wild populations of resident Canada Geese in a suburban setting. Abstract no Q-402. In: Proceedings of the American Society for Microbiology 99th Annual Meeting, June 1999, Chicago, Illinois.
66. Heberer T, Stan H-J. Determination of clofibric acid and N-(phenylsulfonyl)-sarcosine in sewage river and drinking water. *Int J Environ Anal Chem* 67:113-124 (1997).
67. Buser H-R, Müller MD, Theobald N. Occurrence of the pharmaceutical drug clofibric acid and the herbicide mecoprop in various Swiss lakes and in the North Sea. *Environ Sci Technol* 32:188-192 (1998).
68. Jones KC, de Voigt P. Persistent organic pollutants (POPs): state of the science. *Environ Pollut* 100(1-3):209-221 (1999).
69. Stumpf M, Ternes TA, Wilken R-D, Rodrigues SV, Baumann W. Polar drug residues in sewage and natural waters in the state of Rio de Janeiro Brazil. *Sci Total Environ* 225(1):135-141 (1998).
70. Ternes TA, Stumpf M, Schuppert B, Haberer K. Simultaneous Determination of antiseptics and acidic drugs in sewage and river. *Vom Wasser* 90:295-309 (1998).
71. Buser H-R, Poiger T, Müller MD. Occurrence and fate of the pharmaceutical drug diclofenac in surface waters: rapid photodegradation in a lake. *Environ Sci Technol* 32:3449-3456 (1998).
72. Buser H-R, Poiger T, Müller MD. Occurrence and environmental behavior of the pharmaceutical drug ibuprofen in surface waters and in wastewater. *Environ Sci Technol* 33:2529-2535 (1999).
73. Hirsch R, Ternes TA, Haberer K, Kratz K-L. Determination of  $\beta$ -blockers and -sympathomimetics in the aquatic environment. *Vom Wasser* 87:263 (1996).
74. Kulkarni GK, Nagabhushanam R, Amaldoss G, Jaiswal RG, Fingerman M. In vivo stimulation of ovarian development in the red swamp crayfish *Procambarus clarkii* (Girard) by 5-hydroxytryptamine. *Invert Reprod Devel* 21(3):231-240 (1992).
75. Sarojini R, Nagabhushanam R, Fingerman M. In vivo evaluation of 5-hydroxytryptamine stimulation of the testes in the fiddler crab *Uca pugilator*: a presumed action on the neuroendocrine system. *Comparat Biochem Physiol* 106C(2):321-325 (1993).
76. Fong PP. Zebra mussel spawning is induced in low concentrations of putative serotonin reuptake inhibitors. *Biol Bull* 194:143-149 (1998).
77. Couper JM, Leise EM. Serotonin injections induce metamorphosis in larvae of the gastropod mollusc *Lymnaea obsoleta*. *Biol Bull* 191:178-86 (1996).
78. Sarojini R, Nagabhushanam R, Fingerman M. Mode of Action of the Neurotransmitter 5-Hydroxytryptamine in Stimulating Ovarian Maturation in the Red Swamp Crayfish *Procambarus clarkii*: an in vivo and in vitro study. *J Exp Zool* 271:395-400 (1995).
79. Fong PP, Huminski PT, d'Urso LM. Induction of potentiation of parturition in fingernail clams (*Sphaerium striatinum*) by selective serotonin re-uptake inhibitors (SSRIs). *J Exp Zool* 280(3):260-264 (1998).
80. Huber R, Smith K, Delago A, Isaksson K, Kravitz EA. Serotonin and aggressive motivation in crustaceans: altering the decision to retreat. *Proc Natl Acad Sci* 94:5939-5942 (1997).
81. Sacher F, Lochow E, Bethmann D, Brauch H-J. Occurrence of drugs in surface waters. *Vom Wasser* 90:233 (1998).
82. Steger-Hartmann T, Kümmerer K, Hartmann A. Biological degradation of cyclophosphamide and its occurrence in sewage water. *Ecotoxicol Environ Safety* 36:174-179 (1997).
83. Kümmerer K, Steger-Hartmann T, Meyer M. Biodegradability of the anti-tumour agent ifosfamide and its occurrence in hospital effluents and communal sewage. *Wat Res* 31(11):2705-2710 (1997).
84. Kümmerer K, Helmers E, Hubner P, Mascart G, Milandri M, Reinthaler F, Zwakenberg M. European hospital as a source for platinum in the environment in comparison with other sources. *Sci Total Environ* 225(1-2):155-165 (1999).
85. Falter R, Wilken R-D. Determination of carboplatinum and cisplatinum by interfacing HPLC with ICP-MS using ultrasonic nebulisation. *Sci Total Environ* 225(1-2):167-176 (1999).
86. White PA, Rasmussen JB. The genotoxic hazards of domestic wastes in surface waters. *Mutat Res* 410:223-236 (1998).
87. Wilson EK. Impotence drugs: more than Viagra. *Chem Eng News* 29:29-33 (1998).
88. Genicola FA. Personal communication. New Jersey Dept. Environmental Protection Office of Quality Assurance. Communication, 3 June 1999.
89. Maden M. Retinoic acid in development and regeneration. *J Biosciences* 21(3):299-312 (1996).
90. La Clair JJ, Bantle JA, Dumont J. Photoproducts and metabolites of a common insect growth regulator produce developmental deformities in *Xenopus*. *Environ Sci Technol* 32(10):1453-1461 (1998).
91. Kalsch W. Biodegradation of the iodinated X-ray contrast media diatrizoate and iopromide. *Sci Total Environ* 225(1-2):143-153 (1999).
92. Ternes TA, Hirsch R. Occurrence and behavior of iodinated contrast media in the aquatic environment. *Environ Sci Technol*, in press.
93. Steger-Hartmann T, Länge R, Schweinfurth H. Umweltverhalten und ökotoxikologische Bewertung von iodhaltigen Röntgenkontrastmitteln. [Environmental behavior and ecotoxicological assessment.] *Vom Wasser* 91:185-194 (1998).
94. Stevenson C, Davies R.J.H. Photosensitization of guanine-specific DNA damage by 2-phenylbenzimidazole and the sunscreen agent 2-phenylbenzimidazole-5-sulfonic acid. *Chem Res Toxicol* 12(1):38 (1999).
95. Statistisches Bundesamt 1993. Monatlicher Produktionseilbericht - Produktion ausgewählter Körperpflegemittel in Deutschland. [Monthly Production Report - Production of Selected Personal Care Products in Germany.] Wiesbaden, Germany: Statistische Bundesamt,
96. Gatermann R, Hühnerfuss H, Rimkus G, Attar A, Ketrup A. Occurrence of musk xylene and musk ketone metabolites in the aquatic environment. *Chemosphere* 36(11):2535-2547 (1998).
97. Bronaugh RL, Yourick JJ, Havery DC. Dermal exposure assessment for the fragrance musk xylol. Abstract No 274. In: Proceedings of the Society of Toxicology 1998 Annual Meeting.
98. Müller S, Schmid P, Schlatter C. Occurrence of nitro and non-nitro benzenoid musk compounds in human adipose tissue. *Chemosphere* 33(1):17-28 (1996).
99. Kafferlein HU, Göen T, Angerer J. Musk xylene: analysis, occurrence, kinetics, and toxicology. *Crit Rev Toxicol* 28(5):431-476 (1998).
100. Yamagishi T, Miyazaki T, Horii S, Kaneko S. Identification of musk xylene and musk ketone in freshwater fish collected from the Tama River, Tokyo. *Bull Environ Contam Toxicol* 26:656-662 (1981).
101. Yamagishi T, Miyazaki T, Horii S, Akiyama K. Synthetic musk residues in biota and water from Tama River and Tokyo Bay (Japan). *Arch Environ Contam Toxicol* 12:83-89 (1983).
102. Geyer HJ, Rimkus G, Wolf M, Attar A, Steinberg C, Ketrup A. Synthetische Nitromoschus-Duftstoffe und Bromocyclen - neue Umweltchemikalien in Fischen und Muscheln bzw. Muttermilch und Humanfett. [Synthetic nitro musk fragrances and bromocyclen—new environmental chemicals in fish and mussels as well as in breast milk and human lipids.] *Z Umweltchem Ökotox* 6(1):9-17 (1994).
103. Rimkus GG, Butte W, Geyer HJ. Critical considerations on the analysis and bioaccumulation of musk xylene and other synthetic nitro musks in fish. *Chemosphere* 35(7):1497-1507 (1997).
104. Winkler M, Kopf G, Hauptvogel C, Neu T. Fate of artificial musk fragrances associated with suspended particulate matter (SPM) from the River Elbe (Germany) in comparison to other organic contaminants. *Chemosphere* 37(6):1139-1156 (1998).
105. Kirschner EM. Boomers quest for agelessness. *Chem Eng News* 75(16):19-25 (1997).
106. Draisci R, Marchiafava C, Ferretti E, Palleschi L, Catellani G, Anastasio A. Evaluation of musk contamination of freshwater fish in Italy by accelerated solvent extraction and gas chromatography with mass spectrometric detection. *J Chromatogr A* 814:187-197 (1998).
107. Mersch-Sundermann V, Kevekedeso S, Jentero C. Lack of mutagenicity of polycyclic musk fragrances in *Salmonella typhimurium*. *Toxicol in Vitro* 12(4):389-393 (1998).
108. Heberer Th, Gramer S, Stan H-J. Occurrence and distribution of organic contaminants in the aquatic system in Berlin. Part iii: Determination of synthetic musks in Berlin surface water applying solid-phase microextraction (SPME) and gas chromatography-mass spectrometry (GC/MS). *Acta Hydrochim Hydrobiol* 27:150-156 (1999).
109. Kallenborn R, Gatermann R, Planting S, Rimkus GG, Lund M, Schlabach M, Burkow IC. Gas chromatographic determination of synthetic musk compounds in Norwegian air samples. *J Chromatogr A* 846(1-2):295-306 (1999).
110. Kevekedeso S, Mersch-Sundermann V, Diez M, Bolten C, Dunkelberg H. Genotoxicity of polycyclic musk fragrances in the sister-chromatid exchange test. *Anticancer Res* 18:449-452 (1998).
111. Behechti A, Schramm K-W, Attar A, Niederfellner J, Ketrup A. Acute aquatic toxicities of four musk xylene derivatives on *Daphnia magna*. *Water Res* 32(5):1704-1707 (1998).
112. Routledge EJ, Parker J, Odum J, Ashby J, Sümptner JP. Some alkyl hydroxy benzoate preservatives (parabens) are estrogenic. *Toxicol Appl Pharmacol* 153:12-19 (1998).
113. Okumura T, Nishikawa Y. Gas chromatography-mass spectrometry determination of triclosans in water sediment and fish samples via methylation with diazomethane. *Analytica Chimica Acta* 325(3):175-184 (1996).
114. McMurry LM, Oethinger M, Levy SB. Triclosan targets lipid synthesis. *Nature* 394(6693):531-532 (1998).
115. Nagtegaal M, Ternes TA, Baumann W, Nagel R. Nachweis von UV-Filtersubstanzen in Wasser und Fischen aus dem Meerfelder Maar in der Eifel. Detection of UV-sunscreen agents in water and fish of the Meerfelder Maar the Eifel Germany. *UWSF-Z für Umweltchem Ökotox* 9(2):79-86 (1997).
116. Hany J, Nagel R. Nachweis von UV-Filtersubstanzen in Muttermilch. [Detection of UV-sunscreen agents in breast milk.] *Deutsche Lebensmittell* 91:341-245 (1995).
117. Borman S. End run around FDA? *Chem Eng News* 1:45-47 (1998). Also: Zeisel SH. Regulation of "nutraceuticals." *Science* 285:1853-1855 (1999).
118. Pharmanex, Inc. v. Donna Shalala. 2:97 CV 0262 K, 1999. Available <http://www.fjx.com/LJXfiles/fda/cholestin.html> [cited 10 November 1999].
119. Dietary Supplement Health and Education Act of 1994 (DSHEA). Pub L no 103-417, 108 Stat. 4325 (1994). Available: <http://odp.od.nih.gov/ods/about/law.html> [cited 10 November 1999].
120. NIH Office of Dietary Supplements. The International Bibliographic Information on Dietary Supplements (IBIDS) Database. Available: <http://odp.od.nih.gov/ods/databases/ibids.html> [cited 31 August 1999].
121. National Center for Complementary and Alternative Medicine. Home Page. Available: <http://altmed.od.nih.gov/nccam> [cited 31 August 1999].
122. PharmaPrint. Home Page. Available: <http://www.pharmaprint.com> [cited 31 August 1999].
123. Shao Y, He K, Zheng B, Zheng Q. Reversed-phase high-performance liquid chromatographic method for quantitative analysis of the six major kavalactones in *Piper methysticum*. *J Chromatogr A* 825:1-8 (1998).
124. Cytotoxicology Laboratory, Uppsala. Home Page. Available: [http://www.ctlu.se/CTLU\\_HOME.html](http://www.ctlu.se/CTLU_HOME.html). Available: [http://www.ctlu.se/CTLU\\_MEIC.html](http://www.ctlu.se/CTLU_MEIC.html) [cited 31 August 1999].
125. Lilius H, Hästbacka T, Isomaa B. A comparison of the toxicity of 30 reference chemicals to *Daphnia magna* and *Daphnia pulex*. *Environ Toxicol* 14(12):2085-2088 (1995).
126. Calleja MC, Perronne G, Geladi P. The predictive potential of a battery of ecotoxicological tests for human acute toxicity. *Altern Lab Animals* 21:330-349 (1993).
127. Kalbfus W, Kopf W. Erste Ansätze zur ökotoxikologischen Bewertung von Pharmaka in Oberflächengewässern (First Attempts for the Ecotoxicological Assessment of Drugs in Surface Waters). 51. Fachtagung Bayerisches Landesamt für Wasserwirtschaft, 1996 [51st Meeting of Bavarian Ministry of Water Economy, 1996]. Munich, Germany, 1997.
128. Kurelec B. The genotoxic disease syndrome. *Mar Environ Res* 35:341-348 (1993).
129. Weiss B. Environmental health: nickel-and-diming it [Letter]. *Science* 282:1644 (1998).
130. Weiss B. A risk assessment perspective on the neurobehavioral toxicity of endocrine disruptors. *Toxicol Ind Health* 14(1-2):341-359 (1998).
131. Grimes DJ, Singleton FL, Colwell RR. Allogenic succession of marine bacterial communities in response to pharmaceutical waste. *J Appl Bacteriol* 57:247-261 (1984).
132. Azur Environmental. Microtox® Rapid Toxicity Testing System. Available: <http://www.azurenv.com/mtox.htm> [cited 31 August 1999].
133. Tahedl H, Häder D-P. Fast examination of water quality using the automatic biotest ECOTOX based on the movement behavior of a freshwater flagellate. *Wat Res* 33(2):426-432 (1999).
134. OECD (Organisation for Economic Co-operation and Development). OECD's Guidelines for the Testing of Chemicals Currently Available Test Guidelines, Draft Test Guidelines, and Guidance and Review Documents. List Revised, October 1999. Available: <http://www.oecd.org/ehs/test/testlist.htm> [cited 10 November 1999].

135. OECD. (Organisation for Economic Co-operation and Development). Detailed Review Paper on Aquatic Testing Methods for Pesticides and Industrial Chemicals. Part 1: Report, 1998, OECD Series on Testing and Assessment No. 11, ENV/MC/CHEM(98)19/PART1. Available: <http://www.oecd.org/ehs/ehsmono/index.htm#TESTING> [cited 1 September 1999]. Also: Detailed Review Paper on Aquatic Testing Methods for Pesticides and Industrial Chemicals - Part 2: Annexes, 1998, OECD Series on Testing and Assessment No. 11, ENV/MC/CHEM(98)19/PART2. Available: <http://www.oecd.org/ehs/ehsmono/index.htm#TESTING> [cited 1 September 1999].
136. U.S. EPA. Final Guidelines for Ecological Risk Assessment. Risk Assessment Forum. EPA/630/R-95/002F. Washington, DC:U.S. Environmental Protection Agency, April 1998. Available: <http://www.epa.gov/ncea/ecorsk.htm> [cited 1 September 1999].
137. PhRMA (Pharmaceutical Research & Manufacturers Association). Industry Profile 1998. Available: <http://www.phrma.org/publications/industry/profile98/exec.html> [cited 31 August 1999].
138. PhRMA (Pharmaceutical Research & Manufacturers Association). New Drug Approvals in 1998. Available: [http://www.phrma.org/charts/nda\\_c98.html](http://www.phrma.org/charts/nda_c98.html) [cited 31 August 1999].
139. FDA (Food and Drug Administration) Modernization Act of 1997. Available: <http://www.fda.gov/opacom/backgrounders/modact.htm> and <http://www.fda.gov/cdrh/modact/modern.html> [cited 31 November 1999].
140. EudraLex (Rules Governing Medicinal Products in the European Union). Vol 5: Pharmaceutical legislation: Veterinary Medicinal Products. Available: <http://dg3.eudra.org/eudralex/vol-5/home.htm> [cited 31 August 1999].
141. EMEA (The European Agency for the Evaluation of Medicinal Products: Veterinary Medicines Evaluation Unit). Note for Guidance: Environmental Risk Assessment for Veterinary Medical Products other than GMO-Containing and Immunological Products. The European Agency for the Evaluation of Medicinal Products Veterinary Medicines Evaluation Unit. EMEA/CVMP/055/96-Final, 1997. Available: <http://www.eudra.org/vetdocs/PDFs/GUIDE/005596en.pdf> [cited 11 November 1999].
142. OECD (Organisation for Economic Co-operation and Development). Environmental Exposure Assessment Strategies for Existing Industrial Chemicals in OECD Member Countries. 1999, Series on Testing and Assessment No. 17, ENV/JM/MONO(99)10. Available: <http://www.oecd.org/ehs/ehsmono/index.htm#TESTING> [cited 1 September 1999].
143. The European Agency for the Evaluation of Medicinal Products: Veterinary Medicines Evaluation Unit. Guideline on Environmental Impact Assessment (EIAS) for Veterinary Medicinal Products - Phase I. The European Agency for the Evaluation of Medicinal Products Veterinary Medicines Evaluation Unit. Canary Wharf, England. EMEA/CVMP/592/98-consultation, 10 December 1998. Available: <http://www.eudra.org/vetdocs/PDFs/VICH/059298en.pdf> [cited 11 November 1999].
144. Directive of the European Parliament and of the Council No. 98/8/EC (16 February 1998): On the placing of biocidal products on the market. Available: [http://www.retroscreen.com/BiocidalDirective/biocidal\\_directive.htm](http://www.retroscreen.com/BiocidalDirective/biocidal_directive.htm) [cited 10 November 1999].
145. FDA (Food and Drug Administration). Guidance for Industry: Environmental Assessment of Human Drug and Biologics Application. CDER/CBER CMC 6, rev 1, 39 pp, July 1998. Available: <http://www.fda.gov/cber/guidelines.htm> [cited: 1 September 1999].
146. National Research Council, Science and Judgment in Risk Assessment. Washington, DC:National Academy Press, 1994, 672 pp. Available: <http://books.nap.edu/catalog/2125.html> [cited 31 August 1999].
147. Jenkins AL, Uy OM, Murray GM. Polymer-based lanthanide luminescent sensor for detection of the hydrolysis product of the nerve agent soman in water. *Anal Chem* 71(2):373-378 (1999).
148. National Institute of Standards and Technology. NIST/EPA/NIH Mass Spectral Library - NIST 98. Available: <http://www.nist.gov/srd/nist1a.htm> [cited 1 September 1999].
149. Steger-Hartmann T, Kümmerer K, Schecker J. Trace analysis of the antineoplastics ifosfamide and cyclophosphamide in sewage water by two-step solid-phase extraction and gas chromatography-mass spectrometry. *J Chromatogr A* 726:179-184 (1996).
150. Belfroid AC, Van der Horst A, Vethaak AD, Schäfer AJ, Rijs GBJ, Wegener J, Cofino WP. Analysis and occurrence of estrogenic hormones and their glucuronides in surface water and waste water in the Netherlands. *Sci Total Environ* 225(1-2):101-108 (1999).
151. Burhenne J, Ludwig M, Nikoloudis P, Spittler M. Photolytic degradation of fluoroquinolone carboxylic acids in aqueous solution. Part I: Primary photoproducts and half-lives. *Environ Sci Pollut Res* 4(1):10-15 (1997).
152. Burhenne J, Ludwig M, Spittler M. Photolytic degradation of fluoroquinolone carboxylic acids in aqueous solution. Part II: Isolation and structural elucidation of polar photometabolites. *Environ Sci Pollut Res* 4(2):61-67 (1997).
153. Hartmann A, Golet EM, Gattiser S, Alder AC, Koller T, Widmer RM. Primary DNA damage but not mutagenicity correlates with Ciprofloxacin concentrations in German hospital wastewaters. *Arch Environ Contam Toxicol* 36:115-119 (1999).
154. Gatermann R, Hellou J, Hühnerfuss H, Rimkus G, Zitko V. Polycyclic and nitro musks in the environment: a comparison between Canadian and European aquatic biota. *Chemosphere* 38(14):3431-3441 (1999).

資 料 - IV 目 次

(資料 7) 医薬品の環境影響評価のための指針に関する報告(案)  
未定稿

## 医薬品の環境影響評価のための指針に関する報告（案）

1. 医薬品の環境影響評価の必要性
  - 1-1. 化学物質の生態影響評価の背景
    - 1-1-1. 国際的動向
    - 1-1-2. 国内の動向
  - 1-2. 医薬品生態影響評価の背景
    - 1-2-1. 国際的動向
    - 1-2-2. 国内の動向
  - 1-3. 我が国において医薬品の環境影響評価を導入する必要性
    - 1-3-1. 我が国の化学物質の審査制度の概要
    - 1-3-2. 我が国の医薬品審査制度の概要
    - 1-3-3. 医薬品の環境影響評価の導入の必要性
  - 1-4. まとめ
2. 医薬品のリスク管理の枠組み（案）
  - 2-1. 対象物質
    - 2-1-1. 対象物質の範囲
    - 2-1-2. 代謝産物に対する考え方
    - 2-1-3. 易分解性物質に対する考え方
    - 2-1-4. 高濃縮性物質に対する考え方
    - 2-1-5. 排水規制との棲み分けの概念
    - 2-1-6. 新規医薬品と既存医薬品の区分
  - 2-2. 評価の個別要素
    - 2-2-1. PEC/PNEC による評価
    - 2-2-2. 確率的な扱いの PEC/PNEC による評価
    - 2-2-3. 段階的な評価
    - 2-2-4. カテゴリー（グループ）評価の可能性
    - 2-2-5. 例外物質の取り扱い
    - 2-2-6. 総量規制
  - 2-3. 提案する評価法
    - 2-3-1. 提案する評価法の概要
    - 2-3-2. 提案する評価法における課題
  - 2-4. まとめ
3. 環境濃度の推定
  - 3-1. 排出シナリオ
  - 3-2. 排出モデル
  - 3-3. 環境濃度の推定法
  - 3-4. 推定法の妥当性