

Human intake of benzophenones occurs from food and medicine, and these chemicals may accumulate in the body to some extent. Indeed, Hany and Nagel (1995) detected 2-OH-4-MeO-BP and octyl *p*-methoxycinnamate used in as sunscreen in human milk (total 16–417 ng/g fat) from Germans. This level corresponds to  $10^{-7}$  to  $10^{-6}$  M. The contents of 4-hydroxybenzophenones have not been reported. However, when 4-hydroxybenzophenones are applied to human skin as sunscreen (usually 1–2% content), the level may temporarily reach the level used in this study immediately after the dose. Subsequently, the level of 4-hydroxybenzophenones would decrease rapidly owing to metabolism and excretion. These considerations and the *in vivo* results obtained in this study suggest that benzophenones pose a relatively low risk to humans in terms of endocrine disrupting activity, in spite of the *in vitro* activity shown in this study. In contrast, some hydroxylated benzophenones, such as those hydroxylated at the 2- and 4-positions, showed both estrogenic and antiandrogenic activities. Thus, these compounds might exhibit endocrine-disrupting action via both mechanisms in animals. Their *in vivo* effects due to these combined activities should be further investigated.

### Acknowledgments

This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas (13027256) from the Japanese Ministry of Education, Science, Sports and Culture, and a Grant-in-Aid for Scientific Research (C16590092) from the Japan Society for the Promotion of Science.

### References

- Andersen, H.R., Andersson, A.-M., Arnold, S.F., Autrup, H., Barfoed, M., Beresford, N.A., et al., 1999. Comparison of short-term estrogenicity tests for identification of hormone-disrupting chemicals. *Environ. Health Perspect.* 107 (1), 89–108.
- Blair, R.M., Fang, H., Branham, W.S., Hass, B.S., Dial, S.L., Moland, C.L., Tong, W., Shi, L., Perkins, R., Sheehan, D.M., 2002. The estrogen receptor relative binding affinities of 188 natural and xenochemicals: structural diversity of ligands. *Toxicol. Sci.* 54, 138–153.
- Burdock, G.A., Pence, D.H., Ford, R.A., 1991. Safety evaluation of benzophenone. *Food Chem. Toxicol.* 11, 741–750.
- Chen, C.W., Hurd, C., Vorobjekina, D.P., Arnold, S.F., Notides, A.C., 1997. Transcriptional activation of the human estrogen receptor by DDT isomers and metabolites in yeast and MCF-7 cells. *Biochem. Pharmacol.* 53, 1161–1172.
- Dutta, K., Das, M., Rahman, T., 1993. Toxicological impacts of benzophenone on the liver of guinea pig. *Bull. Environ. Contam. Toxicol.* 50, 282–285.
- Fang, H., Tong, W., Perkins, R., Soto, A.M., Prechtel, N.V., Sheehan, D.M., 2000. Quantitative comparisons of *in vitro* assay for estrogenic activities. *Environ. Health Perspect.* 108, 723–729.
- Felix, T., Hall, B.J., Brodbelt, J.S., 1998. Determination of benzophenone-3 and metabolites in water and human urine by solid-phase micro-extraction and quadrupole ion trap GC-MS. *Anal. Chim. Acta* 371, 195–203.
- Gaido, K.W., Leonard, L.S., Lovell, S., Gould, J.C., Babai, D., Portier, C.J., Mcdonell, D.P., 1997. Evaluation of chemicals with endocrine modulating activity in yeast-based steroid hormone receptor gene transcription assay. *Toxicol. Appl. Pharmacol.* 143, 205–212.
- Gray Jr., L.E., Ostby, J., Manosson, E., Kelce, W.R., 1999. Environmental antiandrogens: low dose of the fungicide vinclozolin alter sexual differentiation of the male rat. *Toxicol. Ind. Health* 15, 48–64.
- Hagedorn-Leweke, U., Lippold, B.C., 1995. Absorption of sunscreens and other compounds through human skin *in vivo*: derivation of a method to predict maximum fluxes. *Pharmacol. Res.* 12, 1354–1360.
- Hany, J., Nagel, R., 1995. Detection of sunscreen agents in human breast milk. *Deutsche Lebens.-Rund.* 91, 341–345.
- Hayden, C.G.J., Roberts, M.S., Benson, H.A.E., 1997. Systemic absorption of sunscreen after topical application. *Lancet* 350, 863–864.
- Hershberger, L., Shipley, E., Meyer, R., 1953. Myotrophic activity of 19-nortestosterone and other steroids determined by modified levator ani muscle method. *Proc. Soc. Exp. Biol. Med.* 83, 175–180.
- Hong, H., Tong, W., Fang, H., Shi, L., Xie, Q., Wu, J., Perkins, R., Walker, J.D., Branham, W., Sheehan, D.M., 2002. Prediction of estrogen receptor binding for 58,000 chemicals using an integrated system of a tree-based model with structural alerts. *Environ. Health Perspect.* 110, 29–36.
- Jiang, R., Roberts, M.S., Collins, D.M., Benson, H.A.E., 1999. Absorption of sunscreens across human skin: an evaluation of commercial products for children and adults. *Br. J. Clin. Pharmacol.* 48, 635–637.
- Kawamura, Y., Ogawa, Y., Nishimura, T., Kikuchi, Y., Nishikawa, J., Nishihara, T., et al., 2003. Estrogenic activities of UV stabilizers used in food contact plastics and benzophenone derivatives tested by the yeast two-hybrid assay. *J. Health Sci.* 49, 205–212.
- Kelce, W.R., Stone, C.R., Laws, S.C., Gray, L.E., Kemppainen, J.A., Wilson, E.M., 1995. Persistent DDT metabolite p,p'-DDE is a potent androgen receptor antagonist. *Nature* 375, 581–585.
- Kitamura, S., Suzuki, T., Ohta, S., Fujimoto, N., 2003. Antiandrogenic activity and metabolism of the organophosphorus pesticide fenitrothion and related compounds. *Environ. Health Perspect.* 111, 503–508.
- Klein, K., 1992. Encyclopedia of UV absorbers for sunscreen products. *Cosmetics. Toiletries* 107, 45–64.
- Kupfer, D., Bulger, W.H., 1987. Metabolic activation of pesticides with proestrogenic activity. *Fed. Proc.* 46, 1864–1869.
- Ma, R., Cotton, B., Lichtensteiger, W., Schlumpf, M., 2003. UV filters with antagonistic action at androgen receptors in the MDA-kb2 cell transcriptional-activation assay. *Toxicol. Sci.* 74, 43–50.
- Matsumoto, K., Akao, Y., Kobayashi, E., Ito, T., Ohguchi, K., Tanaka, T., Iinuma, M., Nozawa, Y., 2003. Cytotoxic benzophenone derivatives from *Garcinia* species display a strong apoptosis-inducing effect against human leukemia cell lines. *Biol. Pharm. Bull.* 26, 569–571.
- Nakagawa, Y., Suzuki, T., 2002. Metabolism of 2-hydroxy-4-methoxybenzophenone in isolated rat hepatocytes and xenoestrogenic effects of its metabolites on MCF-7 human breast cancer cells. *Chem. Biol. Inter.* 139, 115–128.
- Nakagawa, Y., Tayama, K., 2001. Estrogenic potency of benzophenone and its metabolites in juvenile female rats. *Mol. Toxicol.* 75, 74–79.
- Okereke, C.S., Kadry, A.M., Abdel-Rahman, M.S., Davis, R.A., Friedman, M.A., 1993. Metabolism of benzophenone-3 in rats. *Drug Metab. Dispos.* 21, 788–791.
- Okereke, C.S., Barat, S.A., Abdel-Rahman, M.S., 1995. Safety evaluation of benzophenone-3 after dermal administration in rats. *Toxicol. Lett.* 80, 61–67.
- Satoh, K., Nagai, F., Aoki, N., 2001. Several environmental pollutants have binding affinities for both androgen receptor and estrogen receptor  $\alpha$ . *J. Health Sci.* 47, 495–501.
- Schauder, S., Ippen, H., 1997. Contact and photocontact sensitivity to sunscreens. Review of a 15 year experience and of the literature. *Contact Dermatitis* 37, 221–232.
- Schlumpf, M., Cotton, B., Conscience, M., Haller, V., Steinmann, B., Lichtensteiger, W., 2001. *In vitro* and *in vivo* estrogenicity of UV screens. *Environ. Health Perspect.* 109, 239–244.

- Schultz, W.T., Seward, J.R., Sinks, G.D., 2000. Estrogenicity of benzophenones evaluated with a recombinant yeast assay: comparison of experimental and rules-based predicted activity. *Environ. Toxicol. Chem.* 19, 301–304.
- Soto, A.M., Chung, K.L., Sonnenschein, C., 1994. The pesticides endosulfan, toxaphene and dieldrin have estrogenic effects on human estrogen-sensitive cells. *Environ. Health Perspect.* 102, 380–383.
- Stresser, D.M., Kupfer, D., 1998. Human cytochrome P450-catalyzed conversion of the proestrogenic pesticide methoxychlor into an estrogen. Role of CYP2C19 and CYP1A2 in *O*-demethylation. *Drug Metab. Dispos.* 26, 868–874.
- Sugihara, K., Kitamura, S., Sanoh, S., Ohta, S., Fujimoto, N., Maruyama, S., Ito, A., 2000. Metabolic activation of the proestrogens trans-stilbene and trans-stilbene oxide by rat liver microsomes. *Toxicol. Appl. Pharmacol.* 167, 46–54.
- Tamura, H., Maness, S.C., Reischmann, K., Dorman, D.C., Gray, L.E., Gaido, K.W., 2001. Androgen receptor antagonism by the organophosphate insecticide fenitrothion. *Toxicol. Sci.* 60, 56–62.
- Yamasaki, K., Takeyoshim, M., Sawaki, M., Imatanaka, N., Shinoda, K., Takatsuki, M., 2003. Immature rat uterotrophic assay of 18 chemicals and Hershberger assay of 30 chemicals. *Toxicology* 183, 93–115.

## COMPARATIVE METABOLISM OF POLYCHLORINATED BIPHENYLS AND TISSUE DISTRIBUTION OF PERSISTENT METABOLITES IN RATS, HAMSTERS, AND GUINEA PIGS

Koichi Haraguchi, Nobuyuki Koga, and Yoshihisa Kato

*Daiichi College of Pharmaceutical Sciences, Fukuoka, Japan (K.H.); Department of Food and Nutrition, Nakamura Gakuen University, Fukuoka, Japan (N.K.); and School of Pharmaceutical Sciences, University of Shizuoka, Shizuoka, Japan (Y.K.)*

Received September 21, 2004; accepted December 16, 2004

### ABSTRACT:

The present study was performed to compare the metabolite profiles of polychlorinated biphenyls (PCBs) in the liver and serum of rats, hamsters, and guinea pigs after exposure to a PCB mixture, Kanechlor 500 (100 mg/kg, i.p.). The percentage of contribution of major PCB residues in the liver 5 days after exposure indicated that nonplanar PCBs with 2,4- or 2,3,4-chlorine substitution were more abundant in the liver in the order rats (43% of total PCBs) > hamsters (20%) > guinea pigs (11%), whereas coplanar PCBs with 4-, 3,4-, or 3,4,5-chlorine substitution were predominant in guinea pigs (61%), followed by hamsters and rats (both 26%). The hepatic concentrations of methylsulfonyl metabolites (MeSO<sub>2</sub>-CBs) were higher in the order guinea pigs > rats > hamsters. Whereas ham-

sters formed minute amounts of MeSO<sub>2</sub>-CBs from 2,5-dichloro-substituted PCBs, guinea pigs formed higher levels of meta-MeSO<sub>2</sub>-CBs derived from 2,3,6-trichloro-substituted PCBs. In contrast, the serum concentrations of phenolic PCBs were higher in the order hamsters > rats > guinea pigs. Metabolites were predominated by 4-OH-2,3,5,3',4'-pentaCB (89% contribution) for rats, 3-OH-2,4,5,2',4'-pentaCB (56%) for guinea pigs, and dihydroxylated metabolites (39%) for hamsters. The reduced elimination of coplanar PCBs and the specific distribution of MeSO<sub>2</sub>- and phenolic PCBs may have implications for the differences in sensitivity to PCB toxicity among rats, guinea pigs, and hamsters.

Metabolism of PCBs proceeds via P450-mediated formation of arene oxide intermediates, which results in the distribution of both OH-PCBs and MeSO<sub>2</sub>-CBs (Letcher et al., 2000). The ratios of formation of these metabolites are dependent on the degree of substrate chlorination (Kato et al., 1980; Haraguchi et al., 1997) as well as the metabolic capacity of the species (Safe, 1994; Koga and Yoshimura, 1996). Some OH-PCBs and MeSO<sub>2</sub>-CBs are persistent and show specific retention in the blood or tissues of laboratory and wild animals (Bergman et al., 1994a,b; Oberg et al., 2002) as well as in humans (Sandau et al., 2000; Chu et al., 2003). Phenolic PCB metabolites cause alterations in thyroid hormone metabolism (Morse et al., 1996; Brouwer et al., 1998), inhibit estrogen sulfation and binding to estrogen receptors for estrogenic or antiestrogenic effects (Connor et al., 1997; Kester et al., 2002), and give rise to potentially cytotoxic dihydroxylated or quinoid PCB metabolites (Amaro et al., 1996). On the other hand, some MeSO<sub>2</sub>-CBs induce expression of several P450 isozymes (Kato et al., 1997), reduce thyroid hormone levels in rats (Kato et al., 1999), and also show antiestrogenic effects

(Letcher et al., 2002). These persistent metabolites may be responsible for the toxic effects of PCBs.

Guinea pigs are the most sensitive experimental animals to the toxicity of coplanar PCBs and polychlorinated dibenzo-*p*-dioxins (Kociba and Cabey, 1985), whereas hamsters are less sensitive (Olson et al., 1980; Wroblewski and Olson, 1988). These differences in sensitivity may be due to metabolic activation by the unique P450-dependent monooxygenase system and substrate specificity in these species (Koga et al., 1998). For example, *in vitro* studies have shown that 3,4,3',4'-tetraCB (CYP1A1) is hydroxylated by rats or hamsters but not by guinea pigs (Koga et al., 1995), whereas 2,4,5,2',4',5'-hexaCB (CYP2B18) is hydroxylated by guinea pigs but hardly by rats or hamsters (Ariyoshi et al., 1997). Therefore, the different metabolic capacities of PCBs may result in the differences in distribution profiles of metabolites in animals exposed to PCBs.

In the present study, we examined the species differences in tissue distribution of PCB residues and persistent PCB metabolites among rats, hamsters, and guinea pigs 5 days after exposure to a technical PCB mixture, Kanechlor 500. This article describes the 1) comparison of the residual PCB profiles (percentage of contribution) in the liver of these three species, 2) species differences in metabolite profiles of MeSO<sub>2</sub>-CBs in the liver and of OH-PCBs in serum, and 3) GC/MS characterization of unknown metabolites retained in the serum of guinea pigs.

The research was partially funded by a grant-in-aid for Scientific Research (C) (no. 16590101, K.H.; 15510058, Y.K.; and 14572119, N.K.) from the Japan Society for the Promotion of Science and a Health and Labor Sciences Research Grant (Y.K.) from the Ministry of Health, Labor, and Welfare of Japan.

Article, publication date, and citation information can be found at <http://dmd.aspetjournals.org>.

doi:10.1124/dmd.104.002444.

**ABBREVIATIONS:** PCB, polychlorinated biphenyl; P450, cytochrome P450; OH, hydroxyl; CB, chlorobiphenyl; OH-PCB, hydroxylated PCB; MeSO<sub>2</sub>-CB, methylsulfonyl PCB; GC/MS, gas chromatography-mass spectrometry; GC/ECD, gas chromatography-electron capture detection; MC, 3-methylcholanthrene.

TABLE 1

Residual patterns of PCBs in the livers of rats, hamsters, and guinea pigs dosed with Kanechlor 500 (100 mg/kg i.p.)

The 41 major isomers of a total of 61 were quantified and values represent the means for two animals.

Type	IUPAC	Structure	Relative Composition (%) of PCBs						
			Rat	Guinea Pig	Hamster	Kanechlor 500			
Group A	4-	28	2,4,4'-	0.08	1.03	0.15	0.86		
		37	3,4,4'-	N.D.	6.01	0.04	0.15		
		60	2,3,4,4'-	0.09	0.57	0.10	0.10		
		74	2,4,5,4'-	0.72	4.33	0.74	0.54		
		81	3,4,5,4'-	N.D.	0.19	0.02	0.007		
	3,4-	114	2,3,4,5,4'-	0.47	2.24	0.33	0.25		
		77	3,4,3',4'-	0.02	1.82	0.06	0.03		
		107	2,3,5,3',4'-	N.D.	2.46	N.D.	0.62		
		118	2,4,5,3',4'-	17.6	32.2	18.6	7.68		
		126	3,4,5,3',4'-	0.20	0.41	0.23	0.02		
	3,4,5-	156	2,3,4,5,3',4'-	4.15	5.87	3.33	1.25		
		157	2,3,4,3',4',5'-	0.91	1.10	0.85	0.25		
		167	2,4,5,3',4',5'-	1.26	2.22	1.74	0.45		
		Sum of group A	25.5	60.5	26.2	12.2			
Group B	2,4-	47	2,4,2',4'-	0.18	0.003	N.D.	0.20		
		49	2,4,2',5'-	0.11	0.04	0.05	0.83		
		85	2,3,4,2',4'-	1.75	0.04	0.05	0.83		
		99	2,4,5,2',4'-	12.6	2.44	3.45	3.22		
		137	2,3,4,5,2',4'-	1.28	0.30	0.49	0.42		
	2,3,4-	147	2,3,5,6,2',4'-	0.27	0.02	0.02	0.11		
		87	2,3,4,2',5'-	0.73	0.34	0.62	3.92		
		105	2,3,4,3',4'-	5.16	3.92	4.44	2.99		
		128	2,3,4,2',3',4'-	3.05	N.D.	N.D.	2.11		
		132	2,3,4,2',3',6'-	0.17	N.D.	0.08	2.34		
		138	2,3,4,2',4',5'-	13.7	2.22	8.81	5.56		
		170	2,3,4,5,2',3',4'-	3.01	1.72	2.13	0.96		
		177	2,3,4,2',3',5',6'-	0.71	0.04	0.08	0.23		
		195	2,3,4,5,6,2',3',4'-	0.21	0.02	0.04	0.05		
				Sum of group B	42.9	11.1	20.3	23.8	
		Group C	2,4,5-	146	2,3,5,2',4',5'-	1.32	1.18	2.41	0.62
				153	2,4,5,2',4',5'-	12.1	10.3	17.0	5.36
				180	2,3,4,5,2',4',5'-	3.79	2.11	8.68	1.34
				183	2,3,4,6,2',4',5'-	1.30	0.35	1.33	0.41
187	2,3,5,6,2',4',5'-			0.92	0.28	1.63	0.36		
2,3,4,5-	203		2,3,4,5,6,2',4',5'-	0.51	0.05	1.21	0.15		
	194		2,3,4,5,2',3',4',5'-	0.57	0.4	0.57	0.19		
	201		2,3,4,5,2',3',5',6'-	0.32	0.27	0.94	0.10		
				Sum of group C	20.8	15	33.8	8.53	
	Group D		2,5-	52	2,5,2',5'-	0.08	0.36	1.40	5.56
95		2,3,6,2',5'-		0.13	0.18	0.73	6.52		
101		2,4,5,2',5'-		1.35	2.15	4.68	10.0		
2,3,6-		141	2,3,4,5,2',5'-	0.20	0.08	0.59	1.13		
		110	2,3,6,3',4'-	0.35	0.16	0.41	7.44		
		149	2,3,6,2',4',5'-	0.58	0.6	0.82	2.76		
				Sum of group D	2.69	3.53	8.63	33.4	
				Total %	91.9	90.0	88.9	77.9	

N.D., not detected (&lt; 0.002%).

### Materials and Methods

**Chemicals.** The standard reference compounds, methoxy-PCBs and MeSO<sub>2</sub>-CBs, were synthesized as described previously (Haraguchi et al., 1987; Bergman et al., 1995). Syntheses of veratrol PCBs will be described elsewhere. The chemical names of MeSO<sub>2</sub>-CBs and OH-PCBs were simplified on the basis of the IUPAC-derived numbering system of the parent PCBs. All solvents (*n*-hexane, acetone, ethanol, and dichloromethane) were of pesticide grade (Kanto Chemical Co., Tokyo, Japan).

**Animals.** Male Wistar rats (b.wt. 190–200 g), male Hartley guinea pigs (280–290 g), and male Golden Syrian hamsters (80–90 g) were housed three or four per cage in the laboratory with free access to commercial chow and tap water. Animals received a single i.p. injection of 100 mg/kg Kanechlor 500 dissolved in corn oil. Control animals received an equivalent volume of vehicle alone. The animals were sacrificed by decapitation 5 days after injection, and the liver, lungs, and serum were removed and stored at –20°C until analysis.

**Isolation of Metabolites.** Tissue samples were treated as described previously (Haraguchi et al., 1998). Briefly, tissues were homogenized and extracted with acetone/*n*-hexane (2:1, v/v). Two internal standards, 2,3,4,5,3',4',5'-heptaCB (70 ng) and 4-methyl-3-MeSO<sub>2</sub>-5,2',3',4',5'-pentaCB (24 ng), were added to each extract, and the mixtures were applied to a gel-permeation column packed with Bio-Beads S-X3 (50 g; Bio-Rad, Hercules, CA). Dichloromethane/*n*-hexane (1:1) was used as a mobile phase at a flow rate of 4 ml/min. The metabolite fraction (120–200 ml) was divided into phenolic and neutral fractions by partitioning with potassium hydroxide (0.5 M in 30% ethanol) and *n*-hexane. The potassium hydroxide phase was acidified with hydrochloric acid (2 M, 10 ml) and extracted with *n*-hexane/diethyl ether (9:1), and then methylated with diazomethane for 30 min at 4°C. The neutral fraction (*n*-hexane layer) was further subjected to silica gel column chromatography (1 g; Wakogel S-1; Wako Pure Chemicals, Osaka, Japan) by elution with 10 ml of *n*-hexane for PCBs and successively with 10 ml of dichloromethane.




Fig. 1. GC/ECD chromatograms of methylsulfonyl PCB metabolites in the livers of rats (A), guinea pigs (B), and hamsters (C) after exposure to Kanechlor 500 (100 mg/kg i.p.).

ane for MeSO<sub>2</sub>-CBs. Serum (0.1–1 ml) was spiked with an internal standard (2,3,4,5,3',4',5'-heptaCB), acidified with 4 volumes of 0.5 M HCl and extracted with *n*-hexane/diethyl ether (1:1). The phenolic PCB fraction was spiked with another internal standard (2,3,4,5,6,3',4',5'-octaCB) and methylated with diazomethane. Recoveries of 4'-hydroxy-2,3,4,5,3',5'-hexaCB added to control serum and liver samples prior to extraction were 89 to 97% (*n* = 4) in the phenolic fraction.

**Identification and Quantification.** Identification of individual PCBs and phenolic PCB metabolites was performed on a GC/MS system (electron ionization-selected ion monitoring mode; QP-5000; Shimadzu, Kyoto, Japan) with a DB-5MS capillary column (60 m × 0.25 mm i.d. and 0.25- $\mu$ m film thickness; J&W Scientific, Folsom, CA). The injection temperature was 250°C. The temperature program was initiated at 100°C (2 min) by 20°C/min to 250°C, and then by 2°C/min to 290°C (20 min). Helium was used as a carrier gas at a constant flow rate of 1.0 ml/min. The transfer line and quadrupole temperatures were 250 and 150°C, respectively. The mono- and dihydroxylated PCBs were monitored at molecular ion (M<sup>+</sup>) and M<sup>+</sup> + 2 for methoxy derivatives of tetra-, penta-, and hexachlorinated congeners.

Quantification of PCBs, OH-PCBs, and MeSO<sub>2</sub>-CBs was performed on a Shimadzu GC 14A equipped with an electron capture detector (<sup>63</sup>Ni-ECD) and a split-splitless injector operated in the splitless mode. Nitrogen was used as a carrier and make-up gas. Column conditions were analogous to those described above for GC/MS. Individual PCBs and metabolites were quantified from the relative peak area to internal standards by comparison with authentic standards.

## Results

**Residual PCB Pattern in the Liver.** Table 1 shows the percentages of composition of PCBs classified into four groups, A to D, by chlorination pattern in the livers of rats, hamsters, and guinea pigs.

The original composition of major PCB congeners in Kanechlor 500 was 2,5,2',5'-tetraCB (5.6% of total PCBs), 2,3,6,2',5'-pentaCB (6.5%), 2,4,5,2',5'-pentaCB (10%); 2,3,6,3',4'-pentaCB (7.4%), 2,4,5,3',4'-pentaCB (7.7%), 2,3,4,2',4',5'-hexaCB (5.6%), and 2,4,5,2',4',5'-hexaCB (5.4%). After exposure of animals to Kanechlor 500, 2,5,2',5'-tetraCB, 2,3,6,2',5'-pentaCB, 2,4,5,2',5'-pentaCB, and 2,3,6,3',4'-pentaCB were eliminated rapidly from the livers of all species. Major PCB residues were 2,3,4,2',4',5'-hexaCB (14% of total PCBs) in rats, 2,4,5,2',4',5'-hexaCB (17%) in hamsters, and 2,4,5,3',4'-pentaCB (32%) in guinea pigs. The coplanar PCBs chlorinated at the 4-, 3,4-, or 3,4,5-position (group A, 12% of total PCBs) were detected in the liver in the following order: guinea pigs (61%) > hamsters (26%) = rats (26%). In contrast, the nonplanar PCBs chlorinated at the 2,4- or 2,3,4-position (group B, 24% of total PCBs) were more abundant in the liver in the following order: rats (43%) > hamsters (20%) > guinea pigs (11%). Highly chlorinated PCBs with 2,4,5-chlorine substitution (group C, 8.5% of total PCBs) were more




Fig. 2. GC/ECD chromatograms of phenolic PCBs (after methylation) in the serum of rats (A), guinea pigs (B), and hamsters (C) after exposure to Kanechlor 500 (100 mg/kg i.p.). The phenolic PCB fraction was spiked with 2,3,4,5,6,3',4',5'-octaCB (internal standard) after separation of PCBs and then methylated with diazomethane. See Table 2 for abbreviation on the peak.

TABLE 2

Levels of MeSO<sub>2</sub>-PCBs in the livers of rats, guinea pigs, and hamsters dosed with Kanechlor 500

Each value represents the mean ± S.E. for three to four animals.

Abbreviation	Structure	Levels (µg/g wet wt)		
		Rat	Guinea Pig	Hamster
<b>2,5-Dichloro-substituted</b>				
3'-49	3'-MeSO <sub>2</sub> -2,4,2',5'-tetraCB	17 ± 2	60 ± 17	N.D.
4'-49	4'-MeSO <sub>2</sub> -2,4,2',5'-tetraCB	8 ± 2	N.D.	N.D.
3-70	3-MeSO <sub>2</sub> -2,5,3',4'-tetraCB	50 ± 11	6 ± 2	6 ± 1
4-70	4-MeSO <sub>2</sub> -2,5,3',4'-tetraCB	41 ± 6	N.D.	20 ± 7
3'-87	3'-MeSO <sub>2</sub> -2,3,4,2',5'-pentaCB	32 ± 8	160 ± 44	7 ± 2
4'-87	4'-MeSO <sub>2</sub> -2,3,4,2',5'-pentaCB	27 ± 4	70 ± 7	5 ± 2
3'-101	3'-MeSO <sub>2</sub> -2,4,5,2',5'-pentaCB	43 ± 10	32 ± 4	42 ± 10
4'-101	4'-MeSO <sub>2</sub> -2,4,5,2',5'-pentaCB	48 ± 5	5 ± 2	25 ± 5
3'-141	3'-MeSO <sub>2</sub> -2,3,4,5,2',5'-hexaCB	2 ± 1	33 ± 4	N.D.
4'-141	4'-MeSO <sub>2</sub> -2,3,4,5,2',5'-hexaCB	3 ± 1	N.D.	N.D.
	Total	271 ± 44	366 ± 72	105 ± 16*
<b>2,3,6-Trichloro-substituted</b>				
5-64	5-MeSO <sub>2</sub> -2,3,6,4'-tetraCB	5 ± 1	130 ± 26	N.D.
4-64	4-MeSO <sub>2</sub> -2,3,6,4'-tetraCB	6 ± 1	N.D.	N.D.
5-91	5-MeSO <sub>2</sub> -2,3,6,2',4'-pentaCB	4 ± 1	77 ± 19	15 ± 4
4-91	4-MeSO <sub>2</sub> -2,3,6,2',4'-pentaCB	2 ± 1	N.D.	9 ± 2
5-110	5-MeSO <sub>2</sub> -2,3,6,3',4'-pentaCB	N.D.	6 ± 2	N.D.
4-110	4-MeSO <sub>2</sub> -2,3,6,3',4'-pentaCB	42 ± 9	N.D.	5 ± 3
5'-132	5'-MeSO <sub>2</sub> -2,3,4,2',3',6'-hexaCB	20 ± 4	146 ± 42	2 ± 1
4'-132	4'-MeSO <sub>2</sub> -2,3,4,2',3',6'-hexaCB	21 ± 3	N.D.	N.D.
5'-149	5'-MeSO <sub>2</sub> -2,4,5,2',3',6'-hexaCB	14 ± 2	110 ± 21	2 ± 1
4'-149	4'-MeSO <sub>2</sub> -2,4,5,2',3',6'-hexaCB	17 ± 3	N.D.	3 ± 1
	Total	131 ± 20	469 ± 82	36 ± 8
	Sum <i>meta</i> -MeSO <sub>2</sub> -CBs	187 ± 50	760 ± 93*	74 ± 16
	Sum <i>para</i> -MeSO <sub>2</sub> -CBs	215 ± 71	75 ± 13.3	67 ± 11
	Total MeSO <sub>2</sub> -CBs	402 ± 90	835 ± 184*	141 ± 42*
Ratio	<i>meta/para</i> -MeSO <sub>2</sub> -CBs	0.87	10.1	1.10
	2,5-/2,3,6-chloro substituted	2.07	0.78	2.92

N.D., not detected (&lt;0.1 µg/g).

\* Significantly different from rats,  $P < 0.01$  (Student's *t* test).

resistant to metabolism in the order hamsters (34%) > rats (21%) > guinea pigs (15%). Nonpersistent PCBs with 2,5- or 2,3,6-chlorine substitution (group D, 33% of total PCBs) were eliminated more rapidly in rats (2.7%) and guinea pigs (3.5%) than in hamsters (8.6%).

**MeSO<sub>2</sub>-CBs in the Liver.** Figure 1 shows the ECD gas chromatograms of the MeSO<sub>2</sub> fraction of liver extracts of the three species treated with Kanechlor 500. GC/MS analysis showed that most of the MeSO<sub>2</sub>-CBs detected were originated from 2,5-di- or 2,3,6-trichloro-substituted congeners. The levels and *meta/para* substitution ratios of MeSO<sub>2</sub>-CB congeners in the liver are shown in Table 2. Rats showed preferential production of *meta*- and *para*-substituted MeSO<sub>2</sub> metabolites of 2,5,3',4'-tetraCB, 2,3,4,2',5'-pentaCB, 2,4,5,2',5'-pentaCB, 2,3,6,3',4'-pentaCB, 2,3,4,2',3',6'-hexaCB, and 2,3,6,2',4',5'-hexaCB, with a similar 3-/4-substitution ratio. Hamsters produced minute amounts of *meta*- and *para*-substituted MeSO<sub>2</sub>-CBs derived from 2,5,3',4'-tetraCB and 2,4,5,2',5'-pentaCB. In contrast, guinea pigs showed selective production of *meta*-substituted MeSO<sub>2</sub>-CBs derived from 2,3,6,4'-tetraCB, 2,3,6,2',5'-pentaCB, 2,4,5,2',5'-pentaCB, 2,3,4,2',3',6'-hexaCB, 2,3,4,5,2',5'-hexaCB, and 2,3,6,2',4',5'-hexaCB, although 2,3,4,2',5'-pentaCB yielded 3'- and 4'-MeSO<sub>2</sub>-2,3,4,2',5'-pentaCBs at a similar concentration ratio (Fig. 1). Total levels of MeSO<sub>2</sub>-CBs were higher in the order guinea pigs > rats > hamsters. The concentration ratios of *meta*- and *para*-MeSO<sub>2</sub>-CBs were the highest (10.1) for guinea pigs, whereas the ratios were 0.87 and 1.10 for rats and hamsters, respectively. On the other hand, marked and selective retention of 4'-MeSO<sub>2</sub>-2,4,5,2',5'-pentaCB was observed in the lungs of rats (4'-/3'-MeSO<sub>2</sub> substitution ratio = 11), although no selective retention of *para*-MeSO<sub>2</sub>-CBs was observed in the lungs of guinea pigs or hamsters (data not shown).

**OH-PCBs in Serum.** Figure 2 shows the retention profiles of phenolic PCBs in serum of the three species after exposure to Kanechlor 500. The concentrations of major OH-PCBs and the metabolite/parent CB ratios are shown in Table 3. Rat serum showed specific retention of 4-OH-2,3,5,3',4'-pentaCB (89% contribution), most likely derived from 2,3,4,3',4'-pentaCB and/or 2,4,5,3',4'-pentaCB. On the other hand, guinea pig serum showed specific retention of 3-OH-2,4,5,2',4'-pentaCB (56%) and 3-OH-2,4,5,3',4'-pentaCB (20%). In contrast, hamster serum retained primarily 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCB, 4,5-(OH)<sub>2</sub>-2,3,6,3',4'-pentaCB, and 4,5-(OH)<sub>2</sub>-2,3,6,2',4',5'-hexaCB, as well as 3- or 4-hydroxy metabolites derived from 2,4,5,2',4'-pentaCB and 2,4,5,3',4'-pentaCB (Fig. 2). Thus, phenolic PCB levels showed the following order: hamsters > rats > guinea pigs. The ratios of 3-OH-2,4,5,2',4'-pentaCB/2,4,5,2',4'-pentaCB were higher in hamsters (12.3) and guinea pigs (12.0) than in rats (0.3). The ratio of (4-OH-2,3,5,3',4'-pentaCB + 3-OH-2,4,5,3',4'-pentaCB)/(2,3,4,3',4'-pentaCB + 2,4,5,3',4'-pentaCB) was higher in rats (6.3) than in guinea pigs (0.6).

**Metabolites in the Serum of Guinea Pigs.** The phenolic fraction in the serum of guinea pigs contained several persistent metabolites with different molecular weights from OH-PCBs or MeSO<sub>2</sub>-CBs, although no such metabolites were detected in rats and were formed at lower levels in hamsters. Figure 3 shows the GC/MS total ion chromatogram of phenolic fraction (M<sub>a</sub>-M<sub>d</sub>, at longer GC retention times than MeSO<sub>2</sub>-CBs) in the serum of guinea pigs after exposure to Kanechlor 500. The electron impact mass spectra of M<sub>a</sub> to M<sub>d</sub> are shown in Fig. 4. By GC/ECD and GC/MS analyses, M<sub>a</sub> was tentatively identified as methoxy-MeSO<sub>2</sub>-tetraCB (M<sup>+</sup>, *m/z* = 398), and M<sub>d</sub> as a methoxy-MeSO<sub>2</sub>-pentaCB (M<sup>+</sup>, *m/z* = 432) due to a frag-

TABLE 3

Serum concentration of phenolic PCBs in rats, guinea pigs, and hamsters after dosing with Kanechlor 500 (100 mg/kg i.p.)

Each value represents the mean  $\pm$  S.E. for three to five animals.

PCB and Metabolite Congeners	Serum Concentration (ng/g wet wt)		
	Rat	Guinea Pig	Hamster
2,4,5,2',4'-pentaCB (CB99)	33 $\pm$ 3	12 $\pm$ 2	8 $\pm$ 2
3-OH-2,4,5,2',4'-pentaCB	9 $\pm$ 2	144 $\pm$ 40	74 $\pm$ 22
4-OH-2,3,5,2',4'-pentaCB	N.D.	N.D.	24 $\pm$ 8
Ratio of OH-PCBs/CB99	0.27 $\pm$ 0.22	12.0 $\pm$ 2.1*	12.3 $\pm$ 2.1*
2,3,4,3',4'-pentaCB (CB105)	15 $\pm$ 3	20 $\pm$ 3	14 $\pm$ 2
2,4,5,3',4'-pentaCB (CB118)	56 $\pm$ 6	87 $\pm$ 14	50 $\pm$ 14
3-OH-2,4,5,3',4'-pentaCB	12 $\pm$ 3	51 $\pm$ 10	37 $\pm$ 6
4-OH-2,3,5,3',4'-pentaCB	342 $\pm$ 75	N.D.	111 $\pm$ 16
Ratio of OH-PCBs/(CB105 + CB118)	6.3 $\pm$ 0.9	0.59 $\pm$ 0.1*	3.0 $\pm$ 0.5*
2,3,4,2',4',5'-hexaCB (CB138)	93 $\pm$ 8	38 $\pm$ 7	37 $\pm$ 3
3'-OH-2,3,4,2',4',5'-hexaCB	14 $\pm$ 4	33 $\pm$ 6	24 $\pm$ 6
4'-OH-2,3,4,2',3',5'-hexaCB	N.D.	N.D.	11 $\pm$ 3
Ratio of OH-PCBs/CB138	0.15 $\pm$ 0.02	0.87 $\pm$ 0.11*	0.95 $\pm$ 0.14*
2,3,5,2',4',5'-hexaCB (CB146)	13 $\pm$ 3	9.1 $\pm$ 2.0	6.2 $\pm$ 2.1
2,4,5,2',4',5'-hexaCB (CB153)	80 $\pm$ 15	74 $\pm$	55 $\pm$
3-OH-2,4,5,2',4',5'-hexaCB	9 $\pm$ 3	28 $\pm$ 8	4 $\pm$ 1
4-OH-2,3,5,2',4',5'-hexaCB	N.D.	N.D.	3 $\pm$ 1
Ratio of OH-PCBs/(CB146 + CB153)	0.11 $\pm$ 0.02	0.38 $\pm$ 0.06*	0.13 $\pm$ 0.02
3',4'-dihydroxy-2,4,5,2',5'-pentaCB (diOH-CB101)	N.D.	9 $\pm$ 3	42 $\pm$ 6
4,5-dihydroxy-2,3,6,3',4'-pentaCB (diOH-CB110)	N.D.	N.D.	84 $\pm$ 10
4,5-dihydroxy-2,3,6,2',4',5'-hexaCB (diOH-CB149)	N.D.	N.D.	61 $\pm$ 9
Total phenolic PCBs	386 $\pm$ 29	256 $\pm$ 32*	475 $\pm$ 49*

N.D., not detected ( $<0.005 \mu\text{g/g}$ ).\* Significantly different from rats,  $P < 0.01$  (Student's *t* test).

ment ion,  $[M - 81]^+$  (due to loss of oxygen and  $\text{OCH}_2\text{Cl}$  from  $M^+$ ), and  $[M - 144]^+$  (loss of  $\text{OCH}_2\text{Cl-O-SOCH}_2$  from  $M^+$ ).  $M_b$  showed a weak molecular ion ( $m/z = 400$ ) and an abundant fragment at  $[M - 32]^+$  (loss of two oxygen atoms from  $M^+$ ), which was characteristic of vicinal methoxy-methylthio-pentaCB.  $M_c$  showed a weak molecular ion ( $m/z = 416$ ), and fragment ion  $[M - 16]^+$  (loss of oxygen) as a base peak that was characteristic of vicinal methoxy-methylsulfinyl pentaCBs. Thus, the PCB metabolites in the serum of guinea pigs were tentatively identified as hydroxy-MeSO<sub>2</sub>-tetraCB for  $M_a$ , hydroxy-methylthio-pentaCB for  $M_b$ , hydroxy-methylsulfinyl-pentaCB for  $M_c$ , and hydroxy-MeSO<sub>2</sub>-pentaCB for  $M_d$ .

### Discussion

Although oxidative metabolism of PCBs by rats has been studied extensively (Kato et al., 1980; Kaminsky et al., 1981), there has been




FIG. 3. Total ion chromatogram of GC/MS (electron impact mode) of phenolic PCBs (after methylation) in guinea pigs after exposure to Kanechlor 500 (100 mg/kg i.p.). The phenolic fraction was spiked with 2,3,4,5,6,3',4',5'-octaCB (internal standard) after separation of PCBs.

only limited comparative analysis of in vivo metabolism of PCBs between hamsters and guinea pigs (Koga and Yoshimura, 1996). The results of the present study indicated that the tissue distribution profiles of PCB residues and OH- and MeSO<sub>2</sub>-CBs were largely dependent on the chlorination patterns classified into four groups. This may have been due to differences in substrate specificity of each species, especially between guinea pigs and hamsters, following administration of Kanechlor 500.

**PCB Residues in the Liver.** The percentage composition of residual PCBs in the liver indicated that guinea pigs had lower elimination ratios of coplanar PCBs with 4-, 3,4-, or 3,4,5-chlorine substitution (group A in Table 1) than rats or hamsters. For example, the composition of 3,3',4,4'-tetraCB, which originally accounted for 0.03% in Kanechlor 500, increased to 1.8% in the livers of guinea pigs 5 days after exposure, whereas the values were 0.02 and 0.06% in the livers of rats and hamsters, respectively (Table 1). These observations suggest that guinea pigs have lower ability to metabolize coplanar PCBs. This was supported by the in vitro observation that 3,3',4,4'-tetraCB can hardly be metabolized by liver microsomes from MC or phenobarbital-treated guinea pigs, whereas it is metabolized effectively by those of rats and hamsters treated with MC (Koga et al., 1995). The reduced elimination of coplanar PCBs may partially explain the higher sensitivity to the toxicity of coplanar PCBs in guinea pigs.

On the other hand, hamsters exhibited reduced elimination of the 2,4,5-trichloro-substituted CB congeners (group C) compared with rats or guinea pigs. This observation suggests that hamsters have a reduced capability to hydroxylate 2,4,5-trichloro-substituted CBs than the other two species. The results of a previous in vitro study indicated that 2,4,5,2',4',5'-hexaCB can hardly be metabolized by rats or hamsters, whereas it is metabolized effectively by guinea pigs (Ariyoshi et al., 1997). This may be due to the lower levels of induction of CYP2B1/2 in rats and hamsters compared with guinea pigs. CYP3A may also be related to the metabolism of these PCBs, depending on species. Furthermore, it is noted that nonpersistent CBs with 2,5- or




FIG. 4. Mass spectra of  $M_a$  through  $M_d$  in total ion chromatogram (Fig. 3) of phenolic metabolites (as methyl derivatives) from the serum of guinea pigs after exposure to Kanechlor 500.

2,3,6-chlorine substitution (group D) tend to be eliminated more slowly in hamsters than in rats or guinea pigs. The reduced elimination of these congeners is probably due to the lower catalytic activity of CYP2B in hamsters than in rats (Ishida et al., 1991; Koga et al., 1995), or to the lower abilities of phase II isozymes, such as glutathione *S*-transferase, to conjugate for subsequent excretion in hamsters.

**MeSO<sub>2</sub>-CBs in Liver.** The MeSO<sub>2</sub> metabolites detected in the present study were consistent with the results of previous studies regarding metabolism of PCB congeners with 2,5- or 2,3,6-chlorine substitution (Haraguchi et al., 1997, 1999). However, the levels and metabolite profiles in guinea pigs and hamsters were quite different from those in rats. Hamsters formed minute amounts of *meta*- and *para*-MeSO<sub>2</sub>-CBs from 2,5,3',4'-tetraCB and 2,4,5,2',5'-pentaCB, whereas guinea pigs produced much higher levels of MeSO<sub>2</sub>-CBs and showed selective distribution as *meta*-substituted-MeSO<sub>2</sub>-CBs derived from CBs with 2,3,6-chlorine substitution. These results indicate that the *meta*-position of the 2,3,6-trichloro-substituted phenyl ring in the molecule is the preferred site for methylsulfonylation and guinea pigs may have substrate-specific phase II isozymes that catalyze the nucleophilic reaction of arene oxide intermediates with sulfhydryl groups. Indeed, the selective retention of *meta*-MeSO<sub>2</sub>-CBs has been observed in the livers of otters, minks, and seals in Canadian and Swedish water (Bergman et al., 1994b). Especially, 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCB was the most abundant in human tissues (Chu et al., 2003). Kato et al. (1997) reported that only *meta*-substituted MeSO<sub>2</sub>-CBs strongly induced CYP2B1/2 and reduced the serum thyroid hormone level in rats. Exposure of minks to a synthetic

mixture of *meta*- and *para*-MeSO<sub>2</sub>-CBs also resulted in the selective retention of *meta*-MeSO<sub>2</sub>-CBs in the liver and increased induction of some hepatic CYP2B isozymes as well as alteration of thyroid hormone levels in blood (Lund et al., 1999). Higher levels of induction of CYP2B1/2 by MeSO<sub>2</sub>-CBs as well as induction of CYP3A-like activity (Schuetz et al., 1986, 1998) may alter the metabolic pathways of individual PCBs in rats and contribute to the high level of toxicity of nonplanar PCBs.

**Phenolic PCBs in Serum.** For the three species examined in the present study, persistent OH-PCBs in serum were derived metabolically from 2,4,5-trichloro-substituted CBs (e.g., 2,4,5,2',4'-pentaCB, 2,4,5,3',4'-pentaCB, 2,3,4,2',4',5'-hexaCB, and 2,4,5,2',4',5'-hexaCB) via direct insertion of a hydroxyl group into the 3-position or via epoxidation and subsequent NIH chlorine shift at the 4-position. Indeed, all the OH-PCBs detected in the present study have been observed in gray seal and in human blood (Bergman et al., 1994a; Sandau et al., 2000); however, the levels of OH-PCBs and metabolite/parent CB ratios were species-specific since different species have different types of P450 enzymes (Koga et al., 1998).

Exposure of animals to Kanechlor 500 resulted in higher levels of elimination of 2,4,5,3',4'-pentaCB in rats than in guinea pigs or hamsters. For this metabolism, rats showed selective formation of 4-OH-2,3,5,3',4'-pentaCB (metabolite/parent CB ratio = 6.3), whereas guinea pigs showed selective formation of 3-OH-2,4,5,3',4'-pentaCB (ratio = 0.59). Hamsters showed formation of 3-OH-2,4,5,3',4'-pentaCB and 4-OH-2,3,5,3',4'-pentaCB in a ratio of 1:3 (ratio = 3.0). Although serum does not necessarily have the same profile as liver, the differences in the levels of OH-PCBs and bio-



transformation ratios in serum may be explained by the species-dependent metabolic capacities. In fact, these results are consistent with the in vitro observation that 2,4,5,3',4'-pentaCB can be metabolized readily to 4-OH-2,3,5,3',4'-pentaCB (CYP1A1/2) by MC-inducible hepatic microsomes of rats, whereas it shows a lesser degree of metabolism to 2- or 3-OH-2,4,5,3',4'-pentaCB (CYP2B18) by guinea pig liver microsomal P450 isozymes (Koga et al., 2002).

For metabolism of 2,4,5,2',4'-pentaCB, both guinea pigs and hamsters showed selective formation of 3-OH-2,4,5,2',4'-pentaCB (transformation ratios = 12.0 and 12.3, respectively), whereas this PCB was relatively resistant to metabolism by rats (ratio = 0.27). On the other hand, the elimination of 2,4,5,2',4',5'-hexaCB and 2,3,4,2',4',5'-hexaCB was delayed in rats and hamsters, resulting in lower distribution of their metabolites in the liver. These findings were supported by those of previous in vitro studies indicating that 2,3,4,2',4',5'-hexaCB and 2,4,5,2',4',5'-hexaCB can be hydroxylated effectively by the guinea pig cytochrome P450 isoform CYP2B18 but are resistant to metabolism by rats or hamsters (Koga et al., 2001).

Other persistent phenolic PCBs were catechol metabolites derived from CBs with 2,5- or 2,3,6-chlorine substitution (e.g., 2,4,5,2',5'-pentaCB, 2,3,6,3',4'-pentaCB, and 2,3,6,2',4',5'-hexaCB), which were retained in the serum of hamsters at concentrations similar to those of monohydroxylated PCBs. The formation of catechol may involve a two-step direct hydroxylation process or a pathway via arene oxide and dihydrodiol intermediates. Thus, the tissue distribution ratios of catechols in hamsters and of MeSO<sub>2</sub>-CBs in guinea pigs would reflect the extreme differences in metabolism of CBs with 2,5- or 2,3,6-chlorine substitution. Although their mechanisms of retention are still unknown, catechols may act as endocrine disruptors, similar to OH-PCBs, with affinity to the thyroid hormone transport protein transthyretin, concomitant with a reduction in circulating thyroid hormone level (Brouwer et al., 1998) or to estrogen receptor with estrogenicity (Garner et al., 1999).

The phenolic PCB fraction in the serum of Kanechlor 500-treated guinea pigs contained several additional metabolites that were observed in GC/MS total ion chromatograms at longer GC retention times than those of MeSO<sub>2</sub>-CBs (Fig. 3). Two of these, M<sub>3</sub> and M<sub>d</sub>, were identified tentatively as hydroxy-MeSO<sub>2</sub>-tetraCB and -hexaCBs, respectively. The other two, M<sub>b</sub> and M<sub>c</sub>, were presumably the precursors of M<sub>d</sub> because when the fraction was oxidized chemically, the two peaks were converted to the corresponding methylsulfones (M<sub>d</sub>). Mio and Sumino (1985) identified 3-hydroxy-4-methylthio-2,5,2',5'-tetraCB in the feces from rats, mice, and guinea pigs dosed with 2,5,2',5'-tetraCB, suggesting that the precursor of the metabolite was 3,4-epoxide. The epoxide may have been opened by nucleophile attack of the sulfhydryl group and subsequently converted to the hydroxylated metabolite by dehydrogenation (Bakke et al., 1982). Therefore, the metabolites retained in the serum of guinea pigs may be congeners containing a hydroxyl group in the position adjacent to the MeSO<sub>2</sub> group.

### Conclusion

In Kanechlor 500 metabolism, guinea pigs showed reduced elimination of coplanar PCBs with 4-, 3,4-, or 3,4,5-chlorine substitution. In contrast, hamsters showed reduced elimination of 2,4,5-trichloro-substituted PCBs. Guinea pigs exhibited a higher ability for selective metabolism of CBs with 2,3,6-chlorine substitution to *meta*-MeSO<sub>2</sub>-CBs, whereas hamsters showed selective metabolism to catechol metabolites. The reduced elimination rate for coplanar PCBs and the differences in distribution of *meta*-MeSO<sub>2</sub>-CBs, and phenolic and catechol PCBs may have implications for the differences in sensitivity to PCB toxicity between guinea pigs and hamsters.

### References

- Amaro AR, Oakley GG, Bauwer U, Spielmann HP, and Pobertson LW (1996) Metabolic activation of PCBs to quinones: reactivity toward nitrogen and sulfur nucleophiles and influence of superoxide dismutase. *Chem Res Toxicol* 9:623-629.
- Ariyoshi N, Koga N, Yoshimura H, and Oguri K (1997) Metabolism of 2,4,5,2',4',5'-hexachlorobiphenyl (PCB153) in the guinea pig. *Xenobiotica* 27:973-983.
- Bakke JE, Bergman Å, and Larsen GL (1982) Metabolism of 2,4',5-trichlorobiphenyl by the mercapturic acid pathway. *Science (Wash DC)* 217:645-647.
- Bergman Å, Klasson-Wehler B, and Kuroki H (1994a) Selective retention of hydroxylated PCB metabolites in blood. *Environ Health Perspect* 102:464-469.
- Bergman Å, Klasson-Wehler B, Kuroki H, and Nilsson A (1995) Synthesis and mass spectrometry of some methoxylated PCB. *Chemosphere* 30:1921-1938.
- Bergman Å, Norstrom RJ, Haraguchi K, Kuroki H, and Bêland P (1994b) PCB and DDE methyl sulfones in mammals from Canada and Sweden. *Environ Toxicol Chem* 13:121-128.
- Brouwer A, Morse DC, Lans MC, Schuur AG, Murk AJ, Klasson-Wehler B, Bergman Å, and Visser TJ (1998) Interactions of persistent environmental organohalogenes with the thyroid hormone system: mechanisms and possible consequences for animal and human health. *Toxicol Ind Health* 14:59-84.
- Chu S, Covaci A, Jacobs W, Haraguchi K, and Schepens P (2003) Distribution of methyl sulfone metabolites of polychlorinated biphenyls and p,p'-DDE in human tissues. *Environ Health Perspect* 111:1222-1227.
- Connor K, Ramamoorthy K, Moore M, Mustain M, Chen I, Safe S, Zacharewski T, Gillesby B, Joyeux A, and Balaguer P (1997) Hydroxylated polychlorinated biphenyls (PCBs) as estrogens and antiestrogens: structure-activity relationships. *Toxicol Appl Pharmacol* 145:111-123.
- Garner CE, Jefferson WN, Burka LT, Matthews HB, and Newbold RR (1999) In vitro estrogenicity of the catechol metabolites of selected polychlorinated biphenyls. *Toxicol Appl Pharmacol* 154:188-197.
- Haraguchi K, Kato Y, Kimura R, and Masuda Y (1997) Comparative study on formation of hydroxy and sulfur-containing metabolites from different chlorinated biphenyls with 2,5-substitution in rats. *Drug Metab Dispos* 25:845-852.
- Haraguchi K, Kato Y, Kimura R, and Masuda Y (1998) Hydroxylation and methylthiolation of mono-ortho-substituted polychlorinated biphenyls in rats: identification of metabolites with tissue affinity. *Chem Res Toxicol* 11:1508-1515.
- Haraguchi K, Kato Y, Kimura R, and Masuda Y (1999) Tissue distribution of methylsulfonyl metabolites derived from 2,2',4,5,5'-penta- and 2,2',3,4',5',6-hexachlorobiphenyls in rats. *Arch Environ Contam Toxicol* 37:135-142.
- Haraguchi K, Kuroki H, and Masuda Y (1987) Synthesis and characterization of tissue-retainable methylsulfonyl polychlorinated biphenyl isomers. *J Agric Food Chem* 35:178-182.
- Ishida C, Koga N, Hanioka N, Saeki HK, and Yoshimura H (1991) Metabolism in vitro of 3,4,3',4'- and 2,5,2',5'-tetrachlorobiphenyl by rat liver microsomes and highly purified cytochrome P-450. *J Pharmacobio-Dyn* 14:276-284.
- Kaminsky LS, Kennedy MW, Adams SM, and Guengerich FP (1981) Metabolism of dichlorobiphenyls by highly purified isozymes of rat liver cytochrome P-450. *Biochemistry* 20:7379-7384.
- Kato S, McKinney JD, and Matthews HB (1980) Metabolism of symmetrical hexachlorobiphenyl isomers in the rat. *Toxicol Appl Pharmacol* 53:389-398.
- Kato Y, Haraguchi K, Shibahara T, Yumoto S, Masuda Y, and Kimura R (1999) Reduction of thyroid hormone levels by methylsulfonyl metabolites of tetra- and pentachlorinated biphenyls in male Sprague-Dawley rats. *Toxicol Sci* 48:51-54.
- Kato Y, Haraguchi K, Tomiyasu K, Saito H, Isogai M, Masuda Y, and Kimura R (1997) Structure-dependent induction of CYP2B1/2 by 3-methylsulfonyl metabolites of polychlorinated biphenyl congeners in rats. *Environ Pharmacol Toxicol* 3:137-144.
- Kester MH, Bulduk S, van Toor H, Tibboel D, Neini W, Glatt H, Falany CN, Coughtrie MW, Schuur AG, Brouwer A, and Visser TJ (2002) Potent inhibition of estrogen sulfotransferase by hydroxylated metabolites of polyhalogenated aromatic hydrocarbons reveals alternative mechanism for estrogenic activity of endocrine disruptors. *J Chin Endocrinol Metab* 87:1142-1150.
- Kociba RJ and Cabely O (1985) Comparative toxicology and biologic activity of chlorinated dibenzo-p-dioxins and furans relative to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Chemosphere* 14:649-660.
- Koga N, Haraguchi K, Kanamaru T, Kato Y, and Kimura R (2002) Metabolism of 2,3',4,4',5-pentachlorobiphenyl (CB118) by guinea pig liver microsomes. *Organohalogen Compd* 56:17-20.
- Koga N, Kanamaru T, Kikuichi N, Oishi N, Kato S, and Yoshimura H (1995) Effect of cytochrome P450 inducers on liver microsomal metabolism of tetrachlorobiphenyls in rats, guinea pigs and hamsters. *Biol Pharm Bull* 18:705-710.
- Koga N, Kanamaru T, Oishi N, Kato Y, Kimura R, Haraguchi K, and Masuda Y (2001) Species differences in the in vitro metabolism of 2,4,5,2',3',4'-hexachlorobiphenyl. *Fukuoka Acta Med* 92:167-176.
- Koga N, Kikuichi N, Kanamaru T, Kuroki H, Matsusue K, Ishida C, Ariyoshi N, Oguri K, and Yoshimura H (1998) Metabolism of 2,3',4',5-tetrachlorobiphenyl by cytochrome P450 from rats, guinea pigs and hamsters. *Chemosphere* 37:1985-1994.
- Koga N and Yoshimura H (1996) Metabolism of PCBs and related compounds and their toxicity, in *Yusho—a Human Disaster Caused by PCBs and Related Compounds* (Kuratsune M ed), pp 105-120, Kyushu University Press, Fukuoka.
- Letcher RJ, Klasson-Wehler E, and Bergman Å (2000) Methyl sulfone and hydroxylated metabolites of polychlorinated biphenyls, in *The Handbook of Environmental Chemistry Vol 3, Part K* (Paasivirta J ed): *New Type of Persistent Halogenated Compounds* pp 315-319, Springer-Verlag, Berlin, Heidelberg.
- Letcher RJ, Lemmen JG, van der Berg B, Brouwer A, Bergman Å, Giesy JP, and van den Berg M (2002) In vitro antiestrogenic effects of aryl methyl sulfone metabolites of polychlorinated biphenyls and 2,2-bis(4-chlorophenyl)-1,1-dichloroethene on 17 $\beta$ -estradiol-induced gene expression in several bioassay systems. *Toxicol Sci* 69:362-372.
- Lund BO, Örborg J, Bergman Å, Larsson C, Bergman A, Bäcklin BM, Håkansson H, Madej A, Brouwer A, and Brunström B (1999) Chronic and reproductive toxicity of a mixture of 15 methylsulfonyl-polychlorinated biphenyls and 3-methylsulfonyl-2,2-bis(4-chlorophenyl)-1,1-dichloroethene in mink (*Mustela vison*). *Environ Toxicol Chem* 18:292-298.
- Mio T and Sumino K (1985) Mechanism of biosynthesis of methylsulfones from PCBs and related compounds. *Environ Health Perspect* 59:129-135.
- Morse DC, Wehler BK, Wesseling W, Koeman JH, and Brouwer A (1996) Alterations in rat brain thyroid hormone status following pre- and postnatal exposure to polychlorinated biphenyls (Aroclor 1254). *Toxicol Appl Pharmacol* 136:269-279.

- Öberg M, Sjödin A, Casabona H, Nordgren I, Klasson-Wehler E, and Håkansson H (2002) Tissue distribution and half-lives of individual polychlorinated biphenyls and serum levels of 4-hydroxy-2,3,3',4',5-pentachlorobiphenyl in the rat. *Toxicol Sci* **70**:171-182.
- Olson JR, Gasiewicz TA, and Neal RA (1980) Tissue distribution, excretion, and metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in the Golden Syrian hamster. *Toxicol Appl Pharmacol* **56**:78-85.
- Safe SH (1994) Polychlorinated biphenyls (PCBs): environmental impact, biochemical and toxic responses, and implications for risk assessment. *Crit Rev Toxicol* **24**:87-149.
- Sandau CD, Ayotte P, Dewailly E, Duffe J, and Norstrom RJ (2000) Analysis of hydroxylated metabolites of PCBs (OH-PCBs) and other chlorinated phenolic compounds in whole blood from Canadian Inuit. *Environ Health Perspect* **108**:611-616.
- Schuetz EG, Brimer C, and Schuetz JD (1998) Environmental xenobiotics and the antihormones cyproterone acetate and spironolactone use the nuclear hormone pregnenolone X receptor to activate the CYP3A23 hormone response element. *Mol Pharmacol* **54**:1113-1117.
- Schuetz EG, Wrighton SA, Safe SH, and Guzelian PS (1986) Regulation of cytochrome P-450p by phenobarbital and phenobarbital-like inducers in adult rat hepatocytes in primary monolayer culture and in vivo. *Biochemistry* **25**:1124-1133.
- Wroblewski VJ and Olson JR (1988) Effect of monooxygenase inducers and inhibitors on the hepatic metabolism of 2,3,7,8-tetrachlorodibenzo-p-dioxin in the rat and hamster. *Drug Metab Dispos* **16**:43-51.

---

**Address correspondence to:** Dr. Koichi Haraguchi, Daiichi College of Pharmaceutical Sciences, 22-1 Tamagawa-cho, Minami-ku, Fukuoka 815-8511, Japan. E-mail: ko-ichi@ka2.so-net.ne.jp

---

## Species differences in the tissue distribution of catechol and methylsulphonyl metabolites of 2,4,5,2',5'-penta- and 2,3,4,2',3',6'-hexachlorobiphenyls in rats, mice, hamsters and guinea pigs

K. HARAGUCHI<sup>1</sup>, Y. KATO<sup>2</sup>, N. KOGA<sup>3</sup>, & M. DEGAWA<sup>2</sup>

<sup>1</sup>*Daiichi College of Pharmaceutical Sciences, Fukuoka, Japan,* <sup>2</sup>*School of Pharmaceutical Sciences, University of Shizuoka, Shizuoka, Japan,* <sup>3</sup>*Faculty of Nutritional Sciences, Nakamura Gakuen University, Fukuoka, Japan*

(Received 9 August 2004)

### Abstract

Polychlorinated biphenyls (PCBs) are metabolized to phenolic or methylsulphonyl PCBs (MeSO<sub>2</sub>-CBs) in animal species. The study determined the species differences in the tissue distribution of persistent PCB metabolites in rats, mice, hamsters and guinea pigs 4 days after exposure to 2,4,5,2',5'-pentachlorobiphenyl (CB101) or 2,3,4,2',3',6'-hexachlorobiphenyl (CB132). For CB101 metabolism, the hydroxylation in rats, mice and hamsters occurred primarily at the 3'-position in the 2',5'-dichlorinated phenyl ring, whereas the hydroxylation in guinea pigs occurred preferentially at the 3-position. Metabolite profiles in tissues of hamsters were dominated by 3',4'-catechol-CB101, whereas metabolite profiles in rats and mice were dominated by 3'- or 4'-MeSO<sub>2</sub>-CBs. For CB132 metabolism, rats and mice produced 4'- and 5'-MeSO<sub>2</sub>-CBs at similar concentration ratios, whereas guinea pigs produced MeSO<sub>2</sub>-CBs at higher levels and selectively retained 5'-MeSO<sub>2</sub>-CB in liver. In contrast, hamsters preferentially produced 4',5'-catechol-CB132 that was retained in serum. Consequently, hamsters produced catechols, whereas guinea pigs produced *meta*-substituted MeSO<sub>2</sub>-CBs, preferentially from CB132. These findings indicate that PCBs with 2,3,6-chlorine substitution are preferred substrates for the formation of catechols or MeSO<sub>2</sub>-CBs and the differences in metabolite profiles are related to species-dependent metabolic capacities.

**Keywords:** *Species difference, tissue distribution, methylsulphonyl, catechol, polychlorinated biphenyl*

---

Correspondence: K. Haraguchi, Daiichi College of Pharmaceutical Sciences, 22-1 Tamagawa-cho, Minami-ku, Fukuoka 815-8511, Japan. Fax: 81-92-553-5698. E-mail: ko-ichi@ka2.so-net.ne.jp

ISSN 0268-8697 print/ISSN 1360-046X online © 2005 Taylor & Francis Ltd  
DOI: 10.1080/00498250400026456

## Introduction

Non-planar polychlorinated biphenyls (PCBs) with *meta/para*-unsubstituted carbon atoms are major components of technical PCB mixtures and are detected frequently in the environment and biota (Brevik et al., 2002). These PCBs are readily metabolized by cytochrome P450 (CYP)-mediated enzymes to phenolic PCBs via arene oxide intermediates (Forgue et al., 1979) or by direct insertion of a hydroxyl group (Preston et al. 1983). The arene oxides are metabolized further to methylsulphonyl PCBs (MeSO<sub>2</sub>-CBs) via the mercapturic acid pathway (Bakke et al., 1982).

MeSO<sub>2</sub>-CBs have been detected in the liver and adipose tissues of both wildlife (Haraguchi et al., 1992, Bergman et al., 1994, Klasson-Wehler et al., 1998) and humans (Chu et al., 2003). These metabolites, particularly *meta*-substituted MeSO<sub>2</sub>-CBs, have been shown to induce extensively CYP2B-related enzymes in rats (Kato et al., 1995) and to elicit hormone-disrupting effects (Letcher et al., 2002).

Catechols have been shown to be produced by *in vivo* and *in vitro* metabolism of selected PCBs in some species (McLean et al., 1996, Garner et al., 1999). More recently, several catechol PCBs were identified in uridine diphosphate glucuronosyltransferase (UGT)-deficient Gunn rats following administration of a PCB mixture, Kanechlor 500 (Haraguchi et al., 2004). The results indicated that CBs with 2,5- or 2,3,6-chlorine substitution could be preferred substrates for the formation of catechols. Several catechols from lower chlorinated biphenyls had potential oestrogenic activities *in vitro* (Garner et al., 1999) and might be related to perturbation of the endocrine systems (Garner et al., 2000).

An *in vitro* study with 2,5,2',5'-tetraCB metabolism has demonstrated that there are marked differences in the catalytic activity of each P450 (phase I) in hydroxylation among animal species (Koga et al., 1998a); 3-hydroxylation is mediated by guinea pig liver microsomal CYP2B18 (Koga et al., 1998b), whereas 4-hydroxylation is mediated by hamster liver microsomal CYP2A8 (Koga et al., 1996). An *in vitro* metabolism study by Tampal et al. (2002) has demonstrated that some persistent phenolic PCBs in the body are poor substrates for UGTs (phase II) and resist conjugation, whereas the present *in vivo* study in Gunn rats indicated that catechol PCBs might be substrates for UGTs (Haraguchi et al. 2004). The findings suggest that the persistent PCB metabolite profiles in tissues vary largely depending on the activities of phase I or II enzymes in different species or strains.

The study was performed to gain insight into the species differences in the distribution of phenolic PCBs, catechols and MeSO<sub>2</sub>-CBs in the liver and serum after *in vivo* exposure to 2,4,5,2',5'-pentachlorobiphenyl (CB101) and 2,3,4,2',3',6'-hexachlorobiphenyl (CB132) in Wistar rats, ddy mice, Syrian hamsters and Hartley guinea pigs.

## Materials and methods

### Chemicals

CB101 and CB132 were synthesized by the Cadogan (1962) coupling reaction. The 3'-methoxy- and 4'-methoxy-2,4,5,2',5'-pentaCBs, and 4'-methoxy- and 5'-methoxy-2,3,4,2',3',6'-hexaCBs were synthesized by the method of Bergman et al. (1995). 3'- and 4'-MeSO<sub>2</sub>-2,4,5,2',5'-penta-CBs, and 4'- and 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCBs were synthesized as described (Haraguchi et al., 1987). 3'-Methoxy-2,4,5,2',5'-pentaCB was synthesized from 2,5-dichloroaniline and 2,3,6-trichloroanisole. 3',4'-Dimethoxy-2,4,5,2',5'-pentaCB and 4',5'-dimethoxy-2,3,4,2',3',6'-hexaCBs were synthesized from 4-aminoveratrole and chlorobenzene as described by Haraguchi et al. (2004). The purities of isomers were determined to be > 99% by gas chromatography. Phenolic metabolites were

methylated by diazomethane prepared from N-methyl-N'-nitro-N-nitroso-guanidine and 6 M sodium hydroxide solution. Note that diazomethane is potentially toxic by inhalation and therefore should be carefully handled in a well-ventilated place.

#### *Animal treatments*

Male Wistar rats (150–200 g), male ddy mice (27–35 g), male Syrian hamsters (95–120 g) and male Hartley guinea pigs (400–540 g) were obtained from Japan SLC., Inc. (Shizuoka, Japan). Animals were housed three or four per cage with free access to commercial chow and tap water, and maintained under a 12-h dark/light cycle (lights on, 08.00–20.00 hours) in an air-controlled room ( $24.5 \pm 1^\circ\text{C}$ ,  $55\% \pm 5\%$  humidity). Animals received an intraperitoneal injection of CB101 ( $11 \text{ mg kg}^{-1}$ ) and CB132 ( $19 \text{ mg kg}^{-1}$ ) dissolved in Panacete 810 ( $5 \text{ ml kg}^{-1}$ ). Food and water were provided *ad libitum*. All animals were killed by decapitation on day 4 after dosing, and the liver, lung, adipose tissue and blood were removed, weighed and stored at  $-20^\circ\text{C}$  before chemical analysis.

#### *Sample clean-up procedure*

Sample clean-up and quantification were carried out as described (Haraguchi et al., 1998). Briefly, tissue samples were homogenized with acetone/*n*-hexane (2:1, v/v). Three internal standards were added to each extract, which were then applied to a gel permeation column packed with Bio-Beads S-X3 (50 g, Bio-Rad Laboratories, Hercules, CA, USA). Dichloromethane/*n*-hexane (1:1) was used as a mobile phase at a flow rate of  $4 \text{ ml min}^{-1}$ . The metabolite fraction (120–200 ml) was collected and concentrated to dryness. The residue was methylated by diazomethane. The methylated fraction was subjected to a silica gel (1 g, Wako gel S-1, Wako Pure Chemical Industries, Osaka, Japan), eluted with dichloromethane (10 ml). The eluate was analysed without any further purification.

#### *Instruments and quantification*

Structural identification of PCBs and their metabolites was carried out on a GC/MS system (AOC-17, GC-17A, QP-5000, Shimadzu, Co., Ltd, Kyoto, Japan) with a DB-5MS capillary column ( $60 \text{ m} \times 0.25 \text{ mm i.d.}$ ; J&W Scientific, Folsom, CA, USA) in electron ionization — scan mode. Temperature programme:  $100^\circ\text{C}$  (2 min),  $100\text{--}250^\circ\text{C}$  at  $20^\circ\text{C min}^{-1}$ ,  $250\text{--}280^\circ\text{C}$  at  $2^\circ\text{C min}^{-1}$ . Quantification of PCBs and PCB metabolites was performed on a GC-14A (Shimadzu) instrument equipped with a  $^{63}\text{Ni}$  electron capture detector (ECD) with column conditions analogous to those described for GC/MS. The three internal standards, 2,3,4,5,6,3',4',5'-octaCB, 4-OH-2,3,5,6,3',4',5'-heptaCB and 4-methyl-3-MeSO<sub>2</sub>-5,2',3',4',5'-pentaCB, were used for quantification. The phenolic metabolites were analysed as their methoxy derivatives by comparison with authentic standards. Recoveries of 4'-hydroxy-2,3,3',4,5,5'-hexaCB spiked into control liver samples before extraction were 88–92% ( $n=4$ ).

## **Results**

### *CB101 metabolism*

After exposure of four species to CB101, four phenolic PCB metabolites (as methylated derivatives) and two MeSO<sub>2</sub>-CBs were isolated from the extracts of all tissues. By GC/MS

comparison with authentic compounds, these metabolites were identified as 3-OH-, 3'-OH-, 4'-OH- and 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCBs, and 3'-MeSO<sub>2</sub>- and 4'-MeSO<sub>2</sub>-2,4,5,2',5'-pentaCBs. These metabolites were characterized by Haraguchi et al. (1999, 2004).

Figure 1 shows the concentrations of six PCB metabolites in liver and serum of the four species dosed with CB101. In rats and mice, 3'-MeSO<sub>2</sub>- or 4'-MeSO<sub>2</sub>-2,4,5,2',5'-pentaCBs were predominant metabolites in the liver, followed by 3'-OH-2,4,5,2',5'-pentaCB, whereas 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCB was a predominant metabolite in serum, followed by 3'-OH-2,4,5,2',5'-pentaCB. In guinea pigs, 3-OH-2,4,5,2',5'-pentaCB was a predominant metabolite in the liver, followed by 3'-OH-2,4,5,2',5'-pentaCB, and their concentrations were higher than MeSO<sub>2</sub>-CBs. In hamsters, 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCB was a predominant metabolite in both liver and serum.

The concentrations of unchanged CB101 and its metabolites in adipose tissue, liver, lung and serum are summarized in Table I. In rats and mice, MeSO<sub>2</sub>-CBs were retained in adipose tissue, and a lesser extent in lung. In guinea pigs, MeSO<sub>2</sub>-CBs were most abundant in adipose tissues. In hamsters, 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCB constituted more than 90% of total phenolic PCBs in all tissues and the concentration was the highest in adipose tissues, followed by serum. The ratios of catechol to parent CB101 were the highest (1.75) in serum of hamsters.

#### *CB132 metabolism*

After treatment of four species with CB132, three phenolic PCBs (as methylated derivatives) and two MeSO<sub>2</sub>-CBs were detected in all tissues. They were identified as 4'-OH-, 5'-OH- and 4',5'-(OH)<sub>2</sub>-2,3,4,2',3',6'-hexaCBs, and as 4'-MeSO<sub>2</sub>- and 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCBs. The characterization of catechol metabolite was described by Haraguchi et al. (2004).

Figure 2 shows the concentrations of five PCB metabolites in liver and serum of the four species dosed with CB132. In rats and mice, persistent metabolites in the liver were dominated by 4'- or 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCBs, followed by 5'-OH-2,3,4,2',3',6'-hexaCB. In guinea pigs, on the other hand, 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCB was selectively distributed in liver (5'-/4'-MeSO<sub>2</sub>-hexaCB ratio = 4.2). In hamsters, 4',5'-(OH)<sub>2</sub>-2,3,4,2',3',6'-hexaCB was a predominant metabolite in both liver and serum.

The tissue concentrations of unchanged CB132 and its metabolites are summarized in Table II. In rats, mice and guinea pigs, MeSO<sub>2</sub>-CBs were predominantly distributed to adipose tissues with less accumulation in liver and lung, and the lowest concentration in serum. Particularly in the liver of guinea pigs, the sum of MeSO<sub>2</sub>-CB levels was higher than the residual CB132 level (the ratio of MeSO<sub>2</sub>-CBs/CB132 = 1.8). In contrast, hamsters retained 4',5'-(OH)<sub>2</sub>-2,3,4,2',3',6'-hexaCB in adipose tissues at higher levels than rats or mice. The concentration ratios of the catechol to parent CB132 were the highest (5.5) in serum, followed by lung.

#### **Discussion**

The study indicated that metabolism of CB101 and CB132 by the four species examined resulted in considerable variations in the tissue distribution profiles of catechols and MeSO<sub>2</sub>-CBs. After treatment with each PCB, rats, mice and guinea pigs preferentially produced two MeSO<sub>2</sub>-CBs, whereas hamsters preferentially produced a catechol metabolite. In addition, guinea pigs exhibited different metabolite profiles for CB101 and




Figure 1. Concentrations of six PCB metabolites present in the liver and serum of four species after exposure to 2,4,5,2',5'-pentachlorobiphenyl (CB101) ( $11 \text{ mg kg}^{-1}$ , i.p.). Values are the mean  $\pm$  standard error (vertical bars) for  $n=4$ ; diOH-CB101 = 3',4'-(OH)<sub>2</sub>-2,4,5,2',5'-pentaCB.

CB132. The structures of major PCB metabolites distributed in tissues of guinea pigs and hamsters are shown in Figure 3.

#### Metabolic pathways

The metabolism of each PCB involved hydroxylation, catechol formation and methylsulfonylation in all species. Hydroxylation proceeds mainly at the *meta*-position in the 2',5'-dichlorinated phenyl ring for CB101, and in the 2',3',6'-trichlorinated phenyl ring for CB132, probably via direct insertion of a hydroxyl group (Preston et al., 1983). Methylsulfonylation proceeds mainly at the *meta/para* unsubstituted position of the phenyl

Table I. Concentrations and ratios of unchanged PCB, hydroxy- and methylsulfonyl metabolites in tissues of rats, mice, guinea pigs and hamsters 4 days after exposure to 2,4,5,2',5'-pentachlorobiphenyl (CB101, 11 mg kg<sup>-1</sup>, i.p.).

	Concentration ( $\mu\text{g g}^{-1}$ wet weight)			
	Adipose tissue	Liver	Lung	Serum
<b>Rat</b>				
Unchanged CB101	58 ± 21	0.87 ± 0.19	0.23 ± 0.10	0.18 ± 0.03
Σ OH-PCB	0.45 ± 0.14	0.09 ± 0.03	0.08 ± 0.02	0.11 ± 0.03
OH-PCBs/CB101	0.01	0.10	0.35	0.61
Σ MeSO <sub>2</sub> -CB	1.9 ± 0.4	0.11 ± 0.02	0.55 ± 0.17	0.03 ± 0.01
MeSO <sub>2</sub> -CBs/CB101	0.03	0.13	2.39	0.17
<b>Mouse</b>				
Unchanged CB101	22 ± 6.5	1.1 ± 0.7	0.31 ± 0.12	0.31 ± 0.13
Σ OH-PCB	0.20 ± 0.10	0.09 ± 0.03	0.04 ± 0.01	0.09 ± 0.02
OH-PCBs/CB101	0.01	0.08	0.13	0.29
Σ MeSO <sub>2</sub> -CB	2.3 ± 0.9	0.22 ± 0.04	0.99 ± 0.22	0.03 ± 0.01
MeSO <sub>2</sub> -CBs/CB101	0.10	0.20	3.19	0.10
<b>Guinea pig</b>				
Unchanged CB101	8.3 ± 0.3	0.38 ± 0.09	0.08 ± 0.03	0.05 ± 0.02
Σ OH-PCB	0.17 ± 0.02	0.17 ± 0.03	0.05 ± 0.02	0.04 ± 0.01
OH-PCBs/CB101	0.02	0.45	0.63	0.80
Σ MeSO <sub>2</sub> -CB	0.62 ± 0.25	0.06 ± 0.01	0.06 ± 0.02	0.01 ± 0.002
MeSO <sub>2</sub> -CBs/CB101	0.07	0.16	0.75	0.20
<b>Hamster</b>				
Unchanged CB101	10 ± 3.5	0.68 ± 0.24	0.13 ± 0.05	0.12 ± 0.04
Σ OH-PCB	0.52 ± 0.12	0.08 ± 0.02	0.08 ± 0.03	0.21 ± 0.03
OH-PCBs/CB101	0.05	0.12	0.62	1.75
Σ MeSO <sub>2</sub> -CB	0.06 ± 0.02	0.01 ± 0.003	0.002 ± 0.001	0.002 ± 0.001
MeSO <sub>2</sub> -CBs/CB101	0.01	0.01	0.02	0.02

Values are means ± standard errors for  $n=4$ ; Σ OH-PCB, sum of four phenolic metabolites including catechol; Σ MeSO<sub>2</sub>-CB, sum of two MeSO<sub>2</sub>-CBs.

ring for both CBs via the mercapturic acid pathway from arene oxide intermediates (Bakke et al., 1982). Catecholation, on the other hand, involves two possible pathways. One possibility is further oxidation of phenolic CBs, in which the major metabolite 3'-OH-2,4,5,2',5'-pentaCB for CB101 or 5'-OH-2,3,4,2',3',6'-hexaCB for CB132 could be a substrate for formation of the catechol. This hypothesis may be supported by the *in vitro* metabolism of 4-chlorobiphenyl by rat liver microsomes via a second hydroxylation step (McLean et al., 1996). The alternate pathway is hydrolysis of the arene oxide intermediate to dihydrodiol, followed by dehydrogenation to the catechol (McLean et al., 1996, Garner et al., 1999). As both 3',4'-dihydrodiol- and 3',4'-dihydroxy-2,4,5,2',5'-pentaCB have been detected in rats dosed with CB101 (Chen et al., 1976), the dihydrodiol may be oxidized by dehydrogenase to the catechol (Penning et al., 1996).

#### Rats and mice

The metabolism of both CBs in rats and mice involved mainly 3' (5')-hydroxylation as well as 3' (5')- or 4'-methylsulfonylation. The *meta*-hydroxylation has been elucidated for the *in vitro* metabolism of 2,5,2',5'- or 2,5,3',4'-tetraCB in rats (Ishida et al., 1991, Koga et al., 1998a), where 3-hydroxylation was increased in liver microsomes from rats treated with






Figure 2. Concentrations of five PCB metabolites present in the liver and serum of four species after exposure to 2,3,4,2',3',6'-hexachlorobiphenyl (CB132) ( $19 \text{ mg kg}^{-1}$ , i.p.). Values are the mean  $\pm$  standard error (vertical bars) for  $n=4$ ; diOH-CB132 = 4',5'-(OH)<sub>2</sub>-2,3,4,2',3',6'-hexaCB.

phenobarbital. Previously, it was found that the *in vivo* formation rates of phenolic and MeSO<sub>2</sub>-CBs from 2,5-chlorinated PCB congeners in rats were largely dependent on the degree of chlorination in the non-metabolized ring (Haraguchi et al. 1997). In the current study, metabolite profiles in rats and mice dosed with CB101 or CB132 were similar to each other, and dominated by MeSO<sub>2</sub>-CBs. The selective retention of *para*-MeSO<sub>2</sub>-CBs from CB101 or CB132 was observed in the lungs of rats and mice (Klasson-Wehler et al., 1996, Haraguchi et al., 1997), although there was no specificity for guinea pig and hamster lung.

#### Guinea pigs

The metabolite profile in guinea pigs differs from those in other species and between CB101 and CB132. For CB101 metabolism, guinea pigs preferentially produced 3-OH-2,4,5,2',5'-pentaCB rather than 3'-OH-2,4,5,2',5'-pentaCB (Figure 1). Such 3-hydroxylation was

Table II. Concentrations and ratios of unchanged PCB, hydroxy- and methylsulfonyl metabolites in tissues in rats, mice, guinea pigs and hamsters 4 days after exposure to 2,3,4,2',3',6'-hexachlorobiphenyl (CB132, 19 mg kg<sup>-1</sup>, i.p.).

	Concentration ( $\mu\text{g g}^{-1}$ wet weight)			
	Adipose tissue	Liver	Lung	Serum
<b>Rat</b>				
Unchanged CB132	87 $\pm$ 27	0.37 $\pm$ 0.12	0.29 $\pm$ 0.09	0.04 $\pm$ 0.01
$\Sigma$ OH-PCB	0.30 $\pm$ 0.11	0.16 $\pm$ 0.05	0.02 $\pm$ 0.005	0.06 $\pm$ 0.01
OH-PCBs/CB132	0.003	0.43	0.07	1.50
$\Sigma$ MeSO <sub>2</sub> -CB	1.49 $\pm$ 0.61	0.19 $\pm$ 0.03	0.22 $\pm$ 0.10	0.01 $\pm$ 0.003
MeSO <sub>2</sub> -CBs/CB132	0.02	0.51	0.76	0.25
<b>Mouse</b>				
Unchanged CB132	36 $\pm$ 16	0.97 $\pm$ 0.25	0.53 $\pm$ 0.17	0.12 $\pm$ 0.04
$\Sigma$ OH-PCB	0.12 $\pm$ 0.01	0.16 $\pm$ 0.04	0.05 $\pm$ 0.01	0.03 $\pm$ 0.01
OH-PCBs/CB132	0.003	0.16	0.09	0.25
$\Sigma$ MeSO <sub>2</sub> -CB	3.1 $\pm$ 1.1	0.58 $\pm$ 0.15	0.24 $\pm$ 0.11	0.02 $\pm$ 0.003
MeSO <sub>2</sub> -CBs/CB132	0.09	0.60	0.45	0.17
<b>Guinea pig</b>				
Unchanged CB132	11 $\pm$ 4	0.34 $\pm$ 0.13	0.20 $\pm$ 0.12	0.07 $\pm$ 0.02
$\Sigma$ OH-PCB	0.09 $\pm$ 0.01	0.11 $\pm$ 0.05	0.06 $\pm$ 0.02	0.04 $\pm$ 0.01
OH-PCBs/CB132	0.01	0.32	0.30	0.57
$\Sigma$ MeSO <sub>2</sub> -CB	6.9 $\pm$ 3.6	0.63 $\pm$ 0.07	0.12 $\pm$ 0.04	0.03 $\pm$ 0.01
MeSO <sub>2</sub> -CBs/CB132	0.63	1.85	0.60	0.43
<b>Hamster</b>				
Unchanged CB132	14 $\pm$ 4	0.38 $\pm$ 0.08	0.16 $\pm$ 0.05	0.09 $\pm$ 0.03
$\Sigma$ OH-PCB	1.14 $\pm$ 0.25	0.36 $\pm$ 0.15	0.36 $\pm$ 0.05	0.50 $\pm$ 0.09
OH-PCBs/CB132	0.08	0.95	2.25	5.56
$\Sigma$ MeSO <sub>2</sub> -CB	0.13 $\pm$ 0.03	0.03 $\pm$ 0.01	0.004 $\pm$ 0.001	0.02 $\pm$ 0.005
MeSO <sub>2</sub> -CBs/CB132	0.01	0.08	0.03	0.22

Values are means  $\pm$  standard errors for  $n=4$ ;  $\Sigma$  OH-PCB, sum of three phenolic metabolites including catechol;  $\Sigma$  MeSO<sub>2</sub>-CB, sum of two MeSO<sub>2</sub>-CBs.

a minor pathway in the other three species (Figure 1) and has been observed in CB101-treated minks (Klasson-Wehler et al., 1996). Ariyoshi et al. (1997) reported that guinea pig livers can metabolize 2,4,5,2',4',5'-hexaCB to 3-OH-2,4,5,2',4',5'-hexaCB. It is therefore likely that the 3-position of the 2,4,5-trichlorinated phenyl ring could be a preferred site of hydroxylation in guinea pigs. On the other hand, guinea pigs readily metabolize CB132 to the corresponding MeSO<sub>2</sub>-CBs, but metabolize CB101 less readily. In addition, 5'-MeSO<sub>2</sub>-2,3,4,2',3',6'-hexaCB formed from CB132 was selectively retained in liver (*meta/para*-substitution ratio = 4.2) rather than other tissues (the ratio = 1.6 for adipose tissues). The results indicate that guinea pigs may have substrate-selective phase II enzymes for 4,5-epoxide-2,3,6-trichlorinated congeners and/or have high affinity binding proteins for *meta*-substituted MeSO<sub>2</sub>-CBs in liver. Specific retention of *meta*-MeSO<sub>2</sub>-CBs has been observed in the livers of wild seals, otters and minks from Canada and Sweden (Bergman et al., 1994) as well as in human adipose tissues (Chu et al., 2003). According to Kato et al. (1995), only *meta*-substituted MeSO<sub>2</sub>-CBs have been shown to induce CYP2B-related enzymes strongly in rats, indicating that the retention of 5-MeSO<sub>2</sub>-2,3,6-trichlorinated PCBs may contribute significantly to the toxicity of the parent CBs in guinea pigs.




Figure 3. Chemical structures of major PCB metabolites distributed in tissues of guinea pigs and hamsters after exposure to 2,4,5,2',5'-pentachlorobiphenyl (CB101) or 2,3,4,2',3',6'-hexachlorobiphenyl (CB132).

### Hamsters

In contrast to guinea pigs, hamsters metabolize CB101 and CB132 primarily to their catechol PCBs rather than MeSO<sub>2</sub>-CBs. Although the formation ratio of the catechol to parent CB for CB101-treated hamsters was comparable with the ratios for the other species, CB132-treated hamsters produced the highest amounts of a catechol, which was retained in tissues, especially in adipose tissues and serum. This suggests that CBs with 2,3,6-chlorine substitution are more preferred substrates than CBs with 2,5-chlorine substitution for the formation of catechols in hamsters. The ratios of 3' (5')- to 4'-hydroxylation for both CBs were about 5:1 in the liver of hamsters. This is in contrast to the ratio of 1:8 in the *in vitro* metabolism of 2,5,3',4'-tetraCB by hamster liver microsomes (Ishida et al., 1991, Koga et al., 1998a).

### Mechanism of catechol retention

In Haraguchi et al. (2004), UGT1A-deficient Gunn rats exposed to CB101 (112 mg kg<sup>-1</sup>, i.p.) showed preferential distribution of a catechol PCB metabolite as compared with Wistar rats, which was present in serum at higher concentrations than parent CB101. The distribution profiles of catechol PCBs in Gunn rats may resemble those in hamsters in this study. The retention of catechols in Gunn rats was explained by poor conjugation ability due to UGT deficiency. However, hamsters exposed to PCBs potentially induce UGTs to a similar extent to Wistar rats (Kato et al., 2003), indicating that UGT activities may not be crucial factors for the increased amounts of catechol PCB metabolite. Another retention

mechanism for catechols, such as binding affinity to transthyretin (Lans et al., 1993) or to oestrogen receptors (Garner et al., 2000), cannot be excluded. Taken together, the formation and retention of catechols and MeSO<sub>2</sub>-CBs may be associated with species-different phase II enzymes. The potential effects of the catechol metabolites in the blood are still unknown, even though the catechols from lower chlorinated congeners have been shown to exhibit oestrogenic activity (Garner et al., 1999) and could be precursors for quinone products that show potent cytotoxicity (Srinivasan et al., 2001).

### Conclusion

Rats and mice preferentially produced MeSO<sub>2</sub>-CBs rather than catechols from both CBs, whereas hamsters and guinea pigs showed different metabolite profiles to each other and between CB101 and CB132. Hamsters preferentially produced a catechol from CB132 rather than from CB101, which was retained in serum or adipose tissues at higher concentrations, whereas guinea pigs formed *meta*-MeSO<sub>2</sub>-CBs selectively from CB132, but not much from CB101. The findings indicate that PCBs with 2,3,6-chlorine substitution could be the preferred substrates to yield catechols or MeSO<sub>2</sub>-CBs, and the differences in the metabolite profiles are related to the species-dependent metabolic capacities in the liver.

### Acknowledgements

Research was partially funded by a Grant-in-Aid for Scientific Research (C) (Grant Nos 16590101 K.H.; 12680549 Y.K.; 14572119 N.K.) from the Japan Society for the Promotion of Science and a Health and Labour Sciences Research Grants (Y. K.) from the Ministry of Health, Labour and Welfare of Japan.

### References

- Ariyoshi N, Koga N, Yoshimura H, Oguri K. 1997. Metabolism of 2,4,5,2',4',5'-hexachlorobiphenyl (PCB153) in guinea pig. *Xenobiotica* 27:973-983.
- Bakke JE, Bergman Å, Larsen GL. 1982. Metabolism of 2,4',5-trichlorobiphenyl by the mercapturic acid pathway. *Science* 217:645-647.
- Bergman Å, Klasson-Wehler E, Kuroki H, Nilsson A. 1995. Synthesis and mass spectrometry of some methoxylated PCBs. *Chemosphere* 30:1921-1938.
- Bergman Å, Norstrom RJ, Haraguchi K, Kuroki H, Béland P. 1994. PCB and DDE methyl sulphones in mammals from Canada and Sweden. *Environmental Toxicology and Chemistry* 13:121-128.
- Brevik K, Sweetman A, Pacyna JM, Jones KC. 2002. Towards a global historical emission inventory for selected PCB congeners — a mass balance approach. 1. Global production and consumption. *Science of the Total Environment* 290:181-198.
- Cadogan JIG. 1962. A convenient new method of aromatic arylation. *Journal of the Chemical Society* 4257-4258.
- Chen PR, McKinney JD, Matthews HB. 1976. Metabolism of 2,4,5,2',5'-pentachlorobiphenyl in the rat. Qualitative and quantitative aspects. *Drug Metabolism and Disposition* 4:362-367.
- Chu S, Covaci A, Jacobs W, Haraguchi K, Schepens P. 2003. Distribution of methyl sulphone metabolites of polychlorinated biphenyls and p,p'-DDE in human tissues. *Environmental Health Perspectives* 111:1222-1227.
- Forgue ST, Preston BD, Hargraves WA, Reich IL, Allen JR. 1979. Direct evidence that an arene oxide is a metabolic intermediate of 2,2',5,5'-tetrachlorobiphenyl. *Biochemical and Biophysical Research Communications* 91:475-483.
- Garner CE, Burka LT, Etheridge AE, Matthews HB. 2000. Catechol metabolites of polychlorinated biphenyls inhibit the catechol-O-methyltransferase-mediated metabolism of catechol estrogens. *Toxicology and Applied Pharmacology* 162:115-123.