

TOX - AH Receptor and AH-Receptor-Dependent Signaling - I

spleen (CFU-S). Aliquots of BM cell suspensions were used to evaluate the number of CFU-S. The number of BM cells was adjusted to that appropriate for producing nonconfluent spleen colonies, and the cells were then transplanted into lethally irradiated mice by injection through the tail vein. Spleens were harvested 9 and 13 days after the injection, and fixed in Bouin's solution. Macroscopic spleen colonies were counted under an inversion microscope at a magnification of x 5.6.

CFU-GM and CFU-E Assay. *in vitro* colony formation was assayed in semisolid methylcellulose culture^{6, 8}. Briefly, 8×10^4 BM cells suspended in 100 ml of medium were added to 3.9 ml of a culture medium containing 0.8% methyl cellulose, 30% fetal calf serum, 1% bovine serum albumin, 10^{-4} M 2-mercaptoethanol, with 10 ng/ml murine granulocyte-macrophage colony-stimulating factor (GM-CSF) for CFU-GM or 1ng/ml murine Interleukin-3 and 2 U/ml erythro-poietin for erythroid CFU (CFU-E). One-ml aliquots containing 2×10^4 BM cells were plated in triplicate in a 35-mm tissue-culture plate, and incubated for six days in a completely humidified incubator at 37 °C with 5% CO₂ in air. Colonies were counted under an inverted microscope at magnifications of x 40 for CFU-GM after 6-day culture and x100 for CFU-E after 3-day culture.

BM repopulation assay⁹. The BM repopulation assay was performed similarly to the assay of CFU-S, except that 10^6 BM cells were injected into lethally irradiated mice. One month after the transfusion of BM cells, the repopulated mice were used in the experiment.

Results and Discussion

As previously reported, AhR-KO mouse showed a significant increase in WBC counts (Figure 1 A). This was also consistent with the high number of myeloid progenitor cells, *i.e.*, CFU-S-9 and CFU-S-13, observed in the AhR-KO mice (Figure 1, B). Thus, steady-state hemopoiesis is presumed to be suppressed by AhR signaling due to the possible presence of a physiological ligand, which is not readily observed in AhR-KO mice. In response to such an AhR-null effect, the AhR-KO mouse reversely shows extensive hemopoiesis in the spleen, although this hemopoietic enhancement is also reflected in another negative hemopoietic regulation in the BM. Accordingly, in the present study, benzene-induced hematotoxicity was evaluated in the Wt mice after a lethal dose of whole-body irradiation followed by the repopulation of BM cells that lack AhR.




Figure 1: Comparison of various blood parameters between Wt mice (open columns) and AhR-KO mice (shaded columns)¹. A. Peripheral blood, bone marrow and spleen weight. * Parameters indicate the counts of peripheral red blood cells (RBCs, $\times 10^6/\text{ml}$) and white blood cells (WBCs, $\times 10^6/\text{ml}$), bone marrow cellularity (BM, $\times 10^5/\text{femur}$), and weight of spleen (SPL, mg). B. Number of colony-forming units in spleen (CFU-S/ 1×10^5 BM cells) observed on days 9 (CFU-S-9) and 13 (CFU-S-13). C. Numbers of *in vitro* granulocyte-macrophage CFUs (CFU-GM/ 5×10^3 BM cells) and erythroid CFU (CFU-E/ 1×10^4 BM cells). †: Significant difference between Wt and AhR-KO mice determined by *t*-test at $p < 0.05$.

Figures 2, A-C, show the RBC (A), WBC (B), and platelet (PLT: C) counts (per mL) in the peripheral blood after repopulation of the BM. In each figure, in the Wt mice repopulated with Wt BM cells (two columns on the left), the groups subjected to intraperitoneal benzene exposure (second from the left) show significant decreases in RBC and PLT counts (92% and 69%; $p=0.010$ and 0.016 , respectively) compared with the sham exposure groups (farthest left in each figure), except 2B, *i.e.*, WBC counts (96%). When the mice repopulated with AhR-KO BM cells (two columns on the right) are exposed to benzene, there are no significant differences between the sham exposure groups (second from the right) and the benzene exposed groups (farthest right) in A through C. Significant decreases

TOX - Ah Receptor and Ah-Receptor-Dependent Signaling - I

observed in the Wt mice repopulated with Wt BM cells were negated when the Wt mice were repopulated with AhR-KO BM cells; thus, the reduction in the number of peripheral blood cells observed in the Wt mice after benzene exposure is assumed to be responsible for the AhR expression in BM cells. Although the two sham exposures (i.e., Wt mice, open column; and AhR-KO mice, solid column) are essentially identical in A and C, there seems to be insufficient recovery of the BM in transplantation in Figure 2B, and the solid column is significantly reduced (see, Figure 3 on CFU-GM).




Figure 2: Comparison of various blood parameters in peripheral blood; A, RBC, B, WBC, and C, platelets (open bars vs lightly shaded bars in Wt mice repopulated with Wt BM cells; solid bars vs heavily shaded bars in Wt mice repopulated with AhR BM cells).





Figure 3: *: Significant difference between sham and exposed determined by *t*-test at $p < 0.05$.

In Figure 3, again, the significant decrease in the number of granulocyte-macrophage colony-forming units *in vitro* (CFU-GM/5 x10³ BM cells) in the BM cells from the Wt mice repopulated with Wt BM cells (82.9% in benzene exposure, lightly shaded column to the right of the sham exposure, open leftmost column; $p=0.041$) is negated in the BM cells from mice repopulated with AhR-KO BM cells (sham exposure, solid column; and benzene exposure, heavily shaded column, respectively). In this figure, the efficiency of repopulation with AhR-KO BM cells (solid column) seems to be insufficient, since the solid column is smaller than the open column ($p=0.025$). The mechanism underlying the incomplete recovery of AhR-KO BM cells, is still unknown; however, the sublethal irradiation of the recipient mice may be the case, where suppressive intrinsic factors may have been released from tissues given the lethal dose of irradiation received by the host animals.

Despite the insufficient recovery of the number of GM-CFU in mice repopulated with AhR-KO BM cells, number of BM cells, regardless of repopulated cell type (either Wt or AhR-KO BM cells) and type of exposure (either benzene or sham exposure), there were no significant differences in number of BM cells among the groups in a homeostatic manner (Figure 4, A; 91.4% and 92.4%, respectively, $p > 0.1$). However, after benzene exposure, a significant decrease in splenic weight was observed in the Wt→Wt group (85.0%, $p=0.022$), but not in the AhR-KO-BM→Wt group (90.0%, $p=0.173$). This supports the notion that AhR-KO negates the suppressive effect on splenic weight after benzene exposure.

Figure 4: Comparison of number of BM cells (A) or weight of spleen (B) with or without benzene exposure, in mice repopulated with Wt BM cells or in mice repopulated with AhR-KO BM cells. (Wt mice repopulated with Wt BM cells, lightly shaded columns, second from left, or without benzene exposure, open columns, farthest left, benzene exposure vs Wt mice repopulated with AhR-KO BM cells, each of the two right columns).



Conclusions

The up-regulation of CYP2E1 after benzene exposure was specifically observed in our previous microarray study of the bone marrow tissue⁴. The analysis of the gene expression specifically derived from the hematopoietic stem cell compartment¹⁰, and the evaluation of the toxicological alteration of such an expression as a measure of stem cell specific toxicological biomarkers are hot issues in the current hematotoxicology¹¹. Mice that have been lethally irradiated and repopulated with BM cells from AhR-KO mice essentially did not show any benzene-induced hematotoxicity, implying that such toxicity is derived from *de novo* metabolisms with CYP2E1 in the BM other than hepatic metabolism. The present study raises two questions on AhR-mediated TCDD-induced hematotoxicity: Do Wt mice repopulated with AhR-KO BM cells show hematotoxicity by TCDD unlike in the case of benzene exposure? If such is the case, what would be the transmitter from the site of xenobiotic metabolic activation to the bone marrow?

References

- 1 Hirabayashi, Y., Li, G., Yoon, B.I., Fujii-Kuriyama, Y., Kaneko, T., Kanno, J., and Inoue, T. (2003) *Organohalogen Compounds*, 64, 270
- 2 Garrett, R.W., and Gasiewicz, T.A. (2005) Meeting abstract [Molecular Regulation of Stem Cell, Keystone symosia], 61
- 3 Yoon, B.I., Hirabayashi, Y., Kawasaki, Y., Kodama, Y., Kaneko, T., Kanno, J., Kim, D.Y., Fujii-Kuriyama, Y., and Inoue, T. (2002) *Toxicol Sci*, 70, 150
- 4 Yoon, B.I., Li, G.X., Kitada, K., Kawasaki, Y., Igarashi, K., Kodama, Y., Inoue, T., Kobayashi, K., Kanno, J., Kim, D.Y., Inoue, T., and Hirabayashi, Y. (2003) *Environ Health Perspect*, 111, 1411
- 5 Mimura, J., Yamashita, K., Nakamura, K., Morita, M., Takagi, T.N., Nakao, K., Ema, M., Sogawa, K., Yasuda, M., Katsuki, M., and Fujii-Kuriyama, Y. (1997) *Genes Cells*, 2, 645
- 6 Yoon, B.I., Hirabayashi, Y., Kawasaki, Y., Kodama, Y., Kaneko, T., Kim, D.Y., and Inoue, T. (2001) *Exp Hematol*, 29, 278
- 7 Till, J.E., and McCulloch, E.A. (1961) *Radiat Res*, 14, 213
- 8 Hirabayashi, Y., Matsuda, M., Aizawa, S., Kodama, Y., Kanno, J., and Inoue, T. (2002) *Exp Biol Med (Maywood)*, 227, 474
- 9 Hirabayashi, Y., Inoue, T., Suda, Y., Aizawa, S., Ikawa, Y., and Kanisawa, M. (1992) *Exp Hematol*, 20, 167
- 10 Ivanova, N.B., Dimos, J.T., Schaniel, C., Hackney, J.A., Moore, K.A., and Lemischka, I.R. (2002) *Science*, 298, 601
- 11 Hirabayashi, Y., Inoue, T. (2005) in: *Handbook of Toxicogenomics* (Borlak J., Ed.), Wiley-VCH Verlag GmbH & Co. Weinheim, 583.



25TH INTERNATIONAL SYMPOSIUM
ON HALOGENATED ENVIRONMENTAL
ORGANIC POLLUTANTS AND POPs

DIOXIN 2005

ISPAC 20

20TH INTERNATIONAL SYMPOSIUM ON
POLYCYCLIC AROMATIC COMPOUNDS

ORGANOHALOGEN COMPOUNDS VOLUME 67

**CD-ROM OF PROCEEDINGS OF
DIOXIN 2005 AND ISPAC-20**

Edited by E. Reiner and M. Alaei

CD production sponsored by



CHEM

ISBN 0-9738972-0-1