







この配置は回転による平面固定が不十分の可能性はあるが、benzimidazole 環と benzene が共役しており、比較的平面性を維持していると考えられる。

Pimozide の酸化反応



Pimozide は主に CYP3A4 によって N-脱アルキル化される。この反応に CYP1A2 も弱いながら関与する。Fluorobenzene と Benzimidazole 誘導体が trigger 候補となる。Piperazine 環は平面から逸脱するので、許容範囲内に収める必要がある。Fluorobenzene が奥に回って配置される構造が想定される。いずれにせよ反応の進行は遅いと予想する。





D. E. 考察と結論

薄い幅の CYP1A2 酸化部位にはまり込んだ下段立体配置で脱アルキル化が起きると予想される。

pyrene 則に基づいてヒト CYP1A2 の代謝可否と酸化部位の特定が可能となった。環境物質、天然物および医薬品について解析したデータから pyrene 周辺の環境を具体的な大きさ、空間としてとらえることができた。今後、ヒトによるマニュアル操作からコンピュータによる自動解析を行う手法の開発に着手する予定である。

G. 研究発表

1. 論文発表

R. Ueda, H. Iketaki, K. Nagata, S. Kimura, F. J. Gonzalez, K. Kusano, T. Yoshimura and Y. Yamazoe. A common regulatory region

functions bi-directionally in transcriptional activation of the human *CYP1A1* and *CYP1A2* genes. *Mol. Pharmacol.*, in press

2. 学会発表

山添康, Chemical Approach to Predict Metabolism of CYP2Cs, ISSX-JXXS meeting, Maui, Oct 23 2005

H. 知的財産権の出願・登録状況

なし

研究成果の刊行に関する一覧表

書籍

著者氏名	論文タイトル名	書籍全体の 編集者名	書 籍 名	出版社名	出版地	出版年	ページ
Ema M, Hirose A.	Reproductive and developmental toxicity of organotin compounds	Golub MS, Ed.	Metals, Fertility, and productive Toxicity	CRC Press	Boca Raton	2006	23-64

雑誌

発表者氏名	論文タイトル名	発表誌名	巻号	ページ	出版年
Hayashi M, Kamata E, Hirose A, Takahashi M, Morite T, Ema M	In silico assessment of chemical mutagenesis in comparison with results of Salmonella microsome assay on 909 chemicals	<i>Muta Res.</i>	588	129-135	2005
Suzuki H, Ikeda N, Kobayashi K, Terashima Y, Shimada Y, Suzuki T, Hagiwara T, Hatakeyama S, Nagaoka K, Yoshida J, Saito Y, Tanaka J, Hayashi M	Evaluation of liver and peripheral blood micronucleus assays with 9 chemicals using young rats: A study by the Collaborative Study Group for the Micronucleus Test (CSGMT)/Japanese Environmental Mutagen Society (JEMS)—Mammalian Mutagenicity Study Group (MMS)	<i>Mutat Res.</i>	583	133-145	2005
Hayashi M	げっ歯類を用いる小核試験の基礎研究ならびにその行政面への応用	<i>Environ Mutagen Res</i>	27	13-20	2005
Sofuni T, Nohmi T, Ohta T, Hayashi M	Genotoxicity: Is a threshold concept applicable to evaluate the mutagenic activity of DNA-targeting substances !?	<i>Environ Mutagen Res</i>	27	61-73	2005
Hamamura M, Hirose A, Kamata E, Katoku K, Kuwasaki E, Oshikata T, Nakahara Y, Ema M and Hasegawa R	Semi-quantitative immunohistochemical analysis of male rat-specific α 2u-globulin accumulation for chemical toxicity evaluation	<i>J Toxicol Sci</i>	31	35-47	2006
Hasegawa R, Hirata-Koizumi M, Takahashi M, Kamata E and Ema M	Comparative susceptibility of newborn and young rats to six industrial chemicals	<i>Congenit Anom</i>	45	137-145	2005
Kawabe K, Saegusa H, Seto K, Urabe H and Motojima K	Peroxisome proliferator-activated receptor α and its response element are required but not sufficient for transcriptional activation of the mouse heart-type fatty acid binding protein gene	<i>Int J Biochem Cell Biol.</i>	37	1534-1546	2005

Nakamura T ^a , Shibata N ^a , Nishimoto-Shibata T, Feng D, Ikemoto M, Motojima K, Iso-O N, Tsukamoto K, Tsujimoto M and Arai H	Regulation of SR-BI protein levels by phosphorylation of its associated protein, PDZK1	<i>Proc Natl Acad Sci U S A</i>	20	13404- 13409	2005
Hirono S, Nakagome I, Imai R, Maeda K, Kusahara H, and Sugiyama Y	Estimation of the Three-Dimensional Pharmacophore of Ligands for Rat Multidrug- Resistance-Associated Protein 2 using Ligand-Based Drug Design Techniques	<i>Pharm. Res</i>	22	260-269	2005
Katsuki M, Chuang V T G, Nishii K, Kawahara K, Nakayama H, Yamaotsu N, Hirono S and Otagiri M.	Use of Photoaffinity Labeling and Site Directed Mutagenesis for Identification of Key Residue Responsible for Extraordinarily High Affinity Binding of UCN-01 in Human Alpha 1-Acid Glycoprotein	<i>J. Biol. Chem.</i>	280	1384-1391	2005
Oda A, Yamaotsu N, and Hirono S	New AMBER Force-Field Parameters of Cytochrome P450s Determined by Quantum Chemical Calculations of Simplified Models	<i>J. Compt. Chem.</i>	26	818-826	2005
Yumiko Osa, Yuka Hikima, Yoko Sato, Koichi Takino, Yoshihiro Ida, Hirono S, and Hiroschi Nagase	Covenient Synthesis of Oxazolidinones by the Use of Halomethyloxirane, Primary Amine, and Carbonate Salt	<i>J. Org. Chem.</i>	70	5737-5740	2005
S.-F. Ma, M. Anraku, Y. Iwao, K. Yamasaki, U. Kragh-Hansen, N. Yamaotsu, S. Hirono, T. Ikeda, M. Otagiri	Hydrolysis of Angiotensin II Receptor Blocker Prodrug Olmesartan Medoxomil by Human Serum Albumin and Identification of Its Catalytic Active Site	<i>Drug Metab. Dispos</i>	33	1911-1919	2005
Gouda H, Sunazuka T, Ui H, Handa M, Sakoh Y, Iwai Y, Hirono S, and Omura S	Stereostructure of luminamicin, an anaerobic antibiotic, via molecular dynamics, NMR spectroscopy, and the modified Mosher method	<i>Proc.Natl.Acad.Sci.US A,</i>	102	18286-182 91	2005

Oda A, Tsuchida K, Takakura T, Yamaotsu N, Hirono S	Comparison of Consensus Scoring Strategies for Evaluating Computational Models of Protein-Ligand Complexes	<i>J.Chem. Inf. Model.</i>	46	380 -391	2006
Tsuchida K, Chaki H, TakakuraT, Kotsubo H, Tanaka T, Aikawa Y, Shiozawa S, Hirono S	Discovery of Nonpeptidic Small-Molecule AP-1 Inhibitors: Lead Hopping Based on Three-Dimensional Pharmacophore Model	<i>J.Med.Chem</i>	49	80-91	2006

研究成果の刊行物・別冊

Metals, Fertility, and Reproductive Toxicity

Edited by
Mari S. Golub



Taylor & Francis

Taylor & Francis Group

Boca Raton London New York

A CRC title, part of the Taylor & Francis imprint, a member of the
Taylor & Francis Group, the academic division of T&F Informa plc.

Published in 2006 by
CRC Press
Taylor & Francis Group
6000 Broken Sound Parkway NW, Suite 300
Boca Raton, FL 33487-2742

© 2006 by Taylor & Francis Group, LLC
CRC Press is an imprint of Taylor & Francis Group

No claim to original U.S. Government works
Printed in the United States of America on acid-free paper
10 9 8 7 6 5 4 3 2 1

International Standard Book Number-10: 0-415-70040-X (Hardcover)
International Standard Book Number-13: 978-0-415-70040-5 (Hardcover)
Library of Congress Card Number 2005049128

This book contains information obtained from authentic and highly regarded sources. Reprinted material is quoted with permission, and sources are indicated. A wide variety of references are listed. Reasonable efforts have been made to publish reliable data and information, but the author and the publisher cannot assume responsibility for the validity of all materials or for the consequences of their use.

No part of this book may be reprinted, reproduced, transmitted, or utilized in any form by any electronic, mechanical, or other means, now known or hereafter invented, including photocopying, microfilming, and recording, or in any information storage or retrieval system, without written permission from the publishers.

For permission to photocopy or use material electronically from this work, please access www.copyright.com (<http://www.copyright.com/>) or contact the Copyright Clearance Center, Inc. (CCC) 222 Rosewood Drive, Danvers, MA 01923, 978-750-8400. CCC is a not-for-profit organization that provides licenses and registration for a variety of users. For organizations that have been granted a photocopy license by the CCC, a separate system of payment has been arranged.

Trademark Notice: Product or corporate names may be trademarks or registered trademarks, and are used only for identification and explanation without intent to infringe.

Library of Congress Cataloging-in-Publication Data

Metals, fertility, and reproductive toxicity / edited by Mari Golub.
p. cm.

Includes bibliographical references.

ISBN 0-415-70040-X (alk. paper)

1. Metals--Toxicology. 2. Reproductive toxicology. I. Golub, Mari S.

RA1231.M52M53 2005
615.9'253--dc22

2005049128

informa
Taylor & Francis Group
is the Academic Division of Informa plc.

Visit the Taylor & Francis Web site at
<http://www.taylorandfrancis.com>

and the CRC Press Web site at
<http://www.crcpress.com>

chapter 3

Reproductive and Developmental Toxicity of Organotin Compounds

Makoto Ema and Akihiko Hirose

*Division of Risk Assessment, Biological Safety Research Center,
National Institute of Health Sciences, Tokyo, Japan*

Contents

Introduction	24
Effects on Aquatic Organisms	25
Imposex on Gastropods	25
Effects on Fish.....	27
Effects on Other Organisms.....	28
Summary of Effects on Aquatic Organisms	28
Effects on Experimental Animals.....	28
Reproductive Toxicity of Phenyltin Compounds.....	28
Reproductive Toxicity of Triphenyltins.....	28
Reproductive Toxicity of Diphenyltin Compounds.....	31
Summary of Reproductive Toxicity of Phenyltin Compounds.....	33
Developmental Toxicity of Phenyltin Compounds.....	33
Summary of Developmental Toxicity of Phenyltin Compounds	35
Reproductive Toxicity of Butyltin Compounds.....	35
Summary of Reproductive Toxicity of Butyltin Compounds.....	39
Developmental Toxicity of Butyltin Compounds.....	40
<i>In Vivo</i> Developmental Toxic Effects of Butyltin Compounds	40
<i>In Vitro</i> Dymorphogenic Effects of Butyltin Compounds	49
Summary of Developmental Toxicity of Butyltin Compounds	51
Developmental Toxicity of Miscellaneous Organotin Compounds	51
Summary of Developmental Toxicity of Miscellaneous Organotin Compounds	54

Conclusions.....	54
References.....	55

Introduction

Organotin compounds are chemicals widely used in agriculture and industry (Piver 1973, World Health Organization 1980). Tetrasubstituted organotin compounds are mainly used as intermediates in the preparation of other organotin compounds. Trisubstituted organotin compounds have biocidal properties and are used in agriculture as fungicides and acaricides, as rodent repellents, and molluscicides, and are widely used as antifoulants in ship paints and underwater coatings. Especially, triphenyltins (TPTs) and tributyltins (TBTs) have been used extensively in antifouling products such as algaecides and molluscicides. Disubstituted organotin compounds are commercially the most important derivatives, and are mainly used in the plastics industry, particularly as heat and light stabilizers for polyvinyl chloride (PVC) plastics to prevent degradation of the polymer during melting and forming of the resin into its final products, as catalysts in the production of polyurethane foams, and as vulcanizing agents for silicone rubbers. Mono-substituted organotin compounds are used as stabilizers in PVC films. Widespread use of organotin compounds has caused increasing amounts to be released into the environment. The most important nonpesticidal route of entry for organotin compounds into the environment is through leaching of organotin-stabilized PVC in water (Quevauviller et al. 1991), and the use in antifouling agents, resulting in the introduction of organotin into the aquatic environment (Maguire 1991). Data are available regarding the detection of butyltin and phenyltin compounds in aquatic marine organisms (Sasaki et al. 1988, Fent and Hunn 1991, Lau 1991) and marine products (Suzuki et al. 1992, Belfroid et al. 2000, Tsuda et al. 1995, Ueno et al. 1999, Toyoda et al. 2000). Food chain bioamplification of butyltin in oysters (Waldock and Thain 1983), mud crabs (Evans and Laughlin 1984), marine mussels (Laughlin et al. 1986), Chinook salmon (Short and Thrower 1986), and dolphin, tuna, and shark (Kannan et al. 1996), and of phenyltin in carp (Tsuda et al. 1987) and horseshoe crab (Kannan et al. 1995) has been reported. These indicate that organotin compounds accumulate in the food chain and are bioconcentrated, and that humans can be exposed to organotin compounds via seafood. The World Health Organization (WHO) reported in 1980 that the estimated mean total daily intake of tin by humans ranged from 200 μg to 17 mg. Recently, Tsuda et al. (1995) reported that the daily intakes in Shiga prefecture in Japan were 0.7 to 5.4 μg in 1991 and 0.7 to 1.3 μg in 1992 for TPT and 4.7 to 6.9 μg in 1991 and 2.2 to 6.7 μg in 1992 for TBT. Toyoda et al. (2000) also showed that the daily intakes in Japanese consumers, based on analysis with the 1998 total diet samples, were 0.09 μg for TPT, 0 μg for diphenyltin (DPT), 1.7 μg for TBT, and 0.45 μg for dibutyltin (DBT). These values are lower than the acceptable daily intake for TPT according to the JMPR (Joint Meetings of the FAO [Food and Agriculture Organization] and World Health Organization

Panel of Experts on Pesticides Residues), 25 µg (World Health Organization 1992), and the guidance value for oral exposure to tributyltin oxide (TBTO), 18 µg (International Programme on Chemical Safety 1999a). Thus, the levels of organotin compounds in seafood are not considered to be sufficiently high to affect human health (Tsuda et al. 1995, Ueno et al. 1999). However, Belfroid et al. (2000) noted that more research on residual TBT levels in seafood is needed before a definitive conclusion on possible health risks can be drawn.

In recent years, adverse effects of environmental chemicals on the reproductive success of wildlife populations have been reported (Colborn et al. 1993). These phenomena may result from interference with the endocrine system. Disturbances of hormonal regulation during pre- and postnatal development may produce deleterious effects on reproduction and development. TPT and TBT are suspected to be endocrine disruptors (Japan Environment Agency 1998). TBT and TPT are known to have strong effects on the development of imposex (imposition of male sex characteristics on females) in the rock shell (Horiguchi et al. 1996, 1997a), and this condition may bring about reproductive failure and a consequent population decline.

Although the toxicity of organotins has been extensively reviewed (World Health Organization 1980, Snoeij et al. 1987, Winship 1988, Boyer 1989, International Programme on Chemical Safety 1999a, b), the reproductive and developmental toxicity of these compounds is not well understood. In this chapter, we summarize the findings of the studies on reproductive and developmental effects of organotin compounds.

Effects on Aquatic Organisms

Imposex on Gastropods

TBT causes reproductive toxic effects in marine gastropods, which were represented by some masculinizing effects including *imposex* or *pseudohermaphroditism*. The imposition of male sex organs (a penis and vas deferens) on female mud snails (*Nassarius ovooietus*) was found in near harbors, and the degree of penis development and frequency of imposex were positively correlated to the seawater TBT concentration (Smith 1981a, b). Imposex has been induced experimentally by treatment with 4.5 to 5.5 µg/L of TBT compounds for 60 days. In field studies in southeastern England, imposex has been reported in declining populations of the common dogwhelk (Bryan et al. 1986, 1987, 1989, Gibbs and Bryan 1986, Davies et al. 1987, Gibbs et al. 1987).

Imposex has not just occurred at a regional level, but worldwide on a global scale. Imposex in dogwhelk was not only reported in England, but in Scotland, the Netherlands, and the coastline of the North Sea. Imposex in other whelk species occurred in Canada, West Africa, New Zealand, Australia, Malaysia, Singapore, Indonesia, and Japan (Fent 1996, Horiguchi et al. 1996). Imposex among prosobranchs is known to occur in around 70 species of 50 genera, although some species are less susceptible to TBT compounds (Fioroni et al. 1991, Fent 1996).

TPT also induced imposex in *Thais clavigera* at the same potency as TBT (Horiguchi et al. 1997a). Although, in *Nucvolla lapillus*, TPT did not induce imposex, tripropyltin (TPrT) had a small effect on the development of imposex (Bryan et al. 1988). DBT and monobutyltin (MBT) did not induce imposex in the gastropod species examined. Three trisubstitution compounds (TBT, TPT, TPrT) and monophenyltin (MPT) easily induced imposex in some species, among the eight organotins, i.e., MBT, DBT, TBT, tetrabutyltin (TeBT), MPT, DPT, TPT, and TPrT. (Bryan et al. 1988, Hawkins and Hutchinson 1990, Horiguchi et al. 1997b).

The early studies in the 1980s reached some common conclusions, which are described below (Eisler 2000). Imposex correlated with the body burden of tributyl- and dibutyltin, but not with the tissue concentration of arsenic, cadmium, copper, lead, silver, or zinc. Forty-one percent of females had male characteristics, when the body burden reached to 1.65 mg Sn/kg of dry soft parts, by exposing with 0.02 µg Sn/L for 120 days. Imposex in immature females is caused above the concentration of around 1 ng/L (Sn) in seawater. At higher concentrations of TBT, the oviduct had been blocked, resulting in sterilization. Declining dogwhelk populations could be caused by aborting capsules, sterility, and premature death, which were characterized by a moderate to high degree of imposex, fewer female functions, fewer juveniles, and scarcity of laid egg capsules.

There is also a great variety of gradations of imposex in different species. The intensity is characterized by a classification system, which distinguishes six stages with a few different types, mainly based on a Vas Deferens Sequence (VDS) index (Oehlmann et al. 1991). Imposex development occurred in three variations: (1) a small penis without penis duct, (2) a short distal vas deferens section, or (3) a short proximal vas deferens section (stage 1). At stages 2 and 3 the male sex characteristics of each type are developed continuously. Stage 4 is characterized by a penis with penis duct and a complete vas deferens, and represents the last stage of fertility. The reproductive failure or sterility is induced in later stages. At stage 5 the vagina is replaced with a small prostate gland, the vagina opening is blocked by vas deferens tissue, or the incompleteness of the pallial oviduct closure occurs. Abortive egg capsules fill the lumen and vestibulum of the capsule gland and evoke an intense swelling of the gland at stage 6 (Bettin et al. 1996). High TBT exposure in the early stages of life induced gametogenesis or sex changes characterized by a suppression of oogenesis and commencement of spermatogenesis in females (Gibbs et al. 1988, Fioroni et al. 1991, Oehlmann et al. 1991, 1996, Horiguchi et al. 2002). It was thought that the initial phases of imposex corresponding to VDS stages 1 and 2 may be reversible; however, advanced phases of imposex and sterilization with gross morphological changes corresponding to VDS stage 5 and 6 would be irreversible (Fent 1996).

Although many morphological aspects of pseudohermaphroditism caused by TBT have been investigated, the biochemical mechanism has been indistinct. It is known that a neurotropic hormone called the penis morphogenic factor (PMF) develops male normal differentiation in mollusks (Féral

and LeGall, 1983). Co-localization of TBT with PMF in ganglia suggested that PMF release through TBT's neurotropic action induced masculinization in females (Bryan et al. 1989). Other studies indicated increased testosterone levels detected in female dogwhelk exposed to TBT, and that testosterone injection without TBT induced penis development in females (Spooner et al. 1991, Stroben et al. 1991). The later studies suggested that TBT disturbed the P-450-dependent aromatization of androgens to estrogen, and a nonsteroidal specific aromatase inhibitor-induced imposex similar to TBT (Bettin et al. 1996). However, the PMF has not been well characterized, and the role of vertebrate sex steroids is not known in gastropods to date. A recent study proposed that the combination of changes in the neuropeptide (APGWamide), which is considered to be a PMF in mud snails, and steroid hormones would lead to imposex induction at extremely low doses of TBT (Oberdörster and McClellan-Green 2002).

Effects on Fish

TBT or TPT exposure in early life stages induces altered embryonic development, and delayed or inhibited hatching in fish. Exposure of TBT or TBT to minnow eggs and larvae at concentrations of 0.2 to 18 $\mu\text{g}/\text{L}$ in the water in which the fish lived induced dose-dependent morphological effects on larvae. Marked body axis deformations were observed at more than about 4 $\mu\text{g}/\text{L}$ exposure, and incomplete hatching occurred at similar concentrations in 10 to 30% of larvae. At 15.9 $\mu\text{g}/\text{L}$ of TPT exposure, hatching was delayed and the hatching rate was reduced significantly (Fent and Meier 1992, 1994). Developmental defects, such as skeletal abnormality and retarded yolk sac resorption, occurred in zebrafish larvae at more than 25 $\mu\text{g}/\text{L}$ of triphenyltin acetate (TPTA) exposure, and hatching delay was found at more than 0.5 $\mu\text{g}/\text{L}$ (Strmac and Braunbeck 1999). These developmental effects in fish were caused not only by organotin compounds, but also by a variety of contaminants (i.e., heavy metals, chlorinated hydrocarbons, altered pH), suggesting that such alteration would be classified as a nonspecific reaction to organic toxicants (Fent 1996, Strmac and Braunbeck 1999).

Some reproductive effects (i.e., reduced fecundity and sperm counts) in fish were reported. Reproductive success of three-spine stickleback with TBT exposure were examined over a 7-month period; no effects were detected in relation to fecundity, number of hatched fry, or frequency of malformed fry. However, no changes were found in the gonad somatic index (GSI; ovary weight ratio to total body weight); by the 7-month TBT treatment (2 $\mu\text{g}/\text{L}$) despite increasing GSI in controls, which suggested a lack of maturation of egg tissue and consequently a potential reduced fecundity (Holm et al. 1991). In sheepshead minnows, reduction in both total and percent viable eggs was found at more than 1.3 $\mu\text{g}/\text{L}$ of TBT exposure, although the reductions were not statistically significant (Manning et al. 1999). TBT exposure to Japanese medaka at 1 mg/kg body weight caused a reduction of the spawning frequency (Nirmala et al. 1999). Additionally, environmentally relevant concentrations of

TBT induced significantly decreased sperm counts in guppies (11.2 to 22.3 ng/L for 21 days), and decreased sperm motility at concentrations less than 1 µg/L (Haubruge et al. 2000, Kime et al. 2001).

Effects on Other Organisms

Despite a great number of studies on imposex in snails and a comparable number of toxicity reports on fish, there is little information on development and reproductive effects on other species by organotin compounds. It was reported that imposex has not only been found in gastropods, but also been induced in Japanese freshwater crabs by TBT (Takahashi et al. 2000). In crabs, imposex has also occurred in males, which is characterized by dual-gender imposex (either a female genital opening or a single ovary occurred in males). Malformations during limb regeneration occurred in fiddler crabs (Weis and Kim 1988) and in axolotl, induced by TBT (Scadding 1990).

Summary of Effects on Aquatic Organisms

TBT or TPT causes the imposition of male sex organs (imposex) on female mud snails above the concentration of about 1 ng/L (Sn) in seawater, but DBT or MPT does not induce imposex. The intensity is characterized by a classification system based on the VDS, and advanced phases of imposex and sterilization with gross morphological changes are irreversible. The biochemical mechanism studies suggested that the induction of either neurotropic hormone or androgen titer would lead to imposex at an extremely low dose of TBT. Also, TBT or TPT exposure in the early life stages of fish causes altered embryonic development, impaired morphological development, and delayed or inhibited hatching, and reduces fecundity and sperm counts. Such reproductive and developmental defects were also found in other species. The impaired reproduction and subsequent population decline in a variety of aquatic organisms by organotins are an important issue in aquatic ecosystems.

Effects on Experimental Animals

Reproductive Toxicity of Phenyltin Compounds

Reproductive Toxicity of Triphenyltins

TPTs have been reported to be insect chemosterilants (Kenaga 1965). Reproductive studies on TPTs are presented in Table 3.1. Several reports on male reproductive toxicity have been published. Male Sharman rats were given a diet containing triphenyltin hydroxide (TPTH) at 50, 100, or 200 ppm and then mated with untreated females repeatedly five times (Gaines and Kimbrough 1968). Reduced fertility, such as decreases in the total number of matings, total number of litters born alive, and ratio of number of litters to number of matings, accompanied by a marked reduction in food consumption

Table 3.1 Reproductive Toxicity of Phenylin Compounds

Compounds	Animals	Dose	Days of Administration	Route	Reproductive and Developmental Effects	Author(s)
TPTH	Sharman rat	100-200 ppm	64-238 days	Diet	Decreased no. of matings Decreased no. of litters born alive Decreased ratio of no. of litters to no. of matings	Gains and Kimbrough (1968)
TPIA or TPICI	Holtzman rat	20 mg/kg	19 days	Diet	Decreased testicular size Change in testicular morphology	Pate and Hays (1968)
TPIA or TPICI	Holtzman rat	20 mg/kg	20 days	Diet	Impairment of spermatogenic process	Snow and Hays (1983)
TPIA	ICR/Ha	2.4-12 mg/kg	1 day	ip	No dominant lethal effect	Epstein et al. (1972)
TPTH	Swiss mouse	6 mg/kg	5 days	Gavage	No dominant lethal effect	Newton and Hays (1968)
		1.3-8.5 mg/kg	1 day	ip	No dominant lethal effect	
TPIA or TPICI	Holtzman rat	11 mg/kg	5 days	Gavage	No dominant lethal effect	Newton and Hays (1968)
		20 mg/kg	4-24 days	Diet	Decreased no. of mature follicles Increased incidence of atresia in early follicle growth	
TPICI	Wistar rat	4.7-6.3 mg/kg	Days 0-3 of pregnancy	Gavage	Decreased no. of corpora lutea	Ema et al. (1997a)
		12.5-25 mg/kg	Days 4-6 of pregnancy	Gavage	Decreased pregnancy rate, decreased fetal wt.	
TPICI	Wistar rat	4.7-6.3 mg/kg	Days 0-3 of pseudopregnancy	Gavage	Suppression of uterine decidualization	Ema et al. (1999a)
DPTCI	Wistar rat	16.5-24.8 mg/kg	Days 0-3 of pregnancy	Gavage	Decreased pregnancy rate, preimplantation loss, decreased fetal wt.	Ema et al. (1999b)
DPTCI	Wistar rat	33.3 mg/kg	Days 4-7 of pregnancy	Gavage	Effects as above, postimplantation loss	Ema and Miyawaki (2002)
		4.1-24.8 mg/kg	Days 0-3 of pregnancy	Gavage	Suppression of uterine decidualization	

and weight gain, were observed at 100 or 200 ppm for 64 days. At these doses, food consumption later improved, and with it, fertility. Dietary exposure to triphenyltin acetate (TPTA) or triphenyltin chloride (TPTCl) at 20 mg/kg for 19 days produced marked effects on body weight, testicle size, and testicular structure in male Holtzman rats (Pate and Hays 1968). Microscopic examinations revealed degenerative changes, such as a decrease in the number of layers per tubule, a depletion of the more advanced cell forms from the tubules, and a closing of the tubule lumina. Effects were more pronounced in rats treated with TPTA. TPTA or TPTCl at 20 mg/kg in feed for 20 days was reported to cause an impairment of the spermatogenic process in male Holtzman rats; complete recovery of the spermatogenesis was observed after feeding a normal diet for 70 days (Snow and Hays 1983). No mutagenicity was detected in dominant lethal assay in which male ICR/Ha Swiss mice were given a single intraperitoneal injection of TPTA at 2.4 or 12 mg/kg or TPTH at 1.3 or 8.5 mg/kg, or given TPTA at 6mg/kg or TPTH at 11 mg/kg by gavage on 5 successive days and then mated with untreated females, and pregnancy outcome was determined on day 13 of pregnancy (Epstein et al. 1972).

Adverse effects on female reproductive toxicity were also reported. Dietary TPTA and TPTCl at 20 mg/kg for 4 days produced significant changes in the ovarian tissue, including a decreased number of mature follicles, an increased incidence of atresia in early follicle growth, and a pronounced decrease in the number of corpora lutea in female Holtzman rats (Newton and Hays 1968). These effects were regarded as a decrease in ovulation, and thus decreased fertility. The adverse effects of TPTCl on the initiation and maintenance of pregnancy were determined after administration to the mother during early pregnancy (Ema et al. 1997a). Following successful mating, female Wistar rats were given TPTCl by gavage on days 0 to 3 of pregnancy at 3.1, 4.7, or 6.3 mg/kg or on days 4 to 6 of pregnancy at 6.3, 12.5, or 25.0 mg/kg, and pregnancy outcome was determined on day 20 of pregnancy. TPTCl totally prevented implantation in a dose-dependent manner. The pregnancy rate was decreased after administration of TPTCl on days 0 to 3 at 4.7 and 6.3 mg/kg and on days 4 to 6 at 12.5 and 25.0 mg/kg. Preimplantation loss was increased after administration of TPTCl on days 0 to 3 at 4.7 mg/kg and higher. In females having implantations, the numbers of implantations and live fetuses, and the incidences of pre- and postimplantation embryonic loss in the TPTCl-treated groups were comparable to the controls. These results indicate that TPTCl during early pregnancy causes failure in implantation and has greater antiimplantation effects when administered during the preimplantation period than the periimplantation period.

The function of the uterine endometrium is one of the principle factors in embryonic survival. Uterine decidualization is required for normal implantation, placentation, and therefore normal gestation in rats. The uterine growth induced by endometrial trauma in pseudopregnant animals mimics the decidual response of the pregnant uterus that occurs after embryo implantation (Cummings 1990, Kamrin et al. 1994). The decidual cell

response (DCR) is a model for maternal physiological events that are associated with implantation (Cummings 1990). This technique can distinguish between the adverse effects of chemical compounds in the maternal and fetal compartments, and has been used to evaluate the reproductive toxicity of chemical compounds (Spencer and Sing 1982, Bui et al. 1986, Cummings 1990, Kamrin et al. 1994, Ema et al. 1998). The effects of TPTCl on the reproductive capability of the uterus, as a cause of implantation failure, were evaluated using pseudopregnant rats (Ema et al. 1999a). Female Wistar rats were given TPTCl by gastric intubation at 3.1, 4.7, or 6.3 mg/kg on days 0 to 3 of pseudopregnancy. Between 11:00 and 13:00 on day 4 of pseudopregnancy, induction of DCR was performed via midventral laparotomy under ether anesthesia, and experimental decidualization was initiated by scratching the antimesometrial surface of the endometrium with a bent needle. The uterine weight on day 9 of pseudopregnancy served as an index of the uterine decidualization (De Feo 1963). A decrease in the uterine weight, which indicates suppression of the uterine decidualization, was detected at 4.7 and 6.3 mg/kg. TPTCl at 4.7 and 6.3 mg/kg also produced a decrease in the serum progesterone levels in female rats on day 4 and on day 9 of pseudopregnancy. These doses caused an increase in implantation failure (preimplantation embryonic loss) in female rats given TPTCl on days 0 to 3 of pregnancy (Ema et al. 1997a). These results suggest that TPTCl causes the suppression of uterine decidualization correlated with the reduction in serum progesterone levels, and these participate in the induction of implantation failure due to TPTCl. Protective effects of progesterone against suppression of uterine decidualization and implantation failure induced by TPTCl were examined (Ema and Miyawaki 2001). The hormonal regimen, consisting of progesterone and estorone supported decidual development in ovariectomized rats given TPTCl. The pregnancy rate and number of implantations in groups given TPTCl at 4.7 or 6.3 mg/kg in combination with progesterone were higher than those in the groups given TPTCl alone. These results indicate that the TPTCl-induced suppression of uterine decidualization is mediated, at least partially, by ovarian hormones, and that progesterone protects against TPTCl-induced implantation failure.

Reproductive Toxicity of Diphenyltin Compounds

Oral TPT is metabolized to DPT, MPT, and further to inorganic tin in rats (Kimmel et al. 1977, Ohhira and Matsui 1993 a, b). Reproductive toxicity studies on DPTs are also published (Table 3.1). The adverse effects of diphenyltin dichloride (DPTCl) on the initiation and maintenance of pregnancy, and the role of DPT in the implantation failure of TPT were evaluated. Following successful mating, DPTCl was given to Wistar rats by gavage on days 0 to 3 of pregnancy at 4.1, 8.3, 16.5, or 24.8 mg/kg or on days 4 to 7 of pregnancy at 8.3, 16.5, 24.8, or 33.0 mg/kg (Ema et al. 1999b). The pregnancy rate was decreased after administration of DPTCl on days 0 to 3 at 24.8 mg/kg and on days 4 to 7 at 33.0 mg/kg. The incidence of preimplantation loss was increased at 16.5 mg (equivalent to 48 μ mol)/kg on days 0 to 3. In

females having implantations, the incidences of pre- and postimplantation embryonic loss in the groups given DPTCl on days 0-3 were comparable to the controls. The incidence of postimplantation embryonic loss was increased after administration of DPTCl on days 4 to 7 at 33.0 mg/kg. These results indicate that DPTCl during early pregnancy causes implantation failure, and that DPTCl has greater effects on reproduction when administered during the preimplantation period rather than the periimplantation period. Following administration on days 0 to 3 of pregnancy, the increased incidence of preimplantation embryonic loss was induced by TPTCl, a parent compound of DPTCl, at 4.7 mg (equivalent to 12 μ mol)/kg and higher (Ema et al. 1997a), or DPTCl at 16.5 mg (equivalent to 48 μ mol)/kg. If, on a mole-equivalent basis, a metabolite is as, or more, effective than the parent compound, this is consistent with the view that the metabolite is the proximate toxicant or at least an intermediate to the proximate toxicant. Thus, it seems unlikely that only DPTCl and/or its further metabolites can be considered the agents responsible for the antiimplantation effects of TPTCl. As for the metabolism of phenyltin, however, Ohhira and Matsui (1993b) showed that TPT compound was formed in the liver of the DPTCl-treated rat by metabolism of DPTCl, and suggested that part of the administered DPT compound has some harmful effect as the TPT compound in rats, and this must be taken into consideration in toxicological research on DPT. Further studies are needed to clarify the difference in the reproductive toxicity induced by TPT and DPT, and to identify the proximate or ultimate toxicant of phenyltins. The effects of DPTCl on the reproductive capability of the uterus were evaluated in pseudopregnant rats according to the procedure described above. Female Wistar rats were given DPTCl by gastric intubation on days 0 to 3 of pseudopregnancy at 4.1, 8.3, 16.5, or 24.8 mg/kg (Ema and Miyawaki 2002). Suppression of uterine decidualization was observed at 16.5 mg/kg and higher. A decrease in the serum progesterone levels in pseudopregnant rats was also found on day 4 and on day 9 of pseudopregnancy at 16.5 mg/kg and higher. These doses induced an increase in preimplantation embryonic loss in female rats given DPTCl on days 0 to 3 of pregnancy (Ema et al. 1999b). No changes in serum estradiol levels in pseudopregnant rats were noted. These results suggest that DPTCl causes the suppression of uterine decidualization correlated with the reduction in serum progesterone levels. These are responsible for the DPTCl-induced implantation failures. The hormonal regimen consisting of progesterone and estrone supported decidual development in ovariectomized rats given DPTCl (Ema and Miyawaki 2002). The pregnancy rate and number of implantations in groups given DPTCl at 16.5 or 24.3 mg/kg in combination with progesterone were higher than those in the groups given DPTCl alone. These results show that the DPTCl-induced suppression of uterine decidualization is mediated, at least partially, by ovarian hormones, and that progesterone protects against the DPTCl-induced implantation failure.

Summary of Reproductive Toxicity of Phenyltin Compounds

TPTs caused a decrease in male fertility due to degenerative changes in testicular tissue, which were associated with a marked decrease in food consumption. Complete recovery of fertility and impairment of the spermatogenesis was noted following withdrawal of treatment. Female reproductive failure induced by TPTs is more prominent. The harmful effects of TPTs on the ovaries were present after 5 days of treatment, before any significant effects on body weight gain. TPTCl during early pregnancy caused implantation failure at relatively low doses, and TPTCl had greater antiimplantation effects when administered during the preimplantation period. The implantation failure due to TPTCl might be mediated by suppression of uterine decidualization and correlated with the reduction in serum progesterone levels. Implantation failure and suppression of uterine decidualization accompanied with decreased levels of serum progesterone were also observed in rats given DPT, a major metabolite of TPT.

Developmental Toxicity of Phenyltin Compounds

Table 3.2 presents the developmental toxicity studies on phenyltin compounds given to female animals during pregnancy. Several reports on the adverse effects of phenyltins on development of offspring following maternal exposure have been published. Female SD rats were given TPTA by gavage at 5, 10, or 15 mg/kg on days 6 to 15 of pregnancy (Giavini et al. 1980). TPTA caused a decrease in maternal body weight gain at 10 mg/kg and higher, an increase in postimplantation loss at 15 mg/kg, and a reduction of fetal ossification at 5 mg/kg and higher. Teratogenic effects of TPTA were not found even at doses resulting in clear maternal toxicity. Depression of maternal body weight gain and food intake at 9.0 mg/kg and higher, and increase in postimplantation embryonic loss and decrease in fetal ossification at 9.0 mg/kg and higher, but not teratogenic effects, were observed in Wistar rats after administration of TPTA at 1.5, 3.0, 6.0, 9.0, or 12.0 mg/kg by gavage on days 7 to 17 of pregnancy (Noda et al. 1991a). Behavioral effects of prenatal exposure to TPTA were reported. A transient increase in spontaneous locomotor activity and increased mortality during the lactation period were found in pups of CFY rats given TPTA by gavage at 6 mg/kg on days 6 to 14 of pregnancy (Lehotzky et al. 1982). In this study, maternal rats were free of any overt signs of toxicity. Disruptions of learning acquisition, as evidenced by low avoidance rate in the Sidman avoidance test, and prolonged swimming time to the goal, and an increased number of errors in a reversed test in the water E-maze, were observed in postnatal offspring of Tokai High Avoiders (THA) rats received TPTA by gavage on days 6 to 20 of pregnancy at 4 or 8 mg/kg (Miyake et al. 1991). Maternal deaths and decreased weight gain were found at 8 mg/kg, no maternal toxicity was observed at 4 mg/kg, and no malformed offspring appeared in any group.

Table 3.2 Developmental Toxicity of Phenyltin Compounds

Compounds	Animals	Dose	Days of Administration	Route	Reproductive and Developmental Effects	Author(s)
TPTA	Wistar rat	5-15 mg/kg	Days 6-15 of pregnancy	Gavage	Postimplantation loss, delayed ossification	Giavini et al. (1980)
TPTA	Wistar rat	9-12 mg/kg	Days 7-17 of pregnancy	Gavage	Postimplantation loss, delayed ossification	Noda et al. (1991a)
TPTA	CFY rat	6 mg/kg	Days 6-14 of pregnancy	Gavage	Postnatal death, transient increase in spontaneous locomotor activity	Lehotzky et al. (1982)
TPTA	THA rat	4-8 mg/kg	Days 6-20 of pregnancy	Gavage	Disruption of learning acquisition	Miyake et al. (1991)
TPTH	SD rat	20 mg/kg	Days 1-7 of pregnancy	Gavage	Decreased pregnancy rate	Winek et al. (1978)
		15 mg/kg	Days 8-14 of pregnancy	Gavage	Postimplantation loss, decreased fetal wt.	
TPTH	SD rat	15 mg/kg	Days 14-12 of pregnancy	Gavage	Effects as above	
		13 mg/kg	Days 6-15 of pregnancy	Gavage	Postimplantation loss	Chernoff et al. (1990)
TPTCl	Wistar rat	6.3-12.5 mg/kg	Days 7-9 of pregnancy	Gavage	Postimplantation loss	Ema et al. (1999c)
		9.4-12.5 mg/kg	Days 10-12 or 13-15 of pregnancy	Gavage	Postimplantation loss, decreased fetal wt.	