

Ethical Committee of Kagoshima University Faculty of Medicine approved this study.

#### Serum, CSF, and genomic DNA preparation

Fresh peripheral blood mononuclear cells (PBMCs) were obtained by Histopaque-1077 (Sigma, Tokyo, Japan) density gradient centrifugation, and washed three times with phosphate buffered saline (PBS) with 1% fetal calf serum (FCS). Isolated PBMCs were cryopreserved in liquid nitrogen until use. Genomic DNA was extracted from PBMCs using a QIAamp blood kit (Qiagen Ltd, Tokyo, Japan) according to the manufacturer's instructions. The CSF and serum samples were also collected, and stored at  $-70^{\circ}\text{C}$  until use.

#### Determination of polymorphism and provirus load measurement

The *aggrecan* gene contains a large exon (exon 12) of 3.5 kb, which encodes the entire glycosaminoglycan (GAG) attachment regions of its core protein (Doerge et al. 1991). This region consists of numerous repeated sequences, including a particularly highly conserved set of repeats in the CS attachment site. The VNTR polymorphism of the *aggrecan* gene in exon 10 has already been reported (Doerge et al. 1997). This VNTR can be detected by PCR as different lengths of PCR products. A genomic PCR was performed with 20 ng of genomic DNA as template, 50 pmol of each primer (forward: 5'-TAG AGG GCT CTG CCT CTG GAG TTG-3' and reverse: 5'-AGG TCC CCT ACC GCA GAG GTA GAA-3'), 20-mM deoxynucleotide triphosphates (dNTPs), 15-mM  $\text{MgCl}_2$ , reaction buffer provided by the manufacturer, and one unit of Takara-Taq DNA polymerase (Takara, Tokyo, Japan) in a final volume of 10  $\mu\text{l}$ . PCR conditions were as follows: initial denaturation at  $94^{\circ}\text{C}$  for 5 min, followed by 35 cycles of denaturation at  $94^{\circ}\text{C}$  for 60 s, annealing at  $58^{\circ}\text{C}$  for 60 s, and elongation at  $72^{\circ}\text{C}$  for 60 s with a final extension at  $72^{\circ}\text{C}$  for 10 min. PCR products were separated on 1% agarose gels, visualized by ethidium bromide staining, after which the products were determined. Several of the alleles differ by only one repeat in size (59 bp), and care was taken to identify these alleles using appropriate gels and size markers. Two independent readers scored the alleles.

Provirus load of the samples was measured by a quantitative PCR method using an ABI Prism 7700 (PE-Applied Biosystems) (Nagai et al. 1998).

#### Quantification of aggrecan in serum and CSF

Serum, as well as CSF aggrecan concentration, was measured in duplicate using a commercial ELISA kit (BiSource Europe S.A., Nivelles, Belgium). According to the manufacturer's instruction, the kit detects aggrecan and aggrecan fragments, and the assay system used is sensitive to detect 0.9 ng/ml of aggrecan in samples. Serum aggrecan levels in

normal adults ranged between 1 and 4.4  $\mu\text{g/ml}$ , whereas no information was available regarding CSF levels. When we needed to separate CSF aggrecan amounts into two groups, we selected 0.9 ng/ml as the cut-off level, as this value was the lowest value of the cut-off range and there was no previous report measuring CSF aggrecan concentration. Optical density at 450 nm was measured on the ImmunoMini NJ-2300 (Nippon Inter Med, Tokyo, Japan) and aggrecan concentration was determined by linear regression from a standard curve using the aggrecan supplied with the kit as standard.

#### Statistical analysis

Statistical analysis was performed using the SPSS for Windows release 7.0, run on an IBM-compatible computer (Analytical software, Version 7, Tallahassee, FL, USA). Comparison of whole-allele distribution between patients with HAM/TSP and HCs was performed using a chi-square test for  $2 \times 9$  contingency table with a significance level  $p < 0.05$ . The distribution of each allele and genotype of the VNTR polymorphism of the *aggrecan* gene in patients with HAM/TSP patients was compared with those in HCs using a chi-square test for a  $2 \times 92$  and  $2 \times 3$  contingency table. Bonferroni multiple adjustments (Motulsky 1995) were made to the level of significance because of the multiple comparisons for VNTR allele frequencies. This level was set at  $p < 0.0057$  ( $p = 1 - 0.95^{(1/9)}$ ).

To assure reproducibility of our study, we have combined  $p$  values from the analysis on two sets of populations and have compared the combined  $p$  values to the BADGE (Manly 2005).

Serum and CSF aggrecan levels in patients and controls in three different groups were compared using either ANOVA or Kruskal-Wallis test. A  $p$  value less than 0.05 was considered statistically significant. When aggrecan levels in three groups were different, multiple comparisons were done by Scheffe's test. We also performed multiple-hypotheses testing when it was needed, and the level was set at  $p < 0.017$  ( $p = 1 - 0.95^{(1/3)}$ ).

## Results

Frequency of aggrecan VNTR allele 28 was significantly higher in HAM/TSP than HCs and normal control

We applied two-step analysis on our cohort. We first typed 100 samples from each group observing nine aggrecan VNTR alleles, and found the difference between the groups [ $\chi^2 = 18.18$  ( $df = 8$ ),  $p = 0.019$ ]. We then proceeded to analyze whole samples in this study (227 HAM/TSP patients and 217 HCs) (Table 1).

Comparison of whole allele distribution among patients with HAM/TSP, in HCs and normal controls was performed using a chi-square test for  $3 \times 9$  contingency table with a significance level  $p < 0.05$ . This analysis has revealed

**Table 1** Distribution of aggrecan VNTR polymorphism in HAM/TSP patients, in healthy carriers (HCs) and in normal control subjects

Allele	Length	Allele*						Genotype**					
		HAM/TSP		HCs		Control		HAM/TSP		HCs		Control	
		Obs	Freq (%)	Obs	Freq (%)	Obs	Freq (%)	Homozygote	Heterozygote	Homozygote	Heterozygote	Homozygote	Heterozygote
32	1858	2	0.4	2	0.4	1	0.6	0	2	0	2	0	1
29	1687	18	3.9	12	2.7	6	3.5	2	14	3	6	0	6
28	1630	115	25.3	69	15.9	20	11.8	23	69	12	45	5	10
27	1573	141	31	155	35.7	57	33.5	32	77	37	81	13	31
26	1516	90	19.8	102	23.5	44	25.9	15	60	22	58	8	28
25	1459	62	13.6	53	12.2	25	14.7	4	54	9	35	1	23
22	1288	23	5	37	8.5	15	8.8	6	11	11	15	3	9
21	1231	2	0.4	3	0.6	1	0.6	1	0	1	1	0	1
18	1060	1	0.2	1	0.2	1	0.6	0	1	0	1	0	1
		454		434		170							

\* Comparison of whole allele distribution among patients with HAM/TSP, HCs and normal controls was performed using a chi-square test for 3×9 contingency table with a significance level  $p < 0.05$ . This analysis has revealed  $\chi^2 = 27.33 (df=16)$ ,  $p = 0.038$ .

Comparisons of whole allele distribution between each two groups were performed using a chi-square test for 2×9 contingency table with a significance level  $p < 0.05$ . This analysis has revealed  $\chi^2 = 17.84 (df=8)$ ,  $p = 0.02$  (HAM/TSP vs HCs),  $\chi^2 = 16.53 (df=8)$ ,  $p = 0.035$  (HAM/TSP vs normal controls), and  $\chi^2 = 3.24 (df=8)$ ,  $p = 0.918$  (HCs vs normal controls). The distribution of each allele of the VNTR polymorphism of the aggrecan gene in patients with HAM/TSP patients was compared with those in HCs using a chi-square test for a 2×2 contingency table. Allele 28 has been detected more frequently in patients group than HCs ( $\chi^2 = 12.02$ ,  $p = 0.0005$ , odds ratio 1.79, 95% C.I. 1.29–2.50).

\*\* The  $p$  value of genotype among three groups was calculated by  $\chi^2$  test with a 3×3 contingency table. This analysis revealed that genotype of the 28 repeat was frequently observed in HAM/TSP than HCs ( $\chi^2 = 19.68$ ,  $p = 0.003$ ,  $df = 6$ ). Then  $p$  values of genotype in each two groups were calculated by  $\chi^2$  test with a 2×3 contingency table. This analysis revealed that genotype of the 28 repeat was frequently observed in HAM/TSP than in HCs ( $\chi^2 = 10.41$ ,  $p = 0.005$ ,  $df = 2$ ) and in HAM/TSP than in normal controls ( $\chi^2 = 14.65$ ,  $p = 0.0007$ ,  $df = 2$ ), but not in HCs and in normal controls ( $\chi^2 = 3.31$ ,  $p = 0.19$ ,  $df = 2$ ).

$\chi^2 = 27.33$  ( $df = 16$ ),  $p = 0.038$ . Comparisons of whole allele distribution between each two groups were performed using a chi-square test for 2×9 contingency table with a significance level  $p < 0.05$ . This analysis has revealed  $\chi^2 = 17.84$  ( $df = 8$ ),  $p = 0.02$  (HAM/TSP vs HCs),  $\chi^2 = 16.53$  ( $df = 8$ ),  $p = 0.035$  (HAM/TSP vs normal controls), and  $\chi^2 = 3.24$  ( $df = 8$ ),  $p = 0.918$  (HCs vs normal controls).

Allele 28 was observed in 25.3% of HAM/TSP patients, whereas, only 15.9% of HCs and 11.8% of normal controls carried this allele. We, therefore, compared the distribution of allele 28 in patients with HAM/TSP and in HCs, and that in normal controls using a chi-square test for a 2×2 contingency table. As nine alleles appeared in our analysis, we set  $p < 0.0057$  ( $p = 1 - 0.95^{(1/9)}$ ) using the Bonferroni adjustment for multiple comparisons. Allele 28 has been detected more frequently in patients group than in HCs ( $\chi^2 = 12.02$ ,  $p = 0.0005$ , odd ratio 1.79, 95% C.I. 1.29–2.50) and than in

normal controls ( $\chi^2 = 13.43$ ,  $p = 0.0002$ , odd ratio 2.54, 95% C.I. 1.52–4.25).

The  $p$  value of genotype in three groups was calculated first by  $\chi^2$  test with a 3×3 contingency table. This analysis revealed that genotype of the 28 repeat was frequently observed in HAM/TSP than in HCs in and normal controls ( $\chi^2 = 19.68$ ,  $p = 0.003$ ,  $df = 6$ ). We, then, calculated the  $p$  values of genotype between two groups by  $\chi^2$  test with a 2×3 contingency table. This analysis revealed that genotype of the 28 repeat was frequently observed in HAM/TSP than in HCs ( $\chi^2 = 10.41$ ,  $p = 0.005$ ,  $df = 2$ ) and in HAM/TSP than in normal controls ( $\chi^2 = 14.65$ ,  $p = 0.0007$ ,  $df = 2$ ), but not in HCs and in normal controls ( $\chi^2 = 3.31$ ,  $p = 0.19$ ,  $df = 2$ ). The observed frequency of alleles other than allele 28 was very similar to the frequency reported in a European population (Doege et al. 1997).

**Table 2** Distribution of aggrecan 1630-bp allele in HAM/TSP patients at different provirus load

Provirus load	Total number of Patients	Patients with allele 1630	Frequency (%)
<100	28	10	35.7
<300	33	12	36.3
<600	48	18	37.5
<1000	34	14	41.2
<2000	46	18	39.1
>2000	16	8	50.0

Provirus load is presented as number of the cells in  $10^4$  PBMC

Mann–Whitney's  $U$  test has revealed that the distribution of allele 1630 positive patients is not different at different provirus load ( $p = 0.402$ )

We assessed the reproducibility of our study by comparing combined  $p$  values and the BADGE system (Manly 2005). We first multiplied the  $p$  value for the  $3 \times 9$   $\chi^2$ -square test from the first population (0.038) and that of the  $2 \times 9$   $\chi^2$ -square test from the second population (0.0001). This yielded the combined  $p$  value of 0.0000038. This estimate suggested that the association of our study should be regarded as a second-class association in the BADGE system. We also tried to assess the reproducibility on the test applied to allele 28. We multiplied the  $p$  value from the first population (0.0005) with that from the second population (0.0009) and applied Bonferroni correction by multiplying 9 on each  $p$  value. This produced the combined  $p$  value of 0.00004 as a third-class association.

The possession of allele 28 was not related to disease progression or HTLV-I provirus load

Of 52 HAM patients with CSF aggrecan analyzed, eight patients with allele 1630 showed rapid progression while ten were without this allele. A chi-square test for  $2 \times 2$  contingency table revealed that disease progression was not correlated with allele possession ( $\chi^2=0.188$ ,  $p=0.66$ , odds ratio 1.29, 95% C.I. 0.41–4.12).




Fig. 1 The amount of aggrecan in serum (normal range 1–4.4  $\mu\text{g/ml}$ , shadowed area) showed similar level among HAM/TSP patients with rapid progression, with slow progression and HTLV-I carriers (HCs)




Fig. 2 The amount of aggrecan in CSF of patients with HAM/TSP showing rapid or slow progression, and other non-inflammatory disease (OND)

The distribution of allele 1630 positive patients in the entire sample of 205 HAM patients

In these samples, the provirus load was measured and was not correlated with the provirus load (Table 2) (Mann-Whitney's  $U$  test,  $p=0.402$ ). We measured the provirus load of samples when we had an approval to measure the provirus load.

The CSF level of aggrecan was significantly higher in rapidly progressive HAM/TSP than in chronic HAM/TSP

We, next, compared the serum aggrecan level in rapidly progressive and chronically progressive HAM/TSP patients and HCs (Fig. 1). However, there was no significant difference among these three groups ( $F=0.78$ ,  $p=0.47$ ). We then compared CSF aggrecan levels among rapidly and chronically progressive HAM/TSP patients as well as OND (Fig. 2). The results showed that aggrecan levels in CSF in the three different groups were different by Kruskal-Wallis test ( $H=13.45$ ,  $df=2$ ,  $p=0.0006$ , corrected  $p=0.0018$ ) and the level in the rapidly progressive HAM/TSP patients was significantly higher than that in the chronically progressive HAM/TSP ( $p=0.0049$ , corrected  $p=0.0145$ ) but not in that of OND ( $p=0.026$ , corrected  $p=0.078$ ) (Scheffe's test).

**Table 3** Distribution of aggrecan VNTR polymorphism in the second group of HAM/TSP patients compared with the second healthy carrier (HCs) group

Allele	Length	Allele*				Genotype**			
		HAM/TSP		HCs		HAM/TSP		HCs	
		Obs	Freq (%)	Obs	Freq (%)	Homozygote	Heterozygote	Homozygote	Heterozygote
32	1858	0	0	2	1.5	0	0	0	2
29	1687	3	2.6	7	5	1	1	2	3
28	1630	31	26.8	15	10.8	7	17	3	9
27	1573	42	36.3	49	35	5	32	8	33
26	1516	17	14.7	34	24.3	0	17	3	28
25	1459	13	11.3	26	18.6	0	13	1	24
22	1288	10	8.4	1	0.8	0	10	0	1
18	1060	0	0	2	1.5	0	0	0	2

\* Comparison of whole allele distribution among patients with HAM/TSP and HCs in the second group was performed using a chi-square test for 2×9 contingency table with a significance level  $p < 0.05$ . This analysis has revealed  $\chi^2 = 31.09 (df=8)$ ,  $p = 0.0001$ . The distribution of each allele of the VNTR polymorphism of the aggrecan gene in patients with HAM/TSP patients was compared with those in HCs using a chi-square test for a 2×2 contingency table. Allele 28 has been detected more frequently in patients group than HCs ( $\chi^2 = 11.03$ ,  $p = 0.0009$ , odds ratio 3.04, 95% C.I. 1.55–5.97)

\*\* The  $p$  value of genotype was calculated by  $\chi^2$  test with a 2×3 contingency table. This analysis revealed that the genotype of 28 repeat was frequently observed in HAM/TSP than HCs ( $\chi^2 = 9.28$ ,  $df = 2$ ,  $p = 0.009$ )

#### Aggrecan VNTR analysis in the different set of patients with HAM/TSP and HCs

We, then, analyzed this aggrecan VNTR polymorphism in the different set of patients with HAM/TSP ( $n = 58$ ) and healthy carriers ( $n = 70$ ) (Table 3). We performed this second analysis to ensure our first observation. Comparisons of whole allele distribution between two groups were performed using a chi-square test for 2×9 contingency table with a significance level  $p < 0.05$ . This analysis has revealed  $\chi^2 = 31.09 (df=8)$ ,  $p = 0.0001$ . Allele 28 was observed in 26.8% of HAM/TSP patients and 10.8% of HCs in this second analysis. We compared the distribution of allele 28 in patients with HAM/TSP and that in HCs using a chi-square test for a 2×2 contingency table. The allele 28 was detected more frequently in HAM/TSP group than in HCs ( $\chi^2 = 11.03$ ,  $p = 0.0009$ , odds ratio 3.04, 95% C.I. 1.55–5.97). The  $p$  value of genotype was calculated by  $\chi^2$  test with a 2×3 contingency table. This analysis revealed that the genotype of 28 repeat was frequently observed in HAM/TSP than HCs ( $\chi^2 = 9.28$ ,  $df = 2$ ,  $p = 0.009$ ).

#### Discussion

In this study, we report three findings. First, allele 28 (1630 bp) of the aggrecan gene was more frequently observed in HAM/TSP patients than in HCs and in normal controls. This frequent distribution of allele 28 was observed also in the different set of HAM/TSP patients and HCs. The reproducibility of our study was assessed by comparing combined  $p$  values and the BADGE system (Manly 2005) and was regarded as a second- or third-class association. Second, possession of allele 28 was not related to the disease progression or HTLV-I provirus load. Finally, the rapidly progressive HAM/TSP patients showed a higher aggrecan

concentration in the CSF than the chronically progressive HAM/TSP patients.

Recent genetic analysis of the aggrecan gene has shown that it has 18 exons and that there is a polymorphic region in the 12th exon, which is the CS attachment site (Doege et al. 1991). This site has a VNTR of 57 bp. Using this VNTR, several reports have analyzed whether there is a correlation between osteoarthritis (OA) of the hand and a particular allele of the aggrecan gene (Horton et al. 1998). Another study of aggrecan gene VNTR polymorphism has shown that individuals with shorter VNTR tend to develop multilevel disc degeneration at an earlier age (Kawaguchi et al. 1999). Even though no disease association of aggrecan VNTR has been shown in chronic inflammatory or immunological disease of the nervous system, the reported nature and function of aggrecan and these association studies prompted us to investigate its relation to HTLV-I-related neurological diseases. Ours is the second report of aggrecan VNTR allele distribution in the Asian population, but the first study to examine the association between aggrecan polymorphism and a neurological disease. Regarding allele 28, Kawaguchi et al. (Kawaguchi et al. 1999) reported that allele frequency was 9.4% in their studied population, whereas we have observed 25.3% in patients with HAM/TSP, 15.9% in HCs and 11.8% in normal controls from our area (Table 1). We have, again, shown that the allele frequency of this allele 28 was 26.8% in HAM/TSP patients and 10.8% in HCs (Table 3). This has shown that the allele 28 is indeed increased in our patient population and there was no possibility for a population stratification artifact. To estimate the reproducibility of our study, we have employed the BADGE system to describe genetic association (Manly 2005). As shown in the results, the association of whole allele distribution of aggrecan gene to HAM/TSP has reached second-class and the association of allele 28 has reached a third-class association. We have, therefore,

assumed that our study suggests reproducibility under conservative assumptions for traits previously mapped to a chromosome or a small region.

Aggrecan was recently reported to be produced by astrocytes and to exist in the perineurial region of the CNS (Matthews et al. 2002). In general, aggrecan degenerates with age and is cleaved between the G1 domain and the KS binding domain by proteolysis with the enzyme aggrecanase (Lark et al. 1997). Fragments of aggrecan are produced by aging, mechanical processes and/or activation of cleaving enzymes. Once these fragments activate T cells, these T cells can infiltrate into the CNS through the blood-brain barrier and initiate inflammatory CNS diseases (Buzas et al. 1995; Lemons et al. 1999; Mikecz et al. 1988; Zhang et al. 1998b). The reported lower concentration of CS and lack of KS in brain aggrecan (Buzas et al. 1995; Glant et al. 1998; Koppe et al. 1997) may be related to this elicited immune response in the CNS, as decrease of CS or KS content are reported to generate T- or B-cell immune response (Glant et al. 1998). Previous studies on the pathological mechanism of HAM/TSP have revealed that the main disease process is T-cell-mediated inflammation of the thoracic spinal cord (Izumo et al. 2000; Umehara et al. 1993). Taking these findings together, it is of interest to know that the length of the CS attachment site determined by VNTR may have a correlation with HAM/TSP.

Next, we were not able to show the correlation between possession of allele 28 and disease progression or HTLV-I provirus load. This may be because the genetic background we have found in this study can be one independent factor in causing HAM/TSP. Our previous studies have revealed that higher provirus load correlates with strong inflammation of the spinal cord and that the load is related to the deterioration of motor disability in 64 HAM/TSP patients followed up for 10 years (Matsuzaki et al. 2001). We also reported that there were HAM/TSP patients with lower provirus load (Nakagawa et al. 1995). From these observations, we speculated that tissue damage during immune inflammation might not only be controlled only by the strength of the inflammation itself but by the strength of the tissue structure as well. Weak inflammation is sufficient when inflammation occurs in a genetically determined weak tissue. In this regard, our present study may open a novel approach in finding the cause of HTLV-I-related neurological diseases.

To investigate whether aggrecan leakage correlates with disease progression, we measured aggrecan concentration in sera of HAM/TSP patients and HCs, and in CSF of HAM/TSP patients and OND. We found higher CSF aggrecan concentration in rapidly progressive HAM/TSP patients than in chronically progressive patients. As our previous clinical analysis of HAM/TSP patients showed that the patients with later disease onset and knee-joint arthritis showed faster progression of the disease (Nakagawa et al. 1995), we speculated that aggrecan that leaked into the CSF was caused by the degradation of spinal cord tissue secondary to inflammation induced by HTLV-I infection. We also showed that the degree of aggrecan degradation was higher in rapidly progressive patients in this study, and would, therefore, like

to propose that the concentration of aggrecan in CSF may be a marker for denaturing in the spinal cord. Although HAM/TSP is reported to occur more frequently in female and we have observed slightly more male cases in rapid progressive group than expected by the reported ratio, age of onset is the only factor, so far, that has been shown to correlate with the disease progression rate (Nakagawa et al. 1995). To find a correlation between sex and disease progression, we may need to measure aggrecan concentration in more cases. To our knowledge, this is the first study to show the presence of aggrecan in CSF. Analysis of CSF aggrecan in other neurological diseases may clarify the significance of this molecule.

As aggrecan and other proteoglycan family molecules play a role in neuronal regeneration and tissue repair after CNS injury (Davies et al. 1997; Gates et al. 1996; Koppe et al. 1997; Lemons et al. 1999), our present observation suggests the possibility that the genetically determined nature of aggrecan determines the efficiency of tissue damage of the spinal cord. This may explain the axonal damage of the spinal cord observed in HAM/TSP patients (Umehara et al. 2000). Profound spinal tissue damage after acute inflammation caused by HTLV-I infected T cells may lead to an acute course of the disease, and insufficient or excessive repair of spinal tissue due to the genetic background may accumulate in a chronic course of the disease. Further studies are necessary to clarify these points.

**Acknowledgements** We thank Ms Tomoko Muramoto and Yoko Nishino of Kagoshima University for their excellent technical assistance.

Part of this study has been presented at the 11th International Conference on Human Retrovirology: HTLV and Related Viruses in San Francisco from June 9 to 12, 2003. All subjects gave written informed consent. The Ethical Committee of Kagoshima University Faculty of Medicine approved this study. The authors do not have commercial or other associations that might pose a conflict of interest.

## References

- Adams LA, Ang LC, Munoz DG (1993) Chromogranin A, a soluble synaptic vesicle protein, is found in cortical neurons other than previously defined peptidergic neurons in the human neocortex. *Brain Res* 602:336-341
- Ang LC, Zhang Y, Cao L, Yang BL, Young B, Kiani C, Lee V, Allan K, Yang BB (1999) Versican enhances locomotion of astrocytoma cells and reduces cell adhesion through its G1 domain. *J Neuropathol Exp Neurol* 58:597-605
- Asher RA, Scheibe RJ, Keiser HD, Bignami A (1995) On the existence of a cartilage-like proteoglycan and link proteins in the central nervous system. *Glia* 13:294-308
- Asher RA, Morgenstern DA, Fidler PS, Adcock KH, Oohira A, Braistead JE, Levine JM, Margolis RU, Rogers JH, Fawcett JW (2000) Neurocan is upregulated in injured brain and in cytokine-treated astrocytes. *J Neurosci* 20:2427-2438
- Becker KG, Barnes KC, Bright TJ, Wang SA (2004) The genetic association database. *Nat Genet* 36:431-432
- Buzas EI, Brennan FR, Mikecz K, Garzo M, Negroiu G, Hollo K, Cs-Szabo G, Pintye E, Glant TT (1995) A proteoglycan (aggrecan)-specific T cell hybridoma induces arthritis in BALB/c mice. *J Immunol* 155:2679-2687

- Cardon LR, Bell JI (2001) Association study design for complex diseases. *Nat Rev Genet* 2:91–99
- Colhoun HM, McKeigue PM, Davey Smith G (2003) Problems of reporting genetic associations with complex outcomes. *Lancet* 361: 865–872
- Davies SJ, Fitch MT, Memberg SP, Hall AK, Raisman G, Silver J (1997) Regeneration of adult axons in white matter tracts of the central nervous system. *Nature* 390:680–683
- Doerge KJ, Sasaki M, Kimura T, Yamada Y (1991) Complete coding sequence and deduced primary structure of the human cartilage large aggregating proteoglycan, aggrecan. Human-specific repeats, and additional alternatively spliced forms. *J Biol Chem* 266:894–902
- Doerge KJ, Coulter SN, Meek LM, Maslen K, Wood JG (1997) A human-specific polymorphism in the coding region of the *aggrecan* gene. Variable number of tandem repeats produce a range of core protein sizes in the general population. *J Biol Chem* 272:13974–13979
- Feng L, Precht P, Balakir R, Horton WE Jr (1998) Evidence of a direct role for Bcl-2 in the regulation of articular chondrocyte apoptosis under the conditions of serum withdrawal and retinoic acid treatment. *J Cell Biochem* 71:302–309
- Furukawa Y, Yamashita M, Usuku K, Izumo S, Nakagawa M, Osame M (2000) Phylogenetic subgroups of human T cell lymphotropic virus (HTLV) type I in the tax gene and their association with different risks for HTLV-I-associated myelopathy/tropical spastic paraparesis. *J Infect Dis* 182:1343–1349
- Gates MA, Fillmore H, Steindler DA (1996) Chondroitin sulfate proteoglycan and tenascin in the wounded adult mouse neostriatum in vitro: dopamine neuron attachment and process outgrowth. *J Neurosci* 16:8005–8018
- Gessain A, Barin F, Vernant JC, Gout O, Maurs L, Calender A, de TG (1985) Antibodies to human T-lymphotropic virus type-I in patients with tropical spastic paraparesis. *Lancet* 2:407–410
- Glant TT, Buzas EI, Finnegan A, Negroiu G, Cs-Szabo G, Mikecz K (1998) Critical roles of glycosaminoglycan side chains of cartilage proteoglycan (aggrecan) in antigen recognition and presentation. *J Immunol* 160:3812–3819
- Grumet M, Flaccus A, Margolis RU (1993) Functional characterization of chondroitin sulfate proteoglycans of brain: interactions with neurons and neural cell adhesion molecules. *J Cell Biol* 120:815–824
- Hirschhorn JN, Lohmueller K, Byrne E, Hirschhorn K (2002) A comprehensive review of genetic association studies. *Genet Med* 4:45–61
- Horton WE Jr, Lethbridge-Cejku M, Hochberg MC, Balakir R, Precht P, Plato CC, Tobin JD, Meek L, Doerge K (1998) An association between an aggrecan polymorphic allele and bilateral hand osteoarthritis in elderly white men: data from the Baltimore Longitudinal Study of Aging (BLSA). *Osteoarthritis Cartilage* 6:245–251
- Ioannidis JP, Ntzani EE, Trikalinos TA, Contopoulos-Ioannidis DG (2001) Replication validity of genetic association studies. *Nat Genet* 29: 306–309
- Izumo S, Umebara F, Osame M (2000) HTLV-I-associated myelopathy. *Neuropathology* 20 Suppl:S65–S68
- Jeffery KJ, Usuku K, Hall SE, Matsumoto W, Taylor GP, Procter J, Bunce M, Ogg GS, Welsh KI, Weber JN, Lloyd AL, Nowak MA, Nagai M, Kodama D, Izumo S, Osame M, Bangham CR (1999) HLA alleles determine human T-lymphotropic virus-I (HTLV-I) proviral load and the risk of HTLV-I-associated myelopathy. *Proc Natl Acad Sci U S A* 96:3848–3853
- Jeffery KJ, Siddiqui AA, Bunce M, Lloyd AL, Vine AM, Witkover AD, Izumo S, Usuku K, Welsh KI, Osame M, Bangham CR (2000) The influence of HLA class I alleles and heterozygosity on the outcome of human T cell lymphotropic virus type I infection. *J Immunol* 165:7278–7284
- Kawaguchi Y, Osada R, Kanamori M, Ishihara H, Ohmori K, Matsui H, Kimura T (1999) Association between an *aggrecan* gene polymorphism and lumbar disc degeneration. *Spine* 24: 2456–2460
- Koppe G, Bruckner G, Hartig W, Delpesch B, Bigl V (1997) Characterization of proteoglycan-containing perineuronal nets by enzymatic treatments of rat brain sections. *Histochem J* 29:11–20
- Lark MW, Bayne EK, Flanagan J, Harper CF, Hoerner LA, Hutchinson NI, Singer II, Donatelli SA, Weidner JR, Williams HR, Mumford RA, Lohmander LS (1997) Aggrecan degradation in human cartilage. Evidence for both matrix metalloproteinase and aggrecanase activity in normal, osteoarthritic, and rheumatoid joints. *J Clin Invest* 100:93–106
- Lemons ML, Howland DR, Anderson DK (1999) Chondroitin sulfate proteoglycan immunoreactivity increases following spinal cord injury and transplantation. *Exp Neurol* 160:51–65
- Levin MC, Lee SM, Kalume F, Morcos Y, Dohan FC Jr, Hasty KA, Callaway JC, Zunt J, Desiderio D, Stuart JM (2002) Autoimmunity due to molecular mimicry as a cause of neurological disease. *Nat Med* 8:509–513
- Manly KF (2005) Reliability of statistical associations between genes and disease. *Immunogenetics* 57:549–558
- Matsuzaki T, Nakagawa M, Nagai M, Usuku K, Higuchi I, Arimura K, Kubota H, Izumo S, Akiba S, Osame M (2001) HTLV-I proviral load correlates with progression of motor disability in HAM/TSP: analysis of 239 HAM/TSP patients including 64 patients followed up for 10 years. *J Neurovirol* 7:228–234
- Matthews RT, Kelly GM, Zerillo CA, Gray G, Tiemeyer M, Hockfield S (2002) Aggrecan glycoforms contribute to the molecular heterogeneity of perineuronal nets. *J Neurosci* 22:7536–7547
- Mikecz K, Glant TT, Baron M, Poole AR (1988) Isolation of proteoglycan-specific T lymphocytes from patients with ankylosing spondylitis. *Cell Immunol* 112:55–63
- Milev P, Maurel P, Chiba A, Mevissen M, Popp S, Yamaguchi Y, Margolis RK, Margolis RU (1998) Differential regulation of expression of hyaluronan-binding proteoglycans in developing brain: aggrecan, versican, neurocan, and brevican. *Biochem Biophys Res Commun* 247:207–212
- Moon LD, Asher RA, Fawcett JW (2003) Limited growth of severed CNS axons after treatment of adult rat brain with hyaluronidase. *J Neurosci Res* 71:23–37
- Moritoyo T, Reinhart TA, Moritoyo H, Sato E, Izumo S, Osame M, Haase AT (1996) Human T-lymphotropic virus type I-associated myelopathy and tax gene expression in CD4+ T lymphocytes. *Ann Neurol* 40:84–90
- Motulsky H (1995) *Multiple comparisons. Intuitive biostatistics.* Oxford University Press, New York
- Nagai M, Usuku K, Matsumoto W, Kodama D, Takenouchi N, Moritoyo T, Hashiguchi S, Ichinose M, Bangham CR, Izumo S, Osame M (1998) Analysis of HTLV-I proviral load in 202 HAM/TSP patients and 243 asymptomatic HTLV-I carriers: high proviral load strongly predisposes to HAM/TSP. *J Neurovirol* 4:586–593
- Nakagawa M, Izumo S, Ijichi S, Kubota H, Arimura K, Kawabata M, Osame M (1995) HTLV-I-associated myelopathy: analysis of 213 patients based on clinical features and laboratory findings. *J Neurovirol* 1:50–61
- Nishioka K, Maruyama I, Sato K, Kitajima I, Nakajima Y, Osame M (1989) Chronic inflammatory arthropathy associated with HTLV-I. *Lancet* 1:441
- Oohira A, Matsui F, Tokita Y, Yamauchi S, Aono S (2000) Molecular interactions of neural chondroitin sulfate proteoglycans in the brain development. *Arch Biochem Biophys* 374: 24–34
- Osame M (1990) Review of WHO Kagoshima meeting and diagnostic guidelines for HAM/TSP. Blattener, W. A. *Human Retrovirology; HTLV.* 191–197. Raven, New York
- Osame M, Usuku K, Izumo S, Ijichi N, Amitani H, Igata A, Matsumoto M, Tara M (1986) HTLV-I associated myelopathy, a new clinical entity. *Lancet* 1:1031–1032
- Perris R, Perissinotto D (2000) Role of the extracellular matrix during neural crest cell migration. *Mech Dev* 95:3–21

- Poole AR (1998) The histopathology of ankylosing spondylitis: are there unifying hypotheses? *Am J Med Sci* 316:228-233
- Redden DT, Allison DB (2003) Nonreplication in genetic association studies of obesity and diabetes research. *J Nutr* 133:3323-3326
- Takahashi-Iwanaga H, Murakami T, Abe K (1998) Three-dimensional microanatomy of perineuronal proteoglycan nets enveloping motor neurons in the rat spinal cord. *J Neurocytol* 27:817-827
- Umehara F, Izumo S, Nakagawa M, Ronquillo AT, Takahashi K, Matsumuro K, Sato E, Osame M (1993) Immunocytochemical analysis of the cellular infiltrate in the spinal cord lesions in HTLV-I-associated myelopathy. *J Neuropathol Exp Neurol* 52:424-430
- Umehara F, Abe M, Koreeda Y, Izumo S, Osame M (2000) Axonal damage revealed by accumulation of beta-amyloid precursor protein in HTLV-I-associated myelopathy. *J Neurol Sci* 176:95-101
- Vine AM, Witkover AD, Lloyd AL, Jeffery KJ, Siddiqui A, Marshall SE, Bunce M, Eiraku N, Izumo S, Usuku K, Osame M, Bangham CR (2002) Polygenic control of human T lymphotropic virus type I (HTLV-I) provirus load and the risk of HTLV-I-associated myelopathy/tropical spastic paraparesis. *J Infect Dis* 186:932-939
- Watanabe H, Yamada Y, Kimata K (1998) Roles of aggrecan, a large chondroitin sulfate proteoglycan, in cartilage structure and function. *J Biochem (Tokyo)* 124:687-693
- Zhang Y, Guerassimov A, Leroux JY, Cartman A, Webber C, Lalic R, de ME, Rosenberg LC, Poole AR (1998a) Arthritis induced by proteoglycan aggrecan G1 domain in BALB/c mice. Evidence for T-cell involvement and the immunosuppressive influence of keratan sulfate on recognition of T and B cell epitopes. *J Clin Invest* 101:1678-1686
- Zhang Y, Guerassimov A, Leroux JY, Cartman A, Webber C, Lalic R, de ME, Rosenberg LC, Poole AR (1998b) Induction of arthritis in BALB/c mice by cartilage link protein: involvement of distinct regions recognized by T and B lymphocytes. *Am J Pathol* 153:1283-1291

## HTLV-I viral escape and host genetic changes in the development of adult T Cell leukemia

Yoshitaka Furukawa<sup>1\*</sup>, Mitsutoshi Tara<sup>2</sup>, Shuji Izumo<sup>3</sup>, Kimiyoshi Arimura<sup>4</sup> and Mitsuhiro Osame<sup>4</sup>

<sup>1</sup>Division of Blood Transfusion Medicine and Cell Therapy, Kagoshima University Hospital, Kagoshima, Japan

<sup>2</sup>Department of Internal Medicine, Kagoshima City Hospital, Kagoshima Japan

<sup>3</sup>Center for Chronic Viral Diseases, Faculty of Medicine, Kagoshima University, Japan

<sup>4</sup>Department of Neurology and Geriatrics, Kagoshima University Graduate School of Medical and Dental Science, Japan

In the pathogenesis of adult T cell leukemia (ATL), an oncogenic role of the human T cell lymphotropic virus type I (HTLV-I) Tax protein, viral escape from the host immune system, and host genetic changes have been proposed as contributory factors. We examined the premature stop codons in *tax* gene as one of the mutations that may lead to escape of HTLV-I from the cytotoxic T lymphocyte (CTL) response in HTLV-I carriers, to test whether a putative CTL escape mutant can emerge in the early stage of ATL development and whether HTLV-I infected cells with such a mutation can proliferate subsequently. We also examined deletion of cyclin-dependent kinase inhibitor 4 (*INK4*) genes and mutation of *p53* gene in combination with changes in the HTLV-I genome in acute type ATL to test whether host genetic changes promoted the malignant transformation of ATL cells that carry putative CTL escape mutations. The premature stop codon in *tax* gene existed in many non-ATL HTLV-I carriers as a minor population but not in the commonest HTLV-I sequence of the individual. This minor population with a premature stop codon did not expand subsequently in 3 asymptomatic carriers tested. There were cases who had a mutation or deletion in HTLV-I who also have either deletion of *INK4* genes or mutation in *p53* gene. Our findings suggest that CTL escape mutation can occur at an early stage of ATL development, and that certain host genetic changes favor the development of the aggressive form of ATL.

© 2005 Wiley-Liss, Inc.

**Key words:** viral escape; *INK4*; *p53*; ATL

Adult T cell leukemia (ATL) is a T cell malignancy with clonal proliferation of human T cell leukemia virus type I (HTLV-I) infected cells.<sup>1,2</sup> HTLV-I is also an etiologic agent for HTLV-I-associated myelopathy/tropical spastic paraparesis (HAM/TSP).<sup>3,4</sup> ATL is subdivided into 4 types (smoldering, chronic, lymphoma and acute).<sup>5</sup> ATL has a long incubation period and smoldering and chronic type ATL sometimes transform into a more aggressive acute/lymphoma type of ATL, suggesting a multistep leukemogenesis model<sup>6</sup> for the development of ATL.

HTLV-I Tax protein is a key regulator for immortalization, transformation and oncogenesis of the HTLV-I infected lymphocytes through its interaction with many cellular proteins. For example, Tax binds to CBP/p300 and determines the accessibility of CBP/p300 to protein complexes on specific DNA elements,<sup>7</sup> resulting in Tax mediated trans-activation of viral genes<sup>8</sup> and growth factors,<sup>9</sup> or trans-repression of *p18*,<sup>10</sup> DNA polymerase  $\beta$  and *bax* genes.<sup>11</sup> Tax also modifies the cell cycle through binding *p16*<sup>INK4A</sup>,<sup>12</sup> *hDLG*<sup>13</sup> and *MAD1*<sup>14</sup> and contributes to the development of ATL.

Tax also plays a role as an immunodominant target antigen for the cytotoxic T lymphocyte response (CTL)<sup>15,16</sup> to HTLV-I and Tax-expressing cells will be rejected by the host immune response. It is possible therefore that immortalized cells that elicit weaker CTL responses are selected during the development of ATL. We have reported previously several mutations and deletions in the *tax* gene in ATL that can escape from the host immune system.<sup>17</sup> A premature stop codon in the *tax* gene (substitution at nt. position 7464 from G–A) that was observed frequently in ATL, is one such escape mutation, because the resulting truncated Tax protein loses its transactivational activity<sup>18</sup> and expression of

HTLV-I related proteins is diminished. This premature stop codon was also observed in the chronic type of ATL as a consensus sequence of the patient,<sup>17</sup> suggesting that Tax is dispensable at least in some chronic ATL cases.

These findings suggest that viral escape from CTL recognition is one of the important steps for the development of ATL; however host genetic changes such as cyclin-dependent kinase inhibitor genes and *p53* gene have also been observed in ATL.

The signaling pathway governed by G1 cyclins, cyclin-dependent kinases (CDK), pRb and E2F plays a major regulatory role during G1 to S transition in the cell cycle.<sup>19,20</sup> The complex formed by CDK4 and D-type cyclins controls the passage of cells through G1 phase, and the function of CDK4/CDK6 complexes is inhibited by a number of inhibitor of CDK4 (*INK4*), i.e., *p15*<sup>INK4B</sup>, *p16*<sup>INK4A</sup>, *p18*<sup>INK4C</sup> and *p19*<sup>INK4D</sup>.<sup>21–25</sup> The human *p16*<sup>INK4A</sup> and *p15*<sup>INK4B</sup> genes are situated within 30kb on chromosome 9p21.<sup>26</sup> *p18*<sup>INK4C</sup> and *p19*<sup>INK4D</sup> proteins also inhibit the activities of D-type CDK.<sup>23–25</sup> Among these *INK4*, however, *p16*<sup>INK4A</sup> is most impaired frequently in tumor cells.<sup>27</sup> There is another tumor suppressor gene named *p14*<sup>ARF</sup><sup>28</sup> encoded in an alternative reading frame (ARF) of the *p16*<sup>INK4A</sup> gene. The *p53* gene is another tumor suppressor gene.<sup>29</sup> Mutations of the *p53* gene have been found in several malignancies including ATL.<sup>30</sup>

We wondered when a premature stop codon in the *tax* gene, a putative CTL escape mutant, emerged during the development of ATL, and whether HTLV-I with this stop codon would subsequently proliferate. To this end, we examined the occurrence of a premature stop codon in the *tax* gene in 219 asymptomatic carriers and 143 HAM/TSP patients. We also examined the proportion of HTLV-I infected cells with this stop codon in 3 asymptomatic carriers at different time points to test whether such HTLV-I infected cells continuously proliferate in asymptomatic HTLV-I carriers without ATL. We also examined the deletion of cyclin-dependent kinase (CDK) 4 inhibitor genes (*p15*<sup>INK4B</sup>, *p16*<sup>INK4A</sup>, *p18*<sup>INK4C</sup>, *p19*<sup>INK4D</sup>) in 23 acute ATL patients and mutations of *p53* gene in 22 ATL patients to investigate whether additional host genetic changes favor the development of the aggressive form of ATL.

Presented in part at the 46th Conference Annual Meeting of the Japanese Society of Hematology, Kyoto, Japan, 19 September 2004.

The authors have no commercial or other associations that might pose a conflict of interest.

Informed consent was obtained from all HTLV-I carriers and patients. This research was approved by the institutional review boards of the author's institutions, and human experimentation guidelines of the US Department of Health and Human Services and those of the author's institutions were followed in the conduct of clinical research.

Grant sponsor: Japan Leukemia Research Fund.

\*Correspondence to: Division of Blood Transfusion Medicine and Cell Therapy, Kagoshima University Hospital, 8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan. Fax: +81-99-275-5641.

E-mail: furukawy@m2.kufm.kagoshima-u.ac.jp

Received 29 December 2004; Accepted after revision 13 May 2005

DOI 10.1002/ijc.21328

Published online 28 July 2005 in Wiley InterScience (www.interscience.wiley.com).



## Material and methods

### Study population

Two hundred and nineteen HTLV-I seropositive asymptomatic blood donors (AC) and 143 cases of HAM/TSP whose *tax* gene had not been sequenced in our previous study<sup>17</sup> were examined for the stop codon in the *tax* gene. Twenty-three cases of acute type ATL whose *tax* genes were sequenced in the previous study<sup>17</sup> were tested for the deletion of cyclin-dependent kinase inhibitor 4 genes and 22 of these cases were tested for a mutation in the *p53* gene. All cases were of Japanese ethnic origin and resided in Kagoshima prefecture (Japan). The diagnosis and clinical subtyping of ATL were made according to Shimoyama's criteria.<sup>5</sup> The diagnosis of HAM/TSP was made according to WHO diagnostic criteria.<sup>31</sup>

### Proviral load measurement

The HTLV-I provirus load in peripheral blood mononuclear cells (PBMC) was measured in HAM/TSP patients and HC as described.<sup>32</sup> A quantitative PCR reaction was carried out using the ABI PRISM 7700 sequence detector (Perkin-Elmer Applied Biosystems, Tokyo, Japan). The amount of HTLV-I proviral DNA was calculated as follows: copy number of HTLV-I (*tax*) per 10<sup>4</sup> PBMC = [copy number of *tax*/(copy number of  $\beta$ -actin/2)]  $\times$  10<sup>4</sup>. The lower limit of detection was 1 copy/10<sup>4</sup> PBMC.

### Statistical analysis

The Mann-Whitney *U*-test was used for statistical analysis of HTLV-I provirus load and the variables were treated as continuous.

### Restriction fragment length polymorphism analysis of the HTLV-I *tax* gene

Substitution at nt. position 7464 from G–A on the *tax* gene created a premature stop codon<sup>17</sup> and also created a Bln I restriction site (CCTGGG–CCTAGG). This stop codon was observed frequently in ATL in the commonest sequence of the individual ATL patients. Restriction fragment length polymorphism (RFLP) analysis using Bln I was done on 219 AC and in 143 HAM/TSP cases. Nested polymerase chain reaction (PCR) was carried out on the extracted DNA to amplify proviral DNA and the amplified product was digested with Bln I. One hundred nanograms of DNA were amplified by 35 cycles of PCR using expand high fidelity PCR system (Boehringer Mannheim, Tokyo, Japan) and 1  $\mu$ M primers (PX01+: 5'-TCGAAACAGCCCTGCAGATA-3' [7257–7276] and PX02–: 5'-TGAGCTTATGATTGTCTCA-3' [8447–8467]). After the first PCR reaction, 1  $\mu$ l aliquots of the amplified products were subjected to further 20 cycles of the second PCR using internal primers (PX11+: 5'-ATACAAAGTTAAC-CATGCTT-3' [7274–7293] and PX11–: 5'-GGGTTCCATG-TATCCATTC-3' [7644–7663]). Each PCR cycle consisted of denaturation at 94°C for 60 sec, annealing at 58°C for 75 sec, extension at 72°C for 90 sec and extension of the final cycle at 72°C for 10 min. Two  $\mu$ l of the nested PCR product was digested with 5 U of Bln I (Takara, Tokyo, Japan) in 10  $\mu$ l volume at 37°C for 18 hr and was then electrophoresed on 1% agarose gel.

### Proportion of HTLV-I infected cells with stop codon in the *tax* gene in asymptomatic carriers at different time points

RFLP analysis showed that there are AC and HAM/TSP patients that have a stop codon in the *tax* gene as a minor subpopulation of HTLV-I infected cells. To test whether such HTLV-I infected cells with a premature stop codon in the *tax* gene that can escape from the host immune response to HTLV-I can expand subsequently as a major population, we carried out RFLP analysis at different time points in 3 asymptomatic carriers. Case 1 was examined with samples taken on June 25 1999 and June 26 2000. Case 2 was examined with samples taken on Jan 14 2000 and Jan 26 2001. Case 3 was examined with samples taken on May 26 2000, Jan 25 2002 and Oct. 22 2004. RFLP analysis suggested that


in Case 1, the proportion of HTLV-I infected cells with the premature stop codon decreased after 1 year. In this case, to quantify the ratio of HTLV-I infected cells with the premature stop codon to HTLV-I infected cells without this premature stop codon, the nested PCR product was cloned into pCR-Blunt II-TOPO vector (Zero Blunt TOPO PCR cloning kit: Invitrogen), transformed into competent *E. coli* cells and spread on LB plates containing 50  $\mu$ g/ml kanamycin. Colonies from the plate were cultured overnight in LB medium containing 50  $\mu$ g/ml kanamycin, and plasmids containing subcloned *tax* genes were extracted. Purified plasmids containing the subcloned *tax* gene were digested with Eco RI and Bln I and then electrophoresed on 1% agarose gel. When the subcloned *tax* gene was cleaved by Bln I, the subclone was judged as having the stop codon, and if uncleaved, the subclone was judged as not having the stop codon. The proportion of HTLV-I cells that carry the stop codon was then calculated.

### Southern blot analysis of p16<sup>INK4A</sup>, p15<sup>INK4B</sup>, p18<sup>INK4C</sup> and p19<sup>INK4D</sup> and HTLV-I

Southern blot analysis of p16<sup>INK4A</sup>, p15<sup>INK4B</sup>, p18<sup>INK4C</sup> and p19<sup>INK4D</sup> was carried out in 23 cases with acute type ATL. Southern blot analysis of HTLV-I was also carried out. High molecular weight DNA was extracted by a standard method using phenol extraction. In Southern blot analysis for cyclin-dependent kinase inhibitor genes (p16<sup>INK4A</sup>, p15<sup>INK4B</sup>, p18<sup>INK4C</sup>, p19<sup>INK4D</sup>), 10  $\mu$ g of genomic DNA was digested with Hind III, separated on a 1% agarose gel, and transferred to a nylon membrane. Probes used in hybridization were a EcoRI–XhoI fragment of p16<sup>INK4A</sup> cDNA, EcoRI–XhoI fragment of p15<sup>INK4B</sup> cDNA, BamHI–BamHI fragment of p18<sup>INK4C</sup> cDNA and EcoRI–EcoRI fragment of p19<sup>INK4D</sup> cDNA. All of these probes were provided from Dr. Hirai (Banyu Tsukuba Research Institute). The same filters were rehybridized successively with the respective probes. Nylon membranes were also hybridized with  $\beta$ -globin probe. Probe DNA fragments were labeled with  $\alpha$ -<sup>32</sup>P-dCTP by random priming. Blots were hybridized at 65°C for 12 hr in a mixture containing 4 $\times$  SSC (1 $\times$  SSC; 0.15 M NaCl, 0.015 M sodium citrate) and 50  $\mu$ g of sonicated and denatured salmon sperm DNA and then washed in 0.1% sodium dodecylsulfate (SDS) and 1 $\times$  SSC at 65°C for 30 min, and autoradiographed, then exposed to a imaging plate and analyzed by a laser image analyzer (MAC-BAS-1000). Southern blot analysis of HTLV-I was also done in our previous study<sup>17</sup> with 10  $\mu$ g of genomic DNA digested with Pst I and hybridized with total sequence of HTLV-I as a probe. The same filters were rehybridized with a <sup>32</sup>P-labeled HTLV-I long terminal repeat (LTR) probe.

### Sequence of p53 gene

The sequence of p53 was examined in 22 ATL cases. Three p53 fragments were amplified using nested PCR: (i) 371 bp encompassing the entire exon 4; (ii) 499 bp encompassing the entire exons 5 and 6; and (iii) 692 bp encompassing the entire exons 7 and 8. The primers used for PCR encompassing exon 4 were sense 5'-AACGTTCTGGTAAGGACAAGGG-3' (p53\_41) and antisense 5'-AAGGGTGAAGAGGAATCCAAA-3' (p53\_42) for the first PCR and sense 5'-AGGACCTGGTCTCTGACTG-3' (p53\_43) and antisense 5'-ATACGGCCAGGCATTGAAGT-3' (p53\_44) for the second PCR. The primers used for PCR encompassing exons 5 and 6 were sense 5'-TAGTGGGTGCAG-GAGGTGCTT-3' (p53\_51) and antisense 5'-GCAGGAGAA-AGCCCCCTACTG-3' (p53\_62) for the first PCR and sense 5'-TATCTGTTCACCTGTGCCCT-3' (p53\_53) and antisense 5'-GGCCACTGACAACCACCCT-3' (p53\_64) for the second PCR. The primers used for PCR encompassing exons 7 and 8 were sense 5'-GACAGAGCGAGATTCCATCTCA-3' (p53\_71) and antisense 5'-GCTGGTGTGTGGGCACTGCT-3' (p53\_82) for the first PCR and sense 5'-AGGTCTCCCCAA-GGCGCACTGG-3' (p53\_73) and antisense 5'-GGCATAACTGCACCCCTGGTCT-3' (p53\_84) for the second PCR. One hundred nanogram DNA was amplified by 35 cycles for the first PCR using the Expand



**FIGURE 1** – RFLP analysis of *tax* gene. Nested PCR products of *tax* was digested by restriction enzyme Bln I. PCR product in ATL case with a premature stop codon at nt. 7464 in HTLV-I genome was completely cut by Bln I (lane 1), whereas in ATL case without the stop codon at nt. 7464 was not cut by Bln I (lane 2). PCR products in HAM case (lane 3) and in AC case (lane 5) was partially cut by Bln I, suggesting existence of a minor population of HTLV-I infected cells with the premature stop codon at nt. 7464 in HTLV-I *tax* genome, whereas in other HAM case (lane 4) and in AC case (lane 6), PCR products were not cut by Bln I. Long arrow indicates the nested PCR product and short arrow indicates the band cut by Bln I. M, 100 base marker.

high-fidelity PCR system (Boehringer Mannheim, Tokyo, Japan) and 1  $\mu$ M of each primers. After the first PCR, 1  $\mu$ l of aliquots of the amplified products were subjected to an additional 20 cycles of the second PCR using internal primers. Each PCR cycle consisted of denaturation at 95°C for 60 sec, annealing at 60°C for 75 sec, extension at 72°C for 120 sec and extension of the final cycle at 72°C for 10 min. Amplified DNA products were purified using QIA quick purification kit (Qiagen, Tokyo, Japan) and 0.1  $\mu$ g of PCR products were sequenced using dye terminator DNA sequencing kit (Applied Biosystems, Tokyo, Japan) with 3.2 pmol of each primers (p53\_43 and p53\_44 for exon 4, p53\_53 and p53\_64 for exons 5 and 6, p53\_73 and p53\_84 for exons 7 and 8) in an automatic sequencer (377 DNA Sequencer, Applied Biosystems).

## Results


### Premature stop codon in the *tax* gene in AC and in HAM/TSP patients

In 219 asymptomatic carriers and in 143 HAM/TSP patients, there was no case that had a premature stop codon in the *tax* gene in the commonest sequence of the individual. There are cases, however, who had HTLV-I infected cells with this premature stop codon in the *tax* gene as a minor population of the HTLV-I infected cells. Figure 1 shows representative results. In the ATL case with a premature stop codon in the HTLV-I *tax* gene, all of the nested PCR product was cut by Bln I (Fig. 1, lane 1). Under the same experimental conditions, there were no AC or HAM/TSP patients whose nested PCR products were completely cut by Bln I, but there was partial cleavage in some cases (Fig. 1., lanes 3,5).

**TABLE I** – HTLV-I PROVIRUS LOAD IN AC AND IN HAM/TSP IN ASSOCIATION WITH OR WITHOUT THE PREMATURE STOP CODON IN *TAX* GENE AS MINOR POPULATION OF HTLV-I INFECTED CELLS<sup>1</sup>

	AC (n = 219)		HAM (n = 143)	
	+	-	+	-
n	79	140	78	65
Median	166*	34*	523	420

<sup>1</sup>HTLV-I copy number per 10<sup>-4</sup> PBMC was represented. n = number of subjects. + Subjects with the premature stop codon in *tax* gene as minor population of HTLV-I infected cells detected by RFLP analysis - Subjects without premature stop codon in *tax* gene; HAM, patients with HAM; AC, asymptomatic carriers. \*p-value < 0.001 by Mann-Whitney U-test.



**FIGURE 2** – RFLP analysis of *tax* gene at different time points. RFLP analysis of nested PCR product of *tax* gene digested by Bln I was done in 3 asymptomatic carriers at different time points. Case 1 (lane 1 at June 25 1999 and lane 2 at June 26 2000). Case 2 (lane 3 at Jan 14 2000 and lane 4 at Jan 26 2001). Case 3 (lane 5 at May 26 2000, lane 6 at Jan 25 2002 and lane 7 at Oct. 22 2004). Long arrow indicates the nested PCR product and short arrow indicates the band cut by Bln I. M, 100 base marker.

There were 79 cases of 219 AC (36.1%) and 78 cases of 143 HAM/TSP patients (54.5%) that had HTLV-I infected cells with the premature stop codon in the *tax* gene as a minor population of the individuals (Table I).

The median provirus load in AC who had the premature stop codon in the *tax* gene as a minor population of HTLV-I infected cells was 166 and the median provirus load in AC who did not have this premature stop codon in the HTLV-I infected cells was 34.5, and this difference was significant ( $p < 0.001$ ). The median provirus load in HAM patients who had the premature stop codon in *tax* gene as a minor population of HTLV-I infected cells was 523 and the median provirus load in HAM patients who did not have this premature stop codon in HTLV-I infected cells was 420, and this difference was not significant ( $p = 0.305$ ) (Table I).

### Proportion of HTLV-I infected cells with a stop codon in the *tax* gene as a minor population in asymptomatic carriers at different time points

In 3 asymptomatic carriers having the premature stop codon in the *tax* gene as a minor population of HTLV-I infected cells, we examined whether this minor population expanded subsequently.




FIGURE 3 – Southern blot analysis of  $p15^{INK4B}$ ,  $p16^{INK4A}$ ,  $p18^{INK4C}$ ,  $p19^{INK4D}$  and  $\beta$ -globin in 23 acute ATL. Ten micrograms of genomic DNA was digested with Hind III and hybridized with each probe. Cases 1, 7–12, 14 and 22 has weak or decreased  $p16^{INK4A}$  band when compared to the density of  $\beta$ -globin band. Cases 1, 8–10, 12, 14, 18 and 22 has weak or decreased  $p15^{INK4B}$  band and Case 7 has a shorter  $p15^{INK4B}$  band. Abbreviations: p15,  $p15^{INK4B}$ ; p16,  $p16^{INK4A}$ ; p18,  $p18^{INK4C}$ ; p19,  $p19^{INK4D}$ .

Figure 2 shows that this minor population with the premature stop codon in the *tax* gene in these 3 asymptomatic carriers did not expand subsequently (Case 1, lane 1,2; Case 2, lane 3,4; Case 3, lanes 5–7). In Case 1, the proportion of the population with premature stop codon decreased subsequently when analyzed by RFLP (Fig. 2, lanes 1,2). To quantify this, we subcloned the PCR product and counted the number of clones that had a stop codon at different time points. The number of subclones that had a premature stop codon in the *tax* gene were 9 of 59 (15.3%) on June 25 1999, and was 2 of 63 (3.2%) on June 26 2000.

#### Deletion of cyclin-dependent kinase 4 inhibitor genes and mutations in HTLV-I provirus in acute type ATL patients

Judged from the density of the  $p16^{INK4A}$  gene band in Southern blot hybridization compared to the density of the band of  $\beta$ -globin using an image analyzer, 9 of 23 (39.1%) acute-type ATL patients deleted the  $p16^{INK4A}$  gene in leukemic cells (Fig. 3). Similarly, an absent or weak hybridization band indicating deletion of the  $p15^{INK4B}$  gene was observed in 8 of 23 acute type ATL patients and a shorter size of the  $p15^{INK4B}$  gene band indicating partial deletion or rearrangement of this gene was observed in one patient. No genetic alteration was detected by Southern blot analysis in  $p18^{INK4C}$  and in  $p19^{INK4D}$ .

Case 9 had the stop codon in the *tax* gene but did not have deletion in the HTLV-I provirus genome by Southern blotting (Fig. 4, lane 1A,1B), and  $p15^{INK4B}$  and  $p16^{INK4A}$  were deleted (Table II). Case 11 had only one LTR band by Southern blotting (data not shown) and  $p16^{INK4A}$  was deleted. Case 22 had a large deletion in HTLV-I genome suggested by the Southern blotting (Fig. 4, lanes 2A,2B), and  $p15^{INK4B}$  and  $p16^{INK4A}$  were deleted.

#### Mutations in p53 gene

Sequencing of p53 was carried out from codons 33–307 (exons 4–8) in 22 acute ATL cases. Mutations were found in 3 cases. All of them were homozygous missense mutations. Case 13 had a mutation at codon 285 (Glu to Lys) in exon 8. This case had only one LTR band by Southern blot analysis of HTLV-I (Fig. 4, lanes 3A,3B). In this case (Case 13), 1.8 kb band was absent when hybridized with a total HTLV-I probe, suggesting a large deletion encompassing the 5' LTR through the first Pst I restriction site. Case 17 had a mutation at codon 266 (Gly to Arg) in exon 8 and had a deletion in HTLV-I genome by Southern blot analysis of HTLV-I. Case 19 had a mutation at codon 193 (His to Leu) in exon 6. There were also ATL cases that had deletion in HTLV-I provirus genome but did not have deletion in INK4 genes and did not have mutation in p53 gene. Case 15 represent one such ATL case. Case 15 had only one LTR band when hybridized with a LTR probe (Fig. 4, lane 4B) and there was a larger size of band instead of 1.8 kb band when hybridized with a total HTLV-I probe (Fig. 4, lane 4A), suggesting a deletion encompassing the 5' LTR through the first Pst I restriction site. The deletions and mutations observed in the HTLV-I provirus genome, INK genes and p53




FIGURE 4 – Southern blot analysis of the HTLV-I provirus genome. Ten micrograms cellular DNA was digested with Pst I and subjected to standard Southern blot analysis. The filter was hybridized with a total HTLV-I probe (a) and then with a LTR probe (b). Arrowheads show the viral-cellular junction bands with LTR probe, in addition to 3 internal bands (2.5kb, 1.8kb, 1.2kb). Schematic illustration of the HTLV-I genome, restriction map, and probes are shown on the upper column. (↓) Pst I site. Lane 1 (Case 9 in Table II): Typical patient with ATL showing 3 major internal bands (2.5kb, 1.8kb, 1.2kb) with additional 2 viral-cellular junction bands (arrow heads). Lane 2 (Case 22 in Table II): Three major internal bands are absent (a), but 2 LTR bands are observed, suggesting a large deletion in HTLV-I proviral genome. Lane 3 (Case 13 in Table II): 1.8kb band is absent when hybridized with a total HTLV-I probe (a) and there is only one LTR band hybridized with a LTR probe (b), suggesting a large deletion encompassing the 5' LTR through the first Pst I restriction site. Lane 4 (Case 15 in Table II): Another example of ATL who have only one LTR band hybridized with a LTR probe (b) and 1.8kb band is absent hybridized with a total HTLV-I probe (a).

gene are summarized in Table II. There were 9 cases of acute ATL who had a putative CTL escape mutation in the HTLV-I provirus genome and 4 of these 9 patients had either a deletion in  $INK4A$  or  $INK4B$  gene or p53 gene.

TABLE II - MUTATION/DELETION IN HTLV-I GENOME, p15, p16 AND p53 GENES IN ACUTE ATL

Case no.	Southern blot/mutation analysis	Southern blot analysis		Mutation analysis		
	HTLV-I	p15	p16	codon	p53 nucleotide substitution	amino acid
1		Del	Del			
2						
3						
4						
5	G7464A <sup>1</sup>					
6						
7		Del	Del			
8		Del	Del			
9	G7464A <sup>1</sup>	Del	Del			
10		Del	Del			
11	1 LTR <sup>3</sup>		Del			
12		Del	Del			
13	1 LTR <sup>3</sup>			285	GAG-AAG	Glu-Lys
14		Del	Del			
15	1 LTR <sup>3</sup>					
16						
17	PD			266	GGA-AGA	Gly-Arg
18		Del				
19				193	CAT-CTT	His-Leu
20						
21	A7337G <sup>2</sup>			not examined		
22	PD	Del	Del			
23	PD					

<sup>1</sup>G7464A: substitution at nt. position 7464 from G-A that creates a premature stop codon in *tax*. <sup>2</sup>A7337G: substitution at nt. position 7337 from A-G that causes an amino acid change from Gly - Arg that is an putative escape mutation. <sup>3</sup>1LTR Cases that show only one LTR band by Southern blot analysis using HTLV-I LTR as a probe. PD, partial deletion; Del, deletion.

## Discussion

The purpose of our present study was to examine when the premature stop codon in the *tax* gene, a putative escape mutation from the anti-HTLV-I CTL response can emerge in the proviral population, and to examine whether such HTLV-I infected cells with escape mutation will subsequently proliferate even in an asymptomatic carrier. We found that the premature stop codon in the *tax* gene exist frequently in HTLV-I carriers as a minor population of the individual carriers; however, such a minor population did not expand subsequently, and deletions in certain host genes still favored the development of ATL.

In our study, we focused on the mutation in *tax* gene especially the premature stop codon of *tax*. Although we could not detect this premature stop codon in the *tax* gene in the majority of ATL patients, there were 4 cases among 55 ATL patients as we reported previously<sup>17</sup> and 4 cases among 47 ATL patients were reported in another study.<sup>33</sup> In addition to this premature stop codon, deletions in the *tax* gene,<sup>17</sup> deletion of 5'-LTR<sup>33</sup> that is a promoter of viral genes, and silencing of the *tax* gene<sup>34</sup> have been reported previously in ATL, suggesting that ATL cells that do not express HTLV-I-encoded proteins are selected by the cytotoxic T cell (CTL) response to HTLV-I, during the development of ATL. A premature stop codon in the *tax* gene is one such escape mutation that is observed in some ATL patient as the commonest sequence in the individual.<sup>17</sup> It was clear that this premature stop codon in the ATL cell emerged after the viral transmission, because the HTLV-I *tax* sequence in the family member was identical with other nucleotide alterations specific for this family except this premature stop codon.<sup>17</sup> It was not clear, however, if this premature stop codon emerged in the early stage of ATL development, or whether the mutation emerged after ATL had developed. To infer at which stage the premature stop codon could have emerged, and to investigate whether there is any HTLV-I carrier with this premature stop codon in the *tax* gene in the commonest proviral sequence of the individual, we examined this premature stop codon in the *tax* gene in 219 AC and in 143 HAM/TSP patients. Although the premature stop codon in the *tax* gene did not exist as

the commonest sequence in any AC or HAM/TSP patients, a stop codon in the *tax* gene was observed as a minor population of the HTLV-I infected cells in many AC and in HAM/TSP patients. This observation was consistent with our previous study by direct sequencing in which we found no AC or HAM/TSP patients that had this premature stop codon as the commonest sequence of the individual.<sup>17</sup> The observation is also consistent with our previous study of a small number of AC and HAM/TSP patients that the premature stop codon in the *tax* gene is present in a minor population in some of the HTLV-I carriers.<sup>35</sup> In our present study, we found that many AC and HAM/TSP patients carried the premature stop codon in the *tax* gene in a minor population of the infected cells. We also reported previously a chronic type of ATL with this premature stop codon as the commonest sequence of the individual.<sup>17</sup> These findings suggest that this premature stop codon emerged in the early stages of ATL development rather than as a consequence of genetic instability after the progression to an aggressive form of ATL. The median HTLV-I provirus load was significantly higher in AC who had a minor population of HTLV-I infected cells with this premature stop codon in the *tax* gene compared to AC who did not have the premature stop codon in the *tax* gene as a minor population, but was not different in HAM patients regardless of the presence of premature stop codon in the *tax* gene in HTLV-I infected cells. We do not know why this difference happens, but it is possible to speculate that the proportion of the role of viral transcription vs. cell division in maintaining the provirus load is different among AC and HAM because, the HTLV-I sequence mutation are caused frequently by the reverse transcriptase, but caused rarely by the host DNA polymerase.

Although the RFLP analysis of the premature stop codon in the *tax* gene was not quantitative, there were HTLV-I carriers with a significant proportion of HTLV-I infected cells with this premature stop codon, as judged from the density of the band cut by the Bln I restriction enzyme. To investigate whether such HTLV-I infected cells were already in the process of developing ATL, we followed 3 asymptomatic carriers who had this premature stop codon in the *tax* gene as minor population (Fig. 2). In these 3 carriers, the population with the premature stop codon in the *tax* gene

did not expand, although the observation time was 1 year in each of 2 cases and 4 years in one case. Interestingly, in one case, in which we quantified the proportion of HTLV-I infected cells with the premature stop codon in the *tax* gene, the proportion of HTLV-I infected cells with this premature stop codon decreased after one year. The provirus load was 64 when the proportion of HTLV-I infected cells had the premature stop codon in *tax* gene was 15.3%, but the provirus load was 5 after a year when the proportion of HTLV-I infected cells with the premature stop codon in *tax* gene decreased to 3.2%. These findings suggest that, although HTLV-I infected cells with escape mutation can accumulate in non-ATL HTLV-I carriers, HTLV-I infected cells with a mutation that abolishes the function of Tax lose their proliferative advantage and progressively disappear. It is likely that an escape mutation that abolishes the function of Tax should appear after the immortalization of HTLV-I infected cells for the subsequent development of ATL.

Next, we examined whether host genetic changes promote the malignant transformation of HTLV-I infected cells even in ATL cells that can escape from the host immune response to HTLV-I. To this end, we examined 4 known cyclin dependent kinase inhibitors (INK) and *p53* gene, in combination with the alterations in the HTLV-I genome. In the case of *p16<sup>INK4A</sup>* and *p15<sup>INK4B</sup>* genes, there is a report that these genes are deleted in many ATL patients,<sup>36</sup> whereas *p18<sup>INK4C</sup>* and *p19<sup>INK4D</sup>* are not deleted.<sup>37,38</sup> This is consistent with our finding that 9 of 23 (39.1%) acute-phase ATL samples had a deletion in *p16<sup>INK4A</sup>* gene and 8 of these patients (36%) who deleted *p16<sup>INK4A</sup>* also had deletion or rearrangement of *p15<sup>INK4B</sup>*, whereas *p18<sup>INK4C</sup>* and *p19<sup>INK4D</sup>* were not deleted. In our ATL cases that deleted *p16<sup>INK4A</sup>* or *p15<sup>INK4B</sup>*, there were 3 cases that could potentially escape from the host CTL response to HTLV-I. One had a premature stop codon in the *tax* gene, one other case had only one LTR region, and the other had a large deletion in the HTLV-I genome. Each of these genomic alterations could lead to escape from the host immune response to HTLV-I. In the case of *p53* gene, we sequenced the entire exons 4–8, where are highly conserved regions of the *p53* gene,<sup>40</sup> and

are also identified as hot spots for mutations in several malignancies including ATL.<sup>30</sup> The missense mutations of *p53* gene observed in our study were at codons 193, 266 and 285 where mutation had been reported in other malignancies (anonymous ftp to ftp.ebi.ac.uk, in the directory/pub/databases/p53) and 2 of them had alterations in the HTLV-I provirus genome. These deletions in the HTLV-I genome could also lead to escape from the host immune response to HTLV-I. Regarding the host genetic changes in the development of ATL, Yamada *et al.*<sup>39</sup> reported that 3 chronic ATL cases who progressed to acute type lost the *p16<sup>INK4A</sup>* gene alone or the *p15<sup>INK4B</sup>* and *p16<sup>INK4A</sup>* genes at their exacerbation phase.<sup>39</sup> Hatta *et al.* reported an ATL patient with a homozygously deleted *p16<sup>INK4A</sup>* gene in the chronic phase who rapidly progressed to acute ATL and died within 6 months of the initial diagnosis.<sup>36</sup> There was also a case reported in which the *p53* gene was intact at chronic stage but was mutated when the disease progressed to acute type ATL.<sup>30</sup> These reports suggest that deletion/mutation of tumor suppressor genes such as *p16<sup>INK4A</sup>* gene and *p53* in ATL are not a result of genetic instability after the exacerbation of ATL, but are factors that predict poor prognosis. We speculate that in the early stage of HTLV-I infection, Tax gives advantage for proliferation of HTLV-I infected cells, but those cells that continuously express HTLV-I viral proteins are likely to be rejected by host immune system. If an escape mutation emerges after the immortalization of the HTLV-I infected cells, cells that carry such mutations are selected by the host immune system and when additional host genetic changes have accumulated, such as in the *p16<sup>INK4A</sup>* or *p53* gene, HTLV-I infected cells will progress further to an aggressive form of ATL.

#### Acknowledgements

We thank Ms. T. Muramoto and Ms. Y. Nishino (Department of Neurology and Geriatrics, Kagoshima University, Japan) for their excellent technical assistance. We also thank Professor C.R.M. Bangham (Immunology Department, Imperial College Faculty of Medicine, UK) for critical reading of the manuscript.

#### References

- Uchiyama T, Yodoi J, Sagawa K, Takatsuki K, Uchino H. Adult T cell leukemia: clinical and hematologic features of 16 cases. *Blood* 1977;50:481–92.
- Yoshida M, Seiki M, Yamaguchi K, Takatsuki K. Monoclonal integration of human T cell leukemia provirus in all primary tumors of adult T cell leukemia suggests causative role of human T cell leukemia virus in the disease. *Proc Natl Acad Sci USA* 1984; 81:2534–7.
- Gessain A, Barin F, Vernant JC, Gout O, Maurs L, Calender A, de Thé G. Antibodies to human T-lymphotropic virus type-I in patients with tropical spastic paraparesis. *Lancet* 1985;2:407–10.
- Osame M, Usuku K, Izumo S, Ijichi N, Amitani H, Igata A, Matsumoto M, Tara M. HTLV-I associated myelopathy, a new clinical entity. *Lancet* 1986;1:1031–2.
- Shimoyama M. Diagnostic criteria and classification of clinical subtypes of adult T cell leukaemia-lymphoma. A report from the Lymphoma Study Group (1984–87). *Br J Haematol* 1991;79:428–37.
- Okamoto T, Ohno Y, Tsugane S, Watanabe S, Shimoyama M, Tajima K, Miwa M, Shimotohno K. Multi-step carcinogenesis model for adult T cell leukemia. *Jpn J Cancer Res* 1989;80:191–5.
- Suzuki T, Uchida-Toita M, Yoshida M. Tax protein of HTLV-I inhibits CBP/p300-mediated transcription by interfering with recruitment of CBP/p300 onto DNA element of E-box or p53 binding site. *Oncogene* 1999;18:4137–43.
- Giebler HA, Loring JE, van Orden K, Colgin MA, Garrus JE, Escudero KW, Brauweiler A, Nyborg JK. Anchoring of CREB binding protein to the human T cell leukemia virus type 1 promoter: a molecular mechanism of Tax transactivation. *Mol Cell Biol* 1997; 17:5156–64.
- Siekvitz M, Feinberg MB, Holbrook N, Wong-Staal F, Greene WC. Activation of interleukin 2 and interleukin 2 receptor (Tac) promoter expression by the trans-activator (tat) gene product of human T cell leukemia virus, type I. *Proc Natl Acad Sci USA* 1987;84:5389–93.
- Suzuki T, Narita T, Uchida-Toita M, Yoshida M. Down-regulation of the INK4 family of cyclin-dependent kinase inhibitors by tax protein of HTLV-I through two distinct mechanisms. *Virology* 1999; 259:384–91.
- Brauweiler A, Garrus JE, Reed JC, Nyborg JK. Repression of bax gene expression by the HTLV-1 Tax protein: implications for suppression of apoptosis in virally infected cells. *Virology* 1997;231: 135–40.
- Suzuki T, Kitao S, Matsushime H, Yoshida M. HTLV-1 Tax protein interacts with cyclin-dependent kinase inhibitor p16INK4A and counteracts its inhibitory activity towards CDK4. *EMBO J* 1996;15:1607–14.
- Suzuki T, Ohsugi Y, Uchida-Toita M, Akiyama T, Yoshida M. Tax oncoprotein of HTLV-1 binds to the human homologue of Drosophila discs large tumor suppressor protein, hDLG, and perturbs its function in cell growth control. *Oncogene* 1999;18:5967–77.
- Jin DY, Spencer F, Jeang KT. Human T cell leukemia virus type 1 oncoprotein Tax targets the human mitotic checkpoint protein MAD1. *Cell* 1998;93:81–91.
- Kannagi M, Shida H, Igarashi H, Kuruma K, Murai H, Aono Y, Maruyama I, Osame M, Hattori T, Inoko H. Target epitope in the Tax protein of human T cell leukemia virus type I recognized by class I major histocompatibility complex-restricted cytotoxic T cells. *J Virol* 1992; 66:2928–33.
- Goon PK, Biancardi A, Fast N, Igakura T, Hanon E, Mosley AJ, Asquith B, Gould KG, Marshall S, Taylor GP, Bangham CR. Human T cell lymphotropic virus (HTLV-I) type-1-specific CD8+ T cells: frequency and immunodominance hierarchy. *J Infect Dis* 2004;189:2294–8.
- Furukawa Y, Kubota R, Tara M, Izumo S, Osame M. Existence of escape mutant in HTLV-I tax during the development of adult T cell leukemia. *Blood* 2001;97:987–93.
- Seiki M, Inoue J, Takeda T, Yoshida M. Direct evidence that p40x of human T cell leukemia virus type I is a trans-acting transcriptional activator. *EMBO J* 1986;5:561–5.
- Sherr CJ. Mammalian G1 cyclins. *Cell* 1993;73:1059–65.
- Sherr CJ, Roberts JM. CDK inhibitors: positive and negative regulators of G1-phase progression. *Genes Dev* 1999;13:1501–12.
- Hannon GJ, Beach D. p15INK4B is a potential effector of TGF-beta-induced cell cycle arrest. *Nature* 1994;371:257–61.

22. Serrano M, Hammon GJ, Beach D. A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* 1993;366:704-7.
23. Guan KL, Jenkins CW, Li Y, Nichols MA, Wu X, O'Keefe CL, Matera AG, Xiong Y. Growth suppression by p18, a p16INK4/MTS1- and p14INK4B/MTS2-related CDK6 inhibitor, correlates with wild-type pRb function. *Genes Dev* 1994;8:2939-52.
24. Hirai H, Roussel MF, Kato JY, Ashmun RA, Sherr CJ. Novel INK4 proteins, p19 and p18, are specific inhibitors of the cyclin D-dependent kinases CDK4 and CDK6. *Mol Cell Biol* 1995;15:2672-81.
25. Chan FK, Zhang J, Cheng L, Shapiro DN, Winoto A. Identification of human and mouse p19, a novel CDK4 and CDK6 inhibitor with homology to p16ink4. *Mol Cell Biol* 1995;15:2682-8.
26. Kamb A, Gruis NA, Weaver-Feldhaus J, Liu Q, Harshman K, Tavitgian SV, Stockert E, Day RS, Johnson BE, Skolnick MH. A cell cycle regulator potentially involved in genesis of many tumor types. *Science* 1994;264:436-40.
27. Ruas M, Peters G. The p16INK4a/CDKN2A tumor suppressor and its relatives. *Biochim Biophys Acta*. 1998;1378:115-77.
28. Stott FJ, Bates S, James MC, McConnell BB, Starborg M, Brookes S, Palmero I, Ryan K, Hara E, Vousden KH. The alternative product from the human CDKN2A locus, p14(ARF), participates in a regulatory feedback loop with p53 and MDM2. *EMBO* 1998;17:5001-14.
29. Michalovitz D, Halevy O, Oren M. p53 mutations: Gains or losses? *J Cell Biochem* 1991;45:22-9.
30. Sakashita A, Hattori T, Miller CW, Suzushima H, Asou N, Takatsuki K, Koeffler HP. Mutations of the p53 gene in adult T cell leukemia. *Blood* 1992;79:477-80.
31. Osame M. HTLV. In: Blattner E, ed. *Human retrovirology*. New York: Raven Press, 1990. p 191-7.
32. Nagai M, Usuku K, Matsumoto W, Kodama D, Takenouchi N, Moritoyo T, Hashiguchi S, Ichinose M, Bangham CR, Izumo S, Osame M. Analysis of HTLV-I proviral load in 202 HAM/TSP patients and 243 asymptomatic HTLV-I carriers: high proviral load strongly predisposes to HAM/TSP. *J Neurovirol* 1998;4:586-93.
33. Tamiya S, Matsuoka M, Etoh K, Watanabe T, Kamihira S, Yamaguchi K, Takatsuki K. Two types of defective human T-lymphotropic virus type I provirus in adult T cell leukemia. *Blood* 1996;88:3065-72.
34. Takeda S, Maeda M, Morikawa S, Taniguchi Y, Yasunaga J, Nosaka K, Tanaka Y, Matsuoka M. Genetic and epigenetic inactivation of tax gene in adult T cell leukemia cells. *Int J Cancer* 2004;109:559-67.
35. Saito M, Furukawa Y, Kubota R, Usuku K, Sonoda S, Izumo S, Osame M, Yoshida M. Frequent mutation in pX region of HTLV-1 is observed in HAM/TSP patients, but is not specifically associated with the central nervous system lesions. *J Neurovirol* 1995;1:286-94.
36. Hatta Y, Hiramata T, Miller CW, Yamada Y, Tomonaga M, Koeffler HP. Homozygous deletions of the p15 (MTS2) and p16 (CDKN2/MTS1) genes in adult T cell leukemia. *Blood* 1995;85:2699-704.
37. Hatta Y, Spirin K, Tasaka T, Morosetti R, Said JW, Yamada Y, Tomonaga M, Koeffler HP. Analysis of p18INK4C in adult T cell leukaemia and non-Hodgkin's lymphoma. *Br J Haematol* 1997;99:665-7.
38. Shiohara M, Spirin K, Said JW, Gombart AF, Nakamaki T, Takeuchi S, Hatta Y, Morosetti R, Tasaka T, Seriu T, Bartram C, Miller CW, et al. Alterations of the cyclin-dependent kinase inhibitor p19 (INK4D) is rare in hematopoietic malignancies. *Leukemia* 1996;10:1897-900.
39. Yamada Y, Hatta Y, Murata K, Sugawara K, Ikeda S, Mine M, Maeda T, Hirakata Y, Kamihira S, Tsukasaki K, Ogawa S, Hirai H, et al. Deletions of p15 and/or p16 genes as a poor-prognosis factor in adult T cell leukemia. *J Clin Oncol* 1997;15:1778-85.
40. Hollstein M, Shomer B, Greeblatt M, Soussi T, Hovig E, Montesano R, Harris CC. Somatic point mutations in the p53 gene of human tumor and cell lines: updated compilation. *Nucleic Acids Res* 1996;24:141-6.

## A prospective uncontrolled trial of fermented milk drink containing viable *Lactobacillus casei* strain Shirota in the treatment of HTLV-1 associated myelopathy/tropical spastic paraparesis

Toshio Matsuzaki<sup>a</sup>, Mineki Saito<sup>a,\*</sup>, Koichiro Usuku<sup>b</sup>, Hirohisa Nose<sup>a</sup>, Shuji Izumo<sup>c</sup>,  
Kimiyoishi Arimura<sup>a</sup>, Mitsuhiro Osame<sup>a</sup>

<sup>a</sup>Department of Neurology and Geriatrics, Kagoshima University Graduate School of Medical and Dental Sciences,  
8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan

<sup>b</sup>Department of Medical Information Science, Kagoshima University Graduate School of Medical and Dental Sciences,  
8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan

<sup>c</sup>Department of Molecular Pathology, Center for Chronic Viral Diseases, Kagoshima University,  
8-35-1 Sakuragaoka, Kagoshima 890-8520, Japan

Received 30 November 2004; received in revised form 20 May 2005; accepted 23 May 2005  
Available online 14 June 2005

### Abstract

Ten patients with human T-cell lymphotropic virus type-1 (HTLV-1)-associated myelopathy/tropical spastic paraparesis (HAM/TSP) were treated in an uncontrolled preliminary trial by oral administration of viable *Lactobacillus casei* strain Shirota (LcS) containing fermented milk. HTLV-1 provirus load, motor function, neurological findings, and immunological parameters were evaluated after 4 weeks. Although LcS did not change the frequencies or absolute numbers of all the examined cell surface phenotypes of peripheral blood mononuclear cells, NK cell activity was significantly increased after 4 weeks of oral administration of LcS preparation. Improvements in spasticity (modified Ashworth Scale scores) and urinary symptoms were also seen after LcS treatment. No adverse effect was observed in all the 10 patients throughout the study period. Our results indicated that LcS may be a safe and beneficial agent for the treatment of HAM/TSP; therefore randomized controlled studies are warranted.

© 2005 Elsevier B.V. All rights reserved.

**Keywords:** Immunomodulation; *Lactobacillus casei* strain Shirota; NK cell activity; HAM/TSP; HTLV-1

### 1. Introduction

Human T-cell lymphotropic virus type 1 (HTLV-1)-associated myelopathy/tropical spastic paraparesis (HAM/TSP) is a chronic progressive myelopathy characterized by spastic paraparesis, sphincter dysfunction, and mild sensory disturbance in the lower extremities [1–3]. Although the precise mechanism causing HAM/TSP is still obscure, virus–host immunological interactions are considered to be the most important cause of this disease, because (1) the

median HTLV-1 provirus load is more than 10 times higher in HAM/TSP patients than in healthy HTLV-1 carriers (HCs), and high provirus load is also associated with an increased risk of progression to disease [4]; (2) the titer of anti-HTLV-1 antibody often reaches a very high level in HAM/TSP patients, and correlates well with a HTLV-1 provirus load [4]; (3) in HAM/TSP patients, large populations of activated T cells both in peripheral blood mononuclear cells (PBMCs) [5] and cerebrospinal fluid (CSF) [6], and spontaneous proliferation of PBMCs in vitro [7–9] have been reported; (4) HTLV-1-specific CD8<sup>+</sup> cytotoxic T lymphocytes (CTL) are abundant and activated in PBMCs in HAM/TSP patients [10], and these CTLs are preferentially accumulated in CSF cells [11,12]; and (5) it

\* Corresponding author. Tel.: +81 99 275 5332; fax: +81 99 265 7164.  
E-mail address: mineki@m3.kufm.kagoshima-u.ac.jp (M. Saito).

has been shown that HTLV-1 Tax<sub>11–19</sub>-specific CD8<sup>+</sup> T cells have the potential to produce proinflammatory cytokines [13], whereas possession of the HLA-A\*02 allele was associated with protection against HAM/TSP as well as a lower provirus load [14]. To regulate such immunologic processes, some therapeutic trials with agents such as prednisolone [15,16], plasma exchange [17], and interferon (IFN)- $\alpha$  [18–22], have been conducted and reported to be effective.

*Lactobacillus casei* strain Shirota (LcS), a type of lactic acid bacteria, was originally isolated from the human intestine and has been used commercially for a long time to produce fermented milk [23]. It was previously reported that LcS activates macrophages [24], NK cells [25], and cytotoxic T cells [26], and also exhibits a marked inhibitory effect against tumors [23,27], type II collagen-induced arthritis in DBA/1 mice [28], and infectious pathogens such as *Listeria monocytogenes* [29], herpes simplex virus [30], and influenza virus [31]. Adverse effects and toxic reactions to LcS are rarely reported. On the other hand, previous reports indicated that the NK cell activity and the percentages of NK cell subsets was significantly decreased in HAM/TSP patient compared with that in uninfected normal controls [32,33]. In our previous study, we also reported that one of the inhibitory NK cell receptor, NKG2A, was significantly decreased in frequency in HAM/TSP patients but not in HCs [34]. Since inhibitory NK cell receptors such as NKG2A can down-regulate antigen-mediated T-cell effector functions including cytotoxic activity and

cytokine release, decreased NK cell function and NKG2A<sup>+</sup> T cells may be associated with impaired regulation of T cell-mediated antiviral immune responses in HAM/TSP patients. These observations therefore raise the possibility that LcS could improve the symptoms of HAM/TSP via activating NK cell subsets.

To conduct a preliminary investigation of the efficacy of LcS for HAM/TSP—and to see if controlled trials are warranted—we evaluated in an uncontrolled trial with 10 HAM/TSP patients. After daily oral administration of LcS preparation that contains approximately  $8 \times 10^{10}$  viable LcS for 4 weeks, the effects of spasticity, motor disabilities, urinary symptoms and immunological parameters were evaluated.

## 2. Patients and methods

### 2.1. Study population

Our study constituted a prospective open evaluation of LcS over a 4-week period. Ten patients with clinical diagnoses of HAM/TSP in accordance with World Health Organization criteria [35] were enrolled in this study. All patients gave informed consent.

### 2.2. Patients and cells

Clinical characteristics of the patients are shown in Table 1. The duration of illness ranged from 7 to 24 years.

Table 1

Summary of demographic and clinical characteristics for 10 patients with human T lymphotropic virus type 1 (HTLV-1)-associated myelopathy/tropical spastic paraparesis (HAM/TSP)

Patient	Age, years	Sex	Disease duration, years	HTLV-1 antibody titer	Provirus load before Tx	Provirus load after Tx <sup>a</sup>	Concomitant Tx <sup>b</sup>	MAS before Tx	MAS after Tx <sup>c</sup>	OMDS before Tx	OMDS after Tx <sup>d</sup>	UDS before Tx	UDS after Tx <sup>e</sup>	Overall evaluation of efficacy <sup>f</sup>
HAM1	34	F	19	×131072	1757	1397	VC 375mg/day	2	1	4	3	2	0	good
HAM2	62	M	14	×32768	634	777	PSL 5mg/day, VC 375mg/day	+1	0	6	6	3	0	good
HAM3	50	F	17	×2048	907	779	None	2	1	5	4	3	0	good
HAM4	45	F	15	×16384	2942	471	PSL 5mg/day, VC 375mg/day	+1	0	5	3	6	3	excellent
HAM5	60	F	7	×8192	204	194	PSL 5mg/day, VC 375mg/day	3	4	4	4	1	0	fair
HAM6	47	M	18	×2048	716	849	PSL 10mg/day, VC 375mg/day	2	1	6	6	5	4	fair
HAM7	46	M	24	×16384	278	361	None	3	1	4	3	2	0	good
HAM8	55	F	10	×65536	2882	524	None	2	1	4	3	6	3	good
HAM9	41	F	17	×65536	1073	1263	None	3	0	2	2	3	1	good
HAM10	57	F	13	×32768	245	387	None	2	1	4	4	2	1	Fair

LcS, *Lactobacillus casei* strain Shirota; OMDS, Osame Motor Disability Score; UDS, urinary disturbance score; MAS, modified Ashworth scale.

<sup>a</sup> At 4 weeks after the first administration of LcS; for the changes in HTLV-1 provirus load,  $p=0.401$ , by the Wilcoxon signed rank test.

<sup>b</sup> VC, vitamin C; PSL, prednisolone.

<sup>c</sup> At 4 weeks after the first administration of LcS; for the change in MAS,  $p=0.015$ , by the Wilcoxon signed rank test.

<sup>d</sup> At 4 weeks after the first administration of LcS; for the change in OMDS,  $p=0.157$ , by the Wilcoxon signed rank test.

<sup>e</sup> At 4 weeks after the first administration of LcS; for the change in UDS,  $p=0.0085$ , by the Wilcoxon signed rank test.

<sup>f</sup> The evaluation was mainly based on improvement in motor dysfunction, urinary disturbances, and neurologic signs, as judged by the attending neurologists (see Patients and methods).



In these patients, motor disability scores rated from 0 to 13 (as described in Table 2), and ranged from 2 to 6 (Table 1). Fresh PBMCs were isolated on a Histopaque-1077 (Sigma) density gradient centrifugation, washed twice in RPMI 1640 with 5% heat-inactivated fetal calf serum (FCS), then stored in liquid nitrogen until use.

### 2.3. Treatment protocol

All patients were given LcS preparation, which contains approximately  $4 \times 10^{10}$  viable LcS, orally twice a day for 4 weeks. The drugs other than LcS such as muscle relaxants or vitamins were continued on the condition that the dosage regimen was kept constant before, during, and after treatment. Four patients (HAM2, HAM4, HAM5 and HAM6) received low-dose (5–10 mg daily) oral prednisolone throughout the study period. None of the patients—with the exception of these four—had received any immunomodulators.

### 2.4. Concomitant therapy

The drugs other than LcS such as vitamin C or low dose (5–10 mg daily) oral *prednisolone* were continued on condition that the dosage regimen was kept constant before, during, and after therapy. Four patients (HAM2, HAM4, HAM5 and HAM6) out of 10 had received low dose oral prednisolone throughout study period. None of the patients except for these four had received any immunomodulators (Table 1).

### 2.5. Clinical evaluation

Assessments were performed on subjective symptoms, degree of spasticity, disability grading, sphincter disturbance and adverse effects before and 1 month after administration of LcS. The clinical findings of each

patient were assessed by the blinded consultant neurologists, i.e. the examining neurologist for post-clinical scores was blinded to the pre-clinical scores. Spasticity of the lower limbs was graded with the modified Ashworth scale (MAS) [36]. Motor dysfunction was evaluated based on the Osame Motor Disability Score (OMDS), which grades motor dysfunction from 0 (normal walking and running) to 13 (complete bedridden) (see Table 2). The severity of the three main symptoms of urinary disturbance, i.e. increased frequency of urination, feeling of residual urine, and urinary incontinence were scored using the following scale: 0=normal; 1=slight; 2=moderate; 3=severe. The urinary disturbance score (UDS) represents the sum of all three symptoms. An overall evaluation of efficacy (improvement of dysfunction) based mainly on the improvement of motor dysfunction, as well as changes in urinary disturbances and neurological signs, was also evaluated according to the following scale, which had been previously used in the multicenter, randomized, double blind, and controlled study for IFN- $\alpha$  [21]: (1) excellent: patients showing two or more grades of improvement on the OMDS; (2) good: patients with one grade of improvement on the OMDS, or patients with an improvement of motor dysfunction with no change on the OMDS, but apparent improvement on more than two items other than motor dysfunction; (3) fair: patients exhibiting an improvement of motor dysfunction with no change on the OMDS, or patients with no improvement in motor dysfunction but with an improvement on one or more of the other items examined; (4) poor: patients with no improvement; and (5) none: patients with continuing clinical deterioration. In this study, the entire HTLV-1 provirus load data was measured after 4 weeks of LcS administration were completed, by an investigator who was not involved in the patients' clinical care. Clinical neurologists did not have access to provirus load data.

### 2.6. Lymphocytes phenotyping by flow cytometric analysis

After thawing, cells were washed three times with phosphate-buffered saline (PBS) and fixed in a PBS containing 2% paraformaldehyde (Sigma) for 20 min, and then resuspended in PBS at 4 °C. Fixed cells were washed with PBS containing 7% of normal goat serum (Sigma). Cells were then incubated for 15 min at room temperature with various combinations of fluorescence-conjugated monoclonal antibodies (mAb) as follows: fluorescein isothiocyanate (FITC)-labeled anti-CD27 (M-T271; Pharmingen, CA), FITC-labeled anti-CD3 (UCHT1; Beckman Coulter, CA), FITC-labeled anti-pan  $\gamma\delta$  T cell (Immu 510; Beckman Coulter, CA), Cy-Chrome-labeled anti-CD45RA (HI100; Pharmingen), phycoerythrin (PE)-labeled anti-CXCR3 (1C6/CXCR3; Pharmingen), PE-labeled anti-CD16 (3G8; Beckman Coulter), PE-labeled anti-NKG2A (Z199; Beckman Coulter), phycoerythrin-Cy5

Table 2

Osame Motor Disability Score (OMDS) for HAM/TSP

Grade	Disability
0	Normal gait and running
1	Normal gait but runs slow
2	Abnormal gait (staggering or spastic)
3	Abnormal gait and unable to run
4	Needs support (handrails) while using stairs but limited to 10 m
5	Needs one-hand support in walking
6	Needs two-hand support in walking
7	Needs two-hand support in walking but is limited to 10 m
8	Needs two-hand support in walking but limited to 5 m
9	Unable to walk even with support but can crawl in hands and knees
10	Crawls with hands
11	Unable to crawl but can turn sideways in bed
12	Unable to crawl but can move the toes
13	Completely bedridden

(PC5)-labeled anti-CD4 (13B8.2; Beckman Coulter), PC5-labeled anti-CD56 (N901; Beckman Coulter), and energy-coupled dye (ECD)-labeled anti-CD8 (SFCI21ThyD3; Beckman Coulter). Isotype-matched mouse immunoglobulins were used as a control. The phenotype was determined by flow cytometry (EPICS<sup>®</sup> XL; Beckman Coulter, Tokyo, Japan) in the lymphocyte gate, based on forward versus side scatter.

### 2.7. NK cell-mediated cytotoxic assays

The erythroleukemia cell line K562 was maintained in RPMI 1640 medium, and supplemented with 10% FBS. Target cells ( $1 \times 10^6$  viable K562 cells) were labeled with  $\text{Na}_2^{51}\text{CrO}_4$  (100  $\mu\text{Ci}$ ) for 1 h at 37 °C, washed, and resuspended at a concentration of  $5 \times 10^4$  cells/ml in a culture medium. Effectors and labeled targets were incubated in triplicates in 0.2 ml volume at E/T ratio of 50:1 in round-bottom 96-well plates, and incubated for 4 h. Percentage of lysis was determined by the following formula: % specific lysis =  $100 \times (\text{exp} - \text{spont}) / (\text{max} - \text{spont})$ , where experimental (exp) release represents the radioactivity from the experimental wells, maximum (max) release represents counts from detergent-lysed targets, and spontaneous (spont) release represents background release from wells with targets alone.

### 2.8. Quantification of HTLV-1 provirus load and anti-HTLV-1 antibody titers

To examine the HTLV-1 provirus load, we carried out a quantitative PCR using ABI Prism 7700<sup>™</sup> (PE-Applied Biosystems) with 100 ng of genomic DNA (roughly equivalent to  $10^4$  cells) from PBMCs as reported previously [12]. We chose the pX gene of HTLV-1, which has four open reading frames and encodes the unique regulatory and accessory proteins, for quantitative PCR because pX gene is retained by defective proviruses that are sometimes present in the infected individuals. Using  $\beta$ -actin as an internal control, the amount of HTLV-1 proviral DNA was calculated through the following formula: copy number of HTLV-1 (pX) per  $1 \times 10^4$  PBMC =  $[(\text{copy number of pX}) / (\text{copy number of } \beta\text{-actin} / 2)] \times 10^4$ . All samples were performed in triplicate. Serum and CSF antibody titers to HTLV-1 were determined by a particle agglutination method (Serodia-HTLV-1<sup>®</sup>, Fujirebio).

### 2.9. Statistical analysis

We made a paired comparison of changes in T cell phenotypes, HTLV-1 provirus load, NK cell activity, OMDS, and UDS scores between LcS treatments (before and 4 weeks after LcS treatment) by using a Wilcoxon signed rank test. The results represent the mean  $\pm$  standard deviation (S.D.) where applicable. Values of  $p < 0.05$  were considered statistically significant.

## 3. Results

### 3.1. Clinical improvement of HAM/TSP with LcS treatment

We studied 10 HAM/TSP patients who received a LcS preparation, which contains approximately  $4 \times 10^{10}$  viable LcS, orally once a day for 4 weeks. All 10 patients showed the rate of excellent to fair response by overall evaluation of efficacy (see Patients and methods). Motor dysfunction (OMDS) was improved after treatment in 5 out of 10 patients ( $p = 0.157$ , Wilcoxon signed rank test). One patient (HAM 4) who had needed a walking cane before treatment became able to walk without assistance after treatment. Urinary symptoms (UDS) were improved in all 10 patients ( $p = 0.0085$ , Wilcoxon signed rank test; Table 1). Nine out of 9 patients showed decreased spasticity on neurological examination ( $p = 0.015$ , Wilcoxon signed rank test; Table 1). No adverse effect and laboratory findings were observed. The concomitant administration of prednisolone was not associated with the specific clinical outcome against LcS treatment (Table 1).

### 3.2. Laboratory findings

Simultaneous staining with CD45RA and CD27 mAbs has been demonstrated to identify subpopulations of human  $\text{CD8}^{\text{high}}$  T cells that were distinct for both phenotypic and functional properties [37]. From this staining, unprimed naïve T cells were defined as  $\text{CD45RA}^+\text{CD27}^+$ , antigen-experienced effector T cells that parallel with CTLs were defined as  $\text{CD45RA}^+\text{CD27}^-$ , and memory T cells were defined as  $\text{CD45RA}^-\text{CD27}^+$ . The  $\text{CD45RA}^-\text{CD27}^-$  population was found to contain both effector and memory type T cells. In accordance with this phenotype,  $\text{CD45RA}^+\text{CD27}^-$  but not  $\text{CD45RA}^-\text{CD27}^+$  T cells have cytolytic activity without previous in vitro stimulation [37]. Our data have shown that the frequency as well as the absolute number of all the T cell phenotypes examined were not significantly changed in patients who received 4 weeks of daily oral LcS administration (Table 3). Interestingly, although the frequency as well as the absolute number of  $\text{CD16}^+\text{CD56}^+/\text{CD3}^-$  NK cells and inhibitory NK receptor  $\text{NKG2A}^+$  cells, which was a selectively decreased on T cells in HAM/TSP patients [34], showed no significant change after 4 weeks of LcS treatment, NK cell activity, which was reported to be decreased in HAM/TSP patients [32,33], was significantly increased after LcS treatment (Table 3). However, the concomitant administration of prednisolone was not associated with specific changes of NK cell activity against LcS administration (Fig. 1).

### 3.3. Provirus load of HTLV-1

The change in the HTLV-1 provirus load was measured by competitive PCR before and 4 weeks after LcS administration. The copy number of HTLV-1 per  $1 \times 10^4$  PBMCs was not significantly changed after 4 weeks of LcS

Table 3

Absolute number, frequency of cell subsets and HTLV-1 provirus copy number in peripheral blood mononuclear cells (PBMCs) from patients with human T lymphotropic virus type 1 (HTLV-1)-associated myelopathy/tropical spastic paraparesis (HAM/TSP) who received *Lactobacillus casei* strain Shirota daily for 4 weeks

Cell type	Before treatment		After treatment		<i>P</i> <sup>a</sup>	
	Absolute count, cells $\times 10^2/\text{mm}^3$	Frequency, %	Absolute count, cells $\times 10^2/\text{mm}^3$	Frequency, %	Absolute count	Frequency
CD4 <sup>+</sup>	5.79 $\pm$ 4.54	26.38 $\pm$ 12.71	5.96 $\pm$ 3.16	27.13 $\pm$ 11.09	0.674	0.575
CD8 <sup>high+</sup>	3.89 $\pm$ 1.61	20.47 $\pm$ 6.93	4.72 $\pm$ 2.02	22.60 $\pm$ 9.01	0.208	0.327
Naive in CD8 <sup>high+</sup>	1.13 $\pm$ 1.71	5.24 $\pm$ 7.11	0.93 $\pm$ 1.06	4.25 $\pm$ 4.08	1.000	0.674
Memory in CD8 <sup>high+</sup>	4.84 $\pm$ 2.60	24.05 $\pm$ 6.44	5.25 $\pm$ 2.13	24.47 $\pm$ 7.37	0.401	0.889
Effector in CD8 <sup>high+</sup>	7.56 $\pm$ 4.46	37.35 $\pm$ 10.81	7.57 $\pm$ 2.56	36.10 $\pm$ 9.93	1.000	0.674
Effector/memory in CD8 <sup>high+</sup>	5.99 $\pm$ 2.36	33.36 $\pm$ 14.54	7.21 $\pm$ 3.47	34.07 $\pm$ 13.78	0.263	0.889
CXCR3 <sup>+</sup>	4.38 $\pm$ 2.38	21.07 $\pm$ 4.28	3.93 $\pm$ 1.16	18.78 $\pm$ 6.57	0.735	0.398
CXCR3 <sup>+</sup> in CD4 <sup>+</sup>	2.03 $\pm$ 2.13	8.26 $\pm$ 5.06	1.60 $\pm$ 0.75	7.46 $\pm$ 3.28	0.866	0.612
$\delta\gamma\text{T}^+$	0.40 $\pm$ 0.25	2.28 $\pm$ 1.46	0.47 $\pm$ 0.34	2.35 $\pm$ 1.83	0.208	0.674
NKG2A <sup>+</sup>	0.72 $\pm$ 0.43	3.68 $\pm$ 1.84	0.74 $\pm$ 0.37	3.84 $\pm$ 2.39	0.674	0.889
CD16 <sup>+</sup> CD56 <sup>+</sup> CD3 <sup>-</sup>	3.14 $\pm$ 1.32	18.63 $\pm$ 11.33	3.38 $\pm$ 1.73	16.13 $\pm$ 8.81	0.575	0.635
NK cell activity (%) <sup>b</sup>	26.54 $\pm$ 16.13		39.43 $\pm$ 15.48		<b>0.015</b>	
HTLV-1 provirus load <sup>c</sup>	867.38 $\pm$ 874.62		641.75 $\pm$ 343.12		0.401	

Data are mean $\pm$ S.D. of 10 HAM/TSP patients (except for NK cell activity, *n*=9). Significant *P* value is in bold.

<sup>a</sup> Calculated using the Wilcoxon signed rank test.

<sup>b</sup> NK cell activity measured against K562 target cells at E/T ratio of 50/1 by the Cr release assay (*n*=9).

<sup>c</sup> HTLV-1 (pX) copy number per  $1 \times 10^4$  PBMC by quantitative PCR.

treatment (Table 3). HTLV-1 antibody titers in sera and CSF also showed no significant change after 4 weeks of LcS treatment (data not shown).




Fig. 1. Increased NK cell activity in PBMCs of HAM/TSP patients after oral administration of fermented milk drink containing *Lactobacillus casei* strain Shirota (LcS). The NK cell activity was significantly increased after LcS treatment (*n*=9, *p*=0.015, Wilcoxon's signed rank test). Each patient was given LcS preparation, which contains approximately  $4 \times 10^{10}$  viable LcS, orally twice a day for 4 weeks. NK cell activity was measured against K562 target cells at E/T ratio of 50/1 by the Cr release assay. Four patients out of ten received low dose oral prednisolone throughout the study period. None of the patients except for these four had received any immunomodulators. ----- Patients who received 5 mg/day oral prednisolone. ----: Patient who received 10 mg/day oral prednisolone.

#### 4. Discussion

In this study, we demonstrated that clinical improvement was observed in all the HAM/TSP patients following 4 weeks of daily oral administration of LcS. Especially, improvement of urinary symptoms (UDS) was observed in all the 10 patients examined (*p*=0.0085, Wilcoxon signed rank test). LcS was also effective for motor dysfunction (OMDS) and spasticity (MAS) on lower extremity. Nine out of 10 patients showed decreased spasticity on neurological examination whereas OMDS score was improved only in 5 out of 10 patients. Thus, differences in OMDS score did not reach statistical significance (*p*=0.157, Wilcoxon signed rank test) whereas MAS scores did (*p*=0.015, Wilcoxon signed rank test). Although transient and mild diarrhea was previously reported in minor population (3 out of 65, 4.6%) after LcS administration [38], no adverse effect was observed in our patients.

Laboratory findings revealed that the NK cell activity in HAM/TSP patients was significantly increased after LcS treatment, but that HTLV-1 provirus load as well as the frequencies and absolute numbers of all the cell phenotypes examined were not significantly changed after treatment. In contrast, we have previously reported that the frequency of memory cells (CD45RA<sup>-</sup>CD27<sup>+</sup>) within CD8<sup>high+</sup>, CXCR3<sup>+</sup> cells and HTLV-1 provirus load were significantly decreased after IFN- $\alpha$  treatment [22]. These findings suggest the possibility that NK cell activity is involved in the mechanism of clinical efficacy in LcS treatment, whereas Th1 type T cells and decreased HTLV-1 proviral load is associated with IFN- $\alpha$  effectiveness. Our present observation is consistent with the previous observation by Nagao et al. [39], who reported that in a healthy human subject, the frequency and number of NK cells, CD4<sup>+</sup> T

cells or CD8<sup>+</sup> T cells and T cell responsiveness to mitogens (Con A and PHA) were not significantly changed after 3 weeks of oral intake of LcS preparation containing approximately  $4 \times 10^{10}$  viable LcS. They also showed that the NK cell activity was significantly increased, and the enhancement of NK cell activity was particularly prominent in the low-NK cell activity individuals. Interestingly, our HAM/TSP case with most obvious clinical improvement (HAM 4) showed the lowest NK activity of all 10 patients before LcS treatment. However, another HAM/TSP case (HAM6) with low levels of NK activity before LcS treatment also showed the marked enhancement of NK cell activity after LcS but this patient did not show the obvious clinical improvement as seen in HAM4. Therefore, the level of enhanced NK cell activity after treatment did not always correlate with the level of clinical improvement. Since both of these patients (HAM 4 and 6) received low dose oral prednisolone throughout study period, this may also suggest that the concomitant administration of prednisolone is not specifically associated with the clinical outcome against LcS administration.

In general, probiotic bacteria such as LcS favorably alter the intestinal microflora balance, inhibit the growth of harmful bacteria, promote good digestion, and may boost immune function, and possibly increase resistance to infection. Although the mechanism of increased NK cell activity after oral administration of LcS is not clear, several studies have reported an effect of LcS on the stimulation of phagocytic activity of circulating immune cell populations which participate in innate immunity. Stimulated phagocytic cells may also play a role in potentiate acquired immunity via presentation of the antigen or synthesis of cytokines. Previous histological examination in an animal study showed that LcS was taken up by M cells in Peyer's patches [40], suggesting that LcS may be degraded in gut-associated lymphoid tissue and their signal from immunocompetent cells leads to a systemic effect such as stimulation of phagocytosis, synthesis of cytokines, IgA secretion, enhancement of NK cell activity, etc. Our data indicates that NK cell numbers were not significantly changed after LcS treatment. Also, we could not detect any significant changes of frequencies and absolute numbers of all the cell phenotypes examined, including the NK inhibitory receptor NKG2A positive cells and  $\gamma\delta$ T cells. It is therefore likely that increased NK cell activity might be regulated by increased expression levels of cytotoxic molecules—rather than the changes of activating receptors and/or inhibitory receptors on NK cells—although further studies are necessary to clarify this point.

Previous studies have indicated that therapies effective for HAM/TSP such as prednisolone [15,16], plasma exchange [17], and interferon- $\alpha$  [18–22] have problems associated with a high frequency of adverse effects; furthermore, IFN- $\alpha$  and plasmapheresis usually require hospitalization, are very expensive and the clinical effects usually last only a few months. Since HAM/TSP is a

chronic and progressive disorder, establishment of a chronic and safe treatment is essential. Also, innate immunity is thought to be important in protecting the host against many viral infections [41], and a combination treatment of acquired immunity oriented agents like IFN- $\alpha$  and innate immunity oriented agents like LcS might be a more desirable approach for HAM/TSP treatment. Since previous reports have indicated that the NK cell activity was significantly decreased in HAM/TSP patients by an unknown mechanism [32,33], the augmentation of a decreased host innate immune system by LcS might be a suitable approach for treating HAM/TSP.

In conclusion, our present results suggest that oral LcS may be comparatively effective, easy and safe, and is therefore a good candidate for maintenance treatment for HAM/TSP. However, in our present study, the number of patients is small and the study was conducted in an open, uncontrolled manner. Randomized controlled studies are warranted for the evaluation of LcS in HAM/TSP treatment.

#### Acknowledgments

The authors thank Ms. Tomoko Muramoto and Yoko Nishino of Kagoshima University for their excellent technical assistance. This work was supported by the Grant in Aid for Research on Brain Science of the Ministry of Health, Labor and Welfare, Japan.

#### References

- [1] Osame M, Usuku K, Izumo S, Ijichi N, Amitani H, Igata A, et al. HTLV-I associated myelopathy, a new clinical entity. *Lancet* 1986; 1:1031–2.
- [2] Gessain A, Barin F, Vernant JC, Gout O, Maurs L, Calender A, et al. Antibodies to human T-lymphotropic virus type-1 in patients with tropical spastic paraparesis. *Lancet* 1985;2:407–10.
- [3] Nakagawa M, Izumo S, Ijichi S, Kubota H, Arimura K, Kawabata M, et al. HTLV-I-associated myelopathy: analysis of 213 patients based on clinical features and laboratory findings. *J Neurovirol* 1998;4:586–93.
- [4] Nagai M, Usuku K, Matsumoto W, Kodama D, Takenouchi N, Moritoyo T, et al. Analysis of HTLV-I proviral load in 202 HAM/TSP patients and 243 asymptomatic HTLV-I carriers: high proviral load strongly predisposes to HAM/TSP. *J Neurovirol* 1998;4:586–93.
- [5] Mori M, Kinoshita K, Ban N, Yamada Y, Shiku H. Activated T-lymphocyte with polyclonal gammopathy in patients with human T-lymphotropic virus type I-associated myelopathy. *Ann Neurol* 1988;24:280–2.
- [6] Ijichi S, Eiraku N, Osame M, Izumo S, Kubota R, Maruyama I, et al. Activated T lymphocytes in cerebrospinal fluid of patients with HTLV-I-associated myelopathy (HAM/TSP). *J Neuroimmunol* 1989; 25:251–4.
- [7] Itoyama Y, Minato S, Kira J, Goto I, Sato H, Okochi K, et al. Spontaneous proliferation of peripheral blood lymphocytes increased in patients with HTLV-I-associated myelopathy. *Neurology* 1988; 38:1302–7.
- [8] Jacobson S, Zaninovic V, Mora C, Rodgers-Johnson P, Sheremata WA, Gibbs CJ, et al. Immunological findings in neurological diseases