

- A., Role of inorganic polyphosphate in promoting ribosomal protein degradation by the Lon protease in *E. coli*. *Science* 2001, **293**:705-708.
25. Kuroda, A., Tanaka, S., Ikeda, T., Kato, J., Takiguchi, N. and Ohtake, H., Inorganic polyphosphate kinase is required to stimulate protein degradation and for adaptation to amino acid starvation in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 1999, **96**:14264-14269.
26. Mandelstam, J., Turnover of protein in growing and non-growing populations of *Escherichia coli*. *Biochem. J.* 1958, **69**: 110-119.
27. Maurizi, M. R., Proteases and protein degradation in *Escherichia coli*. *Experientia* 1992, **48**:178-201.
28. Morohoshi, T., Maruo, T., Shirai, Y., Kato, J., Ikeda, T., Takiguchi, N., Ohtake, H. and Kuroda, A., Accumulation of inorganic polyphosphate in *phoU* mutants of *Escherichia coli* and *Synechocystis sp.* strain PCC6803. *Appl. Environ. Microbiol.* 2002, **68**:4107-4110.
29. Nash, H. A. and Robertson, C. A., Purification and properties of the *Escherichia coli* protein factor required for lambda integrative recombination. *J. Biol. Chem.* 1981, **256**:9246-9253.
30. Nomura, K., Kato, J., Takiguchi, N., Ohtake, H. and Kuroda, A., Effects of inorganic polyphosphate on the proteolytic and DNA-binding activities of Lon in *Escherichia coli*. *J. Biol. Chem.* 2004, **279**:34406-34410.
31. Prentki, P., Chandler, M. and Galas, D. J., *Escherichia coli* integration host factor bends the DNA at the ends of IS1 and in an insertion hotspot with multiple IHF binding sites. *EMBO J.* 1987, **6**:2479-2487.
32. Ross, W., Thompson, J. F., Newlands, J. T. and Gourse, R. L., *E. coli* Fis protein activates ribosomal RNA transcription *in vitro* and *in vivo*. *EMBO J.* 1990, **9**:3733-3742.
33. Rouviere-Yaniv, J., Yaniv, M. and Germond, J. E., *E. coli* DNA binding protein HU forms nucleosomelike structure with circular double-stranded DNA. *Cell* 1979, **17**:265-274.
34. Sambrook, J., Fritsch, E. F. and Maniatis, T., *Molecular cloning: a laboratory manual, 2nd ed.* Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 1989.
35. Surette, M. G. and Chaconas, G., A protein factor which reduces the negative supercoiling requirement in the Mu DNA strand transfer reaction is *Escherichia coli* integration host factor. *J. Biol. Chem.* 1989, **264**:3028-3034.
36. Sussman, A. J. and Gilvarg, C., Protein turnover in amino acid-starved strains of *Escherichia coli* K-12 differing in their ribonucleic acid control. *J. Biol. Chem.* 1969, **244**:6304-6306.
37. Verbeek, H., Nilsson, L., Baliko, G. and Bosch, L., Potential binding sites of the trans-activator FIS are present upstream of all rRNA operons and of many but not all tRNA operons. *Biochim. Biophys. Acta.* 1990, **1050**:302-306.
38. Yen, C., Green, L. and Miller, C. G., Degradation of intracellular protein in *Salmonella typhimurium* peptidase mutants. *J. Mol. Biol.* 1980, **143**:21-33.
39. Zuilanello, L., de la Gorgue de Rosny, E., van Ulsen, P., van de Putte, P. and Goosen, N., The HimA and HimD subunits of integration host factor can specifically bind to DNA as homodimers. *EMBO J.* 1994, **13**:1534-1540.