

“Health and Labour Sciences Research Grant for Research on Medical Devices for Analyzing, Supporting and Substituting the Function of Human Body” from the Ministry of Health, Labour and Welfare of Japan, “Program for Promotion of Fundamental Studies in Health Science” from the National Institute of Biomedical Innovation, a Grant provided by the Ichiro Kanehara Foundation, “Ground-based Research Announcement for Space Utilization” promoted by Japan Space Forum, and “Industrial Technology Research Grant Program in 03A47075” from New Energy and Industrial Technology Development Organization (NEDO) of Japan.

## References

- Borgdorff P, Versteeg PG. An implantable nerve cooler for the exercising dog. *Eur J Appl Physiol Occup Physiol* 1984;53(2):175–9.
- Bowser-Riley F, Cornish M, Kidd C, Zheng F. Role of vagal afferent C fibres in the bradycardiac response to an increase in arterial blood pressure in ferrets. *Exp Physiol* 1990;75(6):859–62.
- Costanzo EM, Barry JA, Ribchester RR. Co-regulation of synaptic efficacy at stable polyneuronally innervated neuromuscular junctions in reinnervated rat muscle. *J Physiol* 1999;521(Pt 2):365–74.
- Cudd TA. Vagal blockade does not prevent adrenocorticotropin, cortisol, and cardiopulmonary responses to thromboxane A<sub>2</sub>. *Can J Physiol Pharmacol* 1998;76(12):1087–94.
- Derksen FJ, Robinson NE, Stick JA. Technique for reversible vagal blockade in the standing conscious pony. *Am J Vet Res* 1981;42(3):523–5.
- Gokin AP, Philip B, Strichartz GR. Preferential block of small myelinated sensory and motor fibers by lidocaine: in vivo electrophysiology in the rat sciatic nerve. *Anesthesiology* 2001;95(6):1441–54.
- Kanje M, Lundborg G, Edstrom A. A new method for studies of the effects of locally applied drugs on peripheral nerve regeneration in vivo. *Brain Res* 1988;439(1–2):116–21.
- Lee LY, Morton RF, Kou YR. Acute effects of cigarette smoke on breathing in rats: vagal and nonvagal mechanisms. *J Appl Physiol* 1990;68(3):955–61.
- Li M, Zheng C, Kawada T, Sato T. Vagal stimulation improved acute-phase survival after myocardial infarction in conscious rats. *Circulation* 2003;108(Suppl. IV):IV-268, 1268.
- Potocnik I, Tomsic M, Bajrovic F. Sensitivity of sensory axons to lidocaine nerve block in rats. *Pflugers Arch* 2001;442(6 Suppl. 1):R193–4.
- Schultz HD, Gardner DG, Deschepper CF, Coleridge HM, Coleridge JC. Vagal C-fiber blockade abolishes sympathetic inhibition by atrial natriuretic factor. *Am J Physiol* 1988;255(1 Pt 2):R6–13.
- Thalhammer JG, Vladimirova M, Bershinsky B, Strichartz GR. Neurologic evaluation of the rat during sciatic nerve block with lidocaine. *Anesthesiology* 1995;82(4):1013–25.
- Vizek M, Frydrychova M, Houstek S, Palecek F. Effect of vagal cooling on lung functional residual capacity in rats with pneumonia. *Bull Eur Physiopathol Respir* 1983;19(1):23–6.