

- residues involved in substrate recognition by a rat H⁺/peptide cotransporter, PEPT1. *FEBS Lett* **394**:196–200.
- Tsuda M, Terada T, Asaka J, Ueba M, Katsura T, and Inui K (2007) Oppositely-directed H⁺ gradient functions as a driving force of rat H⁺/organic cation antiporter MATE1. *Am J Physiol* **292**:F593–F598.
- Uchiyama T, Kulkarni AA, Davies DL, and Lee VH (2003) Biophysical evidence for His57 as a proton-binding site in the mammalian intestinal transporter hPepT1. *Pharm Res (NY)* **20**:1911–1916.
- Urakami Y, Akazawa M, Saito H, Okuda M, and Inui K (2002) cDNA cloning, functional characterization, and tissue distribution of an alternatively spliced variant of organic cation transporter hOCT2 predominantly expressed in the human kidney. *J Am Soc Nephrol* **13**:1703–1710.
- Wright SH (2005) Role of organic cation transporters in the renal handling of therapeutic agents and xenobiotics. *Toxicol Appl Pharmacol* **204**:309–319.
- Yonezawa A, Masuda S, Yokoo S, Katsura T, and Inui K (2006) Cisplatin and oxaliplatin, but not carboplatin and nedaplatin, are substrates for human organic cation transporters (SLC22A1–3 and MATE family). *J Pharmacol Exp Ther* **379**: 879–886.

Address correspondence to: Dr. Professor Ken-ichi Inui, Department of Pharmacy, Kyoto University Hospital, Sakyo-ku, Kyoto 606-8507, Japan.
E-mail: inui@kuhp.kyoto-u.ac.jp
