

Ogawa K, Hozumi T, Sugioka K, Matsumura Y, Nishiura M, Kanda R, Abe Y, Takemoto Y, Yoshiyama M, Yoshikawa J.	Usefulness of automated quantitation of regional left ventricular wall motion by a novel method of two-dimensional echocardiographic tracking.	Am J Cardiol.	98(11)	1531-1537	2006
Nakamura Y, Wang X, Xu C, Asakura A, Yoshiyama M, From AH, Zhang J.	Xenotransplantation of long-term-cultured swine bone marrow-derived mesenchymal stem cells.	Stem Cells	25(3)	612-620	2006
Fujimoto K, Hozumi T, Watanabe H, Tokai K, Shimada K, Yoshiyama M, Homma S, Yoshikawa J.	Acute hyperglycemia induced by oral glucose loading suppresses coronary microcirculation on transthoracic Doppler echocardiography in healthy young adults.	Echocardiography.	23(10)	829-834	2006
Kusuyama T, Omura T, Nishiya D, Enomoto S, Matsumoto R, Takeuchi K, Yoshikawa J, Yoshiyama M.	Effects of treatment for diabetes mellitus on circulating vascular progenitor cells.	J Pharmacol Sci.	102(1)	96-102	2006
Shirai N, Ikura Y, Naruko T, Ohsawa M, Sugama Y, Suekane T, Kitabayashi C, Ehara S, Yoshiyama M, Yoshikawa J, Ueda M.	Endothelin-converting enzyme expression in the neointima after percutaneous coronary intervention.	Osaka City Med J.	52(1)	29-37	2006
Hirai H, Shibata T, Aoyama T, Yoshiyama M, Omura T, Suehiro S.	Activation of mitogen-activated protein kinases, activator protein-1, and nuclear factor-kappaB during acute rejection after heterotopic heart transplantation in rats.	Osaka City Med J.	52(1)	9-19	2006

Takemoto Y, Hozumi T, Sugioka K, Watanabe H, Matsumura Y, Yoshiyama M, Takeuchi K, Yoshikawa J.	Automated three-dimensional analysis of mitral annular dynamics in patients with myocardial infarction using automated mitral annular tracking method.	Echocardiograph y.	23(8)	658-665	2006
Wake R, Takeuchi M, Yoshitani H, Miyazaki C, Otani S, Yoshiyama M, Yoshikawa J.	Role of contrast-enhanced dobutamine stress echocardiography in predicting outcome in patients with known or suspected coronary artery disease.	Echocardiograph y	23(8)	642-649	2006
Okajima K, Kawase Y, Matsushita N, Iwata S, Doi A, Hasegawa T, Hato K, Nishimoto M, Abe Y, Yoshiyama M, Yoshikawa J.	Usefulness of myocardial contrast echocardiography with nicorandil stress for the detection of coronary artery stenosis.	Heart	92(9)	1331-1332	2006
Kusuyama T, Omura T, Nishiya D, Enomoto S, Matsumoto R, Murata T, Takeuchi K, Yoshikawa J, Yoshiyama M.	The effects of HMG-CoA reductase inhibitor on vascular progenitor cells.	J Pharmacol Sci.	101(4)	344-349	2006
Iwata S, Hozumi T, Matsumura Y, Sugioka K, Yoshitani H, Murata E, Takemoto Y, Kobayashi Y, Yoshiyama M, Yoshikawa J.	Cut-off value of coronary flow velocity reserve by transthoracic Doppler echocardiography for the assessment of significant donor left anterior descending artery stenosis in patients with spontaneously visible collaterals.	Am J Cardiol.	98(3)	298-302	2006

Tatsumi H, Takagi M, Nakagawa E, Yamashita H, Yoshiyama M.	Risk stratification in patients with Brugada syndrome: analysis of daily fluctuations in 12-lead electrocardiogram (ECG) and signal-averaged electrocardiogram (SAECG).	J Cardiovasc Electrophysiol.	17(7)	705-711	2006
Wake R, Takeuchi M, Yoshiyama M, Yoshikawa J.	Quantitative assessment of left ventricular function during contrast-enhanced dobutamine stress echocardiography predicts future cardiac events in diabetic patients.	Circ J.	70(7)	868-874	2006
Nishiya D, Omura T, Shimada K, Matsumoto R, Kusuyama T, Enomoto S, Iwao H, Takeuchi K, Yoshikawa J, Yoshiyama M.	Effects of erythropoietin on cardiac remodeling after myocardial infarction.	J Pharmacol Sci.	101(1)	31-39	2006
Murata E, Hozumi T, Matsumura Y, Fujimoto K, Sugioka K, Takemoto Y, Watanabe H, Yamagishi H, Yoshiyama M, Iwao H, Yoshikawa J.	Coronary flow velocity reserve measurement in three major coronary arteries using transthoracic Doppler echocardiography.	Echocardiography.	23(4)	279-286	2006
Zhu H, Hozumi T, Takemoto Y, Takagi Y, Negishi K, Abo K, Sugioka K, Matsumura Y, Otsuka R, Yoshitani H, Nakao M, Yoshiyama M, Yoshikawa	J.Estimation of myocardial ischemia by diastolic strain analysis in exercise stress echocardiography: comparison with exercise thallium-201 single photon emission computed tomography.	J Cardiol.	47(4)	165-172	2006

Asawa K, Kataoka T, Kobayashi Y, Hasegawa T, Nishioka H, Yamashita H, Qiu Z, Ehara S, Hirose M, Kamimori K, Shimada K, Yoshiyama M, Yoshikawa J.	Method analysis for optimal continuous imaging using intravascular optical coherence tomography.	J Cardiol.	47(3)	133-141	2006
Shirai N, Naruko T, Ohsawa M, Ikura Y, Sugama Y, Hirayama M, Kitabayashi C, Ehara S, Inoue T, Itoh A, Haze K, Tanzawa K, Yoshiyama M, Yoshikawa J, Ueda M.	Expression of endothelin-converting enzyme, endothelin-1 and endothelin receptors at the site of percutaneous coronary intervention in humans.	J Hypertens.	24(4)	711-721	2006
Ehara S, Kobayashi Y, Yoshiyama M, Ueda M, Yoshikawa J.	Coronary artery calcification revisited.	J Atheroscler Thromb.	13(1)	31-37	2006
Naruko T, Ueda M, Ehara S, Itoh A, Haze K, Shirai N, Ikura Y, Ohsawa M, Itabe H, Kobayashi Y, Yamagishi H, Yoshiyama M, Yoshikawa J, Becker AE.	Persistent high levels of plasma oxidized low-density lipoprotein after acute myocardial infarction predict stent restenosis.	Arterioscler Thromb Vasc Biol.	26(4)	877-883	2006
Hirata K, Watanabe H, Otsuka R, Fujimoto K, Tokai K, Yamagishi H, Yoshiyama M, Yoshikawa J.	Noninvasive diagnosis of restenosis by transthoracic Doppler echocardiography after percutaneous coronary intervention: comparison with exercise TI-SPECT.	J Am Soc Echocardiogr.	19(2)	165-1712	2006

Hasegawa T, Ehara S, Kobayashi Y, Kataoka T, Yamashita H, Nishioka H, Asawa K, Yamagishi H, Yoshiyama M, Takeuchi K, Yoshikawa J, Ueda M.	Acute myocardial infarction: clinical characteristics and plaque morphology between expansive remodeling and constrictive remodeling by intravascular ultrasound.	Am Heart J.	151(2)	332-337	2006
Fukuda D, Shimada K, Tanaka A, Kusuyama T, Yamashita H, Ehara S, Nakamura Y, Kawarabayashi T, Iida H, Yoshiyama M, Yoshikawa J.	Comparison of levels of serum matrix metalloproteinase-9 in patients with acute myocardial infarction versus unstable angina pectoris versus stable angina pectoris.	Am J Cardiol.	97(2)	175-180	2006
Kim-Mitsuyama S, Izumi Y, Izumiya Y, Namba M, Yoshida K, Wake R, Yoshiyama M, Iwao H.	Dominant-negative c-Jun inhibits rat cardiac hypertrophy induced by angiotensin II and hypertension.	Gene Ther.	13(4)	348-355	2006
Fukuda D, Yoshiyama M, Shimada K, Yamashita H, Ehara S, Nakamura Y, Kamimori K, Tanaka A, Kawarabayashi T, Yoshikawa J.	Relation between aortic stiffness and coronary flow reserve in patients with coronary artery disease.	Heart.	92(6)	759-762	2006
Kubota, T. et al	CYP2A6 polymorphisms are associated with nicotine dependence and influence withdrawal symptoms in smoking cessation.	Pharmacogenomics J.	6	115-119	2006

Matsuda, T. et al	N-cadherin signals through Rac1 determined the localization of connexin 43 in cardiac myocytes.	J. Mol. Cell. Cardiol.	40	495-502	2006
Mohri, T. et al	Leukemia inhibitory factor induces endothelial differentiation in cardiac stem cells.	J. Biol. Chem.	281	6442-6447	2006
Maeda, M. et al	MTHFR gene polymorphism and diabetic retinopathy.	Current Diabetes Review.	2	467-476	2006
Yasui, M. et al	Antisense to cyclin D1 inhibits vascular endothelial growth factor-stimulated growth of vascular endothelial cells: Implication of tumor vascularization.	Clin. Cancer Res.	12	4720-4729	2006
Fukuda T. et al	Warfarin dose requirement for patients with both VKORC1 3673A/A and CYP2C9*3/*3 genotypes.	Clin. Pharmacol. Ther.	80	553-554	2006
Ito, T. et al	Degradation of NFAT5, a transcriptional regulator of osmotic stress-related genes, is a critical event for doxorubicin-induced cytotoxicity in cardiac myocytes.	J. Biol. Chem.	282	1152-1160	2007
Yamane, M. et al	Rac1 activity is required for cardiac myocyte alignment in response to mechanical stress.	Biochem. Biophys. Res. Commun.	353	1023-1027	2007
Hiramoto, Y. et al	Effect of bosentan on plasma endothelin-1 level in patients with pulmonary arterial hypertension.	Circ. J.	71	367-369	2007
Maeda, M. et al	MTHFR gene polymorphism is susceptible to diabetic retinopathy but not to diabetic nephropathy in Japanese type 2 diabetic patients.	J. Diabetes Complications	In press		

Nonen, S. et al	Polymorphisms of norepinephrine transporter and adrenergic receptor α_{1D} are associated with the response to β -blockers in dilated cardiomyopathy.	Pharmacogenomics J.	In press		
Tougou, K. et al	Genetic polymorphism of bile acid CoA: Amino acid N-acetyltransferase in Japanese individuals.	Drug Metab. Pharmacokinet.	In press		