

## References

1. Frishman WH, Grewall P: Serotonin and the heart. *Ann Med* 2000; **32**: 195–209.
2. Watts SW: Serotonin-induced contraction in mesenteric resistance arteries: signaling and changes in deoxycorticosterone acetate–salt hypertension. *Hypertension* 2002; **39**: 825–829.
3. Liolitsa D, Powell JF, Prince M, Lovestone S: Association study of the 5-HT(2A) receptor gene polymorphism, T102C and essential hypertension. *J Hum Hypertens* 2001; **15**: 335–339.
4. Yu BN, Wang A, Zhou G, *et al*: T102C genetic polymorphism of the 5-HT2A receptor in Chinese hypertensive patients and healthy controls. *Clin Exp Pharmacol Physiol* 2004; **31**: 847–849.
5. Abe M, Wu Z, Yamamoto M, *et al*: Association of dopamine beta-hydroxylase polymorphism with hypertension through interaction with fasting plasma glucose in Japanese. *Hypertens Res* 2005; **28**: 215–221.
6. Yang ZH, Richard V, von Segesser L, *et al*: Threshold concentrations of endothelin-1 potentiate contractions to nor-epinephrine and serotonin in human arteries. A new mechanism of vasospasm? *Circulation* 1990; **82**: 188–195.
7. Haugen G, Stray-Pedersen S: Effects of endothelin-1 on vascular tension in human umbilical vessels. *Early Hum Dev* 1991; **27**: 25–32.
8. Pietraszek MH, Takada Y, Takada A: Endothelins inhibit serotonin-induced platelet aggregation via a mechanism involving protein kinase C. *Eur J Pharmacol* 1992; **219**: 289–293.
9. Balligand JL, Godfraind T: Effect of nisoldipine on contractions evoked by endothelin-1 in human isolated distal and proximal coronary arteries and veins. *J Cardiovasc Pharmacol* 1994; **24**: 618–625.
10. Okatani Y, Taniguchi K, Sagara Y: Amplifying effect of endothelin-1 on serotonin-induced vasoconstriction of human umbilical artery. *Am J Obstet Gynecol* 1995; **172**: 1240–1245.
11. Yoshida M, Aizawa H, Hara N: Effect of endothelin-1 on the serotonin-induced contraction of smooth muscle in the guinea pig trachea. *Respiration* 1999; **66**: 59–64.
12. Liras A, Catalan RE, Martinez AM: Synergistic effect of endothelin-1 and serotonin in rabbit platelets: effect on tyrosine phosphorylation. *Thromb Res* 2000; **100**: 325–331.
13. Watanabe T, Pakala R, Katagiri T, Benedict CR: Angiotensin II and serotonin potentiate endothelin-1-induced vascular smooth muscle cell proliferation. *J Hypertens* 2001; **19**: 731–739.
14. Watanabe T, Pakala R, Katagiri T, Benedict CR: Antioxidant *N*-acetylcysteine inhibits vasoactive agents–potentiated mitogenic effect of mildly oxidized LDL on vascular smooth muscle cells. *Hypertens Res* 2002; **25**: 311–315.
15. Yao L, Kobori H, Rahman M, *et al*: Olmesartan improves endothelin-induced hypertension and oxidative stress in rats. *Hypertens Res* 2004; **27**: 493–500.
16. Tanaka C, Kamide K, Takiuchi S, Kawano Y, Miyata T: Evaluation of the Lys198Asn and –134delA genetic polymorphisms of the endothelin-1 gene. *Hypertens Res* 2004; **27**: 367–371.
17. Jin JJ, Nakura J, Wu Z, *et al*: Association of endothelin-1 gene variant with hypertension. *Hypertension* 2003; **41**: 163–167.
18. Holloway JW, Beghe B, Turner S, Hinks LJ, Day IN, Howell WM: Comparison of three methods for single nucleotide polymorphism typing for DNA bank studies: sequence-specific oligonucleotide probe hybridisation, TaqMan liquid phase hybridisation, and microplate array diagonal gel electrophoresis (MADGE). *Hum Mutat* 1999; **14**: 340–347.
19. Ishikawa K, Baba S, Katsuya T, *et al*: T+31C polymorphism of angiotensinogen gene and essential hypertension. *Hypertension* 2001; **37**: 281–285.
20. Nordfors L, Jansson M, Sandberg G, *et al*: Large-scale genotyping of single nucleotide polymorphisms by Pyrosequencing™ and validation against the 5′ nuclease (Taqman®) assay. *Hum Mutat* 2002; **19**: 395–401.
21. de Kok JB, Wiegerinck ET, Giesendorf BA, Swinkels DW: Rapid genotyping of single nucleotide polymorphisms using novel minor groove binding DNA oligonucleotides (MGB probes). *Hum Mutat* 2002; **19**: 554–559.
22. Ioannidis JP, Ntzani EE, Trikalinos TA: ‘Racial’ differences in genetic effects for complex diseases. *Nat Genet* 2004; **36**: 1312–1318.
23. Peroutka SJ: Serotonin receptor variants in disease: new therapeutic opportunities? *Ann NY Acad Sci* 1998; **861**: 16–25.
24. Poleskaya OO, Sokolov BP: Differential expression of the “C” and “T” alleles of the 5-HT2A receptor gene in the temporal cortex of normal individuals and schizophrenics. *J Neurosci Res* 2002; **67**: 812–822.

## Mutation of the Follicle-Stimulating Hormone Receptor Gene 5'-Untranslated Region Associated With Female Hypertension

Tomohiro Nakayama, Nobuhiro Kuroi, Morihiko Sano, Yasuharu Tabara, Tomohiro Katsuya, Toshio Ogihara, Yoshio Makita, Akira Hata, Michiko Yamada, Norio Takahashi, Nobuhito Hirawa, Satoshi Umemura, Tetsuro Miki, Masayoshi Soma

**Abstract**—Inactivating mutations in the follicle-stimulating hormone receptor (FSHR) gene have been reported to cause hereditary hypergonadotropic ovarian failure. It has been found recently that the FSHR knockout mouse exhibits hypertension. The aim of the present study was to investigate the association between polymorphisms in the human FSHR gene and essential hypertension (EH) by using single nucleotide polymorphisms (SNPs). We selected 5 SNPs in the gene (rs1394205, rs2055571, rs11692782, rs1007541, and rs2268361) and performed 2 genetic case-control studies in different populations. A confirmative case-control study was performed using 1035 EH patients and 1058 age-matched controls. Transcriptional activities were measured with a luciferase assay system. The first case-control study found that the A allele of rs1394205 was significantly higher in EH females ( $P=0.010$ ). In addition, in the confirmative case-control study, there was a significant difference for this SNP between female normotensive subjects (44.5%) and EH patients (50.7%) ( $P=0.043$ ). Multiple logistic regression analysis in female subjects also revealed a significant association of subjects with the A allele of rs1394205 with EH ( $P=0.033$ ), with the odds ratio calculated as 1.68 (95% CI: 1.04 to 2.73). Transcriptional activity of the A allele was  $56 \pm 8\%$  (mean  $\pm$  SD) of that observed for the G-type allele ( $P=0.001$ ). Serum estradiol levels were significantly lower in patients with the A/A genotype than in patients without the A/A genotype ( $P=0.004$ ). The SNP in the 5'-untranslated region of the FSHR gene affects levels of transcriptional activity and is a susceptibility mutation of EH in women. (*Hypertension*. 2006;48:512-518.)

**Key Words:** hypertension, essential ■ hormones ■ case-control studies

It is likely that essential hypertension (EH) is a polygenic disorder that results from the inheritance of a number of susceptibility genes. The causal genes identified may contribute from 30% to 50% of the variations in blood pressure seen among individuals.<sup>1</sup> These genetic determinants interact with environmental factors, such as dietary salt, to produce the final disease phenotype. Despite significant recent progress in genomic and statistical tools, the genetic dissection of human EH remains a major challenge.<sup>2</sup>

The effects of follicle-stimulating hormone (FSH) are mediated by its interaction with specific receptors and the activation of  $G_s$ , the stimulatory guanine nucleotide binding protein, which stimulates the enzyme adenylyl cyclase.<sup>3</sup> The FSH receptor (FSHR) belongs to the superfamily of G protein-coupled receptors that are characterized by the common struc-

tural feature of 7 transmembrane domains. They differ structurally from other G protein-coupled receptors in that they contain a large extracellular domain in the amino-terminal part of the receptor protein, which is required for interaction with complex glycoprotein hormones.<sup>4-6</sup> The human FSHR gene is localized to 2p21 to p16 and is composed of 10 exons.<sup>7</sup> The 5'-flanking region of the gene has neither a TATA nor a CCAAT box and exhibits promoter-type features that are seen in housekeeping genes.<sup>8</sup>

Inactivating mutations in the FSHR gene have been reported to cause hereditary hypergonadotropic ovarian failure in women.<sup>9</sup> These mutations have been shown to be associated with a recessive inheritance pattern, and all of the affected subjects have homozygous or compound heterozygous mutations in the coding region of the gene. The phenotype in men for

Received December 11, 2005; first decision January 4, 2006; revision accepted June 15, 2006.

From the Division of Molecular Diagnostics (T.N.), Advanced Medical Research Center, and Division of Nephrology and Endocrinology (M.S.), Department of Medicine, Nihon University School of Medicine (N.K., M.S.), Tokyo, Japan; Departments of Basic Medical Research and Education (Y.T.) and Geriatric Medicine (T.M.), Ehime University Graduate School of Medicine, Ehime, Japan; Department of Geriatric Medicine (T.K., T.O.), Osaka University Graduate School of Medicine, Osaka, Japan; Department of Pediatrics (Y.M.), Asahikawa Medical College, Asahikawa, Japan; Department of Public Health (A.H.), Graduate School of Medicine, Chiba University, Chiba, Japan; Departments of Clinical Studies (M.Y.) and Genetics (N.T.), Radiation Effects Research Foundation; and the Department of Medical Science and Cardiorenal Medicine (N.H., S.U.), Yokohama City University Graduate School of Medicine, Yokohama City, Japan.

Correspondence to Tomohiro Nakayama, Division of Molecular Diagnostics, Advanced Medical Research Center, Nihon University School of Medicine, Ooyaguchi-kamimachi, 30-1 Itabashi-ku, Tokyo 173-8610, Japan. E-mail: tnakayam@med.nihon-u.ac.jp

© 2006 American Heart Association, Inc.

*Hypertension* is available at <http://www.hypertensionaha.org>

DOI: 10.1161/01.HYP.0000233877.84343.d7

this particular FSHR mutation is less clear, with homozygous men found to have normal masculinization and circulating testosterone, normal or slightly elevated luteinizing hormone (LH), moderately elevated FSH, and slightly to severely reduced testicular volume.<sup>10</sup> Although men with inactivating mutations have abnormal semen parameters that range from severe-to-moderate oligozoospermia, normal sperm concentrations with low volumes, or teratozoospermia (reported in 1 individual), none have been found to be azoospermic, and there are reports of such individuals fathering children. These results suggest that FSHR mutations have a greater consequence in women than in men.

The FSHR knockout mice have ovarian insufficiency, low estrogen levels with functionally responsive estrogen receptors, and increased testosterone levels.<sup>11–13</sup> They also exhibit the same changes seen in postmenopausal women, such as osteoporosis, hypercholesterolemia, and weight gains. In 2003, Javeshghani et al<sup>14</sup> reported that these mice have increased blood pressure, indicating that there is vascular remodeling.

There have been no previous studies on the association between the FSHR gene and EH. The aim of the present study was to investigate the association between the human FSHR gene and EH through the use of single nucleotide polymorphisms (SNPs) in the human FSHR gene.

## Methods

### First Case–Control Study

Patients and control subjects from the northern area of Tokyo were recruited for the first case–control screening study. A total of 235 patients were diagnosed with EH according to the following criteria: seated systolic blood pressure (SBP) >160 mm Hg and/or diastolic blood pressure (DBP) >100 mm Hg on 3 occasions within 2 months after the first medical examination. None of the patients were using antihypertensive medications, and subjects diagnosed with secondary hypertension were excluded. We also included 237 normotensive (NT) healthy individuals as controls. None of the NT participants had a family history of hypertension, and all had SBP and DBP <130 and 85 mm Hg, respectively. A family history of hypertension was defined as a previous diagnosis of hypertension in grandparents, uncles, aunts, parents, or siblings. Informed consent was obtained from each individual as per a protocol approved by the Human Studies Committee of Nihon University.<sup>15</sup>

### Confirmative Case–Control Study: Subgroup Collaboration Study With the Hypertensive Section of the Japanese Millennium Project

For confirmation of a possible SNP association, we used a second set of 1035 EH patients and 1058 age-matched control subjects, who were recruited through a subgroup collaboration study with the hypertensive section of the Japanese Millennium Project. Six medical institutes took part in the collaborative study and collected data on hypertensive cases and controls. Hypertensive patients were defined as having SBP  $\geq$ 140 mm Hg or DBP  $\geq$ 90 mm Hg or were patients with chronic antihypertensive medication. To increase the statistical power of the present study, hypertensive subjects additionally had to meet the following criteria: age <60 years old or onset of hypertension <50 years of age, a family history of hypertension, and without obesity (body mass index <26 kg/m<sup>2</sup>). The NT criteria were as follows: SBP/DBP <130/85 mm Hg, without a family history of hypertension, and without obesity. In this study, both groups were recruited from throughout Japan, and informed consent was obtained from each individual as per the protocol approved by each institutions' human studies committee.

### Biochemical Analysis

Serum concentrations of total cholesterol, triglyceride, high-density lipoprotein cholesterol, uric acid, creatinine, and  $\gamma$ -glutamyl transpeptidase were measured using the methods of the clinical laboratory department of each hospital or institution.

### Genotyping

Using information about allelic frequencies of SNPs registered on the web site of the National Center for Biotechnology Information and Celera Discovery System–Applied Biosystems, 5 SNPs with minor allele frequencies >20% were selected. SNPs with relatively high minor allele frequencies have been shown to be very useful as genetic markers for genetic case–control studies.

We examined the association between EH and 5 SNPs in the human FSH receptor gene. All 5 of the SNPs were confirmed using the National Center for Biotechnology Information web site and Applied Biosystems–Celera Discovery System with the accession numbers, rs1394205 (C\_426553\_10), rs2055571 (C\_246842\_10), rs11692782 (C\_228130\_10), rs1007541 (C\_9561251\_1\_1), and rs2268361 (C\_11813031\_1\_1; Figure 1 a). Genotypes were determined using Assays-on-Demand kits (Applied Biosystems) together with TaqMan PCR.<sup>16</sup> When allele-specific fluorogenic probes hybridize to the template during the PCR, the 5' nuclease activity of Taq polymerase can discriminate alleles. Cleavage results in increased emission of a reporter dye that otherwise is quenched by the dye minor groove binder (MGB). Each 5' nuclease assay requires 2 unlabeled PCR primers and 2 allele-specific probes. Each probe is labeled with a reporter dye (VIC [a proprietary dye from Applied Biosystems] and 6-carboxyfluorescein [FAM]) at the 5' end and MGB at the 3' end. Amplification by PCR preceded using TaqMan Universal Master Mix (PE Biosystems) in a 25- $\mu$ l reaction volume containing 50 ng of DNA, 700 nM primer, and 100 nM probe final concentrations. Thermal cycling conditions consisted of 95°C for 10 minutes and then 40 cycles of 92°C for 15 s and 60°C for 1 minute in a GeneAmp 9700 system. Fluorescence levels of the PCR products were measured using an ABI PRISM 7700 Sequence Detector (Applied Biosystems), which resulted in 3 genotypes of 2 alleles being clearly identified. We analyzed *cis* elements in the 5' upstream region of the gene using AliBaba2 software that was obtained from the Biological Databases web site (<http://www.gene-regulation.com/pub/programs.html#alibaba2>).

### Linkage Disequilibrium Analysis


SNPAlyze version 3.2 (DYNACOM Co, Ltd) was used for determining haplotype and linkage disequilibrium (LD) analyses and is available online at <http://www.dynacom.co.jp/products/package/snpyalyze/index.html> (Figure 1b).

### Preparation of Plasmids Including the G-Type or A-Type DNA Fragment

G-type and A-type rs1394205 reporter constructs were made. Because it has been reported that the –250-bp upstream region of this gene has a core promoter activity, 2 oligonucleotide primers, 5'-CCTTAGGTCAGGGTGTAAAGAAACCC-3' (bases –202 to –226) and 5'-GGCCATAATTATGCATCCATCCACC-3' (bases +6 to –19), were designed (Figure 1c).<sup>8</sup> These primers contain the *Kpn*I and *Hind*III restriction sites, respectively. After digestion with *Kpn*I and *Hind*III, PCR products were subcloned into the *Kpn*I and *Hind*III sites of the luciferase reporter gene vector pGV-B2 (Tokyo Ink). The G-type and A-type constructs were verified by sequencing.

### Measurement of Transcriptional Activities of G-Type and A-Type Alleles

For transfection of Chinese hamster ovary (CHO) cells with FSHR gene promoter constructs, cells (60% to 70% confluent) were preincubated in OptiMEM medium (Lipofectamine, Gibco BRL) for 30 minutes at 37°C. FSHR gene promoter plasmids (1  $\mu$ g) and a plasmid containing TK-driven pRL (Toyo Ink, 200 ng, used to normalize for transfection efficiency) were mixed with liposome suspension (Lipofectamine,



**Figure 1.** a, Organization of the human FSH receptor gene and location of the SNPs used for the present case-control study. ■, exons, and —, indicate introns. b, Pairwise LD in the FSH receptor gene, as evaluated by  $D'$  and  $r^2$ . Pairwise LD among the 6 marker pairs studied in the FSH receptor gene were computed, and pairs in LD ( $D' > 0.5$  or  $r^2 < 0.1$ ) are shown as gray-shaded values. c, Nucleotide sequence of the 5' upstream region of the FSH receptor gene. →, the major transcriptional initiation site, and the first nucleotide upstream of the transcriptional initiation site is designated -1. The G/A in bold type indicates the SNP of rs1394205. The italicized ATG indicates the start codon. Underlined nucleotide sequences indicate the PCR primers for making the insert of the plasmid that was used to measure transcriptional activities. The double underline indicates the *cis* element of the transcription factor Ets-1.

Gibco BRL, 6  $\mu$ l/well) and incubated for 20 minutes at room temperature. The lipid-coated DNA was then added to each well containing 0.8 mL of OptiMEM media. After 3 hours, the medium was removed and replaced with complete medium for an additional 48 hours. CHO cells were then lysed (400  $\mu$ L), and extracts were centrifuged to remove intact cells and debris. Extracts (50  $\mu$ L) were used for measurement of luciferase activity. Luciferase activity was measured  $\geq 3$  times in duplicate with a double luciferase assay system (PicaGene Dual Sea-Pansy, Toyo Ink.) and a luminometer (LB-9507, Berthold). All of the data were normalized as relative light units/pRL-TK activity.<sup>17</sup>

## Serum Levels of Estradiol

We measured serum levels of estradiol in female EH patients. Seventeen subjects in the female EH group were selected randomly. All of the subjects were confirmed to be postmenopausal by their own testimony. Six patients had the A/A genotype, and 12 patients had the G/A or G/G genotypes. Serum estradiol levels were measured with the Electro Chemiluminescence Immunoassay (Elecsys Systems Immunoassay, Roche Diagnostics), as reported previously.<sup>18</sup>

## Statistical Analysis

Data are shown as mean  $\pm$  SD. All of the statistical analyses were conducted using StarView 5.0 (SAS Inc) and Dr. SPSS II (SPSS Inc). Hardy-Weinberg equilibrium was assessed by a  $\chi^2$  analysis. The overall distributions of the genotypes or alleles were analyzed by  $\chi^2$  analysis using 2 $\times$ 3 or 2 $\times$ 2 contingency tables between EH patients and NT controls. To assess the quantitative effects of covariates, multiple logistic regression analysis was performed using SPSS II. Statistical significance was established at  $P < 0.05$ .

## Results

### First Case-Control Study

Table 1 shows the distribution of genotypic and allelic frequencies of the 5 SNPs in each group. The observed and expected genotypic frequencies of each of the SNPs in the total subjects, male and female, of the NT group were in good agreement with the predicted Hardy-Weinberg equilibrium values (data not shown). The overall distribution of genotype and allele of all 5 of the SNPs did not significantly differ between men for the EH and NT groups. However, among women, the genotypic and allelic frequency of rs1394205 and the allelic frequency of rs2055571 showed a significant difference between the 2 groups. For the total subjects (male and female) the genotypic and allelic frequency of rs1394205 and the genotypic frequency of rs2268361 were significantly different between the EH and NT subjects. These data indicate that the distribution of rs1394205 in women can affect the overall analysis in the total subject group. Therefore, this could be used as a genetic marker of EH in women, because the probability value for the allelic distribution of this SNP was very significant ( $P = 0.003$ ).

Patterns of LD in the FSHR gene are illustrated by their  $D'$  and  $r^2$  values (Figure 1b). LD analysis for the NT group and the  $D'$  values indicate that rs1394205 and rs2055571 are located in one haplotype block, whereas the other 3 SNPs are located in another block. Because the  $r^2$  values were beyond 0.1 in rs1394205 to rs2055571, it is not advantageous to use these combinations to isolate susceptibility haplotypes.

Based on these results, rs1394205 was selected for the large-size case-control study. We were also interested in the fact that the rs1394205 may influence the transcriptional activity of the gene, because the SNP is located in the 5' upstream region.

### Confirmative Case-Control Study: Subgroup Collaboration Study With the Hypertensive Section of the Japanese Millennium Project

Table 2 shows the clinical features of the EH patients and NT controls. These 2 groups were age-matched for the total, male, and female groups. We performed the confirmative case-control study for the rs1394205 SNP using 1035 EH patients and 1058 NT controls.

TABLE 1. Genotype Distribution in NTs and Patients With EH of First Screening Analysis

Parameters	Total Subjects		<i>P</i>	Male Subjects		<i>P</i>	Female Subjects		
	NT	EH		NT	EH		NT	EH	<i>P</i>
No. of participants	235	237		152	158		83	79	
Variants									
rs1394205									
Genotype									
GG	64 (0.272)	45 (0.190)		34 (0.224)	30 (0.190)		30 (0.361)	15 (0.190)	
GA	123 (0.524)	125 (0.527)		81 (0.533)	84 (0.532)		42 (0.506)	41 (0.519)	
AA	48 (0.204)	67 (0.283)	0.040*	37 (0.243)	44 (0.278)	0.673	11 (0.133)	23 (0.291)	0.010*
Allele									
G	251 (0.534)	215 (0.454)		149 (0.490)	144 (0.456)		102 (0.614)	71 (0.449)	
A	219 (0.466)	259 (0.546)	0.013*	155 (0.510)	172 (0.544)	0.391	64 (0.386)	87 (0.551)	0.003*
rs2055571									
Genotype									
TT	121 (0.515)	140 (0.591)		82 (0.540)	93 (0.589)		39 (0.470)	47 (0.594)	
TC	93 (0.396)	84 (0.354)		59 (0.388)	55 (0.348)		34 (0.410)	29 (0.367)	
CC	21 (0.089)	13 (0.055)	0.156	11 (0.072)	10 (0.063)	0.683	10 (0.120)	3 (0.038)	0.090
Allele									
T	335 (0.713)	364 (0.768)		223 (0.734)	241 (0.763)		112 (0.675)	123 (0.778)	
C	135 (0.287)	110 (0.232)	0.053	81 (0.266)	75 (0.237)	0.404	54 (0.325)	35 (0.222)	0.036*
rs11692782									
Genotype									
AA	54 (0.23)	63 (0.266)		33 (0.217)	39 (0.247)		21 (0.253)	24 (0.304)	
AT	121 (0.515)	108 (0.456)		77 (0.507)	74 (0.468)		44 (0.530)	34 (0.430)	
TT	60 (0.255)	66 (0.278)	0.426	42 (0.276)	45 (0.285)	0.761	18 (0.217)	21 (0.266)	0.446
Allele									
A	229 (0.487)	234 (0.494)		143 (0.470)	152 (0.481)		86 (0.518)	82 (0.519)	
T	241 (0.513)	240 (0.506)	0.843	161 (0.530)	164 (0.519)	0.791	80 (0.482)	76 (0.481)	0.987
rs1007541									
Genotype									
GG	100 (0.426)	80 (0.338)		59 (0.388)	53 (0.335)		41 (0.494)	27 (0.342)	
GA	104 (0.442)	122 (0.514)		72 (0.474)	84 (0.532)		32 (0.386)	38 (0.481)	
AA	31 (0.132)	35 (0.148)	0.143	21 (0.138)	21 (0.133)	0.569	10 (0.120)	14 (0.177)	0.138
Allele									
G	304 (0.647)	282 (0.595)		190 (0.625)	190 (0.601)		114 (0.687)	92 (0.582)	
A	166 (0.353)	192 (0.405)	0.101	114 (0.375)	126 (0.399)	0.544	52 (0.313)	66 (0.418)	0.051
rs2268361									
Genotype									
GG	48 (0.204)	53 (0.224)		30 (0.197)	34 (0.215)		18 (0.217)	19 (0.241)	
GA	102 (0.434)	123 (0.519)		67 (0.441)	82 (0.519)		35 (0.422)	41 (0.518)	
AA	85 (0.362)	61 (0.257)	0.046*	55 (0.362)	42 (0.266)	0.184	30 (0.361)	19 (0.241)	0.238
Allele									
G	198 (0.421)	229 (0.483)		127 (0.418)	150 (0.475)		71 (0.428)	79 (0.500)	
A	272 (0.579)	245 (0.517)	0.056	177 (0.582)	166 (0.525)	0.154	95 (0.572)	79 (0.500)	0.192

\*Significant difference in distribution.

The observed and expected genotypic frequencies of each of the SNPs in the total subjects ( $P=0.553$ ), males ( $P=0.853$ ), and females ( $P=0.426$ ) of the NT group were in good agreement with the predicted Hardy–Weinberg equilibrium values. The overall distribution for the alleles of the SNPs did not signifi-

cantly differ between the total EH and total NT groups. However, among women, the allelic frequency of the A allele of rs1394205 was significantly higher for EH subjects than for NT subjects ( $P=0.042$ ; Table 3). Although the genotype distribution of rs1394205 among women in the first case–control study showed

TABLE 2. Characteristics of Study Participants of Subgroup Analysis of the Hypertension Team in Japanese Millennium Project

Parameters	Total			Male			Female		
	NT	EH	<i>P</i>	NT	EH	<i>P</i>	NT	EH	<i>P</i>
No. of subjects	1058	1035		792	762		266	273	
Age, y	60.4±7.9	59.7±9.5	0.052	59.6±7.3	59.1±8.7	0.269	63.0±9.1	61.3±11.1	0.051
BMI, kg/m <sup>2</sup>	22.4±2.7	23.5±2.8	<0.001*	22.5±2.8	23.4±2.5	<0.001*	22.2±2.7	23.8±3.6	<0.001*
SBP, mm Hg	114.1±10.2	161.5±22.3	<0.001*	114.6±10.4	160.8±21.3	<0.001*	111.4±8.6	164.0±25.2	<0.001*
DBP, mm Hg	70.2±7.0	97.1±14.4	<0.001*	70.6±6.9	97.2±14.0	<0.001*	68.1±7.2	96.8±15.9	<0.001*
Total cholesterol, mg/dL	203.3±36.0	204.2±34.0	0.609	198.4±34.3	198.8±32.2	0.818	215.7±37.1	218.5±34.7	0.364
Triglyceride, mg/dL	116.5±74.2	143.7±97.9	<0.001*	117.4±7.6	146.3±102.2	<0.001*	114.3±67.7	135.2±72.7	0.002*
HDL cholesterol, mg/dL	56.8±14.8	58.4±16.5	0.043*	55.7±14.6	57.7±16.4	0.022*	59.5±15.0	60.0±16.6	0.736
Uric acid, mg/dL	5.3±2.6	5.7±2.4	0.004*	5.6±1.3	6.1±2.7	<0.001*	4.8±4.3	4.8±1.1	0.962
Creatinine, mg/dL	0.81±0.19	0.83±0.27	0.106	0.86±0.18	0.89±0.29	0.015*	0.71±0.17	0.69±0.17	0.372
γ-GTP	35.4±39.7	52.0±71.7	<0.001*	44.4±46.5	68.1±87.1	<0.001*	25.8±27.9	29.5±29.8	0.190
Smoking, %	45	46.6	0.583	66.6	62.1	0.182	4.6	7.3	0.271
Alcohol consumption, %	49.1	43.2	0.045*	62.9	74.3	0.001*	25.2	17.1	0.053

BMI indicates body mass index.

\*Significant difference in distribution.

a significant difference, the genotype distribution in the confirmatory study did not exhibit any significant difference. This discrepancy could have been caused by differences in sample size.

Multiple logistic regression analysis in the female subjects revealed that the significant association of A/A and G/A with EH ( $P=0.033$ ) remained after adjustment for confounding factors (such as age, body mass index, total cholesterol, triglyceride, high-density lipoprotein cholesterol, uric acid, and creatinine), and the calculated odds ratio was 1.68 (95% CI: 1.04 to 2.73).

### Transcriptional Activities

To study transcriptional activity, we transfected CHO cells with promoter constructs of the FSHR gene. The promoter activity of the A allele was  $56\pm 8\%$  (mean±SD) of that for the G-type allele ( $P=0.001$ ; Figure 2). The rs1394205 is located in the Ets-1 site, and when there is substitution of the

A-type allele from the G-type allele, the Ets-1 site vanishes (Figure 1c).

### Serum Levels of Estradiol

All of the values of the serum levels of estradiol were <32 pg/mL, because all of the subjects were postmenopausal. The values of G/G, G/A, and A/A genotypes were  $19.8\pm 2.1$ ,  $20.5\pm 6.2$ , and  $13.3\pm 3.2$ , respectively. Because there was no difference between the G/G and G/A genotypes ( $P=0.807$ ), Figure 3 only shows the serum levels of estradiol in postmenopausal EH patients with and without the A/A genotype. Serum estradiol levels were significantly lower in patients with the A/A genotype than in patients without the A/A genotype ( $P=0.004$ ).


### Discussion

We performed 2 genetic case-control studies in different populations for the FSHR gene and found that the SNP in the

TABLE 3. Genotype Distribution in NTs and Patients With Essential Hypertension of Subgroup Analysis of Hypertension in Japanese Millennium Project

Parameters	Total Subjects			Male			Female		
	NT	EH	<i>P</i>	NT	EH	<i>P</i>	NT	EH	<i>P</i>
No. of participants	1058	1035		792	762		266	273	
Variants									
rs1394205									
Genotype									
GG	297 (0.281)	253 (0.244)		212 (0.268)	184 (0.242)		85 (0.320)	69 (0.253)	
GA	518 (0.489)	531 (0.513)		393 (0.496)	400 (0.525)		125 (0.470)	131 (0.480)	
AA	243 (0.230)	251 (0.243)	0.169	187 (0.236)	178 (0.233)	0.431	56 (0.210)	73 (0.267)	0.139
Allele									
G	1112 (0.526)	1037 (0.501)		817 (0.516)	768 (0.504)		295 (0.555)	269 (0.493)	
A	1004 (0.474)	1033 (0.499)	0.112	767 (0.484)	756 (0.496)	0.509	237 (0.445)	277 (0.507)	0.04*


\*Significant difference in distribution.



**Figure 2.** Transcriptional activities of the reporter constructs. The activities of the G-type constructs were considered as 100%. Values are expressed as the average of 3 independent experiments done on different days. Data are mean ± SD.

5'-untranslated region of the FSHR gene affects levels of transcriptional activity and is a susceptibility mutation of EH in women. Very recently it has been reported that knockout mice can be used as a model of postmenopausal hypertension with low levels of estrogen and high levels of testosterone.<sup>14</sup> This report is consistent with our results that indicate that there is both an association between the polymorphism in the FSHR gene and EH and that the levels of plasma estradiol in female EH patients with the A/A genotype are significantly lower than those in female EH patients without the genotype. Furthermore, because Ets-1 has been reported to be involved in the regulation of angiogenesis and cell growth,<sup>19,20</sup> our findings suggest that the substitution of the A-type allele from the G-type allele of the rs1394205 leads to a decreased transcriptional activity, which is followed by a subsequent loss of the Ets-1 site. Because there have been many studies reporting that low estrogen levels are associated with hypertension in postmenopausal women,<sup>21,22</sup> our results also may provide evidence on the etiology of  $\geq 1$  of the causes of hypertension in postmenopausal women.

In genetic case-control studies, pseudopositive results can sometimes occur. Therefore, to improve reliability, we increased the number of samples in addition to selecting samples strictly for the purpose of removing bias associated with confounding factors. It is especially advantageous to perform case-control studies using large and different populations, because this can lead to results that have a greater likelihood of being correct. The strategy of our study was to first carry out a screening case-control study followed by a confirmative case-control study that included >1000 subjects in each group. The rs1394205 showed significant differences for both case-control studies. It should be noted that the *P* values for the associations determined in the



**Figure 3.** Serum levels of estradiol. Data are mean ± SD.

present study are somewhat weak. However, in accordance with the guidelines issued for specific criteria for acceptability of association study results, the current study provides biological support (transient transfection finding) and plausibility based on knockout data in mice.<sup>23</sup> Therefore, the larger *P* value could be considered to be acceptable in this case.

Some case-control studies have identified gene variants associated with gender-specific susceptibility to EH<sup>24,25</sup>; however, in general, such studies do not explain the reason for the positive gender-specific association findings. Our experiment also found significant differences only for women. In contrast to previous reports, our experiment specifically examined whether the functional mutations of the gene were clearly identified in patients with a hereditary disease in a gender-specific manner. Mutations in the FSHR gene are reported to cause hereditary hypergonadotropic ovarian failure in women but not in men. To the best of our knowledge, our study is the first to report that the causal genes of gender-specific hereditary diseases are associated with EH in a gender-specific manner.

An increasing number of reports of genome-wide scans for hypertension and blood pressure variation have been seen in the past few years.<sup>26</sup> Some of the reports have shown that 2p, on which the FSHR gene is located, is associated with the candidate loci for EH.<sup>27-32</sup> However, these regions are broad, and there has been no definite susceptibility gene identified in this region. Although like our study in which the strategy for identification of susceptibility genes of EH was different between the genome-wide scans and case-control studies, the final goal for identifying the causal mutation for EH was the same. Therefore, because our current findings indicate that a gene variant is related to EH, our results are thought to be worthwhile. In conclusion, an SNP in the 5'-untranslated region of the FSHR gene affected the levels of transcriptional activity and, therefore, is a susceptibility mutation of EH in women.

### Perspectives

We performed 2-step case-control studies and identified that the SNP in the 5'-untranslated region of the FSHR gene affects levels of transcriptional activity, which is a susceptibility mutation of EH in women. However, there are some limitations associated with our study. It is possible that the case-control studies sometimes exhibit pseudopositive results because of sample scales or selection of the genetic markers. Further research is required to perform familial linkage studies and transmission disequilibrium tests for the purpose of confirming the reliability of the rs1394205 SNP.

### Acknowledgments

We thank Naoyuki Sato and Kaoru Sugama for their excellent technical assistance.

### Sources of Funding

This work was supported in part by a Grant-in-Aid for Scientific Research on Priority Areas (15012251) from the Ministry of Education, Culture, Sports, Science and Technology.

### Disclosures

None.

## References

- Dominiczak AF, Negrin DC, Clark JS, Brosnan MJ, McBride MW, Alexander MY. Genes and hypertension: from gene mapping in experimental models to vascular gene transfer strategies. *Hypertension*. 2000; 35:164–172.
- Colhoun H. Confirmation needed for genes for hypertension. *Lancet*. 1999;353:1200–1201.
- Johnson GL, Dhanasekaran N. The G-protein family and their interaction with receptors. *Endocr Rev*. 1989;10:317–331.
- McFarland KC, Sprengel R, Phillips HS, Kohler M, Roseblit N, Nikolic K, Segaloff DL, Seeburg PH. Lutropin-choriogonadotropin receptor: an unusual member of the G protein-coupled receptor family. *Science*. 1989;245:494–499.
- Parmentier M, Libert F, Maenhaut C, Lefort A, Gerard C, Perret J, Van Sande J, Dumont JE, Vassart G. Molecular cloning of the thyrotropin receptor. *Science*. 1989;246:1620–1622.
- Sprengel R, Braun T, Nikolic K, Segaloff DL, Seeburg PH. The testicular receptor for follicle stimulating hormone: structure and functional expression of cloned cDNA. *Mol Endocrinol*. 1990;4:525–530.
- Minegishi T, Nakamura K, Takakura Y, Ibuki Y, Igarashi M. Cloning and sequencing of human FSH receptor cDNA. *Biochem Biophys Res Commun*. 1991;175:1125–1130.
- Gromoll J, Dankbar B, Gudermann T. Characterization of the 5' flanking region of the human follicle-stimulating hormone receptor gene. *Mol Cell Endocrinol*. 1994;102:93–102.
- Aittomaki K, Lucena JL, Pakarinen P, Sistonen P, Tapanainen J, Gromoll J, Kaskikari R, Sankila EM, Lehvaslaiho H, Engel AR, Nieschlag E, Huhtaniemi I, de la Chapelle A. Mutation in the follicle-stimulating hormone receptor gene causes hereditary hypergonadotropic ovarian failure. *Cell*. 1995;82:959–968.
- Tapanainen JS, Aittomaki K, Min J, Vaskivuo T, Huhtaniemi IT. Men homozygous for an inactivating mutation of the follicle-stimulating hormone (FSH) receptor gene present variable suppression of spermatogenesis and fertility. *Nat Genet*. 1997;15:205–206.
- Dierich A, Sairam MR, Monaco L, Fimia GM, Gansmuller A, LeMour M, Sassone-Corsi P. Impairing follicle-stimulating hormone (FSH) signaling in vivo: Targeted disruption of the FSH receptor leads to aberrant gametogenesis and hormonal imbalance. *Proc Natl Acad Sci U S A*. 1998;95:13612–13617.
- Danilovich N, Harada N, Sairam MR, Maysinger D. Age-related neurodegenerative changes in the central nervous system of estrogen-deficient follitropin receptor knockout mice. *Exp Neurol*. 2003;183:559–572.
- Grover A, Sairam MR, Smith CE, Hermo L. Structural and functional modifications of sertoli cells in the testis of adult follicle-stimulating hormone receptor knockout mice. *Biol Reprod*. 2004;71:117–129.
- Javeshghani D, Touyz RM, Sairam MR, Virdis A, Neves MF, Schifffrin EL. Attenuated responses to angiotensin II in follitropin receptor knockout mice, a model of menopause-associated hypertension. *Hypertension*. 2003;42:761–767.
- Nakayama T, Soma M, Sato N, Haketa A, Kosuge K, Aoi N, Sato M, Izumi Y, Matsumoto K, Kanmatsuse K, Kokubun S. An association study in essential hypertension using functional polymorphisms in lymphotoxin- $\alpha$  gene. *Am J Hypertens*. 2004;17:1045–1049.
- Sano M, Kuroi N, Nakayama T, Sato N, Izumi Y, Soma M, Kokubun S. The association study of calcitonin-receptor-like receptor gene in essential hypertension. *Am J Hypertens*. 2005;18:403–408.
- Nakayama T, Soma M, Takahashi Y, Rehemudula D, Kanmatsuse K, Furuya K. Functional deletion mutation of the 5'-flanking region of the type A human natriuretic peptide receptor gene and its association with essential hypertension and left ventricular hypertrophy in the Japanese. *Circ Res*. 2000;86:841–845.
- Saavedra I, Leon J, Prado J, Sanchez MP, Lopez F, Gaete L. A comparative pharmacokinetic study of micronized estradiol valerate administered alone and in combination with medroxyprogesterone acetate in postmenopausal women. *Ther Drug Monit*. 2004;26:482–485.
- Hashiya N, Jo N, Aoki M, Matsumoto K, Nakamura T, Sato Y, Ogata N, Ogihara T, Kaneda Y, Morishita R. In vivo evidence of angiogenesis induced by transcription factor Ets-1: Ets-1 is located upstream of angiogenesis cascade. *Circulation*. 2004;109:3035–3041.
- Sakaguchi H, Fujimoto J, Aoki I, Toyoki H, Sato E, Tamaya T. Expression of E26 transformation specific (ETS-1) related to angiogenesis in ovarian endometriosis. *Fertil Steril*. 2004;82:507–510.
- Aldrighi JM, Alecrin IN, Caldas MA, Gebara OC, Ramires JA, Rosano GM. Effects of estradiol on myocardial global performance index in hypertensive postmenopausal women. *Gynecol Endocrinol*. 2004;19:282–292.
- Fak AS, Erenus M, Tezcan H, Caymaz O, Oktay S, Oktay A. Effects of a single dose of oral estrogen on left ventricular diastolic function in hypertensive postmenopausal women with diastolic dysfunction. *Fertil Steril*. 2000;73:66–71.
- Freimer NB, Sabatti C. Guidelines for association studies in Human Molecular Genetics. *Hum Mol Genet*. 2005;14:2481–2483.
- O'Donnell CJ, Lindpaintner K, Larson MG, Rao VS, Ordovas JM, Schaefer EJ, Myers RH, Levy D. Evidence for association and genetic linkage of the angiotensin-converting enzyme locus with hypertension and blood pressure in men but not women in the Framingham Heart Study. *Circulation*. 1998;97:1766–1772.
- Ono K, Mamami T, Iwai N. Association of a promoter variant of the haeme oxygenase-1 gene with hypertension in women. *J Hypertens*. 2003;21:1497–1503.
- Mein CA, Caulfield MJ, Munroe PB. Selection of candidate genes in hypertension. *Methods Mol Med*. 2005;108:107–129.
- Krushkal J, Ferrell R, Mockrin SC, Turner ST, Sing CF, Boerwinkle E. Genome-wide linkage analyses of systolic blood pressure using highly discordant siblings. *Circulation*. 1999;99:1407–1410.
- Rice T, Rankinen T, Province MA, Chagnon YC, Perusse L, Borecki IB, Bouchard C, Rao DC. Genome-wide linkage analysis of systolic and diastolic blood pressure: the Quebec Family Study. *Circulation*. 2000;102:1956–1963.
- Atwood LD, Samollow PB, Hixson JE, Stern MP, MacCluer JW. Genome-wide linkage analysis of blood pressure in Mexican Americans. *Genet Epidemiol*. 2001;20:373–382.
- Rice T, Rankinen T, Chagnon YC, Province MA, Perusse L, Leon AS, Skinner JS, Wilmore JH, Bouchard C, Rao DC. Genomewide linkage scan of resting blood pressure: HERITAGE Family Study. Health, risk factors, exercise training, and genetics. *Hypertension*. 2002;39:1037–1043.
- Angius A, Petretto E, Maestrone GB, Forabosco P, Casu G, Piras D, Fanciulli M, Falchi M, Melis PM, Palermo M, Pirastu M. A new essential hypertension susceptibility locus on chromosome 2p24–p25, detected by genomewide search. *Am J Hum Genet*. 2002;71:893–905.
- Rao DC, Province MA, Leppert MF, Oberman A, Heiss G, Ellison RC, Arnett DK, Eckfeldt JH, Schwander K, Mockrin SC, Hunt SC; HyperGEN Network. A genome-wide affected sibpair linkage analysis of hypertension: the HyperGEN network. *Am J Hypertens*. 2003;16:148–150.



ARTICLE

# An intronic variable number of tandem repeat polymorphisms of the *cold-induced autoinflammatory syndrome 1 (CIAS1)* gene modifies gene expression and is associated with essential hypertension

Toshinori Omi<sup>1</sup>, Maki Kumada<sup>1</sup>, Toyomi Kamesaki<sup>1</sup>, Hiroshi Okuda<sup>1</sup>, Lkhagvasuren Munkhtulga<sup>1</sup>, Yoshiko Yanagisawa<sup>1</sup>, Nanami Utsumi<sup>1</sup>, Takaya Gotoh<sup>1</sup>, Akira Hata<sup>2</sup>, Masayoshi Soma<sup>3</sup>, Satoshi Umemura<sup>4</sup>, Toshio Ogihara<sup>5</sup>, Norio Takahashi<sup>6</sup>, Yasuharu Tabara<sup>7</sup>, Kazuyuki Shimada<sup>8</sup>, Hiroyuki Mano<sup>9</sup>, Eiji Kajii<sup>1</sup>, Tetsuro Miki<sup>7</sup> and Sadahiko Iwamoto<sup>\*,1</sup>

<sup>1</sup>Division of Human Genetics, Center for Community Medicine, Jichi Medical School, Tochigi, Japan; <sup>2</sup>Department of Public Health, Graduate School of Medicine, Chiba University, Chiba, Japan; <sup>3</sup>Department of Nephrology and Endocrinology, Advanced Medical Research Center, Nihon University School of Medicine, Tokyo, Japan; <sup>4</sup>Second Department of Internal Medicine, Yokohama City University Medical School, Kanagawa, Japan; <sup>5</sup>Department of Genetics, Radiation Effects Research Foundation, Hiroshima, Japan; <sup>6</sup>Department of Geriatric Medicine, Osaka University Graduate School of Medicine, Osaka, Japan; <sup>7</sup>Department of Geriatric Medicine, Ehime University School of Medicine, Ehime, Japan; <sup>8</sup>Division of Cardiovascular Medicine, Department of Medicine, Jichi Medical School, Tochigi, Japan; <sup>9</sup>Division of Functional Genomics, Jichi Medical School, Kawachigun, Tochigi, Japan

*Cold-induced autoinflammatory syndrome 1 (CIAS1)* gene is a member of the NALP subfamily of the CATERPILLER protein family that is expressed predominantly in peripheral blood leukocytes, which is to regulate apoptosis or inflammation through the activation of NF- $\kappa$ B and caspase. Recent genetic analyses suggested an association between inflammation and oxidative stress-related genes in the development of hypertension. This is the first genetic study indicating an association between the *CIAS1* gene and susceptibility to essential hypertension (EH). The frequency of subject with the homozygote of 12 repeat allele was significantly higher in patients with hypertension compared with control subjects (987 cases, 924 controls) ( $P=0.030$ ; odds ratio = 1.24) at a novel VNTR polymorphism of *CIAS1* intron 4 loci. We also found that the mean of systolic blood pressure of homozygotes of 12 repeat allele was 6.4 mmHg higher than those of homozygotes of non-12 repeat allele in male random population ( $P=0.009$ ). The frequency of six SNPs spanning of the *CIAS1* gene was not significantly between patients and controls. The real-time PCR analysis showed that among healthy young adults, 12-12 subjects expressed *CIAS1* mRNA in peripheral leukocytes significantly more abundantly than homozygote of non-12 repeat alleles subjects ( $P<0.05$ ). Reporter gene assay of the *CIAS1*-VNTR in HL60 stimulated by lipopolysaccharides showed that the intronic sequence involving 12 repeat increased the expression of luciferase compared with 9, 7, and 6 repeats.

\*Correspondence: Professor S Iwamoto, Division of Human Genetics, Center for Community, Medicine, Jichi Medical School, Minamikawachi-machi, Tochigi 329-0498, Japan. Tel: 81 285 44 2111; Fax: 81 285 44 4902; E-mail: siwamoto@jichi.ac.jp  
Received 6 October 2005; revised 29 May 2006; accepted 16 June 2006

Thus, we propose here the *CIAS1* is associated with EH through the dominant expression of transcripts, which may depend on the *CIAS1*-VNTR genotype.

*European Journal of Human Genetics* (2006) 0, 000–000. doi:10.1038/sj.ejhg.5201698

**Keywords:** association study; *CIAS1*; hypertension; VNTR polymorphis

Essential hypertension (EH [MIN1455]) is a complex disease that is thought to be influenced by polygenic and multiple environmental factors. Blood pressure is known to be a heritable trait, but the inherited genetic factors appear to have an influence on blood pressure and hypertension only ~50% of the time. Recently susceptibility loci for EH and a healthy variation in blood pressure have been mapped on several human chromosomes by genome-wide scans.<sup>1–9</sup> Although several dozen candidate genes for hypertension have been proposed, understanding the genetic influences on this disease is difficult because, when combining data from different studies, there is a lack of replication of an identified locus.

Recent biochemical and morphological studies suggest that remodeling of large and small arteries is involved in the development and associated complications of hypertension.<sup>10,11</sup> In the past decade, inflammation and oxidative stress have emerged as major players in vascular remodeling of hypertension in addition to known mechanisms, such as salt sensitivity, insulin resistance, and an imbalance in the renin-angiotensin system.<sup>12–15</sup> Polymorphonuclear leukocytes (PMNL) are the main producer of reactive oxygen species (ROS) and have been shown to be involved in hypertension in both human and animal models. There is evidence supporting the presence of activated PMNL in humans and rats with hypertension. In a hypertension model using Sabra hypertension-prone rats, superoxide release from PMNL and inflammation anteceded the onset of hypertension (brought on by salt loading).<sup>16,17</sup> Results indicated that genetic factors contributed to the activation of PMNL and production of ROS. The p22phox is a major component of NADPH oxidase and is essential for O<sup>2-</sup> generation. A polymorphism in the promoter region of the p22phox gene, –930A/G, has been shown to be associated with EH.<sup>18</sup> The present study was designed to identify a novel hypertensive-susceptibility locus using a single-candidate-gene approach on a gene involved in the activation of PMNL.

The CATERPILLER family gene (CARD, transcription enhancer, R (purine)-binding, pyrin, lots of leucine protein family that is expressed predominantly in peripheral blood leukocytes<sup>19–25</sup> in human. In addition, the rat orthologue of a member in CATERPILLER family gene, PYPAF5 (PYRIN-containing apoptotic protease-activating factor -1-like proteins 5), encodes angiotensin II/vasopressin dual receptor as a splicing isoform product<sup>26</sup> and the variation cosegregate with salt-sensitive hypertension in Dahl salt-

sensitive hypertension rat model.<sup>27</sup> Then, we focused the relation between hypertension and the CATERPILLER family gene. The *CIAS1* (MIN 606416; *cold-induced autoinflammatory syndrome 1*) gene mapped 1q44<sup>19</sup> encodes cryopyrin/NALP3/PYPAF1 (NACHT-LRR-PYD-containing protein-3/PYRIN-containing Apaf1-like protein-1) proteins, a member of the NALP subfamily of the CATERPILLER. The function of the proteins within the subfamily is thought to involve the regulation of apoptosis or inflammation through the activation of (nuclear factor) (NF)- $\kappa$ B and caspase. Thus far, many missense mutations in the *CIAS1* gene have been identified in hereditary systemic autoinflammatory disorders.<sup>28</sup> However, an association between the *CIAS1* gene and hypertension has not been reported. We used a case-control study using a single-candidate-gene approach to determine whether the *CIAS1* gene was associated with hypertension.

## Methods

### Subjects for genotyping

Population of panels 1 (100 cases and 109 controls) and 2 (987 cases and 924 controls) for case-control study were from the Tochigi prefecture, Ehime prefecture, Hiroshima prefecture, Osaka, or Kanagawa prefecture, Tokyo, Hokkaido in Japan (the study group from the Group of Millennium Genome Project for Hypertension). All subjects gave informed consent, and the study was approved by the ethics committee of Jichi Medical School. The hypertension group included patients who were diagnosed before 60 years of age; hypertension was defined as a blood pressure (systolic/diastolic) of 160/100 mmHg or higher. We excluded hypertensive individuals who had diabetes, intrinsic renal disease, a self-reported history of secondary hypertension that was corroborated by a family physician, or coexisting illness. We aimed to recruit hypertensive individuals with a body mass index (BMI) less than 25 to exclude the patient including obesity. In addition, the hypertensive individuals were selected with at least one first-degree family history because we considered that the patients with family history much received the genetic influence of hypertension compare to the patients without family history. Control subjects after 50 years included those with a blood pressure of 120/80 mmHg or lower and without hypertensive first-degree relatives. Blood pressure was measured with a mercury sphygmomanometer at least three times in a sitting position in clinic.

Panel 3 from random population (268 men and 162 women) was collected from the individual who came a medical check up at the Health Examination Center in Jichi Medical School. Genotypes were compared with phenotypes; age, BMI, systolic blood pressure (SBP), diastolic blood pressure (DBP), total-cholesterol, high-density lipoprotein (HDL), triglycerides, fasting blood sugar (FBS), and glycosylated hemoglobin (HbA1c).

#### Genotyping of SNPs and a VNTR in the *CIAS1* gene

Four single nucleotide polymorphisms (SNPs) (IMS-JST084196, IMS-JST165517, IMS-JST165520, and IMS-JST084195) selected from seven SNPs registered in the Japanese SNP database,<sup>20</sup> a SNP (rs1539019) of NCBI and a SNP (HCV26052025) of commercial database (Celera Discovery System) were selected to cover the *CIAS1* gene. Each SNP was genotyped by the Taq Man-PCR assay using the ABI PRISM 7900-HT according to the manufacture's instructions (Applied Biosystems, Foster, CA, USA). For an association study, we also used a novel variable number of tandem repeat polymorphisms (VNTR) polymorphism in intron 4 of the *CIAS1* gene that was found through resequencing. To genotype the VNTR polymorphisms, standard PCR was performed in a final reaction volume of 25  $\mu$ l of 1  $\times$  PCR buffer containing 20 ng of the genomic DNA, 0.4  $\mu$ M of a set of primer (forward: 5'-CTGACCTCCCAATGTGCCTT-3' and reverse: 5'-CAGAGCTTCTTCAGATTGCA-3'), 0.2 mM concentrations of each dNTP, and 2.5 U *Taq* DNA polymerase (Roche Diagnostics, GmbH, Mannheim, Germany), using the following temperature profile: 95°C for 3 min, followed by 35 cycles of 95°C for 1 min, 55°C for 1 min, 72°C for 1 min, and a final extension of 72°C for 7 min. The PCR products were electrophoresed on a 3% agarose gel and stained with ethidium bromide.

#### Association between VNTR polymorphisms and gene expression *in vivo*

Total RNA from leukocytes was isolated from the blood of 46 healthy young adults with different genotypes by the Aquapure RNA blood kit (Bio-Rad, CA, USA). The 46 RNA consisted of 21 homozygotes of major repeats (12, 16 heterozygotes of major and minor repeats (9, 7, and 6 repeats), and nine homozygotes of minor repeats. We further analyzed the expression of *CIAS1* in their primary monocytes after lipopolysaccharide (LPS) stimulation. Monocytes were isolated from peripheral leukocytes by standard Ficoll centrifugation procedure followed by adhesion on plastic dishes coated by auto-serum. The attached cells were stimulated with 200 ng/ml LPS (LPS 026:B6; Sigma-Aldrich, St Louis, MO) for 30 or 60 min at 37°C and lysed in Trizol reagent (Life Technologies). For real-time PCR, approximately 100 ng total RNA from each cell was first reverse transcribed (final volume of 12.5  $\mu$ l) into cDNA using the SuperScript II (Life Technologies, Gaithersburg, MD) according to the manufacture's proto-

col. RT product (0.2  $\mu$ l) was used for quantification of gene expression by TaqMan real-time RT-PCR (Applied Biosystems, Foster City, CA, USA) in a reaction volume of 5  $\mu$ l. The cumulative fluorescence for each amplicon was normalized to that seen with VIC-labeled GAPDH (Applied Biosystems, Foster, CA, USA) amplification. A FAM-labeled Hs00918086 probe (Applied Biosystems, Foster, CA, USA) was used for *CIAS1* mRNA amplification.

#### *In vitro* expression study by reporter gene assay of individual VNTR polymorphisms in HL60

We constructed expression vectors using the pGL3 basic (Promega, Madison, WI) and pCAGGS vector. The four alleles (12, 9, 7, and 6 repeats unit) of *CIAS1* 42bp-VNTR were individually amplified from DNA of the subjects encoding the alleles using *NheI*- and *NcoI*-tailed primers (GATTACCATGGGCTAGCAATGTGCCTTGTG, and AGCACCCATGGTGGCGGCCACAGCGCCCCA), and the PCR products were digested by *NheI/NcoI* and ligated into the pGL3 basic vector. After verification of the insert VNTR fragment by sequencing, VNTR-luciferase DNA fragment was extracted from the pGL3 vector by *NheI/XbaI* digestion and inserted in the pCAGGS vector at the *XbaI* site to mimic the splicing acceptor position. Ten micrograms of the constructs were transfected into  $5 \times 10^6$  HL60 cells with 0.2  $\mu$ g of pRL vector (renilla luciferase, Promega) by electroporation procedure using GenePulser II (Bio-Rad Laboratories, Hercules, CA, USA). At 24 h after transfection, the relative light units (firefly/renilla light units) were measured by a dual luciferase reporter assay system (Promega) in a TD-20/20 luminometer (Turner Designs, Sunnyvale, CA, USA). For all transfection assays, at least three independent experiments were performed.


#### Statistical analysis

Values are expressed as mean  $\pm$  standard error of mean (SEM). Values of  $P < 0.05$  were considered statistically significant. Data were analyzed using a contingency table analysis and the Student's *t*-test. Hardy-Weinberg equilibrium was calculated by the  $\chi^2$  test. Statistical analysis included a comparison of the proportions of the prevalence of the allele in cases and controls. Odds ratios (ORs) were generated from two-by-two tables, and statistical significance was assessed using the Fisher's exact.  $D'$  and  $r^2$  values were analyzed to evaluate a linkage disequilibrium (LD). Statistical analysis was performed using the SNPalyze (Dynacom, Mobar, Japan) and SPSS statistical package (SPSS Japan, Tokyo).

#### Results

##### Identification and characterization of a novel VNTR in the *CIAS1* locus

A novel VNTR polymorphism was found in the process of *CIAS1* resequencing, which was identified near the 3' end



**Figure 1** (a) Schema of *CIAS1* gene with position of six SNPs and an identified VNTR polymorphisms for the genetic association study. Four SNPs were selected from the IMS-JST (Institute of Medical Science-Japan Science and Technology Agency) Japanese SNP database (IMS-JST): SNP1 (IMS-JST084196), SNP2 (IMS-JST165517), SNP3 (IMS-JST165520), and SNP4 (IMS-JST084195). SNP5 (rs1539019) and SNP6 (hCV26052025) were selected from a NCBI database and a commercial database (Celera Discovery System), respectively. (b) Nine genotypes of a novel VNTR polymorphism in *CIAS1*, a:12-12, b:12-9, c:12-7, d:12-6, e:9-7, f:9-6, g:7-7, h:7-6, and i:6-6 were detected in this study as shown in the agarose gel electrophoresis. M is 100 bp molecular ladder marker. (c) The nucleotide sequences of the 12 repeat allele of the *CIAS1*-42 bp VNTR. The tandem repeat number in each allele is 12 (GenBank Ac. No AB120959), 9 (GenBank Ac. No AB120960), 7 (GenBank Ac. No AB120961), and 6 (GenBank Ac. No AB120962).

of *CIAS1* intron 4, -527 to -24 bp from the splicing acceptor site of *CIAS1* exon 5 (Figure 1a). The PCR analysis identified nine genotypes derived from four alleles (Figure 1b). Each allele consisted of the 720, 594, 510, and 468 bp nucleotide sequences, respectively. The multiple alignment analysis between each allele showed the existence of the 42 bp tandem repeat sequences in the *CIAS1* gene (*CIAS1* 42 bp-VNTR). The nucleotide sequences of the 12 repeat allele was shown in Figure 1c. This tandem repeat polymorphism consisted of the 12, 9, 7, and 6 repeats units, respectively (GenBank accession number AB120959 to AB120962). The 12 repeat alleles' largest band was similar to the published sequence of the *Homo sapiens* chromosome 1 clone RP11-433K2 in GenBank (accession number AC104335). Estimation of the allele frequencies showed that the 12, 9, 7, and 6 repeat units was 0.577, 0.008, 0.248, and 0.167 in the *CIAS1* 42 bp-VNTR locus ( $n=507$ ). The percentage of the all genotypes was 32.9% (12-12), 1% (12-9), 30% (12-7), 18.5% (12-6), 0% (9-9),

0.2% (9-7), 0.4% (9-6), 6.5% (7-7), 6.5% (7-6), and 4% (6-6), respectively. The observed genotype frequencies of all polymorphisms did not deviate from the predictions of the Hardy-Weinberg equation.

To assess whether the *CIAS1* locus is associated with hypertension, the *CIAS1* 42 bp-VNTR was genotyped for 109 control subjects and 100 patients with EH in panel 1. The frequency of subject with the homozygote of 12 repeat allele was significantly higher in patients with hypertension compared with control subjects ( $P=0.04$ ; odds ratio = 1.84) (Table 1). The allele frequency of 12 repeat units, major allele, also tended to increase in patients compared with controls.

#### Single-locus analyses and case-control haplotype analyses of the *CIAS1* gene

To reveal whether the *CIAS1* 42 bp-VNTR is in tight linkage with other polymorphisms, six SNPs were chosen from the *CIAS1* gene using a public database (JSNP, <http://>

**Table 1** Genotyping data and association test of DNA polymorphisms on the *CIAS1* in panel 1 (100 cases and 109 controls)

	SNP1	SNP2	SNP3	SNP4	VNTR <sup>a</sup>	SNP5	SNP6
<b>Patients</b>							
Major allele	0.830	0.585	0.585	0.825	0.635	0.560	0.510
Minor allele	0.170	0.415	0.415	0.175	0.365	0.440	0.490
Major homozygous	68	31	31	67	40	28	22
Heterozygous	30	55	55	31	47	56	58
Minor homozygous	2	14	14	2	13	16	20
<b>Controls</b>							
Major allele	0.830	0.638	0.638	0.812	0.525	0.555	0.561
Minor allele	0.170	0.362	0.362	0.178	0.475	0.445	0.439
Major homozygous	74	47	47	72	29	37	35
Heterozygous	33	45	45	33	56	47	52
Minor homozygous	2	17	17	4	24	25	22
$\chi^2$ (P)							
Major homozygous vs others	0.0003 (0.986)	3.274 (0.070)	3.274 (0.070)	0.021 (0.885)	4.231 (0.039)	0.860 (0.353)	2.687 (0.101)
Minor homozygous vs others	0.007 (0.930)	0.105 (0.746)	0.105 (0.746)	0.521 (0.470)	2.91 (0.087)	1.59 (0.207)	0.001 (0.974)

\* $P < 0.05$ .

<sup>a</sup>Major allele is 12 repeat and minor alleles are 9, 7 and 6 repeat.

**Table 2** Frequency of the major haploype constructed from SNPs 1 to 6 between 109 controls and 100 cases

Haplotypes	Controls (%)	Case (%)	P
CTACGT	32.9	31.1	NS
CTACTC	16.2	13.3	NS
CCGCTC	13.1	17.5	NS
CTACTT	12.9	10.1	NS
ACGTTC	9.6	11.2	NS
ACGTGT	5.6	2.7	NS
CCGCGT	2.9	4.7	NS
CCGTTC	1.8	2.0	NS
CCGCGC	1.3	5.9	NS

NS: not significance.

**Table 3** Pairwise LD in *CIAS1* gene, evaluated by  $D'$  and  $r^2$  from panel 1

	SNP1	SNP2	SNP3	SNP4	VNTR	SNP5	SNP6
SNP1							
SNP2   $D'$	0.940						
$r^2$	0.286						
SNP3   $D'$	0.940	1.000					
$r^2$	0.286	1.000					
SNP4   $D'$	0.912	1.000	1.000				
$r^2$	0.765	0.351	0.351				
VNTR   $D'$	0.715	0.145	0.145	0.841			
$r^2$	0.142	0.082	0.082	0.214			
SNP5   $D'$	0.294	0.405	0.405	0.348	0.185		
$r^2$	0.014	0.082	0.082	0.021	0.020		
SNP6   $D'$	0.475	0.534	0.534	0.501	0.094	0.810	
$r^2$	0.053	0.209	0.209	0.066	0.007	0.452	

www.snp.ims.u-tokyo.ac.jp, Figure 1a)<sup>29</sup> for genotyping in panel 1 subjects. In the VNTR loci, the subjects were categorized into three genotypes: major allele homozygote (12-12), heterozygote with major and minor alleles (12-9, 12-7, and 12-6), and minor allele homozygote (9-9, 9-7, 9-6, 7-6, and 7-7), respectively. As a results, no significant association was shown between cases and controls in the six SNPs of the *CIAS1* gene (Table 1).

Table 2 showed that the difference of haplotype frequency between the cases and controls. Our set of SNPs generated 10 common haplotypes (frequency > 0.01) constructed from SNPs 1 to 6, and there were no differences in haplotype frequencies between cases ( $n = 100$ ) and controls ( $n = 109$ ). We also analyzed the pairwise linkage disequilibrium (LD) by calculating  $D'$  and  $r^2$  using genotype results (Table 3);  $|D'| > 0.5$  and  $r^2 > 0.5$  were regarded as tight linkage. Among the six SNPs, SNPs 1 (5' flanking region) to 4 (intron 4) tended to be in tight linkage but the LD was not distributed across the *CIAS1* gene. The LD and case-

control haplotype analysis of the *CIAS1* gene suggest that the VNTR locus was possibly independent from other SNPs and a functional polymorphism associated with hypertension.

#### Case-control study of the *CIAS1* 42bp-VNTR for hypertension in a large sample

To confirm the possibility that the homozygote of 12 repeat allele is a primary variant that determines susceptibility to hypertension, genotyping of *CIAS1* 42bp-VNTR was performed in panel 2 (987 cases and 924 controls). The results confirmed the statistical significance between the *CIAS1* 42bp-VNTR and EH (Table 4). The significant difference was  $P = 0.030$  (Odds 1.24) in the comparison of 12-12 vs 12-X + X-X and  $P = 0.006$  (Odds 1.39) in 12-12 + 12-X vs X-X. These results led us the homozygote of 12 repeat alleles was an association with risk of EH. The

**Table 4** Association of the CIAS1-VNTR genotype with essential hypertension in panel 2

Panel 2	Controls	Patients	$\chi^2$	P	Odds
<i>Genotype distribution</i>					
12-12	278 (30.1%)	343 (34.7%)			
12-X	466 (50.4%)	498 (50.5%)			
X-X	180 (19.5%)	146 (14.8%)			
<i>Allele frequency</i>					
12	0.553	0.600			
X	0.447	0.400			
<i>Genotype comparison</i>					
12-12 vs 12-X+X-X			4.735	0.030	1.24
12-12+12-X vs X-X			7.413	0.006	1.39

**Table 5** Clinical characterization of random population of panel 3 to CIAS1 42 bp-polymorphism

Panel 3 (Men and Women)	12-12	Genotype 12-X	X-X	P
N (%)	138 (32.1%)	218 (50.7%)	74 (17.2%)	NS
Age	51.4±8.7	52.1±9.5	53.5±8.6	NS
Height (cm)	163.0±6.8	163.2±8.8	163.3±8.5	NS
Weight (kg)	63.7±9.1	63.1±11.2	62.5±10.2	NS
BMI (kg/m <sup>2</sup> )	24.0±2.7	23.8±3.4	23.3±2.6	NS
SBP (mmHg)	127.7±14.2	124.3±16.4	122.4±13.3	0.018*
DBP (mmHg)	78.6±9.7	77.6±11.1	76.3±9.8	NS
Total-cholesterol (mg/dl)	205.2±34.4	205.6±35.8	198.6±32.3	NS
HDL (mg/dl)	61.8±13.9	60.9±18.2	61.9±14.3	NS
Triglyceride (mg/dl)	109.3±53.8	103.0±51.7	96.2±43.1	NS
Fasting blood glucose (mg/dl)	100.5±8.7	98.9±20.2	101.4±15.8	NS
HbA1c (%)	5.0±0.3	5.1±0.4	5.1±0.4	NS

NS; not significant.

T-Cho, total cholesterol; HDL-Cho, HDL cholesterol; TG, triglyceride, FBS, Fasting blood sugar; and HbA1c, glycohemoglobin, A-1-C.

\*12-12 vs 12-X+X-X.

frequency of the 12-12 subjects was 0.348 and 0.301 in panel 2. The most significant association was shown in the comparison of 12-12 vs X-X ( $\chi^2 = 13.048$ ,  $P = 0.0003$ , odds ratio = 1.60) (data not shown). These genetic results indicate that the *CIAS1* locus contributes to the susceptibility to EH.

#### Association between *CIAS1* 42 bp-VNTR genotype and random population in blood pressure

We further examined the association between the *CIAS1* gene and healthy variations in blood pressure. Panel 3 from random population (268 men and 162 women) was collected at the Health Examination Center in Jichi Medical School, which was independent of panels 1 and 2. Genotypes were compared with phenotypes; age, BMI, SBP, DBP, total-cholesterol, HDL, triglycerides, FBS, and HbA1c. Table 5 showed the genotype effect in three groups of 12-12, 12-X, and X-X to clinical characteristics in *CIAS1* gene. We found the significant difference of the SBP in three genotype groups by one-way ANOVA study in panel 3 ( $P = 0.018$ ). The association between the genotype and the

SBP was revealed in men ( $P = 0.028$ , Table 6) but not in women (Table 7). The mean SBP in 12-12 subjects ( $n = 86$ ) was 6.4 mmHg higher than those of X-X subjects ( $n = 44$ ) in men. The *CIAS1* 42 bp-VNTR genotype was not associated with BMI, DBP, total-cholesterol, HDL, triglycerides, fasting blood glucose, and HbA1c levels. To exclude the influence of age on BP or population stratification among generation, age was put into covariate and analysis of covariance test (ANCOVA) was carried out. After this correction significant difference of SBP was observed again among three genotypes in men ( $P = 0.028$ ), and the significance became distinct ( $P = 0.011$ ), when over 55 years old men were analyzed. In 162 women, we did not find a significant difference of blood pressure among the three groups in ANOVA or ANCOVA study even after age correction, but the mean SBP of 12-12 subjects was slightly higher than that of other groups (Table 7).

#### Genotype and the expression level of *CIAS1* transcript

The transcript from *CIAS1* was predominantly expressed in peripheral blood leukocytes compared with total RNA from

**Table 6** Clinical characterization of random population of panel 3 to CIAS1 42 bp-polymorphism in Men

Panel 3 (Men)	Genotype			P
	12-12	12-X	X-X	
N (%)	86 (32.1%)	138 (51.5%)	44 (16.4%)	NS
Age	53.5±8.5	52.6±9.3	54.1±7.2	NS
Height (cm)	167.0±6.8	167.9.1±6.1	168.7±5.1	NS
Weight (kg)	67.7±9.3	68.4±8.6	67.2±8.6	NS
BMI (kg/m <sup>2</sup> )	24.2±2.7	24.2±2.3	23.6±2.7	NS
SBP (mmHg)	131.7±14.2	126.8±16.0	125.4±13.6	0.028
DBP (mmHg)	80.6±9.7	79.7±11.1	79.0±9.6	NS
Total-cholesterol (mg/dl)	207.3±34.4	206.4±35.1	197.9±32.9	NS
HDL (mg/dl)	58.7±13.9	55.8±13.3	56.8±13.9	NS
Triglyceride (mg/dl)	127.6±53.8	119.1±53.7	112.2±44.9	NS
Fasting blood glucose (mg/dl)	104.2±8.7	103.0±11.5	102.8±11.7	NS
HbA1c (%)	5.0±0.3	5.1±0.4	5.0±0.6	NS

NS; not significant.

T-Cho, total cholesterol; HDL-Cho, HDL cholesterol; TG, triglyceride, FBS, Fasting blood sugar; and HbA1c, glycohemoglobin, A-1-C.

**Table 7** Clinical characterization of random population of panel 3 to CIAS1 42 bp-polymorphism in Women

Panel 3 (Women)	Genotype			P
	12-12	12-X	X-X	
N (%)	52 (32.1%)	80 (49.4%)	30 (18.5%)	NS
Age	47.9±7.8	51.7±9.8	49.5±7.8	NS
Height (cm)	156.3±5.115	5.1±6.7	155±5.7	NS
Weight (kg)	57.1±7.8	54.0±9.1	55.5±8.1	NS
BMI (kg/m <sup>2</sup> )	23.5±3.5	23.1±4.7	22.9±2.4	NS
SBP (mmHg)	121.0±13.5	120.1±16.4	118.1±11.5	NS
DBP (mmHg)	75.3±8.8	74.0±10.2	72.5±8.7	NS
Total-cholesterol (mg/dl)	201.8±27.5	204.2±37.0	199.8±31.4	NS
HDL (mg/dl)	67.0±15.6	67.9±15.6	69.2±11.3	NS
Triglyceride (mg/dl)	79.2±49.0	75.6±33.5	72.3±26.5	NS
Fasting blood glucose (mg/dl)	94.3±10.3	92.0±28.5	99.4±20.2	NS
HbA1c (%)	5.0±0.4	5.1±0.3	5.1±0.4	NS

NS; not significant.


T-Cho, total cholesterol; HDL-Cho, HDL cholesterol; TG, triglyceride, FBS, Fasting blood sugar; and HbA1c, glycohemoglobin, A-1-C.

bone marrow, brain, fetal liver heart, kidney, lung, placenta, prostate, salivary gland, skeletal muscle, spleen, testis, thyroid gland, trachea, uterus, colon, small intestine, spinal cord, and stomach packaged in the master panel (BD Biosciences, CA, USA) by RT-PCR analysis (data not shown). We next tested the genotype effect of CIAS1 42bp-VNTR locus to the expression level of CIAS1 transcript using the total leukocyte RNA extracted from blood samples in healthy young volunteers with different genotypes. Real-time PCR analysis showed that the expression level of the CIAS1 transcript was different in subjects with 12-12 (*n*=21), 12-X (*n*=16), and X-X (*n*=9) VNTR genotypes (Figure 2). The CIAS1 transcript was expressed significantly more in the 12-12 subjects compared with subjects with the X-X genotype. Real-time PCR analysis showed that the expression level of the CIAS1 transcript showed significant difference in subjects with among three genotype groups (*P*=0.009) by One-way ANOVA tests. (Figure 2). Among the three genotype groups, the CIAS1

transcript from 12-12 subjects was much expressed than that of 12-X and X-X subjects.

#### Expression of the transcript from CIAS1 locus in monocytes after LPS stimulation

The CIAS1 expression level in total leukocytes showed remarkably high SD in both 12-12 and 12-X subjects (Figure 2), which was estimated to be dependent on the activated state of the leukocytes. To rule-out the possibility that the deviations simply resulted from the condition of donors or preparation procedure of leukocytes, the transcript level was also analyzed in monocytes under stimulation of LPS. As shown in Figure 3, transcript level in monocytes before stimulation of 12-12 subjects was higher than other genotype groups. One-way ANOVA tests showed the expression level of CIAS1 mRNA on monocyte was the significant difference on 30 min (*P*=0.01) and 60 min (*P*=0.0001) after LPS stimulation. These results




**Figure 2** The *CIAS1* mRNA levels in peripheral leukocytes among the three genotype groups of 12-12 (homozygote of 12 repeat alleles), 12-X (heterozygote of 12 and non-12 repeat alleles), and X-X (homozygote of non-12 repeat alleles) in *CIAS1* 42bp-VNTR. The *CIAS1* expression was analyzed by real-time PCR. The relative amount of *CIAS1* mRNA normalized by GAPDH is shown as fold volume. n: number of subjects. One-way ANOVA tests showed that expression level of the *CIAS1* transcript was significant difference in subjects among three genotype groups ( $P=0.009$ ).

indicated that the 12 repeat allele had an additive effect on the expression level of this gene.

### Functional assay of *CIAS1* VNTR in HL 60

To investigate whether *CIAS1* 42 bp-VNTR directly affected gene expression, we next examined the reporter gene assay. We constructed the pCAGGS-based expression plasmid, containing the *CIAS1* 42 bp-VNTR alleles (12, 9, 7, and 6 repeat units) linked to the luciferase gene at the splicing acceptor site (Figure 4a). The pCAGGS promoter activity of the 12 VNTR repeat construct was higher than other repeat constructs before LPS stimulation in HL60 cells, although the difference was not significant (Figure 4b). The difference, however, became significant after 1 or 2 h of LPS stimulation. These results may confirmed the additive effect of 12 repeat allele on the expression level of *CIAS1* gene (Figure 3) in the functional assay.




**Figure 3** The effect of stimulation of LPS on *CIAS1* mRNA levels in monocytes in different genotypes. Monocytes were isolated from peripheral leukocytes by Ficoll centrifugation followed by adhesion on a plastic dish. Cells were treated with 200 ng/ml LPS for 30 or 60 min and lysed. The relative amount of *CIAS1* mRNA normalized by GAPDH is shown. Black bars show the transcripts with X-X genotype (homozygote of non-12 repeat alleles), hatched bars indicate the transcripts with 12-X genotype (heterozygote of 12 repeat allele and non-12 alleles), and gray bars show the transcript with 12-12 genotype (homozygote of 12 repeat allele). One-way ANOVA tests showed the expression level of *CIAS1* mRNA on monocyte was the significant difference on 30 min ( $P=0.01$ ) and 60 min ( $P=0.0001$ ) after LPS stimulation. Three individuals with each genotypes were tested.

### Discussion

A single-candidate-gene approach for hypertension focusing on the CATERPILLER protein family has not been demonstrated. This is the first genetic study indicating an association between the *CIAS1* gene and susceptibility to EH. Our study of 987 cases and 924 controls showed that subjects with 12-12 genotype of the *CIAS1* 42bp-VNTR experienced hypertension more frequently than controls, with a significant association ( $P=0.006$ ). Furthermore, the





**Figure 4** Functional assay of VNTR in HL60 cells. (a) The genomic region including the VNTR repeats, 12 (720 bp), 9 (594 bp), 7 (510 bp), and 6 (468 bp), was inserted in pGL3 vector. The VNTR-luciferase inserts were directionally subcloned in the pCAGGS vector. (b) HL60 cells were co-transfected with the pRE vector (renilla luciferase) by electroporation procedure. At 24 h after the transfection, HL60 cells were stimulated by LPS. The relative light units (firefly/renilla light units) were measured by dual luciferase reporter assay system. PCAGGS-VNTR-12 (●), vector PCAGGS-VNTR-9 (○), PCAGGS-VNTR-7 (■), and PCAGGS-VNTR-6 (□). The relative luciferase activity with 12 repeat allele was significantly enhanced compared with the activity with 9, 7, and 6 repeat until more than 2 h after LPS.

comparison with genotype and clinical data in a random population showed that SBP was significantly higher in 12-12 subjects compared with 12-X and X-X subjects, although there was no difference in other variables. These results lead us to question whether the *CIAS1* 42 bp-VNTR in intron is a true functional polymorphism in hypertension. We showed that the transcript level of *CIAS1* in peripheral leukocytes and monocytes of healthy young adults was more abundant in 12-12 subjects than in those with the other genotypes. Furthermore, our reporter assay for *CIAS1* 42 bp-VNTR showed that the 12 repeat units

increased the expression of luciferase in the *CIAS1* gene. These results may indicate that the *CIAS1* 42 bp-VNTR directly affects gene expression. Thus, we believe that increased expression of *CIAS1* in leukocytes dependent on the *CIAS1* 42 bp-VNTR length may contribute to the development of EH.

*CIAS1* is a member of the recently discovered NALP/PYPAF subfamily of the CATERPILLER protein family, thought to function in apoptotic and inflammatory signaling pathways.<sup>27</sup> *CIAS1*/NALP3/PYPAF1 protein is composed of three domains; the N-terminal Pyrin domain, the central a specific nucleotide binding (NACHT) domain, and the C-terminal tandem copy of Leucin-rich repeats (LRRs). A large number of nucleotide substitutions have been found in the NACHT domain<sup>19</sup> in patients with autosomal-dominant autoinflammatory disorders; Muckle-Wells syndrome, familial cold urticaria, and chronic infantile neurologic cutaneous and articular syndrome.<sup>19,27</sup> It has been proposed that the missense mutation in the NACHT domain weakens the self-regulation via its interaction with the LRRs and enables spontaneous activation by releasing NACHT domain from LRRs. The activated form recruits an apoptosis-associated speck-like protein containing a CARD (ASC) and Cardinal (TUCAN, CARD8, NDDP1). The caspase recruit domain (CARD) of ASC and Cardinal interact with that of pro-caspase-1. The resulting macrocomplex, called inflammasome, is available to convert pro-IL-1 $\beta$  into matured IL-1 $\beta$ . Moreover, ASC and Cardinal in the inflammasome have been shown to be associated with IKK (I $\kappa$ B kinase) complex and activate NF- $\kappa$ B. The phenotypic heterogeneity (cold sensitivity, sensorineural hearing loss, severe neurologic involvement) has been observed in autoinflammatory syndromes. Although, the phenotypic heterogeneity has been mainly attributed to the mutation position in the NACHT domain, the amount of mutant *CIAS1* mRNA might also contribute to the development of the syndrome.

The physiological function of the wild type *CIAS1* is, however, still under investigation. Opposite results about NF- $\kappa$ B activation have been reported from overexpression analysis of *CIAS1* in culture cell lines. NF- $\kappa$ B was activated by the overexpression of *CIAS1* and ASC in HEK293 cells.<sup>21</sup> The *CIAS1* suppressed the activation in HeLa cells after TNF stimulation and suggested that a molecular balance of *CIAS1* with ASC is critical for the inflammatory response.<sup>23,24</sup> Recently Coas1-knockout mice are developed in two independent research groups. Mariathasan *et al* showed that Cryopyrin produced from *CIAS1* activates the inflammasome in response to toxins and ATP.<sup>30</sup> Kanne-ganti *et al* reports that bacteria RNA and small antiviral compounds activate caspase 1 through cryopyrin produced from *CIAS1* gene.<sup>31</sup> We propose here the *CIAS1* is associated with hypertension through the dominant expression of transcripts, which may depend on the

CIAS1-VNTR genotype. To prove this hypothesis, an animal model overexpressing CIAS1 in leukocytes is desirable, which may also resolve the matter of whether CIAS1 alone activates or suppresses NF- $\kappa$ B signaling.

A recent study showed that chronic inflammation is crucial for the development of hypertension. Plasma C-reactive protein has been shown to be an independent risk factor for hypertension,<sup>32</sup> and PMNL count in peripheral blood and increased superoxide release contributes to hypertension.<sup>33</sup> In the animal model, the activation of NF- $\kappa$ B is increased in numerous tissues in stroke-prone spontaneous hypertension (SHRsp) rats compared with Sprague-Dawley (SD) rats.<sup>34</sup> NF- $\kappa$ B promotes proinflammatory genes or induces apoptosis in many organs. In leukocytes, NF- $\kappa$ B can induce iNOS, cell adhesion molecules, IL-6, and other cytokines and chemokines.<sup>35</sup> The induced inflammation generates large amount of oxidative agents, and the oxidative stress activates NF- $\kappa$ B.<sup>36,37</sup> Thus, we predict that CIAS1 might play an important role in the development of hypertension based on inflammation brought on by NF- $\kappa$ B activation.

#### Acknowledgements

The study was supported by a grant in aid (No.13204075) from Japanese Ministry of Education, Culture, Sports, Science and Technology. We thank Oyamada T and Nagashima K for technical assistance. We thank Drs Kario, Hoshida S, Sugimoto K, Ishikawa K, Sakamoto A, (Jichi Medical School), Sekiguchi T, and Hara K (Nasu minami) Ohishi T (Ohishi medical clinic) for sample correction.

#### References

- Hsueh WC, Mitchell BD, Schneider JL *et al*: QTL influencing blood pressure maps to the region of PPH1 on chromosome 2q31–34 in Old Order Amish. *Circulation* 2000; 101: 2810–2816.
- Levy D, DeStefano AL, Larson MG *et al*: Evidence for a gene influencing blood pressure on chromosome 17. Genome scan linkage results for longitudinal blood pressure phenotypes in subjects from the Framingham Heart Study. *Hypertension* 2000; 36: 477–483.
- Perola M, Kainulainen K, Pajukanta P *et al*: Genome-wide scan of predisposing loci for increased diastolic blood pressure in Finnish siblings. *J Hypertens* 2000; 18: 1579–1585.
- Rice T, Rankinen T, Province MA *et al*: Genome-wide linkage analysis of systolic and diastolic blood pressure: the Quebec Family Study. *Circulation* 2000; 102: 1956–1963.
- Atwood LD, Samollow PB, Hixson JE, Stern MP, MacCluer JW: Genome-wide linkage analysis of blood pressure in Mexican Americans. *Genet Epidemiol* 2001; 20: 373–382.
- Angius A, Petretto E, Maestrale GB *et al*: A new essential hypertension susceptibility locus on chromosome 2p24–p25, detected by genomewide search. *Am J Hum Genet* 2002; 71: 893–905.
- Caulfield M, Munroe P, Pembroke J *et al*: Genome-wide mapping of human locus for essential hypertension. *Lancet* 2003; 361: 2118–2123.
- Rice T, Rankinen T, Chagnon YC *et al*: Genomewide linkage scan of resting blood pressure: HERITAGE family study. Health, risk factors, exercise training, and genetics. *Hypertension* 2002; 39: 1037–1043.
- de Lange M, Spector TD, Andrew T: Genome-wide scan for blood pressure suggests linkage to chromosome 11, and replication of locus on 16, 17, and 22. *Hypertension* 2004; 44: 872–877.
- Intengan HD, Schiffrin EL: Vascular remodeling in hypertension: roles of apoptosis, inflammation, and fibrosis. *Hypertension* 2001; 38: 581–587. Review.
- Rodriguez-Iturbe B, Vaziri ND, Herrera-Acosta J, Johnson RJ: Oxidative stress, renal infiltration of immune cells, and salt-sensitive hypertension: all for one and one for all. *Am J Physiol Renal Physiol* 2004; 286: F606–F616. Review.
- Russo C, Olivieri O, Girelli D *et al*: Anti-oxidant status and lipid peroxidation in patients with essential hypertension. *J Hypertens* 1998; 16: 1267–1271.
- Berry C, Brosnan MJ, Fennell J, Hamilton CA, Dominiczak AF: Oxidative stress and vascular damage in hypertension. *Curr Opin Nephrol Hypertens* 2001; 10: 247–255.
- Romeo JC, Reckelhoff JF: Role of angiotensin and oxidative stress in arterial hypertension. *Hypertension* 1999; 34: 943–949.
- Wilcox CS: Reactive oxygen species: roles in blood pressure and kidney function. *Curr Hypertens Rep* 2002; 4: 160–166. Review.
- Moreno MU, San Jose G, Orbe J *et al*: Preliminary characterisation of the promoter of the human p22(phox) gene: identification of a new polymorphism associated with hypertension. *FEBS Lett* 2003; 542: 27–31.
- Sela S, Mazor R, Amsalam M, Yagil C, Yagil Y, Kristal B: Primed polymorphonuclear leukocytes, oxidative stress, and inflammation antecede hypertension in the Sabra rat. *Hypertension* 2004; 44: 764–769.
- San Jose G, Moreno MU, Olivan S *et al*: Functional effect of the p22phox -930A/G polymorphism on p22phox expression and NADPH oxidase activity in hypertension. *Hypertension* 2004; 44: 163–169.
- Hoffman HM, Mueller JL, Broide DH, Wanderer AA, Kolodner RD: Mutation of a new gene encoding a putative pyrin-like protein causes familial cold autoinflammatory syndrome and Muckle-Wells syndrome. *Nat Genet* 2001; 29: 301–305.
- Mao M, Fu G, Wu JS *et al*: Identification of genes expressed in human CD34+ hematopoietic stem/progenitor cells by expressed sequence tags and efficient full-length cDNA cloning. *Proc Nat Acad Sci* 1998; 95: 8175–8180.
- Manji GA, Wang L, Geddes BJ *et al*: PYPAF1, a PYRIN-containing Apaf1-like protein that assembles with ASC and regulates activation of NF-kappa B. *J Biol Chem* 2002; 277: 11570–11575.
- Koonin EV, Aravind L: The NACHT family – a new group of predicted NTPases implicated in apoptosis and MHC transcription activation. *Trends Biochem Sci* 2000; 25: 223–224.
- Harton JA, Linhoff MW, Zhang J, Ting JP: Cutting edge: CATERPILLER: a large family of mammalian genes containing CARD, pyrin, nucleotide-binding, and leucine-rich repeat domains. *J Immunol* 2002; 169: 4088–4093.
- O'Connor Jr W, Harton JA, Zhu X, Linhoff MW, Ting JP: Cutting edge: CIAS1/cryopyrin/PYPAF1/NALP3/CATERPILLER 1.1 is an inducible inflammatory mediator with NF-kappa B suppressive properties. *J Immunol* 2003; 171: 6329–6333.
- Neven B, Callebaut I, Prieur AM *et al*: Molecular basis of the spectral expression of CIAS1 mutations associated with phagocytic cell-mediated autoinflammatory disorders CINCA/NOMID, MWS, and FCU. *Blood* 2004; 103: 2809–2815.
- Albrecht M, Domingues FS, Schreiber S, Lengauer T: Structural localization of disease-associated sequence variations in the NACHT and LRR domains of PYPAF1 and NOD2. *FEBS Lett* 2003; 554: 520–528.
- Ruiz-Opazo N, Lopez LV, Herrera VL: The dual AngII/AVP receptor gene N119S/C163R variant exhibits sodium-induced dysfunction and cosegregates with salt-sensitive hypertension in the Dahl salt-sensitive hypertensive rat model. *Mol Med* 2002; 8: 24–32.
- Touitou I, Lesage S, McDermott M *et al*: Infervers: an evolving mutation database for auto-inflammatory syndromes. *Hum Mutat* 2004; 24: 194–198.

- 29 Hirakawa M, Tanaka T, Hashimoto Y, Kuroda M, Takagi T, Nakamura Y: JSNP: a database of common gene variations in the Japanese population. *Nucleic Acids Res* 2002; 30: 158–162.
- 30 Bautista LE, Vera LM, Arenas IA, Gamarra G: Independent association between inflammatory markers (C-reactive protein, interleukin-6, and TNF-alpha) and essential hypertension. *J Hum Hypertens* 2005; 19: 149–154.
- 31 Mariathasan S, Weiss DS, Newton K *et al*: Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 2006 (published online Jan 11 2006).
- 32 Kanneganti TD, Ozoren N, Body-Malapel M *et al*: Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature* 2006 (published online Jan 11 2006).
- 33 Kristal B, Shurtz-Swirski R, Chezdar J *et al*: Participation of peripheral polymorphonuclear leukocytes in the oxidative stress and inflammation in patients with essential hypertension. *Am J Hypertens* 1998; 11: 921–928.
- 34 Wu L, Noyan Ashraf MH, Facci M *et al*: Dietary approach to attenuate oxidative stress, hypertension, and inflammation in the cardiovascular system. *Proc Natl Acad Sci USA* 2004; 101: 7094–7099.
- 35 Cooke CL, Davidge ST: Peroxynitrite increases iNOS through NF-kappaB and decreases prostacyclin synthase in endothelial cells. *Am J Physiol Cell Physiol* 2002; 282: C395–C402.
- 36 Sanlioglu S, Williams CM, Samavati L *et al*: Lipopolysaccharide induces Rac1-dependent reactive oxygen species formation and coordinates tumor necrosis factor-alpha secretion through IKK regulation of NF-kappa B. *J Biol Chem* 2001; 276: 30188–30198.
- 37 Asehnoune K, Strassheim D, Mitra S, Kim JY, Abraham E: Involvement of reactive oxygen species in Toll-like receptor 4-dependent activation of NF-kappa B. *J Immunol* 2004; 172: 2522–2529.
- 38 Kanneganti TD, Ozoren N, Body-Malapel M *et al*.

## Appendix

The Study Group of Millennium Genome Project for Hypertension:

Chairman: Tetsuro Miki (Department of Geriatric Medicine, Ehime University School of Medicine, Ehime)

Michio Yasunami, Akinori Kimura (Department of Molecular Pathogenesis, Medical Research Inst., Tokyo Medical and Dental University), Akira Hata (Chiba University Graduate School of Medicine, Chiba), Toshio Saruta

(Department of Internal Medicine, School of Medicine, Keio University), M Yokota (Department of Clinical Pathophysiology, Nagoya University, Graduate School of Medicine), Tomohiro Katsuya, Toshio Ogihara (Department of Geriatric Medicine, Osaka University Graduate School of Medicine), Hirotsugu Ueshima (Department of Health Science, Shiga University of Medical Science), Katsuhiko Kohara, Yasuharu Tabara, Jun Nakura, Tetsuro Miki (Department of Geriatric Medicine, Ehime University School of Medicine), Sumio Sugano (Human Genome Center, Institute of Medical Science, University of Tokyo), Masayoshi Soma, Tomohiro Nakayama (Second Department of Internal Medicine, and Division of Receptor Biology, Advanced Medical Research Center, Nihon University School of Medicine), Norihiro Kato (Department of Gene Diagnostics and Therapeutics, Research Institute, International Medical Center of Japan), Ichiro Kishimoto, Kazuwa Nakao (Department of Medicine and Clinical Science, Kyoto University Graduate School of Medicine), Yuhei Kawano, Kei Kamide (Division of Hypertension and Nephrology, Department of Medicine, National Cardiovascular Center), Naoharu Iwai, Toshiyuki Miyata (Research Institute, National Cardiovascular Center), Takayuki Morisaki (Department of Bioscience, National Cardiovascular Center Research Institute), Hitonobu Tomoike (Division of Preventive Cardiology, National Cardiovascular Center), Tsutomu Yamazaki (Department of Pharmacoepidemiology, Graduate School of Medicine, Faculty of Medicine, University of Tokyo), Katsushi Tokunaga (Graduate School of Medicine Faculty, International Health, International Biomedical Sciences, University of Tokyo), Satoshi Umemura, Nobuhito Hirawa (Department of Medicine II, Yokohama City University School of Medicine), Hiroyuki Mano (Division of Functional Genomics, Jichi Medical School), Sadahiko Iwamoto (Division of Human Genetics, Center for Community Medicine, Jichi Medical School).

## Original Article

# Relationship between Metabolic Syndrome and Trp64Arg Polymorphism of the $\beta$ 3-Adrenergic Receptor Gene in a General Sample: The Shigaraki Study

Shinji TAMAKI<sup>1)</sup>, Yasuyuki NAKAMURA<sup>2)</sup>, Yasuharu TABARA<sup>3)</sup>, Tomonori OKAMURA<sup>4)</sup>,  
Yoshikuni KITA<sup>4)</sup>, Takashi KADOWAKI<sup>4)</sup>, Yasuyuki TSUJITA<sup>5)</sup>, Minoru HORIE<sup>5)</sup>,  
Tetsuro MIKI<sup>6)</sup>, and Hirotsugu UESHIMA<sup>4)</sup>

It has been reported that the  $\beta$ 3-adrenergic receptor gene (*ADRB3*) is associated with abnormal metabolic risk factors. Therefore, we examined whether the Trp64Arg polymorphism of *ADRB3* affects the occurrence of metabolic syndrome (MS). The participants were 2,395 subjects who underwent a medical examination in Shigaraki in Shiga, Japan. Among them, 1,416 subjects who gave informed consent for genetic analysis and were not receiving treatment for hypertension, diabetes, or hyperlipidemia were enrolled in this study. MS was diagnosed in 86 (16.0%) of 537 men, and 8 (0.9%) of 879 women. There was no significant relationship between *ADRB3* polymorphism and the frequency of MS. Multiple logistic regression analysis including smoking, sex, and age as confounding factors showed no interaction between MS and *ADRB3* polymorphism (odds ratio: 0.94; 95% confidence interval: 0.59–1.49;  $p=0.78$ ). Subjects were also analyzed according to differences in the number of abnormal metabolic risk factors. However, there was no significant relationship between *ADRB3* polymorphism and the number of such factors. In conclusion, in a general sample, the frequency of MS was 16.0% in men, and 0.9% in women. There was no relationship between *ADRB3* polymorphism and MS. (*Hypertens Res* 2006; 29: 891–896)

**Key Words:**  $\beta$ 3-adrenergic receptor gene, Trp64Arg polymorphism, metabolic syndrome, hypertension

## Introduction

The  $\beta$ 3-adrenergic receptor is expressed in visceral adipose tissue and is thought to contribute to lipolysis, energy metab-

olism and regulation of the metabolic rate (1–3). A missense mutation of *ADRB3* (Trp64Arg) in the first intracellular loop of the receptor has been reported. This mutation has been reported to be associated with diabetes mellitus (4, 5), abdominal obesity (5), obesity (6), insulin resistance (5) and the

From the <sup>1)</sup>Division of Cardiology, Department of Medicine, Kohka Public Hospital, Kohka, Japan; <sup>2)</sup>Cardiovascular Epidemiology, Kyoto Women's University, Kyoto, Japan; <sup>3)</sup>Division of EcoGenetics, Department of Environmental Health and Social Medicine and <sup>4)</sup>Department of Geriatric Medicine, Ehime University School of Medicine, Ehime, Japan; and <sup>5)</sup>Department of Health Science and <sup>6)</sup>Department of Cardiovascular and Respiratory Medicine, Shiga University of Medical Science, Otsu, Japan.

This study represents joint research of Shiga University of Medical Science and Ehime University. The study was supported in part by the Japanese Ministry of Education, Culture, Sports, Science and Technology (Grant-in-Aid for Scientific Research on Priority Areas (C): 12204059; Grant-in-Aid for Scientific Research (B): 02454211; Grant-in-Aid for Scientific Research (C): 06670414) and the Japan Society for the Promotion of Science (Grant-in-Aid for Scientific Research (C): 16590500).

Address for Reprints: Yasuyuki Nakamura, M.D., Ph.D., Cardiovascular Epidemiology, Kyoto Women's University, Imakumano, Kitahiyoshi-cho, Higashiyama-ku, Kyoto 605–8501, Japan. E-mail: nakamury@kyoto-wu.ac.jp

Received April 26, 2006; Accepted in revised form July 28, 2006.