

basal metabolic rate (4).

In Japan, Kadowaki *et al.* (7) reported that the frequency of the mutant allele in obese subjects was significantly higher than that in non-obese subjects. The presence of this mutation was also accompanied by significantly higher fasting and 2-h serum insulin levels during an oral glucose tolerance test (OGTT). Thus, they suggested that *ADRB3* might contribute to obesity and hyperinsulinemia/insulin resistance in Japanese subjects.

On the other hand, there have been several reports that the Trp64Arg mutation of *ADRB3* was not associated with glucose metabolism, non-insulin-dependent diabetes mellitus (NIDDM) (8), insulin resistance syndrome (8), obesity (9) or insulin sensitivity (10). We also reported that the Trp64Arg polymorphism of *ADRB3* did not appear to have any pathophysiological significance in Japanese samples (11).

People with metabolic syndrome (MS) are at high risk of developing type 2 diabetes (12) and cardiovascular disease (13). The Third Report of the National Cholesterol Education Program Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (NCEP-ATP III) proposed a new definition of MS (14). In 2005, the International Diabetes Federation (IDF) (the IDF consensus worldwide definition of the metabolic syndrome; available from: <http://www.idf.org/home/index.cfm?unode=1120071E-AACE-41D2-9FA0-BAB6E25BA072>) and a group of eight Japanese societies (15) announced new and separate diagnostic criteria for MS.

In our previous study employing a group of hypertensive patients, the prevalence of MS by the new Japanese criteria was 44.0% in men and 23.3% in women (16).

However, there have been no reports about the correlation between the new Japanese criteria of MS and *ADRB3* polymorphism. Therefore, in this study, we evaluated the relationship between the new Japanese criteria of MS and the Trp64Arg polymorphism of *ADRB3* in Japanese residents in a rural community.

## Methods

### Study Population

The Shigaraki Study was based on a medical examination undertaken in 1999 in Shigaraki, a farming community near Kyoto, western Japan (17–22). A total of 2,902 subjects underwent the examination, of whom 2,395 were enrolled in this genetic study after receiving a full explanation and providing informed consent. Of these subjects, 403 were excluded for the following reasons: undetermined genotype,  $n=127$ ; a serum aspartate aminotransferase (AST [GOT]) or alanine aminotransferase (ALT [GPT]) level of over 100 IU/l,  $n=13$ ; and/or a history of transient ischemic attack, stroke, angina pectoris, or myocardial infarction,  $n=263$ . Moreover, 576 (221 men, and 355 women) were excluded because they were receiving treatment for hypertension, dia-

betes or hyperlipidemia. The remaining 1,416 subjects (537 men and 879 women) were analyzed. Subjects were between the ages of 30 and 79. This study was approved by the Institutional Review Board of Shiga University of Medical Science (Nos. 11-15, 1999).

### Measurement of Waist Circumference

The waist was measured at the level of the umbilicus using a tape measure; subjects wore an undergarment and stood during the measurement. Waist measurements were taken by the same person. Each measurement was performed in triplicate and the average value was used.

### Blood Pressure and Biochemical Examinations

Systolic blood pressure (SBP) and diastolic blood pressure (DBP) were measured twice using a standard sphygmomanometer on the right arm while the subject was seated after having rested for at least 5 min. Korotkov's first and fifth points were regarded as the SBP and DBP, respectively, and the blood pressure (BP) was measured by a well-trained nurse. The mean of 2 measurements from each subject was used for data analysis. The body mass index (BMI) was calculated as the weight (kg) divided by the square of the height (m). Non-fasting blood was drawn and serum total cholesterol (TC) and high density lipoprotein-cholesterol (HDL-C) levels were determined in one laboratory (Medic, Shiga, Japan). The measurement precision and accuracy of these serum lipids were certified through the lipid standardization program by Osaka Medical Center for Cancer and Cardiovascular Disease, which is a member of the Cholesterol Reference Method Laboratory Network (CRMLN) controlled by the centers for Disease Control and Prevention (Atlanta, USA) (23).

### Criteria of Metabolic Syndrome

We used the 2005 Japanese criteria of MS (15). Briefly, men with an abdominal circumference of  $\geq 85$  cm and women with an abdominal circumference of  $\geq 90$  cm were defined as abdominally obese. MS is defined as abdominal obesity plus at least 2 of the following abnormalities: hypertriglyceridemia (triglyceride [TG]  $\geq 150$  mg/dl), low HDL-C (HDL-C  $< 40$  mg/dl), hypertension (SBP  $\geq 130$  mmHg and/or DBP  $\geq 85$  mmHg), and impaired fasting glucose (fasting blood glucose level  $\geq 110$  mg/dl).

### Determination of the Trp64Arg Polymorphism of *ADRB3*

DNA was isolated from peripheral leukocytes and *ADRB3* genotypes were determined as previously reported (1). Genotypes, determined by the polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) method for 75 random samples consisting of 25 PCR products for each

**Table 1. Characteristics of the Study Population According to the Trp64Arg Polymorphism of *ADRB3* in Men and Women, Shigaraki Study in 1999 (Not Receiving Treatment for Hypertension, Diabetes or Hyperlipidemia)**

Risk characteristics	Polymorphism of <i>ADRB3</i>					
	Men (537)			Women (879)		
	Trp/Trp	Trp/Arg + Arg/Arg	<i>p</i> -value	Trp/Trp	Trp/Arg + Arg/Arg	<i>p</i> -value
<i>N</i> (1,416)	368	157+12		585	257+37	
Age (years)	55.5±15.5	53.6±16.1	0.1921	51.3±15.2	53.3±15.9	0.0789
BMI (kg/m <sup>2</sup> )	22.4±2.8	22.2±3.0	0.3827	21.9±3.0	22.1±3.0	0.4327
Waist (cm)	80.2±8.3	79.4±9.0	0.3100	70.5±7.5	71.6±8.3	0.0404
Abdominally obese (men waist ≥ 85 cm, women waist ≥ 90 cm)	109/259 (29.6%)	33+5/124+7 (22.5%)	0.0851	6/579 (1.0%)	6+0/251+37 (2.0%)	0.2211
Alcohol (P/N)	282/86	115+9/42+3	0.4143	157/428	75+10/182+27	0.5160
Smoking (P/N)	279/89	119+10/38+2	0.8966	48/537	24+3/233+34	0.6242
130/85 (P/N)	167/201	73+7/84+5	0.6726	110/475	51+5/206+32	0.9305
TG/HDL (P/N)	145/223	63+5/94+7	0.8543	205/380	93+15/164+22	0.6211
TG (mg/dl)	145.3±99.8	140.3±94.3	0.5815	106.2±62.0	108.6±65.9	0.6041
HDL (mg/dl)	52.8±14.3	55.0±15.1	0.1059	61.9±14.1	60.4±13.8	0.1276
BG ≥ 110 (P/N)	97/271	38+6/119+6	0.9370	118/467	58+6/199+31	0.5812
MS (P/N)	61/307 (16.6%)	20+5/137+7 (14.8%)	0.6008	4/581 (0.7%)	4+0/253+37 (1.4%)	0.3188

*N*: number of subjects. Values are the means±SD. *ADRB3*, β<sub>3</sub> adrenergic receptor gene; BMI, body mass index; Waist, waist circumference; Alcohol, alcohol drinking habit; P/N, positive/negative; Smoking, smoking habit; 130/85, systolic blood pressure ≥ 130 mmHg and/or diastolic blood pressure ≥ 85 mmHg; TG/HDL, triglyceride ≥ 150 mg/dl and/or high density lipoprotein cholesterol < 40 mg/dl; BG ≥ 110, fasting blood glucose level ≥ 110 mg/dl; MS, frequency of metabolic syndrome.

genotype, were confirmed by direct sequencing. Briefly, after fractionation of PCR-RFLP products on 1% agarose gels (Nippon Gene, Toyama, Japan), the desired DNA bands were excised, and DNA was purified using a QIAquick Gel Extraction Kit (QIAGEN, Valencia, USA), amplified with the above 5' primer, and analyzed with an ABI PRISM 310 Genetic Analyzer (Perkin-Elmer, Norwalk, USA).

**Statistical Analysis**

Data are expressed as the mean±SD. All statistical analyses were performed using the SAS statistical package. Two-way ANOVA and the χ<sup>2</sup> test were used to assess differences among subjects with different genotypes. Since few subjects carried Arg/Arg genotypes, we combined the Trp/Arg and Arg/Arg genotypes for assessment.

To examine the independent contribution of *ADRB3* polymorphism to the risk of MS, while adjusting for the effects of other clinical characteristics, logistic analysis was used. For the risk of MS, a logistic model was developed with the following variables as covariates: sex, age, smoking, alcohol, and *ADRB3* polymorphisms.

**Results**

Table 1 summarizes the characteristics of subjects according to the genotype of *ADRB3*. The Trp64Arg polymorphism of *ADRB3* was not significantly associated with age, BMI, waist, drinking and smoking habits, tendency toward hyper-

tension, high TG and/or low HDL-C, blood glucose (BG) over 110 mg/dl, or the frequency of MS. The waist circumference of subjects with the Arg64 allele of Trp64Arg was significantly larger than that of subjects without the Arg64 allele (*p*=0.0404). In the assessment of genotypes not associated with the Trp64Arg polymorphism, the mean waist circumferences of subjects were 70.5±7.5 cm for subjects with Trp/Trp, 71.8±8.4 for subjects with Trp/Arg, and 70.6±7.6 for subjects with Arg/Arg (*p*=0.0835). There was no significant association between the waist circumference and *ADRB3* polymorphism.

The frequency of MS was 16.0% in men and 0.9% in women. There was no significant relationship between *ADRB3* polymorphism and the frequency of MS.

Multiple logistic regression analysis including smoking, sex, and age as confounding factors showed that each of these factors was associated with MS; however, there was no interaction between MS and *ADRB3* polymorphism (odds ratio: 0.94; 95% confidence interval: 0.59–1.49; *p*=0.78).

Tables 2 and 3 summarize the correlation between the Trp64Arg polymorphism of *ADRB3* and the number of abnormal metabolic risk factors in men and women, respectively. There was no relationship between *ADRB3* polymorphism and the number of abnormal metabolic factors in either sex (*p*=0.925 for men; *p*=0.765 for women).

**Discussion**

Kadowaki et al. (7) and Yoshida et al. (24) reported that the

**Table 2. Relationship between the Trp64Arg Polymorphism of *ADRB3* and the Number of Abnormal Metabolic Factors in Men (Not Receiving Treatment for Hypertension, Diabetes or Hyperlipidemia)**

Number of abnormal risk factors	Polymorphism of <i>ADRB3</i>		N	
	Trp/Trp	Trp/Arg+Arg/Arg		
0	102 (68.0%)	44+4 (32.0%)	150	<i>p</i> =0.925
1	150 (70.1%)	63+1 (29.9%)	214	
2	89 (67.4%)	39+4 (32.6%)	132	
3	27 (65.9%)	11+3 (34.1%)	41	
	368	157+12	537	

*N*: number of subjects. Abnormal metabolic factors: hypertriglyceridemia: triglycerides  $\geq 150$  mg/dl and/or high density lipoprotein cholesterol  $< 40$  mg/dl; hypertension: systolic blood pressure  $\geq 130$  mmHg and/or diastolic blood pressure  $\geq 85$  mmHg; and impaired fasting glucose: fasting blood glucose level  $\geq 110$  mg/dl.

**Table 3. Relationship between the Trp64Arg Polymorphism of *ADRB3* Gene and the Number of Abnormal Metabolic Factors in Women (Not Receiving Treatment for Hypertension, Diabetes or Hyperlipidemia)**

Number of abnormal risk factors	Polymorphism of the <i>ADRB3</i>		N	
	Trp/Trp	Trp/Arg+Arg/Arg		
0	325 (67.7%)	136+19 (32.3%)	480	<i>p</i> =0.765
1	144 (65.2%)	65+12 (34.8%)	221	
2	59 (62.8%)	31+4 (37.2%)	94	
3	57 (67.9%)	25+2 (32.1%)	84	
	585	257+37	879	

*N*: number of subjects. Abnormal metabolic factors: hypertriglyceridemia: triglycerides  $\geq 150$  mg/dl and/or high density lipoprotein cholesterol  $< 40$  mg/dl; hypertension: systolic blood pressure  $\geq 130$  mmHg and/or diastolic blood pressure  $\geq 85$  mmHg; and impaired fasting glucose: fasting blood glucose level  $\geq 110$  mg/dl.

Trp64Arg mutation had pathological significance in Japanese populations. However, in our study of 1,416 subjects taking no medications for hypertension, diabetes or hyperlipidemia, the *ADRB3* polymorphism did not appear to have any significant influence on the various parameters associated with MS. Similarly, Yuan *et al.* (25) reported that *ADRB3* polymorphism was not a major genetic determinant of obesity or diabetes in the Japanese general population. Moreover, Masuo *et al.* (26) reported that the Trp64 allele was linked to BP elevation in a 5-year longitudinal study; however, our study indicated no significant relation between *ADRB3* polymorphism and BP. These discrepancies may be difficult to reconcile; however, it is possible that this polymorphism may have only a small effect on MS, and this may be masked by differences in the genetic background or environmental factors, including lifestyle. Another possibility is that the Trp64Arg polymorphism of *ADRB3* may not have any functional significance itself, but may be linked to an as-yet-unidentified mutation in this gene. Indeed, two previous studies have reported that expression of the Trp64Arg polymorphism of *ADRB3* *in vitro* was associated with normal lipolytic function (9, 27). However, Pietri-Rouxel *et al.* (28) reported a biochemical dysfunction induced by the Trp64Arg mutation in *ADRB3*, which could be important in pathophysiological situations pertaining to obesity. Indeed, although *ADRB3* exists on chromo-

some 8, Kissebah *et al.* (29) reported that trait loci on chromosomes 3 and 17 influence MS phenotypes. Thus the Trp64Arg polymorphism of *ADRB3* and its functional significance remain controversial.

Essential hypertension is reported to be associated with insulin resistance; however, Castellano *et al.* (30) reported that the Trp64Arg polymorphism of *ADRB3* has no relevant influence on BP levels or other features of insulin resistance syndrome. In Japan, Fujisawa *et al.* (31) reported that the Trp64Arg polymorphism of *ADRB3* did not play a major role in the susceptibility to essential hypertension or insulin resistance. In their study, as in our normal population in this study, the Trp64Arg polymorphism of *ADRB3* had no significant influence on various parameters associated with MS. Also, in their study, visceral obesity was reported to significantly affect glucose metabolism; however, the Trp64Arg polymorphism of *ADRB3* did not appear to have any significant effect on MS in this study. In the present study, it seemed that women with the Arg64 allele of Trp64Arg had a significantly larger waist circumference than those without the Arg64 allele (*p*=0.0404). However, there was no difference in waist circumference between subjects with and those without Trp64Arg polymorphism of *ADRB3* (*p*=0.0835).

In our sample, the frequency of MS was 16.0% in men and 0.9% in women. In the Tanno and Sobetsu Study, the fre-

quency of MS in a rural Japanese sample was 25.3% in men (32), and in a study in Okinawa, the frequency of MS was 30.2% in men and 10.3% in women (33). Our present data in men were comparable with the data from these previous studies, but our data in women were lower. This is possibly because the criteria for MS in the other studies were NCEP-ATP III and a waist girth over 85 cm. If we use a waist girth over 85 cm for women in this study, the frequency of MS was 5.8% in women in our sample. Compared with a previous study (16), the prevalence of MS in this study was much lower. This may be because our study subjects were not being treated, but the previous subjects were all being treated for hypertension. In addition, in the standard examination in this study, there was no relationship between *ADRB3* polymorphism and MS.

There are some limitations of our study. First, we only analyzed *ADRB3* as a candidate gene of MS. Although there are many other candidate genes of MS, such as the leptin, insulin receptor substrate-1, and adiponectin genes, these were not examined in this study. In particular, for adiponectin located at chromosome 3, Kissebah et al. (29) reported that trait loci on chromosomes 3 and 17 influenced MS phenotypes. Further investigation will need to be carried out using a large-scale sample. Second, not all the blood samples collected were fasting samples. If we use fasting blood collection, the frequency of MS will decrease and will not be related to MS. Third, the sample power was weak. A much larger number of subjects may be required to confirm the association.

In conclusion, in this general sample, the frequency of MS was 16.0% in men and 0.9% in women. There was no relationship between *ADRB3* polymorphism and MS.

### Acknowledgements

We received excellent technical assistance from the staff at Ehime University for the determination of the *ADRB3* polymorphism.

### References

- Emorine LJ, Marullo S, Briend-Sutren MM, et al: Molecular characterization of the human  $\beta 3$  adrenergic receptor. *Science* 1989; **245**: 1118–1121.
- Spronsen AV, Nahmias C, Krief S, Briend-Sutren MM, Strosberg AD, Emorine LJ: The promoter and intron/exon structure of the human and mouse  $\beta 3$  adrenergic receptor genes. *Eur J Biochem* 1993; **213**: 1117–1124.
- Krief S, Lonngqvist F, Raimbault S, et al: Tissue distribution of  $\beta 3$  adrenergic receptor m-RNA in man. *J Clin Invest* 1993; **91**: 344–349.
- Walston J, Silver K, Bogardus C, et al: Time of onset of non-insulin-dependent diabetes mellitus and genetic variation in the  $\beta 3$  adrenergic-receptor gene. *N Engl J Med* 1995; **333**: 343–347.
- Widen E, Lehto M, Kanninen T, Walston J, Shuldiner AR, Groop LC: Association of a polymorphism in the  $\beta 3$  adrenergic-receptor gene with features of the insulin resistance syndrome in Finns. *N Engl J Med* 1995; **333**: 348–351.
- Clement K, Vaisse C, Manning BS, et al: Genetic variation in the  $\beta 3$  adrenergic-receptor and an increased capacity to gain weight in patients with morbid obesity. *N Engl J Med* 1995; **333**: 352–354.
- Kadowaki H, Yasuda K, Iwamoto K, et al: A mutation in the  $\beta 3$  adrenergic-receptor gene is associated with obesity and hyperinsulinemia in Japanese subjects. *Biochem Biophys Res Commun* 1995; **215**: 555–560.
- Rissanen J, Kuopusjarvi J, Pihlajamaki J, et al: The Trp64Arg polymorphism of the  $\beta 3$  adrenergic receptor gene. *Diabetes Care* 1997; **20**: 1319–1322.
- Li LS, Lonngqvist F, Luthman H, Arner P: Phenotypic characterization of the Trp64Arg polymorphism in the beta3 adrenergic receptor gene in normal weight and obese. *Diabetologia* 1996; **39**: 857–860.
- Urhammer SA, Clausen JO, Hansen T, Pedersen O: Insulin sensitivity and body weight changes in young white carriers of the codon 64 amino acid polymorphism of the  $\beta 3$ -adrenergic receptor gene. *Diabetes* 1996; **45**: 1115–1120.
- Tamaki S, Iwai N, Tsujita Y, Nakamura Y, Ohmichi N, Kinoshita M: Variant of the  $\beta 3$ -adrenergic receptor gene and coronary atherosclerosis in Japanese subjects. *Int J Cardiol* 1999; **69**: 309–311.
- Lorenzo C, Okoloise M, Williams K, Stern MP, Haffner SM: The metabolic syndrome as predictor of type 2 diabetes: the San Antonio heart study. *Diabetes Care* 2003; **26**: 3153–3159.
- Sattar N, Gaw A, Scherbakova O, et al: Metabolic syndrome with and without C-reactive protein as a predictor of coronary heart disease and diabetes in the West of Scotland Coronary Prevention Study. *Circulation* 2003; **108**: 414–419.
- Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults: Executive Summary of the Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *JAMA* 2001; **285**: 2486–2497.
- The Examination Committee of Criteria for Metabolic Syndrome: Definition and criteria of metabolic syndrome. *J Jpn Soc Intern Med* 2005; **94**: 794–809 (in Japanese).
- Tamaki S, Nakamura Y, Yoshino T, et al: The association between morning hypertension and metabolic syndrome in hypertensive patients. *Hypertens Res* 2006; **29**: 783–788.
- Nakamura Y, Amamoto K, Tamaki S, et al: Genetic variation in aldehyde dehydrogenase 2 and the effect of alcohol consumption on cholesterol levels. *Atherosclerosis* 2002; **164**: 171–177.
- Amamoto K, Okamura T, Tamaki S, et al: Epidemiologic study of the association of low- $K_m$  mitochondrial acetaldehyde dehydrogenase genotypes with blood pressure level and the prevalence of hypertension in a general population. *Hypertens Res* 2002; **25**: 857–864.
- Tamaki S, Nakamura Y, Tsujita Y, et al: Polymorphism of the angiotensin converting enzyme gene and blood pressure in a Japanese general population (the Shigaraki Study). *Hypertens Res* 2002; **25**: 843–848.
- Tamaki S, Nakamura Y, Tabara Y, et al: Combined analysis of polymorphisms in angiotensinogen and adducin genes

- and their effect on hypertension in a Japanese population (the Shigaraki Study). *Hypertens Res* 2005; **28**: 645–650.
21. Tsujita Y, Nakamura Y, Zhang Q, *et al*: The association between high-density lipoprotein cholesterol level and cholesteryl ester transfer protein TaqIB gene polymorphism is influenced by alcohol drinking in a population-based sample. *Atherosclerosis* (in press).
  22. Ozawa T, Ito M, Tamaki S, *et al*: Gender and age effects on ventricular repolarization abnormality in Japanese general carriers of a G643S common single nucleotide polymorphism for the KCNQ1 gene. *Circ J* 2006; **70**: 645–650.
  23. Myers GL, Cooper GR, Henderson LO, Hassemer DJ, Kimberly MM: Standardization of lipid and lipoprotein measurement, in Rifai N, Warnick GR, Dominiczak MH (eds): *Handbook of Lipoprotein Testing*. Washington, DC, AACC, 1997, pp 223–250.
  24. Yoshida T, Sakane N, Umekawa T, Sakai M, Takahashi T, Kondo M: Mutation of  $\beta$ 3 adrenergic receptor gene and response to treatment of obesity. *Lancet* 1995; **346**: 1433–1434.
  25. Yuan X, Yamada K, Koyama K, *et al*: Beta 3-adrenergic receptor gene polymorphism is not a major genetic determinant of obesity and diabetes in Japanese general population. *Diabetes Res Clin Pract* 1997; **37**: 1–7.
  26. Masuo K, Katsuya T, Fu Y, Rakugi H, Ogihara T, Tuck ML: Beta2- and beta3-adrenergic receptor polymorphisms are related to the onset of weight gain and blood pressure elevation over 5 years. *Circulation* 2005; **28**: 3429–3434.
  27. Candelore MR, Deng L, Tota LM, Kelly LJ, Cascieri MA, Strader CD: Pharmacological characterization of a recently described human  $\beta$ 3-adrenergic receptor mutant. *Endocrinology* 1996; **6**: 2638–2641.
  28. Pietri-Rouxel F, Manning STJ, Gros J, Strosberg AD: The biochemical effect of the naturally occurring Trp64-Arg mutation on human  $\beta$ 3-adrenoceptor activity. *Eur J Biochem* 1997; **247**: 1174–1179.
  29. Kissebah AH, Sonnenberg GE, Myklebust J, *et al*: Quantitative trait loci on chromosomes 3 and 17 influence phenotypes of the metabolic syndrome. *Proc Natl Acad Sci U S A* 2000; **97**: 14478–14483.
  30. Castellano M, Muiesan ML, Cinelli A, *et al*: Lack of association of the Trp64Arg polymorphism of the  $\beta$ 3 adrenergic-receptor gene with blood pressure. *Circulation* 1997; **96** (Suppl 1): 226.
  31. Fujisawa T, Ikegami H, Yamato E, *et al*: Trp64Arg mutation of beta3-adrenergic receptor in essential hypertension: insulin resistance and the adrenergic system. *Am J Hypertens* 1997; **10**: 101–105.
  32. Takeuchi H, Saitoh S, Takagi S, *et al*: Metabolic syndrome and cardiac disease in Japanese men: applicability of the concept of metabolic syndrome defined by the National Cholesterol Education Program—Adult Treatment Panel III to Japanese men—the Tanno and Sobetsu Study. *Hypertens Res* 2005; **28**: 203–208.
  33. Tanaka H, Shimabukuro T, Shimabukuro M: High prevalence of metabolic syndrome among men in Okinawa. *J Atheroscler Thromb* 2005; **12**: 284–288.