

- 634. Blood supplements
- 635. Reagent for biological test
- 636. Mixed biological agent
- 639. Other biological agent

- 64. Drug against parasitic disease
 - 641. Protozoicide
 - 642. Parasiticide
 - 649. Other drug against parasitic disease

- 69. Other drugs for pathogenic organism

- 7. Drugs used not for treatment
 - 71. Dispensing drug
 - 711. Diluents
 - 712. Ointment base
 - 713. Solution
 - 714. Corrective, tinction
 - 715. Emulsifier
 - 719. Other dispensing drug

 - 72. Drugs used for diagnosis
 - 721. X-ray contrast medium
 - 722. Reagent for functional examination
 - 729. Other drugs for diagnosis

 - 73. Drugs used for public health
 - 731. Antiseptic
 - 732. Pesticide for prevention of epidemics
 - 733. Insect repellent
 - 734. Insecticide
 - 735. Rodenticide
 - 736. Other drug used for public health

 - 74. Reagent for extracorporeal diagnosis

- 741. Reagent for general examination
- 742. Reagent for blood analysis
- 743. Reagent for biochemical analysis
- 744. Reagent for immunologic or serologic test
- 745. Reagent for microbiological examination
- 746. Reagent for histopathological examination
- 747. Radiopharmaceutical for extracorporeal diagnosis
- 749. Other reagent for extracorporeal diagnosis

- 791. Adhesive bandage
- 799. Other drug not for treatment

8. Narcotics

- 81. Alkaloid agent
 - 811. Opiate alkaloid agent
 - 812. Coca alkaloid agent
 - 819. Other alkaloid agent

82. Non-alkaloid narcotics

- 821. Synthetic narcotic

9. Unclassifiable drug

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
General anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hypogenic sadative agent	Male	5	1.7	11	3.7	10	3.3	24	9.4	6	11.3	56	4.7
	Female	2	0.7	10	3.5	26	9.5	54	18.9	8	13.6	100	8.4
	Total	7	1.2	21	3.6	36	6.3	78	14.4	14	12.5	156	6.5
Antiepileptic agent	Male	1	0.3	0	0.0	5	1.7	4	1.6	0	0.0	10	0.8
	Female	0	0.0	1	0.4	2	0.7	3	1.1	2	3.4	8	0.7
	Total	1	0.2	1	0.2	7	1.2	7	1.3	2	1.8	18	0.8
Antipyretic analgesic agent	Male	24	8.4	23	7.8	30	10.0	62	24.3	19	35.8	158	13.3
	Female	60	20.4	42	14.8	39	14.3	66	23.2	22	37.3	229	19.2
	Total	84	14.5	65	11.2	69	12.0	128	23.7	41	36.6	387	16.2
Stimulant drug, antihypnotic drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antiparkinson drugs	Male	0	0.0	0	0.0	2	0.7	3	1.2	0	0.0	5	0.4
	Female	0	0.0	1	0.4	0	0.0	1	0.4	0	0.0	2	0.2
	Total	0	0.0	1	0.2	2	0.3	4	0.7	0	0.0	7	0.3
Antipsychotic drug	Male	6	2.1	7	2.4	12	4.0	8	3.1	5	9.4	38	3.2
	Female	5	1.7	6	2.1	11	4.0	40	14.0	6	10.2	68	5.7
	Total	11	1.9	13	2.2	23	4.0	48	8.9	11	9.8	106	4.4
Drug for common cold	Male	20	7.0	24	8.1	18	6.0	15	5.9	2	3.8	79	6.6
	Female	28	9.5	18	6.4	19	7.0	12	4.2	1	1.7	78	6.5
	Total	48	8.3	42	7.3	37	6.5	27	5.0	3	2.7	157	6.6
Other agent acting on CNS	Male	0	0.0	0	0.0	0	0.0	3	1.2	1	1.9	4	0.3

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other agent acting on CNS	Female	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	1	0.1
	Total	0	0.0	1	0.2	0	0.0	3	0.6	1	0.9	5	0.2
Local anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Muscle relaxant	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for autonomic nervous system	Male	1	0.3	2	0.7	0	0.0	2	0.8	0	0.0	5	0.4
	Female	1	0.3	0	0.0	0	0.0	3	1.1	1	1.7	5	0.4
	Total	2	0.3	2	0.3	0	0.0	5	0.9	1	0.9	10	0.4
Antispasmodic agent	Male	0	0.0	1	0.3	2	0.7	2	0.8	1	1.9	6	0.5
	Female	2	0.7	4	1.4	5	1.8	10	3.5	1	1.7	22	1.8
	Total	2	0.3	5	0.9	7	1.2	12	2.2	2	1.8	28	1.2
Diaphoretic, antiperspirants	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for PNS	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ophthalmic solution (others)	Male	52	18.2	58	19.7	54	18.0	53	20.8	18	34.0	235	19.8
	Female	48	16.3	70	24.7	75	27.5	89	31.2	25	42.4	307	25.7
	Total	100	17.2	128	22.1	129	22.5	142	26.3	43	38.4	542	22.7
Ophthalmic solution (for cataract)	Male	1	0.3	2	0.7	12	4.0	25	9.8	5	9.4	45	3.8
	Female	0	0.0	3	1.1	20	7.3	46	16.1	10	16.9	79	6.6

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Ophthalmic solution (for cataract)	Total	1	0.2	5	0.9	32	5.6	71	13.1	15	13.4	124	5.2
Ophthalmic solution (pilocarpine)	Male	0	0.0	0	0.0	3	1.0	0	0.0	0	0.0	3	0.3
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	3	0.5	0	0.0	0	0.0	3	0.1
Ophthalmic solution (β blocker)	Male	2	0.7	3	1.0	7	2.3	13	5.1	3	5.7	28	2.4
	Female	1	0.3	2	0.7	5	1.8	13	4.6	2	3.4	23	1.9
	Total	3	0.5	5	0.9	12	2.1	26	4.8	5	4.5	51	2.1
Ophthalmic solution (epinephrine)	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Ophthalmic solution (for glaucoma)	Male	2	0.7	0	0.0	9	3.0	6	2.4	3	5.7	20	1.7
	Female	1	0.3	1	0.4	4	1.5	8	2.8	1	1.7	15	1.3
	Total	3	0.5	1	0.2	13	2.3	14	2.6	4	3.6	35	1.5
Ophthalmic solution (steroid)	Male	0	0.0	3	1.0	6	2.0	6	2.4	5	9.4	20	1.7
	Female	6	2.0	6	2.1	9	3.3	16	5.6	6	10.2	43	3.6
	Total	6	1.0	9	1.6	15	2.6	22	4.1	11	9.8	63	2.6
Ophthalmic solution (unknown)	Male	0	0.0	1	0.3	0	0.0	3	1.2	0	0.0	4	0.3
	Female	2	0.7	0	0.0	1	0.4	10	3.5	0	0.0	13	1.1
	Total	2	0.3	1	0.2	1	0.2	13	2.4	0	0.0	17	0.7
Agent for ENT use	Male	4	1.4	1	0.3	2	0.7	3	1.2	0	0.0	10	0.8
	Female	1	0.3	5	1.8	3	1.1	1	0.4	0	0.0	10	0.8
	Total	5	0.9	6	1.0	5	0.9	4	0.7	0	0.0	20	0.8
Anti-vertigeneous drug	Male	2	0.7	0	0.0	0	0.0	2	0.8	2	3.8	6	0.5
	Female	3	1.0	2	0.7	1	0.4	7	2.5	3	5.1	16	1.3
	Total	5	0.9	2	0.3	1	0.2	9	1.7	5	4.5	22	0.9

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other agent acting on sensory organ	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Digitalis agent	Male	0	0.0	0	0.0	4	1.3	6	2.4	3	5.7	13	1.1
	Female	0	0.0	2	0.7	1	0.4	3	1.1	0	0.0	6	0.5
	Total	0	0.0	2	0.3	5	0.9	9	1.7	3	2.7	19	0.8
Xanthine agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other inotropic agent	Male	1	0.3	0	0.0	0	0.0	1	0.4	2	3.8	4	0.3
	Female	0	0.0	0	0.0	0	0.0	1	0.4	1	1.7	2	0.2
	Total	1	0.2	0	0.0	0	0.0	2	0.4	3	2.7	6	0.3
β blocker	Male	2	0.7	7	2.4	12	4.0	12	4.7	1	1.9	34	2.9
	Female	3	1.0	8	2.8	7	2.6	15	5.3	2	3.4	35	2.9
	Total	5	0.9	15	2.6	19	3.3	27	5.0	3	2.7	69	2.9
Agents for membrane-stabilizing activity	Male	0	0.0	2	0.7	4	1.3	4	1.6	1	1.9	11	0.9
	Female	1	0.3	0	0.0	3	1.1	6	2.1	0	0.0	10	0.8
	Total	1	0.2	2	0.3	7	1.2	10	1.9	1	0.9	21	0.9
Other antiarrhythmic agent	Male	0	0.0	0	0.0	1	0.3	1	0.4	0	0.0	2	0.2
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	2	0.4	0	0.0	3	0.1
Thyazide diuretic	Male	0	0.0	0	0.0	1	0.3	0	0.0	1	1.9	2	0.2
	Female	0	0.0	1	0.4	0	0.0	7	2.5	1	1.7	9	0.8
	Total	0	0.0	1	0.2	1	0.2	7	1.3	2	1.8	11	0.5
Loop diuretic	Male	0	0.0	1	0.3	2	0.7	6	2.4	2	3.8	11	0.9

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Loop diuretic	Female	0	0.0	3	1.1	2	0.7	6	2.1	4	6.8	15	1.3
	Total	0	0.0	4	0.7	4	0.7	12	2.2	6	5.4	26	1.1
Potassium sparing diuretic	Male	0	0.0	0	0.0	1	0.3	4	1.6	0	0.0	5	0.4
	Female	0	0.0	0	0.0	0	0.0	5	1.8	1	1.7	6	0.5
	Total	0	0.0	0	0.0	1	0.2	9	1.7	1	0.9	11	0.5
Carbonate dehydratase inhibitor	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Other diuretic	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	4	1.4	1	1.7	5	0.4
	Total	0	0.0	0	0.0	0	0.0	5	0.9	1	0.9	6	0.3
ACE inhibitor	Male	8	2.8	19	6.4	38	12.7	44	17.3	11	20.8	120	10.1
	Female	4	1.4	14	4.9	30	11.0	67	23.5	13	22.0	128	10.7
	Total	12	2.1	33	5.7	68	11.9	111	20.6	24	21.4	248	10.4
Ca antagonist	Male	7	2.4	33	11.2	69	23.0	89	34.9	13	24.5	211	17.7
	Female	8	2.7	19	6.7	40	14.7	101	35.4	24	40.7	192	16.1
	Total	15	2.6	52	9.0	109	19.0	190	35.2	37	33.0	403	16.9
Other antihypertensive agent	Male	2	0.7	6	2.0	10	3.3	8	3.1	0	0.0	26	2.2
	Female	0	0.0	0	0.0	1	0.4	10	3.5	1	1.7	12	1.0
	Total	2	0.3	6	1.0	11	1.9	18	3.3	1	0.9	38	1.6
Vascular reinforcing agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vasoconstrictor	Male	1	0.3	0	0.0	0	0.0	1	0.4	0	0.0	2	0.2
	Female	0	0.0	0	0.0	0	0.0	2	0.7	0	0.0	2	0.2

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Vasoconstrictor	Total	1	0.2	0	0.0	0	0.0	3	0.6	0	0.0	4	0.2
Vasodilator	Male	0	0.0	3	1.0	9	3.0	19	7.5	7	13.2	38	3.2
	Female	1	0.3	0	0.0	4	1.5	16	5.6	6	10.2	27	2.3
	Total	1	0.2	3	0.5	13	2.3	35	6.5	13	11.6	65	2.7
Antihyperlipidemic agent	Male	2	0.7	19	6.4	26	8.7	33	12.9	5	9.4	85	7.1
	Female	3	1.0	20	7.1	55	20.1	71	24.9	13	22.0	162	13.6
	Total	5	0.9	39	6.7	81	14.1	104	19.3	18	16.1	247	10.4
Other agent on cardiovascular system	Male	0	0.0	3	1.0	6	2.0	15	5.9	8	15.1	32	2.7
	Female	1	0.3	6	2.1	4	1.5	25	8.8	4	6.8	40	3.4
	Total	1	0.2	9	1.6	10	1.7	40	7.4	12	10.7	72	3.0
Respiratory stimulator	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antitussive agent	Male	2	0.7	1	0.3	2	0.7	5	2.0	0	0.0	10	0.8
	Female	2	0.7	7	2.5	7	2.6	7	2.5	1	1.7	24	2.0
	Total	4	0.7	8	1.4	9	1.6	12	2.2	1	0.9	34	1.4
Expectorant	Male	4	1.4	5	1.7	6	2.0	12	4.7	2	3.8	29	2.4
	Female	3	1.0	6	2.1	8	2.9	6	2.1	2	3.4	25	2.1
	Total	7	1.2	11	1.9	14	2.4	18	3.3	4	3.6	54	2.3
Antitussive agent and expectorant	Male	2	0.7	1	0.3	3	1.0	7	2.7	0	0.0	13	1.1
	Female	4	1.4	4	1.4	5	1.8	1	0.4	1	1.7	15	1.3
	Total	6	1.0	5	0.9	8	1.4	8	1.5	1	0.9	28	1.2
Bronchodilator	Male	3	1.0	5	1.7	5	1.7	9	3.5	3	5.7	25	2.1
	Female	3	1.0	3	1.1	4	1.5	4	1.4	1	1.7	15	1.3
	Total	6	1.0	8	1.4	9	1.6	13	2.4	4	3.6	40	1.7

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Gargle	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent acting on respiratory system	Male	1	0.3	3	1.0	1	0.3	1	0.4	0	0.0	6	0.5
	Female	2	0.7	2	0.7	1	0.4	2	0.7	1	1.7	8	0.7
	Total	3	0.5	5	0.9	2	0.3	3	0.6	1	0.9	14	0.6
Antidiarrheal drug	Male	6	2.1	8	2.7	7	2.3	7	2.7	2	3.8	30	2.5
	Female	5	1.7	8	2.8	7	2.6	11	3.9	3	5.1	34	2.8
	Total	11	1.9	16	2.8	14	2.4	18	3.3	5	4.5	64	2.7
Antiulcerative agent	Male	11	3.8	23	7.8	34	11.3	64	25.1	14	26.4	146	12.3
	Female	10	3.4	23	8.1	31	11.4	71	24.9	18	30.5	153	12.8
	Total	21	3.6	46	8.0	65	11.3	135	25.0	32	28.6	299	12.5
Stomachic	Male	3	1.0	1	0.3	4	1.3	9	3.5	6	11.3	23	1.9
	Female	0	0.0	4	1.4	2	0.7	16	5.6	4	6.8	26	2.2
	Total	3	0.5	5	0.9	6	1.0	25	4.6	10	8.9	49	2.1
Antacid	Male	1	0.3	1	0.3	2	0.7	2	0.8	1	1.9	7	0.6
	Female	1	0.3	1	0.4	0	0.0	1	0.4	1	1.7	4	0.3
	Total	2	0.3	2	0.3	2	0.3	3	0.6	2	1.8	11	0.5
Laxative	Male	3	1.0	4	1.4	4	1.3	24	9.4	11	20.8	46	3.9
	Female	9	3.1	13	4.6	7	2.6	27	9.5	5	8.5	61	5.1
	Total	12	2.1	17	2.9	11	1.9	51	9.4	16	14.3	107	4.5
Cholagogue	Male	1	0.3	2	0.7	5	1.7	5	2.0	1	1.9	14	1.2
	Female	1	0.3	2	0.7	1	0.4	2	0.7	0	0.0	6	0.5
	Total	2	0.3	4	0.7	6	1.0	7	1.3	1	0.9	20	0.8
Combined digestive drug	Male	15	5.2	8	2.7	16	5.3	7	2.7	0	0.0	46	3.9

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Combined digestive drug	Female	12	4.1	13	4.6	5	1.8	6	2.1	2	3.4	38	3.2
	Total	27	4.7	21	3.6	21	3.7	13	2.4	2	1.8	84	3.5
Other agent acting on alimentary system	Male	3	1.0	3	1.0	6	2.0	9	3.5	2	3.8	23	1.9
	Female	2	0.7	5	1.8	3	1.1	15	5.3	5	8.5	30	2.5
	Total	5	0.9	8	1.4	9	1.6	24	4.4	7	6.3	53	2.2
Pituitary hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Hormone drug of salivary gland	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Thyroid parathyroid hormone drug	Male	2	0.7	1	0.3	1	0.3	2	0.8	2	3.8	8	0.7
	Female	4	1.4	5	1.8	3	1.1	9	3.2	1	1.7	22	1.8
	Total	6	1.0	6	1.0	4	0.7	11	2.0	3	2.7	30	1.3
Anabolic steroid drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Adrenocortical hormone drug	Male	5	1.7	2	0.7	7	2.3	7	2.7	0	0.0	21	1.8
	Female	3	1.0	7	2.5	9	3.3	2	0.7	2	3.4	23	1.9
	Total	8	1.4	9	1.6	16	2.8	9	1.7	2	1.8	44	1.8
Androgen drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Estrogen or progesterone drug	Male	0	0.0	0	0.0	0	0.0	3	1.2	0	0.0	3	0.3
	Female	2	0.7	6	2.1	0	0.0	0	0.0	0	0.0	8	0.7

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Estrogen or progesterone drug	Total	2	0.3	6	1.0	0	0.0	3	0.6	0	0.0	11	0.5
Mixed hormone drug	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other hormone drug	Male	0	0.0	0	0.0	2	0.7	5	2.0	3	5.7	10	0.8
	Female	0	0.0	0	0.0	2	0.7	7	2.5	2	3.4	11	0.9
	Total	0	0.0	0	0.0	4	0.7	12	2.2	5	4.5	21	0.9
Agent acting on urinary system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent acting on genital system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Uterine constrictor	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Contraceptive	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	2	0.7	0	0.0	0	0.0	0	0.0	0	0.0	2	0.2
	Total	2	0.3	0	0.0	0	0.0	0	0.0	0	0.0	2	0.1
Drugs for hemorrhoids	Male	0	0.0	1	0.3	2	0.7	1	0.4	0	0.0	4	0.3
	Female	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
	Total	0	0.0	1	0.2	3	0.5	1	0.2	0	0.0	5	0.2
Other agent acting on urogenital or anus	Male	0	0.0	1	0.3	6	2.0	20	7.8	4	7.5	31	2.6
	Female	1	0.3	0	0.0	1	0.4	7	2.5	4	6.8	13	1.1
	Total	1	0.2	1	0.2	7	1.2	27	5.0	8	7.1	44	1.8

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Integumentary microbiosite	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Protective agent for wound	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pyostatic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Analgenic (external application)	Male	1	0.3	1	0.3	0	0.0	1	0.4	0	0.0	3	0.3
	Female	1	0.3	0	0.0	1	0.4	2	0.7	0	0.0	4	0.3
	Total	2	0.3	1	0.2	1	0.2	3	0.6	0	0.0	7	0.3
Integumentary Parasiticide	Male	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Emollient	Male	1	0.3	0	0.0	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	1	0.2	0	0.0	0	0.0	0	0.0	0	0.0	1	0.0
Hair tonic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Bath liquid	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other surface-acting agent	Male	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.1

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other surface-acting agent	Female	2	0.7	0	0.0	0	0.0	1	0.4	0	0.0	3	0.3
	Total	2	0.3	1	0.2	0	0.0	1	0.2	0	0.0	4	0.2
Dental local anesthetic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for devitalization of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Analgesic or sedative for dental use	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0
Agent for mummification of the pulp	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for pulp capping	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics for dental use	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for dental use	Male	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Female	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	1	0.2	0	0.0	2	0.1
Vitamin A or D	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Vitamin A or D	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Vitamin B1	Male	0	0.0	1	0.3	2	0.7	2	0.8	2	3.8	7	0.6
	Female	0	0.0	0	0.0	1	0.4	3	1.1	1	1.7	5	0.4
	Total	0	0.0	1	0.2	3	0.5	5	0.9	3	2.7	12	0.5
Vitamin B complex	Male	4	1.4	2	0.7	12	4.0	15	5.9	9	17.0	42	3.5
	Female	4	1.4	5	1.8	1	0.4	30	10.5	5	8.5	45	3.8
	Total	8	1.4	7	1.2	13	2.3	45	8.3	14	12.5	87	3.7
Vitamin C	Male	0	0.0	1	0.3	0	0.0	1	0.4	0	0.0	2	0.2
	Female	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	1	0.1
	Total	0	0.0	2	0.3	0	0.0	1	0.2	0	0.0	3	0.1
Vitamin E	Male	0	0.0	0	0.0	1	0.3	2	0.8	1	1.9	4	0.3
	Female	0	0.0	1	0.4	0	0.0	0	0.0	0	0.0	1	0.1
	Total	0	0.0	1	0.2	1	0.2	2	0.4	1	0.9	5	0.2
Vitamin K	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.4	4	1.4	1	1.7	6	0.5
	Total	0	0.0	0	0.0	1	0.2	4	0.7	1	0.9	6	0.3
Mixed vitamin	Male	1	0.3	1	0.3	1	0.3	7	2.7	2	3.8	12	1.0
	Female	2	0.7	1	0.4	2	0.7	12	4.2	3	5.1	20	1.7
	Total	3	0.5	2	0.3	3	0.5	19	3.5	5	4.5	32	1.3
Other vitamin	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Calcium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	1	0.4	7	2.6	4	1.4	1	1.7	13	1.1
	Total	0	0.0	1	0.2	7	1.2	4	0.7	1	0.9	13	0.5

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Mineral preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	1.9	1	0.1
	Female	2	0.7	0	0.0	0	0.0	3	1.1	2	3.4	7	0.6
	Total	2	0.3	0	0.0	0	0.0	3	0.6	3	2.7	8	0.3
Saccaride preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Organic acid preparation	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
protein or amino acid preparation	Male	0	0.0	2	0.7	0	0.0	0	0.0	0	0.0	2	0.2
	Female	0	0.0	0	0.0	1	0.4	0	0.0	1	1.7	2	0.2
	Total	0	0.0	2	0.3	1	0.2	0	0.0	1	0.9	4	0.2
Hormone drugs made from internal organ	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for infant	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other tonic	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Blood substitute	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Styptic	Male	3	1.0	2	0.7	3	1.0	2	0.8	0	0.0	10	0.8

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Styptic	Female	3	1.0	2	0.7	4	1.5	2	0.7	1	1.7	12	1.0
	Total	6	1.0	4	0.7	7	1.2	4	0.7	1	0.9	22	0.9
Anticoagulant	Male	0	0.0	2	0.7	4	1.3	2	0.8	1	1.9	9	0.8
	Female	0	0.0	1	0.4	2	0.7	4	1.4	0	0.0	7	0.6
	Total	0	0.0	3	0.5	6	1.0	6	1.1	1	0.9	16	0.7
Other blood substitute	Male	0	0.0	2	0.7	9	3.0	23	9.0	7	13.2	41	3.4
	Female	0	0.0	0	0.0	6	2.2	20	7.0	3	5.1	29	2.4
	Total	0	0.0	2	0.3	15	2.6	43	8.0	10	8.9	70	2.9
Agent for hemodialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for peritoneal dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other agent for dialysis	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Agent for liver disease	Male	1	0.3	2	0.7	3	1.0	5	2.0	0	0.0	11	0.9
	Female	0	0.0	0	0.0	2	0.7	2	0.7	0	0.0	4	0.3
	Total	1	0.2	2	0.3	5	0.9	7	1.3	0	0.0	15	0.6
Antidote	Male	0	0.0	0	0.0	0	0.0	2	0.8	0	0.0	2	0.2
	Female	0	0.0	0	0.0	2	0.7	1	0.4	0	0.0	3	0.3
	Total	0	0.0	0	0.0	2	0.3	3	0.6	0	0.0	5	0.2
Agent for habitual intoxication	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Agent for habitual intoxication	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for gout	Male	8	2.8	13	4.4	14	4.7	21	8.2	3	5.7	59	5.0
	Female	0	0.0	0	0.0	2	0.7	6	2.1	1	1.7	9	0.8
	Total	8	1.4	13	2.2	16	2.8	27	5.0	4	3.6	68	2.9
Enzyme preparation	Male	2	0.7	1	0.3	3	1.0	4	1.6	1	1.9	11	0.9
	Female	2	0.7	4	1.4	6	2.2	6	2.1	0	0.0	18	1.5
	Total	4	0.7	5	0.9	9	1.6	10	1.9	1	0.9	29	1.2
Antidiabetics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Other agent on metabolic system	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Unclassified agent on metabolic system	Male	2	0.7	2	0.7	3	1.0	6	2.4	2	3.8	15	1.3
	Female	3	1.0	3	1.1	3	1.1	5	1.8	2	3.4	16	1.3
	Total	5	0.9	5	0.9	6	1.0	11	2.0	4	3.6	31	1.3
Chlorophyll agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Pigment agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other cell activation agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	1	0.4	0	0.0	1	0.1
	Total	0	0.0	0	0.0	0	0.0	1	0.2	0	0.0	1	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Alkylating agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antimetabolite	Male	0	0.0	0	0.0	0	0.0	1	0.4	1	1.9	2	0.2
	Female	0	0.0	0	0.0	1	0.4	1	0.4	0	0.0	2	0.2
	Total	0	0.0	0	0.0	1	0.2	2	0.4	1	0.9	4	0.2
Antibiotics for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Plant component for antitumor agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Other antitumor agent	Male	1	0.3	0	0.0	1	0.3	2	0.8	2	3.8	6	0.5
	Female	1	0.3	0	0.0	1	0.4	6	2.1	0	0.0	8	0.7
	Total	2	0.3	0	0.0	2	0.3	8	1.5	2	1.8	14	0.6
Antihistaminic drug	Male	9	3.1	7	2.4	5	1.7	5	2.0	1	1.9	27	2.3
	Female	15	5.1	8	2.8	6	2.2	8	2.8	2	3.4	39	3.3
	Total	24	4.1	15	2.6	11	1.9	13	2.4	3	2.7	66	2.8
Agent for stimulus therapy	Male	0	0.0	0	0.0	0	0.0	2	0.8	0	0.0	2	0.2
	Female	0	0.0	2	0.7	1	0.4	0	0.0	0	0.0	3	0.3
	Total	0	0.0	2	0.3	1	0.2	2	0.4	0	0.0	5	0.2
Agent for nonspecific immunogenicity	Male	0	0.0	0	0.0	0	0.0	0	0.0	1	1.9	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	1	0.9	1	0.0
Other antiallergic drug	Male	14	4.9	14	4.7	14	4.7	11	4.3	1	1.9	54	4.5

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Other antiallergic drug	Female	17	5.8	15	5.3	15	5.5	14	4.9	1	1.7	62	5.2
	Total	31	5.3	29	5.0	29	5.1	25	4.6	2	1.8	116	4.9
Crude drug	Male	0	0.0	1	0.3	0	0.0	0	0.0	0	0.0	1	0.1
	Female	0	0.0	1	0.4	1	0.4	1	0.4	1	1.7	4	0.3
	Total	0	0.0	2	0.3	1	0.2	1	0.2	1	0.9	5	0.2
Herb medicine	Male	31	10.8	23	7.8	37	12.3	36	14.1	7	13.2	134	11.3
	Female	37	12.6	38	13.4	34	12.5	40	14.0	9	15.3	158	13.2
	Total	68	11.7	61	10.6	71	12.4	76	14.1	16	14.3	292	12.3
Other drugs by a herb medicine	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(+))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(-))	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics (G(+), G(-))	Male	2	0.7	2	0.7	6	2.0	6	2.4	0	0.0	16	1.3
	Female	8	2.7	6	2.1	10	3.7	5	1.8	0	0.0	29	2.4
	Total	10	1.7	8	1.4	16	2.8	11	2.0	0	0.0	45	1.9
Antibiotics (G(+), G(-), Mycoplasma)	Male	2	0.7	3	1.0	3	1.0	4	1.6	1	1.9	13	1.1
	Female	3	1.0	2	0.7	0	0.0	2	0.7	0	0.0	7	0.6
	Total	5	0.9	5	0.9	3	0.5	6	1.1	1	0.9	20	0.8
Antibiotics (Rickettsia, Chlamidia)	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0

(Continued)

Prevalence of taking medicine in previous 2 weeks

		40-49yr		50-59yr		60-69yr		70-79yr		80yr-		Total	
		N	%	N	%	N	%	N	%	N	%	N	%
Antibiotics (Rickettsia, Chlamidia)	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics against acid- fastbacterium	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Antibiotics against fungus	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Other antibiotics	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	1	0.4	0	0.0	0	0.0	1	0.1
	Total	0	0.0	0	0.0	1	0.2	0	0.0	0	0.0	1	0.0
Sulfa drug	Male	0	0.0	0	0.0	1	0.3	0	0.0	0	0.0	1	0.1
	Female	2	0.7	0	0.0	0	0.0	2	0.7	0	0.0	4	0.3
	Total	2	0.3	0	0.0	1	0.2	2	0.4	0	0.0	5	0.2
Antituberculous agent	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Drugs for leprosy	Male	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Female	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	Total	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
Synthetic antibiotics	Male	2	0.7	0	0.0	2	0.7	1	0.4	0	0.0	5	0.4
	Female	3	1.0	2	0.7	3	1.1	8	2.8	0	0.0	16	1.3
	Total	5	0.9	2	0.3	5	0.9	9	1.7	0	0.0	21	0.9
Antiviral agent	Male	1	0.3	1	0.3	0	0.0	1	0.4	0	0.0	3	0.3
	Female	0	0.0	1	0.4	1	0.4	0	0.0	0	0.0	2	0.2
	Total	1	0.2	2	0.3	1	0.2	1	0.2	0	0.0	5	0.2

(Continued)