



Fig. 1 - Square stepping exercise.

### Exercise intervention

Based on the mechanism and direction of falls, and on ladder training drills designed for athletes, we developed our SSE program and used it for the intervention described here. SSE was performed on a thin mat of 250 cm by 100 cm, partitioned into 40 small squares (25 cm each side). A smaller mat, 200 x 80 cm, also partitioned with smaller squares (20 cm each side), was prepared for shorter participants (i.e., those less than 150 cm in height). Participants were required to step in the length direction (250 or 200 cm), basically without treading on the lines making up the squares. The SSE program included forward, backward, lateral and oblique steps, and step patterns were progressively made more complicated (Fig. 1).

One hundred and nineteen step patterns were developed and categorized into Junior (8 patterns), Basic (19), Semi-Regular (17), Regular (26), Senior (35) and Master (14) patterns, according to difficulty. Examples of Junior, Basic, Regular and Master patterns are shown in

Fig. 2. Junior patterns consisted of easy steps, i.e., forward steps only. For Basic patterns, lateral steps were added to the Junior patterns. Further, oblique and backward steps were added in the Semi-Regular patterns; somewhat complicated oblique and backward steps were added in the Regular patterns; complicated lateral, oblique and backward steps were added in the Senior patterns; and much more complicated steps were added in the Master patterns.

There were some options, so that step patterns in each category were more complicated. Many patterns required a wide stance over one or two squares. Some patterns had bilateral symmetry, which led participants to grasp the pattern image easily. Each pattern consisted of two to 16 steps; participants were asked to repeat the step pattern until they got to the end of the mat. Once a pattern was performed, a mirror-imaged pattern was then also performed.

The exercise group participated in a six-month regimen of SSE once a week. Each SSE class consisted of 10-min warm-up activities, such as walking and mild stretching, 30-min SSE, 10-min whole-body light resistance training, and 10-min cool-down. To familiarize participants with SSE, only Junior patterns were used for the first month. From the second month, Junior patterns were performed as the warm-up. Participants were encouraged to try some new patterns in the Basic or more complicated categories. Essentially, a one-step pattern was repeated 3-5 times and then the mirror-image pattern was repeated the same number of times. If participants could not step a pattern smoothly, they repeated it until it was accomplished, and then they progressed to a new step pattern. The Basic and more complicated category patterns which included lateral, oblique and backward steps were carefully introduced, because many participants needed a long time to learn each new pattern. During the SSE program, step patterns drawn on leaflets or boards were not provided. Participants were asked to learn the pattern by watching an instructor's movements.

Participants were also asked to go forward on the mat one-way, to pass back by the mat walking normally, and then to stand in a line for the next stepping exercise: this meant that the number of steps did not differ among participants. Raising the heels was a recommended option: this made it more difficult for participants to step the patterns and ensured improvements in agility, locomotion speed and balance. Although step cadence was not determined, participants stepped slowly at first and then were able to step faster after some practice. For example, in the Basic pattern shown in Fig. 2, participants stepped at approximately 1 step/sec. They needed some time to remember where they should place their feet during the steps, and so took 20 to 30 seconds in one direction. After some practice, they could step out the patterns at approximately 2 steps/s: and took 20 or fewer seconds to complete.

Junior				Basic				Regular				Master			

Fig. 2 - Examples of Junior, Basic, Regular and Master patterns of square stepping exercise. Steps were performed according to numbers, from lower end of each figure to upper end. Odd and even numbers indicate location of right and left feet, respectively.

In each SSE class, participants gathered in a room in a Juroku Bank branch building; no transportation was provided. The floor of the venue in which classes was held was fitted with a thin carpet. Two instructors concurrently taught all SSE participants.

*Statistical analysis*

Data are presented as means  $\pm$ SD. The overall design employed a 2 (group) by 2 (time) repeated-measures format. Repeated-measures techniques were conducted to determine time-by-group interactions. Before this, the Kolmogorov-Smirnov test was applied to determine if data on each test significantly differed from normal distribution. The Levene test was then used to verify the assumption that variances were equal across groups. When significances in these tests were found, the Wilcoxon signed-rank test for paired data was used. Two kinds of correlation coefficients in the SSE group were also calculated: the first between baseline data and the pre-post change for

each test; the second between the change in one test and the change in another test. These coefficients were calculated by: (i) Pearson's correlation analysis, when data were similar to normal distribution, and (ii) Spearman's correlation analysis when data were significantly different from normal distribution. All statistics were computed using SPSS version 11.5.1 J. The level of significance was set at  $\alpha = 0.05$ .

**RESULTS**

The baseline characteristics of the two groups were very similar (Table 1). Data of the SSE group in agility, leg power, locomotion speed, flexibility and balance measures were comparable with those of the control group at the beginning of the study (Table 2). Average attendance in the SSE program was 92% ( $\pm 7.9$ , range= 75-100%). In the exercise group, no participants discontinued involvement or claimed knee or back pains that had not already existed at the start of the study. No falls or injuries

Table 1 - Baseline characteristics of participants.

Measure	Control (n=26)	Exercise (n=26)
Age (yrs)	68.3±6.3	67.5±4.9
Height (cm)	151.4±6.3	148.5±4.3
Weight (kg)	52.5±9.3	53.2±7.0

resulted directly from the exercises. Of all the participants in the exercise group (n=26), one achieved only the Basic pattern, ranked as the second easiest category, by the time the six-month intervention was over. He could step Basic patterns easily and safely, but had difficulty in stepping the more complicated ones. Two other participants achieved Semi-regular patterns. Of the remaining participants, 14, 4 and 5 could step Regular, Senior and Master patterns, respectively, at the end of the six-month regimen. Compliance among controls was 100%, i.e., no controls engaged in any new form of exercise, they all maintained their lifestyles, and attended pre- and post-measurement sessions.

A repeated-measure ANOVA revealed a significant group-by-time interaction for standing up from a lying position ( $p<0.01$ ). Univariate follow-up analysis revealed that the significant agility increases were attributable to the SSE program (Table 2). The exercise group experienced

a decrease of 0.39 seconds in this measurement. Walking round two cones and sit-and-reach had significant group-by-time interactions ( $p<0.05$ ). Participants in the exercise group had a significant decrease in walking time (-1.4 s) and a significant increase in sit-and-reach (+5.8 cm). Kolmogorov-Smirnov tests revealed that data for chair-stand in 10 seconds and single-leg balance with eyes closed could not be assumed to be normally distributed. Therefore, Wilcoxon signed-rank tests were applied to these data, in which scores for chair stand (+1.8 n/10 s) and single-leg balance (+4.6 s) increased significantly in the exercise group. The control group exhibited no significant improvement in any of the five measures.

Correlation coefficients between baseline data and changes in each test are shown in Table 3a. They range from -0.45 to 0.27, accounting for 1.4-20.3% of variance; none achieved significance level ( $p<0.05$ ) except for sit-and-reach. Correlation coefficients between the change in one test and the change in another test were not statistically significant. All coefficients were 0.37 or less in absolute value (Table 3b), which accounted for approximately 13.7% or less of each variance.

## DISCUSSION

These results demonstrate improvements in agility, leg power, locomotion speed, flexibility and balance resulting from the six-month regimen of our square-stepping

Table 2 - Functional fitness at pre- and post-treatment and comparison of changes in variables.

Functional fitness related to risk of falls	Measurement item	Control group		Exercise group	
		Pre	Post	Pre	Post
Agility	Standing up from a lying position (s)	3.26±0.93	3.70±1.42	3.39±1.20	3.00±0.90 <sup>a</sup>
Leg power	Chair-stand in 10 seconds (n)	8.8±2.6	7.6±4.3	7.5±2.2	9.3±2.6 <sup>b</sup>
Locomotion speed	Walking round two cones (s)	22.6±3.4	22.9±4.8	21.2±3.5	19.8±3.6 <sup>a</sup>
Flexibility	Sit-and-reach (cm)	3.2±8.8	4.2±11.0	3.7±13.9	9.5±11.0 <sup>a</sup>
Balance	Single-leg balance with eyes closed (s)	11.0±15.5	6.7±9.1	9.3±11.7	13.9±15.6 <sup>b</sup>

Significances by Kolmogorov-Smirnov test were found in leg power and balance; Group-by-time interactions were found in agility, locomotion speed and flexibility by ANOVA; <sup>a</sup>Significant main effect of time; <sup>b</sup>Significant difference by Wilcoxon signed-rank test.

Table 3 - Correlation coefficients: (a) between baseline data and change in each test and (b) between change in one test and change in other test.

	(a)	(b)				
		1	2	3	4	5
1. Standing up from a lying position	0.27	1	0.27	0.21	0.37	0.16
2. Chair-stand in 10 seconds	-0.12		1	0.17	0.30	-0.14
3. Walking round two cones	0.24			1	0.17	0.15
4. Sit-and-reach	-0.45 <sup>*</sup>				1	0.06
5. Single-leg balance with eyes closed	0.12					1

Spearman's correlation coefficients were calculated when sit-and-reach or single-leg balance with eyes closed were included as independent variables; Other coefficients calculated by Pearson's analysis; <sup>\*</sup> $p<0.05$ .

exercise (SSE) program. In addition, the attendance rate was relatively high and there were no accidents during the SSE program. Therefore, we believe that it is an effective, practical means of decreasing the risk factors for falling in apparently healthy elderly people.

Standing up from a lying position, which we defined as agility, was significantly improved. This task requires a combination of quick response and whole-body movement. Whole-body movement during the test needs control over the center of gravity of the body, or dynamic balance. There are no published reports of changes in time or dynamic balance in standing up from a lying position as a result of an exercise intervention in the elderly, but it has been hypothesized that reaction time (2, 7) and dynamic balance (6, 16, 17) are major determinants of this test item. In previous studies, increases in reaction time are associated with high risk factors for falls (2, 7). The twelve-month prospective study of Nevitt et al. (2) found that those who had an increase in 0.75 or more seconds in hand-reaction time showed 1.8 times the relative risk of falls than those who had an increase of 0.74 seconds or less. In addition, anterior body mass carriage following a trip needs dynamic balance to control the whole body to avoid falling (6, 16, 17). Therefore, improving agility (standing up from a lying position) probably provided some reduction in risk factors for falls.

The increase in the test of chair-stand in 10 seconds probably reflected the improvement in participants' leg power (14) even though it was not assessed directly. A number of studies have identified a lack of muscular strength or power as risk factors for falling (18, 19), especially trip-induced falls (20). For example, being unable, or taking more than 2 seconds, to rise from a chair without using one's arms was associated with an increased risk of injury (2); fallers show a significant decrease in the strength of knees and ankles compared with non-fallers (21). Hip flexion joint torque from a sitting to standing position, which is related with leg power, may also be an important factor for quick chair-stand movement (22).

A low walking speed is also considered as a risk factor for falling (23-25). In the prospective study of Taylor et al. (25), a decrease of 0.22 m/sec in maximal walking speed over ten years was associated with an increase in hip fracture risk (RR 1.25, 95% CI 1.17-1.33). In the present study, the speed of walking round two cones changed from 0.79 m/sec before the program to 0.85 m/sec at the end of the program. Although a decrease in falls in itself cannot be attributed to a change of only 0.06 m/sec, the improvement in this test probably provided some improvement in risk factors for falls. Also, this improvement may play a distinct role in falls, by improving neuromuscular function, so that the SSE program influences an individual's speed, coordination and protective responses during a fall (23).

Increases in hip joint flexibility, which was measured by the sit-and-reach test in this study, should result in decreasing the risk factors for falls (13). The increase in the present study (about 6 cm) was slightly higher than our expectations, because some recent reports have found that exercise intervention that did not focus on flexibility showed an increase of about 3 cm; in these, exercise frequency or duration were not very different from those of our study (26, 27). This improvement may be due to the leg and hip stretching exercise that was provided in every warm-up period. We also told participants that doing sufficient stretching could prevent falls and injuries from occurring in the SSE program, which made them concentrate on stretching and thus improving sit-and-reach scores. A previous study (13) showed that those who had fallen three or more times in the previous year had significantly lower hip-joint flexibility than those who had had no falls. It is probable that hip-joint flexibility is related to stability and mobility in the elderly. Those with lower hip joint flexibility could not walk smoothly with a certain stance, and had great difficulty in maintaining their stability and placing the foot as a support for their bodies when a trip occurred.

Postural instability has also been cited as a risk factor for falls in the elderly (2, 12, 13, 16, 25, 28). It was hypothesized that single-leg balance with eyes closed, defined as balance, would be improved due to the stepping imposed by the SSE program with raised heels. Raised heels, which for safety reasons was not added until participants were familiar with a step pattern, probably afforded neuromuscular stimulation to the exercisers. Shigematsu et al. (29) demonstrated increases in single-leg balance with eyes closed in women aged 72-87 years who participated in a three-month dance-based aerobic exercise. They performed movements that extended, flexed, abducted, adducted and rotated the leg and foot, such as side-stepping, fast walking, forward-backward stepping and heel rises. Participants in the present study performed steps similar to those of the dance movements of Shigematsu et al.

These findings indicate that individuals in the exercise group were able to move their whole body more quickly and accurately after the exercise program. This may have clinical meaning, given that the leg movement in the front, side or oblique directions after tripping or slipping is appropriately carried out, and the center of gravity of the body is controlled to avoid falls. Since only 1.4-20.3% of the variance of the changes in each test could be explained by the initial levels, it is suggested that all persons, independently of their initial performance on any test, could benefit from our SSE program in at least one fitness element.

The present study was conducted with non-randomized allocation of participants. However, it may be difficult for persons who are transitioning to frailty or who are already frail to attend an SSE program. Therefore, our results may

not be generalizable to a wider population, although they do appear to demonstrate that the SSE program is applicable as a primary prevention strategy. Only data concerning some functional fitness elements according to risk factors were used, so that other functional fitness elements, such as maximum step length or rapid stepping ability (30) and the effect of the SSE program on fall occurrence were not determined. However, these results clearly indicate that apparently healthy individuals aged 60-80 with no regular exercise habits can reduce their risk factors for falling by undertaking a square stepping exercise program once a week for six months.

### ACKNOWLEDGEMENT

This study was supported by the Mitsui Sumitomo Insurance Welfare Foundation (principal investigator R. Shigematsu).

### REFERENCES

1. Orimo H, Hashimoto T, Sakata K, Yoshimura N, Suzuki T, Hosoi T. Trends in the incidence of hip fracture in Japan, 1987-1997: The third nationwide survey. *J Bone Miner Metab* 2000; 18: 126-31.
2. Nevitt MC, Cummings SR, Hudes ES. Risk factors for injurious falls: A prospective study. *J Gerontol* 1991; 46: M164-70.
3. Blake AJ, Morgan K, Bendall MJ, et al. Falls by elderly people at home: Prevalence and associated factors. *Age Ageing* 1998; 17: 365-72.
4. Smeesters C, Hayes WC, McMahon TA. Disturbance type and gait speed affect fall direction and impact location. *J Biomech* 2001; 34: 309-17.
5. Greenspan SL, Myers ER, Kiel DP, Parker RA, Hayes WC, Resnick NM. Fall direction, bone mineral density, and function: Risk factors for hip fracture in frail nursing home elderly. *Am J Med* 1998; 104: 539-45.
6. Pavol MJ, Owings TM, Foley KT, Grabiner MD. Mechanisms leading to a fall from an induced trip in healthy older adults. *J Gerontol* 2001; 56A: M428-37.
7. Bogert AJ, Pavol MJ, Grabiner MD. Response time is more important than walking speed for the ability of older adults to avoid a fall after a trip. *J Biomech* 2002; 35: 199-205.
8. Herala M, Kivela SL, Honkanen R, Koski K, Laippala P, Luukinen H. Recent decline in heavy outdoor work activity predicts occurrence of fractures among the home-dwelling elderly. *Osteoporos Int* 2002; 13: 42-4.
9. Luukinen H, Koski K, Laippala P, Kivela SL. Factors predicting fractures during falling impacts among home-dwelling older adults. *J Am Geriatr Soc* 1997; 45: 1302-9.
10. Sherrington C, Lord SR. Increased prevalence of fall risk factors in older people following hip fracture. *Gerontology* 1998; 44: 340-4.
11. Koepell TD, Wolf ME, Buchner DM, et al. Footwear style and risk of falls in older adults. *J Am Geriatr Soc* 2004; 52: 1495-501.
12. Chu LW, Pei CK, Chiu A, et al. Risk factors for falls in hospitalized older medical patients. *J Gerontol* 1999; 54: M38-43.
13. Gehlsen GM, Whale MH. Falls in the elderly: Part II: Balance, strength, and flexibility. *Arch Phys Med Rehabil* 1990; 71: 739-41.
14. Rikli RE, Jones CJ. Development and validation of a functional fitness test for community-residing older adults. *J Aging Phys Act* 1999; 7: 129-61.
15. Shigematsu R, Tanaka K. Age scale for assessing functional fitness in older Japanese ambulatory women. *Aging Clin Exp Res* 2000; 12: 256-63.
16. Rogers ME, Rogers NL, Takeshima N, Islam MM. Methods to assess and improve the physical parameters associated with fall risk in older adults. *Prev Med* 2003; 36: 255-64.
17. Wolfson LI, Whipple R, Amerman P, Kleinberg A. Stressing the postural response. A quantitative method for testing balance. *J Am Geriatr Soc* 1986; 34: 845-50.
18. American Geriatrics Society, British Geriatric Society, and American Academy of Orthopaedic Surgeons Panel on Falls Prevention. Guideline for the prevention of falls in older persons. *J Am Geriatr Soc* 2001; 49: 664-72.
19. Moreland JD, Richardson JA, Goldsmith CH, Clase CM. Muscle weakness and falls in older adults: A systematic review and meta-analysis. *J Am Geriatr Soc* 2004; 52: 1121-9.
20. Pavol MJ, Owings TM, Foley KT, Grabiner MD. Influence of lower extremity strength of healthy older adults on the outcome of an induced trip. *J Am Geriatr Soc* 2002; 50: 256-62.
21. Whipple RH, Wolfson LI, Amerman PM. The relationship of knee and ankle weakness to falls in nursing home residents: An isokinetic study. *J Am Geriatr Soc* 1987; 35: 13-20.
22. Mak MK, Levin O, Mizrahi J, Hui-Chan CW. Joint torques during sit-to-stand in healthy subjects and people with Parkinson's disease. *Clin Biomech* 2003; 18: 197-206.
23. Dargent-Molina P, Favier F, Grandjean H, et al. Fall-related factors and risk of hip fracture: the EPIDOS prospective study. *Lancet* 1996; 348: 145-9.
24. Rekeneire N, Visser M, Peila R, et al. Is a fall just a fall: correlates of falling in healthy older persons. The Health, Aging and Body Composition Study. *J Am Geriatr Soc* 2003; 51: 841-6.
25. Taylor BC, Schreiner PJ, Stone KL, et al. Long-term prediction of incident hip fracture risk in elderly white women: Study of Osteoporotic Fractures. *J Am Geriatr Soc* 2004; 52: 1479-86.
26. Adams KJ, Swank AM, Berning JM, Sevens-Adams PG, Barnard KL, Shimp-Bowerman J. Progressive strength training in sedentary, older African American women. *Med Sci Sports Exerc* 2001; 33: 1567-76.
27. Ransdell LB, Taylor A, Oakland D, Schmidt J, Moyer-Mileur L, Shultz B. Daughters and mothers exercising together: effects of home- and community-based programs. *Med Sci Sports Exerc* 2003; 35: 286-96.
28. Lord SR, Ward JA, Williams P, Anstey KJ. Physiological factors associated with falls in older community-dwelling women. *J Am Geriatr Soc* 1994; 42: 1110-7.
29. Shigematsu R, Chang M, Yabushita N, et al. Dance-based aerobic exercise may improve indices of falling risk in older women. *Age Ageing* 2002; 31: 261-6.
30. Mendell JL, Alexander NB. A clinical measure of maximal and rapid stepping in older women. *J Gerontol* 2000; 55A: M429-33.

## Cutoff and Target Values for Intra-Abdominal Fat Area for Prevention of Metabolic Disorders in Pre- and Post-Menopausal Obese Women Before and After Weight Reduction

Ryosuke Shigematsu, PhD; Tomohiro Okura, PhD\*; Syuzo Kumagai, PhD\*\*;  
Yuko Kai, PhD\*\*,\*\*; Teruo Hiyama, MD†; Haruka Sasaki, MD††;  
Hitoshi Amagai, MSc‡; Kiyoji Tanaka, PhD\*.§

**Background** The Japan Society for the Study of Obesity originally proposed a cutoff value of  $>100\text{ cm}^2$  for the intra-abdominal fat area (IFA) as a definition for “visceral fat obesity” in Japanese adults. There are no studies on the cutoff or target values after weight reduction in pre- and post-menopausal women.

**Methods and Results** In the present study 149 pre-menopausal obese women (PreM, 43.3 years,  $27.3\text{ kg/m}^2$ ) and 58 post-menopausal women (PostM, 53.9 years,  $27.7\text{ kg/m}^2$ ) participated in a 14-week weight reduction program. The IFA was measured by computed tomography. The program induced significant reductions in body weight (8.6 kg in PreM and 7.8 kg in PostM). The IFA decreased significantly from  $80.4\pm 41.3$  to  $50.7\pm 23.8$  (PreM) and from  $115.4\pm 38.0$  to  $75.7\pm 30.5$  (PostM).

**Conclusions** The receiver-operating characteristic curve analyses revealed that the appropriate cutoff values were  $80\text{ cm}^2$  (PreM) and  $110\text{ cm}^2$  (PostM) before the program, and after the program the appropriate target values were determined as 60 and  $70\text{ cm}^2$ , respectively. (Circ J 2006; 70: 110–114)

**Key Words:** Diet; Exercise; Fat body; Menopause; Metabolic syndrome

The “visceral fat obesity” refers to the condition of excess intra-abdominal fat (IF), which places people having this type of excess fat at high risk for obesity-related metabolic disorders, such as hyperglycemia and dyslipidemia. The Japan Society for the Study of Obesity (JASSO)<sup>1</sup> originally defined visceral fat obesity in Japanese as having an IF area (IFA)  $>100\text{ cm}^2$  and indicated that such people tend to have 1 or more metabolic disorders<sup>1</sup>. Nakamura et al reported that approximately 62% of patients with coronary artery disease have an IFA  $\geq 100\text{ cm}^2$  or more<sup>2</sup> and Banno et al found that sleep-disordered breathing was closely associated with obesity<sup>3</sup>.

JASSO used a cross-sectional study design to validate the cutoff value for IFA of  $100\text{ cm}^2$  for the diagnosis of visceral fat obesity<sup>1</sup>, but intervention studies for assessing an appropriate target value that can be used for people who reduce their IF significantly have been lacking, and it is unclear whether, or at what point, decreasing IF improves metabolic disorders.

There are several studies of the effects of menopause on the relationship of IF with metabolic diseases. Excess IF deposition is more prevalent in post-menopausal women than in pre-menopausal women<sup>4</sup> although it occurs more frequently in males of all ages<sup>5</sup>. Hunter et al<sup>6</sup> and Gower et al<sup>7</sup> showed that the IFA and the risk of coronary heart disease (CHD) were positively correlated and that each average in post-menopausal women was higher than that in pre-menopausal women. The results of the study by Rebuffe-Scrive et al<sup>8</sup> suggest that one of the reasons for this phenomenon is the more pronounced activation of lipoprotein lipase in the omental adipose tissue of post-menopausal women than in that of pre-menopausal women. The cutoff value for the IFA derived by JASSO<sup>1</sup> was defined using a combination of pre- and post-menopausal women; the standards were, therefore, not established while considering the presence of menopause.

Based on these results, the current study assesses JASSO’s visceral fat obesity IFA cutoff value of  $100\text{ cm}^2$  in pre- and post-menopausal women and also assesses the IFA target value after a weight reduction program. We tested 2 related hypotheses: (1) the cutoff value would be valid when applied to a group consisting of only pre- or post-menopausal women and (2) it would remain valid in each group after reducing the IFA.

### Methods

#### Participants

Advertisements were placed in local newspapers and on bulletin boards in Toride City in Ibaraki Prefecture and Abiko City in Chiba Prefecture in Japan to locate potential

(Received August 26, 2005; revised manuscript received October 19, 2005; accepted November 4, 2005)

Faculty of Education, Mie University, Tsu, \*Graduate School of Comprehensive Human Sciences, University of Tsukuba, Tsukuba, \*\*Institute of Health Science, Kyushu University, Fukuoka, †Division of Cardiology, Higashi Toride Hospital, Toride, ††Department of Internal Medicine, Chikushi Hospital, Fukuoka University, Chikushino, ‡Department of Orthopedic Surgery, Moriya Keiyu Hospital, Moriya, §§Physical Fitness Research Institute Meiji Yasuda Life Foundation of Health and Welfare, Tokyo and §Tsukuba Advanced Research Alliance, University of Tsukuba, Tsukuba, Japan  
Mailing address: Ryosuke Shigematsu, PhD, Faculty of Education, Mie University, 1577 Kurimamachiya, Tsu, Mie 514-8507, Japan.  
E-mail: rshige@edu.mie-u.ac.jp



participants with a desire to lose weight. Those who responded to the advertisements were interviewed by telephone. The participants supplied information on demographics, menstrual status, and medical history. They were excluded from the study if their weight had been unstable for the past 6 months, if they had attended any weight reduction programs in the past year, or if they were breast feeding or pregnant. A study physician confirmed if participants were possibly pregnant. Further, the study staff and radiologic technologist explained to all participants that computed tomography (CT) can have deleterious effects. After applying the exclusion criteria to potential participants, the selected participants (n=220) received the details of the study's purpose and protocol. Oral informed consent, following the Helsinki Declaration principles and approved by the Higashi Toride Hospital Review Board, was obtained from each person. We defined "menopause" as the status of no menses for 1 year prior to the study. "Pre-menopause" was used to define individuals who were not experiencing menopause. Therefore, the pre-menopausal group consisted of women who declared having menses in the year prior to the study (PreM). The post-menopausal group included those women who had not had menses for more than 1 year prior to the beginning of the study (PostM).

#### IFA

We measured the IFA and subcutaneous fat area (SFA) at the level of the umbilicus using cross-sectional CT (SCT-6800TX; Shimadzu, Japan). Scans were performed with the participants in the supine position. Details of the scanning have been reported by Tokunaga et al<sup>9</sup> and Yoshizumi et al<sup>10</sup>. Measurements taken before and after the program were conducted at the same time of day by the same technician to minimize technical error. The IFA and SFA were calculated using a computer-software program (FatScan; N2system, Japan).<sup>10</sup> The intra-class correlation for repeated IFA determinations in the laboratory (Institute of Health and Sport Sciences, University of Tsukuba) is 0.99 (n=30).

#### Obesity-Related Metabolic Disorders

The obesity-related metabolic disorders were defined as follows: accumulation of IF (waist circumference  $\geq 90$  cm in female) plus 2 or more co-morbidities consisting of (i) triacylglycerol (TG)  $\geq 150$  mg/dl and/or high-density lipoprotein cholesterol (HDL-C)  $< 40$  mg/dl, (ii) systolic blood pressure (SBP)  $\geq 130$  mmHg or diastolic blood pressure (DBP)  $\geq 85$  mmHg, or (iii) fasting plasma glucose  $\geq 110$  mg/dl.<sup>11,12</sup> These biochemical assays were performed on approximately 10 ml of blood drawn from each participant after an overnight fast. The blood assays were analyzed by technicians at the Koto Biken Research Institute in Tsukuba, Japan. Total body composition was assessed by bioelectrical impedance methods.<sup>13</sup> We used the Tanaka formula<sup>13</sup> to estimate the total body density (Db) and the Brozek formula<sup>14</sup> to determine the percentage of body fat. The Tanaka formula accurately predicts the total Db in obese Japanese women (R=0.903, SEE=0.0061 g/cm<sup>3</sup>, with the hydrodensitometrically determined Db). SBP and DBP were taken from the right arm using a mercury manometer after at least a 20-min rest while seated. Cuff sizes were selected based on upper arm girth and length.

#### Weight Reduction Program

A 14-week weight reduction program was monitored by

a physician, dietician, exercise instructors, and graduate school students majoring in exercise intervention. After the baseline assessment, participants received instruction on the diet program, which comprised weekly 90-min diet consultations, at which a diet-recording notebook and several handouts were given to participants to help them adhere to the principles of the daily diet. They were asked to take a well-balanced supplemental food product (MicroDiet; Sunny Health Co, Ltd, Japan) daily as 1 of their meals, preferably as lunch or dinner. The MicroDiet, which includes various amino acids, vitamins, and minerals, was developed for very low-energy diets. To prevent boredom, the MicroDiet was served in 7 flavors: coffee, milk tea, cocoa, yogurt, banana, strawberry, and apple. Participants received packages consisting of 7 meals (each flavor) once a week. The nutritional values for each flavor were slightly different (ie, there was a range for protein (20.6–21.5 g), carbohydrate (15.0–18.1 g), fat (1.6–3.0 g), and energy (169–173 kcal) for each meal). The diet records were obtained from 86 participants (60 in the PreM group, 26 in the PostM group), who were randomly selected. One week before the study, the participants were asked to record everything they had eaten for the 3 days prior to the study. Furthermore, they were asked to record their diets for 3 days during week 7, the midpoint of the intervention.

The exercise program included 3 weekly 45-min sessions. During the first and second weeks of the 14-week program, exercise sessions consisted mainly of walking and stretching, with the gradual addition of a bench-stepping exercise<sup>15</sup> as the main element. Thereafter, the exercise session consisted of a 10-min warm-up, 25-min bench stepping, and a 10-min cool-down. The bench stepping targeted an exercise intensity in which the participant's heart rate reached a level 10–15% higher than the level corresponding to her lactate threshold (LT). The LT was defined as the point at which blood lactate concentration maintained a non-linear increase above the level at rest.<sup>16</sup> To determine LT, a series of venous blood samples (1 ml each) was drawn from the antecubital vein every minute during a maximal cycling exercise test, which was done with an accompanying electrocardiogram as a baseline assessment. All blood samples were analyzed by the electrochemical enzymatic method using a lactate analyzer (model 23L, YSI Inc, OH, USA). For establishing LT, the log (oxygen uptake)–log (lactate) transformation method was used.<sup>16</sup>

Exercise was consistently performed for 45 min throughout the 14 weeks, but the intensity was progressively increased. In the first 2 weeks, the bench-stepping instructor targeted the intensity as described. After the 3<sup>rd</sup> week, the instructor progressively increased the intensity by increasing the cadence of the step and adding more dynamic movements. Ratings of the perceived exertion (RPE)<sup>17</sup> by all participants were also monitored during the bench stepping. Based on their RPE, the instructor moderated the intensity as "somewhat hard" to "hard," which corresponded to LT or a little above LT!<sup>18</sup>

#### Statistical Analysis

Differences in variables between the beginning and end of the program were tested in each group by using Student's paired t-tests. Data were analyzed with the SPSS 11.01J statistical software package (SPSS, Chicago, IL, USA), and P-values less than 0.05 were considered statistically significant.

To assess the cutoff value (before weight reduction) and

**Table 1** Baseline Characteristics of Participants

	PreM + PostM (n=207)	PreM (n=149)	PostM (n=58)
Age (years)	46.2±8.1	43.3±6.7 (24–57)	53.9±6.0 (45–62)
Height (cm)	157.0±5.2	157.9±5.1 (146.1–171.8)	154.6±4.9 (145.6–165.4)
Weight (kg)	67.6±8.2	68.1±7.6 (53.6–87.6)	66.3±9.7 (50.0–111.3)
Body mass index (kg/m <sup>2</sup> )	27.4±3.0	27.3±2.9 (21.8–37.3)	27.7±3.3 (20.9–40.7)
Percent body fat (%)	34.6±4.9	34.1±4.2 (24.9–46.7)	35.9±6.2 (24.1–51.9)
Intra-abdominal fat area (cm <sup>2</sup> )	90.2±43.3	80.4±41.3 (12.2–222.9)	115.4±38.0 (32.3–191.2)
Subcutaneous fat area (cm <sup>2</sup> )	252.2±82.1	250.9±75.4 (103.5–548.0)	255.5±97.9 (90.5–684.0)
Abdominal circumference (cm)	95.7±8.6	95.1±8.4 (73.8–118.0)	97±8.9 (80.5–131.5)

Values are means ± standard deviations (minimum–maximum).

PreM, pre-menopausal obese group; PostM, post-menopausal obese group.

**Table 2** Effects of a 14-Week Weight Reduction Program on Anthropometric Variables, Abdominal Fat Area, Metabolic Variables, and Blood Pressures

	PreM + PostM (n=207)		PreM (n=149)		PostM (n=58)	
	Before	After	Before	After	Before	After
Weight (kg)	67.6±8.2	59.3±7.4* (–12%)	68.1±7.6	59.6±6.9* (–12%)	66.3±9.7	58.5±8.5* (–12%)
Body mass index (kg/m <sup>2</sup> )	27.4±3.0	24.0±2.7* (–12%)	27.3±2.9	23.9±2.6* (–12%)	27.7±3.3	24.4±2.9* (–12%)
Percent body fat (%)	34.6±4.9	29.4±4.6* (–15%)	34.1±4.2	28.8±4.0* (–15%)	35.9±6.2	31.1±5.5* (–13%)
Intra-abdominal fat area (cm <sup>2</sup> )	90.2±43.3	57.7±28.1* (–32%)	80.4±41.3	50.7±23.8* (–31%)	115.4±38.0	75.7±30.5* (–34%)
Subcutaneous fat area (cm <sup>2</sup> )	252.2±82.1	181.6±76.9* (–29%)	250.9±75.4	176.2±73.2* (–31%)	255.5±97.9	195.5±84.6* (–24%)
Abdominal circumference (cm)	95.7±8.6	85.2±8.6* (–6%)	95.1±8.4	84.9±8.0* (–5%)	97.3±8.7	86.2±10.0* (–7%)
Fasting plasma glucose (mmol/L)	5.41±1.13	4.94±0.68* (–7%)	5.25±0.86	4.88±0.68* (–6%)	5.84±1.57	5.10±0.67* (–10%)
Total cholesterol (mmol/L)	5.71±0.95	5.17±0.89* (–9%)	5.59±0.95	5.02±0.83* (–9%)	6.00±0.89	5.57±0.92* (–7%)
Triacylglycerol (mmol/L)	1.18±0.59	0.80±0.41* (–23%)	1.13±0.59	0.76±0.38* (–24%)	1.30±0.58	0.91±0.48* (–21%)
HDLC (mmol/L)	1.70±0.38	1.65±0.33* (–1%)	1.72±0.36	1.65±0.32* (–2%)	1.66±0.44	1.65±0.35 (+2%)
SBP (mmHg)	132.4±18.8	120.6±16.5* (–8%)	129.9±17.9	118.6±15.5* (–8%)	138.7±19.7	125.8±17.7* (–9%)
DBP (mmHg)	82.1±11.7	74.4±11.0* (–9%)	81.0±11.5	74.2±10.5* (–8%)	84.9±11.9	74.8±12.4* (–12%)

Values are means ± standard deviations (relative change, %).

PreM, pre-menopausal obese group; PostM, post-menopausal obese group; HDLC, high-density lipoprotein cholesterol; SBP, systolic blood pressure; DBP, diastolic blood pressure.

\*Significant intra-group difference ( $P < 0.05$ ).

**Table 3** Number and Percentage of Participants That Exceeded Each Criterion of the Metabolic Disorders Before and After Weight Reduction Program

	PreM (n=149)		PostM (n=58)	
	Before	After	Before	After
High abdominal circumference	112 (75%)	37 (25%)	50 (86%)	18 (31%)
High triacylglycerol and/or low HDLC	27 (18%)	6 (4%)	11 (19%)	6 (10%)
High triacylglycerol	26 (17%)	5 (3%)	9 (16%)	5 (9%)
Low HDLC	3 (2%)	3 (2%)	4 (7%)	2 (3%)
High systolic and/or diastolic blood pressure	80 (54%)	36 (24%)	41 (71%)	23 (40%)
High systolic blood pressure	74 (50%)	33 (22%)	39 (67%)	23 (40%)
High diastolic blood pressure	52 (35%)	22 (15%)	28 (48%)	8 (14%)
High fasting plasma glucose	11 (7%)	6 (4%)	14 (24%)	6 (10%)

Abbreviations see in Table 2.

the target value (after weight reduction) for IFA, receiver-operating characteristic (ROC) curve analysis was applied to the data derived from the IFA and the number of metabolic disorders. By provisionally varying the cutoff/target values of IFA, we calculated the sensitivities and specificities for each value. Sensitivity was defined as the proportion of participants having a given disorder who also had an IFA equal to or greater than the provisional value to all participants having a given disorder. Specificity was defined as the proportion of participants having no disorders who had an IFA that fell below the provisional value to all participants having no disorders. The sensitivities and specificities were calculated for every 10 cm<sup>2</sup> of IFA from 30 to 140 cm<sup>2</sup>. At each 10 cm<sup>2</sup> provisional value, the sensitivity was multiplied by the specificity, and the point having the maximum

product of sensitivity × specificity was considered to be the most valid cutoff/target value.

## Results

Of the 220 women originally enrolled in this study, 13 dropped out because they moved out of the area, needed to care for a family member, or felt fatigued. Consequently, 207 women completed the study (Table 1), and attendance averaged 92% (range 83–100%).

There were significant decreases in the anthropometric variables, IFA, SFA, metabolic variables, and blood pressures in each group (Table 2). Total body composition analysis revealed that the reduction in body weight was mostly from loss of body fat. The reduction in fat-free mass



**Table 4** Sensitivities and Specificities From Each Provisional Cutoff/Target Value of Intra-Abdominal Fat Area (IFA)

Cutoff/target value (IFA, cm <sup>2</sup> )	PreM				PostM			
	Before		After		Before		After	
	Sensitivity	Specificity	Sensitivity	Specificity	Sensitivity	Specificity	Sensitivity	Specificity
30	0.96	0.04	0.93	0.22	1.00	0.00	1.00	0.06
40	0.91	0.23	0.87	0.42	1.00	0.06	1.00	0.11
50	0.89	0.37	0.73	0.57	0.98	0.12	1.00	0.3
60	0.80	0.47	<u>0.67</u>	<u>0.71</u>	0.95	0.18	0.91	0.40
70	0.72	0.55	0.33	0.82	0.93	0.29	<u>0.91</u>	<u>0.63</u>
80	<u>0.63</u>	<u>0.74</u>	0.27	0.89	0.90	0.41	0.82	0.66
90	0.53	0.81	0.07	0.96	0.88	0.59	0.64	0.72
100	0.45	0.89	0.07	0.98	0.85	0.59	0.64	0.79
110	0.40	0.95	0.07	0.99	<u>0.73</u>	<u>0.71</u>	0.45	0.96
120	0.27	0.96	0.07	0.99	0.56	0.82	0.36	0.98
130	0.21	0.97	0.07	1.00	0.41	0.94	0.18	1.00
140	0.16	0.97	0.00	1.00	0.29	0.94	0.09	1.00

Abbreviations see in Table 1.

Underlined values indicate the most valid cutoff/target values.

was significant, but the absolute change was less than the change in fat mass.

The daily average energy intake in the PreM group was 2,100±354 kcal at 1 week before the study and it decreased significantly to 1,163±242 kcal. The PostM group significantly reduced their energy intake from 1,870±394 kcal to 1,029±152 kcal. The daily protein intake in the PreM group was 78.1±15.1 g, and it decreased significantly to 70.3±14.2 g. In the PostM group, it decreased significantly from 86.1±33.5 g to 65.1±9.0 g. The daily fat intake decreased significantly from 66.3±14.9 g to 33.1±9.9 g in the PreM group and from 56.6±20.4 g to 27.4±6.2 g in the PostM group. The daily carbohydrate intake also decreased significantly from 285.5±61.2 g to 147.5±31.3 g in the PreM group and from 272.3±94.1 g to 136.1±23.8 g in the PostM group.

The percentage of participants that exceeded each criterion of the metabolic disorders is shown in Table 3. More than 50% of the participants had a high abdominal circumference before the program (PreM, 75%; PostM, 86%). The most frequent disorder in both groups was hypertension, with hyper-SBP (PreM, 50%; PostM, 67%) and hyper DBP (PreM, 35%; PostM, 48%). After the program, the percentages of all disorders, except for hypo-HDLc in the PreM group, decreased.

The characteristics of the 12 provisional cutoff/target values for IFA from 30 cm<sup>2</sup> to 140 cm<sup>2</sup> are presented in Table 4. Sensitivities before the program ranged from 0.16 to 0.96 in the PreM group and from 0.29 to 1.00 in the PostM group. Specificities ranged from 0.04 to 0.97 for the PreM group and from 0.00 to 0.94 for the PostM group. The products obtained by multiplying the sensitivity by the specificity at each provisional value ranged from 0.04 to 0.47 in the PreM group and from 0.00 to 0.52 in the PostM group. The largest products of sensitivity and specificity were found at 80 cm<sup>2</sup> (0.47) for the PreM group and 110 cm<sup>2</sup> (0.52) for the PostM group. Therefore, the cutoff values with the best equilibrium between sensitivity and specificity approached 80 cm<sup>2</sup> in the PreM group and 110 cm<sup>2</sup> in the PostM group before weight reduction. Using the same method of analysis, the most valid target values after the weight reduction program were determined to be 60 cm<sup>2</sup> for the PreM group and 70 cm<sup>2</sup> for the PostM group.

## Discussion

In only a few studies, attempts have been made to determine the cutoff or target value for obesity-related metabolic disorders!<sup>19,20</sup> In the present study the cutoff values of IFA were 80 cm<sup>2</sup> for pre-menopausal women and 110 cm<sup>2</sup> for post-menopausal women before weight reduction, which are similar to the 100 cm<sup>2</sup> value considered appropriate by JASSO<sup>1</sup> in a study that did not differentiate between pre- and post-menopausal women. Williams et al, in a combined study of both pre- (n=133) and post-menopausal women (n=87), concluded that 110 cm<sup>2</sup> was the cutoff value for IFA above which the risk of metabolic disorders increases!<sup>20</sup> Despres and Lamarche indicated that 130 cm<sup>2</sup> of IFA was the point at which the metabolic risks increase significantly, derived from a sample of 115 males and 72 females!<sup>19</sup> Considering those findings, the cutoff values in the current study seem to be reasonable.

A difference of 30 cm<sup>2</sup> in the cutoff values was noted between the PreM women (80 cm<sup>2</sup>) and the PostM women (110 cm<sup>2</sup>) before weight reduction. Williams et al reported that menopause has little effect on the risks of metabolic disorders, such as HDLC, TG, SBP, DBP, and TC:HDLC ratio,<sup>20</sup> although in a review by Knopp,<sup>21</sup> post-menopausal women were found to have elevated risks because of decreased estrogen contributing to increased low-density lipoprotein cholesterol (LDLC) and decreased HDLC concentrations. Hunter et al<sup>6</sup> have also reported that post-menopausal women showed a greater IFA than pre-menopausal women and that menopausal status was significantly related to an increased risk for CHD risk factors (ie, LDLc, TC:HDLC ratio). Therefore, in the current study, the cutoff values were expected to differ according to the menopausal status of the participants. Because estrogen decreases the risk of CHD during the pre-menopausal period, perhaps counterbalancing some of the CHD risks brought on by excess IF<sup>21</sup> further assumptions were made that the cutoff value for PreM women would be the same or even greater than that of PostM women. The ROC analyses revealed a difference of 30 cm<sup>2</sup> between the cutoff values in each group, but the value of the PreM group was lower than that of the PostM group. The study from the Women's Health Initiative also showed that estrogen would not confer benefits for preventing CHD among women with estrogen plus progestin therapy relative to women given a placebo!<sup>22</sup>

There seem to be other factors in addition to estrogen affecting the risk of metabolic disease; for example, aging, which correlates to an increase in IFA<sup>2,6,23</sup> and adiponectin<sup>24</sup> may be a factor.

In previous studies, a cross-sectional design was used to determine an IFA cutoff value<sup>1,19,20</sup> but because it is also important to determine a target IFA value for reducing the risk of metabolic disease, an intervention design was used in the current study. The IFA relates to the risk of obesity-related metabolic disorders; therefore, we assumed that the target values after weight reduction would remain the same as before the program, but they were lower. Although the reasons for this are unclear, we speculate that once a person is suffering from a metabolic disorder, a significant reduction in IFA may not be enough in itself to ameliorate the situation.

#### Study Limitations

The reasons for the relatively low sensitivities and specificities derived from IFA and metabolic disorders are unclear. Some unmeasured factors, such as diet and the genetic effect of metabolic disorders, may play a part. Furthermore, homeostasis was not maintained during and just after the weight loss. Another limitation is that the number of participants was small and that the mean body mass index or IFA was not very high, although most participants were obese. Future studies should include a larger number of extremely obese participants to verify the target values for risk of IFA after weight reduction. A significant decrease in HDLC in the PreM group was found after weight reduction, which may have been caused by the diet. Hagan et al<sup>25</sup> reported that HDLC decreased as middle-aged women lost body weight during a 12-week diet program. The significant decrease in TC could be attributed to the fact that TC includes HDLC.

In conclusion, this study presents the cutoff values for IFA in both pre- and post-menopausal obese women, as well as the target values after weight reduction, which are useful for the diagnosis of obesity-related metabolic disorders. Before weight reduction, the cutoff values with the best equilibrium were 80 cm<sup>2</sup> for pre-menopausal women and 110 cm<sup>2</sup> for post-menopausal women. After weight reduction, the target values shifted to 60 cm<sup>2</sup> and 70 cm<sup>2</sup>, respectively. Using these values, persons diagnosed with visceral fat obesity can clearly see the benefits of engaging in a diet and exercise program. Furthermore, awareness of a target value makes adherence to the program more likely.

#### Acknowledgments

This work was supported in part by Grants-in-Aid from the Japanese Society of Physical Fitness and Sports Medicine (1998–2000), by the Tanaka Project of the Japanese Society of Physical Fitness and Sports Medicine (1998–2000), by the Tanaka Project (1999–2002) of TARA (Tsukuba Advanced Research Alliance) at University of Tsukuba, and by the 21st century COE (Center of Excellence) program, Ministry of Education, Culture, Sports, Science and Technology (2002–2003 Nishihira Project: Promotion of health and sport scientific research). The MicroDiet was provided by the Sunny Health Co Ltd.

#### References

- Examination Committee of Criteria for 'Obesity Disease' in Japan, Japan Society for the Study of Obesity. New criteria for 'obesity disease' in Japan. *Circ J* 2002; **66**: 987–992.
- Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K, et al. Contribution of visceral fat accumulation to the development of coronary artery disease in non-obese men. *Atherosclerosis* 1994; **107**: 239–246.
- Banno K, Shiomi T, Sasanabe R, Otake K, Hasegawa R, Maekawa M, et al. Sleep-disordered breathing in patients with idiopathic cardiomyopathy. *Circ J* 2004; **68**: 338–342.
- Ley CH, Lees B, Stevenson JC. Sex- and menopause-associated changes in body-fat distribution. *Am J Clin Nutr* 1992; **55**: 950–954.
- Björntorp PA. Sex differences in the regulation of energy balance with exercise. *Am J Clin Nutr* 1989; **49**: 958–961.
- Hunter GR, Kekes-Szabo T, Treuth MS, Williams MJ, Goran M, Pichon C. Intra-abdominal adipose tissue, physical activity and cardiovascular risk in pre- and post-menopausal women. *Int J Obes Relat Metab Disord* 1996; **20**: 860–865.
- Gower BA, Nagy TR, Goran MI, Toth MJ, Poelman ET. Fat distribution and plasma lipid-lipoprotein concentrations in pre- and post-menopausal women. *Int J Obes Relat Metab Disord* 1998; **22**: 605–611.
- Rebuffe-Scrive M, Andersson B, Olbe L, Björntorp P. Metabolism of adipose tissue in intraabdominal depots of nonobese men and women. *Metabolism* 1989; **38**: 453–458.
- Tokunaga K, Matsuzawa Y, Ishikawa K, Tarui S. A novel technique for the determination of body fat by computed tomography. *Int J Obesity* 1983; **7**: 437–445.
- Yoshizumi T, Nakamura T, Yamane M, Islam AH, Menju M, Yamasaki K, et al. Abdominal fat: Standardized technique for measurement at CT. *Radiology* 1999; **211**: 283–286.
- Igarashi K, Fujita K, Yamase T, Norita N, Okita K, Satake K, et al. Sapporo Fitness Club Trial (SFCT): Design, recruitment and implementation of a randomized controlled trial to test the efficacy of exercise at a fitness club for the reduction of cardiovascular risk factors. *Circ J* 2004; **68**: 1199–1204.
- The Examination Committee of Criteria for 'Metabolic Syndrome' in Japan. Criteria for 'metabolic syndrome' in Japan. *J Jpn Soc Intern Med* 2005; **94**: 188–203 (in Japanese).
- Tanaka K, Nakadomo F, Watanabe K, Inagaki A, Kim HK, Matsuura Y. Body composition prediction equations based on bioelectrical impedance and anthropometric variables for Japanese obese women. *Am J Hum Biol* 1992; **4**: 739–745.
- Brozek J, Grande F, Anderson JT, Keys A. Densitometric analysis of body composition: Revision of some quantitative assumptions. *Ann NY Acad Sci* 1963; **110**: 113–140.
- Olson MS, Williford HN, Blessing DL, Greathouse R. The cardiovascular and metabolic effects of bench stepping exercise in females. *Med Sci Sports Exerc* 1991; **23**: 1311–1317.
- Beaver WK, Wasserman K, Whipp BJ. Improved detection of lactate threshold during exercise using a log-log transformation. *J Appl Physiol* 1985; **59**: 1936–1940.
- Borg G. Perceived exertion: A note on "history" and methods. *Med Sci Sports* 1973; **5**: 90–93.
- Hill DW, Cureton KJ, Grisham C, Collins MA. Effect of training on the rating of perceived exertion at the ventilatory threshold. *Eur J Appl Physiol* 1987; **56**: 206–211.
- Despres JP, Lamarche B. Effects of diet and physical activity on adiposity and body fat distribution: Implications for the prevention of cardiovascular disease. *Nutr Res Rev* 1993; **6**: 137–159.
- Williams MJ, Hunter GR, Kekes-Szabo T, Treuth MS, Snyder S, Berland L, et al. Intra-abdominal adipose tissue cut-points related to elevated cardiovascular risk in women. *Int J Obes Relat Metab Disord* 1996; **20**: 613–617.
- Knopp RH. Risk factors for coronary artery disease in women. *Am J Cardiol* 2002; **89**: 28E–35E.
- Writing Group for the Women's Health Initiative Investigators. Risks and benefits of estrogen plus progestin in healthy postmenopausal women. *JAMA* 2002; **288**: 321–333.
- Hernandez-Ono A, Monter-Carreola G, Zamora-Gonzalez J, Cardoso-Saldana G, Posadas-Sanchez R, Torres-Tamayo M, et al. Association of visceral fat with coronary risk factors in a population-based sample of postmenopausal women. *Int J Obes Relat Metab Disord* 2002; **26**: 33–39.
- Ryo M, Nakamura T, Kihara S, Kumada M, Shibazaki S, Takahashi M, et al. Adiponectin as a biomarker of the metabolic syndrome. *Circ J* 2004; **68**: 975–981.
- Hagan RD, Upton SJ, Wong L, Whittam J. The effects of aerobic conditioning and/or caloric restriction in overweight men and women. *Med Sci Sports Exerc* 1986; **18**: 87–94.

日本臨牀 64 卷 増刊号 9 (2006 年 12 月 28 日発行) 別刷

# メタボリックシンドローム

—病因解明と予防・治療の最新戦略—

## VI. 予防・治療・管理

ライフスタイルへの介入によるメタボリックシンドロームの  
予防と治療

### 運動療法

メタボリックシンドローム診断における  
運動療法の基本コンセプト

田中喜代次 林 容市 中田由夫 大藏倫博

## ライフスタイルへの介入によるメタボリックシンドロームの予防と治療 運動療法

### メタボリックシンドローム診断における 運動療法の基本コンセプト

Exercise prescription for people with metabolic syndrome

田中喜代次<sup>1</sup> 林 容市<sup>2</sup> 中田由夫<sup>2</sup> 大藏倫博<sup>2</sup>

**Key words** : 内臓脂肪, 運動療法, 食事療法, エネルギー消費量

#### はじめに

メタボリックシンドロームとは、内臓脂肪の過剰蓄積を基盤として、血圧の上昇(正常高値)や糖・脂質代謝の障害が顕在化しつつある状態を指す<sup>1)</sup>。したがって、治療や改善のための介入に際しては、内臓脂肪を中心とした体脂肪の減少および血圧や種々の代謝障害の正常化が基本目標となる。

メタボリックシンドロームの診断基準は、①内臓脂肪型肥満を必須とするもの<sup>1,2)</sup>、②インスリン抵抗性を必須とするもの<sup>3)</sup>、③必須条件を作らないもの<sup>4,5)</sup>、の3つに大きく分類できる。そのうち、日本人を対象にしたものでは、内臓脂肪型肥満がメタボリックシンドロームの源泉として強調されている<sup>1)</sup>。これらのことから、メタボリックシンドロームの改善を目的とした場合、食事療法(主に食事制限による摂取エネルギー量の減少)とともに、運動の習慣化を通して総消費エネルギー量を増大させる組み合わせ介入が第一選択肢としてあげられよう。

#### 1. 運動だけで内臓脂肪を減らすことは可能か?

肥満を解消するための運動プログラムがテレビや新聞、雑誌などのメディアを通じて頻繁に発信されているが、果たして多くの一般人にとって運動だけで肥満を改善する(内臓脂肪を減らす)ことが可能であろうか。図1において、自転車エルゴメータ運動(またはトレッドミル歩行やウォーキング)と食事制限による体重減少量(1カ月当たり)の理論値を比較した。体重80kgの肥満者が1回30分間、4METs強度で週1回、3回、7回自転車エルゴメータ運動を行った場合、1カ月間の減量の理論値はそれぞれ-0.1kg, -0.2kg, -0.5kgである。一方、減量前の食事による1日総摂取エネルギーを2,500kcalと仮定した場合、この量の10%(250kcal/日)、20%(500kcal/日)、30%(750kcal/日)を減らしたときの1カ月間の減量の理論値はそれぞれ-1.1kg, -2.1kg, -3.2kgである。つまり、週1~3回程度の運動では体重減少効果は極めて小さく、計算上は毎日(週7回)の運動であっても1日の食事量を10%減らした場合の約半分の効果でしかない。例えば、'週3回

<sup>1</sup>Kiyoji Tanaka: Department of Sports Medicine, Graduate School of Comprehensive Human Sciences, University of Tsukuba 筑波大学大学院人間総合科学研究科 スポーツ医学 <sup>2</sup>Yoichi Hayashi, Yoshio Nakata, Tomohiro Okura: Department of Health and Sport Sciences 同 体育科学




図1 ウォーキングと食事制限による体重減少量(1カ月あたり)の理論値の比較

ウォーキングによる理論値算出に際しては、体重80kgの肥満者が1回30分間、4METs強度で行うものと仮定(条件設定)し、週1回、3回、7回の場合について1カ月間で消費されるエネルギー量に基づき、体脂肪1kgの燃焼エネルギーを7,000kcalとして計算した。また、食事制限による理論値算出は、減量前の1日総摂取エネルギーを2,500kcalと仮定し、この量の10%(250kcal/日)、20%(500kcal/日)、30%(750kcal/日)を減らした場合の体重減少量を推定した。

の自転車エルゴメータ運動またはウォーキングは臨床現場で頻繁になされる運動処方であるが、この場合の体重減少量は1日の食事量を20%減少させた場合の約1/10にすぎない。恐らく、週3回自転車エルゴメータ運動を行った程度の体重減少量では内臓脂肪の減少量も小さく、メタボリックシンドロームの根本的な改善策としての効果には疑問を呈さずにはいられない。

図2は実際に著者らが介入研究を通して収集したデータであり、運動の効果はそれほど大きくないことが理解できよう。メタボリックシンドロームの改善を企図した運動は、有酸素性運動(40~90分の低強度での運動または20~60分の中程度~やや高めの強度での運動)を週に3~7日(低強度・短時間なら7日、高め強度・長時間なら3~4日)実践することである。強度の目安は、乳酸、心拍数、血圧、二重積などから

探るのが一般的だが、多人数に効率良く運動を勧めるためには、自覚的運動強度(RPE)を利用しながら本人と指導者による協議に基づいて選定する方法も有用である。その場合、AT pointの処方ではなく、AT zoneを示すのがよい。ここでいうAT zoneとは、RPEで11(‘楽である’)~15(‘きつい’)くらいの範囲を指す。運動時間については、10分×3回と30分×1回を比べると、強度が同じ(低~中)であれば後者による効果が大きい、前者では障害を起こす確率が低いというメリットがある。このことにこだわる必要はない。

“内臓脂肪”(腹囲、中性脂肪)や“血糖”を効率良く減らすには、1日当たり1,200~1,800kcal程度の食事療法を併用することが有効かつ安全で、個人差が大きいとはいえ少量の運動だけでは限界がある。“血圧”を改善するには、食塩感受性、遺伝体質、肥満度、性格などを考慮に入




図2 減量プログラム前後における内臓脂肪面積の変化

れながら、個人に合った運動を処方することが肝要で、薬物療法との併用が最も効果的であるとの報告が増えている。“HDLコレステロール”はランナーなどの(最大酸素摂取量の高い)スポーツ選手で最も高く、運動不足の肥満男性で低く、虚血性心疾患の男性で最も低い。HDLコレステロールを高めるには、ジョギングやウォーキング(週に15km程度)、水泳、エアロビクス、ダンスなどの有酸素性運動を習慣化・日常化することである。早い人で数カ月、遅い人で数年後に数値が上昇してくる。しかし、運動量が減少し、体脂肪が増えてくると、再び低値に戻りやすい。運動は、有酸素性運動、レジスタンス運動、ストレッチ(または柔軟体操、ヨガ、ピラテスなど)とレクリエーションを上手く組み合わせることが最も適切であろう。世代、季節、天候、仲間の有無、屋内外といった条件に合わせて運動を楽しむ方法を習得することが理想である。

## 2. 運動と減量は量-反応関係にあるのか?

Ross ら<sup>6)</sup>は、1966~2000年の文献を対象に、‘weight loss’および‘exercise’をキーワードとしたMEDLINEによる検索を行い、ヒットした36の研究についてレビューを行っている。介入期間が16週未満の短期的介入研究におけるエネルギー消費量(運動量)と体重および体脂肪の変化量との関係(相関係数)は、それぞれ $r = -0.84$ と $r = -0.76$ であり、いずれも統計学的に有意(有意確率 $< 0.001$ )であったことから、短期間の介入では運動量に応じて体重が減少する量-反応(dose-response)関係にあることがうかがえる。一方、介入期間が20週以上の比較的長期間の介入においては、体重および体脂肪ともに運動量との運動はみられなかった(図3)。

Mertens ら<sup>7)</sup>は、食事制限をせずに週当たり1,500~2,000kcalの運動を数年間にわたって処方したところ、年平均の体重減少量は1%(1




図3 1週当たりのエネルギー消費量と体重および体脂肪変化量との関係(介入期間:20週以上)  
(Ross Rら(2001)<sup>9)</sup>の報告に基づき著者らが作成)

kg)にも満たなかったと報告している。彼らはこの理由として、食事によるエネルギー摂取量が介入前に比べて8%増加していたことをあげている。日本人を対象とした著者らの取り組みにおいても、同様のことを観察している<sup>8)</sup>。また、Leonら<sup>9)</sup>は運動を処方することにより介入期間中の日常生活の活動量が減少することを観察しており、結果的に食事制限を伴わない運動処方は減量に成功せず、冠危険因子を改善することができなかったと述べている。

### 3. 減量介入に性差を考慮すべきか?

食事制限と運動による体重減少量の違いに目を向けてみよう。1969~84年までに行われた減量に関する研究(メタアナリシス)<sup>10)</sup>によると、食事制限による体重減少量は平均-10.7kgであるのに対し、運動のみだと-2.9kgでしかなかった。一方、男性肥満者のみに絞って検討した研究<sup>11)</sup>では、食事制限-7.8kgに対して運動-4.6kgと大差は少なく、2つの介入方法による体重減少の違いには性差が存在することも示唆されている。日本人を対象とした著者らの長年の介入研究でも、同様のことを観察している<sup>8)</sup>。男性が本格的に運動に取り組んだ場合、運動による総消費エネルギーが大きく増え、か

つ暴飲暴食習慣をもっていた人では食事制限の効果も大きくなる。最近では、食事制限と運動の併用介入によって、女性において体重減少の累積効果(cumulative effect)が得られることも報告されており<sup>12)</sup>、少なくとも日本人女性を対象として内臓脂肪の減少を目的とした減量介入を行う際には、運動だけでなく食事制限を併用することの必要性がうかがえる。

メタボリックシンドロームの改善、特に内臓脂肪の減少を目的とする場合には、女性については運動の効果を過大評価せず、食事指導(制限)と運動(主に有酸素性運動)を適切に組み合わせる方法を今後詳細に検討していく必要があるといえる。

### 4. 効果的な運動とは?

一般に脂質燃焼を目的とした場合、最大下(submaximal)強度で行う有酸素性運動をある一定時間継続することが推奨されているが、特に内臓脂肪の減少に関しては、やや高い強度(70~80%  $\dot{V}O_2\max$ )での運動実践が有効であることが推察される<sup>13,14)</sup>。一過性の有酸素性運動時においては、50%  $\dot{V}O_2\max$ と比較して、70%  $\dot{V}O_2\max$ に相当する強度での運動時に、内臓脂肪に由来する脂質の燃焼効率が增大する可能性



が示唆されている<sup>15)</sup>。多段階漸増負荷やランブ負荷という特殊なプロトコルでの運動時には、強度が高まるにつれて脂質によるエネルギー供給の割合が低下することは周知の事実であるが、ある程度の時間、運動を継続するという一般的な様式での有酸素性運動時には、高めの強度の方が、総消費エネルギー量(kcal)は大きくなり、脂質の分解・利用の効率が多少低下しても、高い脂肪燃焼効率・少ない消費カロリーの運動時と比較して、実質の脂肪消費量(g)は多くなる。運動中に脂質や糖質が動員される割合の比較にだけ焦点を当てるのではなく、消費されるエネルギーの総量に着目した運動強度の設定や運動処方重要である。

内臓脂肪型肥満と同様に、メタボリックシンドロームのリスク要因であるインスリン抵抗性などの耐糖能異常は、有酸素性運動だけでなく、筋収縮を強調したレジスタンス運動によっても改善されることが指摘されている。最近の著者らの研究<sup>16)</sup>では、一過性の運動においてはやや高めの強度で糖代謝への効果はより増幅されることが明らかとなった。したがって、有酸素性運動とレジスタンス運動の両方を上手く組み合わせることで、より大きな改善が期待できる。また従来より、HDLコレステロールを高めるためには、中程度～高め(最低75% HRmaxまたは60%  $\dot{V}O_2\text{max}$ 以上)の強度で週に12.8~16 km(3日/週走るとすると1日約4.3~5.4 km)以上のジョギングを続けること、すなわち大きな運動量を確保することの必要性が報告されている。ここでいう高めの強度や大きな運動量とは、アスリートの競技レベルを意味するのではなく、一般の人々が実践可能な範囲内の中での‘高いレベル’を指しており、ACSMが推奨する有酸素性運動時の強度範囲を超えるものではない。一般健常者だけでなく、メタボリックシンドロームと診断された場合でも、‘高めの強度’は多くの人に適応可能なレベルといえる。

運動だけで効果を求める場合、大きな運動量を多頻度で実践することが必要となり、コンプライアンスとアドヒアランスのいずれを考える

にしても、対象者にとって継続が困難となる。冒頭で述べたように、運動のみでメタボリックシンドロームを改善しようとするのではなく、食習慣の改善と運動習慣の定着の両者を踏まえた日常生活自体の改善の気づき(脳のスイッチ‘オン’)が肝要である。

## 5. 運動処方パラダイムシフト

専門の運動指導者による導きは有効であるが、運動実践者(クライアント)の体の中で起こっている異常・違和感を運動指導者や医師が常に的確に感じ取ることは困難である。できるだけ運動に伴うケガを起こさず、習慣化につなげていくためには、周囲の者(高齢者体力づくり支援士や健康運動指導士、理学療法士などのコメディカル)が運動実践者に対して自分のからだ対話するよう、繰り返しアドバイスすることが肝要である。自分のからだの内なる声に耳を傾けるといった運動処方<sup>17)</sup>はまさに主観的であって、客観性に欠けるとの指摘を受けるが、安価で、随時可能で、時間的ズレがないということを考えれば、これ以上の事故予防チェック方法はほかにない。

また、公衆衛生分野でいわれている informed choice(多くの情報の中から個人が自分に合ったものを選択すること)を円滑に進めるための多方面からの情報提供や、自らのからだとの対話の勧めが重要である一方で、専門家の別視(蔑視、結果的に差別)に基づく不適切な運動の禁忌指令(逆支援)は避けなければならない。そのためには、専門家は有益な情報を与えつつも、それを選択するかどうかについては不必要に口を挟まないアプローチが望まれる。すなわち、個人の主体性を最優先し、‘押しつけない’あるいは‘過度に規制しない’、そして‘大いに楽しませる’柔軟な運動処方指針の啓発が必要である。

## おわりに

運動療法によってメタボリックシンドロームの改善介入を行う場合、一般に運動によって得られる2つの効果(①一時的な急性効果、②慢

性的な持続効果), および2つの生理的変化(①体脂肪の減少に代表される器質的変化, ②分子レベルへの働きかけによる機能的変化)の両側面に関する考察が必要であろう。しかし, 本稿では誌面の制約があり, 直ちに臨床現場で役立つ食事療法の有効性を強調するとともに, 運動療法の基本コンセプトに焦点を絞って述べた。

運動それ自体が内臓脂肪の減少にどれほど有効であるかについて解説しながら, 血圧や糖・脂質代謝(メタボリックシンドローム)へ与える直接的な効果, およびメタボリックシンドロームの改善にこだわらない広義の健康支援策についても言及した。

## 参考文献

- 1) メタボリックシンドローム診断基準検討委員会: メタボリックシンドロームの定義と診断基準. 日内会誌 94: 188-203, 2005.
- 2) International Diabetes Federation: Worldwide definition of the metabolic syndrome. Available at: [http://www.idf.org/webdata/docs/IDF\\_Meta\\_syndrome\\_definition.pdf](http://www.idf.org/webdata/docs/IDF_Meta_syndrome_definition.pdf), accessed March 3, 2006.
- 3) World Health Organization: Definition, Diagnosis and Classification of Diabetes Mellitus and its Complications, Part 1: Diagnosis and classification of diabetes mellitus, World Health Organization, Geneva, 1999.
- 4) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults: Executive Summary of The Third Report of The National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, And Treatment of High Blood Cholesterol In Adults (Adult Treatment Panel III). JAMA 285: 2486-2497, 2001.
- 5) Gale EA: The myth of the metabolic syndrome. Diabetologia 48: 1679-1683, 2005.
- 6) Ross R, Janssen I: Physical activity, total and regional obesity: dose-response considerations. Med Sci Sports Exerc 33(Suppl 6): S521-S527, 2001.
- 7) Mertens DJ, et al: Exercise without dietary restriction as a means to long-term fat loss in the obese cardiac patient. J Sports Med Phys Fitness 38: 310-316, 1998.
- 8) 大河原一憲ほか: 食事制限を用いた減量介入が睡眠呼吸障害と活力年齢に及ぼす効果について. 診断と治療 92: 145-149, 2004.
- 9) Leon AS, et al: Effects of 2,000 kcal per week of walking and stair climbing on physical fitness and risk factors for coronary heart disease. J Cardiopulm Rehabil 16: 183-192, 1996.
- 10) Wood PD, et al: Changes in plasma lipids and lipoproteins in overweight men during weight loss through dieting as compared with exercise. N Engl J Med 319: 1173-1179, 1988.
- 11) Miller WC, et al: A meta-analysis of the past 25 years of weight loss research using diet, exercise or diet plus exercise intervention. Int J Obes Relat Metab Disord 21: 941-947, 1997.
- 12) Dunn CL, et al: The comparative and cumulative effects of a dietary restriction and exercise on weight loss. Int J Obes Relat Metab Disord 30: 112-121, 2006.
- 13) 大藏倫博, 田中喜代次: 内臓脂肪と運動療法. 肥満研究 6: 125-129, 2000.
- 14) 佐々木誠一ほか: 高い強度の運動は脂肪利用に有効か? 肥満研究 11(増刊号): 129, 2005.
- 15) Numao S, et al: Effects of obesity phenotype on fat metabolism in obese men during endurance exercise. Int J Obes(Lond) 30: 1189-1196, 2006.
- 16) Hayashi Y, et al: A single bout of exercise at higher intensity enhances glucose effectiveness in sedentary men. J Clin Endocrinol Metab 90: 4035-4040, 2005.
- 17) 田中喜代次, 牧田 茂: 事故予防のためのセルフチェックの大切さ. 高齢社会における運動支援実践ガイド. 臨床スポーツ医学 22(臨時増刊号): 413-415, 2005.

# 肥満度と介入方法の違いが内臓脂肪型肥満者の 減量効果に及ぼす影響

魏 丞完\*<sup>1</sup>                      大藏 倫博\*<sup>2,3</sup>                      中田 由夫\*<sup>2,3</sup>  
大河原 一憲\*<sup>1</sup>                      沼尾 成晴\*<sup>1</sup>                      片山 靖富\*<sup>1</sup>  
山中喜代次\*<sup>2,3</sup>

\*<sup>1</sup>筑波大学大学院博士課程人間総合科学研究科

\*<sup>2</sup>筑波大学大学院人間総合科学研究科

\*<sup>3</sup>筑波大学先端学際領域研究センター



「肥満研究」Vol.12 No.1 47-53 2006 別刷

原一著

# 肥満度と介入方法の違いが内臓脂肪型肥満者の減量効果に及ぼす影響

魏 丞完<sup>\*1</sup>, 大藏 倫博<sup>\*2,3</sup>, 中田 由夫<sup>\*2,3</sup>, 大河原一憲<sup>\*1</sup>, 沼尾 成晴<sup>\*1</sup>, 片山 靖富<sup>\*1</sup>, 山中喜代次<sup>\*2,3</sup>

<sup>\*1</sup>筑波大学大学院博士課程人間総合科学研究科 <sup>\*2</sup>筑波大学大学院人間総合科学研究科 <sup>\*3</sup>筑波大学先端学際領域研究センター

索引用語: BMI, 減量方法, 内臓脂肪型肥満, 冠危険因子

内臓脂肪型肥満者を対象として肥満度の違い(BMI25以上30未満群とBMI30以上群)および介入方法(食事制限のみのDO群と食事制限に運動を併用したDE群)の違いが内臓脂肪減少や冠危険因子の改善に与える影響を検討することを目的とした。対象者はBMI25以上および内臓脂肪面積100cm<sup>2</sup>以上の21歳から67歳までの肥満女性145名であり, BMIに応じてDO群とDE群をそれぞれDO I群とDE I群(25≤BMI<30), DO II群とDE II群(BMI≥30)の4群に分けた。CTスキャンを用いて内臓脂肪面積(VFA)と皮下脂肪面積(SFA)を計測した。冠危険因子は安静時収縮期血圧, 安静時拡張期血圧, 総コレステロール, 中性脂肪, 高比重リポ蛋白コレステロール, 低比重リポ蛋白コレステロールと空腹時血糖を測定した。これらの項目は14週間の介入期間の前後に測定した。VFAの変化量については運動の有無とBMIの違いの交互作用が見られたが(p=0.01), SFAの変化量においては運動の有無とBMIの違いの交互作用は見られなかった。各冠危険因子の変化量に運動の有無とBMIの違いの交互作用が見られたのは空腹時血糖のみであった(p=0.02)。これらの結果から, BMIの高い内臓脂肪型肥満者は食事制限に運動を加えることによって, より効果的に内臓脂肪の減少や血糖の改善がもたらされることが示唆された。

## はじめに

世界保健機構(World Health Organization: WHO)は「成人の体重過多に対するbody mass index(BMI)による分類」をグローバルスタンダードとして, BMI25以上をPreobese, 30以上をObeseとする基準を提唱した<sup>1)</sup>。一方, 日本肥満学会による「新しい肥満の判定と肥満症の診断基準」の中では, わが国の肥満に関する民族的・人種的特異性を考慮に入れた上で, BMI25以上30未満を肥満(1度)と判定している。また, 内臓脂肪の蓄積が肥満合併症の

大きな要因であることから, BMI25以上かつcomputed tomography(CT)画像分析による内臓脂肪面積(visceral fat area: VFA)100cm<sup>2</sup>以上の内臓脂肪型肥満を「肥満症」と定義している<sup>2)</sup>。

内臓脂肪の減少を目的とするいくつかの介入研究<sup>3-5)</sup>によると, 食事制限のみの群でも食事制限に有酸素性運動またはレジスタンス運動を併用した群でも体重減少に応じて内臓脂肪が減少することが報告されている。一方, 食事制限は行わずに有酸素性運動を継続させたところ, 運動が内臓脂肪の減少に効果があったとする報告<sup>6)</sup>となかつ

たとする報告<sup>7)</sup>の両方があり, 内臓脂肪の減少に対する運動の有用性は明らかではない。

BMIは体脂肪量をよく反映することが知られている<sup>8)</sup>。先行研究<sup>9)</sup>によると, 運動による消費エネルギーが同じであっても, ベースラインの体脂肪量が多いほど体脂肪はより多く減少することが報告されている。また, 日本肥満学会は腹部CTにて計測した皮下脂肪面積(subcutaneous fat area: SFA)およびVFAとBMIとの関連について, BMIとSFAには高い相関が認められているが, BMIとVFAとの相関は弱く,

特にBMI25以上の肥満者では相関がなく、大きな個人差が存在することを示している<sup>2</sup>。しかしながら、BMIは身長と体重を計測するだけで値を求めることのできる簡便な指標であり、対象に適した介入方法を提案するための簡便なスクリーニングとして活用できる可能性がある。

そこで、本研究ではVFA 100cm<sup>2</sup>以上の肥満症女性を対象として、肥満度の違い(BMI25以上30未満群とBMI30以上群)および食事療法に運動を加えること(食事制限のみ群と食事制限に運動を併用する群)がVFA減少や冠危険因子の改善にいかなる影響を与えるかを検討することとした。

## 対象と方法

### 1. 対象者

対象者は地域情報誌を用いて募集し、14週間の食事制限または食事制限に運動実践を加えた減量プログラムを受けた368名であり、その中から日本肥満学会が設定する肥満の判定基準BMI25以上とCTスキャンによるVFA 100cm<sup>2</sup>以上の内臓脂肪型肥満であること条件を満たした21歳から67歳までの女性145名をデータ解析の対象とした。対象者に重篤な疾患を有する者は含まれておらず、本研究の目的および検査内容に関する説明を口頭および文書により行い、研究参加への同意(署名)を得た。なお、これらの研究手続については筑波大学倫理委員会において承認を得た。

本研究の対象者145名は、食事制限のみ(diet only: DO)の群51名と有酸素性運動に食事制限を加えた(diet + exercise: DE)群94名に分けられた。月経の有無については、DO群は閉経前63%、閉経後37%であり、DE群は閉経前51%、閉経後49%であった。γ検定の結果、DO群とDE群の間に有意

差はなかった。DO群およびDE群はBMIの初期値によってDO I群とDE I群(25 ≤ BMI < 30)、DO II群とDE II群(BMI ≥ 30)に分けられた。

### 2. 測定項目および測定方法

#### 1) 身体組成

体重は0.1kg単位、身長は0.1cm単位で測定し、BMIは体重(kg)を身長(m)の二乗で除することで算出した。体脂肪の測定には、Sekisui製インピーダンス計(Bio impedance SS103)を用い、測定された電気抵抗値からTanaka et al.<sup>15</sup>の成人肥満女性用の式により身体密度を求め、Brozek et al.<sup>16</sup>の式により体脂肪量(fat mass)、体脂肪率(% fat mass)と除脂肪量(fat-free mass)を算出した。

#### 2) 腹部脂肪面積

VFAおよびSFAは仰臥位でCTスキャン(SCT-6800TX, Shimadzu)を用いて膈高位(およそL4-L5)を撮影し、内臓脂肪計測ソフトFat Scan(ver. 2.0, N2システム)を用いて算出した。

#### 3) 冠危険因子

安静時収縮期血圧(systolic blood pressure: SBP)と安静時拡張期血圧(diastolic blood pressure: DBP)は20分以上安静にした後で計測した。12時間以上の絶食絶飲状態で採血し、総コレステロール(total cholesterol: TC)、中性脂肪(triglycerides: TG)、高比重リポ蛋白コレステロール(high-density lipoprotein-cholesterol: HDL-C)、低比重リポ蛋白コレステロール(low-density lipoprotein-cholesterol: LDL-C)、空腹時血糖(fasting plasma glucose: FPG)を測定した。

#### 4) 最大酸素摂取量

最大酸素摂取量( $\dot{V}O_{2max}$ )および無酸素性代謝閾値(anaerobic threshold: AT)を測定するための運動負荷テストは、自転車エルゴメータ(818E, Monark)を使用し、1分ごとに摩擦負

荷を0.25kpずつ高める多段階漸増負荷法にて行った。ペダルの回転数は60rpmで一定とした。運動中の各呼気ガス指標の分析には、Mijnhardt製代謝測定装置(Oxycon Alpha)を用いた。 $\dot{V}O_{2max}$ は症候性限界を呈した時点の酸素摂取量と定義した。DE群に対する運動指導における強度の目安となるATは、漸増負荷運動時1分ごとに正中肘皮静脈より血液を約0.5mlずつ採取し、血中乳酸濃度の分析(YSI製乳酸分析器1500L)により決定される乳酸閾値(lactate threshold: LT)とした。採血が困難であった際は、 $\dot{V}O_{2}$ に対する $\dot{V}CO_{2}$ の上昇開始点(V-slope法)により決定し、その他の呼気ガス指標とともに、Okura & Tanaka<sup>17</sup>による全身持久性体力推定式を補助的に利用した。

### 3. 食事指導および運動指導内容

#### 1) 食事指導

DO群とDE群ともに1食あたり400~600kcalのバランスのとれた食事を4群点数法<sup>18</sup>を用いて指導した。4群点数法は、食品に含まれている栄養素と体内での働きによって、食品を4つの群に分類し、各群から食品を選択することによって、必要な栄養素が過不足なく満たされるようにする栄養計算法である。1群は卵や乳製品、2群は肉や魚介類や豆製品、3群は緑黄色野菜、淡色野菜や芋類、果物、4群は穀物や油、砂糖や調味料である。計算単位は点数で表し、80キロカロリー(kcal) = 1点として計算させた。

本研究は1食(400kcal = 5点)ごとに「1群: 2群: 3群: 4群」の点数比率を「1: 1: 1: 2」になるよう指導し、1日あたりの総摂取エネルギーを1,200kcalとした。対象者には1食ごとの食事内容を記録させ、教室参加時に提出させた。食事記録をもとに摂取エネルギー量を確認するとともに、食事のとり方や栄養バランス、食習慣など