

we utilize CGA more intensively for the care of elderly patients in our hospitals, CGA will have a more beneficial effect on outcomes.<sup>5</sup> Although CGA for inpatients is very important to improve the outcomes of elderly patients, the focus of recent investigations has been shifted to outpatient CGA due to high cost of inpatient care in the United States.<sup>6</sup> One study showed that outpatient CGA helps maintain functioning and the ability to perform daily activities.<sup>7</sup> However, its benefits are not consistently demonstrated or recognized.

Although survival is one of the most commonly reported outcomes in clinical studies, Stuck *et al.* reported in a meta-analysis that outpatient CGA did not improve survival compared to usual care despite the fact that significant survival benefits were observed in inpatients and home-based CGA.<sup>8</sup> Two of the four trials of outpatient CGA in the meta-analysis were, however, criticized because the subjects were relatively healthy and not at high risk. Additional studies of outpatient CGA were therefore conducted with greater attention to targeting frail subjects, resulting in improved outcomes in elderly patients including mental health<sup>9-11</sup> and functional status.<sup>6,7</sup> Although outpatient CGA has, so far, no demonstrable benefit for the survival of older, frail patients compared to usual care, outpatient CGA should be a good way to assess elderly patients, to diagnose patients with mild cognitive impairment or depressive symptoms, and to eventually prevent functional decline. However, additional measurement might be required to improve the survival.

Cognitive impairment in elderly patients is sometimes hard to diagnose in the outpatient clinic. Therefore, screening elderly patients by Mini-Mental State Examination (MMSE) is useful to diagnose the initial phase of dementia or mild cognitive impairment, although it takes time to screen all the patients with MMSE. Screening depression is also helpful for the care of elderly patients and a 15-item Geriatric Depression Scale (GDS-15) is commonly used for that purpose. This test also takes time in the outpatient clinic. Finding a marker for early diagnosis of cognitive impairment or depressive mood therefore would be important to select high-risk patients.

Chronic brain inflammation characterizes Alzheimer's disease (AD), the most of common neurodegenerative disease associated with progressive cognitive decline. Certain cytokines, such as interleukin (IL)-1, IL-6, and tumor necrosis factor (TNF)- $\alpha$ , are shown to influence a number of different mechanisms that can induce or accelerate the development of neurodegeneration, indicating a correlation of inflammation with cognition.<sup>12,13</sup> Because high-sensitivity C-reactive protein (hs-CRP) can reflect the presence of inflammation and can be induced by these cytokines, we chose hs-CRP as a candidate marker for screening patients with cognitive impairment.

Androstenedione and dehydroepiandrosterone (DHEA) are produced in the biosynthetic pathway of androgen and estrogen. The sulfate ester of DHEA is DHEA-S. The decline of DHEA has been pursued as a major factor in the development of age-associated disorders.<sup>14</sup> Among the studies investigating the effect of DHEA-S on mood in the elderly, some studies show that DHEA-S improves mood<sup>15,16</sup> while others do not.<sup>17,18</sup> Therefore, the effect of these steroids on neurodegenerative diseases remains inconclusive. Malondialdehyde-low density lipoproteins (MDA-LDL) are associated with oxidatively-modified products of LDL and can be associated with atherosclerotic disease.<sup>19</sup> Therefore, if MDA-LDL is associated with cognitive impairment or depressive mood, controlling risk factors for atherosclerotic disease might be important.

From these findings we assessed the activities of daily living (ADL), cognitive functions and depressive symptoms in elderly patients who visited our outpatient clinic for the first time and examined the correlation with various serum markers listed above. We also used 'Get up and go', 'Button scores' and 'Functional reach' to assess neurobehavioral functions in these patients. By these measurements, we should be able to select high-risk patients for cognitive and functional decline and eventually would be able to use outpatient CGA to improve survival of elderly patients.

## Methods

### Subjects

All elderly (basically 65 years or older) patients who came to Kyoto University Hospital for the first time or had not been seen for the past 6 months in this hospital were asked to attend for health problem screening. We started outpatient CGA in May 2001. One hundred and forty-five consecutive patients aged 62 and older (mean age  $\pm$  SD: 75.6  $\pm$  0.56) who visited the outpatient clinic from May 2001 through March 2004 were enrolled for this study after written informed consent was taken from each patient or his/her family member. The study protocol was approved by the Ethical Committee of Kyoto University School of Medicine.

### Measurements

Comprehensive Geriatric Assessment (CGA) was done on the day of patient visit by experienced speech therapists after history taking and physical examination were done. The CGA included height, body weight, blood pressure, basic activities of daily living (BADL), which was measured with the Barthel Index. For higher-level functional capacity, each subject's independence was rated by the Tokyo Metropolitan Institute of Gerontology (TMIG) Index of competence.<sup>20</sup> This

assessment consists of a 13-item index including three sublevels of competence: (i) instrumental self-maintenance; (ii) intellectual activities; and (iii) social role. MMSE was used to assess cognitive functions. Neurobehavioral functions were assessed by the Kohs block design (KBD) test,<sup>21</sup> 'Get up and go' and 'Button scores'. A cutoff point of 12 for KBD test was used as described.<sup>21</sup> Functional reach was also determined as described.<sup>22</sup> Briefly, each subject was positioned next to the wall with one arm raised 90° with fingers extended, and a yardstick was mounted on the wall at shoulder height. The distance in centimeters that a subject was able to reach forward from an initial upright posture to the maximal anterior leaning posture without moving or lifting the feet was measured by visual observation of the third finger tip against the mounted yardstick. The distances of two trials were averaged as the functional reach score, with a greater distance indicating better balance ability.

We screened depressive symptoms using the Japanese version of GDS-15.<sup>23</sup> Higher scores of GDS-15 indicate a greater degree of depressive mood. In this study we used a cutoff point of 9. Therefore, we defined depression as a GDS-15 score of 9 or more.

#### 'Get up and go'

This test of balance is commonly used to examine functional mobility in elderly subjects.<sup>24</sup> The test requires the subject to stand up, walk 3 m (10 ft), turn, walk back and sit down. The time to complete the test is strongly correlated with functional mobility. Elderly people who can complete the test in less than 20 s are independent in transfer tasks, which are normal activities in daily living.

#### 'Button scores'

'Button scores' evaluate manual dexterity using a panel with combinations of 10 hooks, 10 big buttons and five small buttons. There were three discrete measurements of time recorded for each participant (10 hook-ons, 10 big button-on-and-offs, and five small button-on-and-offs). Total manual dexterity time in seconds, defined as the Button Score, was calculated by adding the average times for one hook-on and one big or small button-

on-and-off.<sup>25,26</sup> A cutoff point of 17 was used for the analysis.

#### Serum marker measurement

Serum levels of DHEA, DHEA-S, MDA-LDL and hs-CRP were measured by SRL (Tokyo, Japan). DHEA and DHEA-S were measured by radioimmunoassay. MDA-LDL was measured by enzyme-linked immunosorbent assay (ELISA). Hs-CRP was measured with CardioPhase kit (Dade Behring, Tokyo, Japan).

#### Statistical analysis

Differences in continuous variables among the disease groups were determined by one-way analysis of variance (ANOVA). A *P*-value of less than 0.05 was considered significant. Multiple regression analysis was used to assess the involvement of age and sex.

## Results

Table 1 summarizes the patient characteristics in the study population. The mean age in this study group was 75.6 years and the percentage of males was 40%. There was no statistical difference in age between males and females. The ADL of the patients was relatively well preserved. The mean Barthel index (0–100) was 98.3 and was not statistically different between males and females. Instrumental ADL was assessed by the Tokyo Metropolitan Institute of Gerontology Index (TMIG Index) (0–13). The mean value was 10.3 and was not statistically significant between males and females either. We assessed depression by GDS-15 and found that the mean score was 5.23. The GDS scores were slightly higher in females than in males, but the difference was not statistically significant. The mean score was almost comparable to that of community-dwelling elderly people in Japan.<sup>27</sup>

We then determined the cognitive function by MMSE and found that the mean scores were 25.2 (Table 2). We also determined the KBD test to assess spatial recognition and found that the mean score was 22.3. 'Get up and go' and 'Button scores' were assessed and the mean time to be required was 14.9 and 12.1 s, respectively. The mean length of functional reach in these patients

**Table 1** Mean age, Barthel index, Tokyo Metropolitan Institute of Gerontology (TMIG) index and Geriatric Depression Scale (GDS) scores in males and females

	<i>n</i>	Age	Barthel index	TMIG index	GDS
Total	145	75.6 ± 0.56	98.3 ± 0.61	10.3 ± 0.25	5.23 ± 0.31
Male	58	74.8 ± 0.90	99.5 ± 0.38	10.5 ± 0.37	4.51 ± 0.45
Female	87	75.9 ± 0.72	98.3 ± 0.98	10.1 ± 0.33	5.70 ± 0.41

Data are expressed as mean ± SEM.

Table 2 Mean MMSE, KBS, button scores, Get up and go, and functional reach in males and females in this population

	MMSE	KBS	Button score	'Get up and go'	Functional reach
<i>n</i>	142	133	134	131	125
Total	25.2 ± 0.46	22.3 ± 1.13	12.1 ± 0.45	14.9 ± 0.46	23.4 ± 0.66
Male	25.7 ± 0.77	24.0 ± 1.88	12.8 ± 0.62	14.1 ± 0.74	26.2 ± 0.84*
Female	25.0 ± 0.56	21.0 ± 1.39	11.6 ± 0.62	15.5 ± 0.57	21.3 ± 0.66*

Data are expressed as mean ± SEM. \**P* < 0.01. *n*, number of patients studied.

Table 3 Mean levels of dehydroepiandrosterone (DHEA), DHEA-S, malondialdehyde-low density lipoproteins (MDA-LDL) and high-sensitivity C-reactive protein (hs-CRP). Difference in male and female patients

	DHEA (ng/ml)	DHEA-S (ng/ml)	MDA-LDL (U/L)	hs-CRP (ig/ml)
Total	2.08 ± 0.10	777 ± 49.3	147 ± 5.84	4.98 ± 1.56
Male	2.02 ± 0.15	995 ± 93.8*	128 ± 8.31**	3.42 ± 1.66
Female	2.12 ± 0.13	625 ± 45.3*	158 ± 7.62**	5.79 ± 2.22

Data are expressed as mean ± SEM. \**p* < 0.01, \*\**p* < 0.01, male vs female.

was 23.4 cm. These values were also comparable to the data of community-dwelling elderly in Japan.<sup>28</sup>

We next measured the serum levels of DHEA, DHEA-S, MDA-LDL and hs-CRP in this population. The mean value of DHEA, DHEA-S, MDA-LDL and hs-CRP were 2.08 ng/mL, 777 ng/mL, 147 U/L and 4.98 µg/mL, respectively (Table 3). DHEA-S was higher and MDA-LDL was lower in males than in females. However, there was no statistical difference in DHEA or hs-CRP in males and females. Figure 1 A shows the age-dependent decrease of DHEA-S in this population. DHEA-S and age were negatively correlated (the coefficient was -0.4). DHEA also showed an age-dependent decline in this population, but the coefficient was -0.2 (Fig. 1b). MDA-LDL and hs-CRP did not show age-dependent changes in this population (data not shown).

To determine the association of hs-CRP with the cognitive function in the elderly, we examined the correlation to MMSE, KBD and 'Button scores'. We divided the patients into two groups according to the points of MMSE (cutoff; 24), KBD (cutoff; 12), and Button scores (cutoff; 17). We found that the level of hs-CRP was significantly higher in the patients with lower MMSE and KBD, and higher button scores (Fig. 2a). These differences were significant by multiple regression analysis after adjusting for age and sex. These results indicate the association of hs-CRP with cognitive and functional impairment. However, the level of total cholesterol, high-density cholesterol (HDL-C) or albumin was not statistically different between each group studied (data not shown). Although the level of hs-CRP was also higher in the patients who took longer time to complete 'Get up and go', the difference was not statistically significant. The levels of DHEA, DHEA-S or MDA-LDL





Figure 1 Age-dependent decrease of dehydroepiandrosterone (DHEA)-S and DHEA in elderly patients. Relationship between age and serum levels of (A) DHEA-S or (B) DHEA in the study patients is shown. The Y-axis is shown as natural log of (A) DHEA-S or (B) DHEA.



**Figure 2** Levels of high-sensitivity C-reactive protein (hs-CRP) and DHEA-S in study patients. (A) Hs-CRP and (B) DHEA-S were measured in patients at the first visit to Kyoto University hospital after informed consent was taken. Patients were divided into two groups according to the level of each test. Patients were divided into two groups according to the score of Mini-Mental State Examination (MMSE); 24 and more, and less than 24, time for Kohs block design (KBD); less than 12 and 12 and more, 'Button scores'; less than 17 and 17 and more, the time required for 'Get up and go'; less than 20 and 20 and more, Geriatric Depression Scale (GDS); less than 9 and 9 and more. Values are the mean  $\pm$  SEM. Number of the patients in each group is shown in each column. \* $P < 0.05$ , \*\* $P < 0.01$ .

were, however, not associated with these tests (Table 4, Fig. 2b).

In contrast, the levels of DHEA-S were significantly lower in the patients with higher GDS scores (9 or over). These differences were also significant by multiple regression analysis after adjusting for age and sex ( $P < 0.05$ ). In contrast, the other markers, including hs-CRP, were not associated with GDS scores (Fig. 2b).

Among the patients with lower MMSE (less than 24), 52.6% had dementia while only 4.1% had dementia among the patients with normal MMSE (24 or over). As

**Table 4** Mean dehydroepiandrosterone (DHEA) and malondialdehyde-low density lipoproteins (MDA-LDL) levels in each group of patients

	MMSE		KBD		'Button score'		'Get up and go'		GDS	
	$\leq 24$	$> 24$	$\leq 12$	$> 12$	$\geq 17$	$< 17$	$> 20$	$\leq 20$	$> 9$	$\leq 9$
DHEA (ng/ml)	2.07 $\pm$ 0.11	2.11 $\pm$ 0.22	2.11 $\pm$ 0.11	2.12 $\pm$ 0.23	2.08 $\pm$ 0.10	1.98 $\pm$ 0.16	2.14 $\pm$ 0.11	1.78 $\pm$ 0.18	2.10 $\pm$ 0.11	2.04 $\pm$ 0.21
MDA-LDL (U/L)	146 $\pm$ 6.67	150 $\pm$ 12.0	151 $\pm$ 7.00	134 $\pm$ 11.3	150 $\pm$ 6.54	145 $\pm$ 17.2	152 $\pm$ 7.38	141 $\pm$ 12.0	147 $\pm$ 6.56	145 $\pm$ 13.1

Data are expressed as mean  $\pm$  SEM. MMSE, Mini-Mental State Examination; KBD, Kohs block design; GDS, Geriatric Depression Scale.

a risk factors for stroke, hypertension was found in 26.3% of the patients with lower MMSE, while 32.0% of the patients with normal MMSE had hypertension. Other risk factors, such as diabetes mellitus and hyperlipidemia were found in less than 5% of the patients in both groups. In terms of GDS scores, 37.9% of the patients with high scores (nine or over) were diagnosed with depression, while only 5.4% of the patients with low scores (less than 9) were diagnosed with depression. The incidence of dementia was 20.7% and 15.2% in each group, respectively.

## Discussion

In this study we demonstrate that hs-CRP could be a marker to predict the cognitive impairment in elderly patients in outpatient clinic. Our study also indicates that DHEA-S is lower in patients with depressive mood in the elderly. Thus, measuring these markers in the outpatient clinic might be very useful to assess cognitive and functional impairment as well as depressive mood in elderly patients in addition to the assessment by CGA.

Comprehensive Geriatric Assessment is a very effective way to assess cognitive and functional impairment in the elderly and to find geriatric problems to improve their quality of life (QOL). However, most of hospitals have not utilized this assessment at their outpatient clinics because it is time consuming and unprofitable. Therefore, most geriatricians assess inpatients with CGA, which is getting more and more popular in Japan. Studies with outpatient CGA have not been successful in terms of survival so far. Therefore, by utilizing outpatient CGA and serum markers we would be able to select patients with potential risk for the future decline of cognitive functions and to eventually improve survival of frail elderly patients, although Bradly *et al.* indicated that the improvement of mental health may be an appropriate and realistic goal for outpatient CGA.<sup>29</sup>

Findings from epidemiological studies and some small clinical trials that non-steroidal anti-inflammatory drug (NSAID) users have a lower risk of AD, with indications of dose effects, has drawn much interest in inflammatory mechanisms in AD.<sup>30,31</sup> As our data show that the patients with cognitive impairment or potential decline have higher levels of hs-CRP, we might be able to select those patients to treat with NSAID to prevent the progression of cognitive impairment. To rationalize this treatment, we need a larger scale of study to prove whether or not the decline in cognitive function is faster in patients with higher hs-CRP levels.

Plasma DHEA shows a progressive age-related decline in men and women. DHEA and androstenedione have been shown to inhibit IL-6 secretion from human mononuclear cells *in vitro*,<sup>32</sup> suggesting a connection between aging of endocrine and immune sys-

tems. DHEA has also been shown to suppress IL-4, IFN- $\gamma$  and astrocytic TNF- $\alpha$  and IL-6 production.<sup>33,34</sup> Despite its interesting inverse association with IL-6 levels and beneficial effects on senescence and cognition, a recent Cochrane Systematic Review found only limited evidence of an improved sense of well-being with DHEA supplementation.<sup>35</sup> Clinical benefit of DHEA supplementation should wait for other ongoing trials.

Association between DHEA-S levels and degenerative disorders of the nervous system, such as dementia and cognitive decline, have been controversial.<sup>17,36-38</sup> Some reports did not show the association of low serum DHEA-S levels with AD and other forms of cognitive dysfunction,<sup>39,40</sup> while others suggest a role of DHEA-S in depression, dementia and impaired cognitive performances in the elderly.<sup>41,42</sup> Although our study did not show the association of DHEA or DHEA-S with MMSE, KBD, 'Get up and go' or functional reach, a significant association of low DHEA-S with depressive mood was shown in our patient group. Our study is a cross-sectional study and the number of the patients is relatively small. Therefore, a longitudinal study will be necessary to determine whether or not the patients with low DHEA-S have a higher risk for the development of depression and whether or not treatment of those patients with DHEA-S can prevent the development of depression. Since the levels of DHEA-S declined according to age, the age-related increase of depression might be explained by a decrease of sex hormone, such as DHEA-S. In this study, we used 8/9 as cutoff for GDS. We used this cutoff point because it was appropriate in terms of sensitivity and specificity (Wada *et al.* unpubl. data). When we used 5/6 as cutoff for GDS, we found a lower level of DHEA-S in patients with GDS scores of 6 or over, but could not find a statistical significance.

Our data indicate higher incidence of dementia in patients with low MMSE and higher hs-CRP. We also demonstrated higher incidence of depression in patients with higher GDS scores and lower DHEA-S levels in elderly patients with relatively preserved ADL. Risk factors for stroke such as hypertension did not seem to be involved in these markers. With these cross-sectional data in hand, we think that it is important to follow these patients to determine whether or not high levels of hs-CRP results in a decrease in cognitive and functional impairment and whether or not low levels of DHEA-S predicts future depression. If low levels of DHEA-S are associated with the development of depression in the elderly, supplementation of DHEA might be beneficial to improve their QOL. It is also important to determine the cutoff point of these markers to select patients with high risk for cognitive decline or depression. A larger scale of study is necessary to address this issue.

In summary, our study indicates that measuring serum markers such as hs-CRP and DHEA-S would be useful to assess elderly patients along with CGA.

## Acknowledgments

We thank Akiko Masaki, Atsuko Kokuryu, Marie Kinjo and Emiko Matsuyama for assessment of the patients and Nobuo Shirahashi for advice on statistical analysis. This study was supported by research grants from Osaka Gas Group Welfare Foundation and Foundation for Total Health Promotion.

## References

- Epstein AM, Hall JA, Besdine R *et al.* The emergence of geriatric assessment units. The 'new technology of geriatrics'. *Ann Intern Med* 1987; **106**: 299–303.
- Campion EW. The value of geriatric interventions. *N Engl J Med* 1995; **332**: 1376–1378.
- Rubenstein LZ, Josephson KR, Wieland GD, English PA, Sayre JA, Kane RL. Effectiveness of a geriatric evaluation unit. A randomized clinical trial. *N Engl J Med* 1984; **311**: 1664–1670.
- Applegate WB, Miller ST, Graney MJ, Elam JT, Burns R, Akins DE. A randomized, controlled trial of a geriatric assessment unit in a community rehabilitation hospital. *N Engl J Med* 1990; **322**: 1572–1578.
- Applegate WB, Burns R. Geriatric medicine. *JAMA* 1996; **275**: 1812–1813.
- Boult C, Boult LB, Morishita L, Dowd B, Kane RL, Urdangarin CF. A randomized clinical trial of outpatient geriatric evaluation and management. *J Am Geriatr Soc* 2001; **49**: 351–359.
- Reuben DB, Frank JC, Hirsch SH, McGuigan KA, Maly RC. A randomized clinical trial of outpatient comprehensive geriatric assessment coupled with an intervention to increase adherence to recommendations. *J Am Geriatr Soc* 1999; **47**: 269–276.
- Stuck AE, Siu AL, Wieland GD, Adams J, Rubenstein LZ. Comprehensive geriatric assessment: a meta-analysis of controlled trials. *Lancet* 1993; **342**: 1032–1036.
- Rubin CD, Sizemore MT, Loftis PA, de Mola NL. A randomized, controlled trial of outpatient geriatric evaluation and management in a large public hospital. *J Am Geriatr Soc* 1993; **41**: 1023–1038.
- Burns R, Nichols LO, Graney MJ, Cloar FT. Impact of continued geriatric outpatient management on health outcomes of older veterans. *Arch Intern Med* 1995; **155**: 1313–1318.
- Cohen HJ, Feussner JR, Weinberger M *et al.* A controlled trial of inpatient and outpatient geriatric evaluation and management. *N Engl J Med* 2002; **346**: 905–912.
- Kalaria RN, Harshbarger-Kelly M, Cohen DL, Premkumar DR. Molecular aspects of inflammatory and immune responses in Alzheimer's disease. *Neurobiol Aging* 1996; **17**: 687–693.
- Akiyama H, Barger S, Barnum S *et al.* Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000; **21**: 383–421.
- Morrison MF, Redei E, TenHave T *et al.* Dehydroepiandrosterone sulfate and psychiatric measures in a frail, elderly residential care population. *Biol Psychiatry* 2000; **47**: 144–150.
- Morales AJ, Nolan JJ, Nelson JC, Yen SS. Effects of replacement dose of dehydroepiandrosterone in men and women of advancing age. *J Clin Endocrinol Metab* 1994; **78**: 1360–1367.
- Wolkowitz OM, Reus VI, Roberts E *et al.* Dehydroepiandrosterone (DHEA) treatment of depression. *Biol Psychiatry* 1997; **41**: 311–318.
- Wolf OT, Neumann O, Hellhammer DH *et al.* Effects of a two-week physiological dehydroepiandrosterone substitution on cognitive performance and well-being in healthy elderly women and men. *J Clin Endocrinol Metab* 1997; **82**: 2363–2367.
- Wolf OT, Naumann E, Hellhammer DH, Kirschbaum C. Effects of dehydroepiandrosterone replacement in elderly men on event-related potentials, memory, and well-being. *J Gerontol A Biol Sci Med Sci* 1998; **53**: M385–M390.
- Tanaga K, Bujo H, Inoue M *et al.* Increased circulating malondialdehyde-modified LDL levels in patients with coronary artery diseases and their association with peak sizes of LDL particles. *Arterioscler Thromb Vasc Biol* 2002; **22**: 662–666.
- Ishizaki T, Watanabe S, Suzuki T, Shibata H, Haga H. Predictors for functional decline among nondisabled older Japanese living in a community during a 3-year follow-up. *J Am Geriatr Soc* 2000; **48**: 1424–1429.
- Matsubayashi K, Okumiya K, Wada T, Osaki Y, Doi Y, Ozawa T. Cognitive and functional status of the Japanese oldest old. *J Am Geriatr Soc* 1997; **45**: 385–386.
- Weiner DK, Duncan PW, Chandler J, Studenski SA. Functional reach: a marker of physical frailty. *J Am Geriatr Soc* 1992; **40**: 203–207.
- Yesavage JA. Geriatric Depression Scale. *Psychopharmacol Bull* 1988; **24**: 709–711.
- Podsiadlo D, Richardson S. The timed 'Up & Go': a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; **39**: 142–148.
- Matsubayashi K, Okumiya K, Wada T, Osaki Y, Doi Y, Ozawa T. Secular improvement in self-care independence of old people living in community in Kahoku, Japan. *Lancet* 1996; **347**: 60.
- Okumiya K, Matsubayashi K, Wada T, Kimura S, Doi Y, Ozawa T. Effects of exercise on neurobehavioral function in community-dwelling older people more than 75 years of age. *J Am Geriatr Soc* 1996; **44**: 569–572.
- Wada T, Ishine M, Sakagami T *et al.* Depression in Japanese community-dwelling elderly – prevalence and association with ADL and QOL. *Arch Gerontol Geriatr* 2004; **39**: 15–23.
- Okumiya K, Matsubayashi K, Nakamura T *et al.* The timed 'Up & Go' test and manual button score are useful predictors of functional decline in basic and instrumental ADL in community-dwelling older people. *J Am Geriatr Soc* 1999; **47**: 497–498.
- Bradley EH, Bogardus ST Jr, van Doorn C, Williams CS, Cherlin E, Inouye SK. Goals in geriatric assessment: are we measuring the right outcomes? *Gerontologist* 2000; **40**: 191–196.
- Bertozzi B, Barbisoni P, Franzoni S, Frisoni GB, Rozzini R, Trabucchi M. Association of chronic non-steroidal anti-inflammatory drugs use and cognitive decline in nondemented elderly patients admitted to a geriatric evaluation and rehabilitation unit. *Arch Gerontol Geriatr* 1996; **23**: 71–79.
- McGeer PL, Schulzer M, McGeer EG. Arthritis and anti-inflammatory agents as possible protective factors for Alzheimer's disease: a review of 17 epidemiologic studies. *Neurology* 1996; **47**: 425–432.
- Straub RH, Konecna L, Hrach S *et al.* Serum dehydroepiandrosterone (DHEA) and DHEA sulfate are negatively correlated with serum interleukin-6 (IL-6), and DHEA inhibits IL-6 secretion from mononuclear cells in man in

- vitro: possible link between endocrinosenescence and immunosenescence. *J Clin Endocrinol Metab* 1998; 83: 2012-2017.
- 33 Danenberg HD, Ben-Yehuda A, Zakay-Rones Z, Friedman G. Dehydroepiandrosterone (DHEA) treatment reverses the impaired immune response of old mice to influenza vaccination and protects from influenza infection. *Vaccine* 1995; 13: 1445-1448.
  - 34 Kipper-Galperin M, Galilly R, Danenberg HD, Brenner T. Dehydroepiandrosterone selectively inhibits production of tumor necrosis factor alpha and interleukin-6 [correction of interleukin-6] in astrocytes. *Int J Dev Neurosci* 1999; 17: 765-775.
  - 35 Huppert FA, Van Niekerk JK, Herbert J. Dehydroepiandrosterone (DHEA) supplementation for cognition and well-being. *Cochrane Database Syst Rev* 2000; CD000304.
  - 36 Barrett-Connor E, von Muhlen D, Laughlin GA, Kripke A. Endogenous levels of dehydroepiandrosterone sulfate, but not other sex hormones, are associated with depressed mood in older women: the Rancho Bernardo Study. *J Am Geriatr Soc* 1999; 47: 685-691.
  - 37 Cruess DG, Antoni MH, Kumar M *et al.* Cognitive-behavioral stress management buffers decreases in dehydroepiandrosterone sulfate (DHEA-S) and increases in the cortisol/DHEA-S ratio and reduces mood disturbance and perceived stress among HIV-seropositive men. *Psychoneuroendocrinology* 1999; 24: 537-549.
  - 38 Yaffe K, Ettinger B, Pressman A *et al.* Neuropsychiatric function and dehydroepiandrosterone sulfate in elderly women: a prospective study. *Biol Psychiatry* 1998; 43: 694-700.
  - 39 Herndon JG, Lacreuse A, Ladinsky E, Killiany RJ, Rosene DL, Moss MB. Age-related decline in DHEAS is not related to cognitive impairment in aged monkeys. *Neuroreport* 1999; 10: 3507-3511.
  - 40 Watson RR, Huls A, Araghinikou M, Chung S. Dehydroepiandrosterone and diseases of aging. *Drugs Aging* 1996; 9: 274-291.
  - 41 Barrett-Connor E, Ferrara A. Dehydroepiandrosterone, dehydroepiandrosterone sulfate, obesity, waist-hip ratio, and noninsulin-dependent diabetes in postmenopausal women: the Rancho Bernardo Study. *J Clin Endocrinol Metab* 1996; 81: 59-64.
  - 42 Kalmijn S, Launer LJ, Stolk RP *et al.* A prospective study on cortisol, dehydroepiandrosterone sulfate, and cognitive function in the elderly. *J Clin Endocrinol Metab* 1998; 83: 3487-3492.

ORIGINAL ARTICLE

# Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: VII. Khon Khen in Thailand

Masayuki Ishine,<sup>1</sup> Teiji Sakagami,<sup>2</sup> Ryouta Sakamoto,<sup>3</sup> Taizo Wada,<sup>1</sup> Kovit Khampitak,<sup>4</sup> Mutsuko Fushida,<sup>5</sup> Toshiko Kawakita,<sup>5</sup> Kiyohito Okumiya,<sup>6</sup> Toru Kita<sup>7</sup> and Kozo Matsubayashi<sup>1,8</sup>

<sup>1</sup>Department of Field Medicine, <sup>2</sup>Department of Psychiatry, <sup>3</sup>Department of Public Health, Kyoto University, Kyoto, Japan; <sup>4</sup>Department of Obstetrics and Gynecology, Khon Khen University, Khon Khen, Thailand, <sup>5</sup>Kyoto Preventive Medical Center Foundation, <sup>6</sup>Research Institute for Humanity and Nature, <sup>7</sup>Department of Cardiovascular Medicine, and <sup>8</sup>The Center for South-east Asian Studies, Kyoto University, Kyoto, Japan

**Objective:** To compare the findings of comprehensive geriatric assessment between community-dwelling elderly aged 60 years and older living in Thailand and Japan.

**Design:** A cross-sectional, interview- and examination-based study.

**Setting:** The community-dwelling elderly living in rural Thang Kwang villages, in the city Khon Kaen, Khon Kaen province, Thailand and in the town Sonobe, Kyoto, Japan.

**Methods:** The subjects consisted of 198 and 209 community-dwelling elderly aged 60 years or older in two developmentally different areas in Khon Kaen province in Thailand and 411 community-dwelling elderly aged 65 years or older in Japan, respectively, which were examined using a common comprehensive geriatric assessment tool. Interviews pertaining to activities of daily living (ADL), medical and social history, quality of life (QOL) and the 15-item Geriatric Depression Scale as well as anthropometrical and blood chemical examinations were included in the assessment. Using ANOVA and the  $\chi^2$  test, the findings of the three groups were compared.

**Results:** In comparison with Comprehensive Geriatric Assessment (CGA) our findings among communities dwelling in the rural Thang Kwang villages, the city Khon Kaen, Thailand and in the highly-developed city Kyoto, Japan, were that ADL (except social role), QOL (except family relationship), mean systolic blood pressure, serum total and high-density cholesterol levels, hemoglobin concentrations and prevalence of hypertension were lower in the elderly in rural Thang Kwang villages than those in Khon Kaen city or in Kyoto. In contrast, a prevalence of anemia defined by World Health Organization criteria was higher in the elderly in rural Thang Kwang villages or Khon Kaen city than in Kyoto. It is noteworthy that a prevalence of suspected impaired glucose tolerance or diabetes mellitus in Thailand was extraordinarily higher than in Japan, compared to nearby South-Asian countries.

Accepted for publication 13 June 2005.

Correspondence: Masayuki Ishine, MD, Department of Field Medicine, Kyoto University Graduate School of Medicine, 56 Kawahara-machi, Shogoin, Sakyo-ku, Kyoto 606-8105, Japan.  
Email: ishine@kuhp.kyoto-u.ac.jp; masayuki-ishine@hotmail.co.jp



**Conclusion:** It is supposed that economical and social development might bring a better CGA and a better nutritional state to Thailand. However, we should pay more attention to over-nutrition, modified lifestyle and appropriate controls for global risk factors in the community-dwelling elderly in this economically developmental country in Asia.

**Keywords:** activities of daily living (ADL), community-dwelling elderly, comprehensive geriatric assessment, Khon Kaen, quality of life (QOL), Thailand.

## Introduction

Asian countries have diverse characteristics in terms of their geographic ecology, population and ethnicity, histories and cultural backgrounds, economical and industrial developments. However, the aging population is now rapidly growing equally in each country in Asia. This seventh paper in a consecutive series of articles dealing with the comparison of comprehensive geriatric assessment between community-dwelling elderly in six Asian countries (i.e. Singapore, Korea, Vietnam, Laos, Indonesia Myanmar)<sup>1-6</sup> and in Japan addresses the elderly in Thailand.

Thailand is one of the most economically developed and noticeable countries in South-east Asia. The gross domestic product (GDP)-per capita is increasing to \$US8100 and the GDP-per growth rate is increasing by 6.1% year by year. Thailand is located in central South-east Asia and its total population is 64 865 523 people (November 2004), living in a total of 76 provinces. The rate of population aged 65 years and older was 7.3% in 2004. In 2004, life expectancy in Thailand was a mean age of 71.4 years, or 69.2 years in males and 73.7 years in females.

Regarding ethnicity, the population is 75% Thai, 15% Chinese, and 10% other ethnic groups.<sup>7</sup>

Since 1990, we have carried out a comprehensive assessment of the geriatric functioning of community-dwelling elderly and have provided efficient education to promote a healthy state of the elderly population living in several towns in Japan.<sup>8-21</sup> In the present study, we have applied the common method of comprehensive geriatric assessment (CGA) to the community-dwelling elderly population living in two areas (urban and rural) in Khon Kaen province in Thailand compared with the CGA findings of community-dwelling elderly in Kyoto in Japan.

## Methods

### Subjects

Our study population consisted of 198 community-dwelling elderly subjects aged 60 years and older (male : female = 92 : 106; mean age: 68.2 years) living in rural Thang Kwang villages, and 209 subjects (male : female = 50 : 159; mean age: 68.7 years) living in

the city Khon Kaen, Khon Kaen province, located 440 km north-east of the capital, Bangkok, Thailand (Fig. 1). In both sets, approximately 60% of the elderly subjects were retired. The comparative control subjects consisted of 411 community-dwelling elderly (male : female = 174 : 237; mean age: 71.7 years) living in the town Sonobe, Kyoto, Japan. The geriatric survey for community-dwelling elderly living in Thailand and in Japan was carried out in March 2005 and in April 2003,




Figure 1 Map of Thang Kwang villages, Khon Kaen city in Khon Kaen province in Thailand.

respectively. The rural Thang Kwang villages in Waeng Noi district had a population of 2098 people, 222 of whom were elderly (60 years old or older). We examined 198 of the elderly (89.1% of eligible subjects). Khon Khan city in Khon Khan province had a population of 127 470 people, 10 382 of whom were elderly (60 years old or older; male : female = 4875 : 5507) and we examined 209 elderly subjects who were randomized volunteers (2.0% of eligible subjects). Sonobe, Kyoto had a population of 16 700, 3340 of whom were elderly (65 years or older). We examined 411 of the elderly (12.3% of eligible subjects). The proportion aged 65 years or older in Sonobe was 20.0%. All elderly subjects living in this town were given a self-rating questionnaire by mail, and were then invited to get their health status checked, including blood chemical examination and blood pressure measurement by consultation with a physician at several community houses.

#### *Items of comprehensive geriatric assessment*

Items of the CGA included activities of daily living (ADL), screening of depression, quantitative assessing of quality of life (QOL) as well as medical and anthropometrical indicators.

#### *Activities of daily living*

For basic ADL assessment, each subject rated his/her independence on seven items (walking, ascending and descending stairs, eating, dressing, going to the toilet, bathing and grooming) as to the help needed, and rated them from 3 to 0 (score 3 = completely independent, score 2 = need some help, score 1 = need much help, score 0 = completely dependent). The items were added together to give scores ranging 0–21, with low scores indicating disability.<sup>10,16,19</sup> The information-related function was defined as scores we summed the scores for four item functions (visual activity, hearing activity, conversation and memory in 1 day) using a rating scale from 0 (cannot at all) to 3 (completely independent) adding together to a score ranging 0–12. For higher level functional capacity, each subject rated his/her independence on the Tokyo Metropolitan Institute of Gerontology (TMIG) index of competence.<sup>22,23</sup> This assessment consists of a 13-item index including three sublevels of competence: (i) instrumental ADL (five items: the ability to use public transport, buy daily necessities, prepare a meal, pay bills, handle banking matters, rated on a yes/no basis); (ii) intellectual ADL (four items: the ability to fill out forms, read newspapers, read books or magazines, and interest in television programs or news articles on health-related matters, rated on a yes/no basis); and (iii) social ADL (four items: the ability to visit friends, give advice to relatives and friends who confide in them, visit someone at the hospital and initiate

conversation with younger people, rated on a yes/no basis).

#### *Depression and QOL*

We screened for depressive symptoms using the English version of the 15-item-Geriatric Depression Scale (GDS-15).<sup>24,25</sup> We defined depression as a GDS-15 score of 6 or more, with a score of 6–9 indicating 'mild depression', and a score of 10 or more indicating 'severe depression'. QOL were assessed using a 100 mm visual analog scale (worst QOL on the left end of the scale, best on the right) in the following five items: subjective sense of health, relationship with family, relationship with friends, financial satisfaction and subjective happiness.<sup>26–28</sup>

#### *Social, anthropometrical and medical assessments*

Living conditions, lifestyle (current exercise, drinking alcohol, smoking and so on), and medical histories (histories of stroke, heart diseases and osteoarthropathy, as well as taking antihypertensive drugs) were also assessed. Blood pressure levels were measured twice in a sitting position by auto-sphygmomanometer (HEM 757, Omron, Japan) and then averaged. Blood chemical analysis (total cholesterol, high-density cholesterol [HDL-C], creatinine, blood sugar, hemoglobin, uric acid) was performed by auto-blood chemical analyzer (SP-4410, 4420, Arkrey, Japan) in the Thailand survey, and was done by a laboratory company (SRL, Tokyo, Japan) in the Kyoto survey. Each blood test was taken in a casual setting (someone in fasting and someone in non-fasting condition) in both Thailand and Japan.

#### *Statistical analysis*

Statistical analysis was performed using Stat View version 5 for Macintosh (SAS Institute, Cary, NC, USA). ANOVA was used for continuous variables and  $\chi^2$  test was used for categorical variables. *P*-values less than 0.05 were used to indicate statistical significance. Where *P*-values were less than 0.05 by one-factor ANOVA, Fisher's test was performed to compare the scores between the two groups.

#### **Results**

Table 1 shows the comparison of baseline characteristics among the elderly subjects living in the Thang Kwang villages and Khon Kaen city in Thailand, and in Sonobe, Kyoto in Japan. There was a significant difference in mean age; the elderly subjects in the Thang Kwang villages (68.2 years) and Khon Kaen city (68.7 years) in Thailand, were significantly younger than

**Table 1** Baseline characteristics between the community-dwelling elderly living in Thailand and Japan

	Thang Kwang villages in Thailand ( <i>n</i> = 198)	Khon Kaen city in Thailand ( <i>n</i> = 209)	Sonobe, Kyoto in Japan ( <i>n</i> = 411)	<i>P</i>
Age	68.2 ± 7.3*	68.7 ± 5.9*	71.7 ± 4.8	<0.0001
Male/Female	92/106	50/159	174/237	<0.0001
Marital status				<0.0001
Widowed (%)	30.3	37.8	18.8	
Unmarried (%)	3.0	2.4	0.2	
Divorced (%)	1.0	4.3	0.7	
Lifestyle				
Living (%)				<0.0001
With children or parents	73.7	67.5	53.4	
With spouse only	11.6	12.9	38.2	
Alone	0.5	8.1	7.6	
Work or do gardening every day (%)	57.6	46.4	81.4	<0.0001
Walk and exercise every day (%)	84.3	87.5	45.5	<0.0001
Habits				
Drinking alcohol every day (%)	3.0	2.4	22	<0.0001
Current smoking (%)	14.6	6.7	13.8	<0.0001
Medical				
Taking anti-hypertensive medication (%)	6.6	31.1	31.5	<0.0001
History of stroke (%)	1	3.3	0.5	0.0045
History of heart disease (%)	3	9.2	5.8	<0.0001
History of osteoarthopathy (%)	51.0	34.9	59.4	<0.0001
History of fall (%)	44.9	34.1	12.8	<0.0001

SD, standard deviation. *P* value based on ANOVA for continuous variables and  $\chi^2$  test for categorical variables. \**P* < 0.05 vs Sonobe in Fisher's PLSD.

those in Kyoto (71.7 years old). There were also differences in the male : female ratio among the three populations. The percentage of widows in Khon Kaen city (37.8%) was higher than in the Thang Kwang villages (30.3%) or in Kyoto (18.8%). The percentage of the elderly living with children or parents in the Thang Kwang villages (73.7%) was higher than in Khon Kaen city (67.5%) or in Kyoto (53.4%). The percentage of the elderly living alone in the Thang kwang villages (0.5%) was much lower than in Khon Kaen city (8.1%) or in Kyoto (7.6%). The percentage of the elderly working or gardening everyday in Kyoto (81.4%) was higher than in the Thang Kwang villages (57.6%) or in Khon Kaen city (46.4%). The percentage of the elderly walking or exercising everyday in Kyoto (81.4%) was lower than in the Thang Kwang villages (84.3%) or in Khon Kaen city (87.5%). The percentage of the elderly drinking alcohol everyday was higher in Kyoto (22%) than in the Thang Kwang villages (3.0%) or in Khon Kaen city (2.4%). The percentage of the elderly currently smoking in Khon Kaen city (6.7%) was lower than in the Thang Kwang villages (14.6%) or in Kyoto (13.8%). In a medical situation, the rate of elderly subjects taking antihypertensive medication in Kyoto (31.5%) and in Khon Kaen city (31.1%) was higher than in the Thang Kwang villages

(6.6%). The rate of subjects recognizing their apparent history of stroke in Khon Kaen city (3.3%) was higher than in the Thang Kwang villages (1.0%) or in Kyoto (0.5%). The rate of subjects recognizing their apparent history of heart disease in Khon Kaen city (9.2%) was higher than in Kyoto (5.8%) or in the Thang Kwang villages (3.0%). Past history of osteoarthopathy in the elderly subjects in the Thang Kwang villages (34.9%) was lower than in Khon Kaen city (51.0%) or in Kyoto (59.4%). Past history of falls in the elderly subjects in the Thang Kwang villages (44.9%) was higher than in Khon Kaen city (34.1%) or in Kyoto (12.9%).

Table 2 shows the comparison of scores in ADL, GDS-15 score and quantitative QOL among community-dwelling elderly living in the two areas in Thailand and in Japan. There were significant differences in scores of basic ADL, information-related function, instrumental ADL, intellectual ADL, social role and TMIG Index. The mean scores of all ADL except social role were significantly highest in Japanese elderly followed by subjects in Khon Kaen city, and those in the Thang Kwang villages. However, the mean score of social role was not significantly different among the three. The mean score of GDS-15 in elderly subjects in Kyoto were significantly lower than in the Thang Kwang

**Table 2** Comparison of activities of daily living (ADL) and quality of life (QOL) among the community-dwelling elderly living in two areas in Thailand and in Japan

	Thang kwang villages in Thailand ( <i>n</i> = 198)	Khon Kaen City in Thailand ( <i>n</i> = 209)	Sonobe, Kyoto in Japan ( <i>n</i> = 411)	<i>P</i>
Age	68.2 ± 7.3*	68.7 ± 5.9*	71.7 ± 4.8	<0.0001
ADL				
Scores of basic ADL (0–21)	20.5 ± 1.1*	20.5 ± 0.7*	20.8 ± 0.7	<0.0001
% of independence of basic ADL	63.6	52.1	89.2	<0.0001
Information-related function (0–12)	10.9 ± 1.2*	10.9 ± 1.3*	11.7 ± 0.8	<0.0001
% of independence of information-related function	37.9	59.5	81.1	<0.0001
Scores of instrumental ADL (0–5)	3.9 ± 1.4*†	4.3 ± 1.0*	4.9 ± 0.5	<0.0001
% of independence of instrumental ADL	47	53.6	92.6	<0.0001
Scores of intellectual ADL (0–4)	2.2 ± 1.4*†	2.9 ± 1.2*	3.8 ± 0.6	<0.0001
% of independence of intellectual ADL	23.2	43.5	80.2	<0.0001
Scores of social role (0–4)	3.5 ± 0.8	3.6 ± 0.8	3.5 ± 1.0	NS
% of independence of social ADL	68.7	71.8	69.2	<0.0001
Scores of TMIG (0–13)	9.7 ± 2.8*†	10.7 ± 2.3*	12.1 ± 1.6	<0.0001
% of independence of TMIG	21.5	29.7	58.3	<0.0001
Depression				
Scores of Geriatric Depression Scale (0–15)	4.1 ± 3.0*	4.2 ± 3.2*	3.4 ± 3.2	<0.0001
% of GDS ≥ 6	27.3	27.3	22.7	NS
% of GDS ≥ 10	7	9.6	5.1	NS
QOL (0–100)				
Subjective health	54.9 ± 24.1*†	61.0 ± 20.9*	67.2 ± 17.8	<0.0001
Family relationship	84.1 ± 15.9*†	75.6 ± 21.9*	81.5 ± 16.9	<0.0001
Friend relationship	77.2 ± 19.9	72.9 ± 19.7*	79.8 ± 16.7	<0.0001
Financial satisfaction	49.9 ± 22.9*†	51.6 ± 22.3	64.0 ± 21.5	<0.0001
Subjective happiness	68.2 ± 25.8*†	61.0 ± 24.7*	73.2 ± 17.6	<0.0001

SD, standard deviation. *P* value based on ANOVA for continuous variables and  $\chi^2$  test for categorical variables. \**P* < 0.05 vs Sonobe in Fisher's PLSD. †*P* < 0.05 between Khon Kaen city and Thang kwang villages in Fisher's PLSD.

villages or in Khon Kaen city. But the prevalence of mild depression (GDS ≥ 6) and severe depression (GDS ≥ 10) in elderly subjects in Khon Kaen city were higher than in the Thang Kwang villages or in Kyoto. All scores in QOL items, including subjective sense of health, financial satisfaction and subjective happiness except in family relationships, were higher in Japanese elderly subjects than in Thailand's elderly. The scores in QOL items of family relationship in elderly subjects in the Thang Kwang villages were significantly higher than in Khon Kaen city or Kyoto.

Table 3 shows the comparison of anthropometrical indicators among three elderly groups. Body mass index (BMI) was significantly highest in the elderly subjects in Khon Kaen city followed by the ones in the Thang Kwang villages and ones in Kyoto. Mean systolic blood pressure was significantly highest in elderly subjects in Kyoto, followed by those in Khon Kaen city and the Thang Kwang villages. Diastolic blood pressure was not significantly different in elderly subjects among the three elderly groups. The prevalence of hypertension

defined as systolic pressure ≥ 140 mmHg or diastolic pressure ≥ 90 mmHg based on the measurements of casual blood pressure was also highest in Kyoto (49.9%), followed by the Thang Kwang villages (39.4%) and Khon Kaen city (34.5%). The ratio of those taking antihypertensive drugs in the elderly in Kyoto and Khon Kaen city were higher than in the Thang Kwang villages. Therefore, prevalence of hypertension was highest in Kyoto (59.9%), followed by Khon Kaen city (43.4%) and the Thang Kwang villages (40.5%) according to World Health Organization (WHO) criteria for hypertension (systolic pressure ≥ 140 mmHg or diastolic pressure ≥ 90 mmHg or taking hypertensive drugs).

Table 4 shows the comparison of blood chemical findings among the three elderly groups. Both levels of serum total cholesterol and high-density cholesterol (HDL-C) were higher in Japanese subjects than Thailand's subjects. Atherogenic index, blood sugar levels, uric acid and glutamic-pyruvic transaminase were higher in Thailand's elderly than in Japanese elderly. Glucose intolerance was defined as casual blood glucose

**Table 3** Comparison of anthropometrical indicators and blood pressure measurement among the community-dwelling elderly in Thailand and Japan

	Thang Kwang villages in Thailand ( <i>n</i> = 198)	Khon Kaen City in Thailand ( <i>n</i> = 209)	Sonobe, Kyoto in Japan ( <i>n</i> = 411)	<i>P</i>
Anthropometrical				
Height (mean ±SD) (cm)	155.3 ±8.0	154.5 ±7.2	154.2 ±8.6	NS
Weight (mean ±SD)(kg)	54.9 ±11.8 <sup>†</sup>	58.7 ±11.7*	53.9 ±8.7	<0.0001
Body mass index (mean ±SD)	22.7 ±4.2 <sup>†</sup>	24.5 ±4.3*	22.6 ±2.7	<0.0001
Blood pressure				
Systolic blood pressure (mean ±SD) (mmHg)	134.8 ±22.9*	131.9 ±20.3*	140.6 ±20.3	<0.0001
Diastolic blood pressure (mean ±SD) (mmHg)	78.0 ±13.0	77.4 ±12.8	78.3 ±10.6	NS
Prevalence of hypertension (%) (SBP = 140 or DBP ≥ 90)	39.4	34.5	49.9	0.0005
Prevalence of hypertension (%) (WHO criteria)	41.9	49.3	61.3	<0.0001

SD, standard deviation. *P* value based on ANOVA for continuous variables and  $\chi^2$  test for categorical variables. \**P* < 0.05 vs Sonobe in Fisher's PLSD. <sup>†</sup>*P* < 0.05 between Khon Kaen city and Thang kwang villages in Fisher's PLSD.

**Table 4** Comparison of blood chemical findings among the community-dwelling elderly in Thailand and Japan

	Thang Kwang villages in Thailand ( <i>n</i> = 198)	Khon Kaen City in Thailand ( <i>n</i> = 209)	Sonobe, Kyoto in Japan ( <i>n</i> = 411)	<i>P</i>
Total cholesterol (mean ±SD) (mg/dl)	180.0 ±41.6* <sup>†</sup>	189.1 ±37.7*	209.5 ±35.0	<0.0001
HDL-cholesterol (mean ±SD)(mg/dl)	33.4 ±9.3* <sup>†</sup>	27.2 ±10.3*	65.8 ±17.5	<0.0001
Atherogenic index (mean ±SD)	4.8 ±2.1* <sup>†</sup>	7.0 ±3.6*	2.4 ±0.9	<0.0001
Creatinine (mean ±SD)(mg/dl)	1.2 ±1.1* <sup>†</sup>	1.0 ±0.7	0.9 ±0.2	<0.0001
Uric acid (mean ±SD)(mg/dl)	6.2 ±2.0*	6.5 ±5.5*	5.0 ±1.3	<0.0001
GOT (mean ±SD)(IU/L)	28.8 ±14.1*	26.6 ±14.4	25.5 ±16.2	<0.0001
GPT (mean ±SD) (IU/L)	27.9 ±17.6*	25.5 ±16.0*	20.8 ±22.3	0.041
Casual blood sugar (mean ±SD) (mg/dl)	142.4 ±73.7*	151.7 ±85.8*	99.3 ±21.1	<0.0001
% of glucose intolerance (BS ≥ 140mg/dl)	34.3	38.3	4.4	<0.0001
Hemoglobin (mean ±SD) (g/dl)	11.6 ±1.3* <sup>†</sup>	12.1 ±1.3*	13.6 ±1.4	<0.0001
% of anemia (men: Hb < 13g/dl, women: Hb < 12g/dl)	77	59.4	14.2	<0.0001

SD, standard deviation. *P* value based on ANOVA for continuous variables and  $\chi^2$  test for categorical variables. \**P* < 0.05 vs Sonobe in Fisher's PLSD. <sup>†</sup>*P* < 0.05 between Khon Kaen city and Thang kwang villages in Fisher's PLSD.

levels ≥140 mg/dL. Glucose intolerance in this study was much higher in Thailand elderly than in Japanese elderly. Hemoglobin concentration was highest in Japanese elderly, followed by subjects in Khon Kaen city and those in the Thang Kwang villages. According to the WHO criteria of anemia (men: Hb < 13 g/dL; women: Hb < 12 g/dL), percentage of anemia in the elderly in the Thang Kwang villages (77.0%) and Khon Kaen city (59.4%) were extraordinarily higher than in Kyoto (14.2%).

## Discussion

In comparison between community-dwelling elderly in Thailand and in Japan, scores in all ADL items except

social role and in quantitative QOL except family relationship were higher in Japanese elderly subjects than in Thailand's elderly subjects. Especially, the ratios of independence in basic, information-related instrumental, intellectual ADL and TMIG Index were lower in Thailand elderly subjects than in Japanese elderly subjects. These higher functions may be more historically influenced by differences in economic status, infrastructure and educational level between the two countries than basic ADL, because instrumental ADL was associated with economic activities, such as public transportation and shopping. Intellectual activities depend on the levels of literacy of old people.

However, it is noteworthy that it is the family relationship scores in quantitative QOL were higher in the

elderly in the Thang Kwang villages than in Khon Kaen city in Thailand or in Japan, partly because over 70% of the elderly in the Thang Kwang villages lived with children or parents. Kanchanakitsaku reported that senior citizens with children in Thailand were affected by levels of satisfaction including their life support.<sup>29</sup>

In the present study, it was shown that the mean score of GDS-15 in elderly subjects in Kyoto was significantly lower than those in Thailand, so the elderly subjects in Thailand might be depressive. However, the prevalence of mild and severe depression was not significantly different between the two countries. Analysis of other contributing factors to this factor in Thai elderly, such as detailed spiritual or mental factors, remain to be determined due to the limitation of this study.

In blood chemical examinations, elderly people in Thailand had higher atherogenic index scores, and lower total cholesterol, HDL-C levels and hemoglobin concentrations than Japanese subjects, probably due to the differences in caloric intake of nutrition, lifestyles, prevalence of infectious diseases, especially the parasites in Khon Kaen district, or other environmental, economic and hereditary factors and rapid progress in changing their life styles. Mean blood pressure measurements and prevalence of hypertension were higher in Japanese elderly subjects than in Thailand's elderly subjects. But the ratio of prevalence of hypertension (WHO criteria) and the ratio of those taking antihypertensive medication were greatly different between the elderly subjects in Khon Kaen city and in the Thang Kwang villages, probably because hypertensive elderly in the rural Thang Kwang villages leave untreated similarly to Indonesia.<sup>4</sup> And there was a significant difference in mean casual blood sugar levels between the two countries. In this study, the ratio of glucose intolerance in Thai elderly was much higher than in Japanese elderly, which might indicate their own poor recognition and loose control of impaired glucose tolerance or diabetes mellitus among community-dwelling elderly in Thailand and the habitual high-carbohydrate diets similar to Lao PDR in comparison to nearby Asian countries.<sup>3-6</sup>

Also of particular note are the comparative CGA findings among the three elderly populations. CGA findings in the elderly in urban Khon Kaen city lay between the CGA findings of the rural Thang Kwang villages and those in highly developed Kyoto in ADL scores except social roles, serum cholesterol levels, hemoglobin concentrations and prevalence of anemia and hypertension defined based on the WHO criteria. It is also important to note that BMI, atherogenic index and glucose intolerance in Khon Kaen city were much higher among the three elderly groups. This might be because the differences in levels of caloric intake of nutrition, better

socioeconomic status and rapid progress in changing their life styles. Otherwise the energy from cereal, per capita protein and fat supply were almost the same as in WHO's 2000 regional report and the Ministry of Health, Labour and Welfare's 2003 report in Japan.<sup>30,31</sup> Thailand's non-elderly population had a higher caloric intake (2330 kcal/per capita energy supply) than non-elderly Japanese adults (2138 kcal/per capita energy supply). Therefore, it is supposed that economic and social development might bring an over-nutritional state to Thailand.

From the comparison of CGA findings of the community-dwelling elderly among the three different areas, it is clear that appropriate controls for global burden risk factors in the community-dwelling elderly population are needed. In a cross-sectional comparison of CGA findings of the elderly between the rural Thang Kwang villages and urban Khon Kaen city in Thailand, we might be able to find the longitudinal process that Japan has traced during the past several decades. This international comparative study may have some limitations including translated interview questionnaires on ADL or QOL. Although there are differences in history, culture, habits and medical development between Thailand and Japan, this preliminary international cross-sectional comparison of CGA for community-dwelling elderly may contribute in clarifying a fragment of the actual health situation of elderly Thailand subjects compared with that of elderly Japanese subjects.

## Conclusion

In comparison of CGA findings among community-dwelling elderly in the rural Thang Kwang villages, urban Khon Kaen city in Thailand and in highly developed Kyoto in Japan, ADL except social role, QOL except family relationship, mean systolic blood pressure, serum total and HDL-C levels, hemoglobin concentrations and prevalence of hypertension were lower in the elderly in the Thang Kwang villages than in Khon Kaen city or Kyoto. In contrast, prevalence of anemia defined by WHO criteria was higher in the elderly in the Thang Kwang villages or urban Khon Kaen city, the same as close south-Asian countries around Thailand have higher levels than in Kyoto. It is noteworthy that a prevalence of suspected impaired glucose tolerance or diabetes mellitus in Thailand was extraordinarily higher than in Japan, compared to nearby South Asian countries.<sup>3-6</sup> It is supposed that economic and social development might bring a better CGA and a better nutritional state to Thailand. However, we should pay more attention to over-nutrition, modified lifestyle and appropriate controls for global risk factors in the community-dwelling elderly in this economically developmental country in Asia.

## Acknowledgments

We would like to thank all the elderly people who participated in this study both in Thailand and in Japan. We also thank the public health-nurses in Sonobe town for preparing examinations; Sayaka Katori, Chizuka Seki in Shiga Medical School; Prof Shigeyuki Abe in Doshisya University; Prof Kosuke Mizuno in South-east Asian Studies; Chamchan Chalermopol, MD, a foreign student of the Graduate School of Asian African Area Studies in Kyoto University, Japan; Thirawathana Kharntong, MD; Tueanjit Khampitak, MD; Nattia Suwannsri, MD; Thitinat Tassakhon; Prasit Suwannalert; Wanida Wung-pai; Nuapha Inghon; Maturrote Patship; Watcharawit Saensupoh; Nawapote Keeratrakha; and Supagan Thongchon in Khon Kaen University; Pranoonsri Srechompbo and Sauwanan Bumrerraj in Thailand who helped us as interpreters and cooperators. This study was partially supported by the Over-Sea Scientific Research Grant no. 14241005, 15406031 from the Ministry of Education, Science, Culture and Sports, Tokyo, Japan.

## References

- 1 Kozo M, Han KH, Kiyohito O *et al.* Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: I. Singapore. *Geriatr Gerontol Int* 2005; 5: 99–106.
- 2 Teiji S, Kiyohito O, Masayuki I *et al.* Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: II. Hongchon in Korea. *Geriatr Gerontol Int* 2005; 5: 107–114.
- 3 Masayuki I, Taizo W, Teiji S *et al.* Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: IV. Phuto in Vietnam. *Geriatr Gerontol Int* 2005; 5: 115–121.
- 4 Wada T, Ishine M, Okumiya K, Kita T, Mizuno Y, Matsubayashi K. Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: III. West Java in Indonesia. *Geriatr Gerontol Int* 2005; 5: 168–175.
- 5 Okumiya K, Ishine M, Wada T, Bounngong B, Matsubayashi K. Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: V. Savannakhet in Laos. *Geriatr Gerontol Int* 2005; 5: 159–167.
- 6 Taizo W, Kiyohito O, Kentaro S *et al.* Comprehensive geriatric assessment for community-dwelling elderly in Asia compared with those in Japan: VI. Maubin in Myanmar. *Geriatr Gerontol Int* 2005; 5: 276–285.
- 7 CIA. *The World Fact book*. Available from URL: <http://www.cia.gov/cia/publications/factbook/fields/2004.html>, 2004.
- 8 Ishine M, Wada T, Kita T, Kannagi T, Fushida M, Matsubayashi K. Depression, age and ADL in community-dwelling elderly. *Geriatr Gerontol Int* 2003; 3: 262–264.
- 9 HO HK, Matsubayashi K, Wada T, Kimura M, Kita T, Saijoh K. A comparative study of residential care home and community-dwelling elderly in Japan. *Geriatr Gerontol Int* 2002; 2: 80–86.
- 10 Matsubayashi K, Okumiya K, Wada T, Osaki Y, Doi Y, Ozawa T. Secular improvement in self-care independence of old people living in community in Kahoku, Japan. *Lancet* 1996; 347: 60.
- 11 Wada T, Matsubayashi K, Okumiya K *et al.* Serum cholesterol levels and cognitive function assessed by P300 latencies in an older population living in the community. *J Am Geriatr Soc* 1997; 45: 122–123.
- 12 Matsubayashi K, Okumiya K, Wada T, Osaki Y, Doi Y, Ozawa T. Cognitive and functional status of the Japanese oldest old. *J Am Geriatr Soc* 1997; 45: 385–386.
- 13 Matsubayashi K, Okumiya K, Wada T, Doi Y, Ozawa T. Home blood-pressure control in Japanese hypertensive population. *Lancet* 1997; 350: 290–291.
- 14 Matsubayashi K, Okumiya K, Nakamura T, Fujisawa M, Osaki Y. Global burden of disease. *Lancet* 1997; 350: 144.
- 15 Matsubayashi K, Okumiya K, Wada T *et al.* Postural dysregulation in systolic blood pressure is associated with worsened scoring on neurobehavioral function tests and leukoaraiosis in the older elderly living in a community. *Stroke* 1997; 28: 2169–2173.
- 16 Matsubayashi K, Okumiya K, Osaki Y, Fujisawa M, Doi Y. Frailty in elderly Japanese. *Lancet* 1999; 353: 1445.
- 17 Okumiya K, Matsubayashi K, Nakamura T *et al.* The timed 'Up & Go' test and manual button score are useful predictors of functional decline in basic and instrumental ADL in community-dwelling older people. *J Am Geriatr Soc* 1999; 47: 497–498.
- 18 Okumiya K, Matsubayashi K, Wada T *et al.* A U-shaped association between home systolic blood pressure and four-year mortality in community-dwelling older men. *J Am Geriatr Soc* 1999; 47: 1415–1421.
- 19 Matsubayashi K, Okumiya K, Wada T *et al.* Improvement in self-care may lower the increasing rate of medical expenses for community-dwelling older people in Japan. *J Am Geriatr Soc* 1998; 46: 1484–1485.
- 20 Wada T, Ishine M, Kita T, Fujisawa M, Matsubayashi K. Depression screening of elderly community-dwelling Japanese. *J Am Geriatr Soc* 2003; 51: 1328–1329.
- 21 Wada T, Ishine M, Sakagami T *et al.* Depression in Japanese community-dwelling elderly—prevalence and association with ADL and QOL. *Arch Gerontol Geriatr* 2004; 39: 15–23.
- 22 Koyano W, Shibata H, Nakazato K, Haga H, Suyama Y. Measurement of competence: reliability and validity of the TMIG-index of competence. *Arch Gerontol Geriatr* 1991; 13: 103–116.
- 23 Ishizaki T, Watanabe S, Suzuki T, Shibata H, Haga H. Predictors for functional decline among nondisabled older Japanese living in a community during a 3-year follow-up. *J Am Geriatr Soc* 2000; 48: 1424–1429.
- 24 Sheikh JI, Yesavage JA. Geriatric Depression Scale (GDS); recent evidence and development of a shorter version. In: Brink TL, ed. *Clinical Gerontology: a Guide to Assessment and Intervention*. New York: Haworth Press, 1986; 165–173.
- 25 Yesavage JA. Geriatric Depression Scale. *Psychopharmacol Bull* 1988; 24: 709–711.
- 26 Morrison DP. The Crichton Visual Analogue Scale for the assessment of behaviour in the elderly. *Acta Psychiatr Scand* 1983; 68: 408–413.
- 27 Matsubayashi K, Okumiya K, Osaki Y, Fujisawa M, Doi Y. Quality of life of old people living in the community. *Lancet* 1997; 350: 1521–1522.
- 28 Matsubayashi K, Wada T, Okumiya K *et al.* Comparative study of quality of life in the elderly between in Kahoku and in Yaku. *Nippon Ronen Igakkai Zasshi* 1994; 31: 790–799.

- 29 Kanchanakitsakul M. Factors affecting satisfaction of Thai senior citizens living with their children. *Warasan Prachakon Lae Sangkhom* 1999; 8: 143-162.
- 30 Nutrition of South-East Asia. World Health Organization, Regional Office for South-East Asia, New Delhi, 2000.
- 31 Annual Report on Health and Welfare. Available from URL: <http://www.mhlw.go.jp/toukei/saikin/index.html>, 2003.
- Available from URL: <http://www.whosea.org/nhd/book.htm>



ORIGINAL ARTICLE

# Comprehensive Geriatric Assessment for outpatients is important for the detection of functional disabilities and depressive symptoms associated with sensory impairment as well as for the screening of cognitive impairment

Eriko Sumi,<sup>1</sup> Hajime Takechi,<sup>1</sup> Taizo Wada,<sup>1</sup> Masayuki Ishine,<sup>1</sup>  
Yoshio Wakatsuki,<sup>1</sup> Toshinori Murayama,<sup>2</sup> Masayuki Yokode,<sup>2</sup> Makoto Tanaka,<sup>3</sup>  
Toru Kita,<sup>4</sup> Kozo Matsubayashi<sup>5</sup> and Hidenori Arai<sup>1</sup>

*Departments of <sup>1</sup>Geriatric Medicine, <sup>2</sup>Clinical Innovative Medicine, <sup>3</sup>Social Service and <sup>4</sup>Cardiovascular Medicine, Kyoto University Graduate School of Medicine, and <sup>5</sup>Center for South-east Asian Studies, Kyoto University, Kyoto, Japan*

**Background:** The Comprehensive Geriatric Assessment (CGA) for inpatients is very useful to improve the outcomes of elderly patients. However, most of the elderly patients are provided their care by general practitioners in primary care settings without comprehensive assessment. Concise and practical assessment is necessary for the detection of geriatric problems and sufficient care in the outpatient clinic.

**Methods:** The CGA was introduced in the outpatient clinic for elderly people in Kyoto University Hospital and 309 patients participated in the study, where cognitive impairment, depressive symptoms, activities of daily living, and self-reported hearing and visual impairment were assessed.

**Results:** In the patients studied, the most prevalent chief complaint was memory loss (19%). Among the patients complaining of memory loss, two-thirds of the patients were diagnosed as cognitively impaired by the Mini-Mental State Examination. Multiple logistic regression analysis showed that hearing and visual impairment was significantly associated with functional disabilities and that hearing impairment was significantly associated with depressive symptoms.

**Conclusions:** Thus, the CGA for outpatients is useful for the detection of functional disabilities and depressive symptoms by asking about their sensory impairment as well as for the detection of cognitive impairment in elderly patients. Therefore, concise and practical assessment should be introduced in the primary care settings to improve the quality of life of elderly people.

---

Accepted for publication 17 October 2005.

Correspondence: Hidenori Arai, MD, PhD, Department of Geriatric Medicine, Kyoto University Graduate School of Medicine, 54 Kawahara-cho, Shogoin, Sakyo-ku 606-8105, Kyoto, Japan. Email: harai@kuhp.kyoto-u.ac.jp

**Keywords:** cognitive function, Comprehensive Geriatric Assessment (CGA), depression, hearing impairment, visual impairment.

## Introduction

During the last 50 years, the survival rates and consequently the demographic profile of the Japanese population have dramatically changed. The average lifespan of Japanese men and women is 78.3 and 85.2 years, respectively, which is the longest in the world. The proportion of people aged over 65 years reached 18% of the total population in 2002, and is expected to reach 25% by 2020.

Because it is known that a number of undetected problems exist in elderly people, a systematic approach to detect geriatric problems, the Comprehensive Geriatric Assessment (CGA) emerged and started in the 1980s.<sup>1-3</sup> Today, evidence from randomized controlled trials or systematic reviews on the CGA is increasingly available, supporting the use of hospital-based programs with extended ambulatory follow up.<sup>4,5</sup> In primary care settings, however, time and staffs are limited, which makes it difficult to adopt the CGA. Therefore, the CGA for outpatients should be concise, meet the demand of the patients, and be a guide for the subsequent intervention according to the results of CGA.

The first aim of this study was to validate the usefulness of the CGA in primary clinical settings by examining functional and sociomedical problems in elderly patients. Through the analysis, we studied which factor can affect activities of daily living (ADL) and depressive moods in elderly patients. We also tried to find the incidence of patients with cognitive impairment or depressive moods in the patients who had visited our clinic for the first time.

Sensory impairment, cognitive impairment, functional disability and depression are common problems affecting aged people. However, few studies have directly compared their associations in outpatients. The second purpose of the study therefore was to examine associations among these factors and to determine whether sensory impairment is associated with cognitive function, functional disability or depression in elderly patients.

## Methods

### *Subjects*

All elderly (basically 65 years or older) patients who came to Kyoto University Hospital for the first time or had not been seen for the past 6 months in this hospital

were asked to attend the health promotion clinic. This was a general clinic and screened many problems in elderly patients, not specialized to memory loss. We started an outpatient CGA in this clinic in May 2001. Three hundred and nine consecutive patients aged 65 and older (mean age  $\pm$  SD: 75.5  $\pm$  6.6) who visited the outpatient clinic from March 2002 through June 2004 were enrolled for this study after the written informed consent was taken. The study protocol was approved by the Ethical Committee of Kyoto University School of Medicine.

### *Measurements*

The CGA was performed on the day of patient visit by trained speech therapists in the room next to the consultation room after history taking and physical examination were performed. Intensive training and close supervision were provided to these speech therapists to increase interviewer reliability. Blood pressure was measured twice in the sitting position and hypertension was defined based on the World Health Organization (WHO) criteria as the mean pressure level over 140 mmHg in systolic or 90 mmHg in diastolic or in those taking antihypertensive drugs. Demographic data including age, gender, marital status, living conditions, working status, past medical history including cerebrovascular diseases, heart diseases, fracture and arthropathy, were determined by interviewing the patients themselves or their family. Uncertain data was recorded blank. Hearing and vision were assessed by the question: "Do you have difficulty hearing or seeing (even while wearing aids)?" Answers were scored as: 3 (no trouble in hearing or seeing in daily living); 2 (need loud voices or large letters, cannot hear low voices or read newspaper); or 1 (hardly or unable to hear or see). The scores 1 and 2 were defined as hearing or visually impaired. Cognitive status was assessed using the Mini-Mental State Examination (MMSE). Patients with an MMSE score of 23 or less were defined as cognitively impaired. In five of the patients we gave up performing MMSE because of sensory impairment or other reasons. We screened depressive symptoms using the Japanese version of the 15-item Geriatric Depression Scale (GDS-15).<sup>6</sup> Higher scores of GDS-15 indicate a greater degree of depressive mood. In this study, we used a cut-off point of 5/6. Therefore, we defined depression as a GDS-15 score of 6 or more.

We used the Barthel Index for assessment of basic activities of daily living (BADL) and used a cutoff point of 100/95 as most of patients were full score. For higher levels of functional capacity, each subject's independence was rated by the Tokyo Metropolitan Institute of Gerontology Index of Competence (TMIG-IC).<sup>7</sup> This assessment consists of a 13-item index including three sublevels of competence: (i) instrumental self-maintenance; (ii) intellectual activities; and (iii) social role. In this study, we defined decline of ADL as a TMIG-IC score of 9 or less, slightly lower score than the age-specific mean of TMIG-IC, 10.7.<sup>7</sup> Mobility was assessed using timed 'up and go'.

### Timed 'up and go'

This test of balance is commonly used to examine functional mobility in elderly subjects.<sup>8</sup> The test requires the subject to stand up, walk 3 m (10 ft), turn, walk back and sit down. The time to complete the test is strongly correlated to functional mobility. Elderly people who can complete the test in less than 17 s are independent in transfer tasks, which are normal activities in daily living.<sup>9</sup> We then defined completion in more than 17 s as gait difficulty.

### Statistical analysis

Commercially available statistical software, STAT View (SAS Institute, Cary, NC, USA) was used. Continuous data were analyzed by the Mann-Whitney's *U*-test. Dichotomous data were analyzed by analysis. Multiple logistic regression analysis was used to determine the relationships between the GDS-15 and TMIG-IC and screened items in the CGA including age and sex. Associations were considered statistically significant at a level of  $P < 0.05$ .

## Results

Table 1 summarizes the patient characteristics in the study population. The mean age of the patients was 75.5 and the percentage of males was 36.6%. Memory loss was the most frequent chief complaint in our outpatient clinic. Among the 309 patients, 59 patients (19.0%) came to the hospital complaining of memory loss and three patients with hallucinations or delusions, which typically occurs later in the course of dementia.<sup>10</sup> Thirty-six of the 59 patients who complained of memory loss were given a diagnosis of probable Alzheimer's disease (AD) according to diagnostic standards developed by the National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's Disease and Related Disorders Association.<sup>11</sup> Five of the 59 patients with a chief complaint of memory loss were diagnosed with depression.

**Table 1** Baseline characteristics of study patients

Parameter	Value
Total subjects ( <i>n</i> )	309
Age, mean (SD)	75.5 (6.6)
Male subjects	113 (36.6%)
Cerebrovascular diseases ( <i>n</i> = 308)	
Yes	21 (6.8%)
No	287 (93.2%)
Heart diseases ( <i>n</i> = 301)	
Yes	46 (15.3%)
No	255 (84.7%)
Fracture or joint disease ( <i>n</i> = 304)	
Yes	108 (35.5%)
No	196 (64.5%)
Anti-hypertensive drug ( <i>n</i> = 302)	
Yes	93 (30.8%)
No	209 (69.2%)
Chief complaints ( <i>n</i> = 309)	
Memory loss	59 (19.1%)
Psychiatric symptoms	3 (1.0%)
Others	247 (80.0%)

The numbers do not always add up to 309 because of unapplicable values on some variables.

Table 2 shows the diagnostic tests and cutoff points used in this assessment and the prevalence of each screened problem. The problem with highest prevalence was hypertension. Approximately 30% of the patients were taking antihypertensive drugs, while 63.8% of total patients were hypertensive at the physical examination at our clinic. Although it might be expected, patients taking antihypertensive drugs tended to be hypertensive. The proportion of depressive patients was 41.9% in this study when we used the cutoff point 5/6 on the GDS-15. The incidence of impaired ADL and gait disturbance was similar in the TMIG-IC using the cutoff point of 9/10 or in timed 'up and go' using the cutoff point of 17 s.

Patients with cognitive impairment were older, complaining of memory loss more frequently than those without cognitive impairment (Table 3). Among 76 patients who had scores of 23 or less in the MMSE, 38 patients had a chief complaint of memory loss, while 38 patients had chief complaints other than memory loss, such as headache, dizziness and so forth.

We then performed multiple logistic regression analysis to determine which factor screened in the CGA can affect the GDS-15 (Table 4) or TMIG-IC (Table 5). Among the factors studied, hearing impairment was significantly associated with high GDS scores. Impaired ADL (lower TMIG index), female gender, and the presence of hypertension were also correlated with GDS scores. However, age, MMSE, Barthel Index, timed 'up

**Table 2** Prevalence of each screened problem

	Problem	Diagnostic test	Cut-off point	No of patients	Prevalence
Cognition	Cognitive impairment	MMSE ( <i>n</i> = 304)	Score = 23	59	19.4%
Mood	Depression	GDS-15 ( <i>n</i> = 309)	Score = 6	130	42.1%
Activity	Decline of BADL	Barthel Index ( <i>n</i> = 307)	Score = 95	29	9.4%
	Decline of ADL	TMIG-IC ( <i>n</i> = 309)	Score = 9	83	26.9%
	Gait disturbance	Timed 'up and go' ( <i>n</i> = 283)	Score = 17	78	27.6%
Sensory	Visual impairment	'Do you have difficulty seeing?' ( <i>n</i> = 309)	Yes	31	10.0%
	Hearing impairment	'Do you have difficulty hearing?' ( <i>n</i> = 309)	Yes	37	12.0%
Physical	Hypertension	Blood pressure (mmHg) ( <i>n</i> = 301)	SBP = 140 or DBP = 90	191	63.5%

ADL, activities of daily living; BADL, basic activities of daily living.

**Table 3** Characteristics of patients with or without cognitive impairment

MMSE score	≤23 ( <i>n</i> = 76)	≥24 ( <i>n</i> = 227)	<i>P</i>
Mean age (SD), years	78.1 (6.4)	74.6 (6.5)	<0.001*
Chief complaints of memory loss			
Yes	38	21	
No	38	206	<0.001†
Regular medical treatment			
Yes	58	191	
No	18	36	0.1229†

\*The Mann-Whitney's *U*-test was used. †The  $\chi^2$  test was used. MMSE, the Mini-Mental State Examination.

**Table 4** Independent significant associations of each screened problem with depression: multiple logistic regression analysis

Variable	Adjusted Odds ratio	95%CI	<i>P</i> -value
Age	1	1.0–1.0	n.s.
Male gender	0.5	0.3–1.0	<i>P</i> < 0.05
MMSE	1	1.0–1.1	n.s.
Barthel Index	1	0.7–1.5	n.s.
Timed 'Up & Go'	1	0.9–1.1	n.s.
TMIG	1.3	1.1–1.5	<i>P</i> < 0.01
Hearing impairment	5	1.5–16.1	<i>P</i> < 0.01
Visual impairment	1.2	0.5–3.1	n.s.
Hypertension	2	2.1–3.6	<i>P</i> < 0.05

95%CI, 95% confidence interval; n.s., not significant.

**Table 5** Independent significant associations of each screened problem with lower TMIG-IC score: multiple logistic regression analysis

Variable	Adjusted odds ratio	95%CI	<i>P</i> -value
Age	1	1.0–1.0	n.s.
Female gender	1.6	0.8–3.0	n.s.
MMSE	1.1	1.0–1.2	<i>P</i> < 0.01
GDS-15	0.9	0.8–1.0	<i>P</i> < 0.05
Barthel Index	1.4	0.9–2.4	n.s.
Timed 'up and go'	0.9	0.8–1.0	n.s.
Hearing impairment	3.2	1.2–8.8	<i>P</i> < 0.05
Visual impairment	3.2	1.0–10.0	<i>P</i> < 0.05
Hypertension	1.4	0.7–2.5	n.s.

GDS-15, 15-item Geriatric Depression Scale.

and go' or visual impairment were not associated with GDS scores. On the other hand, hearing and visual impairment was significantly associated with a lower TMIG index. MMSE and GDS scores were also signif-

icantly associated with a lower TMIG index, while age, gender, Barthel Index, timed 'up and go' or the presence of hypertension was not associated with a lower TMIG index.