

erative chemoradiotherapy, which has recently been suggested as the standard of care treatment in the United States after a curative resection of gastric adenocarcinoma.³⁴ Because only 10% of these patients had the advised D2 lymph node dissection and 54% of the patients in that trial had a D0 lymph node dissection, the question has raised whether the adjuvant treatment given in that trial only compensates for inadequate surgery. Five-year survival rates of the group that received adjuvant chemoradiotherapy resemble those of the Dutch Gastric Cancer Trial, where no adjuvant treatment was given. Although the population of the INT 0116 trial³⁴ had more advanced stages of disease compared with our trial, we believe that this conclusion seems justified. Many comments on this trial support our opinion.³⁵⁻³⁷ The effect of a limited lymph node dissection on survival was also reported by the study group itself.³⁸ It is therefore doubtful if any survival advantage of chemoradiotherapy would have been found if patients would have had adequate surgery.

We conclude that there is no long-term overall survival benefit from an extended lymph node dissection in Western patients with gastric cancer. The associated higher postoperative mortality offsets its long-term effect in survival. For pa-

tients with N2 disease, an extended lymph node dissection may offer cure, but it remains difficult to identify patients who have N2 disease. Morbidity and mortality are greatly influenced by the extent of lymph node dissection, pancreatectomy, splenectomy, and age. Extended lymph node dissections may be of benefit if morbidity and mortality can be reduced.

Acknowledgment

We are indebted to the participating surgeons and pathologists of the Dutch Gastric Cancer Group and to the data center of the Surgery Department at the Leiden University Medical Center for its contribution to this trial.

Appendix

The appendix is included in the full-text version of this article, available on-line at www.jco.org. It is not included in the PDF (via Adobe® Acrobat Reader®) version.

Authors' Disclosures of Potential Conflicts of Interest

The authors indicated no potential conflicts of interest.

REFERENCES

- Macintyre IMC, Akoh JA: Improving survival in gastric cancer: Review of operative mortality in English language publications from 1970. *Br J Surg* 78:771-776, 1991
- Akoh JA, Macintyre IMC: Improving survival in gastric cancer: Review of 5-year survival rates in English language publications from 1970. *Br J Surg* 79:293-299, 1992
- Dent DM, Madden MV, Price SK: Randomized comparison of R1 and R2 gastrectomy for gastric carcinoma. *Br J Surg* 75:110-112, 1988
- Robertson CS, Chung SCS, Woods SDS, et al: A prospective randomized trial comparing R1 subtotal gastrectomy with R3 total gastrectomy for antral cancer. *Ann Surg* 220:176-182, 1994
- Cuschieri A, Weeden S, Fielding J, et al: Patient survival after D1 and D2 resections for gastric cancer: Long-term results of the MRC randomized surgical trial. *Br J Cancer* 79:1522-1530, 1999
- Bonenkamp JJ, Sasako M, Hermans J, et al: Extended lymph-node dissection for gastric cancer. *N Engl J Med* 340:908-914, 1999
- Bunt AMG, Hermans J, Boon MC, et al: Evaluation of the extent of lymphadenectomy in a randomized trial of Western versus Japanese type surgery in gastric cancer. *J Clin Oncol* 12:417-422, 1994
- Kajitani T: Japanese Research Society for the Study of Gastric Cancer. The general rules for gastric cancer study in Surgery and Pathology. *Jpn J Surg* 11:127-145, 1981
- American Joint Committee of Cancer. Manual for staging of cancer (ed 4). Philadelphia, PA, Lippincott Company, 1992
- International Union Against Cancer: TNM Classification of Malignant Tumors (ed 4). Berlin, Springer, 1992
- International Union Against Cancer: TNM Classification of Malignant Tumors (ed 6). New York, NY, Wiley-Liss, 2002
- Sano T, Katai H, Sasako M, et al: One thousand consecutive gastrectomies without operative mortality. *Br J Surg* 89:123, 2002
- Siewert JR, Böttcher K, Stein HJ, et al: Relevant prognostic factors in gastric cancer: Ten year results of the German gastric cancer study. *Ann Surg* 228:449-461, 1998
- Marubini E, Bozzetti F, Miceli R, et al: Lymphadenectomy in gastric cancer: Prognostic role and therapeutic implications. *Eur J Surg Oncol* 28:406-412, 2002
- Sue-Ling HM, Johnston D, Martin IG, et al: Gastric cancer: A curable disease in Britain. *BMJ* 307:591-596, 1993
- Griffith JP, Sue-Ling HM, Martin I, et al: Preservation of the spleen improves survival after radical surgery for gastric cancer. *Gut* 36:684-690, 1995
- Roukos DH, Lorenz M, Encke A: Evidence of survival benefit of extended (D2) lymphadenectomy in Western patients with gastric cancer based on a new concept: A prospective long-term follow-up study. *Surgery* 123:573-578, 1998
- Bozzetti F, Marubini E, Bonfanti G, et al: Subtotal versus total gastrectomy for gastric cancer: Five-year survival rates in a multicenter randomised Italian trial. *Ann Surg* 230:170-178, 1999
- Kodera Y, Yamamura Y, Shimizu Y, et al: Lack of benefit of combined pancreaticosplenectomy in D2 resection for proximal-third gastric carcinoma. *World J Surg* 21:622-628, 1997
- Kitamura K, Nishida S, Ichikawa D, et al: No survival benefit from combined pancreaticosplenectomy and total gastrectomy for gastric cancer. *Br J Surg* 86:119-122, 1999
- Csendes A, Burdiles P, Rojas J, et al: A prospective randomised study comparing D2 total gastrectomy versus D2 total gastrectomy plus splenectomy in 187 patients with gastric carcinoma. *Surgery* 131:401-407, 2002
- Sano T, Yamamoto S, Sasako M: Randomized controlled trial to evaluate splenectomy in total gastrectomy for proximal gastric carcinoma. *Jpn J Clin Oncol* 32:363-364, 2002
- Edwards P, Blackshaw PG, Barry J, et al: Randomised comparison of D1 versus modified D2 gastrectomy for gastric cancer. *Br J Surg* 90:30, 2003 (suppl 1)
- Damhuis RA, Tilanus HW: The influence of age on resection rates and postoperative mortality in 2773 patients with gastric cancer. *Eur J Cancer* 31A:928-931, 1995
- Klein Kranenbarg E, van de Velde CJH: Gastric cancer in the elderly. *Eur J Surg Oncol* 24:384-390, 1998
- Hermanek P, Altendorf-Hofmann A, Mansmann U, et al: Improvements in staging of gastric carcinoma using the new edition of TNM classification. *Eur J Surg Oncol* 24:536-541, 1998
- Katai H, Yoshimura K, Maruyama K, et al: Evaluation of the new international union against cancer TNM staging for gastric cancer. *Cancer* 88:1796-1800, 2000
- Klein Kranenbarg EK, Hermans J, van Krieken JHJM, et al: Evaluation of the fifth edition of the TNM classification for gastric cancer: Improved prognostic value. *Br J Cancer* 84:64-71, 2001
- Songun I, Hermans J, van de Velde CJH, et al: Expression of oncoproteins and eosinophilic and lymphocytic infiltrates can be used as

D2 Dissection Beneficial for Some Patients

prognostic factors in gastric cancer. *Br J Cancer* 74:1783-1788, 1996

30. Songun I, van de Velde CJH, Arends JW, et al: Classification of gastric carcinoma using the Goseki system provides prognostic information additional to TNM staging. *Cancer* 85:2114-2118, 1999

31. Weiss M, Kuipers E, Postma C, et al: Genomic profiling of gastric cancer predicts lymph node status and survival. *Oncogene* 22:1872-1879, 2003

32. Gunderson LL, Sosin H: Adenocarcinoma of the stomach: Areas of failure in a re-operation series (second or symptomatic look) clinicopathologic correlation and implications for adjuvant

therapy. *Int J Radiat Oncol Biol Phys* 8:1-11, 1982

33. Nashimoto A, Nakajima T, Furukawa H, et al: Randomised trial of adjuvant chemotherapy with mitomycin, fluorouracil, and cytosine arabinoside followed by oral fluorouracil in serosa-negative gastric cancer: Japan clinical oncology group 9206-1. *J Clin Oncol* 21:2282-2287, 2003

34. Macdonald JS, Smalley SR, Benedetti J, et al: Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med* 345:725-730, 2001

35. Schwarz RE: Postoperative adjuvant chemoradiation therapy for patients with resected

gastric cancer: Intergroup 116. *J Clin Oncol* 19:1879, 2001

36. Cuschieri A: Does chemoradiotherapy after intended curative surgery increase survival of gastric cancer patients? *Gut* 50:751, 2002

37. Roukos DH: Adjuvant chemoradiotherapy in gastric cancer: Wave goodbye to extensive surgery? *Ann Surg Oncol* 9:220-221, 2002

38. Hundahl SA, Macdonald JS, Benedetti J, et al: Surgical treatment variation in a prospective, randomized trial of chemoradiotherapy in gastric cancer: The effect of undertreatment. *Ann Surg Oncol* 9:278-286, 2002

FROM THE ASCO-JSCO JOINT SYMPOSIUM

Mitsuru Sasako

Role of surgery in multidisciplinary treatment for solid cancers

Received: June 7, 2004

Abstract In the evolution of solid cancer, there are four steps: noninvasive tumor, local invasive cancer without metastasis, local invasive cancer with lymph node metastasis, and eventually systemic disease. For the first three phases, local treatment, including lymph node dissection, may cure the disease. The choice of local treatment depends on the tumor characteristics, but surgery remains important in many of these cancers. Gastric cancer is one of the typical tumors which remain locally invasive, with or without nodal metastasis, but without systemic metastasis for a rather long period. Metastasis to lymph nodes occurs, frequently even in T1 tumors, but seldom to other sites until the late stage. Thus, the target of local control is the regional lymph nodes. The Intergroup study IT-0116 proved the effect of chemoradiotherapy (CRT) for curable gastric cancer, and thus proved the insufficiency of limited surgery (D0/1). The conventional method of local control for gastric cancer is surgery, including regional lymph node dissection (D2). However, the superiority of D2 has not been proven by randomized controlled trials (RCTs). But all RCTs so far have a crucial problem in the quality of treatment given in the D2 arm. D2 is not a dangerous procedure if done by specialists in large-volume hospitals. D0/1 plus CRT is better than D0/1 alone, but it may be worse than D2 alone. The survival benefit of CRT after D2 is an open question. Establishing standard adjuvant chemotherapy after D2 is a more urgent clinical issue, and there is no reason to abandon D2 gastrectomy for curable gastric cancer in Japan.

Key words Role of surgery · Gastric cancer · Chemoradiotherapy · Local control

M. Sasako (✉)
National Cancer Center Hospital, Tokyo, 5-1-1 Tsukiji, Chuo-ku,
Tokyo 104-0045, Japan
Tel. +81-3-3542-2511; Fax +81-3-3543-9321
e-mail: msasako@gan2.ncc.go.jp

Presentation made at the ASCO-JSCO Joint Symposium in Sapporo, Japan, on October 22, 2003.

The role of surgery in multidisciplinary treatment for cancer

We believe that solid cancers evolve as follows: lesions without invasion, then locally invasive cancer, which will soon metastasize to regional lymph nodes and then to other organs as systemic disease. The initial lesion of cancer is sometimes noninvasive, and is therefore called dysplasia, in spite of cellular or structural atypia, in the West. There are many arguments about dysplasia and early noninvasive cancer between the West and Japan, including, recently, lung cancer. Due to the development of helical computed tomography (CT), very early cancers, i.e., possible noninvasive cancers, are now being diagnosed in many countries, including the United States and Japan. For a long time, in Japan, we have diagnosed these lesions (which are called dysplasia in the West) in the stomach or in the colon, as cancer. It is well known that many of these dysplastic lesions will invade in a rather short time, at which time they are locally invasive cancers (at this point, a diagnosis of cancer is made in the West). The lesions then start to show metastasis to the regional lymph nodes, and then finally, become systemic disease, with metastases in many distant organs. For noninvasive cancer or dysplasia, just observation or limited resection, such as endoscopic mucosal resection (EMR), is the best way to manage them. For locally invasive cancer, just a wide excision could be sufficient. However, as it is impossible to discriminate exactly between locally invasive lesions with and without regional lymph node metastasis, these lesions are often treated by a wide excision plus lymph node dissection. Recently, sentinel-node biopsy has been used to discriminate those lesions with or without nodal metastasis and to minimize the level of aggressive surgery for these tumors. If the tumor becomes systemic disease, local control plus systemic treatment is mandatory if we aim to cure the disease. As the weapon for local treatment, surgery is most frequently used, but radiation can also be used, depending on the tumor characteristics. Different cancers have different patterns of tumor development or evolution. For example, small-cell

lung cancer has a very short span of limited disease, and most of the lesions of this cancer are already local regional disease plus systemic metastasis when diagnosed. At the opposite extreme is gastric cancer. In Japan, more than half of newly diagnosed lesions are T1, early gastric cancers. Advanced lesions of gastric cancer still have only local invasion and regional lymph node metastasis, which can often be cured by surgery alone. Squamous cell cancer of the esophagus would be situated between these two extremes.

Focus on gastric cancer

Table 1 shows the pattern and incidence of metastasis from gastric cancer, according to the tumor depth.¹ Lymph nodes, liver, and peritoneum are the three frequently involved sites. Other sites in the body, such as lung, bone, brain or skin, may have metastasis from gastric cancer, but only at the end of the disease development, at the terminal stage in these patients.

Table 1. Biological behavior of gastric cancer: incidence of metastasis and 5-year survival

Depth	n	LN	Liver	Peritoneum	5-Year survival
pT1					
M	1063	3.3	0.0	0.0	93.3
SM	881	17.4	0.1	0.0	88.9
pT2					
MP	436	46.4	1.1	0.5	81.3
SS	325	63.7	3.4	2.2	65.8
pT3					
SE	1232	78.9	6.3	17.8	35.5
pT4					
SI	724	89.8	15.5	41.6	10.1
Overall	4683	47.8	4.5	11.5	60.3

Patients operated on between 1972 and 1991, at the National Cancer Center Hospital (NCCH), including those with exploratory laparotomy: there were 22 non-resected patients, in whom T was unknown

As shown in Table 1, metastasis occurs almost exclusively to lymph nodes until the primary tumor becomes T3. Liver metastasis occurs in just 6% of the patients with T3 tumor, and in 15.5% of those with T4 tumor. Peritoneal metastasis occurs only after the tumor has reached the serosa, becoming a T3 tumor; the incidence remains at less than 20% in T3 tumors. On the other hand, the incidence of lymph node metastasis is rather high, even in the early stage of disease evolution. Even T1 submucosal invasive tumors have nodal metastasis in nearly 20% of cases. If the tumor becomes T2, over 50% of patients have regional lymph node metastasis. If these nodal metastases were to be left behind after surgery, they would metastasize and eventually become systemic disease.

So, if the patients are treated by D2 or more extensive surgery, which is the standard treatment in Japan, local regional recurrence is not common, as shown in Table 2.¹ This means that D2 dissection can provide rather good local control. By far the commonest site of recurrence is the peritoneum, and systemic and hematogenous metastases are rare (just 7% of all treated patients). Therefore, in patients with gastric cancer, local control can lead to a fairly high success rate for cure. Only 28% of patients developed recurrence; thus, over 70% of patients survived without recurrence. If these tumors are treated by very limited surgery, local regional recurrence could be a big problem.

Dr. Gunderson² reported the pattern of failure after limited surgery with curative intent at his institute. Fifty-four percent of recurrences occurred only in the gastric bed, and recurrences reached nearly 90% if all those with local regional failure were included regardless of other type of recurrence. This shows the importance of local control for gastric cancer.

In gastric cancer, the lymph nodes are the most important metastatic site. Table 3 shows the topographical pN stage according to the tumor depth.¹ The deeper the tumor, the more frequently lymph nodes are metastatic and the more frequently distant regional nodes become metastatic. If the tumor becomes T3, three-fourths of patients have nodal metastasis. If the tumor remains as T1 or T2, we do not see distant regional lymph node metastasis very often.

Table 2. Primary site of recurrence after \geq D2

Depth	n	Recurrence	LN + RF	Peritoneum	Hematogenous (%)
pT1					
M	1063	2	0	0	2 (0.2)
SM	881	18	6	3	9 (1.0)
pT2					
MP	436	45	10	9	26 (5.9)
SS	325	74	15	28	31 (9.5)
pT3					
SE	1232	625	146	330	149 (12.1)
pT4					
SI	724	562	173	283	106 (14.6)
Overall	4683	1326 (28.3%)	330 (7.0%)	635 (13.6%)	323 (6.9%)

Patients operated on between 1972 and 1991, at the NCCH, including those with exploratory laparotomy

A large proportion of patients have N2 disease; even in T2 tumor, over 20% of patients have N2 disease, and in the T3 tumors, over 40% of patients have N2 disease. This means that main target of local control in gastric cancer is lymph node metastasis. There are several grounds for saying that good local control is essential to cure this cancer. First, Professor Siewert reported that R0 resection is by far the most important prognostic factor after curative operation.³ Second, the results of the Intergroup study (IT-0116) showed that adding irradiation to adjuvant chemotherapy could improve the results of limited surgery alone, which could not be achieved by adjuvant chemotherapy alone.⁴ Good local control by radiation, together with chemotherapy, could improve the results of treatment remarkably. The researchers of the Intergroup study also carefully analyzed the prognostic factors in the patients treated in that trial, and found that surgical under-treatment was an independent prognostic factor. This theory can be applied to some other solid cancers as well.

The preferred method of local control depends on the efficacy of treatment other than surgery. If we see a non-Hodgkin's lymphoma in the stomach, we do not operate on

the patients now, and chemotherapy alone can often control both the primary site and the metastasis. Of course, chemoradiotherapy does work, too. Regarding squamous cell carcinoma of the esophagus, chemoradiotherapy can often control the primary tumor and the nodal metastasis, although the local recurrence rate is as high as 20%–30% after chemoradiotherapy. For gastric cancer, even chemoradiotherapy can seldom control an advanced primary tumor, but it may well control nodal disease. Based on the results of the IT-0116 study, if gastric cancer is treated by limited surgery plus chemoradiation (CRT), the primary lesion is controlled by the surgery, and micrometastases in lymph nodes are controlled by the chemoradiation. If gastric cancer is treated by D2 surgery, both the primary and these metastases are controlled by surgery.

Table 4 shows a comparison of two studies, the IT-0116 study, and the Japan Clinical Oncology Group (JCOG) 9501 study.⁵ The JCOG 9501 study is a trial organized by the Gastric Surgery Division of JCOG to evaluate the role of paraaortic lymph node dissection, which is quite extensive surgery. There are remarkable differences between these two trials: in the IT-0116, surgery was rather limited (D0; very limited resection) in 54% of patients, and D1 surgery was done in 36%, while so-called Japanese-type surgery was done in only 10%. But in the JCOG 9501 study, half of the patients underwent D2 dissection, the standard surgery in Japan. The other half underwent much more extensive surgery (D3 dissection). Regarding adjuvant treatment, those allocated to the test arm in the IT-0116 study underwent 45-Gray radiotherapy together with chemotherapy (5-fluorouracil [5-FU] and leucovorin). In the JCOG 9501 trial, none of the patients underwent adjuvant treatment until they developed recurrence. There was no difference in tumor locations between these two trials, although researchers in the United States always say that they have more proximal tumors than antral tumors. Unlike the pattern of tumor location in the general population, a much larger proportion of patients in this American trial had antral tumors, while more tumors of the body were seen in the Japanese trial. Tumor depth is shown in Table 4: 14 T1, 74 T2, 175 T3, and 18 T4 in the IT-0116 study; and 23 T1, 257

Table 3. Lymph node metastasis according to the depth of tumor invasion

Depth	No.	pN+ (%)	pN0	pN1	pN2 (%)	pN3	pN4
T1							
M	619	14 (2)	605	9	5 (0.8)	0	0
SM	499	89 (18)	410	60	29 (5.8)	0	0
T2							
MP	276	126 (46)	150	74	47 (17)	5	0
SS	207	130 (63)	77	65	57 (28)	3	5
T3							
SE	646	484 (75)	162	171	266 (41)	28	19
T4							
SI	152	121 (80)	31	31	65 (43)	12	13
Total	2399	964 (40)	1435	410	469 (20)	48	37


In gastric cancer, the main target of local control is lymph node metastasis

Table 4. Comparison of the results of IT-0116 and JCOG 9501

	IT-0116	JCOG 9501
Surgery	D0/D1/D2-54% : 36% : 10%	D2/D3-50% : 50%
Adjuvant	Radiation 45 Gy Chemotherapy 5-FU + LV	None
No of patients	281 (Test arm)	523
Tumor location	Antrum, 53%; corpus, 24%; cardia, 21%; multiple, 2%	Lower third, 41%; middle third 39%; upper third, 19%
pT stage (1:2:3:4)	14:74:175:18	23:257:230:13
Treatment-related deaths	3 (1.1%) + Postop.	4 (0.8%)
Survival	3-Year: 50% 5-Year: 42%	5-Year: 71.4 (66.5%–76.3%)

Table 5. Estimated 5-year survival of the IT-0116 patients if they would have undergone D2-3 surgery

IT-0116 patients	NCCCH ^a 5-Year survival	Calculated survival proportion	CIH ^b 5-Year survival	Calculated survival proportion
T1, 14	92.2	12.9	96.6	13.5
T2, 74	77.5	57.4	80.6	59.6
T3, 175	47.1	82.4	40.2	70.4
T4, 18	29.9	5.4	17.4	3.1
42%		56.3%		52.2%

^aResults of National Cancer Center Hospital⁶^bResults of Cancer Institute Hospital⁷**Fig. 1.** D1 vs D2 for males and females. High postoperative mortality did not confound comparison in female patients

T2, 230 T3, and 13 T4 in the JCOG 9501. As to the treatment-related death rate (TRD), 1.1% was reported in IT-0116, and 0.8% in JCOG 9501. However, if the total population that could be candidates in this trial is considered, the TRD should be higher in IT-0116, because some postoperative deaths that occurred before enrolment in this trial were not counted. The survival results of IT-0116 are 50% at 3 years and 42% at 5 years, while the overall survival rate at 5 years is 71.4% in the JCOG 9501 study, although the observation time is not sufficient. As there is a non-negligible difference of T-stage distribution between the two trials, this survival comparison is not fair. It is possible, however, to calculate the survival proportion by applying the survival rates of Japanese institutes by pT stage. The hypothetically estimated survival rates are then over 52%, which is about 10% better than the actual survival rate of the patients in the IT-0116 study (Table 5).

The results of the IT-0116 trial are interpreted as follows: (1) D0/1 surgery is proven to be inadequate treatment in terms of local control, (2) the results achieved are worse than the standard level of those treated by D2 surgery, (3) surgical under-treatment clearly undermined survival, (4) whether D0/1 + CRT can be as good as D2 alone should be tested by a RCT, (5) whether CRT after D2 can improve

the results of this type of surgery alone is another question. At the same time, another question arose. Why was D2 not better than D1 in the western RCTs?

In fact, the Dutch and Medical Research Council (MRC) trials did not prove the effect of D2 dissection.^{8,9} However, the quality of D2 dissection in these trials was questionable, with quite high postoperative mortality with extremely small hospital volume. The TRD rate of D2 was as high as 10% and the quality of postoperative care to avoid operative deaths was very poor, due to the small hospital volume. Not only in these trials but also in several other RCTs in surgery, a high TRD rate offsets the long-term effect of treatment. In the two trials on squamous cell carcinoma of the esophagus reported at the 39th annual meeting of the American Society of Clinical Oncology (ASCO), i.e., the German¹⁰ and French¹¹ trials, a benefit of surgery after CRT was not seen in long-term survival, with a remarkable difference of the TRD rates between CRT alone versus CRT plus surgery. Based on the experience in these RCTs, we may say that proper D2 dissection is technically demanding surgery, requiring experience and specific postoperative care, and it should be carried out at specialist centers in the west.

In the Dutch trial, D2 started with a handicap of about 6%, within 3 months, but caught up with the curve of D1,

Table 6. Morbidity and mortality after D2 dissection for gastric cancer

Trial	Type	Number of patients	Number of D2 dissections per hospital/year	Mortality	Morbidity	Reference
Hong Kong ¹²	RCT	30	7.5	3%	57%	Ann Surg
MRC ⁷	RCT	200	1.5	13%	46%	Lancet
Dutch ⁶	RCT	331	1.0	10%	43%	Lancet
Italian ¹³	Phase II	191	8.0	3%	21%	JCO
Sue-Ling ¹⁴	Retrospective	142	14.2	5%	17%	BMJ
Pacelli ¹⁵	Retrospective	157	15.7	4%	22%	Br J Surg

Table 7. Mortality after major postoperative complications

Complications	Dutch trial (n = 711)		NCCH (1980s) (n = 1197)		P Value
Leakage	19/46	41.3%	12/84	14.3%	0.0005
Distal	9/22	40.1%	2/23	8.7%	0.012
Total	10/24	41.7%	10/60	16.7%	0.0047
Abscess/pancreatic fistula	19/91	20.9%	2/75	2.7%	0.0004

Experience is needed to manage major adverse effects to avoid treatment-related deaths TRD, which occur slightly more often in surgery than in chemotherapy. Hospital volume is a concern

although the difference never reached statistical significance. The hospital mortality for D2 and D1 showed a large difference, at nearly 10% for D2, and 4% for D1. But this difference was seen only in male patients, in whom hospital mortality was 4.2% for D1 versus 14% for D2. There was no difference in mortality between D1 and D2 in female patients. Accordingly, the hazard ratio between D1 and D2 by time for each sex is completely different. In female patients, the hazard ratio is almost constant. The survival curves by procedure by sex are shown in Fig. 1. As we would expect, the survival curves of the female patients do not cross, as typical model curves of survival showing a constant hazard, and the *P* value is 0.04. We can confirm that high immediate mortality easily offsets the long-term effect of any cancer treatment.

Table 6 shows the relation between the hospital volume and the TRD rates in many trials or consecutive series of D2 dissection for gastric cancer. The Dutch and MRC trials show extremely low numbers of patients treated per year, per hospital, and show extremely high hospital mortality, compared with other reports.

Table 7 shows the mortality after major complications, comparing the results of the Dutch trial and those of the National Cancer Center Hospital (NCCH) in the 1980s.¹ Even in a high-volume hospital, major complications, such as anastomotic leakage or intraabdominal abscess, were not rare. However, in the Dutch trial, over 40% of patients died when they developed anastomotic leak, while only 14% of such patients died in the NCCH. As to mortality after abdominal abscess, a difference of nearly ten times was observed. Experience is needed to manage major adverse effects to avoid TRD, which occurs slightly more often in surgery than in chemotherapy or CRT. In this regard, hospital volume is a concern.

The Japanese perspective of the role of D2 dissection in multidisciplinary treatment for advanced gastric carcinoma

can be summarized as follows. The superiority of D2 has not been proven by RCTs. But all RCTs so far have a crucial problem in regard to the quality of treatment given in the D2 arm. D2 is not a dangerous procedure if it is done by specialists in large-volume hospitals. D0/1 plus CRT is better than D0/1 alone, but it may be worse than D2 alone. The survival benefit of CRT after D2 is an open question. Establishing standard adjuvant chemotherapy after D2 is a more urgent clinical issue. There is no reason to abandon D2 gastrectomy for curable gastric cancer in Japan.

References

- Sasako M (2003) Principles of surgical treatment for curable gastric cancer. *J Clin Oncol* 23s(December 1 supplement):274–275
- Gunderson LL, Sosin H (1982) Adenocarcinoma of the stomach: Area of failure in a re-operation series (second or symptomatic look) clinicopathologic correlation and implication for adjuvant therapy. *Int J Radiat Oncol Biol Phys* 8:1–11
- Siewert JR, Boettcher K, Roder JD, et al. (1993) Prognostic relevance of systematic lymph node dissection in gastric carcinoma. *Br J Surg* 80:1015–1018
- Macdonald JS, Smalley SR, Benedetti J, et al. (2001) Chemoradiotherapy after surgery compared with surgery alone for adenocarcinoma of the stomach or gastroesophageal junction. *N Engl J Med* 345:725–730
- Sano T, Sasako M, Yamamoto S, et al. (2004) Gastric cancer surgery: morbidity and mortality results from a prospective randomized controlled trial comparing D2 and extended para-aortic lymphadenectomy – Japan Clinical Oncology Group Study 9501. *J Clin Oncol* 22:2767–2773
- Sasako M, Sano T, Katai H, et al. (1997) Radical surgery in gastric cancer. In: Sugimura T, Sasako M (eds) *Gastric Cancer*. Oxford University Press, pp 223–248
- Nakajima T, Yamaguchi T (1999) Appendix: Tubular analysis of Cancer Institute Hospital Data Base. In: *Multimodality therapy for gastric cancer*, pp 256
- Bonenkamp JJ, Hermans J, Sasako M, et al. (1999) Extended lymph-node dissection for gastric cancer. *N Engl J Med* 340:908–914

9. Cuschieri A, Weeden S, Fielding J, et al. (1999) Patient survival after D1 and D2 resection for gastric cancer: long-term results of the MRC randomized surgical trial. *Br J Cancer* 79:1522-1530
10. Stahl M, Wilke M, Walz MK, et al. (2003) Randomized phase III trial in locally advanced squamous cell carcinoma (SCC) of the esophagus: chemoradiation with and without surgery. *J Clin Oncol* 22 (Proceedings of ASCO meeting 2003): 250
11. Bonnetain F, Bedenne L, Michel P, et al. (2003) Definitive results of a comparative longitudinal quality of life study using the Spitzer index in the randomized multicentric phase III trial FFCD 9102 (surgery vs radiochemotherapy in patients with locally advanced esophageal cancer). *J Clin Oncol* 22 (Proceedings of ASCO meeting 2003): 250
12. Robertson CS, Chung SCS, Woods SDS, et al. (1994) A prospective randomized trial comparing R1 subtotal gastrectomy with R3 total gastrectomy for antral cancer. *Ann Surg* 220:175-182
13. Degiuli M, Sasako M, Ponti A, et al. (1998) Morbidity and mortality after D2 gastrectomy for gastric cancer: results of the Italian Gastric Cancer Study Group prospective multicenter surgical study. *J Clin Oncol* 16:1490-1493
14. Sue-Ling H, Johnston D, Martin IG, et al. (1993) Gastric cancer: a curable disease in Britain. *BMJ* 307:591-596
15. Pacelli F, Doglietto GB, Bellantone R, et al. (1993) Extensive versus limited lymph node dissection for gastric cancer: a comparative study of 320 patients. *Br J Surg* 80:1153-1156

Gastric Cancer Surgery: Morbidity and Mortality Results From a Prospective Randomized Controlled Trial Comparing D2 and Extended Para-Aortic Lymphadenectomy—Japan Clinical Oncology Group Study 9501

Takeshi Sano, Mitsuru Sasako, Seiichiro Yamamoto, Atsushi Nashimoto, Akira Kurita, Masahiro Hiratsuka, Toshimasa Tsujinaka, Taira Kinoshita, Kuniyoshi Arai, Yoshitaka Yamamura, and Kunio Okajima

From the Gastric Surgery Division, National Cancer Center Hospital; Cancer Information and Epidemiology Division, National Cancer Center Research Institute; Department of Surgery, Tokyo Metropolitan Komagome Hospital, Tokyo; Department of Surgery, Niigata Cancer Center Hospital, Niigata; Department of Surgery, National Shikoku Cancer Center, Matsuyama; Department of Surgery, Osaka Medical Center for Cancer and Cardiovascular Diseases; Department of Surgery, Osaka National Hospital; Department of Surgery, Osaka Medical College, Osaka; Department of Surgery, National Cancer Center Hospital East, Kashiwa; Department of Surgery, Aichi Cancer Center, Nagoya, Japan; the Gastric Cancer Surgical Study Group of Japan Clinical Oncology Group.

Submitted October 24, 2003; accepted January 16, 2004.

This study was supported by the Grant-in-Aid for Cancer Research from the Ministry of Health and Welfare, and the Second Term Comprehensive 10-Year Strategy for Cancer Control by the Ministry of Health and Welfare, Japan.

Presented in part at the 38th Annual Meeting of the American Society of Clinical Oncology, Orlando, FL, May 18-21, 2002 (abstract 697).

Authors' disclosures of potential conflicts of interest are found at the end of this article.

Address reprint requests to Takeshi Sano, MD, Gastric Surgery Division, National Cancer Center Hospital, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan; e-mail: tksano@ncc.go.jp.

© 2004 by American Society of Clinical Oncology

0732-183X/04/2214-2767/\$20.00

DOI: 10.1200/JCO.2004.10.184

ABSTRACT

Purpose

Radical gastrectomy with regional lymphadenectomy is the only curative treatment option for gastric cancer. The extent of lymphadenectomy, however, is controversial. The two European randomized trials only reported an increase in operative morbidity and mortality, but failed to show survival benefit, in the D2 lymphadenectomy group. We conducted a randomized controlled trial to compare the Japanese standard D2 and D2 + para-aortic nodal dissection.

Patients and Methods

Only experienced surgeons in both procedures from 24 Japanese institutions participated in the study. Patients with potentially curable gastric adenocarcinoma (T2-subserosa, T3, or T4) who were surgically fit were intraoperatively randomized. Postoperative morbidity and hospital mortality were recorded prospectively in a fixed format and were compared between the two groups in this study.

Results

A total of 523 patients were randomized between July 1995 and April 2001. Postoperative complications were reported in 24.5% of all patients. Although the morbidity for the extended surgery group (28.1%) was slightly higher than the standard group (20.9%), there was no difference in the incidence of four major complications (anastomotic leak, pancreatic fistula, abdominal abscess, pneumonia) between the two groups. Hospital mortality was reported at 0.80%: one patient in each group died of operative complications, while one from each group died of rapid progressive cancer while inpatient.

Conclusion

Specialized surgeons could safely perform gastrectomy with D2 lymphadenectomy in patients with low operative risks. Para-aortic lymphadenectomy could be added without increasing major surgical complications in this setting.

J Clin Oncol 22:2767-2773. © 2004 by American Society of Clinical Oncology

INTRODUCTION

Gastric cancer is the second most common malignancy in the world, and surgical resection remains the only curative treatment option. Lymph node metastases occur during the early stages of this disease, and regional lymphadenectomy is recommended as part of radical gastrectomy. However, the extent of lymphadenectomy to achieve the optimal

result is controversial, and there is no worldwide consensus.

Japanese surgeons first introduced the extended lymphadenectomy procedure, known today as D2, in the 1960s.¹ This technique requires the systematic dissection of lymph nodes in the first tier (perigastric) and the second tier (along the celiac artery and its branches). Early studies have reported that between 30% to 40% of patients

Table 1. Eligibility Criteria of the Study

<p>Before operation</p> <p>Entry criteria</p> <ul style="list-style-type: none"> Histologically proven adenocarcinoma 75 years or younger Forced expiratory volume in 1 second \geq 50% Arterial oxygen pressure in room air \geq 70 mm Hg Creatinine clearance \geq 50 mL/min Written consent <p>Exclusion criteria</p> <ul style="list-style-type: none"> Carcinoma in the remnant stomach Borrmann type 4 (linitis plastica) Synchronous or metachronous malignancy in other organs except for cervical carcinoma in situ and colorectal focal cancer in adenoma Past history of myocardial infarction or positive results of exercise ECG Liver cirrhosis or chronic liver disease with indocyanine green test \geq 10% <p>During operation</p> <ul style="list-style-type: none"> Macroscopic T staging is T2-subserosa, T3, or T4 Potentially curative operation is possible No gross metastasis in para-aortic nodes (frozen section diagnosis not allowed) Peritoneal lavage cytology is negative for cancer cells

with positive lymph node metastases including the second tier lymph nodes, have survived longer than 5 years with D2 lymphadenectomy.² However, D2 gastrectomy has a steep learning curve,³ and may be associated with a higher-than-expected operative morbidity and mortality.

Two European randomized controlled trials comparing D1 and D2 gastrectomy revealed a high operative mortality exceeding 10% in the D2 group.^{4,5} Based on these reports, the British National Health Service Cancer Guidance discourages the use of D2 technique in routine clinical practice.⁶ In contrast, D2 gastrectomy is considered a standard and safe procedure in Japan, where 100,000 cases of gastric cancers are diagnosed every year. General surgeons are taught this technique early during their surgical training.⁷ The Japanese nationwide registry reported an operative mortality of less than 2%, and in specialized institutions, less than 1% for D2 gastrectomy.^{8,9}

Since the eighties, even more radical extended lymphadenectomy procedures had been practiced in many Japanese specialized centers. It was reported that 20% to 30% of patients with nonearly gastric cancer had microscopic metastasis present in the para-aortic nodes.¹⁰⁻¹³ The 5-year survival for these patients has reached 14% to 30% after extended systematic dissection. In addition to D2 lymphadenectomy, lymph nodes around the upper abdominal aorta were dissected, primarily for ultimate local tumor control. However, this extended dissection may not only increase operative morbidity but also may effect the function of other abdominal organs.

There has never been a prospective study to assess the perioperative morbidity and mortality in Japanese patients after D2 gastrectomy or more extended surgery. To evaluate the survival benefit and operative complications of D2 gas-

trectomy and extended para-aortic dissection in gastric cancer surgery, a multi-institutional randomized controlled trial was conducted on behalf of the Japan Clinical Oncology Group (JCOG). The accrual closed with 523 patients. We hereby present the data on the operative morbidity and mortality, which are the secondary end points of this trial. Survival analysis is scheduled to take place in August 2006.

PATIENTS AND METHODS

Objectives and End Points of the Study

A prospective randomized controlled trial was designed to compare the two surgical techniques: the standard lymphadenectomy and the standard lymphadenectomy with the addition of para-aortic node dissection for gastric cancer. Only surgeons with sufficient experience of para-aortic dissection for gastric cancer participated in the trial. Since the role of neoadjuvant and adjuvant chemotherapy was not established, no patients received chemotherapy until recurrent disease was diagnosed.

The primary end point was the overall survival, while the secondary end points were the relapse-free survival, operative morbidity, hospital mortality, and quality of life. Randomization and data handling for this study was performed by the Data Centre of the JCOG, a government-sponsored organization for multi-institutional clinical trials.¹⁴

Eligibility Criteria

Eligibility criteria for this study are shown in Table 1. Patients with advanced gastric cancer deemed curable and fit for surgery were recruited into the trial following informed consent. Borrmann type 4 tumors (linitis plastica) were excluded because of their very poor prognosis after surgery. Liver cirrhosis and ischemic heart disease were important risk factors for mortality after surgery and hence were excluded from the study. Para-aortic lymph node metastasis is extremely rare in T1 (invasion confined

to the mucosa or submucosa) and T2-MP tumors (invasion confined to the muscularis propria); hence, these patients were not eligible for randomization. Only patients diagnosed with T2-SS (subserosal invasion) or deeper tumors at the time of laparotomy were included in the study. T2-SS is clinically recognized as a white discoloration on the serosal surface, without overt tumor serosal exposure.

During the operation, the para-aortic nodes were inspected to exclude patients with gross metastasis (enlarged and/or hard nodes) in this region. Frozen section diagnosis of the para-aortic nodes was forbidden to avoid technical contamination between the two groups of patients. Peritoneal lavage cytology was performed immediately after initial laparotomy, and absence of free cancer cells was confirmed before enrollment.

Random Assignment

While waiting for the result of lavage cytology, the surgeon examined the above eligibility criteria and started the D2 procedure. When the negative cytology result was obtained 30 to 60 minutes later, he informed the JCOG Data Centre for enrollment. Patients were then randomly assigned either to receive standard lymphadenectomy (group A) or extended lymphadenectomy (group B). The sizes of the groups were balanced according to T stage (T2 v T3/T4), tumor growth pattern (expansive v infiltrative growth), and institution. The randomization arm was notified to the surgeon immediately, who then completed the operation according to the allocated protocol.

Surgical Methods

Group A: Standard D2 gastrectomy. Patients were treated with gastrectomy and D2 lymphadenectomy. Depending on the location of the primary tumor, the surgeon performed either a total, proximal subtotal, or distal subtotal gastrectomy. D2 lymphadenectomy was a standard procedure for dissection of tumors located in the upper two thirds of the stomach as defined in the 12th edition of the Japanese Classification (1993)¹⁵ when the study was initially designed. An extended D2 lymphadenectomy was performed for tumors located in the lower third of the stomach, which involves further dissecting the hepatoduodenal nodes (No.12a), retropancreatic nodes (No.13) and nodes along the superior mesenteric vein (No.14v). This technique was frequently performed as a standard procedure in the specialized centers, and thus adopted in this study (all except No.13 have been integrated as "D2" in the 13th edition of Japanese classification¹⁶).

In total or proximal subtotal gastrectomy for proximal tumors, the spleen was removed in principle for splenic hilar lymphadenectomy, while it was preserved in distal subtotal gastrectomy for distal tumors.

Group B: D2 gastrectomy combined with para-aortic lymphadenectomy. Patients in this group had similar procedure to group A, but with additional para-aortic lymph node dissection. The area to be dissected was defined in the Japanese classification (Fig 1). Proximal tumors were treated with the standard D2 lymphadenectomy, and also all "No.16-a2" (para-aortic nodes between the level of the celiac axis and the left renal vein) and "No.16-b1" (para-aortic nodes between the left renal vein and the inferior mesenteric artery) were removed. Standard distal subtotal gastrectomy was performed for the distal tumors including the "No.16-a2" and "No.16-b1" nodes; however, dissection of the left upper lateral nodes ("No.16-a2-lat") was optional.

Both group A and group B patients were followed up according to a fixed schedule, without receiving adjuvant chemotherapy.




Fig 1. Anatomic definitions of para-aortic lymph nodes.¹⁵ The nodes No.16a2 and No.16b1 are defined as "regional nodes" and were dissected in the extended surgery group. Ao, aorta; CA, celiac artery; Eso, esophagus; IMA, inferior mesenteric artery; IVC, inferior vena cava; LRV, left renal vein; SMA, superior mesenteric artery.

Evaluation of Operative Morbidity and Mortality

Operative methods and pathology results were recorded according to the 12th edition of the Japanese Classification of Gastric Carcinoma.¹⁵ The following information was included on the case report form for prospective data collection concerning the four major groups of operative morbidity: presence or absence of anastomotic leak, pancreatic fistula, abdominal abscess, and pneumonia. Anastomotic leak was diagnosed radiologically either on routine postoperative contrast swallow or based on clinical suspicion, and was recorded regardless of its clinical significance. Pancreatic fistula was usually diagnosed when fluid with a high amylase concentration drained from the peripancreatic area for more than 7 days.

Other complications were recorded on a free format. The duration of surgery, blood loss, blood transfusion requirement and reoperation details were also recorded. Hospital mortality was defined as postoperative death of any cause within 30 days, or death within the same hospitalization.

Sample Size

The projected 5-year survival rates for groups A and B patients were 50% and 62%, respectively, and we initially planned to recruit 412 patients (206 each group) to detect this difference with one-sided α error of .05 and statistical power of 80%. At first, the recruitment was slow, but it improved as the study progressed. When the planned recruitment was almost achieved, the JCOG Clinical Trial Review Committee approved the amendment to increase the number of patients to 520 (260 each group) to

Table 2. Patients' Demographics and Tumor Characteristics

	Group A (n = 263)	Group B (n = 260)	Total (N = 523)
Male-female ratio	176/87 = 2.02	182/78 = 2.33	358/165 = 2.17
Age, years			
Median	60	61	61
Range	25-75	27-75	25-75
Tumor diameter, cm			
Median	5.5	5.5	5.5
Range	2-17	2-15.2	2-17
T-stage (macroscopic)			
T2-SS	99	93	192
T3	150	159	309
T4	14	8	22
Tumor location			
Upper 1/3	53	47	100
Middle 1/3	103	103	206
Lower 1/3	107	110	217

NOTE. All data are numbers of patients except where otherwise indicated. Abbreviation: SS, subserosal invasion.

enforce the statistical power to detect 8% difference in the 5-year survival rates, with a 5.5-year accrual period and an additional 5-year follow-up.

Institutions and Quality Control of Surgery

The approval of the institutional review board from all participating institutions was obtained. Initially, the 12 institutions of the Gastric Cancer Surgical Study Group of the JCOG participated in the trial. Twelve institutions were added to increase patient recruitment before February 1999.

All participating surgeons agreed to the technical details for surgery during the planning stages of this trial. Significant experience in gastric cancer surgery, especially experience in extended lymphadenectomy, was a prerequisite for a surgeon's participation in the trial. Surgeons with experience of more than 100 D2 gastrectomies, or institutions with a specialized unit with annual gastrectomy volume of 80 cases or more were selected.

During the recruitment period, participating surgeons and Data Centre representatives met three times per year to monitor the study. In each meeting, videos of para-aortic dissection were presented for critique from four or five institutions, and the technical details were discussed. To assess compliance with lymphadenectomy, dissection, node recovery status in all nodal "stations," and the number of dissected nodes in the para-aortic area were recorded in the case report form, and the results were monitored.

Statistical Methods

The operative morbidity and mortality rates were based on the proportion of the number of cases divided by all registered patients based on the intention-to-treat principle. The differences in proportion between groups were evaluated using Fisher's exact test. Differences in length of hospital stay and blood loss were compared by Wilcoxon test. All *P* values are two-sided, and statistical analysis was done using SAS (SAS Institute, Cary, NC) version 8.12.

Recruitment

Recruitment commenced in July 1995, and closed in April 2001. A total of 523 patients were enrolled: 263 in group A and 260 in group B. A large variance was observed for the number of patients recruited between the institutions. Fifty-three percent of all patients were recruited by the five major hospitals.

The JCOG site-visit audit reported that written consent was available for all except nine patients from one institution. In another institution, an additional six patients had informed consent submitted by a family member.

Patients and Surgery

Patient demographics and tumor characteristics are presented in Table 2. The two groups were well balanced, as there were no significant differences in their baseline data.

The operative details are shown in Table 3. Total gastrectomy was performed in 38% of all patients, and the vast majority of total gastrectomies (186 of 199 cases) were accompanied by splenectomy. Pancreatectomy was confined to those patients whose pancreas was involved by tumor, accounting for 11% of all total gastrectomies. In four cases, proximal subtotal gastrectomy with splenectomy was performed instead of total gastrectomy. Para-aortic lymphadenectomy required longer operation time (median, 63 minutes) and resulted in greater blood loss (median, 230 mL) than the standard D2. Blood transfusion was required approximately twice as often.

Protocol Violation and Ineligible Cases

There were 10 cases of protocol violation (1.9%). In one case, the para-aortic nodes were examined by frozen

Table 3. Operative Details

	Group A (n = 263)	Group B (n = 260)	Total (N = 523)	P
Gastrectomy, No. of patients				.62
Total	102	97	199	
Distal subtotal	160	160	320	
Proximal subtotal	1	3	4	
Splenectomy, No. of patients	98	93	191	.79
Pancreatectomy, No. of patients	9	13	22	.39
Operation time, minutes				< .001
Median	237	300	270	
Range	127-625	153-600	127-625	
Blood loss, mL				< .001
Median	430	660	530	
Range	32-1,810	60-2,885	32-2,885	
Blood transfusion				< .001
No. of cases	37	78	115	
%	14.1	30.0	22.0	
No. of retrieved nodes				< .001
Median	54	74	61	
Range	14-161	30-235	14-235	

section before registration. In another case, the surgeon performed para-aortic dissection despite the allocation to group A because after randomization, he found a positive node behind the common hepatic artery, believed to be strongly suggestive of metastasis in the para-aortic area. The postoperative course of this patient, who was allocated to group A but treated as group B, was uneventful, and analyzing this patient as either group A or group B had no effect on the results in this study. We left this case in group A based on intention-to-treat analysis. In the other eight patients, nodal stations No.13 and/or No.14v were not dissected in distal third tumors.

In another case, the initial histological diagnosis following endoscopic biopsy was poorly differentiated adenocarcinoma but the final histology of the resected stomach revealed gastric lymphoma. We included this patient in the morbidity/mortality analysis, but will exclude their data from the final survival analyses.

Operative Morbidity

The overall operative morbidity rate was 24.5%. The morbidity for group B patients was higher than group A (28.1% and 20.9%, respectively), but the difference did not reach statistical significance ($P = .067$). The incidence of the four major surgical complications was not different between the two groups (Table 4).

There were various other complications reported, and the incidence was significantly higher in group B than group A patients. Paralytic ileus causing significant delay of recommencement of oral feeding, abdominal and/or left pleural lymphorrhea requiring prolonged drainage for more than 1 week, and severe diarrhea, were specific to the extended para-aortic dissection group (Table 4). Reoperation was needed in 12 patients (2.3%), and there was no

difference in the reoperation rate between the two groups. Median hospital stay after surgery was 21 days in group A, and 24 days in group B ($P < .01$).

Hospital Mortality

There were four hospital deaths (0.8%)—two in each group. Each group had one patient who died of postoperative complications, and one died of rapidly progressive cancer. All other patients recovered from surgery and were discharged from hospital.

DISCUSSION

In this randomized controlled trial, the role of para-aortic dissection will be evaluated in terms of survival benefit,

Table 4. Operative Morbidity and Hospital Mortality

	Group A (n = 263)		Group B (n = 260)		P
	No. of Patients	%	No. of Patients	%	
Any complication	55	20.9	73	28.1	.067
Anastomotic leak	6	2.3	5	1.9	.99
Pancreatic fistula	14	5.3	16	6.2	.71
Abdominal abscess	14	5.3	15	5.8	.85
Pneumonia	12	4.6	4	1.5	.072
Others	24	9.1	52	20.0	< .001
Obstruction or ileus	5		11		
Lymphorrhea	0		10		
Left pleural effusion	1		6		
Severe diarrhea	0		3		
Reoperation	5	1.9	7	2.7	.57
Hospital death	2	0.8	2	0.8	.99

operative morbidity/mortality, and quality of life. The results will provide important information and should guide decision making regarding the choice of operative methods. The quality of life and survival among these patients are still in the follow-up phase, and the analyses will take place in 2004 and 2006, respectively. This report compares the morbidity and mortality rates of D2 plus para-aortic node dissection with standard D2 dissection.

There is a wide variation in operative morbidity and mortality following gastric cancer surgery among countries and institutions. The presence of comorbid disease that affects patient fitness for surgery, surgical experience of the operator, and the workload volume seem to be important factors.^{17,18} The mortality for gastrectomy in Western countries often exceeds 5% and approaches 16% in some series.¹⁹⁻²¹ Conversely, Japanese studies have consistently reported a mortality rate of lower than 2% in retrospective observations. To date, the present study is the first large-scale prospective randomized controlled trial in Japan to compare surgical techniques under strict quality control and data management. The extremely low hospital death rate after extended para-aortic lymphadenectomy (0.8%) in this multi-institutional setting confirms the findings from previous retrospective reports.

This trial is a striking contrast to the the Dutch⁴ and British⁵ D1/D2 trials, in which D2 lymphadenectomy was associated with operative mortality rates of 10% and 13%, respectively. One important criticism of the European randomized trials was the issue of learning curve, as many British and Dutch surgeons participating in the trials were new to the D2 procedure. Surgical experience, specific anatomic knowledge, and careful postoperative managements by experienced teams are crucial to the success of this type of surgery. An Italian group appropriately carried out a phase 2 study of D2 lymphadenectomy in selected institutions²² until an acceptable operative mortality rate was achieved, before conducting a randomized controlled trial comparing D1 and D2 gastrectomies.

The D2 gastrectomy procedure is known as "extended lymphadenectomy" in Western countries, while Japanese surgeons employ D2 as a standard technique, and reserve the term "extended" for para-aortic dissection. Lymphatic drainage from the stomach flows to the perigastric nodes and then to the nodes around the celiac axis and its main branches. From here it enters the para-aortic nodes before joining the systemic circulation via the thoracic duct. Hence, the para-aortic nodes may be regarded as the final station of nodes that can be dissected to remove the threat of systemic metastases originating from the lymphatic system. Many Japanese surgeons in specialized centers who performed para-aortic dissection found microscopic metastases in this region, and believe that this type of surgery may be potentially worthwhile. However, the risk associated with para-aortic dissection dictates advanced operative skills and intensive postoperative care.

Therefore, scientific evidence supporting a survival benefit must be obtained before employing this technique in routine gastric cancer surgery.

The very low operative morbidity and mortality achieved in this JCOG trial can be attributed to several factors: (1) we selected a group of fit patients who could tolerate para-aortic dissection in the study. (2) Only specialist surgeons with an established track record of extended lymphadenectomy participated in the trial. (3) High-throughput centers were selected for their operative skills and standardized postoperative management. (4) Pancreatectomy was avoided whenever possible, while splenectomy accompanied total gastrectomy in most cases. We report that there was no significant difference in the overall complications between the two groups; however, the para-aortic dissection group had significantly higher "other" complications (on free format) compared with standard D2. Lymphorrhea and paralytic ileus were more specific to this operation. This observation may be biased because of the surgeon's awareness of the patient's randomization arm of para-aortic dissection.

In the British and Dutch trials, splenectomy with or without distal pancreatectomy was highlighted as a major risk factor for operative morbidity and mortality.^{5,23} Total gastrectomy for proximal tumor requires more advanced surgical skill and is associated with a higher morbidity compared to distal gastrectomy. Proximal gastric tumors are rapidly increasing in number in the western countries,^{24,25} while the incidence remains stable in Japan,²⁶ and this may partly explain the superior results obtained in Japanese studies. However, no difference was observed in the distribution of the primary tumor location between the Dutch⁴ and the Japanese cohort. The proportion of total to distal gastrectomy was also very similar. Therefore, variation in tumor location and type of gastrectomy could not account for the difference in morbidity/mortality, at least between these trials. JCOG recently launched a randomized controlled trial to evaluate the role of splenectomy combined with total gastrectomy in proximal tumors.²⁷

Gastric cancer, though decreasing in incidence worldwide, remains a major health problem in many countries. R0 (no residual disease) resection is the only curative measure; but the more extended the surgery, it is believed the greater is the risk of operative morbidity and mortality. The type of gastrectomy and the extent of lymphadenectomy must be carefully planned for each individual patient with gastric cancer. The Japanese guidelines clearly define D2 gastrectomy as standard surgery²⁸ based on the excellent results in Japanese studies, while the British cancer guidance⁶ discourages D2 based on the poor results of their randomized trial. This contrast should be addressed by surgeons' efforts, such as establishment of specialized standard training systems or production of evidence by high-quality randomized trials in specialized centers.

In conclusion, this study has shown that specialized surgeons could safely perform gastrectomy with D2 lymphadenectomy in patients with low operative risks. Extending the surgery to para-aortic lymphadenectomy did not increase the major operative complications and hospital deaths. However, compared with the D2 procedure, para-aortic dissection requires a longer operation time, leads to a larger volume of blood loss, and longer hospital stay. Until survival benefits are clarified when the data mature sufficiently, para-aortic lymphadenectomy for gastric cancer should be regarded as experimental surgery²⁸ and only performed in special-

ized institutions within the context of a well-designed clinical trial.

Appendix

The appendix is included in the full-text version of this article, available on-line at www.jco.org. It is not included in the PDF (via Adobe® Acrobat Reader®) version.

Authors' Disclosures of Potential Conflicts of Interest

The authors indicated no potential conflicts of interest.

References

- Kajitani T: The general rules for the gastric cancer study in surgery and pathology: Part 1—Clinical classification. *Jpn J Surg* 11:127-139, 1981
- Sasako M, McCulloch P, Kinoshita T, et al: New method to evaluate the therapeutic value of lymph node dissection for gastric cancer. *Br J Surg* 82:346-351, 1995
- Parikh D, Johnson M, Chaglia L, et al: D2 gastrectomy: Lessons from a prospective audit of the learning curve. *Br J Surg* 83:1595-1599, 1996
- Bonenkamp JJ, Songun I, Hermans J, et al: Randomised comparison of morbidity after D1 and D2 dissection for gastric cancer in 996 Dutch patients. *Lancet* 345:745-748, 1995
- Cuschieri A, Fayers P, Fielding J, et al: Postoperative morbidity and mortality after D1 and D2 resections for gastric cancer: Preliminary results of the MRC randomised controlled surgical trial—The Surgical Cooperative Group. *Lancet* 347:995-999, 1996
- NHS Executive: Guidance on Commissioning Cancer Services: Improving outcomes in uppergastro-intestinal cancers, 2001. <http://www.dh.gov.uk/assetRoot/04/08/02/78/04080278.pdf>
- Moriwaki Y, Kobayashi S, Kunisaki C, et al: Is D2 lymphadenectomy in gastrectomy safe with regard to the skill of the operator? *Dig Surg* 18:111-117, 2001
- Fujii M, Sasaki J, Nakajima T: State of the art in the treatment of gastric cancer: From the 71st Japanese Gastric Cancer Congress. *Gastric Cancer* 2:151-157, 1999
- Sano T, Katai H, Sasako M, et al: One thousand consecutive gastrectomies without operative mortality. *Br J Surg* 89:123, 2002
- Takahashi S: Study of para-aortic lymph node metastasis of gastric cancer subjected to superextensive lymph node dissection. *Nippon Geka Gakkai Zasshi* 91:29-35, 1990
- Kitamura M, Arai K, Iwasaki Y: Clinicopathological studies and problems on para-aortic lymph node dissection: D4 dissection. *Nippon Geka Gakkai Zasshi* 97:302-307, 1996
- Isozaki H, Okajima K, Fujii K, et al: Effectiveness of paraaortic lymph node dissection for advanced gastric cancer. *Hepatogastroenterology* 46:549-554, 1999
- Baba M, Hokita S, Natsugoe S, et al: Para-aortic lymphadenectomy in patients with advanced carcinoma of the upper-third of the stomach. *Hepatogastroenterology* 47:893-896, 2000
- Shimoyama M, Fukuda H, Saijo N, et al: Japan Clinical Oncology Group (JCOG). *Jpn J Clin Oncol* 28:158-162, 1998
- Japanese Research Society for Gastric Cancer: Japanese Classification of Gastric Carcinoma (1st English ed). Tokyo, Japan, Kanehara, 1995
- Japanese Gastric Cancer Association: Japanese classification of gastric cancer: 2nd English edition. *Gastric Cancer* 1:8-24, 1998
- Davis P, Sano T: The difference in gastric cancer between Japan, USA and Europe: What are the facts? What are the suggestions? *Crit Rev Oncol Hematol* 40:77-94, 2001
- Birkmeyer JD, Siewers AE, Finlayson EV, et al: Hospital volume and surgical mortality in the United States. *N Engl J Med* 346:1128-1137, 2002
- Msika S, Benhamiche AM, Tazi MA, et al: Improvement of operative mortality after curative resection for gastric cancer: Population-based study. *World J Surg* 24:1137-1142, 2000
- Wanebo HJ, Kennedy BJ, Chmiel J, et al: Cancer of the stomach: A patient care study by the American College of Surgeons. *Ann Surg* 218:583-592, 1993
- McCulloch P: Should general surgeons treat gastric carcinoma? An audit of practice and results, 1980-1985. *Br J Surg* 81:417-420, 1994
- Degiuli M, Sasako M, Ponti A, et al: Morbidity and mortality after D2 gastrectomy for gastric cancer: Results of the Italian Gastric Cancer Study Group prospective multicenter surgical study. *J Clin Oncol* 16:1490-1493, 1998
- Sasako M: Risk factors for surgical treatment in the Dutch Gastric Cancer Trial. *Br J Surg* 84:1567-1571, 1997
- Blot WJ, Devesa SS, Kneller RW, et al: Rising incidence of adenocarcinoma of the oesophagus and gastric cardia. *JAMA* 265:1287-1289, 1991
- Powel J, McConkey CC: Increasing incidence of adenocarcinoma of the gastric cardia and adjacent sites. *Br J Cancer* 62:440-443, 1990
- Ozawa S, Ando N, Kitagawa Y, et al: Does incidence of carcinoma of the oesophagogastric junction increase? *Nippon Geka Gakkai Zasshi* 99:542-546, 1998
- Sano T, Yamamoto S, Sasako M, et al: Randomized controlled trial to evaluate splenectomy in total gastrectomy for proximal gastric carcinoma: Japan Clinical Oncology Group study JCOG 0110-MF. *Jpn J Clin Oncol* 32:363-364, 2002
- Nakajima T: Gastric cancer treatment guidelines in Japan. *Gastric Cancer* 5:1-5, 2002



Morbidity and mortality after D1 and D2 gastrectomy for cancer: Interim analysis of the Italian Gastric Cancer Study Group (IGCSG) randomised surgical trial[☆]

M. Degiuli^{a,*}, M. Sasako^b, M. Calgaro^c, M. Garino^d, F. Rebecchi^e,
M. Mineccia^a, D. Scaglione^f, D. Andreone^g, A. Ponti^h, F. Calvo^a

^aDivision of Surgery, Department of Oncology, Ospedale San Giovanni Antica Sede, ASO Molinette, Via Cavour 31, 10123 Turin, Italy

^bNational Cancer Centre Hospital, Tokyo, Japan

^cDivision of Surgery, Ospedale Mauriziano, Turin, Italy

^dDivision of Surgery, Department of Gastroenterology, ASO Molinette, Turin, Italy

^eDivision of Emergency Surgery, ASO Molinette, Turin, Italy

^fDivision of Surgery, Ospedale Evangelico Valdese, Turin, Italy

^gDivision of Surgery, Ospedale San Luigi Gonzaga, Orbassano, Turin, Italy

^hDivision of Epidemiology, Department of Oncology, Ospedale San Giovanni Antica Sede, ASO Molinette, Turin, Italy

Accepted for publication 19 November 2003

KEYWORDS

Gastric cancer; Extended lymph node dissection; Randomised trial; D1 resection; D2 resection

Summary Background. The disadvantages of D2 gastrectomy have been mostly related to splenopancreatectomy. Unlike two large European trials, we have recently showed the safety of D2 dissection with pancreas preservation in a one-arm phase I-II trial. This new randomised trial was set up to compare post-operative morbidity and mortality and survival after D1 and D2 gastrectomy among the same experienced centres that participated into the previous trial.


Methods. In a prospective multicenter randomised trial, D1 gastrectomy was compared to D2 gastrectomy. Central randomisation was performed following a staging laparotomy in 162 patients with potentially curable gastric cancer.

Findings. Of 162 patients randomised, 76 were allocated to D1 and 86 to D2 gastrectomy. The two groups were comparable for age, sex, site, TNM stage of tumours, and type of resection performed. The overall post-operative morbidity rate was 13.6%. Complications developed in 10.5% of patients after D1 and in 16.3% of patients after D2 gastrectomy. This difference was not statistically significant ($p < 0.29$). Reoperation rate was 3.4% after D2 and 2.6% after D1 resection. Post-operative mortality rate was 0.6% (one death); it was 1.3% after D1 and 0% after D2 gastrectomy.

* For the IGCSG (T. Allone, D. Andreone, M. Calgaro, F. Calvo, L. Capussotti, M. Degiuli, G. R. Fronda, M. Garino, L. Locatelli, P. Mello Teggia, M. Morino, A. Ponti, F. Rebecchi, D. Scaglione)

Corresponding author. Fax: +39-11-8174180.

E-mail address: mdegiuli@hotmail.com



Interpretation. Our preliminary data confirm that in very experienced centres morbidity and mortality after extended gastrectomy can be as low as those showed by Japanese authors. They also suggest that D2 gastrectomies with pancreas preservation are not followed by significantly higher morbidity and mortality than D1 resections.

© 2004 Elsevier Ltd. All rights reserved.

Introduction

Large retrospective Japanese series have shown impressive survival results after D2 gastrectomy (gastric resection together with the removal of level-2 lymph nodes as standardized by the Japanese Society for Research in Gastric Cancer—JSGC) for potentially curable gastric cancer.^{1,2}

Although some non-Japanese series have also reported favourably,^{3,4} these extended lymphadenectomies are still mostly avoided in western countries due to the related increase of post-operative morbidity and mortality.

During the last decade, two European prospective randomised trials have reported that D2 gastric resections are followed by higher morbidity and mortality than D1 resections, and offer no survival benefit over D1 procedures.^{5,6}

The disadvantages of D2 resections have been mostly related to pancreatico-splenectomy, which had been described as an integral part of D2 gastrectomy for all proximal tumours by the JSGC until the 1990s, and consequently was routinely adopted for middle and upper third tumours in the D2 arm of European trials.⁷

Unlike these two European trials, we have recently shown that D2 dissection with pancreas preservation is safe in a one-arm phase I-II trial with a very strict quality control system.⁸

There is not yet evidence from randomised controlled trials that D2 resections give better long-term survival results than standard D1. For this reason our new IGCSG phase III multicentre randomised trial was set up involving the same centres that had already participated into the previous phase 1-2 trial, in order to maintain a homogeneous level of experience among all surgeons.

Patients and methods

Design of the Italian gastric cancer trial

Goals of the trial

- To evaluate whether extending the lymph node dissection to N2 level can improve the survival rate.

- To evaluate whether extending the lymph node dissection to N2 level can decrease the recurrence rate.
- To evaluate morbidity and mortality rates after surgery in both groups of patients.
- To determine the prognostic value of D2 dissection.

Patient selection

Patients less than 80-year-old with histologically proven and potentially curable gastric cancer were eligible for enrolment in the IGCSG trial. Patients undergoing emergency surgery or with severe cardio respiratory, renal or metabolic disease (ASA \geq 4) precluding extended resections were excluded, as were those with a co-existing cancer or distant metastases at preoperative staging. ASA assessment was performed by an experienced consultant anaesthetist in all cases.

Written informed consent was required.

Criteria of curability at laparotomy included:

- Absence of macroscopic involvement of liver and peritoneum (HO, PO).
- Absence of macroscopic involvement of adjacent organs ($T < 4$).
- Absence of macroscopic massive involvement of N2 nodes (enlarged nodes at celiac area).
- Absence of malignant cells in para-aortic nodes (16B1) at biopsy and frozen section.
- Absence of malignant cells in peritoneal washing fluid, during intraoperative fresh examination.
- Absence of macroscopic residual tumour (RO).
- No involvement of the oesophagus, cardia or duodenum.

Surgical definitions

The study was performed according to the rules of the JSGC as regards the extent of stomach removal and the technique of lymph-node dissection,⁹ and to the Japanese Classification of Gastric Carcinoma—second English Edition by the Japanese Gastric Cancer Association,¹² particularly as concern the definitions of classifications and grouping of regional lymph nodes, the extent of lymph node metastasis (N) and the curative potential of gastric resection (Resection A, B or C). In this new classification the regional lymph nodes are

classified into three groups (compartments or levels 1-3), depending upon the location of the primary tumour.

Treatment details

The operative details of the two procedures respected the general rules for gastric cancer study, as described by the Japanese Research Society for Gastric Cancer in 1981. D1 resection entailed removal of the nodes usually defined as perigastric nodes 'en bloc' with the specimen, according to the JGCA. In the D2 arm, during total gastrectomy, the pancreas was removed only when it was suspected to be involved by the tumour. When required (clinical $T > 1$ on the greater curvature of the proximal and middle thirds of the stomach), splenectomy was performed with the pancreas preservation technique as described by Maruyama.¹⁰

Quality control

Only surgeons who participated in the previous one-arm phase 1-2 study on D2 gastrectomy were allowed to participate in this new randomised trial. This restriction permitted maintenance of an homogeneous level of acquired experience among all participating surgeons, as in our previous trial a strict system for quality control had been set up and documented.⁸ Since there is evidence that the learning curve for D2 gastrectomy may be between 20 and 25 cases,¹¹ the randomised part of the study was restricted to the five centres at which more than 25 D2 dissections had been performed during the earlier study. A minimum number of 25 retrieved nodes were required for definition of proper D2 dissection.

Data about post-operative course (hospital stay, blood transfusions, bowel transit, drainage) and early or late morbidity (< or >30 days) and treatment were reported on patient-cards. Hospital mortality (not 30 days mortality) was reported.

Registration and treatment data were regularly collected and sent to the Reference Centre within 30 days of compilation. Follow-up data were sent every 6 months.

Registration and randomisation

Centralised randomisation was performed from the Department of Oncology, Division of Surgery, San Giovanni Antica Sede Hospital, Turin.

The randomisation was performed using random permuted blocks, stratified according to the different operative units. Patients who fulfilled the eligibility criteria during laparotomy were registered by phone call to the randomisation centre. The operator at the randomisation centre completed

the patient-form data on the patient operative unit, time and date of randomisation, then opened the envelope with the randomisation code and immediately communicated it to the operative unit.

In order to document strict adherence to the recruitment procedures, and to prove the absence of selection bias, all patients with a gastric cancer undergoing surgery in each operative unit (eligible or non-eligible) were registered.

Size of the study

The size of the study was calculated on the basis of the effects D1 and D2 surgery on 5 year survival rate. To detect an increase in survival of 15% (from 30% after D1 to 45% of D2 group) 5 years after curative surgery, 160 patients will have to be randomised to each arm ($\alpha = 0.05$ one-sided, $\text{power} = 0.80$).

Results

From January 1999 to December 2002, 296 patients were registered from five participating centres out of the nine centres which participated in our previous trial. Of these, 134 were found not to be eligible for randomisation. Causes of non-eligibility are shown in Table 1. One hundred and sixty-two patients were randomised either to D1 (76) or D2 (86). The two groups were comparable with respect to median age, sex and location of the tumour, as reported in Table 2. They were also similar as regard the extent of gastric resection and stage of disease. Early gastric cancer accounted for 33 per cent of the tumours. The spleen was removed in only 16 patients, four times during a D1 and 12 times during a D2 gastrectomy. A distal pancreatectomy was required in only four patients, when the pancreas was suspected of being involved by the

Table 1. Reasons for exclusion

	No. of patients (%)
Total patients registered	296 (100)
Patients randomised	162 (54.7)
Patients non-eligible	134 (46.3)
No informed consent	8 (6.0)
Metastases/second tumour	14 (10.4)
Nodal spread (N2, N3)	34 (25.2)
Peritoneal spread	26 (19.3)
T4	25 (18.4)
Physical conditions/age	26 (19.3)
No adenocarcinoma (lymphoma)	1 (1.4)

Table 2. Patient characteristics

	D1	D2	Total
No. of patients	76	86	162
Median age (range)	64 (34-81)	61.46 (25-80)	62.01 (29-81)
≥70	22	18	40
Sex (m/f)	39/37	48/38	87/75
<i>Location</i>			
Upper	9	11	20
Medium	18	22	40
Distal	49	52	101
Diffuse	-	1	1
<i>Tumour stage</i>			
pT1	27	27	54
pT2	23	28	51
pT3	26	31	57
<i>Type of resection</i>			
Total gastrectomy	19	22	41
Distal gastrectomy	57	64	121
Splenectomy (S)	3	9	12
Distal pancr + S	1	3	4

tumour. The mean number of nodes removed was 27.0 during a D1 gastrectomy and 36.6 during a D2.

Post-operative course

Table 3 gives data on post-operative course. Overall, the post-operative hospital morbidity was 13.6%. The rate was higher in the D2 group (16.3%) than in D1 group (10.5%), but this difference was not statistically significant. In both groups there were more complications after total than after distal gastrectomy, but again this difference was not significant.

As regards major abdominal infections, no anastomotic dehiscence occurred and only one case of duodenal stump leakage was registered,

while two pancreatic leakages and two cases of acute pancreatitis were observed.

Reoperation was necessary after five major surgical complications (Table 3). The overall hospital mortality was 1/163. This death occurred after a D1 gastrectomy (1/76) and was due to an intraoperative stroke; obviously no significant difference could be observed between D1 and D2 group as concerns mortality.

Post-operative hospital stay

The data on hospital stay excluded the early death (intraoperative), and consequently were based upon 161 patients. The median time of hospital stay was 12 days for D1 groups (mean 13.75, range 8-78) and 12 days for D2 group (mean 13.15, range 8-27). The effect of splenectomy on duration of hospital stay was not clear: patients having received splenectomy stayed in hospital half-a-day more (12.5 days, mean 13.49, range 9-17) than patients without splenectomy 12 days, (mean 12.87, range 8-78, see Table 4).

Discussion

Despite its recent decline, gastric cancer is still a common lethal disease in western countries. For apparently resectable cancers, surgery offers the best loco regional control; but unfortunately, average 5-year survival rates for treated patients remain low in the western world, ranging from 15 to 30%.^{11,13} Over the years, Japanese surgeons have performed radical procedures involving extended lymphadenectomy, and have reported impressive survival figures with extremely low morbidity and mortality.^{1,2,14} Two recent European randomised trials, however, failed to demonstrate a significant

Table 3. Post-operative complications and mortality

	D1 (76)	D2 (86)	Global (162)
Non-surgical complications	4 (5.26%) Cardiac 2 Pulmonary 2	7 (8.13%) Pulmonary 4 Pleural 3	11 (6.79%)
Surgical complications	4 (5.26%) Pancreatic leakage 1 Intrapert. haemorrhage 2 Colonic perforation 1 ^a	7 (8.13%) Pancreatic leakage 1 Intrapert. haemorrhage 1 ^a Duodenal leakage 1 ^a Acute pancreatitis 2 ^a Abdominal abscess 2	11 (6.79%)
Total morbidity	8 (10.52%)	14 (16.27%)	22 (13.58%)
Mortality	1 (1.31%)		1 (0.61%)

^a Requiring reoperation.

Table 4 Lengths of hospital stay

	D1	D2	S0	S+
Days median (range)	12 (8-78)	12 (8-27)	12 (8-78)	12.5 (9-17)
Days mean	13.75	13.15	12.87	13.49

S0, splenectomy not performed; S+, splenectomy performed.

survival benefit of radical D2 gastrectomy over standard D1 resection.^{5,6} The benefit of D2 gastrectomy's potential for reducing loco regional recurrence may be nullified by the significant increase of post-operative morbidity and mortality. These unfavourable results have been attributed to many factors, including the lack of technical experience of surgeons dealing with extended gastrectomy, the large number of elderly patients presenting with associated vascular and cardio respiratory diseases, the large number of centres involved in randomised trials with consequent low quality control, and particularly the distal pancreatico-splenectomy routinely performed during total gastrectomy in the D2 arms of randomised trials. Subset analysis of the MRC and Dutch randomised trials has recently indicated that the poorer outcomes in D2 resections are largely due to pancreas and spleen removal.^{7,15}

We performed a previous prospective multi-centre phase 1-2 study on feasibility and safety of D2 gastrectomy with pancreas preserving technique, involving only a few surgeons. In this study, distal pancreatico-splenectomy was not performed unless the pancreas was suspected of being involved by the tumour. We observed that, when performed in specialized centres, with a strict quality control system, by experienced surgeons, D2 gastrectomy with pancreas preservation could be safe in Western countries. Our morbidity and mortality rates were not only absolutely comparable to those observed after standard resections but also very close to those shown by Japanese surgeons.⁸

Compared to the patients in the Dutch and British trials our patients were younger, and had a higher proportion of early and distal cancers, and these factors may help to partially explain the striking difference between our morbidity and mortality results and those in these trials.

Having reached a good standard of experience in D2 procedures, we planned a new trial, randomising patient to either D1 or D2 gastrectomy.

To maintain a homogenous level of acquired technical experience in D2 procedures, only surgeons already involved in our previous study were allowed to participate in this new trial; this should avoid bias associated with new surgeons who have not yet completed their learning curve. After

careful review of the safety results obtained in the first trial, four out of the nine surgical teams did not join this new randomised trial because completion of their learning curve could not be proven (see above).

These preliminary data seem to confirm our previous reports. Overall morbidity is around 14%; although this figure is a slight underestimate due to the fact that the majority of centres have registered in their database major and minor non-surgical but only major surgical complications, it is very low, and comparable to the best results shown by Japanese authors.¹⁴ The overall morbidity is higher in D2 gastrectomy, but the difference between the two groups of patients is not statistically significant. Moreover, the rate of complications after D2 gastrectomy (16.35%) is considerably better than the rates of both arms (D1 and D2) in the English and Dutch trials.^{5,6}

The ASA grade is a fairly crude and subjective measure of patient fitness, and it is not possible to make realistic comparisons of comorbid pathology and organ functional reserve between our patients and those in the Dutch and British trials. We cannot exclude the possibility that difference between these populations contributed to the difference in morbidity and mortality results. In support of our belief that proper surgical training and quality control played the leading part in our low morbidity, we observed very few 'technical' complications requiring re-operation, such as anastomotic leakage (seen in only one duodenal stump leak).

The importance of pancreatic complications after extended gastric surgery, was confirmed by our data. Although the pancreas was not removed routinely during D2 total gastrectomies, three out of the seven complications registered after a D2 procedure were related to the pancreas (two acute pancreatitis and one pancreatic leakage), and two of these required a reoperation.

Overall mortality was very low, at 0.6%. This rate is comparable to those shown by eastern authors in series from experienced centres, and is strikingly different from the rates of both arms reported in MRC and Dutch trials. Our study was powered to detect a difference in 5 year survival between D1 and D2 surgery: detecting a morbidity or mortality difference would require a larger number of