

chemotherapy, but the disease remains incurable. Usually a platinum-containing regimen is chosen. For ED SCLC, the MST is less than one year with currently available chemotherapy, and long-term survivors are still rare [4,5].

Furthermore, the prognosis is exceedingly poor for patients who receive second-line therapy after relapse. Response is influenced by the time to progression after cessation of first-line therapy. Patients who relapse less than three months after the completion of first-line therapy are termed refractory; they have response rates that are lower than for those patients who relapse more than three months after therapy, who are termed sensitive. The objective for these patients is palliation and increased quality of life, and therefore salvage therapy should be limited to patients with a good performance status (PS) and without significant comorbidities [3].

A water soluble derivative of camptothecin, irinotecan hydrochloride (CPT-11), a topoisomerase I inhibitor, has been synthesised for use in chemotherapy. The chemical structures of irinotecan and its major metabolites found in plasma are shown in Figure 1. Irinotecan is converted by hepatic and peripheral carboxylesterase to its active metabolite 7-ethyl-10-hydroxycamptothecin (SN38). This is subsequently glucuronidated by hepatic uridine diphosphate glucuronosyl transferase-1A1 (UGT 1A1), the enzyme responsible for bilirubin glucuronidation with multi-genetic variants, to SN38-glucuronide (SN38G) [6]. The patient with UGT1A1*28 has an impaired capacity for glucuronidation of SN-38, increased exposure to SN-38, and there is increased clinical toxicity when treated with irinotecan. To measure UGT1A1*28, in August 2005, FDA in the US cleared the Invader Molecular Assay for irinotecan dosing. However, irinotecan activity is not determined by the product of one gene [7]. Irinotecan, SN-38 and SN-38 glucuronide (SN-38G) may be shunted out of the cell via members of the ATP-binding cassette transporters [8]. The metabolism and pharmacogenetics of irinotecan is beyond the scope of this review, but there are some excellent reviews on this subject [9-11].

It should be cautioned that there are drug-drug interactions [12] with irinotecan. Exposure to irinotecan and its active metabolite SN-38 is substantially reduced in patients receiving the CYP3A4 enzyme-inducing anticonvulsants phenytoin, phenobarbital or carbamazepine [13]. Rifampin, rifabutin and St. John's Wort are also CYP 3A4 inducers [14,15]. St. John's Wort is contraindicated during irinotecan therapy. Ketoconazole, a strong inhibitor of CYP3A4 [16], and contraindicated during irinotecan therapy, should be discontinued in patients at least one week prior to starting irinotecan therapy.

In Japan, 1245 cancer patients received irinotecan as a single agent in Phase I or Phase II trials that were conducted to obtain approval for commercial use from the Ministry of Health, Labour and Welfare. Of the 1245 patients, 55 (4.4%) died from toxicities of irinotecan, mainly myelosuppression and/or diarrhoea [17].

The onset of diarrhoea can occur early or be delayed beyond 24 h after injection of irinotecan. Early-onset diarrhoea is a cholinergic effect. Anticholinergic drugs, such as atropine, seem to easily reverse this side effect. Late-onset diarrhoea represents the dose-limiting toxicity (DLT) of irinotecan; it can be severe and life-threatening, especially in combination with neutropenia. Late-onset diarrhoea is treated with loperamide, and identification of high-dose loperamide as an effective remedy for this toxic effect greatly facilitated development of irinotecan [18,19]. These studies established the usefulness of high-dose loperamide. Patients should be instructed to take high-dose loperamide at the first onset of any irinotecan-associated late-onset diarrhoea that has occurred at least 12 h after drug administration. This therapy has been widely used for the management of diarrhoea caused by irinotecan.

For the treatment of SCLC, initial irinotecan is usually administered on days 1 and 8 every 3 weeks or on days 1, 8 and 15 every 4 weeks. The dose ranges from 50 to 70 mg/m² when administered weekly. As an example of the dose modification of irinotecan, Kudoh *et al.* [20] used the following dose modification: irinotecan is not given on days 8 or 15 if the leukocyte or platelet counts were < 3000/μl or < 75,000/μl, respectively. It is also withheld if the patient develops diarrhoea of grade 2 (increase of 4 – 6 stools/day, or nocturnal stools) or worse (grade 3: increase of > 6 stools/day or incontinence; grade 4: physiological consequences requiring intensive care). The next course of treatment can only be initiated if the leukocyte count is ≥ 4000/μl, the platelet count is ≥ 10,000/μl, serum creatinine is less than the upper limit of normal, and diarrhoea has been resolved. There is no dose modification for the leukocyte count, platelet count or diarrhoea during the same course. The dose of irinotecan in the next course was reduced by 10 mg/m² if the leukocyte count was < 2000/μl, the platelet count was < 50,000/μl, or diarrhoea was grade 3 to 4. This dose modification was applied in most studies with minor variation. For example, in some studies, the delay in the irinotecan doses was applied when the leukocyte count was < 2000/μl [21] instead of 3000/μl.

Another available topoisomerase-I inhibitor, topotecan, has achieved response rates of up to 22% in previously treated patients with SCLC and survival almost double that achieved with other single agents. Compared with cyclophosphamide/doxorubicin/vincristine (CAV), single-agent topotecan achieved a higher response rate, longer survival and statistically significant improvements in dyspnoea, hoarseness, fatigue, anorexia and interference with daily activities [22,23]. The incidence of grade 3 – 4 diarrhoea was extremely low (1%). The clinical comparison of these two topoisomerase-I inhibitors has not been tried. This review focuses mainly on the recent results of irinotecan in the treatment of SCLC in connection with patient safety considerations.




Figure 1. Metabolism of irinotecan. Chemical structures of CPT-11 and its major metabolites.

2. Irinotecan containing regimens as front-line treatment

2.1 Irinotecan plus cisplatin for ED SCLC

Clinically, irinotecan was proved to be effective against SCLC [24]. Negoro *et al.* have demonstrated that 13 (37%) out of 35 patients responded, including 33% of previously treated patients and 50% of chemotherapy-naïve patients. In a Phase II trial of irinotecan for previously treated SCLC, the response rate was 47% out of 16 patients [25].

IP was tested in a Phase II trial for patients with previously untreated SCLC [20]. A total of 40 patients (53%) had LD and 35 patients (47%) had ED. Initially, irinotecan 80 mg/m² over 90-minutes infusion was given on days 1, 8 and 15, and cisplatin 60 mg/m² was given every 4 weeks. After 3 of the initial 10 patients experienced severe haematological toxicity, diarrhoea and hepatic toxicity, and one patient died of diarrhoea and neutropenia, the irinotecan dose was reduced to 60 mg/m². The response rate was 84%, with a complete response rate of 29%. The MST was 14.3 months for LD

patients and 13.0 months for ED patients, an encouraging result. Although the survival of LD was not increased significantly, this may be due to the small number of LD SCLC patients accrued. This study prompted a Phase III study of the Japan Clinical Oncology Group (JCOG 9511).

The JCOG conducted a multi-centre, randomised, Phase III study which compared irinotecan plus cisplatin with etoposide plus cisplatin (EP) in patients with ED SCLC (JCOG 9511) (Figure 2) [26]. IP consisted of four 4-week cycles of 60 mg/m² of irinotecan on days 1, 8 and 15, and 60 mg/m² of cisplatin on day 1. The regimen of etoposide and cisplatin consisted of four 3-week cycles of 100 mg/m² of etoposide on days 1, 2 and 3, and 80 mg/m² of cisplatin on day 1. The delivered dose intensity for irinotecan was 80%. The results are listed in Table 1. This study was terminated early because an interim analysis found a statistically significant difference in survival between the two arms. The MST was 12.8 months in the IP arm and 9.4 months in the EP arm ($p = 0.002$). At two years, the proportion of patients surviving was 19.5% in the IP group and 5.2% in the EP

Table 1. IP versus EP in phase III studies.

	IP	EP	p-value	IP	EP	p-value
	JCOG9511 study [26]			Hanna's study [27]		
	(n = 75)	(n = 77)		(n = 210)	(n = 104)	
Irinotecan: delivered dose intensity	80%			90%		
Survival						
Median survival time (months)	12.8	9.4	0.002	9.3	10.2	0.6226
1-year survival (%)	58.4	37.7		35.4	36.7	
2-year survival (%)	19.5	5.2		8.0	7.9	
Haematological						
Neutropenia	65.3	92.2	< 0.001	36.2	86.5	< 0.0001
Anaemia	26.7	29.9	0.72	4.8	11.5	< 0.0268
Thrombocytopenia	5.3	18.2	0.002	4.3	19.2	< 0.0001
Nonhaematological						
Diarrhoea	16	0	< 0.001	21.3	0	0.0001
Response						
Complete response	2.6	9.1		3.6	2.7	
Partial response	81.8	58.4		44.3	40.9	
Overall response	84.4	67.5	0.02	48.0	43.6	
Stable disease	2.6	20.8		4.1	7.3	
Progressive disease	3.9	11.7		20.0	20.0	
Not evaluable	6.5	0		28.1	29.1	

EP: Etoposide and cisplatin; IP: Irinotecan and cisplatin.




Figure 2. Two Phase III randomised trials.

LDH: Lactate dehydrogenase.

group. This was the first study to show the superiority of any one regimen over etoposide plus cisplatin for the front-line treatment of ED SCLC, and IP has become one of the standard regimens for ED SCLC in Japan. Severe myelosuppression

was more frequent in the EP group than in the IP group. On the other hand, severe diarrhoea was more frequent in the IP arm than in the EP arm. Despite the dose modifications, major deviations from the protocol resulted in failure to reduce the dose of chemotherapy (in 6 patients); administration of irinotecan despite the presence of grade 1 (increase of < 4 stools/day) or 2 diarrhoea (in 9 patients); continuation of the study treatment despite grade 2 to 3 pulmonary toxicity (in 3 patients); and continuation of the treatment despite grade 3 hepatic toxicity (in 1 patient). There were 3 treatment-related deaths in the IP arm; one patient died of bleeding from a metastatic site in the lung, another patient died of sepsis associated with neutropenia and diarrhoea, and the third patient died of pneumonia associated with neutropenia. These three treatment-related deaths in the IP arm occurred during the first or second cycle of treatment and were attributed to haematological toxicities of the first cycle. This may indicate that severe haematological toxicities, as well as diarrhoea, during the first cycles of chemotherapy should be managed carefully. All cases of grade 1 to 4 diarrhoea occurred during the first and second cycles of the IP arm but early suspension of treatment may have prevented death associated with diarrhoea in all but one patient, which

involved a protocol violation because the patient was given irinotecan on day 8 of the first cycle despite the presence of grade 1 diarrhoea. This suggests that irinotecan should not be administered to patients with any degree of ongoing diarrhoea above their baseline.

Confirmatory studies are underway; currently, there is only one concluded study showing IP superiority, but it had a small sample size. Additionally, pharmacogenomic differences may exist between Japanese and Western populations.

Hanna *et al.* presented a Phase III trial comparing IP with EP in patients with previously untreated ED SCLC at the ASCO meeting in 2005 (Figure 2, Table 1) [27]. This was designed to confirm the JCOG9511 trial. However, the dose and schedule were modified to increase dose intensity.

The IP arm consisted of cisplatin 30 mg/m² and irinotecan 65 mg/m² on days 1 and 8 every 3 weeks. The EP arm was cisplatin 60 mg/m² on day 1, and etoposide 120 mg/m² on days 1–3 every 3 weeks for 4 cycles, or disease progression, or intolerable toxicity. This was planned to improve tolerability, achieve greater dose intensity and maintain or improve efficacy. The 336 patients were stratified by gender, lactate dehydrogenase level and age, and were randomised in a 2:1 fashion, with 221 treated with IP (median age, 63 years; range, 37–82 years; male, 57.5%) and 109 to EP (median age, 62 years; range, 38–83 years; male, 57.3%). Baseline characteristics were well balanced across the 2 arms, with a high representation of PS of 0 or 1 (IP, 92.3%; EP, 88.2%). After 30 patients with PS 2 were enrolled, study amendment excluded PS 2 patients. Delivered dose intensity of irinotecan was 39 mg (94%), higher than that of the JCOG9511 trial (80%). In both arms, 65% of patients received 4 or more cycles. Selected grade 3 or 4 toxicities in IP versus EP arm were: diarrhoea (21 versus 0%), neutropenia (35 versus 84%), febrile neutropenia (4 versus 11%). Grade 3 or 4 haematological toxicities were significantly more common with EP than IP. There was a trend towards more febrile neutropenia in the EP arm (10 versus 4%), and significant differences were seen in rates of dehydration (13 versus 3%; $p = 0.15$), vomiting (13 versus 4%; $p = 0.0445$), and diarrhoea (21 versus 0%; $p < 0.0001$). The survival of EP in both trials was similar (MST: 10.2 months in this study and 9.4 months in the JCOG9511 trial). However, the MST of IP was 9.3 months in this trial and 12.8 months in the JCOG9511 trial. Differences in outcome of this study from the JCOG trial may be due to pharmacogenomic or patient characteristic differences, or a change in the dose/schedule of IP. Pharmacogenomic studies among ethnic populations are needed to address this issue. It is likely that IP will prove to be at least as effective as other treatments for patients with ED SCLC.

Other Phase III trials will clarify these issues, including a SWOG S0124-randomised Phase III trial with the dose and schedule of each arm the same as the JCOG9511 trial, and a Phase III study started in June 2002 – (NCT00143455) sponsored by Pfizer. In this second study, IP consists of irinotecan

65 mg/m² on days 1 and 8 and cisplatin 80 mg/m² on day 1. EP consists of etoposide 100 mg/m² on days 1–3 and cisplatin 80 mg/m² on day 1 every 3 weeks. The results of these studies are awaited.

The debate continues regarding the optimal dose of combination chemotherapy as related to improvement of the outcome of SCLC. However, the author can state that too low a dose intensity may lead to poor results. Takigawa *et al.* used fractionated administration of IP in 15 patients with ED SCLC [28]. Both irinotecan at a dose of 50 mg/m² and cisplatin at a dose of 60 mg/m² were given on days 1 and 8, and repeated every 4 weeks up to 4 cycles. Although objective response rates were 80%, no complete response (CR) were obtained. The MST was 9.4 months and one-year survival was 40.0%. They stopped enrollment because of no CR and poor survival compared to Kudoh's data [20]. The dose intensity may be low because this regimen had a lower dose of irinotecan (50 mg/m²) and a two-week rest period.

Han *et al.* reported a Phase II study of dose-intensified weekly IP in chemo-naïve patients with ED SCLC [29]. The initial six patients received cisplatin 50 mg/m² followed by irinotecan 90 mg/m² on day 1 and 8 of a 21-day cycle (level I), with one treatment death and three febrile neutropenias. The doses of cisplatin and irinotecan were then reduced to 40 mg/m² and 80 mg/m², respectively (level II). The overall response rate was 97%, with a complete response rate of 26%. The MST was 11.1 months and 1- and 2-year survival rates were 44.1% and 11.8%, respectively. Major grade 3 or 4 toxicities included neutropenia (89%), anaemia (59%) and diarrhoea (27%). There were three treatment-related deaths, occurring in elderly patients aged > 60 years and/or relative poor baseline PS 2 or 3. Although they adopted the oral alkalinisation and control of defecation to prevent irinotecan-induced side effects, especially delayed diarrhoea, they are uncertain whether or not this preventive treatment reduced the observed incidence of severe delayed diarrhoea.

2.2 Irinotecan plus carboplatin for ED SCLC

Schmittel *et al.* studied the DLT and maximum tolerated dose (MTD) of a dose escalation of carboplatin to a fixed dose of irinotecan (IC) in Caucasian patients [30]. They demonstrated that the maximum tolerated dose is irinotecan 50 mg/m² administered on day 1, 8 and 15, and carboplatin at an area under the concentration–time curve (AUC) of 5 mg/ml x min, on day 1 of a 4-week cycle. DLT (neutropenia, thrombocytopenia and diarrhoea) was comparable to the results of the Japanese trial at a dose of 60 mg/m² of irinotecan and AUC = 5 of carboplatin [31].

Subsequently, Schmittel *et al.* presented a randomised Phase II trial comparing IC and etoposide plus carboplatin (EC) in ED SCLC [32]. Chemotherapy-naïve ED SCLC patients were randomly assigned to receive carboplatin AUC = 5 either in combination with 50 mg/m² of irinotecan on days 1, 8 and 15 or with etoposide 140 mg/m² on days 1–3. In the IC arm, treatment was repeated every four weeks; in the EC arm, every

three weeks. IC improved response rate (10% CR and 61% partial response (PR) in IC, 0% CR and 50% PR in EC) and progression free survival (9 months, $p = 0.03$) over standard EC (6 months). The MST was 12 months in the IC arm and 10 months in the EC arm, but with no significant difference. Patients with EC had significantly higher incidence of grade 3 to 4 leucopenia, neutropenia and thrombocytopenia. Grade 3 – 4 diarrhoea developed more frequently in the IC arm (11 versus 6%), but with no significant difference. Haematotoxicity was favourable in the IC arm. They extended into a randomised Phase III trial to assess impact on overall survival, and concluded this study showed that even when carboplatin is used instead of cisplatin, the survival of IC and EC was not significantly different, and that myelosuppression was more frequent in EC than IC.

2.3 Irinotecan plus etoposide for ED SCLC

A Phase II study of irinotecan and etoposide (IE) for chemotherapy-naïve ED SCLC was recently conducted without platinum by the West Japan Thoracic Oncology Group (WJTOG) [33]. A total of 50 patients were enrolled. This regimen consisted of irinotecan 60 mg/m² on days 1, 8 and 15, and etoposide 80 mg/m² on days 2 – 4. The overall response rate was 66% with a complete response rate of 10%. The MST was 11.5 months and the 1-year survival rate was 43.2%. Grade 3 – 4 neutropenia, thrombocytopenia and diarrhoea were 62.9, 4 and 2%, respectively. There was no treatment-related death. This regimen seems to be equal to the EP regimen. The dose intensity of irinotecan and etoposide achieved with this regimen was not adequate. This may be the reason for the low incidence of diarrhoea (2%). A schedule of irinotecan administered on days 1 and 8 at 3-week intervals may be preferred.

2.4 Triplets including irinotecan for ED SCLC

JCOG9902-DI was a randomised Phase II trial to compare two kinds of three-drug combinations of cisplatin, etoposide and irinotecan (PEI regimens) for the treatment of ED SCLC [34]. A total of 60 patients were randomised to receive either arm A (cisplatin 25 mg/m² on day 1, on weeks 1, 3, 5, 7 and 9 and etoposide 60 mg/m² on days 1 – 3, on weeks 2, 4, 6, 8), or arm B (cisplatin 60 mg/m² on day 1, irinotecan 60 mg/m² on days 1, 8, 15 and etoposide 50 mg/m² on days 1 – 3, every week for 4 cycles). Prophylactic G-CSF support was provided in both arms. This study suggested that the PEI combinations in both schedules have significant activity against ED SCLC with acceptable toxicity. The CR rate of 17% and MST of 12.9 months in arm B were much more promising compared with the CR rate of 7% and MST of 8.9 months in arm A. They concluded that arm B should be selected for future Phase III studies. However, because irinotecan administration often needed to be skipped, especially on day 15, they suggested a 3-week schedule in which irinotecan is administered only on days 1 and 8.

Briasoulis *et al.* showed that irinotecan can be safely combined with cisplatin and etoposide in a convenient and simple

schedule of administration over three days [35]. They treated 36 patients with irinotecan on day 1 in combination with fixed doses of cisplatin (20 mg/m²) and etoposide (75 mg/m²), both for 3 consecutive days. Irinotecan dose was escalated from 60 mg/m² by increments of 40 mg/m² in this Phase I trial. The MTD of irinotecan was 140 mg/m² and the recommended optimal dose 120 mg/m². DLTs were febrile neutropenia and grade 3 diarrhoea. This same regimen is being studied with concurrent TRT in a total dose of 54 Gy in 30 fractions (1.8 Gy once daily) [36].

Thompson *et al.* reported a Phase II trial of the Minnie Pearl Cancer Research Network at the 2005 ASCO meeting [37]. They added a molecular targeted agent, imatinib (60 mg/day, per os) to chemotherapy of irinotecan (60 mg/m² on days 1, 8 and 15) and carboplatin (AUC = 4) every 4 weeks. Imatinib targets c-kit expression. Grade 3/4 haematological toxicity included: neutropenia (29%/16%), anaemia (13%/1%) and thrombocytopenia (7%/0%). The response rate was 66% with 10% CR. Grade 3 diarrhoea was observed in 21%. There were no treatment-related deaths. The MST was 8.5 months. This suggests that C-kit expression did not correlate with survival and that imatinib offers no efficacy at a cost of increased toxicity when combined with irinotecan and carboplatin in the treatment of ED SCLC.

3. Irinotecan-containing regimens for relapsed or refractory SCLC

Huisman *et al.* have summarised 21 Phase II studies and 3 randomised trials of second-line chemotherapy in patients with SCLC reported from 1989 to 1999 [38]. They found a cumulative response rate of 21% for multi-drug regimens and 19% for single agents. As yet there is no standard second-line treatment established for patients with SCLC who fail or relapse after front-line treatment.

Irinotecan was combined with various anticancer drugs in doublet or triplet. As doublets, these include cisplatin [39], weekly or every three weeks carboplatin [40,41], etoposide [42], gemcitabine [43,44], ifosfamide [45] and paclitaxel [46]. The responses vary from 10 to 94%, and the MST ranges from 5.8 to 8.9 months. As described earlier on triplet including irinotecan [34,47], a three-drug combination Phase II study of irinotecan, cisplatin and etoposide (PEI regimen) was conducted only for sensitive relapsed SCLC (40 patients) [48]. This Phase II regimen consisted of cisplatin 25 mg/m² weekly for 9 weeks, etoposide 60 mg/m² for 3 days on weeks 1, 3, 5, 7 and 9, and irinotecan 90 mg/m² on weeks 2, 4, 6 and 8 with G-CSF support after day 1 on week 2. The results showed a response rate of 78% (CR rate of 13%) and the MST of 11.8 months. A total of 39 patients (98%) had a good PS of 0 or 1. Grade 3 – 4 neutropenia, thrombocytopenia, and diarrhoea were observed in 73, 33, and 8%, respectively. Nonhaematological toxicities were mild and transient.

Another three-drug combination of cisplatin, ifosfamide and irinotecan with G-CSF was conducted by Fujita *et al.* [49].

The response rate was 94.4% and the MST was 11.1 months, encouraging result. Because of patient selections, it is difficult to make wholly valid conclusions about the most effective regimen based only on Phase II results. However, three-drug combinations containing irinotecan with G-CSF support may have better survival and feasibility than the doublets. The disadvantage is that triplet regimens require G-CSF support, which may make out-patient treatment difficult.

4. Irinotecan containing regimen for LD SCLC

Two meta-analyses showed that the addition of TRT to chemotherapy in patients with LD SCLC improves survival at two and three years by 5.4% [50,51]. In these meta-analyses, non-platinum-based combination chemotherapies were commonly used, with only a few trials using platinum-based chemotherapy. Cisplatin and etoposide plus TRT is now widely regarded as the standard regimen for LD SCLC, and presents acceptable toxicity [52]. Turrisi *et al.* reported results of once-daily versus twice-daily (b.i.d) TRT with four cycles of cisplatin and etoposide. Results showed that the MST was significantly superior in the b.i.d arm (23 versus 19 months) [53].

Irinotecan showed potent radiosensitising effects in human lung tumour xenografts which were related to the cell cycle [54]. Kubota *et al.* reported a pilot study of concurrent etoposide and cisplatin plus accelerated hyperfractionated TRT followed by irinotecan and cisplatin for LD SCLC (JCOG9903) [21]. Treatment consisted of etoposide 100 mg/m² on days 1 – 3, cisplatin 80 mg/m² on day 1, and concurrent b.i.d TRT of 45 Gy beginning on day 2. The IP regimen started on day 29 and consisted of irinotecan 60 mg/m², days 1, 8, 15 and cisplatin 60 mg/m² on day 1, with three 28-day cycles. A total of 31 patients were accrued. Although a pilot study, the MST was 20.2 months and 1-, 2- and 3-year survival rates were 76%, 41%, and 38%, respectively. This encouraging regimen proved safe with acceptable toxicities. A randomised Phase III trial comparing EP with IP following EP plus concurrent TRT for LD SCLC is now underway (JCOG0202).

The WJTOG also conducted a similar regimen [55]. Treatment included cisplatin 80 mg/m² on day 1 and etoposide 100 mg/m² on days 1 – 3 with concurrent TRT (1.5 Gy/b.i.d, a total dose of 45 Gy) followed by 3 cycles irinotecan 60 mg/m² on days 1, 8 and 15 and cisplatin 60 mg/m². The results of 51 patients were almost identical to JCOG9903; overall response and CR rate was 87.8% and 40.8%, respectively; Grade 4 toxicity included neutropenia (83.7%), anaemia (10.2%), thrombocytopenia (0%), diarrhoea (2%) and infection (2%); the MST was 21.5 months and 2-year survival rate was 45.7%.

A Phase II study of IP induction followed by concurrent b.i.d TRT with EP chemotherapy for LD SCLC was conducted [56] and also showed encouraging results. Treatment consisted of two cycles of cisplatin 40 mg/m² and irinotecan 80 mg/m² on days 1 and 8 of a 3-week cycle. This was followed by two 3-week cycles of cisplatin 60 mg/m² on days 43

and 64, and etoposide 100 mg/m² on days 43 – 45 and 64 – 66, with concurrent b.i.d TRT total of 45 Gy beginning on day 43. Thirty-five patients were accrued. The MST was 25 months (but it should be noted that this is a single institution Phase II study).

In these studies, irinotecan was used on an induction or adjuvant setting, and both regimens were encouraging. However, randomised study in which both modalities are compared has not been conducted.

There have been a few trials of concurrent chemoradiotherapy including irinotecan for patients with SCLC as well as NSCLC. Recently, a combined modality treatment of IC and TRT followed by bevacizumab (antiangiogenic anti-VEGF antibody) in the treatment of LD SCLC was conducted in a Phase II trial by the Minnie Pearl Cancer Research Network [57]. Induction therapy consisted of irinotecan 50 mg/m² on days 1 and 8, carboplatin AUC = 5 on day 1, TRT 1.8 Gy single daily dose to total dose of 61.2 Gy (34 fractions), beginning with the 3rd cycle. Chemotherapy was repeated every three weeks for four cycles. As a maintenance therapy, bevacizumab 10 mg/kg i.v. every 2 weeks was given until disease progression, or a maximum of 10 doses (20 weeks) were administered. The response rate was 81% with 28% CR. This regimen was well tolerated with rare grade 4 toxicity and no treatment-related deaths. One-year progression-free and overall survival were 68% and 71%, respectively. These results suggest that irinotecan can be safely administered with TRT concurrently.

Sohn *et al.* also reported a Phase II study of IP with concurrent TRT in LD SCLC [58]. Chemotherapy of irinotecan 60 mg/m² on days 1, 8 and 15 and cisplatin 40 mg/m² on days 1 and 8 were repeated every 4 weeks until a maximum of 6 cycles. TRT of 2 Gy/day was commenced on day 1 of the second chemotherapy cycle up to a total of 54 Gy. The results are not concluded at this time.

Langer *et al.* reported a Phase I study of IP and either b.i.d TRT (45 Gy) or once daily RT (70 Gy) to determine if irinotecan can be safely integrated with concurrent TRT and cisplatin in LD SCLC [59]. Acute DLT was defined as grade 4 oesophagitis, pneumonitis, or diarrhoea; grade 4 neutropenic fever; or any attributable grade 5 fatal toxicity (\leq 90 days after RT). Although preliminary, there has been no attributable DLT in the 26 patients that have been enrolled. In combination with cisplatin 60 mg/m² every 3 weeks x 4 and either b.i.d TRT or once daily TRT, irinotecan 40 mg/m² on days 1 and 8 was safe and feasible. Irinotecan at 50 mg/m² on days 1 and 8 every 3 weeks x4 was also feasible in combination with cisplatin and b.i.d TRT. These reports allow us to conclude that irinotecan can be administered with radiotherapy sequentially or concurrently.

5. Expert opinion and conclusion

Irinotecan is effective against SCLC. For the treatment of ED SCLC, IP regimen is at least comparable to EP regimen.

The degree of myelosuppression in the IP regimens was less than that of the EP regimens. However, diarrhoea was more often observed in the IP than the EP regimens, and can lead to severe side effects when the IP regimen is used incautiously. Pharmacogenetic study of irinotecan may prompt one to use the drug in a safer way to avoid severe toxicities. McLeod suggests that at the least, irinotecan 300 – 350 mg/m² every 3 weeks should not be given to patients with a known UGT1A1*28 genotype until more definitive guidelines are established [60]. However, the use of UGT1A1*28 genotyping to predict toxicity is controversial and its clinical implications are unclear. Furthermore, whether or not these recommendations are also applicable to patients with SCLC should be

elucidated upon because a lower dose of irinotecan is usually administered weekly for the treatment of SCLC, rather than the every 3 or 4 weeks for colorectal cancers. In the coming decade, we must confront the metabolic and pharmacogenomic differences in various populations for the treatment of cancer. For this, international cooperative studies are warranted and indeed of immense importance.

Considering patient safety, irinotecan can indeed be administered relatively safely in patients with SCLC, provided there is careful monitoring of patients, especially regarding diarrhoea and myelosuppression. Further studies to avoid severe toxicities are needed to advance the safe use of this otherwise promising drug.

Bibliography

Papers of special note have been highlighted as either of interest (*) or of considerable interest (**) to readers.

- PISANI P, PARKIN DM, BRAY F, FERLAY J: Estimates of the worldwide mortality from 25 cancers in 1990. *Int. J. Cancer* (1999) 83:18-29.
- PAGE N, READ W, TIERNEY R, ARQUETTE M, PICCIRILLO J, GOVINDAN R: The epidemiology of small cell lung carcinoma. *Proc. Am. Soc. Clin. Oncol.* (2002) 21:305 (Abstract 1216).
- STUPP R, MONNERAT C, TURRISI AT, 3rd, PERRY MC, LEYVRAZ S: Small cell lung cancer: state of the art and future perspectives. *Lung Cancer* (2004) 45(1):105-117.
- State of art of SCLC.
- COMIS RL, FRIEDLAND DM, GOOD BC: Small-cell lung cancer: a perspective on the past and a preview of the future. *Oncology* (1998) 12(1 Suppl 2):44-50.
- WOLF M, TEBBE S, FINK T: First-line chemotherapy in metastatic small-cell lung cancer (SCLC). *Lung Cancer* (2004) 45(Suppl. 2):S223-S234.
- IYER L, KING CD, WHITTINGTON PF *et al.*: Genetic predisposition to the metabolism of irinotecan (CPT-11). Role of uridine diphosphate glucuronosyltransferase isoform 1A1 in the glucuronidation of its active metabolite (SN-38) in human liver microsomes. *J. Clin. Invest.* (1998) 101(4):847-854.
- Role of UGT1A1 in the glucuronidation of SN-38.
- MCLEOD HL: Drug pathways: moving beyond single gene pharmacogenetics. *Pharmacogenomics* (2004) 5(2):139-141.
- SPARREBOOM A, DANESI R, ANDO Y, CHAN J, FIGG WD: Pharmacogenomics of ABC transporters and its role in cancer chemotherapy. *Drug Resist. Updat.* (2003) 6(2):71-84.
- MARSH S, MCLEOD HL: Pharmacogenetics of irinotecan toxicity. *Pharmacogenomics* (2004) 5(7):835-843.
- MA MK, MCLEOD HL: Lessons learned from the irinotecan metabolic pathway. *Curr. Med. Chem.* (2003) 10(1):41-49.
- DE JONG FA, DE JONGE MJ, VERWEIJ J, MATHIJSSSEN RH: Role of pharmacogenetics in irinotecan therapy. *Cancer Lett.* (2005) Epub ahead of print.
- TAKIMOTO CH: Chapter 15: *Pharmacology of Cancer Chemotherapy*. Section 7: Topoisomerase Interactive Agents. 7th Ed., Philadelphia: Lippincott Williams & Wilkins (2005):375-390.
- KUHN JG: Influence of anticonvulsants on the metabolism and elimination of irinotecan. A North American Brain Tumor Consortium preliminary report. *Oncology* (2002) 16(8 Suppl. 7):33-40.
- YONEMORI K, TAKEDA Y, TOYOTA E, KOBAYASHI N, KUDO K: Potential interactions between irinotecan and rifampin in a patient with small-cell lung cancer. *Int. J. Clin. Oncol.* (2004) 9(3):206-209.
- MATHIJSSSEN RH, VERWEIJ J, DE BRUIJN P, LOOS WJ, SPARREBOOM A: Effects of St. John's wort on irinotecan metabolism. *J. Natl. Cancer Inst.* (2002) 94(16):1247-1249.
- KEHRER DF, MATHIJSSSEN RH, VERWEIJ J, DE BRUIJN P, SPARREBOOM A: Modulation of irinotecan metabolism by ketoconazole. *J. Clin. Oncol.* (2002) 20(14):3122-3129.
- OHE Y: Treatment-related death from chemotherapy and thoracic radiotherapy for advanced cancer. *Panminerva Med.* (2002) 44(3):205-212.
- ABIGERGES D, ARMAND JP, CHABOT GG *et al.*: Irinotecan (CPT-11) high-dose escalation using intensive high-dose loperamide to control diarrhea. *J. Natl. Cancer Inst.* (1994) 86(6):446-449.
- Effect of high-dose loperamide for diarrhoea induced by irinotecan.
- ARMAND JP, TERRET C, COUTEAU C, RIXE O: CPT-11. The European experience. *Ann. N Y Acad. Sci.* (1996) 803:282-291.
- KUDOH S, FUJIWARA Y, TAKADA Y *et al.*: Phase II study of irinotecan combined with cisplatin in patients with previously untreated small-cell lung cancer. West Japan Lung Cancer Group. *J. Clin. Oncol.* (1998) 16(3):1068-1074.
- Activity of irinotecan and cisplatin for SCLC as front-line.
- KUBOTA K, NISHIWAKI Y, SUGIURA T *et al.*: Pilot study of concurrent etoposide and cisplatin plus accelerated hyperfractionated thoracic radiotherapy followed by irinotecan and cisplatin for limited-stage small cell lung cancer: Japan Clinical Oncology Group 9903. *Clin. Cancer Res.* (2005) 11(15):5534-5538.
- IP following chemoradiotherapy for LD SCLC.
- VON PAWEL J, SCHILLER JH, SHEPHERD FA *et al.*: Topotecan versus cyclophosphamide, doxorubicin, and vincristine for the treatment of recurrent small-cell lung cancer. *J. Clin. Oncol.* (1999) 17(2):658-667.

23. VON PAWEL J: The role of topotecan in treating small cell lung cancer: second-line treatment. *Lung Cancer* (2003) 41(Suppl. 4):S3-S8.
24. NEGORO S, FUKUOKA M, NIITANI H, TAGUCHI T: Phase II study of CPT-11, new camptothecin derivative, in small cell lung cancer (SCLC) (abstract). *Proc. Am. Soc. Clin. Oncol.* (1991) 10:214.
25. MASUDA N, FUKUOKA M, KUSUNOKI Y *et al.*: CPT-11: a new derivative of camptothecin for the treatment of refractory or relapsed small-cell lung cancer. *J. Clin. Oncol.* (1992) 10(8):1225-1229.
 - Activity of irinotecan for relapsed SCLC.
26. NODA K, NISHIWAKI Y, KAWAHARA M *et al.*: Irinotecan plus cisplatin compared with etoposide plus cisplatin for extensive small-cell lung cancer. *N. Engl. J. Med.* (2002) 346(2):85-91.
 - First report of superiority of IP to EP.
27. HANNA NH, EINHORN L, SANDLER A *et al.*: Randomized, phase III trial comparing irinotecan/cisplatin (IP) with etoposide/cisplatin (EP) in patients (pts) with previously untreated, extensive-stage (ES) small cell lung cancer (SCLC). *Proc. Am. Soc. Clin. Oncol.* (2005) 23:622s.
 - IP is equivalent to EP.
28. TAKIGAWA N, FUJIWARA K, UEOKA H *et al.*: Fractionated administration of irinotecan and cisplatin for treatment of extensive-disease small-cell lung cancer: a phase II study. *Anticancer Res.* (2003) 23(1B):557-560.
29. HAN JY, LEE DH, LEE SY *et al.*: A phase II study of dose-intensified weekly concomitant administration of cisplatin and irinotecan in chemo-naïve patients with extensive-disease small-cell lung cancer. *Med. Oncol.* (2005) 22(3):281-290.
30. SCHMITTEL A, SCHULZE K, HUTTER G, KREBS P, THIEL E, KEILHOLZ U: Phase I dose escalation study of carboplatin to a fixed dose of irinotecan as first-line treatment of small cell lung cancer. *Onkologie* (2004) 27(3):280-284.
31. FUKUDA M, OKA M, SODA H *et al.*: Phase I study of irinotecan combined with carboplatin in previously untreated solid cancers. *Clin. Cancer Res.* (1999) 5(12):3963-3969.
32. SCHMITTEL A, FISCHER VON WEIKERSTHAL L, SEBASTIAN M *et al.*: Irinotecan plus carboplatin versus etoposide plus carboplatin in extensive disease small cell lung cancer: A randomized phase II trial. *Proc. Am. Soc. Clin. Oncol.* (2005) 23:632s.
33. KUDOH S, NAKAMURA S, NAKANO T *et al.*: Irinotecan and etoposide for previously untreated extensive-disease small cell lung cancer: a phase II trial of West Japan Thoracic Oncology Group. *Lung Cancer* (2005) 49(2):263-269.
34. SEKINE I, NISHIWAKI Y, NODA K *et al.*: Randomized phase II study of cisplatin, irinotecan and etoposide combinations administered weekly or every 4 weeks for extensive small-cell lung cancer (JCOG9902-DI). *Ann. Oncol.* (2003) 14(5):709-714.
 - Three drug combination for ED SCLC.
35. BRIASOULIS E, SAMANTAS E, KALOFONOS H *et al.*: Phase I study of etoposide, cisplatin and irinotecan triplet in patients with advanced-stage small-cell lung cancer. *Cancer Chemother. Pharmacol.* (2005) 56(5):521-528.
36. SAMANTAS E, SYRIGOS K, BRIASOULIS E *et al.*: Phase II study of cisplatin plus etoposide over 3 days combined with irinotecan and early thoracic radiotherapy (TRT) in patients with limited disease (LD) small cell lung cancer (SCLC), preliminary results. *Lung Cancer* (2005) 49(Suppl. 2):S327.
37. THOMPSON DS, HAINSWORTH JD, SPIGEL DR *et al.*: Irinotecan (I), carboplatin (C), and imatinib (IM) in the first-line treatment of extensive-stage small cell lung cancer (SCLC): a phase II trial of the Minnie Pearl Cancer Research Network. *Proc. Am. Soc. Clin. Oncol.* (2005) 23:632s.
38. HUISMAN C, POSTMUS PE, GIACCONE G, SMIT EF: Second-line chemotherapy and its evaluation in small cell lung cancer. *Cancer Treat. Rev.* (1999) 25(4):199-206.
39. ANDO M, KOBAYASHI K, YOSHIMURA A *et al.*: Weekly administration of irinotecan (CPT-11) plus cisplatin for refractory or relapsed small cell lung cancer. *Lung Cancer* (2004) 44(1):121-127.
40. NAKA N, KAWAHARA M, OKISHIO K *et al.*: Phase II study of weekly irinotecan and carboplatin for refractory or relapsed small-cell lung cancer. *Lung Cancer* (2002) 37(3):319-323.
41. HIROSE T, HORICHI N, OHMORI T *et al.*: Phase II study of irinotecan and carboplatin in patients with the refractory or relapsed small cell lung cancer. *Lung Cancer* (2003) 40(3):333-338.
42. MASUDA N, MATSUI K, NEGORO S *et al.*: Combination of irinotecan and etoposide for treatment of refractory or relapsed small-cell lung cancer. *J. Clin. Oncol.* (1998) 16(10):3329-3334.
43. AGELAKI S, SYRIGOS K, CHRISTOPHYLAKIS C *et al.*: A multicenter phase II study of the combination of irinotecan and gemcitabine in previously treated patients with small-cell lung cancer. *Oncology* (2004) 66(3):192-196.
44. SCHUETTE W, NAGEL S, JUERGENS S *et al.*: Phase II trial of gemcitabine/irinotecan in refractory or relapsed small-cell lung cancer. *Clin. Lung Cancer* (2005) 7(2):133-137.
45. ICHIKI M, GOHARA R, RIKIMARU T *et al.*: Combination chemotherapy with irinotecan and ifosfamide as second-line treatment of refractory or sensitive relapsed small cell lung cancer: a phase II study. *Chemotherapy* (2003) 49(4):200-205.
46. RUSHING DA: Phase I/II study of weekly irinotecan and paclitaxel in patients with SCLC. *Oncology* (Huntingt) (2000) 14(7 Suppl. 5):63-66.
47. SEKINE I, NISHIWAKI Y, KAKINUMA R *et al.*: Phase I/II trial of weekly cisplatin, etoposide, and irinotecan chemotherapy for metastatic lung cancer: JCOG 9507. *Br. J. Cancer* (2003) 88(6):808-813.
48. GOTO K, SEKINE I, NISHIWAKI Y *et al.*: Multi-institutional phase II trial of irinotecan, cisplatin, and etoposide for sensitive relapsed small-cell lung cancer. *Br. J. Cancer* (2004) 91(4):659-665.
 - Three drug combination for sensitive relapsed SCLC.
49. FUJITA A, TAKABATAKE H, TAGAKI S, SEKINE K: Combination of cisplatin, ifosfamide, and irinotecan with rhG-CSF support for the treatment of refractory or relapsed small-cell lung cancer. *Oncology* (2000) 59(2):105-109.
50. PIGNON JP, ARRIAGADA R, IHDE D *et al.*: A meta-analysis of thoracic radiotherapy for small-cell lung cancer. *N. Engl. J. Med.* (1992) 327:1618-1624.

51. WARDE P, PAYNE D: Does thoracic irradiation improve survival and local control in limited-stage small-cell carcinoma of the lung? *J. Clin. Oncol.* (1992) 10:890-895.
52. MURRAY N: Treatment of small cell lung cancer: the state of the art. *Lung Cancer* (1997) 17(Suppl 1):S75-S89.
53. TURRISI AT, KIM K, BLUM R *et al.*: Twice-daily compared with once-daily thoracic radiotherapy in limited small-cell lung cancer treated concurrently with cisplatin and etoposide. *N. Engl. J. Med.* (1999) 340:265-271.
54. TAMURA K, TAKADA M, KAWASE I *et al.*: Enhancement of tumor radio-response by irinotecan in human lung tumor xenografts. *Jpn. J. Cancer Res.* (1997) 88(2):218-223.
55. KASHII T, SAITO H, NEGORO S *et al.*: Phase II study of cisplatin plus etoposide with concurrent thoracic radiotherapy (TRT) followed by irinotecan plus cisplatin in limited stage small-cell lung cancer (SCLC); A West Japan Thoracic Oncology Group Trial. *Lung Cancer* (2005) 49(Suppl. 2):S322.
56. HAN JY, CHO KH, LEE DH *et al.*: Phase II study of irinotecan plus cisplatin induction followed by concurrent twice-daily thoracic irradiation with etoposide plus cisplatin chemotherapy for limited-disease small-cell lung cancer. *J. Clin. Oncol.* (2005) 23(15):3488-3494.
57. RAEFSKY EL, SPIGEL DR, GRECO FA *et al.*: Irinotecan (I), carboplatin (C), and radiotherapy (RT) followed by bevacizumab (B) in the treatment of limited-stage small cell lung cancer (SCLC): a phase II trial of the Minnie Pearl Cancer Research Network. *Proc. Am. Soc. Clin. Oncol.* (2005) 23:633s.
 - Irinotecan, carboplatin and concurrent RT for LD SCLC.
58. SOHN J, MOON Y, LEE C *et al.*: Phase II trial of irinotecan and cisplatin with concurrent radiotherapy in limited-disease small cell lung cancer. *Proc. Am. Soc. Clin. Oncol.* (2005) 23:662s.
 - Irinotecan, cisplatin and concurrent RT for LD SCLC.
59. LANGER C, SWANN S, WERNER-WASIK M *et al.*: Phase I study of combination irinotecan and cisplatin and either twice daily thoracic radiation (45Gy) or once daily thoracic radiotherapy (70Gy) in patients with limited small cell lung carcinoma (SCLC): Early toxicity analysis of RTOG 0241. *Lung Cancer* (2005) 49(Suppl. 2):S323.
60. MCLEOD HL, WATTERS JW: Irinotecan pharmacogenetics: is it time to intervene? *J. Clin. Oncol.* (2004) 22(8):1356-1359.
 - Editorial for irinotecan pharmacogenetics.

Affiliation

Masaaki Kawahara MD
Director, Department of Medical Services,
National Hospital Organization Kinki-chuo
Chest Medical Center, 1180 Nagasone,
Sakai, Osaka, 591-8555, Japan
Tel: +81 72 252 3021; Fax: +81 72 250 4034;
E-mail: kawaharam@kch.hosp.go.jp

EGFR and ErbB2 mutation status in Japanese lung cancer patients

Hidefumi Sasaki^{1*}, Shigeki Shimizu², Katsuhiko Endo¹, Minoru Takada³, Masaaki Kawahara³, Hisaichi Tanaka³, Akihide Matsumura³, Keiji Iuchi³, Hiroshi Haneda¹, Eriko Suzuki¹, Yoshihiro Kobayashi¹, Motoki Yano¹ and Yoshitaka Fujii¹

¹Department of Surgery II, Nagoya City University Medical School, Nagoya, Japan

²Department of Pathology II, Nagoya City University Medical School, Nagoya, Japan

³National Hospital Organization, Kinki-chuo Chest Medical Center, Osaka, Japan

Much evidence has accumulated that the epidermal growth factor receptor (*EGFR*) and its family members are strongly implicated in the development and progression of lung cancers. Somatic mutations of the *EGFR* gene were found in about 25–40% of Japanese lung cancer patients. More recently, *erbB2* mutations are found in about 4% of European-derived lung cancer patients. We have investigated *EGFR* and *erbB2* mutation status in 95 surgically treated nonsmall cell lung cancer (NSCLC) cases from Nagoya City University Hospital. Seventy-five adenocarcinoma cases were included. The presence or absence of *EGFR* and *erbB2* mutations of kinase domains were analyzed by reverse transcription polymerase chain reaction (RT-PCR) amplifications and direct sequences. We have also investigated *erbB2* mutation status in 27 surgically treated NSCLC cases followed by treatment with gefitinib from Kinki-chuo Chest Medical Center. *EGFR* mutations (CTG→CGG; L858R) were found from 14 of 95 lung cancer patients. We also detected the deletion 1a-type mutations from 9 patients and deletion 4-type mutations from 6 patients in exon 19. In exon 20, 4 mutations including 2 novel mutations were found. Total *EGFR* mutations were present in 35 patients (36.8%). These mutation statuses were significantly correlated with gender (women 73.3% vs. men 20%, $p < 0.0001$), smoking status (never smoker 69.4% vs. smoker 16.9%, $p < 0.0001$), pathologic subtypes (adenocarcinoma 45.1% vs. nonadenocarcinoma 12.5%, $p = 0.0089$) and differentiation status of the lung cancers (well 51% vs. moderately or poorly 18.4%, $p = 0.0021$). On the other hand, *erbB2* mutation was only found from 1 of 95 patients, at exon 20. This patient was female and a never smoker with adenocarcinoma. This 12 nucleotide insertion mutation (2324–2325 ins ATACGTGATGGC) was located in the exon 20 at kinase domain (775–776 ins YVMA). There was no *erbB2* mutation in 27 gefitinib-treated NSCLC patients. In total, we have found only 1 *erbB2* mutation from 122 (0.8%) Japanese NSCLC patients. There was a significantly higher *erbB2* positive (2+/3+) ratio in *EGFR* mutant patients (13/25, 52.0%) compared to *EGFR* wild-type patients (10/62, 16.1%; $p = 0.0247$). The NSCLC specimen with *erbB2* mutation showed 1+ immunoreactivity. The *EGFR* mutation status might correlate with the clinicopathologic features related to good response to gefitinib, such as gender, smoking history and pathologic subtypes of lung cancers. However, *erbB2* mutation is rare from Japanese lung cancer and is of limited value for molecular target therapy.

© 2005 Wiley-Liss, Inc.

Key words: EGFR; lung cancer; mutations; erbB2; Japanese

Lung cancer is a major cause of death from malignant diseases, due to its high incidence, malignant behavior and lack of major advancements in treatment strategy.¹ Lung cancer was the leading indication for respiratory surgery (42.2%) in 1998 in Japan.² More than 15,000 patients underwent surgical operations at Japanese institutions in 1998.² The clinical behavior of the lung cancer is largely associated with its stage. The cure of the disease by surgery is only achieved in cases representing an early stage of lung cancer.³

There is much accumulated evidence that epidermal growth factor receptor (*EGFR*) and its family members are strongly implicated in the development and progression of numerous human tumors, including lung cancer.^{4,5} The erbB family comprises 4 structurally related receptors: ErbB1 (*EGFR*), ErbB2 (*HER2-neu*), ErbB3 and ErbB4. On ligand stimulation, the receptor forms either

homodimers or heterodimers, which activate their cytoplasmic domain. This tyrosine-auto-phosphorylated region functions as a docking site for messenger proteins, which initiate cascades of cytoplasmic and nuclear mitogenic pathways.⁶ Inhibition of this pathway is facilitated by several newly developed compounds that have shown promising results in preclinical and clinical trials.⁷ The *EGFR* tyrosine kinase inhibitor, gefitinib, was approved in Japan for the treatment of nonsmall cell lung cancer (NSCLC) since 2002. Trastuzumab is a recombinant DNA-derived monoclonal antibody that selectively binds to p185 HER2, the protein product of *erbB2*. Trastuzumab was approved for breast cancer⁸ and clinical trials for NSCLC is underway.^{9,10}

Recently, we have found that novel *EGFR* mutations' status at ATP binding pockets in Japanese NSCLC patients were correlated with the clinicopathologic features related to good response to gefitinib.¹¹ These *EGFR* mutations are predominantly found in Japanese lung cancer patients (about 25%) when compared to USA patients (about 8%^{12–14} to 10%¹⁵). Kasaoka *et al.* have reported that the *EGFR* mutation ratio is 40% of Japanese lung cancer patients.¹⁶ Actually, *EGFR* mutations in lung cancer have been correlated with clinical response to gefitinib therapy *in vivo* and *in vitro*.^{11–13} More recently, it has been reported that novel *erbB2* mutations at kinase domain were found in 4% of European-derived NSCLC patients.¹⁷

To determine the *EGFR* and *erbB2* mutation status in Japanese lung carcinoma for screening purposes, we investigated *EGFR* and *erbB2* mutation status by the RT-PCR amplifications and direct sequences. The findings were compared to the clinicopathologic features of lung cancer.

Material and methods

Study subjects

The study group included 95 lung cancer patients who had undergone surgery (but did not receive gefitinib) at the Department of Surgery II, Nagoya City University Medical School between 1997 and 2002. We have also investigated *erbB2* mutation status for 27 NSCLC patients who had undergone surgery followed by treatment with gefitinib at the National Hospital Organization, Kinki-chuo Chest Medical Center. The lung tumors were classified according to the general rule for clinical and pathologic record of lung cancer in Japan.¹⁸ All tumor samples were immediately frozen and stored at -80°C until assayed.

The clinical and pathologic characteristics of the 95 lung cancer patients are as follows: 52 cases at stage I, 9 at stage II and 34 at stage III–IV. The mean age was 64.9 years (range, 42–82). Among the 95 lung cancer patients, 71 (74.7%) were diagnosed as adeno-

Grant sponsor: Grant-in-Aid for Cancer Research (16-1) from the Ministry of Health, Labour and Welfare; Grant sponsor: AstraZeneca Research Grant 2004.

*Correspondence to: Department of Surgery II, 1 Kawasumi, Mizuhocho, Mizuho-ku, Nagoya 467-8601, Japan. Fax: +81-52-853-6440. E-mail: hisasaki@med.nagoya-cu.ac.jp

Received 11 March 2005; Accepted after revision 18 April 2005

DOI 10.1002/ijc.21301

Published online 7 July 2005 in Wiley InterScience (www.interscience.wiley.com).

carcinoma, 17 (17.9%) were squamous cell carcinoma and 4 (4.2%) were adenocarcinoma. The samples from these patients had never been sequenced for *EGFR* before.

PCR assays for *EGFR* and *erbB2*

Total RNA was extracted from lung cancer tissues and adjacent nonmalignant lung tissues using Isogen kit (Nippon Gene, Tokyo, Japan) according to the manufacturer's instructions. RNA concentration was determined by spectrophotometer and adjusted to a concentration of 200 ng/ml. About 10 cases were excluded because tumor cells were too few to sufficiently extract tumor RNA. RNA (1 μ g) was reverse transcribed by Superscript II enzyme (Gibco BRL, Gaithersburg, MD) with 0.5 μ g oligo (dT)₁₂₋₁₆ (Amersham Pharmacia Biotech, Piscataway, NJ). The reaction mixture was incubated at 42°C for 50 min and then at 72°C for 15 min. We then used 1 μ l of each DNA for PCR analyses. The PCR reactions were performed using LA-Taq kit (Takara Bio, Shiga, Japan) in a 25 μ l reaction volume. The primer sequences for *EGFR* gene for kinase domain (exons 18–21) were as follows: the forward primer, 5-CTCTTACACCCAGTGGAGAA-3 and the reverse primer, 5-CATCCACTTGATAGGCACTT-3 (572 bp). The cycling conditions were as follows: initial denaturation at 94°C for 5 min, followed by 35 cycles at 94°C for 40 sec, 60°C for 40 sec, 72°C for 45 sec. The primer sequences for *erbB2* gene for kinase domain (exons 19–22) were as follows: the forward primer, 5-CGCTTTTGGCACAGTCTACA-3 and the reverse primer, 5-GGGATCCCATCGTAAGGTTT-3 (594bp). The cycling conditions were as follows: initial denaturation at 94°C for 5 min, followed by 35 cycles at 94°C for 40 sec, 60°C for 40 sec, 72°C for 45 sec. The products were purified by Qiagen PCR purification kit (Qiagen, Valencia, CA). Amplified cDNAs were separated on 1% agarose gels, and the bands were visualized by ethidium bromide and photographed under ultraviolet transillumination. These samples were sequenced by ABI prism 3100 analyzer (Applied Biosystems Japan, Tokyo, Japan) and analyzed by BLAST and chromatograms by manual review.

Immunohistochemistry

Tissue blocks were cut into 4 mm sections and mounted on silane-coated slides. The slides were then deparaffinized in xylene, dehydrated in a grade alcohol series and blocked for endogenous peroxidase with 3% H₂O₂ in absolute methanol. After microwave pretreatment in Blockace solution, immunostaining was done at 4°C overnight with a rabbit polyclonal c-*erbB2* oncoprotein antibody (A04085, DakoCytomation, Glostrup, Denmark) at a 1:200 dilution. The expression of *erbB2* was scored as follows: –, no discernible staining or <10% of cell stained; 1+, >10% of cytoplasmic staining or plasma membrane staining with weak intensity; 2+, >10% of plasma membrane staining with moderate intensity; and 3+, >10% of plasma membrane staining with strong intensity.

Statistical methods

Statistical analyses were done using the Mann-Whitney U-test for unpaired samples and Wilcoxon's signed rank test for paired samples. Linear relationships between variables were determined by means of simple linear regression. Correlation coefficients were determined by rank correlation using Spearman's test and χ^2 test. The overall survival of lung cancer patients was examined by the Kaplan-Meier methods, and differences were examined by the log-rank test. All analysis was done using the Stat-View software package (Abacus Concepts, Berkeley, CA) and was considered significant when the *p*-value was less than 0.05.

Results

EGFR gene mutation status in Japanese lung cancer patients

Using the primer sets for *EGFR* kinase domain, a PCR product of 572 bp was obtained. When we visualized the PCR products




FIGURE 1 – Novel *EGFR* mutation at exon 20. Top: a male well-differentiated adenocarcinoma patient had the novel 2312–2313 insertion CAA. Bottom: a female, well-differentiated adenocarcinoma patient had the novel 2319–2320 insertion AACCCCCAC.

with 1% agarose gel, these samples were further studied. In exon 18, there was no G719S mutation found from this study. In exon 19, 9 patients had the del 1a type mutation, 6 patients had the deletion 4 type mutation and 1 patient had the del 1b type mutation. Seven were male and 10 were female. Thirteen were nonsmokers and 4 were smokers. Fifteen patients had adenocarcinoma, 1 had squamous cell carcinoma and 1 had adenocarcinoma. Three of the tumors were moderately differentiated, 2 were poorly differentiated and 11 were well differentiated. Five of 15 adenocarcinomas showed bronchioloalveolar carcinoma (BAC) pattern at the edge of tumor. Thus *EGFR* mutation status at exon 19 was significantly correlated with gender ($p = 0.0172$) and tobacco-smoking ($p = 0.0008$) but not with pathologic stages (stage I vs. II–IV, $p = 0.9144$), subtypes (adenocarcinoma vs. non-adenocarcinoma, $p = 0.2675$) and differentiation of lung cancer (well vs. moderately or poorly differentiated, $p = 0.3812$).

In exon 20, 3 patients had the heterozygous in-frame insertion mutations. Two were male and 1 was female. All 3 were smokers. A female, well-differentiated adenocarcinoma patient had the novel 2319-2320 insertion AACCCCCAC. A male well-differentiated adenocarcinoma patient had the novel 2312-2313 insertion CAA (Fig. 1). We have found one point mutation, C2369T (T790M). This patient also has the predominant L858R mutation (Fig. 2).

For exon 21, 14 patients had the L858R mutation and 1 patient had the L861Q mutation. Four were male and 11 were female. Twelve were nonsmokers and 3 were smokers. All 15 patients had adenocarcinoma, 1 was moderately differentiated and 14 were well differentiated. Six of 15 adenocarcinomas exhibited the BAC pattern at the edge of the tumor. Thus, exon 21 mutation status was significantly correlated with gender ($p = 0.0005$), smoking status ($p = 0.0007$), pathologic stages ($p = 0.0152$), the pathologic subtypes ($p = 0.0329$) and differentiation of lung cancer ($p = 0.0033$).

The mutations detected in lung cancer specimens from 95 lung cancer patients are summarized in Table I. Taken together, 36 mutations were found from 35 lung cancer samples in our analysis. Total *EGFR* mutations were present in 35 patients (36.8%). These mutation statuses were significantly correlated with gender (women 73.3% vs. men 20%, $p < 0.0001$), smoking status (never smoker 69.4% vs. smoker 16.9%, $p < 0.0001$), pathologic subtypes (adenocarcinoma 45.1% vs. nonadenocarcinoma 12.5%, $p = 0.0089$) and differentiation status of the lung cancers (well 51% vs. moderately or poorly 18.4%, $p = 0.0021$).

The overall survival of 95 lung cancer patients from Nagoya City University, with follow-up through December 30, 2003, was studied in reference to the *EGFR* mutation status. The patient with the mutation in the *EGFR* gene ($n = 35$, 4 were dead) had a significantly better prognosis than the patient with wild-type *EGFR* ($n = 60$, 20 were dead; log-rank test $p = 0.0143$, Breslow-Gehan-Wilcoxon test $p = 0.0220$), although the observation period was short (Fig. 3). But a multivariate analysis revealed that pathologic




FIGURE 2 – The premoninant L858R (2573 T to G) mutation in exon 21 (top) and T790M (2369 C to T) mutation at exon 20 (bottom) within the EGFR kinase domain.

TABLE I – CLINICOPATHOLOGIC DATA OF 95 LUNG CANCER PATIENTS

Factors	EGFR gene status		p-value
	Mutation patients	Wild-type patients	
Mean age (years) 64.9 ± 9.0	35	60	
Stage			
I	25 (72.4%)	27 (45.8%)	0.0274
II–IV	10 (28.6%)	32 (54.2%)	
Lymph node metastasis			
N0	8 (22.9%)	21 (35.0%)	0.3119
N+	27 (77.1%)	39 (65.0%)	
BI			
Never smoker	25 (71.4%)	11 (34.0%)	0.001
Smoker	10 (28.6%)	49 (66.0%)	
Differentiation			
Well	26 (78.8%)	23 (42.6%)	0.0021
Moderately or poorly	7 (21.2%)	31 (57.4%)	
Pathologic subtypes			
Adeno	32 (91.4%)	39 (74.7%)	0.0089
Nonadeno	3 (8.6%)	21 (25.3%)	
Age			
≤ 65	19 (54.3%)	29 (48.3%)	0.7269
> 65	16 (45.7%)	31 (51.7%)	
Gender			
Male	13 (37.1%)	52 (86.7%)	< 0.0001
Female	22 (62.9%)	8 (13.3%)	

N+, lymph node metastasis positive; Adeno, adenocarcinoma.

stage ($p = 0.0006$) was the only significant factor but not *EGFR* mutation ($p = 0.1824$).

ErbB2 gene mutation status in Japanese lung cancer patients

We identified only one *erbB2* mutation from 95 NSCLC patients. This 12-nucleotide insertion mutation (2324–2325 ins ATACGTGATGGC) was located in the exon 20 at kinase domain (775–776 ins YVMA) (Fig. 4). This patient was a female non-smoker with well-differentiated adenocarcinoma, without *EGFR* mutation. Adjacent normal lung tissue exhibited a wild-type sequence for the *erbB2* gene, suggesting that this mutation was somatic. We have also done sequencing for 27 gefitinib-treated NSCLC patients. Among 27 patients, 9 patients had *EGFR* mutations (data not shown). However, no *erbB2* mutation was found




FIGURE 3 – The patient with a mutation in the *EGFR* gene ($n = 35$, 4 were dead) had a significantly better prognosis than the patient with wild-type *EGFR* ($n = 60$, 20 were dead) (log-rank test, $p = 0.0143$; Breslow-Gehan-Wilcoxon test, $p = 0.0220$), although the observation period was short.




FIGURE 4 – Detection of the insertion mutation in the *erbB2* gene in genomic DNA extracted from lung cancer. (a) The 12 nucleotide insertion mutation (2324–2325 ins ATACGTGATGGC) was located in the exon 20 at kinase domain (775–776 ins YVMA). (b) Reverse sequence was performed and confirmed. (c) Adjacent normal lung tissue showed a wild-type sequence for the *erbB2* gene.

within the kinase domain. Totally, we have found only 1 *erbB2* mutation from 122 (0.8%) Japanese NSCLC patients.

Immunohistochemistry

The immunohistochemical evaluation was done according to the scoring system described in Material and Methods. Immunohistochemistry was done only for 87 patients because the tissue blocks were not available for other patients. The *erbB2*-positive (2+/3+) ratio was 26.4% (23/87). There was a significantly higher *erbB2*-positive ratio in *EGFR*-mutant patients (13/25, 52.0%) compared to *EGFR* wild-type patients (10/62, 16.1%) ($p = 0.0247$). The patient with *erbB2* mutation exhibited 1+ immunoreactivity (Fig. 5).

Discussion

We obtained findings that *EGFR* mutation status was significantly correlated with gender and smoking history of lung cancers. This was in agreement with the recent reports that *EGFR* gene mutations are common in lung cancers from never smokers^{13,14} and females with adenocarcinoma.^{11,14} However, our analysis also suggested that *erbB2* mutation might be less common in Japanese NSCLC patients.




FIGURE 5 – Immunohistochemistry for ErbB2. Left: erbB2-positive (3+) section. Right: the NSCLC specimen with *erbB2* mutation exhibited 1+ immunoreactivity.

Overexpression of EGFR/ErbB2 and ErbB ligands is correlated with advanced diseases and poor patient prognosis.¹⁹ Although *EGFR* is more abundantly expressed in lung carcinoma,^{20,21} *erbB2* overexpression is less common; it is found in <35% of patients with non-small cell lung cancers, mainly in those with adenocarcinoma.²¹ Amplification of *EGFR* and *erbB2* mRNA²² or overexpression of their proteins²³ has been found to relate to survival in patients with NSCLC, although contradictory results have also been reported.^{24,25} The drug trastuzumab, a humanized antibody against the extracellular domain of erbB2, has been approved for treatment of metastatic breast cancer and is most effective in breast cancer with *erbB2* amplification. Preliminary results suggested that the combination of chemotherapy and trastuzumab is well tolerated for NSCLC.²¹ However, results from phase II trials of trastuzumab as a treatment for NSCLC have not shown any advantage for most patients²² and have provided insufficient evidence to proceed to phase III trials.²³ Because the presence of a mutation appears to be a determination of response to therapy, as is the case with gefitinib and *EGFR* mutations, we therefore investigated the *erbB2* and *EGFR* gene mutation status. However, we have found only 1 *erbB2* mutation from 122 Japanese lung cancer patients. More recently, Shigematsu *et al.* reported that *erbB2* mutations were found in 3% (8/269) of Japanese NSCLC.²⁶ The single *erbB2* mutation we have found was the same as the one repeatedly found by Shigematsu *et al.*²⁶ Because very few NSCLC patients have gene amplification of *erbB2*, trastuzumab in the treatment of NSCLC might have a limited role.⁹ Lung cancers that coexpress both EGFR and erbB2 appear to have more virulent behavior.²⁷ In addition, EGFR-erbB2 heterodimers are associated with a stronger and more sustained proliferative signal than EGFR homodimers.^{22,28} Blockade of a signaling pathway may in theory be overcome by compensatory activation of a separate pathway in the same tumor cell. Because there was a significantly higher erbB2-positive ratio in *EGFR*-mutant patients, blockade of both may ultimately yield superior results.

Because so many *EGFR* mutation phenotypes were discovered, it would be of interest to determine whether resistance to *EGFR* inhibition emerges through secondary mutation as is the case in imatinib-treated chronic myelogenous leukemia.²⁹ In our analysis, a female never smoker adenocarcinoma patient had the predomi-

nant L858R mutation as well as T790M mutation. Actually, this case was untreated with EGFR kinase inhibitors. Threonine 315 to isoleucine substitution in the Abl kinase domain was a critical structural determinant controlling inhibitor sensitivity of STI571.²⁹ Introduction of bulkier hydrophobic side chains at the Thr-790 position fully preserved the cellular kinase activity of the *EGFR* in the presence of selective kinase inhibitors, indicating potential mechanisms of molecular resistance formation as previously found for BCR-Abl at T315I. Previous *in vitro* study showed that mutation of T790M in the *EGFR* revealed a hotspot for resistance formation against gefitinib,³⁰ also *in vivo*.³¹

Over the decades, the incidence of lung adenocarcinoma has increased worldwide. Most individuals with lung adenocarcinoma (especially women) are nonsmokers,³² who are corresponding with the sensitive population to gefitinib. In Taiwan, *EGFR* mutation ratio from adenocarcinoma was also high (55%, 38 of 69), and all of the adenocarcinomas with *EGFR* mutation were well to moderately differentiated.³³ These data were compatible for our results. Because well-differentiated adenocarcinoma patients had a better prognosis,³⁴ *EGFR* mutant patients showed better prognosis in our univariate analysis. The reason why many mutations were especially found in Asian, female nonsmoker adenocarcinoma remains unknown. Human papilloma virus type 16/18 infections,³⁵ cooking oil fume,³⁶ nutritional status, genetic susceptibility, immunologic infection, tuberculosis and asthma³² have been investigated as causes of lung cancer occurring in nonsmoking women.

The findings of the breakdown of *EGFR* mutations among the 3 exons were interesting. The exon 21 mutations correlated with pathologic stage and subtype, unlike mutations in exon19. Since exon 21 mutations are more closed to the activation loop of *EGFR*, these may be more correlated with gefitinib sensitivity. Especially since 3 patients with exon 20 mutations were smokers, all of the mutations might not be equally correlated with sensitivity for gefitinib.

Acknowledgements

The authors thank Mr. N. Hosono, Mr. R. DeBiasi and Dr. J. Lee for their excellent technical assistance.

References

- Ginsberg RJ, Kris K, Armstrong G. Cancer of the lung. In: Principles and practice of oncology, 4th ed. Philadelphia: Lippincott, 1993:673–82.
- Yasuda K, Ayabe H, Ide H, Uchida Y on behalf of the Japanese Association for Thoracic Surgery. Thoracic and cardiovascular surgery in Japan during 1998. Annual report by the Japanese Association for Thoracic Surgery. *Jpn J Cardiothorac Surg* 1998;48:401–15.
- Postus PE on behalf of the Lung Cancer Cooperative Group of the EORTC. The experience of the Lung Cancer Cooperative Group of

- the European Organization for Research and Treatment of Cancer. *Chest* 1997;113(Suppl):28S-31S.
4. Nicolson RI, Gee JM, Harper ME. EGFR and cancer prognosis. *Eur J Cancer* 2001;37:S9-15.
 5. Onn A, Correa AM, Gilcrease M, Isobe T, Massarelli E, Bucane CD, O'Reilly MS, Hong WK, Fidler IJ, Putnum JB, Herbst RS. Synchronous overexpression of epidermal growth factor receptor and HER2-neu protein is a predictor of poor outcome in patients with stage I non-small cell lung cancer. *Clin Cancer Res* 2004;10:136-43.
 6. Prenzel N, Fischer OM, Streit S, Hart S, Ullrich A. The epidermal growth factor receptor family as a central element for cellular signal transduction and diversification. *Endocr Relat Cancer* 2001;8:11-31.
 7. Mendelson J. Targeting the epidermal growth factor receptor for cancer therapy. *J Clin Oncol* 2002;20:1S-13S.
 8. Slamon DJ, Leyland-Jone B, Shak S, Fuchs H, Paton V, Bajamonde A, Fleming T, Eiermann W, Wolter J, Pegram M, Baselga J, Norton L. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpress HER2. *N Engl J Med* 2001;344:783-92.
 9. Langer CJ, Stephenson P, Thor A, Vangel M, Johnson DH. Trastuzumab in the treatment of advanced non-small-cell lung cancer: is there a role? Focus on Eastern Cooperative Oncology Group study 2598. *J Clin Oncol* 2004;22:1180-7.
 10. Rosell R. Toward customized trastuzumab in HER-2/neu-overexpressing non-small-cell lung cancers. *J Clin Oncol* 2004;22:1171-3.
 11. Paez JG, Janne PA, Lee JC, Tracy S, Greulich H, Gabriel S, Herman P, Kaye FJ, Lindeman N, Boggon TJ, Naoki K, Sasaki H, et al. EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* 2004;304:1499-1504.
 12. Lynch TJ, Bell DW, Sordella R, Gurubhagavatula S, Okimoto RA, Brannigan BW, Harris PL, Haserlat SM, Supko JG, Haluska FG, Louis DN, Christiani DC, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2004;353:1179-1192.
 13. Pao W, Miller V, Zakowski M, Doherty J, Politi K, Sarkaria I, Singh B, Heelan R, Rush V, Fulton L, Mardis E, Kupfer D, et al. EGF receptor gene mutations are common in lung cancers from 'never smokers' and are associated with sensitivity of tumors to gefitinib and erlotinib. *Proc Natl Acad Sci* 2004;101:13306-11.
 14. Shigematsu H, Lin L, Takahashi T, Nomura M, Suzuki M, Witsub II, Fong KM, Lee H, Toyooka S, Shimizu N, Fujisawa T, Feng Z, et al. Clinical and biological features associated with epidermal growth factor receptor gene mutation in lung cancers. *J Natl Cancer Inst* 2005;97:339-46.
 15. Marchetti A, Martella C, Felicioni L, Barassi F, Salvatore S, Chella A, Campese PP, Iarussi T, Mucilli F, Mezzetti A, Cuccurullo F, Sacco R, et al. EGFR mutations in non-small-cell lung cancer: analysis of a large series of cases and development of a rapid and sensitive method for diagnostic screening with potential implications on pharmacologic treatment. *J Clin Oncol* 2005;23:857-65.
 16. Kasaoka T, Yatabe Y, Endoh H, Kuwano H, Takahashi T, Mitsudomi T. Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. *Cancer Res* 2004;64: 8919-23.
 17. Stephens P, Hunter C, Bignell G, Edkins S, Davies H, Teague J, Stevens C, O'Meara S, Smith R, Parker A, Barthorpe A, Blow M, et al. Intragenic erbB2 kinase mutations in tumors. *Nature* 2004;431:525-6.
 18. Japan Lung Cancer Society. General rule for clinical and pathological record of lung cancer, 5th ed. Tokyo: Japan Lung Cancer Society, 1999.1-177.
 19. Pegram M, Slamon D. Biological rationale for HER2/neu (c-erbB2) as a target for monoclonal antibody therapy. *Semin Oncol* 2000;5(Suppl 9):13-9.
 20. Rusch V, Baselga J, Cordon-Cardo C, Orazem J, Zamen M, Hoda S, McIntoch J, Kurie J, Dmitrovsky E. Differential expression of the epidermal growth factor receptor and its ligands in primary non-small cell lung cancers and adjacent benign lung. *Cancer Res* 1993;53: 2379-85.
 21. Hirsch FR, Franklin WA, Veve R, Varella-Garcia M, Bunn PA Jr. HER2/neu expression in malignant lung tumors. *Semin Oncol* 2002; 29:51-8.
 22. Brabender J, Danenberg KD, Metzger R, Schneider PM, Park J, Salonga D, Holscher AH, Danenberg PV. Epidermal growth factor receptor and HER2-neu mRNA expression in non-small cell lung cancer is correlated with survival. *Clin Cancer Res* 2001;7:1850-5.
 23. Tateishi M, Ishida T, Kohdono S, Hamatake M, Fukuyama Y, Sugimachi K. Prognostic influence of the co-expression of epidermal growth factor receptor and c-erbB-2 protein in human lung adenocarcinoma. *Surg Oncol* 1994;3:109-13.
 24. Lai WW, Chen FF, Wu M, Chow NH, Su WC, Ma MC, Su PF, Chen H, Lin MY, Tseng YL. Immunohistochemical analysis of epidermal growth factor receptor family member in stage I non-small-cell lung cancer. *Ann Thorac Surg* 2001;72:1868-76.
 25. Fontanini G, De Laurentis M, Vignati S, Chine S, Lucchi M, Silvestri V, Mussi A, DePlacido S, Tortora G, Bianco AR, Gullick W, Angelletti GA, et al. Evaluation of epidermal growth factor-related growth factor and receptors and of neoangiogenesis in completely resected stage I-IIIa non-small-cell lung cancer: amphiregulin and microvessel count are independent prognostic indicators of survival. *Clin Cancer Res* 1998;4:241-9.
 26. Shigematsu H, Takahashi T, Nomura M, Majumdar K, Suzuki M, Lee H, Wistuba II, Fong KM, Toyooka S, Shimizu N, Fujisawa T, Minna JD, et al. Somatic mutations of the HER2 kinase domain in lung adenocarcinoma. *Cancer Res* 2005;65:1642-6.
 27. Johnson DH, Arteaga CL. Gefitinib in recurrent non-small cell lung cancer: a IDEAL trial? *J Clin Oncol* 2003;21:2227-9.
 28. Yarden Y, Sliwkowski MX. Untangling the erbB signaling network. *Nat Rev Mol Cell Biol* 2001;2:127-37.
 29. Gorre ME, Mohammed M, Ellwood K, Hsu N, Paquette R, Rao PN, Sawyers CL. Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. *Science* 2001;293:876-80.
 30. Blencke S, Ullrich A, Daub H. Mutation of threonine 766 in the epidermal growth factor receptor reveals a hotspot for resistance formation against selective tyrosine kinase inhibitors. *J Biol Chem* 2003; 278:15435-40.
 31. Kobayashi S, Boggon TJ, Dayaram T, Janne PA, Kocher O, Meyerson M, Johnson B, Eck EJ, Tenen DG, Halmos B. EGFR mutation and resistance of non-small-cell lung cancer to gefitinib. *N Engl J Med* 2005;352:786-92.
 32. Chen YC, Chen JH, Richard K, Chen PY, Christiani DC. Lung adenocarcinoma and human papillomavirus infection. *Cancer* 2004;101: 1428-36.
 33. Huang SF, Liu HP, Li LH, Ku Y-C, Fu Y-N, Tsai H-Y, Chen Y-T, Lin Y-F, Chang W-C, Kuo H-P, Wu Y-C, Chen Y-R, et al. High frequency of epidermal growth factor receptor mutations with complex patterns in non-small cell lung cancer related to gefitinib responsiveness in Taiwan. *Clin Cancer Res* 2004;10:8195-203.
 34. Moldvay J, Scheid P, Wild P, Nabil K, Siat J, Borrelly J, Marie B, Farre G, Labib T, Pottier G, Sesboue R, Bronner C, et al. Predictive survival markers in patients with surgically resected non-small cell lung carcinoma. *Clin Cancer Res* 2000;6:1125-34.
 35. Cheng YW, Chiou HL, Sheu GT, Hsieh LL, Chen JT, Chen CY, So JM, Lee H. The association of human papillomavirus 16/18 infection with lung cancer among nonsmoking Taiwanese women. *Cancer Res* 2001;61:2799-803.
 36. Ko YC, Cheng LS, Lee CH, Huang JJ, Huang MS, Kao EL, Wang HZ, Lin HJ. Chinese food cooking and lung cancer in women non-smokers. *Am J Epidemiol* 2000;151:140-7.

Second Primary Cancers in Patients with Stage III Non-Small Cell Lung Cancer Successfully Treated with Chemo-radiotherapy

Tomoya Kawaguchi^{1,2}, Akihide Matsumura^{1,3}, Keiji Iuchi^{1,3}, Seiji Ishikawa^{1,4}, Hajime Maeda^{1,5},
Shimao Fukai^{1,6}, Hikotaro Komatsu^{1,7} and Masaaki Kawahara^{1,2}

¹National Hospital Study Group for Lung Cancer in Japan, ²Department of Internal Medicine, National Hospital Organization Kinki-chuo Chest Medical Center, Sakai, Osaka, ³Department of Surgery, National Hospital Organization Kinki-chuo Chest Medical Center, Sakai, Osaka, ⁴National Hospital Organization Okinawa Hospital, Ginowan, Okinawa, ⁵National Hospital Organization Toneyama Hospital, Toyonaka, Osaka, ⁶National Hospital Organization Ibaragi-Higashi Hospital, Naka-gun, Ibaraki and ⁷National Hospital Organization Matsumoto Hospital, Matsumoto, Nagano, Japan

Received August 11, 2005; accepted October 20, 2005; published online December 20, 2005

Background: Patients successfully treated for non-small cell lung cancer (NSCLC) remain at risk for developing second primary cancer (SPC). The purpose of the current study is to assess the incidence of SPC and the impact of smoking status on the SPC in long-term survivors with stage III NSCLC after chemo-radiotherapy.

Methods: Using the database from the Japan National Hospital Lung Cancer Study Group between 1985 and 1995, information was obtained on 62 patients who were more than 3 years disease-free survivors. Details of clinical information and most smoking history were available from the questionnaire.

Results: Nine of the 62 patients developed SPC 3.9–12.2 years (median, 6.2 years) after the initiation of the treatment. The site of SPC was 2 lung, 1 esophagus, 2 stomach, 1 colon, 1 breast, 1 skin and 1 leukemia. Among these nine, three cancers occurred inside the radiation field. The relative risk of any SPC was 2.8 [95% confidence interval (CI) 1.3–5.3]. The risk changed with the passage of time and it increased significantly (5.2 times at or beyond 7 years) after the treatment. In univariate analysis, the patients who were male, had more cumulative smoking and continued smoking, had an increased risk of SPC [relative risk (RR) 2.7, CI 1.1–5.3; RR 3.0, CI 1.2–6.2; RR 5.2, CI 1.6–11.7, respectively]. In multivariate analysis, factors including smoking status and histological type had no effect on the development of a SPC.

Conclusion: The patients with stage III NSCLC successfully treated with chemo-radiotherapy were at risk for developing SPC and this risk increased with time.

Key words: second primary cancer – non-small cell lung cancer – chemo-radiotherapy

INTRODUCTION

The introduction of combined modality therapy as chest radiotherapy (RT) and chemotherapy for patients with stage III non-small cell lung cancer (NSCLC) has resulted in achieving ~15% long time survivors (123). However, patients successfully treated for NSCLC as well as small cell lung cancer (SCLC) remain at risk for developing second primary cancer (SPC) (4). The risk of SPC in patients with NSCLC has been studied mainly in cohorts of surgically resected patients for stage I NSCLC (567). These reports suggest that the risk of developing SPC and second primary lung cancer (SPLC) is

1–4% and 1–2% per patient per year, respectively, and it appears to increase with the passage of time. Another study including stages I and II patients treated with chest RT confirmed a similar trend that the risk of developing SPC and SPLC is 4.3 and 1.4% per patient per year, respectively (8). Unlike the studies of the patients with SCLC (9–11), these did not provide adequate follow-up information to determine relative risk. Also, there has been no report to date to evaluate the risk of SPC associated with the treatment of RT with chemotherapy as well as smoking status in stage III NSCLC patients.

PATIENTS AND METHODS

Information was obtained on 1643 patients with stage III NSCLC between 1985 and 1995, using the database from the National Hospital Study Group for Lung Cancer, including

For reprints and all correspondence: Tomoya Kawaguchi, Department of Internal Medicine, National Hospital Organization Kinki-chuo Chest Medical Center 1180 Nagasone-cho, Sakai, Osaka 591-8555, Japan. E-mail address: t-kawaguchi@kch.hosp.go.jp

National Hospital Organization Kinki-chuo Chest Medical Center, National Hospital Organization Toneyama Hospital and National Hospital Organization Okinawa Hospital. Among them, 547 patients were treated with chemo-radiotherapy with or without surgery. Of the 547, the 62 patients were more than 3 years disease-free survivors. The patients who relapsed within the 3 years were excluded in this study. Details of clinical information after the treatment and smoking history of the patients were obtained by a questionnaire, which was completed by directly interviewing the patients or the relatives of deceased patients, or by checking the patient's medical records.

Smoking cessation was defined as completely stopping smoking within 6 months after initiation of treatment. Smoking-related cancers include cancer of the lung, larynx and oral cavity, including pharynx, esophagus, pancreas, bladder, kidney, stomach and uterine cervix. A second primary lung cancer was diagnosed according to the criteria provided by Martini and Melamed in 1975 (12). The period of the study was taken as starting from the first day of therapy, and the date of second cancer was taken as the day of histological or cytological documentation of cancer.

For estimation of the expected values of SPC development, the period of risk began 3 years after initiation of treatment and ended with the date of death, date of last follow-up or date of diagnosis of a SPC, whichever occurred first. Age, gender and period-specific rates for cancer incidence within the period 1985–98 obtained from the Research Group for Population-based Cancer Registration in Japan were applied to the appropriate person-years of observation (13). Statistical methods for risk estimation were based on the assumption that observed number of second cancers followed a Poisson distribution (14). To calculate excess risks per 10 000 patients per year in subgroups with significant relative risks, the expected number of cases was subtracted from the number observed. The difference was divided by person-years of observation, and multiplied by 10 000. The risk of a SPC with a specific exposure as smoking was estimated by comparing the patients without the specific exposure, using Poisson regression methods adjusting for gender, histology (squamous cell carcinoma versus non-squamous cell carcinoma) and cumulative smoking amount before the treatment of NSCLC (40 pack-years > versus ≥ 40 pack-years) (15).

RESULTS

The 62 questionnaires completed for each patient showed that none of the patients had past history of cancer of any site nor received previous chemotherapy or RT. The patient characteristics are summarized in Table 1. The end of observation to count the person-years was 31 December 1998. The median follow-up from initiation of therapy was 6.2 years (range 3.1–12.2 years). Of the 62 patients, nine developed SPC in 435 person-years of follow-up. Forty-six patients have remained free of cancer since initial treatment. Three other patients relapsed with NSCLC and still remain alive

Table 1. Patient characteristics ($n = 62$)

Gender	
Male	50
Female	12
Age (median, range)	61, 34–80
Histology	
Squamous cell carcinoma	30
Adenocarcinoma	21
Large cell carcinoma	10
Adenosquamous carcinoma	1
Stage	
IIIA	32
IIIB	30
Surgery	
Yes	24
No	38
Smoking (median, range)	40 pack-years, 0–120
Stop smoking	
Yes	29
No	16
Unknown	17

receiving second line chemotherapy. Of the 62 patients, 13 have died: 5 from recurrent NSCLC, 4 from SPC, 4 from other causes. Regarding chemotherapy for initial treatment, 39 patients were treated with cisplatin (CDDP) + mitomycin (MMC) + vindesine (VDS), 16 with CDDP + VDS, 4 with carboplatin, 2 with CDDP + irinotecan, with 1 with CDDP + MMC + inorelbine. In the treatment of RT, 66 Gy were given to 5 patients, 60 Gy to 10, 56 Gy to 28, 50 Gy to 15 and 40 Gy to 4. Of the 62 patients, surgery was performed in 24 patients after the chemo-radiotherapy.

For smoking status, information was obtained for all the 62 patients before the treatment, but was available for 45 patients after the treatment. Of the 45 patients treated in the analysis, 16 patients continue to smoke and 19 patients stopped smoking. For assessment, 10 never smokers were also added to the 19 stopped patients, and the 29 patients were categorized to the stop smoking group.

Details of nine patients who developed SPC out of the 62 patients are shown in Table 2. There has been no SPC among the ten never smokers. Two patients (cases 5 and 9) developed a SPC in different lobes from the original NSCLC. Both tumors arose from the ipsilateral side and both patients continued to smoke after the treatment. One of the two lung cancers developed inside the radiation field. The other malignancies consisted of carcinoma of the esophagus, stomach, colon, skin, breast and acute myelogenous leukemia. Two SPC with skin and breast cancer (cases 6 and 8) also developed inside the radiation field.

Table 3 shows the relative and absolute risks of SPC after initiation of therapy for NSCLC. The risk for development of any SPC increased significantly to 2.8 [95% confidence interval (CI) 1.3–5.3]. In spite of the overall increase in risk there was no significant increase in relative risk of developing a particular cancer. When smoking-related cancers are combined, there was still no significant increased relative risk in the development of SPC.

Table 2. Characteristics of nine patients with second primary cancers

Patient	Age	Gender	CFI (years)	P	His	SPT/His
1	70	M	3.9		LA	Stomach/AD
2	69	M	11.5		AD	Colon/AD
3	61	M	6.3		SQ	Esophagus/SQ
4	65	M	4.5		SQ	Stomach/AD
5	62	M	5.6		SQ	Lung/SQ
6	58	M	4.5		AD	Skin/SQ inside RT field
7	66	M	8.1		SQ	AML
8	54	F	10.4		LA	Breast/AD inside RT field
9	66	M	7.9		AD, SQ	Lung/Undiff inside RT field

CFI, cancer-free interval; P, Primary; His, Histology; AD, adenocarcinoma; LA, large cell carcinoma; SQ, squamous cell carcinoma; Undiff, undifferentiated carcinoma; AML, Acute myeloid leukemia; RT, radiotherapy.

Table 3. Risk of second primary cancers

Site	Obs	E	O/E	95% CI	Absolute risk*
All cancers	9	3.23	2.8	1.3-5.3	238.9
Esophagus	1	0.12	8.6	0.1-47.7	
Stomach	2	0.81	2.5	0.3-8.9	
Colon	1	0.39	2.5	0.1-14.1	
Lung	2	0.50	4.0	0.4-7.2	
Skin	1	0.03	36.2	0.4-201.3	
Breast	1	0.03	36.7	0.4-204.1	
Leukemia	1	0.03	30.9	0.4-171.5	
Smoking-related	5	1.81	2.8	0.9-6.4	

Obs, observed; E, expected.
*Excess risk per 10 000 persons per year.

Next, the effect of the passage of time was evaluated. The relative risk for 3-4 years after the treatment was 2.2 (95% CI 0.1-23.9) and 1.8 (95% CI 0.1-23.9) for 5-6 years, and 5.2 (95% CI 1.4-13.2) for at or beyond 7 years. The risk changed with the passage of time and it increased significantly (5.2 times at or beyond 7 years) after the treatment. The absolute risk was 600.1 per 10 000 persons per years.

Table 4 shows the results of univariate analysis on the relative risk for a SPC. The risk was significant but modestly increased relative to the general population in male and more cumulative smoking amount (2.7 times; 95% CI 1.1-5.3 and 3 times; 95% CI 1.2-6.2, respectively). Among those who continued to smoke, there was a significantly increased relative risk (5.2 times; 95% CI 1.6-11.7). In contrast, those who stopped smoking showed only a 1.8-fold increase (95% CI 0.3-5.9), which was not significantly different from the general population.

Finally, we assessed multivariate analysis and examined the relationship between continued smoking habits and the risk of a SPC, adjusted for gender, histology type and

Table 4. Risk of second primary cancers by histology, gender and smoking status

	Obs	O/E	95% CI	Absolute risk*
Histology				
SQ	4	2.7	0.7-6.9	
Non-SQ	5	2.6	0.9-6.7	
Gender				
Male	8	2.7	1.1-5.3	246.7
Female	1	4.3	0.1-23.9	
Surgery				
Yes	4	3.6	0.9-9.2	
No	5	2.3	0.7-5.4	
Smoking				
≤40 pack-years	2	2.2	0.2-8.0	
≥40 pack-years	7	3.0	1.2-6.2	324.2
Intercurrent smoking				
Yes	3	1.8	0.3-5.9	
No	5	5.2	1.6-11.7	430.5

SQ, squamous cell carcinoma; Obs, observed.
*Excess risk per 10 000 persons per year.

Table 5. Relative risk of second primary cancers estimated by multivariate analysis

Risk factor	Relative risk	95% CI
Cumulative smoking (<40 pack-years/≥40 pack-years)	1.4	0.2-8.4
Intercurrent smoking (yes/no)	2.3	0.5-10.8
Histology (SQ/non-SQ)	3.3	0.2-3.3
Gender (male/female)	1.0	0.1-11.2

SQ, squamous cell carcinoma.

cumulative smoking amount. The results are shown in Table 5. We could not demonstrate that factors such as continued smoking habits, gender, histology type and cumulative smoking amount had effect on the development of a SPC.

DISCUSSION

There has been a large body of work that evaluated the risk of SPC in the patients with NSCLC in the treatment of surgery or RT alone (5678). Although the number of survivors in patients with stage III NSCLC has increased by combined modality therapy as chemotherapy and RT, there has been no report to date to evaluate the risk of SPC in these patients. Additionally, Ng and co-workers (16) reported that the relative risk of SPC was 6.1 with the combined chemotherapy and RT and 4.0 with the RT alone, showing a significant difference ($P = 0.03$) in the surviving patients in Hodgkin's disease. Given that, we focused on the NSCLC patients treated with chemo-radiotherapy.

In our study, 9 patients out of 62 long-term survivors of stage III NSCLC treated with chemo-radiotherapy had a SPC. The relative risk for any SPC (2.8; 95% CI 1.3–5.3) compared with the general population was significantly increased. Instead of many reports examining the risk, these do not provide adequate follow-up information to determine relative risk in the patients with NSCLC. Most studies only show a percent risk per patient per year (5–8). In the current study, the overall rate of developing SPC is estimated at 2.9% per patient per year, which is in agreement with the rates in most surgical series. Ginsberg and Rubinstein (5) reported that SPC occurrence rate was 1.7% per patient per year on 247 patients operated for T1 N0 NSCLC. Other studies showed the rate of 2.8% by Martini et al. (6) and 2.4–3.6% by Thomas and Rubinstein (7). In the current study, we also confirmed the effect of the passage of time on developing SPC. Thomas and Rubinstein (7) reported that the rate of SPC increased from 2.4% for the first 5 years after surgical resection to 3.6% after the fifth year.

We previously studied the relative risk of SPC in the SCLC patient successfully treated with chemotherapy with or without RT (9). Our results showed a similar trend as previous studies (10,11) and demonstrated that the patient had a significantly increased relative risk of 3.6 (95% CI 2.0–5.9) and that the patients who continued to smoke demonstrated a significantly increased risk for a SPC (4.3, 95% CI 1.1–15.9, $P = 0.03$) compared with those who stopped smoking.

Unlike the results of SCLC patients study, the risk of SPC in NSCLC patients was lower, and the impact of continued smoking on developing SPC in the patients was less significant, but the reason for this observation is not completely understood. According to the case-control study from Japan (17), lung cancer risk reduction due to smoking cessation appeared to be greater in SCLC than squamous cell carcinoma or adenocarcinoma, and SCLC seems to be more smoking-related than NSCLC. However, there have been a couple of germline polymorphism as cytochrome P 450 1A1 (CYP1A1) and glutathione S-transferase class mu (GSTM1), reported, which is implicated in smoking-related carcinogenesis (18,19). Therefore, SCLC patients are speculated to have a higher potential to develop a SPC, particularly smoking-related cancers.

Among NSCLC patients, there seems to be a special group of roentgenographically occult early stage squamous cell carcinoma of the lung. In this patient group, the rate of occurrence of SPC, particularly SPLC was estimated at 3–4% per patient per year (20,21). The risk for SPLC seemed to be substantially higher than that of 1–2% in the NSCLC patients treated with surgery or RT from the previous study and treated with chemo-radiotherapy from our study. Therefore, the group should be given a special focus and be divided from the general population of NSCLC patients in the research of risk of SPC. Most of the patients can be cured by surgery, photodynamic therapy, brachytherapy and chest RT because of its early clinical stage (22), and are not included in our study. Roentgenographically occult early stage squamous cell carcinoma of the lung is associated with the concept of

field cancerization (23), and smoking status seems to be very important to evaluate the risk of SPC, which awaits further examination.

A relatively small sample size and rare events such as SPC in this study resulted in large confidence intervals for the estimates. It is still difficult to conclude the effect of continued smoking on the development of SPC. Cigarette smoking causes not only developing cancers but also cardiovascular and lung damage as well (24,25). It may be speculated that continued smokers died off early when interpreting the results. The cessation of smoking is still warranted among patients with stage III NSCLC treated by chemo-radiotherapy.

In conclusion, stage III NSCLC patients treated with chemo-radiotherapy were at risk of developing SPC and this risk increased with time. A large sample size study in a longer follow-up period may be required in further research to conclude the effect of continued smoking on the development of SPC. SPC in another particular group such as roentgenographically occult early stage squamous cell carcinoma of bronchus also awaits further studies.

Acknowledgments

We thank Mrs Chihiro Horii, Mr Toshiyuki Ijima and Dr Satoshi Teramukai for their statistical assistance; Dr Mitsumasa Ogawara, Dr Toshi Hashizume and Dr Yuka Fujita for their support of the study; Dr Minoru Takada for his comments on the manuscript. This work was supported in part by a Grand-in-Aid for Cancer Research from the Ministry of Health and Welfare, Japan.

References

- Ohe Y, Ishizuka N, Tamura T, Sekine I, Nishiwaki Y, Saijo N. Japan Clinical Oncology Group. Long-term follow-up of patients with unresectable locally advanced non-small cell lung cancer treated with chemoradiotherapy: a retrospective analysis of the data from the Japan Clinical Oncology Group trials (JCOG0003A). *Cancer Sci* 2003;94:729–34.
- Komaki R, Seiferheld W, Ettinger D, Lee JS, Movsas B, Sause W. Randomized phase II chemotherapy and radiotherapy trial for patients with locally advanced inoperable non-small-cell lung cancer: long-term follow-up of RTOG 92-04. *Int J Radiat Oncol Biol Phys* 2002;53:548–57.
- Jeremic B, Shibamoto Y, Acimovic L, Milicic B, Milisavljevic S, Nikolic N, et al. Hyperfractionated radiation therapy and concurrent low-dose, daily carboplatin/etoposide with or without weekend carboplatin/etoposide chemotherapy in stage III non-small-cell lung cancer: a randomized trial. *Int J Radiat Oncol Biol Phys* 2001;50:19–25.
- Johnson BE. Second lung cancers in patients after treatment for an initial lung cancer. *J Natl Cancer Inst* 1998;90:1335–45.
- Ginsberg RJ, Rubinstein LV. Randomized trial of lobectomy versus limited resection for T1 N0 non-small cell lung cancer. Lung Cancer Study Group. *Ann Thorac Surg* 1995;60:615–22.
- Martini N, Bains MS, Burt ME, Zakowski MF, McCormack P, Rusch VW, et al. Incidence of local recurrence and second primary tumors in resected stage I lung cancer. *J Thorac Cardiovasc Surg* 1995;109:120–9.
- Thomas PA Jr, Rubinstein L. Malignant disease appearing late after operation for T1 N0 non-small-cell lung cancer. *J Thorac Cardiovasc Surg* 1993;106:1053–8.
- Jeremic B, Shibamoto Y, Acimovic L, Nikolic N, Dagovic A, Aleksandrovic J, et al. Second cancers occurring in patients with early stage non-small-cell lung cancer treated with chest radiation therapy alone. *J Clin Oncol* 2001;19:1056–63.

9. Kawahara M, Ushijima S, Kamimori T, Kodama N, Ogawara M, Matsui K, et al. Second primary tumours in more than 2-year disease-free survivors of small-cell lung cancer in Japan: the role of smoking cessation. *Br J Cancer* 1998;78:409-12.
10. Tucker MA, Murray N, Shaw EG, Ettinger DS, Mabry M, Huber MH, et al. Second primary cancers related to smoking and treatment of small-cell lung cancer. *J Natl Cancer Inst* 1997;89:1782-8.
11. Richardson GE, Tucker MA, Venzon DJ, Linnoila RI, Phelps R, Phares JC, et al. Smoking cessation after successful treatment of small-cell lung cancer is associated with fewer smoking-related second primary cancers. *Ann Intern Med* 1993;119:383-90.
12. Martini N, Melamed MR: Multiple primary lung cancers. *J Thorac Cardiovasc Surg* 1975;70:606-12.
13. The Research Group for Population-based Cancer Registration in Japan. Cancer incidence and incidence rates in Japan in 1998: estimates based on data from 12 population-based cancer registries. *Jpn J Clin Oncol* 2003;33:241-5.
14. Boice J, Lubin J, Preston D. Epidemiologic analysis with a personal computer (EPITOME). NIH Publication (91-380) 1991.
15. SAS Institute SAS/STAT User's Guide, Version 6, Vol. 2, 4th edn. Cary, NC: SAS Institute 1989;1070-26.
16. Ng AK, Bernardo MV, Weller E, Backstrand K, Silver B, Mauch PM, et al. Second malignancy after Hodgkin disease treated with radiation therapy with or without chemotherapy: long-term risks and risk factors. *Blood* 2002;100:1989-96.
17. Sobue T, Suzuki T, Fujimoto I, Matsuda M, Doi O, Mori T, et al. Lung cancer risk among exsmokers. *Jpn J Cancer Res* 1991;82:273-279.
18. To-Figueras J, Gene M, Gomez-Catalan J, Galan MC, Fuentes M, Ramon JM, et al. Glutathione S-transferase M1 (GSTM1) and T1 (GSTT1) polymorphisms and lung cancer risk among Northwestern Mediterraneans. *Carcinogenesis* 1997;18:1529-33.
19. Kihara M, Kihara M, Noda K. Risk of smoking for squamous and small cell carcinomas of the lung modulated by combinations of CYP1A1 and GSTM1 gene polymorphisms in a Japanese population. *Carcinogenesis* 1995;16:2331-6.
20. Woolner LB, Fontana RS, Cortese DA, Sanderson DR, Bernatz PE, Payne WS, et al. Roentgenographically occult lung cancer: pathologic findings and frequency of multicentricity during a 10 year period. *Mayo Clin Proc* 1984;59:453-66.
21. Saito Y, Nagamoto N, Ota S, Sato M, Sagawa M, Kamma K, et al. Results of surgical treatment for roentgenographically occult bronchogenic squamous cell carcinoma. *J Thorac Cardiovasc Surg* 1992;104:401-7.
22. Kawaguchi T, Yamamoto S, Naka N, Okishio K, Atagi S, Ogawara M, et al. Immunohistochemical analysis of bcl-2 protein in centrally located early stage lung cancer treated with photodynamic therapy. *Br J Cancer* 2000;82:418-23.
23. Braakhuis BJ, Tabor MP, Kummer JA, Leemans CR, Brakenhoff RH. A genetic explanation of Slaughter's concept of field cancerization: evidence and clinical implications. *Cancer Res* 2003;63:1727-30.
24. Howard G, Wagenknecht LE, Burke GL, Diez-Roux A, Evans GW, McGovern P, et al. Cigarette smoking and progression of atherosclerosis: The Atherosclerosis Risk in Communities (ARIC) Study. *J Am Med Assoc* 1998;279:119-24.
25. Fujisawa T, Iizasa T, Saitoh Y, Sekine Y, Motohashi S, Yasukawa T, et al. Smoking before surgery predicts poor long-term survival in patients with stage I non-small-cell lung carcinomas. *J Clin Oncol* 1999;17:2086-91.

Standard Thoracic Radiotherapy With or Without Concurrent Daily Low-dose Carboplatin in Elderly Patients with Locally Advanced Non-small Cell Lung Cancer: a Phase III Trial of the Japan Clinical Oncology Group (JCOG9812)

Shinji Atagi¹, Masaaki Kawahara¹, Tomohide Tamura², Kazumasa Noda³, Koshiro Watanabe⁴, Akira Yokoyama⁵, Takahiko Sugiura⁶, Hiroshi Senba⁷, Satoshi Ishikura⁸, Hiroshi Ikeda², Naoki Ishizuka⁹ and Nagahiro Saijo²

¹National Hospital Organization Kinki-Chuo Chest Medical Center, Osaka, ²National Cancer Center Hospital, Tokyo, ³Kanagawa Cancer Center, Yokohama, ⁴Yokohama Municipal Citizen's Hospital, Yokohama, ⁵Niigata Cancer Center Hospital, Niigata, ⁶Aichi Cancer Center, Nagoya, ⁷Kumamoto Regional General Hospital, Kumamoto, ⁸National Cancer Center Hospital East, Kashiwa, Chiba and ⁹The Statistics and Cancer Control Division, National Cancer Center, Tokyo, Japan

Received October 20, 2004; accepted February 14, 2005

Background: The purpose of this study was to evaluate whether radiotherapy with carboplatin would result in longer survival than radiotherapy alone in elderly patients with unresectable stage III non-small cell lung cancer (NSCLC).

Methods: Eligible patients were 71 years of age or older with unresectable stage III NSCLC. Patients were randomly assigned to the radiotherapy alone (RT) arm, irradiation with 60 Gy; or the chemoradiotherapy (CRT) arm, the same radiotherapy and additional concurrent use of carboplatin 30 mg/m² per fraction up to the first 20 fractions.

Results: This study was terminated early when 46 patients were registered from November 1999 to February 2001. Four patients (one in the RT arm, three in the CRT arm) were considered to have died due to treatment-related causes. The JCOG Radiotherapy Committee assessed these treatment-related deaths (TRDs) and the compliance with radiotherapy in this trial. They found that 60% of the cases corresponded to protocol deviation and 7% were protocol violation in dose constraint to the normal lung, two of whom died due to radiation pneumonitis. As to the effectiveness for the 46 patients enrolled, the median survival time was 428 days [95% confidence interval (CI) = 212–680 days] in the RT arm versus 554 days (95% CI = 331 to not estimable) in the CRT arm.

Conclusions: Due to the early termination of this study, the effectiveness of concurrent use of carboplatin remains unclear. We re-planned and started a study with an active quality control program which was developed by the JCOG Radiotherapy Committee.

Key words: non-small cell lung cancer – elderly patients – carboplatin – chemoradiotherapy

INTRODUCTION

Lung cancer is the leading cause of cancer-related deaths in the USA, Europe and Japan. In Japan, the number of elderly is increasing dramatically. In 2001, the proportion of Japanese population older than 65 years was 18%; in other words, the number of people older than 65 years exceeded 22 million (1). Lung cancer death rates for men and women aged 75 or more have increased to ~531 and 138 per 100 000 population, respectively (1). To establish the effective treatment for

the elderly with lung cancer has thus become of greater importance.

Until recently, the standard treatment for locally advanced non-small cell lung cancer (NSCLC) was radiotherapy alone. However, the 5-year survival rate of patients with stage III remained under 10% (2–4). To improve the survival rates, many clinical trials comparing radiotherapy with chemoradiotherapy have been conducted (5–11). A recent meta-analysis suggested that the combination of chemotherapy containing cisplatin (CDDP) and radiation could improve the survival rate compared with radiotherapy alone (12,13). However, it is still unclear whether the combined chemoradiotherapy is also suitable for elderly patients. This is partly because the elderly had been considered inappropriate as study patients.

For reprints and all correspondence: Shinji Atagi, National Hospital Organization Kinki-Chuo Chest Medical Center, 1180 Nagasone, Sakai, Osaka. 591-8555, Japan. E-mail: s-atagi@kch.hosp.go.jp