

recommended as an initial treatment for MCHM when CV in tumour volume is above 1.8. Those patients should be treated by systemic chemotherapy and surgical resection; then should be considered when the disease responds to the therapy.

Selecting the appropriate strategy according to the tumour staging has been an issue for patients with MCHM. Several studies proposed clinical risk scores incorporating the prognostic factors for predicting recurrence or prognosis after resection of MCHM to answer the clinical question.^{4,8,11,23,24} However, those scoring systems and the factors consisting of each scoring system are not identical among the studies. The standard algorithm of treatment for MCHM is still obscure. Definitive and universal factors or a scoring system for predicting recurrence and prognosis after resection of MCHM is needed, and CV might be one of those factors in cases of MCHM.

The multistep process involving mutational events on both oncogenes and tumour suppressor genes is accepted for development of colorectal cancer, and several studies have shown that allelic imbalance correlated with staging and prognosis in colorectal cancer.^{25,26} High CV might be caused by some genetic alterations. Genetic analysis such as evaluations of difference between the allelic imbalance of small tumours and that of large tumours in patients with high CV or evaluations of difference between the allelic imbalance of tumours in patients with low CV and that of tumours in patients with high CV may indicate particularly sensitive genomic regions and offer new information about the development of colorectal cancer. Further study is warranted to verify the prognostic significance of CV in tumour volume.

In conclusion, hepatic resections for MCHM sometimes contribute to long-term survival. Coefficient of variation in tumour volume above 1.8 might predict poor survival after hepatectomy of MCHM and be useful in planning the therapeutic strategy for patients with MCHM.

Acknowledgement

This work was supported in part by grants from Ministry of Health, Labor and Welfare, Japan.

References

- Fortner JG, Silva JS, Golbey RB, Cox EB, Maclean BJ. Multivariate analysis of a personal series of 247 consecutive patients with liver metastases from colorectal cancer. I. Treatment by hepatic resection. *Ann Surg* 1984;199:306–16.
- Steele Jr G, Bleday R, Mayer RJ, Lindblad A, Petrelli N, Weaver D. A prospective evaluation of hepatic resection for colorectal carcinoma metastases to the liver: Gastrointestinal Tumor Study Group Protocol 6584. *J Clin Oncol* 1991;9:1105–12.
- Jamison RL, Donohue JH, Nagorney DM, Rosen CB, Harmsen WS, Ilstrup DM. Hepatic resection for metastatic colorectal cancer results in cure for some patients. *Arch Surg* 1997;132:505–10.
- Fong Y, Fortner J, Sun RL, Brennan MF, Blumgart LH. Clinical score for predicting recurrence after hepatic resection for metastatic colorectal cancer: analysis of 1001 consecutive cases. *Ann Surg* 1999;230:309–18.
- Choti MA, Sitzmann JV, Tiburi MF, Sumetchotimetha W, Rangsin R, Schulick RD, et al. Trends in long-term survival following liver resection for hepatic colorectal metastases. *Ann Surg* 2002;235:759–66.
- Bolton JS, Fuhrman GM. Survival after resection of multiple bilobar hepatic metastases from colorectal carcinoma. *Ann Surg* 2000;231:743–51.
- Scheele J, Stangl R, Altendorf-Hofmann A. Hepatic metastases from colorectal carcinoma: impact of surgical resection on the natural history. *Br J Surg* 1990;77:1241–6.
- Nordlinger B, Guiguet M, Vaillant JC, et al. Surgical resection of colorectal carcinoma metastases to the liver. A prognostic scoring system to improve case selection, based on 1568 patients. Association Française de Chirurgie. *Cancer* 1996;77:1254–62.
- Nakamura S, Suzuki S, Baba S. Resection of liver metastases of colorectal carcinoma. *World J Surg* 1997;21:741–7.
- Cady B, Jenkins RL, Steele Jr GD, et al. Surgical margin in hepatic resection for colorectal metastasis: a critical and improvable determinant of outcome. *Ann Surg* 1998;227:566–71.
- Iwatsuki S, Dvorchik I, Madariaga JR, et al. Hepatic resection for metastatic colorectal adenocarcinoma: a proposal of a prognostic scoring system. *J Am Coll Surg* 1999;189:291–9.
- Bradley AL, Chapman WC, Wright JK, et al. Surgical experience with hepatic colorectal metastasis. *Am Surg* 1999;65:560–6.
- Minagawa M, Makuuchi M, Torzilli G, et al. Extension of the frontiers of surgical indications in the treatment of liver metastases from colorectal cancer: long-term results. *Ann Surg* 2000;231:487–99.
- Resection of the liver for colorectal carcinoma metastases: a multi-institutional study of indications for resection. Registry of hepatic metastases. *Surgery* 1988;103:278–88.
- Jass JR, Sobin LH. Histological typing of intestinal tumors. In: Jass JR, Sobin LH, editors. *World Health Organization. International histological classification of tumors*. 2nd ed. Berlin: Springer-Verlag; 1989.
- Kaplan EL, Meier P. Non-parametric estimation from incomplete observations. *J Am Stat Assoc* 1958;53:457–81.
- Cox DR. Regression models and life tables (with discussion). *J R Stat Soc B* 1972;34:187–220.
- Couinaud C. Bases anatomiques des hepatectomies gauche et droite réglées. *J Chirurgie* 1954;70:933–66.
- Weber SM, Jarnagin WR, DeMatteo RP, Blumgart LH, Fong Y. Survival after resection of multiple hepatic colorectal metastases. *Ann Surg Oncol* 2000;7:643–50.
- Ostle B. *Statistics in research basic concepts and techniques for research workers*. 1st ed. Ames: Iowa State College Press; 1954.
- Steel RGD, Torrie JH, Dickey DA. *Principles and procedures of statistics, a biometrical approach*. 3rd ed. New York: McGrawHill Book Co.; 1997.
- Kato T, Yasui K, Hirai T, et al. Therapeutic results for hepatic metastasis of colorectal cancer with special reference to effectiveness of hepatectomy: analysis of prognostic factors for 763 cases recorded at 18 institutions. *Dis Colon Rectum* 2003;46:S22–31.
- Nagashima I, Takada T, Matsuda K, et al. A new scoring system to classify patients with colorectal liver metastases: proposal of criteria to select candidates for hepatic resection. *J Hepatobiliary Pancreat Surg* 2004;11:79–83.
- Ueno H, Mochizuki H, Hatsuse K, Hase K, Yamamoto T. Indicators for treatment strategies of colorectal liver metastases. *Ann Surg* 2000;231:59–66.
- Weber JC, Schneider A, Rohr S, et al. Analysis of allelic imbalance in patients with colorectal cancer according to stage and presence of synchronous liver metastases. *Ann Surg* 2001;234:795–802. discussion 802–3.
- Beau-Faller M, Weber JC, Schneider A, et al. Genetic heterogeneity in lung and colorectal carcinoma as revealed by microsatellite analysis in plasma or tumor tissue DNA. *Cancer* 2003;97:2308–17.

Positron Emission Tomography with F-18 Fluorodeoxyglucose in Evaluating Colorectal Hepatic Metastasis Down-staged by Chemotherapy

SHINICHIRO TAKAHASHI¹, YOSHIHUMI KUROKI², KATSUHIRO NASU², SHIGERU NAWANO², MASARU KONISHI¹, TOSHIO NAKAGOHRI¹, NAOTO GOTOHDA¹, NORIO SAITO³ and TAIRA KINOSHITA¹

¹Department of Hepato-biliary Pancreatic Surgery, ²Department of Diagnostic Radiology and ³Department of Colorectal Surgery, National Cancer Center Hospital East, Chiba, Japan

Abstract. *Background:* The efficacy of positron emission tomography with ¹⁸F fluoro-2-deoxy-D-glucose (FDG-PET) is obscure in evaluating viability or the extent of colorectal hepatic metastasis (CHM), down-staged by chemotherapy. *Patients and Methods:* A retrospective lesion-by-lesion analysis was performed for seven consecutive patients, who had received rescue hepatectomy for initially unresectable CHM, in order to evaluate the correlation between results of imaging modalities and the corresponding pathology. *Results:* The sensitivity and positive predictive value of the conventional modalities (CT and MRI) were 92% and 42%, respectively, while the sensitivity, specificity, positive predictive value and negative predictive value of FDG-PET were 58%, 100%, 100% and 75% respectively. The sensitivity of FDG-PET was 100% in evaluating the viability of tumors >2 cm, however, this fell to 17% in tumors <2 cm. *Conclusion:* FDG-PET is effective in assessing the viability of tumors >2 cm, but not those <2 cm, in patients with CHM down-staged by chemotherapy.

Unresectable hepatic metastasis is one of the major obstacles in the treatment of colorectal cancer. Systemic chemotherapy for unresectable colorectal cancer has improved recently with the introduction of new effective agents (1-4). Even so, chemotherapy is rarely curative. Rescue hepatectomy after down-staging by chemotherapy is a potential treatment for unresectable colorectal hepatic metastasis (CHM) (5-8). However, pre-operative evaluation of tumor extent or

viability is one of the issues in this strategy. The indication for rescue hepatectomy is based on the interpretation of the imaging of tumor extent or viability after down-staging by chemotherapy, however, evaluating the viability of hepatic tumors treated with chemotherapy is sometimes difficult by conventional imaging methods, such as computed tomography (CT) or magnetic resonance imaging (MRI), especially when the tumors have shrunk remarkably or exhibit notable calcification, after chemotherapy. Therefore, an effective diagnostic imaging modality to evaluate the viability of hepatic tumors after chemotherapy is needed.

Positron emission tomography with ¹⁸F fluoro-2-deoxy-D-glucose (FDG-PET) is a functional imaging technique based on the increased utilization of glucose by tumor cells, and is effective for staging colorectal cancer (9, 10). Furthermore, the ability of FDG-PET to assess the pathological response to chemotherapy has been suggested in various malignant tumors (11-15). The correlation in CHM between FDG-PET findings and those of corresponding pathology has not been fully examined.

The present study was conducted to examine whether FDG-PET was able to assess tumor viability, before rescue surgery, for initially unresectable CHMs down-staged by chemotherapy and indicate which tumors should be resected.

Patients and Methods

Patient population. Seventy-four patients underwent hepatic resection for CHM at the National Cancer Center Hospital East, Japan, between January 2004 and July 2005, and 10 out of these underwent hepatic resection, after down-staging by chemotherapy. Since January 2004, all patients about to undergo rescue surgery are examined using FDG-PET pre-operatively, when informed consent is granted. Consequently, seven consecutive patients, who had been examined by CT, MRI and FDG-PET, were included in the present study. The patients were four men and three women, ranging from 44 to 70 years old. The location of the primary colorectal tumor was the colon in five patients and the rectum in two patients. All the

Correspondence to: Shinichiro Takahashi, MD, Department of Hepato-biliary Pancreatic Surgery, National Cancer Center Hospital East, 6-5-1 Kashiwanoha, Kashiwa 277-8577, Chiba, Japan. Tel: +81-471-33-1111, Fax: +81-471-31-4724, e-mail: shtakaha@east.ncc.go.jp

Key Words: Colorectal cancer, hepatic metastasis, chemotherapy, hepatic resection, FDG-PET.

primary tumors were well- or moderately-differentiated adenocarcinomas. The primary tumors were staged as II (n=2), III (n=2), and IV (n=3) according to the TNM classification.

Chemotherapy. The reasons for choosing chemotherapy instead of curative resection, initially, were multiple bilobar tumors in five patients, invasion to the bilateral bile ducts and portal veins in one and invasion to the 3 major hepatic veins in one. The chemotherapies performed for the unresectable tumors were 5-fluorouracil-leucovorin combined with irinotecan in four patients, 5-fluorouracil-leucovorin alone in one, oral uracil/tegafur in one and capecitabine in one. Five patients experienced a partial response and two patients had stable disease, based on the Response Evaluation Criteria in Solid Tumors.

Conventional imaging (CT, MRI). All patients underwent contrast-enhanced CT and MRI before hepatectomy. Multi slice CT with 16 DAS was used for this study (Aquilion, Toshiba Medical Systems, Japan). CT images were obtained using 5 mm collimation after administration of 100 ml of nonionic iodine intravenous contrast medium injected at 3 ml/sec with a 70-sec delay (portal-dominant phase). Images were reconstructed at 5 mm intervals using a standard soft-tissue algorithm.

MR images were acquired using a 1.5-T MR imager (Gyroscom Intera, Philips Medical Systems, Netherlands) with a phased array coil. A section thickness of 7 mm with a 1 mm gap was used for all sequences. T1-weighted fast field-echo image, T2-weighted fast spin-echo image and diffusion-weighted image with b factor 500 sec/mm² were performed. After gadodiamide injection, T1-weighted fast field-echo dynamic image were also obtained during the hepatic arterial, portal venous and delayed phases.

FDG-PET. Whole body FDG-PET was performed in five patients using a GE Advance Scanner (General Electric Medical System, Milwaukee, WI, USA), which has an axial field of view of 15 cm and a spatial resolution of 4.5 mm full-width-half-maximum. All patients fasted for at least 4 h prior to scanning. Sixty min after intravenous injection of 300 MBq of F18-FDG, emission scanning was performed in 5 min and transmission scanning in 1 min. Data acquisition was performed in 7 bed positions.

In the remaining two patients, PET/CT scanning was performed using a Discovery LS PET/CT system (General Electric Medical Systems, Waukesha, WI, USA) because our PET system had been replaced by PET/CT. The CT component was performed using a multi-detector scanner. The parameters were 140 kV, 80 Ma, 0.8 s/CT rotation, a pitch of 6, and a table speed of 22.5 mm/s. Scans were acquired from the skull base to mid-thigh level, in 7 bed positions, with a total acquisition time of 31.9 sec to 37 sec. CT data was resized from a 512x512 matrix to a 128x128 matrix to match the PET data, to allow for image fusion and generation of CT transmission maps. The PET data were also acquired in the same anatomic positions; in 7 bed positions at 5 min per position.

All PET studies were performed at least 4 weeks after completion of chemotherapy.

Rescue hepatectomy. At the National Cancer Center Hospital East, Japan, all lesions considered positive for malignancy, by any pre-operative diagnostic imaging evaluation, were resected by rescue hepatectomy. During this operation, all the tiny suspicious lesions that were definitive metastases before chemotherapy were resected.

In our patients, a careful search was performed after laparotomy for local recurrence, extrahepatic metastases and peritoneal dissemination in the abdominal cavity. Any suspicious lesions were examined by biopsy. Intra-operative bimanual liver palpation and ultrasonography were performed to confirm tumor location and size, in all seven patients and all of the resections were ultrasound-guided procedures. Hepatic resection was performed with tumor-free resection margins, by the forceps fracture method, under inflow occlusion (Pringle's maneuver). When small lesions that had been detected by pre-operative imaging could not be recognized by either palpation or intra-operative ultrasonography, the estimated part or segment of the liver occupied by the lesions was resected.

Extended lobectomy and multiple partial resections were performed on two patients each, and central bi-segmentectomy, lobectomy, and segmentectomy were performed on one patient each, according to Couinaud's anatomical classification.

Assessment of tumor viability with FDG-PET. In our hospital, when interpreting conventional CT and MRI findings, any hepatic lesions are classified according to the degree of confidence that a metastatic tumor is present, as follows: definitely present, probably present, possibly present, probably absent and definitely absent. Lesions that fall into the definitely present, probably present and possibly present categories are considered positive for malignancy, while lesions that fall into the other categories are considered negative. Lesions which are positive, by either CT or MRI, are considered positive by conventional examination.

In the present study, the category "possibly present" included the small lesions, detected using conventional imaging, that were used to indicate definitive metastasis, but which then could not be determined as viable or otherwise, due to a reduction in size or remarkable calcification after chemotherapy (Figure 1).

All lesions considered positive by conventional examination were compared with their corresponding pathology findings, as the standard reference and the sensitivity and positive predictive value were then calculated. Furthermore, these lesions were also assessed by FDG-PET according to the degree of confidence of malignancy, *i.e.* definitely present, probably present, possibly present, probably absent, and definitely absent. A discrete focus with increased FDG accumulation, markedly greater than that in the hepatic parenchyma, was interpreted as malignancy, being definitely present or probably present. Focally increased FDG uptake, minimally greater than in the liver, was considered possibly positive for malignancy, but heterogeneous uptake in the hepatic parenchyma without a focal lesion was considered to indicate that malignancy was probably absent. Lesions in the definitely present, probably present and possibly present categories were considered positive for malignancy, while those in the probably absent and definitely absent categories were considered negative.

FDG-PET findings were also compared with the pathology findings, as the reference, and then sensitivity, specificity, positive predictive value and negative predictive value were calculated according to tumor size: <2 cm and >2 cm.

Finally, the sensitivities, specificities, positive predictive values and negative predictive values were compared between the subgroups according to tumor size. The results of each imaging test were interpreted by at least two experienced radiologists.

Pathological examination. The resected hepatic specimens were fixed in 10% phosphate-buffered formalin, sliced at 5 mm intervals and embedded in paraffin. The findings of all lesions, considered positive




Figure 1. a) A 45-year-old male with initially unresectable multiple colorectal hepatic metastases. (a-1) CT before chemotherapy. (a-2) CT after chemotherapy. Significant calcification was seen in the tumors. After rescue hepatectomy, histological examination revealed adenocarcinoma in most of the calcificated tumors. b) A 65-year-old female with multiple colorectal hepatic metastases. (b-1) CT before chemotherapy. (b-2) CT after chemotherapy. Tumors had reduced markedly in size.

by any diagnostic imaging or intra-operative examination, were confirmed macroscopically and the lesions were then examined microscopically to evaluate viability. Serial sections 3 μ m thick were stained with hematoxylin and eosin (H&E) for morphological examination. Histological diagnoses were based on the World Health Organization classification (16).

Statistical analysis. χ^2 analysis was used to assess sensitivity, specificity, positive predictive value and negative predictive value between subgroups according to tumor size. A *p*-value of less than 0.05 was considered to denote statistical significance.

Results

In the seven patients with CHMs down-staged by chemotherapy, 27 lesions were resected, all of which were considered positive for malignancy by pre-operative diagnostic imaging or intra-operative examination. Among the 27 lesions, 26 were deemed positive by at least one imaging modality, while the other could not be evaluated using imaging and was diagnosed by intra-operative examination. Twelve lesions were

Table I. Comparisons of interpretations of CT, MR images and FDG-PET with pathology.

Imaging finding	Pathological findings	
	Malignancy (12)	No malignancy (15)
CT and MR Imaging		
Positive (26)	11	15
Negative (1)	1	0
FDG-PET		
Positive (7)	7	0
Negative (20)	5	15

histologically diagnosed as adenocarcinoma, but no malignancies were demonstrated in the remaining 15 lesions.

CT and MRI led to 26 lesions being diagnosed as malignant, while only 7 of these 26 lesions were considered positive by FDG-PET (Figure 2). No lesion was positive only by FDG-PET.




Figure 2. A 44-year-old female with multiple colorectal hepatic metastases. (a-1, 2 and 3) CT before chemotherapy. Tumors were initially unresectable because of invasion to the hepatic hilum. (b-1, 2 and 3) CT after chemotherapy. Tumors in segments III, IV / V and VII showed significant reduction in size. (c-1, 2 and 3) FDG-PET after chemotherapy. By FDG-PET, tumors located in segments IV / V and VIII were considered positive, but tumors in segments III and VII were negative. Extended left lobectomy plus partial resections were performed. Tumors in segments III, IV / V and VIII proved histologically to be adenocarcinoma, but the tumor in segment VII had no viable cells.

The interpretations of the conventional (CT/MRI) findings and FDG-PET findings were compared with the results of pathology in Table I. The sensitivity and positive predictive value of the conventional modalities were 92% (95% CI=0.73-1.00) and 42% (0.22-0.63), respectively, while the sensitivity, specificity, positive predictive value and negative predictive value of FDG-PET were 58% (95% CI=0.26-0.91), 100% (1.00-1.00), 100% (1.00-1.00) and 75% (0.54-0.96), respectively.

An assessment of the accuracy of conventional imaging and FDG-PET, according to tumor size (Table II), revealed that interpretation of the conventional findings and FDG-PET was highly accurate in lesions >2 cm. However, in lesions <2 cm, the positive predictive value for conventional imaging was only 26%, while the sensitivity for FDG-PET was only 17%, both

of which were significantly lower than in lesions >2 cm. The specificity and negative predictive value of conventional imaging were not calculated, because lesions negative by both CT and MRI, were not resected, except for one which was diagnosed as positive by intra-operative examination. The viability of the 27 tumors could not be determined by any cut-off value of tumor-size (data not shown).

Discussion

We assessed the efficacy of FDG-PET in evaluating the tumor viability of CHM down-staged by chemotherapy, before rescue surgery. Our results indicate that FDG-PET is only effective in assessing the viability of hepatic lesions >2 cm after chemotherapy.

Table II. Accuracies of CT, MRI, and FDG-PET in evaluation of tumor viability according to tumor size.

	Sensitivity (%)	Specificity (%)	Positive predictive value (%)	Negative predictive value (%)	P-value
CT, MRI tumor size					
<2 cm	83 (5/6; 0.41, 1.00)	—	26 (5/19; 0.05, 0.48)*	—	<0.01*
≥2 cm	100 (6/6; 1.00, 1.00)	—	86 (6/7; 0.51, 1.00)*	—	
FDG-PET tumor size					
<2 cm	17 (1/6; 0.00, 0.60)†	100 (14/14; 1.00, 1.00)	100 (1/1; —, —)	74 (14/19; 0.52, 0.96)	<0.01†
≥2 cm	100 (6/6; 1.00, 1.00)†	100 (1/1; 1.00, 1.00)	100 (6/6; 1.00, 1.00)	100 (1/1; —, —)	

Numbers in parentheses are the data used to determine the percentages and the 95% confidence intervals. *Difference between positive predictive value of CT and MRI in tumor <2 cm and that in tumor ≥2 cm. †Difference between sensitivity of FDG-PET in tumor <2 cm and that in tumor ≥2 cm.

The ability to use FDG-PET to assess the pathological tumor response to pre-operative chemotherapy, radiation or other treatment has been suggested in several tumors (11-15). In addition, correlation between decreased uptake in FDG-PET and histopathological response in the resected specimen was observed in patients, who underwent pre-operative chemotherapy or chemoradiotherapy for esophageal squamous cell carcinoma (17, 18), rectal cancer (19) and gastric carcinoma (20). However, the correlation between uptake in FDG-PET and the corresponding pathological findings has not been studied fully in CHM.

In the present study, portal phase helical CT and MRI were used as the conventional modalities. In our institution, diagnosis based on either, is routinely performed before hepatic resection for CHM, because portal phase helical CT has shown excellent sensitivity in detecting CHM and is considered as the standard pre-operative examination for CHM (21-23). Furthermore, SPIO-enhanced MRI and diffusion-weighted sensitivity encoding MRI have demonstrated high sensitivity equal to that of portal phase helical CT, and have excellent specificity in detecting CHM (24-26).

The present lesion-by-lesion analysis demonstrated that only 11 out of the 26 lesions that had pathological malignancy, considered positive for malignancy by conventional imaging. In the conventional examinations, the sensitivity of 92% for detecting hepatic viable CHMs was similar to that in the aforementioned studies, but the positive predictive value in tumors <2 cm was only 26%. Thus, CT and MRI were able to detect even small tumors, but could hardly evaluate the viability of small tumors after down-staging by chemotherapy.

On the other hand, FDG-PET showed excellent specificity (100%) and positive predictive value (100%), irrespective of tumor size. In tumors >2 cm, the sensitivity and negative predictive value for FDG-PET were both

100%. However, in tumors <2 cm, sensitivity was extremely low (17%). Thus, many of the tumors that shrunk to <2 cm by chemotherapy were undetectable by FDG-PET.

The reasons for this low sensitivity of FDG-PET in tumors <2 cm may be low spatial resolution, the partial volume effect and decreased FDG uptake in tumor tissue, after chemotherapy. Several groups reported that the sensitivity of FDG-PET for CHM was related to tumor size (27, 28). Lower sensitivity to smaller tumors was shown to be caused by the relatively low spatial resolution of FDG-PET and decreased measured activity concentration owing to the partial volume effect (29, 30). The partial volume effect reduces the measured activity concentration to a greater extent in smaller tumors.

Low sensitivity in FDG-PET has also been ascribed to the effects of chemotherapy itself. Chemotherapy may alter FDG uptake in two ways. First, the chemotherapy may reduce FDG uptake, by causing functional changes in tumor glucose metabolism. Spaepen *et al.* demonstrated that changes in tumor glucose metabolism occurred rapidly after chemotherapy, as demonstrated using transplants of Daudi cells in SCID mice (31). Second, a decrease in viable cells by necrosis or apoptosis induced by chemotherapy may diminish FDG uptake in the tumor. Swisher *et al.* studied the correlation between the percentage of residual tumor and standardized uptake value (SUV) of FDG-PET after pre-operative chemoradiation in patients with esophageal cancer (18). They found that patients with >50% histological tumor viability had a significantly higher average SUV compared with those with <50% histological tumor viability. However, the SUV of tumors with no viability was similar to that of tumors with 10-50% tumor viability. Accordingly, when chemotherapy shows excellent efficacy in CHMs, it is extremely difficult to detect the resulting low FDG uptake by PET.

Our results suggest that the viability of small tumors that have been down-staged by chemotherapy can hardly be evaluated by CT, MRI or FDG-PET. Thus, at the moment, lesions considered positive by either CT or MRI should be treated by surgical resection in order to avoid leaving viable metastases in the residual liver. When surgical resection is not suitable for the specific lesion, perhaps due the small amount of residual liver, locoregional therapy, such as radiofrequency ablation or cryosurgery may become the preferred treatment options (6, 7).

The present study has nevertheless some limitations. The number of subjects in our study was relatively small, although the results have significant implications for rescue surgery for CHMs after chemotherapy. Furthermore, only two of the seven patients underwent PET/CT. The recent introduction of combined FDG-PET and CT has improved imaging accuracy by allowing accurate anatomical localization of FDG uptake. The sensitivity of PET/CT may be superior to that of FDG-PET for the detection of viable CHMs after chemotherapy. However, a major improvement in sensitivity is not expected, because the fundamental problem of detecting small viable tumors in PET is not resolved even using PET/CT.

New functional imaging with higher sensitivity for detecting small viable tumors is necessary to improve rescue surgery for CHM.

FDG-PET can be used to accurately assess the viability of CHMs >2 cm, before rescue surgery for initially unresectable CHMs down-staged by chemotherapy, but cannot be used for hepatic tumors <2 cm, because of the extremely low sensitivity of FDG-PET for such tumors.

Acknowledgements

This work was supported in part by grants from Ministry of Health, Labor and Welfare, Japan.

References

- Goldberg RM, Sargent DJ, Morton RF, Fuchs CS, Ramanathan RK, Williamson SK, Findlay BP, Pitot HC and Alberts SR: A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 22: 23-30, 2004.
- Hurwitz H, Fehrenbacher L, Novotny W, Cartwright T, Hainsworth J, Heim W, Berlin J, Baron A, Griffing S, Holmgren E, Ferrara N, Fyfe G, Rogers B, Ross R and Kabbinavar F: Bevacizumab plus irinotecan, fluorouracil, and leucovorin for metastatic colorectal cancer. *N Engl J Med* 350: 2335-2342, 2004.
- Tournigand C, Andre T, Achille E, Lledo G, Flesh M, Mery-Mignard D, Quinaux E, Couteau C, Buyse M, Ganem G, Landi B, Colin P, Louvet C and de Gramont A: FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: a randomized GERCOR study. *J Clin Oncol* 22: 229-237, 2004.
- Souglakos J, Androulakis N, Syrigos K, Polyzos A, Ziras N, Athanasiadis A, Kakolyris S, Tsousis S, Kouroussis Ch, Vamvakas L, Kalykaki A, Samonis G, Mavroudis D and Georgoulas V: FOLFOXIRI (folinic acid, 5-fluorouracil, oxaliplatin and irinotecan) vs. FOLFIRI (folinic acid, 5-fluorouracil and irinotecan) as first-line treatment in metastatic colorectal cancer (MCC): a multicentre randomised phase III trial from the Hellenic Oncology Research Group (HORG). *Br J Cancer* 94: 798-805, 2006.
- Fowler WC, Eisenberg BL and Hoffman JP: Hepatic resection following systemic chemotherapy for metastatic colorectal carcinoma. *J Surg Oncol* 51: 122-125, 1992.
- Rivoire M, De Cian F, Meeus P, Negrier S, Sebban H and Kaemmerlen P: Combination of neoadjuvant chemotherapy with cryotherapy and surgical resection for the treatment of unresectable liver metastases from colorectal carcinoma. *Cancer* 95: 2283-2292, 2002.
- Adam R, Delvart V, Pascal G, Valeanu A, Castaing D, Azoulay D, Giacchetti S, Paule B, Kunstlinger F, Ghemard O, Levi F and Bismuth H: Rescue surgery for unresectable colorectal liver metastases downstaged by chemotherapy: a model to predict long-term survival. *Ann Surg* 240: 644-657, 2004.
- Delaunoit T, Alberts SR, Sargent DJ, Green E, Goldberg RM, Krook J, Fuchs C, Ramanathan RK, Williamson SK, Morton RF and Findlay BP: Chemotherapy permits resection of metastatic colorectal cancer: experience from Intergroup N9741. *Ann Oncol* 16: 425-429, 2005.
- Ruers TJ, Langenhoff BS, Neeleman N, Jager GJ, Strijk S, Wobbes T, Corstens FH and Oyen WJ: Value of positron emission tomography with [¹⁸F]fluorodeoxyglucose in patients with colorectal liver metastases: a prospective study. *J Clin Oncol* 20: 388-395, 2002.
- Fernandez FG, Drebin JA, Linehan DC, Dehdashti F, Siegel BA and Strasberg SM: Five-year survival after resection of hepatic metastases from colorectal cancer in patients screened by positron emission tomography with F-18 fluorodeoxyglucose (FDG-PET). *Ann Surg* 240: 438-447, 2004.
- Findlay M, Young H, Cunningham D, Iveson A, Cronin B, Hickish T, Pratt B, Husband J, Flower M and Ott R: Noninvasive monitoring of tumor metabolism using fluorodeoxyglucose and positron emission tomography in colorectal cancer liver metastases: correlation with tumor response to fluorouracil. *J Clin Oncol* 14: 700-708, 1996.
- Stroobants S, Goeminne J, Seegers M, Dimitrijevic S, Dupont P, Nuyts J, Martens M, van den Borne B, Cole P, Sciort R, Dumez H, Silberman S, Mortelmans L and van Oosterom A: 18FDG-Positron emission tomography for the early prediction of response in advanced soft tissue sarcoma treated with imatinib mesylate (Glivec). *Eur J Cancer* 39: 7012-2020, 2003.
- Juwaid ME, Wiseman GA, Vose JM, Ritchie JM, Menda Y, Wooldridge JE, Mottaghy FM, Rohren EM, Blumstein NM, Stolpen A, Link BK, Reske SN, Graham MM and Cheson BD: Response assessment of aggressive non-Hodgkin's lymphoma by integrated International Workshop Criteria and fluorine-18-fluorodeoxyglucose positron emission tomography. *J Clin Oncol* 23: 4652-4661, 2005.
- Loi S, Ngan SY, Hicks RJ, Mukesh B, Mitchell P, Michael M, Zalberg J, Leong T, Lim-Joon D, Mackay J and Rischin D: Oxaliplatin combined with infusional 5-fluorouracil and concomitant radiotherapy in inoperable and metastatic rectal cancer: a phase I trial. *Br J Cancer* 92: 655-661, 2005.

- 15 Avril N, Sassen S, Schmalfeldt B, Naehrig J, Rutke S, Weber WA, Werner M, Graeff H, Schwaiger M and Kuhn W: Prediction of response to neoadjuvant chemotherapy by sequential F-18-fluorodeoxyglucose positron emission tomography in patients with advanced-stage ovarian cancer. *J Clin Oncol* 23: 7445-7453, 2005.
- 16 Jass JR and Sobin LH: Histological typing of intestinal tumors. *In: World Health Organization. International Histological Classification of Tumors*, 2nd ed, Jass JR and Sobin LH (eds.). Berlin, Springer-Verlag, 1989.
- 17 Wieder HA, Brucher BL, Zimmermann F, Becker K, Lordick F, Beer A, Schwaiger M, Fink U, Siewert JR, Stein HJ and Weber WA: Time course of tumor metabolic activity during chemoradiotherapy of esophageal squamous cell carcinoma and response to treatment. *J Clin Oncol* 22: 900-908, 2004.
- 18 Swisher SG, Erasmus J, Maish M, Correa AM, Macapinlac H, Ajani JA, Cox JD, Komaki RR, Hong D, Lee HK, Putnam JB Jr, Rice DC, Smythe WR, Thai L, Vaporciyan AA, Walsh GL, Wu TT and Roth JA: 2-Fluoro-2-deoxy-D-glucose positron emission tomography imaging is predictive of pathologic response and survival after preoperative chemoradiation in patients with esophageal carcinoma. *Cancer* 101: 1776-1785, 2004.
- 19 Guillem JG, Moore HG, Akhurst T, Klimstra DS, Ruo L, Mazumdar M, Minsky BD, Saltz L, Wong WD and Larson S: Sequential preoperative fluorodeoxyglucose-positron emission tomography assessment of response to preoperative chemoradiation: a means for determining longterm outcomes of rectal cancer. *J Am Coll Surg* 199: 1-7, 2004.
- 20 Ott K, Fink U, Becker K, Stahl A, Dittler HJ, Busch R, Stein H, Lordick F, Link T, Schwaiger M, Siewert JR and Weber WA: Prediction of response to preoperative chemotherapy in gastric carcinoma by metabolic imaging: results of a prospective trial. *J Clin Oncol* 21: 4604-4610, 2003.
- 21 Kuszyk BS, Bluemke DA, Urban BA, Choti MA, Hruban RH, Sitzmann JV and Fishman EK: Portal-phase contrast-enhanced helical CT for the detection of malignant hepatic tumors: sensitivity based on comparison with intraoperative and pathologic findings. *Am J Roentgenol* 166: 91-95, 1996.
- 22 Valls C, Andia E, Sanchez A, Guma A, Figueras J, Torras J and Serrano T: Hepatic metastases from colorectal cancer: preoperative detection and assessment of resectability with helical CT. *Radiology* 218: 55-60, 2001.
- 23 Soyer P, Pocard M, Boudiaf M, Abitbol M, Hamzi L, Panis Y, Valleur P and Rymer R: Detection of hypovascular hepatic metastases at triple-phase helical CT: sensitivity of phases and comparison with surgical and histopathologic findings. *Radiology* 231: 413-420, 2004.
- 24 Seneterre E, Taourel P, Bouvier Y, Pradel J, Van Beers B, Daures JP, Pringot J, Mathieu D and Bruel JM: Detection of hepatic metastases: ferumoxides-enhanced MR imaging versus unenhanced MR imaging and CT during arterial portography. *Radiology* 200: 785-792, 1996.
- 25 Vogl TJ, Schwarz W, Blume S, Pietsch M, Shamsi K, Franz M, Lobeck H, Balzer T, del Tredici K, Neuhaus P, Felix R and Hammerstingl RM: Preoperative evaluation of malignant liver tumors: comparison of unenhanced and SPIO (Resovist)-enhanced MR imaging with biphasic CTAP and intraoperative US. *Eur Radiol* 13: 262-272, 2003.
- 26 Nasu K, Kuroki Y, Nawano S, Kuroki S, Tsukamoto T, Yamamoto S, Motoori K and Ueda T: Hepatic metastases: diffusion-weighted sensitivity-encoding *versus* SPIO-enhanced MR imaging. *Radiology* 239: 122-130, 2006.
- 27 Rohren EM, Paulson EK, Hagge R, Wong TZ, Killius J, Clavien PA and Nelson RC: The role of F-18 FDG positron emission tomography in preoperative assessment of the liver in patients being considered for curative resection of hepatic metastases from colorectal cancer. *Clin Nucl Med* 27: 550-555, 2002.
- 28 Fong Y, Saldinger PF, Akhurst T, Macapinlac H, Yeung H, Finn RD, Cohen A, Kemeny N, Blumgart LH and Larson SM: Utility of 18F-FDG positron emission tomography scanning on selection of patients for resection of hepatic colorectal metastases. *Am J Surg* 178: 282-287, 1999.
- 29 Geworski L, Knoop BO, de Cabrejas ML, Knapp WH and Munz DL: Recovery correction for quantitation in emission tomography: a feasibility study. *Eur J Nucl Med* 27: 161-169, 2000.
- 30 Weber WA: Use of PET for monitoring cancer therapy and for predicting outcome. *J Nucl Med* 46: 983-995, 2005.
- 31 Spaepen K, Stroobants S, Dupont P, Bormans G, Balzarini J, Verhoef G, Mortelmans L, Vandenberghe P and De Wolf-Peeters C: [(18)F]FDG PET monitoring of tumour response to chemotherapy: does [(18)F]FDG uptake correlate with the viable tumour cell fraction? *Eur J Nucl Med Mol Imaging* 30: 682-688, 2003.

Received September 4, 2006
Accepted September 22, 2006

The Role of a Protease Inhibitor against Hepatectomy

Naoto Gotohda¹, Hiromi Iwagaki², Michitaka Ozaki³, Masaru Konishi¹, Toshio Nakagohri¹, Shinichiro Takahashi¹, Takahito Yagi², Taira Kinoshita¹, Noriaki Tanaka²

¹Department of Surgery, National Cancer Center Hospital East, Kashiwa;

²Department of Gastroenterological Surgery, Okayama University Graduate School of Medicine and Dentistry

Okayama; ³Department of Organ Transplantation and Regenerative Medicine

Hokkaido University Graduate School of Medicine, Faculty of Medicine, Sapporo, Japan

Corresponding Author: Naoto Gotohda, MD, Department of Surgery, National Cancer Center Hospital East

6-5-1 Kashiwanoha, Kashiwa, Chiba 277-8577, Japan

Tel: +81 4 7133 1111, Fax: +81 4 7131 4724, E-mail: ngotohda@east.ncc.go.jp

ABSTRACT

Background/Aims: Nafamostat Mesilate (NM) is a synthetic serine protease inhibitor that is capable of inhibiting the various coagulation factors. To determine whether NM may also be useful in attenuating operative invasiveness, we investigated the effects of perioperative administration of NM on postoperative serum levels of proinflammatory cytokine IL-6 and hepatocyte growth factor (HGF).

Methodology: Thirty patients undergoing hepatectomy with hepatocellular carcinoma, biliary carcinoma and metastatic colorectal cancer were enrolled in this study. These patients were separated into two groups; high invasive group (resected liver volume: 1000cm³ <) and less invasive group (resected liver volume: 1000cm³ >). The high invasive group of 11 patients received perioperative administration of NM (Group NM), while the less invasive group of 19 patients did not (Group C). Serum levels of IL-6, HGF and soluble IL-6 receptor (sIL-6R) were simultaneously measured on preoperative and postoperative day ('day 0', 'day 7').

Results: Serum IL-6 levels on day 0 were significantly elevated and returned to preoperative levels on day 7 in both groups, and the serum IL-6 level in Group NM on day 0 was significantly lower than that in Group C on day 0. Serum HGF levels on day 0 and day 7 were significantly higher in Group NM than those in Group C. Compared with healthy control subjects, the higher serum level of HGF on the preoperative day in all patients was attributable to tumor-burden. The sIL-6R levels on day 0 decreased in both groups, and their levels in Group NM were significantly lower than those in Group C, suggesting that increased synthesis of IL-6/sIL-6R complex which could accelerate liver regeneration.

Conclusions: These results suggested that perioperative administration of NM may attenuate surgical stress by decreasing production of proinflammatory cytokine IL-6, and may accelerate liver regeneration through stimulation with the IL-6/sIL-6R complex and possible involvement of increased production of HGF.

KEY WORDS:

Protease inhibitor; Nafamostat mesilate; Hepatectomy; Interleukin-6; soluble Interleukin-6 receptor; Hepatocyte growth factor

ABBREVIATIONS:

Nafamostat Mesilate (NM); Hepatocyte Growth Factor (HGF); Soluble Interleukin-6 Receptor (sIL-6R); Hepatocellular Carcinoma (HCC); C-Reactive Protein (CRP)

INTRODUCTION

Nafamostat mesilate (6-amino-2-naphthyl p-guanidinobenzoate dimethansulfonate, NM) is a synthetic protease inhibitor generated during the coagulation cascade as well as in the inflammatory process (1,2). NM inhibits coagulation factors such as factor VIIa and thrombin (3), and has been found to be effective in treating animals with induced sepsis (4,5). Moreover, regarding its anticoagulant effects, recent reports demonstrated that NM had inhibitory effects on the production of polymorphonuclear leukocyte elastase and Interleukin-6 (IL-6) and IL-8 in human monocytes (6,7).

In liver, IL-6 is secreted by non-parenchymal cells such as Kupffer cells. Secreted IL-6 acts on neighboring hepatocytes to stimulate liver regeneration and repair (8). It also acts on cells by binding the soluble IL-6 receptor (sIL-6R), either in its membrane-bound or soluble form. And this IL-6/sIL-6R complex binds and

induces liver regeneration and hepatoprotection (9).

To determine whether NM may also be useful in attenuating surgical stress and has a hepatoprotective effect or stimulates liver regeneration, we investigated the effects of perioperative administration of NM on postoperative serum levels of proinflammatory cytokine against hepatectomy.

METHODOLOGY

Thirty consecutive patients undergoing hepatectomy with hepatocellular carcinoma (HCC), biliary tract carcinoma and metastatic colorectal cancer were enrolled in this study.

All liver resections were performed under conditions of continuous hepatic pedicle occlusion. Hepatic parenchymal transection was performed with clamp crushing methods. Liver resections were included in 9 major hepatectomies (2 ≤ segment), 11 minor hepatectomies (1 ≥ segment) and 10 partial resections.

These patients were separated into two groups; high invasive group (resected liver volume: $1000\text{cm}^3 <$) and less invasive group (resected liver volume: $1000\text{cm}^3 >$). The high invasive group of 11 patients received perioperative administration of NM (Group NM), while the less invasive group of 19 patients did not (Group C). In group NM, NM was continuously administered at a dosage of 0.1mg/kg/hr through a peripheral vein starting 30 minutes before the operation for 24 hours.

Eleven patients in group NM included 6 with HCC, 1 with biliary tract carcinoma and 4 with metastatic colorectal carcinoma. Nineteen patients in

group C included 13 with HCC, 3 with biliary tract carcinoma and 3 with metastatic colorectal carcinoma (Table 1). There were no significant differences in age and gender between groups.

In all patients peripheral blood samples were collected before operation ('preop'), immediately after operation ('day 0'), and on day 7 after operation ('day 7'). Plasma concentrations of IL-6 (Fujirebio Inc., Tokyo, Japan), sIL-6R (R&D systems, Minneapolis, MN), hepatocyte growth factor (HGF, Otsuka Pharmaceutical Co. Ltd., Tokyo, Japan) were simultaneously measured on preoperative and postoperative day ('day 0' and 'day 7') using an enzyme-linked immunosorbent assay (ELIZA) kit. Moreover, the preoperative serum levels of HGF in all patients were compared with 109 healthy control subjects.

Percent recovery of liver size was calculated with resected liver mass and volumetric analysis by computed tomography of preoperative and postoperative (4 wk after hepatectomy) liver mass. A: [surgically resected liver volume] was considered as 'an expected liver mass of full recovery', and B: [resected liver mass - (preoperatively measured liver mass - postoperatively measured liver mass)] was considered as 'a real recovery of liver'. B divided by A provides recovery ratio of liver, which was expressed as percent recovery of liver.

Statistical Analysis

All data were expressed as means \pm SE. The Mann-Whitney's U test and the Chi-square test were used to analyze samples, respectively. P values less than 0.05 were considered statistically significant.

RESULTS

General Comparison of Patients in Preoperative Liver Function, Surgical Stresses, Postoperative Liver Function and Liver Recovery Ratio

There was no significant difference between group C and group NM in preoperative liver function. All factors of surgical stress in group NM were significantly higher than those in group C (Table 2). The postoperative rise of C-reactive protein (CRP) and postoperative morbidity in group C was significantly lower than those in group NM. No significant differences were observed between the two groups in postoperative other liver function, mortality and hospitalization (Tables 2 and 3). In liver recovery ratio, there was no significant difference between both groups (Figure 1).

Comparison of Serum Levels IL-6 and sIL-6R between Group C and Group NM

Serum IL-6 levels on day 0 were significantly elevated and returned to preoperative levels on day 7 in both groups, and the serum IL-6 level in group NM on day 0 was significantly lower than that in group C on day 0. The sIL-6R levels on day 0 decreased in both groups, and their levels in group NM were significantly lower than those in group C (Table 4).

TABLE 1 Patient Profile

	Group C	Group NM	P
No. of patients	19	11	
Age (years)	68 ± 1.8	68 ± 2.5	*NS
Sex (M/F)	15/4	7/4	§NS
Viral Hepatitis	10	5	§NS
	(Type B: 1, Type C: 9)	(Type B: 4, Type C: 1)	§NS
Disease			
Hepatocellular carcinoma	13	6	
Biliary tract carcinoma	3	1	
Liver metastases of colorectal carcinoma	3	4	
Liver Resection			§P=0.0021
Partial Resection	10	0	
Minor $1 \geq$ Segment	7	4	
Major $2 \leq$ Segment	2	7	

NM: Nafamostat Mesilate.

Values represent (mean \pm SE. NS: Not significant); *Statistical significance between groups was analyzed by Mann-Whitney's U test. §Statistical significance between groups was analyzed by Chi-square test.

TABLE 2 General Comparison of Patients in Preoperative, Surgical and Postoperative States

	Group C	Group NM	P
Preoperative liver function			
Alanine aminotransferase (ALT) (IU/dL)	68 ± 16	50 ± 10	*NS
Total bilirubin (T-Bil) (mg/dL)	0.9 ± 0.1	1.0 ± 0.1	*NS
Albumin (Alb) (g/dL)	3.7 ± 0.1	3.8 ± 0.1	*NS
Prothrombin time (PT) (%)	12 ± 0.1	12 ± 0.2	*NS
ICGR ₁₅ (%)	15 ± 2.3	13 ± 1.6	*NS
Platelet ($\times 10^4/\mu\text{L}$)	17 ± 2.2	21 ± 1.4	*NS
C-reactive protein (mg/dL)	0.2 ± 0.1	0.3 ± 0.1	*NS
Surgical stress			
Operative time (min)	200 ± 32	370 ± 49	*0.0055
Hepatic ischemic time (min)	52 ± 6.0	98 ± 15	*0.0034
Blood loss (mL)	880 ± 190	3500 ± 1300	*0.0133
Resected liver weight (g)	110 ± 30	430 ± 68	*0.0002
Resected liver volume (cm^3)	250 ± 59	1400 ± 83	* <0.0001
Postoperative liver function (day 7)			
Alanine aminotransferase (ALT) (IU/dL)	71 ± 7.9	91 ± 17	*NS
Total bilirubin (T-Bil) (mg/dL)	1.3 ± 0.4	1.3 ± 0.3	*NS
Albumin (Alb) (g/dL)	3.4 ± 0.1	3.3 ± 0.1	*NS
Prothrombin time (PT) (%)	12 ± 0.1	12 ± 0.2	*NS
Platelet ($\times 10^4/\mu\text{L}$)	20 ± 2.0	21 ± 2.2	*NS
C-reactive protein (mg/dL)	3.2 ± 0.6	7.4 ± 1.5	*0.0059

Values represent (mean \pm SE. NS: Not significant), *Statistical significance between groups was analyzed by Mann-Whitney's U test.

Serum Level HGF in All Patients with Each Disease

In preoperative serum level of HGF, there was a significant difference between all patients enrolled in this study and healthy control subjects. All serum HGF levels on preop, day 0 and day 7 in group NM were higher than those in group C (Table 5).

DISCUSSION

A synthetic protease inhibitor has an inhibitory effect on various serine proteases such as trypsin, thrombin, activated factor X, kallikrein, neutrophil elastase, and activated complement components (1). Effects of synthetic protease inhibitors on microcirculatory and coagulation disorder and on organ failure such as septic ARDS have also been reported (10-14).

Recently, preoperative administration of protease inhibitor substantially ameliorated hepatocyte injury induced by ischemia and reperfusion in human patients when liver resection was performed under continuous inflow occlusion (15). Moreover, there were some reports that serine protease inhibitors may play an important role in liver regeneration (16,17).

These many studies using animal models have demonstrated that hepatically-derived cytokines are produced in the process of hepatic ischemia and reperfusion (18-21). IL-6 is a proinflammatory cytokine that mediates the acute-phase inflammatory response to tissue injury. The greater the surgical stresses, the higher the elevation of serum IL-6 (22-24). Although the precise role of IL-6 in hepatocyte injury remains to be clarified, at least excess production of IL-6 may be associated with adverse events after liver resection in humans because there were many reports that the over-production of plasma IL-6 is followed by major postoperative elevation of plasma transaminase activities after prolonged hepatic inflow occlusion (25-27). However, recent reports have demonstrated that HGF and IL-6 could promote hepatic survival by stimulating liver regeneration and providing hepatoprotection in a variety of liver injury models, including Fas-mediated injury, toxic damage caused by hepatotoxins (such as tetrachlorocarbonide), and ischemic liver injury (28-32). Particularly, IL-6 is a critical proregenerative factor and acute-phase inducer in the liver that also confers resistance to liver injury by hepatic toxins, ischemia, and Fas. Its effects are mediated almost exclusively on hepatocytes within the liver (33). Secreted IL-6 acts on neighboring hepatocytes in a paracrine fashion to stimulate liver regeneration and repair (8). IL-6 bound to the sIL-6R signals via gp130 and Janus kinase-1 (JAK-1), leading to activation of the Stat3 transcription factor and the MAPK signal transduction cascade (33,34). In this way, the IL-6 signal transduction pathway in liver injury and regeneration was clarified. Our results would also suggest that increased synthesis of IL-6/sIL-6R complex which could accelerate liver regeneration.

HGF could be produced in many organs or released from the extracellular matrix of the liver. HGF is a potent stimulator of DNA synthesis in hepa-

TABLE 3 General Comparison of Patients in Postoperative Outcomes

	Group C	Group NM	P
Postoperative mortality	1/19	0/11	*NS
Postoperative morbidity	1/19	4/11	*0.0288
Wound infection	1	1	
Pulmonary infarction	0	1	
Bile leakage	0	1	
Paralytic ileus	0	1	
Postoperative hospitalization (days)	15±3.9	20±4.6	*NS

Values represent (mean ± SE. NS: Not significant), *Statistical significance between groups was analyzed by Mann-Whitney's U test. *Statistical significance between groups was analyzed by Chi-square test.




FIGURE 1 Percent recovery of liver size 4 weeks after hepatectomy. Percent recovery of liver size was calculated by volumetric analysis by resected liver volume, computed tomography of liver preoperatively and 4 wk after hepatectomy. *Statistical significance between groups was analyzed by Mann-Whitney's U test.

TABLE 4 Comparison of Serum Levels of IL-6, sIL-6R, HGF With/Without Nafamostat Mesilate

		Group C	Group NM	P
IL-6 (pg/mL)	Preop	4.3±1.2	3.0±0.7	NS
	Postop (day 0)	240±170	180±51	0.0454
	Postop (day 7)	23±6.1	49±15	NS
sIL-6R (pg/mL)	Preop	29000±2000	24000±1700	NS
	Postop (day 0)	26000±1500	17000±850	0.0007
	Postop (day 7)	28000±1800	24000±2100	NS

Values represent mean ± SE. Statistical significance between groups was analyzed by Mann-Whitney's U test.

toocytes and interacts with other growth factors. After partial hepatectomy, the increased tyrosine phosphorylation of c-MET/HGF-receptor is seen in the hepatocytes. The HGF/c-MET system is also involved in liver regeneration. There is increased synthesis of HGF by nonparenchymal cells after partial hepatectomy (35,36), and therefore, increased HGF production is expected on liver injury and has been described as stimulating liver regeneration factor. And its supportive roles in liver regeneration are expected in many clinical situations.

In the other hand, HGF may well be an important factor in tumor propagation and development of distant metastases because of its profound impact on cell proliferation and motility (37). And then, it has already been reported that HGF has been a useful

TABLE 5 Comparison of Serum Level of HGF in Preoperative Patients and Normal Control, Comparison of Serum Levels of HGF With/Without Nafamostat Mesilate in Postoperative Patients

HGF (pg/mL)	All patients (n=30)	Normal control (n=109)	P
Preop	445±61	293±5.2	0.0002
	Group C	Group NM	P
Postop (day 0)	500±80	900±390	NS
Postop (day 7)	460±44	640±80	0.0210

Values represent mean ± SE. Statistical significance between all patients and normal control was analyzed by Mann-Whitney's U test.

indicator for predicting the status of HCC, colorectal cancer and cholangiocellular carcinoma (36,38-42). Compared with group C, our results demonstrated that the serum level of HGF was higher in group NM. However, the preoperative serum level of HGF was high in group NM, and we wondered whether that result was reflected by the activity of cancer cell. We evaluated the difference between the serum levels of HGF in cancer patients and healthy control subjects. We found a significant difference between patients enrolled in this study and healthy control subjects, which was consistent with a previous report that the serum HGF concentration of patients with colorectal cancer was significantly higher than that in healthy normal control level of 174 ± 14 pg/mL as means ± SE (40). The HGF level in the healthy control in our study of 293 ± 5.2 pg/mL was higher than that in a previous report (40), which was attributable to ELISA kits used.

Excluding the role of the serum HGF concentration as predicting index of tumor propagation, we considered that the effect to liver regenerative rate was reflected by the serum levels of HGF on day 0 and day 7. In the serum levels of HGF on day 0 and day 7, the

level of HGF in group NM was higher than that in group C, we considered that these levels of HGF were increased because the synthesis of HGF was stimulated by the administration of NM. Shimomura *et al.* (43) also reported that blood-coagulation factor XIIa and HGF activator activated HGF, and the HGF-converting activity of the HGF activator was not prevented by the serine protease inhibitors. Though there were some reports that liver regeneration was influenced by the loss of liver volume by liver resection (44,45), these results might insinuate that the higher level of HGF, the better the liver regeneration. And then, protease inhibitor may have a good role in the liver regeneration.

The present study demonstrated that although no significant difference was observed between both groups in the liver regenerative rate, this result seemed to be affected by the high incidence of postoperative morbidity in group NM. Indeed, the postoperative CRP level was significantly higher in group NM in the present research. Weiss *et al.* (46) reported that the growth rate of hepatic regeneration was lower in fulminant sepsis than in mild sepsis. The postoperative morbidity is very susceptible to surgical stresses, all indicators of the surgical stress were higher in group NM than in group C. Despite the high incidence of postoperative morbidity and high surgical stresses in group NM, postoperative recovery rate of liver in group NM was equivalent to that of liver in group C. These results suggested that the perioperative administration of NM may attenuate surgical stress by decreasing production of proinflammatory cytokine IL-6, and may accelerate liver regeneration through increased production of HGF and stimulation with the IL-6/sIL-6R complex. It would be possible that much better liver regeneration is induced by the administration of NM if we could improve our surgical techniques and keep the lower incidence of postoperative morbidity in the liver surgery.

REFERENCES

- 1 Paques EP, Romisch J: Comparative study on the in vitro effectiveness of antithrombotic agents. *Thromb Res* 1991; 64:11-21.
- 2 Okajima K, Uchiba M, Murakami K: Nafamostat mesilate. *Cardiovasc Drug Rev* 1995; 13:51-65.
- 3 Uchiba M, Okajima K, Abe H, Okabe H, Takatsuki K: Effect of nafamostat mesilate, a synthetic protease inhibitor, on tissue factor-factor VIIa complex activity. *Thromb Res* 1994; 74:155-161.
- 4 Takahashi H, Takizawa S, Tatewaki W, Nagai K, Wada K, Hanano M, et al: Nafamostat mesilate (FUT-175) in the treatment of patients with disseminated intravascular coagulation. *Thromb Haemost* 1989; 62:372.
- 5 Yoshikawa T, Murakami M, Furukawa Y, Kato H, Takemura S, Kondo M: Effects of FUT-175, a new synthetic protease inhibitor on endotoxin-induced disseminated intravascular coagulation in rats. *Hemostasis* 1983; 13:374-378.
- 6 Kikuchi M, Endo S, Inada K, Yamashita H, Takakuwa T, Nakae H, et al: Inhibitory effect of FUT-175 on the production of interleukin-8 and polymorphonuclear leukocyte elastase. *Res Comm Mole Path & Pharm* 1995; 87:269-274.
- 7 Sugita H, Ishiko T, Ikei S, Hirota M, Ogawa M: FUT-175 inhibits the production of IL-6 and IL-8 in human monocytes. *Res Comm Mole Path & Pharm* 1999; 103:57-64.
- 8 Zimmers TA, Mckillop IH, Pierce RH, Yoo JY, Koniaris LG: Massive liver growth in mice induced by systemic interleukin 6 administration. *Hepatology* 2003; 38:326-334.
- 9 Bowman T, Garcia R, Turkson J, Jove R: STATs in oncogenesis. *Oncogene* 2000; 19:2474-2488.
- 10 Shimada M, Matsumata T, Shirabe K, Kamakura T, Taketomi A, Sugimachi K: Effect of nafamostat mesilate on coagulation and fibrinolysis in hepatic resection. *J Am Coll Surg* 1994; 178:498-502.
- 11 Suzuki M, Suematsu M, Miura S, Oshio C, Oda M, Tsuchiya M: Microcirculatory disturbances in endotoxin-induced disseminated intravascular coagulation. The effects of heparin and gabexate mesilate on locomotive and metabolic changes of neutrophils. *Adv Exp Med Biol* 1988; 242:135-141.
- 12 Tamaki S, Wada H, Hiyoyama K, Shimura M, Minamikawa K, Wakita Y, et al: Treatment of disseminated intravascular coagulation with gabexate mesilate. *Clin Ther* 1993; 15:1076-1084.
- 13 Okumura Y, Inoue H, Fujiyama Y, Bamba T: Effects of

- serine protease inhibitors on accumulation of polymorphonuclear leukocytes in the lung induced by acute pancreatitis in rats. *J Gastroenterol* 1995; 30:379-386.
- 14 Iwagaki H, Yagi T, Urushihara N, Kobayashi K, Morimoto Y, Isozaki H, et al: Modulatory effect of a serine protease inhibitor on surgical stress: its clinical implications. *Acta Med Okayama* 1999; 53:239-244.
 - 15 Kim YI, Hwang YJ, Song KE, Yun YK, Lee JW, Chun BY: Hepatocyte protection by a protease inhibitor against ischemia/reperfusion injury of human liver. *J Am Coll Surg* 2002; 195:41-50.
 - 16 Fausto N, Laird AD, Webber EM: Liver regeneration. 2. Role of growth factors and cytokines in hepatic regeneration. *FASEB J* 1995; 9:1527-1536.
 - 17 Bhanumathy CD, Tang Y, Monga SP, Katuri V, Cox JA, Mishra B, et al: Itih-4, a serine protease inhibitor regulated in interleukin-6-dependent liver formation: role in liver development and regeneration. *Dev Dyn* 2002; 223:59-69.
 - 18 Colletti LM, Remick DG, Burtch GD, Kunkel SL, Strieter RM, Campbell DA: Role of tumor necrosis factor- α in the pathophysiologic alterations after hepatic ischemia/reperfusion injury in the rat. *J Clin Invest* 1990; 85:1936-1943.
 - 19 McCurry KR, Campbell DA Jr, Scales WE, Warren JS, Remick DG: Tumor necrosis factor, interleukin 6, and the acute phase response following hepatic ischemia/reperfusion. *J Surg Res* 1993; 55:49-54.
 - 20 Kawano K, Kim YI, Kai T, Ishii T, Tatsuma T, Morimoto A, et al: Evidence that FK506 alleviates ischemia/reperfusion injury to the rat liver. In vivo demonstration for suppression of TNF- α production in response to endotoxemia. *Eur Surg Res* 1994; 26:108-115.
 - 21 Suzuki S, Toledo-Pereyra LH: Interleukin 1 and tumor necrosis factor production, as the initial stimulants of liver ischemia and reperfusion injury. *J Surg Res* 1994; 57:253-258.
 - 22 Ozawa A, Konishi F, Nagai H, Okada M, Kanazawa K: Cytokine and hormonal responses in laparoscopic-assisted colectomy and conventional open colectomy. *Surg Today* 2000; 30:107-111.
 - 23 Iwagaki H, Yagi T, Urushihara N, Morimoto Y, Jikuhara A, Isozaki H, et al: Blood transfusion and post-operative plasma cytokine antagonist levels in colorectal cancer patients. *Hepatogastroenterology* 2001; 48:1351-1354.
 - 24 Sadamori H, Yagi T, Kanaoka Y, Morimoto M, Inagaki T, Ishikawa H, et al: The analysis of the usefulness of laparoscopic microwave coagulation therapy for hepatocellular carcinoma in patients with poor hepatic reserve by serial measurements of IL-6, cytokine antagonists, and C-reactive protein. *Surg Endosc* 2003; 17:510-514.
 - 25 Man K, Fan ST, Ng IO, Lo CM, Liu CL, Yu WC, et al: Tolerance of the liver to intermittent pringle maneuver in hepatectomy for liver tumors. *Arch Surg* 1999; 134:533-539.
 - 26 Sugawara Y, Kubota K, Ogura T, Esumi H, Inoue K, Takayama T, et al: Increased nitric oxide production in the liver in the perioperative period of partial hepatectomy with Pringle's maneuver. *J Hepatol* 1998; 28:212-220.
 - 27 Kim YI, Song KE, Ryeon HK, Hwang YJ, Yun YK, Lee JW, et al: Enhanced inflammatory cytokine production at ischemia/reperfusion in human liver resection. *Hepatogastroenterology* 2002; 49:1077-1082.
 - 28 Taub R, Greenbaum LE, Peng Y: Transcriptional regulatory signals define cytokine-dependent and independent pathways in liver regeneration. *Semin Liver Dis* 1999; 19:117-127.
 - 29 Kovalovich K, DeAngelis RA, Li W, Furth EE, Ciliberto G, Taub R: Increased toxin-induced liver injury and fibrosis in interleukin-6-deficient mice. *Hepatology* 2000; 31:149-159.
 - 30 Kovalovich K, Li W, DeAngelis R, Greenbaum LE, Ciliberto G, Taub R: Interleukin-6 protects against Fas-mediated death by establishing a critical level of anti-apoptotic hepatic proteins FLIP, Bcl-2, and Bcl-xL. *J Biol Chem* 2001; 276:26605-26613.
 - 31 Galun E, Zeira E, Pappo O, Peters M, Rose-John S: Liver regeneration induced by a designer human IL-6/sIL-6R fusion protein reverses severe hepatocellular injury. *FASEB J* 2000; 14:1979-1987.
 - 32 Ozaki M, Haga S, Zhang H, Irani K, Suzuki S: Inhibition of hypoxia/reoxygenation-induced oxidative stress in HGF-stimulated antiapoptotic signaling: role of PI3-K and Akt kinase upon racl. *Cell Death Differ* 2003; 10:508-515.
 - 33 Rebecca T: Hepatoprotection via the IL-6/Stat3 pathway. *J Clin Invest* 2003; 112:978-980.
 - 34 Wuestefeld T, Klein C, Streetz KL, Betz U, Lauber J, Buer J, et al: Interleukin-6/Glycoprotein 130-dependent pathways are protective during liver regeneration. *J Biol Chem* 2003; 278:11281-11288.
 - 35 Fausto N: Liver regeneration. *J Hepatol* 2000; 32:19-31.
 - 36 Dluzniewska J, Zolich D, Polanski J, Zajac L, Sitkiewicz D, Lukomska B: Hepatocyte growth factor levels in liver and blood, and post-operative liver cell proliferation in patients with benign and malignant liver tumors after partial hepatectomy. *Med Sci Monit* 2002; 8:CR690-696.
 - 37 Boros P, Miller CM: Hepatocyte growth factor: a multi-functional cytokine. *Lancet* 1995; 345:293-295.
 - 38 Yamagami H, Moriyama M, Matsumura H, Aoki H, Shimizu T, Saito T, et al: Serum concentrations of human hepatocyte growth factor is a useful indicator for predicting the occurrence of hepatocellular carcinomas in C-viral chronic liver diseases. *Cancer* 2002; 95:824-834.
 - 39 Wu T, Han C, Lunz JG 3rd, Michalopoulos G, Sheldhamer JH, Demetris AJ: Involvement of 85-kd cytosolic phospholipase A2 and cyclooxygenase-2 in the proliferation of human cholangiocarcinoma cells. *Hepatology* 2002; 36:363-373.
 - 40 Fukuura T, Miki C, Matsumoto K, Suzuki H: Serum hepatocyte growth factor as an index of disease status of patients with colorectal carcinoma. *Br J Cancer* 1998; 78:454-459.
 - 41 Imai Y, Kubota Y, Yamamoto S, Tsuji K, Shimatani M, Shibatani N, et al: Neutrophils enhance invasion activity of human cholangiocellular carcinoma and hepatocellular carcinoma cells: an in vitro study. *J Gastroenterol Hepatol* 2005; 20:287-293.
 - 42 Daveau M, Scotte M, Francois A, Coulouarn C, Ros G, Tallet Y, et al: Hepatocyte growth factor, transforming growth factor α , and their receptor as combined markers of prognosis in hepatocellular carcinoma. *Mol Carcinog* 2003; 36:130-141.
 - 43 Shimomura T, Miyazawa K, Hiraoka H, Naka D, Morimoto Y, Kitamura N: Activation of hepatocyte growth factor by two homologous proteases, blood-coagulation factor XIIa and hepatocyte growth factor activator. *Eur J Biochem* 1995; 229:257-261.
 - 44 Nagino M, Ando M, Kamiya J, Uesaka T, Sano T, Nimura Y: Liver regeneration after major hepatectomy for biliary cancer. *Br J Surg* 2001; 88:1084-1091.
 - 45 Yamanaka N, Okamoto E, Kawamura E, Kato T, Oriyama T, Fujimoto J, et al: Dynamics of normal and injured human liver regeneration after hepatectomy as assessed on the basis of computed tomography and liver function. *Hepatology* 1993; 18:79-85.
 - 46 Weiss YG, Bellin L, Kim PK, Andrejko KM, Haaxma CA, Raj N, et al: Compensatory hepatic regeneration after mild, but not fulminant, intraperitoneal sepsis in rats. *Am J Physiol Gastrointest Liver Physiol* 2001; 280:G968-973.

Comparison of Indocyanine Green Clearance with Child's-Pugh Score and Hepatic Histology: A Multivariate Analysis

Sandeep Mukherjee MD, Mary AM Rogers PhD, Borys Buniak MD

Liver Gallbladder Pancreas Center, Section of Gastroenterology and Hepatology, Department of Medicine
State University of New York Upstate Medical University, Syracuse, New York, USA

Corresponding Author: Sandeep Mukherjee, MD, Section of Gastroenterology and Hepatology
983285 Nebraska Medical Center, Omaha, NE 68198-3285, USA

Tel: +1 402 559 8859, Fax: +1 402 559 3434, E-mail: smukherj@unmc.edu

KEY WORDS:

Child's-Pugh score; Indocyanine green; Hepatic histology; Multivariate analysis

ABBREVIATIONS:

Indocyanine Green (ICG); Deoxyribonucleic Acid (DNA); Percentage Disappearance Rate (PDR); Ribonucleic Acid (RNA); Histological Activity Index (HAI); Polymerase Chain Reaction (PCR)

ABSTRACT

Background/Aims: Indocyanine green clearance, measured by percentage disappearance rate, detects alterations in liver function and may be used as a non-invasive determinant of hepatic reserve. The aims of this study were to compare liver histology and Child's-Pugh score with percentage disappearance rate and determine which variables correlated with PDR.

Methodology: Child's-Pugh score, liver function tests, liver biopsies and indocyanine green testing (0.5mg/kg) were performed in 102 consecutive patients with cirrhosis of diverse etiologies. Indocyanine green concentration was determined using spectrophotometric analysis (806nm) and plotted logarithmically with Michaelis-Menten kinetics to calculate the percentage disappearance rate. Liver biopsies were graded using the modified Knodell score to

obtain a histological activity index.

Results: In bivariate analysis, percentage disappearance rate significantly correlated with Child's-Pugh score, albumin, bilirubin, prothrombin time and histological activity index. Albumin, prothrombin time and histological activity index were independent predictors of percentage disappearance rate in the final model (albumin $p < 0.01$, prothrombin time $p < 0.046$, histological activity index $p = 0.033$), accounting for 46.2% of variability in percentage disappearance rate measurements.

Conclusions: Percentage disappearance rates correlated with Child's-Pugh scores in this series of cirrhotic patients. However, 46.2% of its variability was accounted for by albumin, prothrombin time and histological activity index.

INTRODUCTION

Chronic liver disease is a major health problem worldwide and for patients with end-stage liver disease, liver transplantation remains the treatment of choice. The Child's-Pugh score has traditionally been used to assess the severity of underlying disease although it is not a sensitive indicator of liver dysfunction (1). Patients may deteriorate before appreciable changes in their scores are recognized and to predict the need for liver transplantation or monitor hepatic response to therapy, a complementary test for quantification of liver function may be helpful.

Indocyanine green (ICG) is a tricarbo-cyanine dye strongly bound to albumin with high hepatic clearance by first order kinetics (70-90%). ICG does not undergo biotransformation or extrahepatic metabolism, is excreted unchanged in bile and does not undergo renal excretion (2,3). It has been used to measure liver blood flow and to quantify hepatic reserve in individuals with chronic liver disease (4-7). The aims of this study were to compare ICG clearance with the Child's-Pugh score and hepatic histology and determine which of these variables correlated with ICG clearance.

METHODOLOGY

Patients: One hundred and two consecutive patients with cirrhosis were referred to a University hospital liver center for evaluation between 1995 and 1997. The mean age of patients was 49.7 ± 13.4 years, of whom 51% were male and 49% female. Causes of cirrhosis in the study population are reported in **Table 1**.

Laboratory investigations: Blood samples were obtained for measurement of standard hepatic biochemical evaluations which included alanine aminotransaminase (ALT), aspartate aminotransaminase (AST), alkaline phosphatase, albumin, prothrombin time and total bilirubin. Hepatitis B surface antibody, hepatitis B e antigen, hepatitis B e antibody, hepatitis B core antibody, hepatitis B surface antibody were tested using commercially available enzyme-linked or radioimmunoassay test kits (Abbott Laboratories, North Chicago, IL). Hepatitis B DNA was detected using semiquantitative hybridization assay (Abbott Laboratories, North Chicago, IL). Hepatitis C antibody was obtained using second-generation enzyme-linked immunoassay (Abbott Laboratories, North

Selection Criteria for Reduction Hepatectomy in Multiple Advanced Hepatocellular Carcinoma

NAOTO GOTOHDA¹, HIROSHI ISHII², MASARU KONISHI¹,
TOSHIO NAKAGOHRI¹, SHINICHIRO TAKAHASHI¹, JUNJI FURUSE²,
MASAHIRO YOSHINO² and TAIRA KINOSHITA¹

¹Department of Hepatobiliary Pancreatic Surgery
and ²Department of Hepatobiliary Pancreatic Medical Oncology,
National Cancer Center Hospital East, Chiba, 277-8577, Japan

Abstract. *Background:* Few studies have compared the prognostic impact of reduction hepatectomy (RH) for advanced hepatocellular carcinoma (HCC) with that of non-surgical treatment or curative hepatectomy. *Patients and Methods:* The treatment outcome of 30 RH patients was compared to that of two control groups: control group A, including 71 patients who underwent curative hepatectomy; control group B, including 106 patients who did not receive definitive local therapy or best supportive care. *Results:* In patients with tumor extension in >50% of the liver, 1-year survival rates for patients according to treatment (RH, control A and B) were 50%, 90% and 11% and 3-year survival rates were 42%, 60% and 0%, respectively. There was no significant difference between RH and control A, but there was a significant difference between RH and control B. *Conclusion:* RH could be recommended to patients with multiple advanced HCC extending to >50% of the whole liver.

Hepatic resection is the only treatment that offers a hope of cure with long-term survival in patients with HCC. However, the indications for radical hepatectomy for hepatocellular carcinoma (HCC) remain limited for patients with multiple intrahepatic metastases (1). If hepatectomy with curative intent for multiple advanced HCCs cannot be performed, the possibility of volume reduction hepatectomy (RH) for those HCCs is often considered, with the aim of decreasing target tumor cells so that effective post-operative treatment can be carried out. The efficacy of RH for advanced HCC has been

reported by several studies (2-5), and we have also performed RH for multiple advanced HCC.

The aim of this study was to evaluate the long-term results of RH for patients with multiple HCC, and to validate the indications for RH compared to other treatments and optimize patient selection.

Patients and Methods

From July 1992 to May 2005, 1,001 patients with a diagnosis of HCC with no distant (extrahepatic) metastases were examined consecutively at the National Cancer Center Hospital East, Japan. Of these HCC patients, 30 underwent RH. Their treatment outcome was compared retrospectively with that of two control groups selected among the remaining 971 patients with a similar background to the RH patients. The selection criteria (background factors) were: total bilirubin <2.4 mg/dl, albumin >2.7 g/dl, no ascites, Child-Pugh score <9 points, multiple tumors, main tumor >44 mm in diameter, Okuda stage I or II (6), and BCLC (Barcelona Clinic Liver Cancer) stage B or C (7). Control A included 71 patients who underwent hepatectomy with curative intent. Control B included 106 patients who did not receive definitive local therapy (hepatectomy, percutaneous ablation therapy or proton beam radiotherapy) or best supportive care.

The indications for hepatectomy were determined according to criteria based on tumor extension and hepatic functional reserve (8, 9). As our treatment strategy for advanced HCC, whether or not hepatectomy with curative intent could be performed was considered first. When hepatectomy with curative intent could not be performed, the possibility of RH was considered. RH was defined at our institute as resection of the main tumor or main tumor plus satellite tumors around the main tumor, with satellite tumors in the remnant liver classified as unresectable. If this surgical treatment for advanced HCC was not feasible, radiotherapy (including proton beam radiotherapy) or transcatheter arterial chemoembolization/infusion chemotherapy (TACE/TAI) tolerance was considered. If liver function and portal vein flow were poor due to tumor thrombi, only TAI with farmorubicin was administered. If TAI was not feasible, best supportive care was considered. After RH, TACE or TAI treatment was repeated until the residual tumor had completely disappeared on CT.

Correspondence to: Naoto Gotohda, MD, Department of Hepatobiliary Pancreatic Surgery, National Cancer Center Hospital East, 6-5-1 Kashiwanoha, Kashiwa, Chiba, 277-8577, Japan. Tel: +81 4 7133 1111, Fax: +81 4 7131 4724, e-mail: ngotohda@east.ncc.go.jp

Key Words: Reduction hepatectomy, advanced hepatocellular carcinoma, selection criteria.

Table I. Patient characteristics.

	RH	Control A	p-value	Control B	p-value
No. of patients	30	71		106	
Patient factors					
Age (yr)	60 (27-76)	65 (28-84)	0.013*	63 (39-86)	0.010*
Gender (M/F)	29/1	62/9	0.274 [§]	90/16	0.119 [§]
HBsAg positive	7	18	1.000 [§]	14	0.250 [§]
Anti-HCVAb positive	15	40	1.000 [§]	74	0.052 [§]
Total bilirubin (mg/dl)	0.9 (0.4-2.3)	0.8 (0.4-1.8)	0.390*	1.0 (0.4-2.2)	0.100*
Albumin (g/dl)	3.6 (2.8-4.3)	3.7 (2.8-4.7)	0.088*	3.5 (2.8-4.3)	0.351*
AFP (pg/ml)	540 (1.8-970,000)	71 (1.7-380,000)	0.126*	400 (1.7-350,000)	0.937*
Child-Pugh stage (A/B)	26/4	63/8	0.746 [§]	83/23	0.438 [§]
Okuda classification (I/II)	18/12	60/11	0.010[§]	84/22	0.054 [§]
BCLC staging (B/C)	14/16	56/15	0.002 [§]	74/32	0.029 [§]
Tumor factors					
Size of main tumor (mm)	120 (45-200)	80 (45-220)	0.001*	70 (45-150)	<0.001*
Vascular invasion positive	16	15	0.002[§]	32	0.029[§]
Tumor extension in >50%	12	10	0.007[§]	11	<0.001*

Values represent median (range). RH: reduction hepatectomy, HBsAg: hepatitis B surface antigen, Anti-HCVAb: anti-hepatitis C virus antibody, AFP: α -fetoprotein, BCLC staging: Barcelona Clinic Liver Cancer staging. Statistical significance was analyzed by Chi-square test[§] and Mann-Whitney's U-test*.

Results

The patient characteristics are listed in Table I. As patients with similar background factors to those in RH patients were selected, no significant differences were found between the RH, control A and B groups. Cumulative 1-year survival rates for patients in RH, control A and B groups were 29%, 85% and 61%, and cumulative 3-year survival rates were 17%, 63% and 16%, respectively. Significant differences were found in survival between control group A and RH, and between control group A and control group B.

In cases where the main tumor was greater than 100 mm in diameter, cumulative 1- and 3-year survival rates for RH, control A and B groups were 32 and 26%, 82 and 45% and 31 and 0%, respectively. Significant differences were found between control A and RH, and between control A and control B group. In patients with vascular invasion, 1-year survival rates for RH, control A and B groups were 13%, 73% and 29%, while 3-year survival rates were 13%, 38% and 3%, respectively. In patients with tumor extension in >50% of the liver, 1-year survival rates for RH, control A and B groups were 50%, 90% and 11%, while 3-year survival rates were 42%, 60% and 0%, respectively. There was no significant difference between RH and control group A, but there was a significant difference between RH and control B (Figure 1). The characteristics of patients with tumor extension in >50% of the liver are shown in Table II. Total bilirubin was higher in control group B compared to RH group, while the other patient factors were similar

between the three groups. Although larger HCCs were included in the RH group compared to control B group, the vascular invasion factor was similar.

Discussion

The background factors of patients treated with RH in our hospital were investigated first, using various prognostic scoring systems (Okuda stage, CLIP (10), JIS (11), Tokyo score (12) and BCLC). There were no RH patients with Okuda stage III or BCLC stage A or D. As the RH patient distribution by the Okuda and BCLC staging systems was characteristic and not biased, they were used to unify the background factors of HCC patients. Several reports, after validation of the various prognostic staging systems for HCC, have indicated that the BCLC staging system provided the best independent prediction of survival (13, 14).

Patient factors were unified as much as possible, and two comparison groups were used (curative hepatectomy group and other local therapy group). After overall survival rates in each group had been compared, RH was definitively inferior to curative hepatectomy, and similar to other local therapies (including best supportive care).

However, although all patients in this study had multiple HCCs, three tumor factors (tumor size, vascular invasion and tumor extension) were further advanced in RH group compared to control A and B groups. These tumor factors were then estimated separately, and survival rates in each group with the same tumor condition were compared.

Table II. Characteristics of patients with tumor extension in >50% of liver.

	RH	Control A	<i>p</i> -value	Control B	<i>p</i> -value
No. of patients	12	10		11	
Patient factors					
Age (yr)	60 (27-74)	66 (45-70)	0.147*	61 (50-70)	0.389*
Gender (M/F)	12/0	9/1	0.455§	8/3	0.093§
HBsAg positive	3	2	1.000§	2	1.000§
Anti-HCVAb positive	4	4	1.000§	8	0.100§
Total bilirubin (mg/dl)	0.9 (0.5-1.2)	0.8 (0.5-1.0)	0.230*	1.4 (0.8-2.2)	0.003*
Albumin (g/dl)	3.5 (2.8-4.0)	3.7 (3.0-4.3)	0.186*	3.6 (2.8-4.0)	0.687*
AFP (pg/ml)	6,400 (2.5-240,000)	290 (1.7-380,000)	0.210*	19,000 (3.5-92,000)	0.854*
Child-Pugh stage (A/B)	10/2	8/2	1.000§	6/5	0.193§
Okuda classification (I/II)	0/12	0/10		0/11	
BCLC staging (B/C)	5/7	7/3	0.231§	6/5	0.684§
Tumor factors					
Size of main tumor (mm)	160 (130-200)	130 (53-220)	0.079*	130 (90-150)	0.006*
Vascular invasion positive	7	3	0.231§	5	0.684§

Values represent median (range). RH: reduction hepatectomy, HBsAg: hepatitis B surface antigen, Anti-HCVAb: anti-hepatitis C virus antibody, AFP: α -fetoprotein, BCLC staging: Barcelona Clinic Liver Cancer staging. Statistical significance was analyzed by Chi-square test[§] and Mann-Whitney's *U*-test*.




Figure 1. Kaplan-Meier estimated survival curves according to treatment in patients with tumor extension in >50% of liver. *RH: reduction hepatectomy. Statistical significance was analyzed by log-rank test.

After those comparisons, the survival benefit of RH for advanced multiple HCCs with tumor extension in >50% of the liver was found to be equivalent to that of curative hepatectomy and superior to that of other local therapy (including best supportive care). Since RH is intended to decrease the tumor volume in the liver, this result could suggest that RH is more effective for prolonged survival benefit if tumor extension in the liver is larger.

With regard to the effectiveness of RH for advanced HCC, Yamamoto *et al.* (4) reported that RH was effective in patients with fewer and smaller tumors left in the residual liver and no extrahepatic metastasis. However, only surgically treated patients were compared in the report. Wakabayashi *et al.* (5) compared two groups of patients (RH group and non-surgical group), and showed that RH had survival benefit. However, their background factors differed from ours, and the authors commented that unintended patient selection bias may have affected the results of the study. Although our evaluated patient factors were selected retrospectively, as shown in Table II, our treatment outcome was compared in groups selected to have equivalent factors, and, thus, our results were considered to be unbiased.

In conclusion, the present study suggested that RH for multiple advanced HCC with tumor extension in >50% of the liver might hold promise of survival benefit.

References

- 1 The Liver Cancer Study Group of Japan: Predictive factors for long term prognosis after partial hepatectomy for patients with hepatocellular carcinoma in Japan. *Cancer* 74: 2772-2780, 1994.
- 2 Asahara T, Itamoto T, Katayama K *et al*: Indications for palliative reduction surgery for hepatocellular carcinoma with multiple intrahepatic metastases. *Hiroshima J Med Sci* 47: 115-120, 1998.
- 3 Inoue K, Nakamura T, Kinoshita T *et al*: Volume reduction surgery for advanced hepatocellular carcinoma. *J Cancer Res Clin Oncol* 130: 362-366, 2004.
- 4 Yamamoto K, Takenaka K, Kawahara N *et al*: Indications for palliative reduction surgery in advanced hepatocellular carcinoma. The use of a remnant tumor index. *Arch Surg* 132: 120-123, 1997.
- 5 Wakabayashi H, Ushiyama T, Ishimura K *et al*: Significance of reduction surgery in multidisciplinary treatment of advanced hepatocellular carcinoma with multiple intrahepatic lesions. *J Surg Oncol* 82: 98-103, 2003.
- 6 Okuda K, Ohtsuki T, Obata H *et al*: Natural history of hepatocellular carcinoma and prognosis in relation to treatment. Study of 850 patients. *Cancer* 56: 918-928, 1985.
- 7 Llovet J M, Bru C, and Bruix J: Prognosis of hepatocellular carcinoma: the BCLC staging classification. *Semin Liver Dis* 19: 329-338, 1999.
- 8 Imamura H, Seyama Y, Kokudo N *et al*: One thousand fifty-six hepatectomies without mortality in 8 years. *Arch Surg* 138: 1198-1206; discussion 1206, 2003.
- 9 Makuuchi M, Kosuge T, Takayama T *et al*: Surgery for small liver cancers. *Semin Surg Oncol* 9: 298-304, 1993.
- 10 The Cancer of the Liver Italian Program (CLIP) Investigators: A new prognostic system for hepatocellular carcinoma: a retrospective study of 435 patients. *Hepatology* 28: 751-755, 1998.
- 11 Kudo M, Chung H, Haji S *et al*: Validation of a new prognostic staging system for hepatocellular carcinoma: the JIS score compared with the CLIP score. *Hepatology* 40: 1396-1405, 2004.
- 12 Tateishi R, Yoshida H, Shiina S *et al*: Proposal of a new prognostic model for hepatocellular carcinoma: an analysis of 403 patients. *Gut* 54: 419-425, 2005.
- 13 Cillo U, Bassanello M, Vitale A *et al*: The critical issue of hepatocellular carcinoma prognostic classification: which is the best tool available? *J Hepatol* 40: 124-131, 2004.
- 14 Marrero J A, Fontana R J, Barrat A *et al*: Prognosis of hepatocellular carcinoma: comparison of 7 staging systems in an American cohort. *Hepatology* 41: 707-716, 2005.

Received June 23, 2006

Revised October 12, 2006

Accepted October 17, 2006

EXTRAHEPATIC PORTAL VEIN OCCLUSION WITHOUT RECURRENCE AFTER PANCREATICOUDENECTOMY AND INTRAOPERATIVE RADIATION THERAPY

SHUICHI MITSUNAGA, M.D.,* TAIRA KINOSHITA, M.D.,* MITSUHIKO KAWASHIMA, M.D.,†
MASARU KONISHI, M.D.,* TOSHIO NAKAGOHRI, M.D.,* SHINICHIRO TAKAHASHI, M.D.,*
AND NAOTO GOTOHDA, M.D.*

Departments of *Hepatobiliary-Pancreatic Surgery and †Radiation Oncology, National Cancer Center Hospital East, Kashiwa, Japan

Purpose: Although there are no definitive studies that characterize the survival benefit of intraoperative radiation therapy (IORT), the therapy does not seem to produce significant complication. In our institution, pancreaticoduodenectomy (PD) and IORT are often complicated by the development of extrahepatic portal vein occlusion (EHPO). The aim of this study was to characterize the phenomenon of EHPO after PD and IORT.

Methods and Materials: Between September 1992 and December 2001, 107 patients received macroscopic curative PD for periampullary disease in our institution. IORT (radiation dose: 20 Gy) was performed in 53 of these patients. Criteria for diagnosis of EHPO were as follows: (1) computerized tomography findings of occlusive extrahepatic portal vein, (2) symptoms of portal hypertension, and (3) confirmation to exclude tumor recurrence from origin of EHPO, because this study examined whether EHPO was a complication of PD and IORT.

Results: EHPO was diagnosed in 12 patients. Among patient and operative variables, IORT was the only statistically significant factor associated with a diagnosis of EHPO ($p = 0.0052$). The median developed time to EHPO and overall survival after surgery in EHPO patients were 358 days and 2,562 days, respectively. Eight patients (67%) with EHPO died during the follow-up period. At 5 years after therapy, EHPO was diagnosed in 67% of survivors who had received IORT.

Conclusions: Patients undergoing IORT and PD have a relatively high incidence of EHPO, and patients who develop postoperative EHPO have poor prognoses. © 2006 Elsevier Inc.

Intraoperative radiation therapy, Extrahepatic portal vein occlusion, Late complication, Pancreaticoduodenectomy.

INTRODUCTION

Intraoperative radiation therapy (IORT) has been used for the treatment of pancreatic cancer. Although there are no definitive studies that characterize the survival benefit of IORT, the therapy does not seem to produce significant disorder in normal tissue (1–3). In our institution, pancreaticoduodenectomy (PD) and IORT are often complicated by the development of extrahepatic portal vein occlusion (EHPO). Although studies have demonstrated that EHPO can be induced by inflammation of periportal structures (4), abdominal surgery (5, 6), or pylephlebitis (7), there are no case reports or systematic studies describing the development of EHPO after IORT. Thus, the goal of the present study was to characterize the phenomenon of EHPO after pancreaticoduodenectomy and intraoperative radiation therapy. Cases of patients undergoing pancreaticoduodenectomy with or without IORT for various diagnoses were

retrospectively reviewed, and their clinical courses, including development of EHPO, were investigated.

METHODS AND MATERIALS

Between September 1992 and December 2001, 107 patients (median age, 62 years; 39 females, 68 males) underwent macroscopic curative PD for periampullary disease at our institution.

Patient characteristics and early outcomes, clinical course after surgery, and follow-up computerized tomography (CT) findings were retrospectively reviewed. These parameters included age, gender, preoperative biliary drainage, preoperative total bilirubin, requirements for additional resection, requirement for regional lymph node resection, complicated resection of the portal vein, use of IORT, surgical complications, postoperative hospital stay duration, and pathologic diagnosis (Table 1).

Additional resection included colectomy in 2 patients and partial hepatectomy in 4 patients when solitary small liver metastasis was

Reprint requests to: Shuichi Mitsunaga, M.D., Department of Hepatobiliary-Pancreatic Surgery, National Cancer Center Hospital East, Kashiwanoha, 6-5-1, Kashiwa, Chiba 277-8577 Japan. Tel: (+81) 471-34-6855; Fax: (+81) 471-34-6865; E-mail:

smitsuna@east.ncc.go.jp

Received June 20, 2005, and in revised form Aug 15, 2005. Accepted for publication Aug 15, 2005.

Table I. Patient characteristics and early outcomes in EHPO-positive and -negative patients

	All patients	EHPO-negative group	EHPO-positive group	<i>p</i> value
Patient number (<i>n</i>)	107	95	12	NA
Age (y)				
Median	62	64	56	0.0616
Range	25–81	25–81	47–67	
Gender (male/female) (<i>n</i>)	68/39	60/35	8/4	0.9359
Preoperative biliary drainage (<i>n</i>)				
(–)	59	53	6	0.7174
(+)	48	42	6	
Preoperative total bilirubin level (mg/dL)				
Median	1.5	1.5	1.65	0.7468
Range	0.2–16.2	0.2–16.2	0.5–6.5	
Pathologic diagnosis (<i>n</i>)				
Invasive malignancy	91	81	10	0.9492
Pancreatic cancer	53	49	4	
Lower bile duct cancer	16	15	1	
Cancer of the ampulla of Vater	17	14	3	
Duodenal cancer	4	3	1	
Retroperitoneal leiomyosarcoma	1	0	1	
Benign or minimally invasive	16	14	2	
Chronic pancreatitis	6	4	2	
Intraductal papillary mucinous tumor	5	5	0	
Islet cell tumor	2	2	0	
Solid and cystic tumor	1	1	0	
Serous cyst adenoma	1	1	0	
Lower bile duct adenoma	1	1	0	
Additional resection (<i>n</i>)				
Partial colectomy	2	2	0	0.5328
Partial hepatectomy	4	3	1	0.9338
Regional lymph node dissection (<i>n</i>)				
(–)	5	5	0	0.9297
(+)	102	90	12	
Resection of portal vein (<i>n</i>)				
(–)	79	71	8	0.8019
(+)	28	24	4	
Combined with intraoperative radiation therapy (<i>n</i>)				
(–)	54	53	1	0.0052*
(+)	53	42	11	
Surgical complications (<i>n</i>)				
Abdominal abscess	15	13	2	0.8722
Pancreatorrhea	53	46	7	0.7330
Bile leakage	5	5	0	0.9354
Intra-abdominal hemorrhage	8	7	1	0.6436
Postoperative hospital stay (d)				
Median	50	48	50.5	0.4218
Range	23–139	23–139	32–113	

Abbreviation: EHPO = extrahepatic portal vein occlusion.

* $p < 0.05$ chi square test.

detected on the surface of the liver intraoperatively. All 4 patients undergoing partial hepatectomy were included in the final analysis, because the procedure did not affect the portal venous system. A total of 102 patients underwent regional lymph node dissection. Complicated portal vein resection was performed in 28 patients. Perioperative portal vein thrombosis was not found. Surgical complications included abdominal abscess ($n = 15$), pancreatorrhea ($n = 53$), bile leakage ($n = 5$), and intra-abdominal bleeding ($n = 8$). All 107 patients were enrolled in this study. None of the patients received adjuvant or neoadjuvant therapy.

Intraoperative radiation therapy was performed in 53 patients, as follows. During operation, an acrylic cylinder of 6–8 cm diameter was introduced to encompass the tumor bed. The irradiation field

included the celiac axis, superior mesenteric artery, and the surgical manipulated portal vein, but excluded the stump of the pancreas. In each case, 20 Gy with electron-beam energies of 6–12 MeV was delivered.

Pathologic diagnosis was established according to the World Health Organization classification (8), and patients were assigned to one of two groups based on pathologic diagnosis: the invasive malignancy group ($n = 93$) and the benign or minimally invasive group ($n = 14$). The invasive malignancy group included those patients with pancreatic cancer, bile duct cancer, cancer of the ampulla of Vater, duodenal cancer, and retroperitoneal leiomyosarcoma. The benign or minimally invasive group included those patients with chronic pancreatitis, intraductal papillary mucinous-