

of 197 minutes from symptom onset in the 114 patients. Intra-arterial UK infusion was started at the mean of 227 minutes from symptom onset in the 56 patients in the UK group. One patient in the UK group was not treated with intra-arterial UK infusion because of technical reason. One patient in the control group underwent intra-arterial UK infusion because of human error. The UK and control groups were well matched with regard to the mean entry NIHSS score, presumed source of embolus, sex, and age (Table 1). Early ischemic change, defined as subtle sulcal effacement or loss of gray-white matter distinction in the insular cortex, frontal and temporal opercula, or lenticular nuclei, was observed on the initial CT scan in 27 patients in both UK and control groups.

Primary and secondary endpoints at 90 days

The primary and secondary end points at 90 days are summarized in Table 2. For the primary endpoint, 28 patients (49.1%) in the UK group had mRS 0 to 2 at 90 days compared with 22 patients (38.6%) in the control group ($p=0.345$). Nevertheless, for the secondary endpoints, the UK group showed a 15.8% to 21.1% absolute increase in excellent functional outcome (mRS 0-1) compared to the control group at 90 days ($p=0.045$). There were significantly more patients with NIHSS 0 or 1 at 90 days in the UK group than the control group ($p=0.017$). The UK group included more patients with Barthel index of 95 or more than the control group, but the difference did not reach at the significance level ($p=0.065$). The 90-day cumulative mortality was 5.3% in the UK group and 3.5% in the control group ($p=0.647$). Death was attributed to medical complications associated with the initial stroke in 2 patients, hemorrhagic transformation in 1, recurrence in 1, and malignant tumor in 1 (Table 3). There was no significant difference between the groups.

Forty patients received the full dose of UK (600,000 IU) and 16 patients received UK less than 600,000 IU. Partial or complete recanalization was achieved in 42 of 57 patients (73.7%) treated with intra-arterial UK infusion. Mechanical clot disruption was performed in 39 of the 57 patients (68%). Recanalization was complete in 3 patients, partial 50 % and more in 27, partial less than 50% in 12, and not achieved in 15.

Severe adverse events

Severe adverse events are summarized in Table 4. Intracerebral hemorrhage within 24 hours of treatment occurred in 5 patients (9%) in the UK group, including one patient with hematoma caused by perforation by the guide wire, compared with 1 patient (2%) in the control group (not significant). Severe brain edema followed by air embolism in one patient in the UK group. There was no significant difference in the frequency of severe adverse events between the UK and control groups.

Early ischemic change expanded from that of the inclusion criteria was observed on the initial CT scan in 9 of 114 patients (7.9%). Among these patients, 5 patients received intra-arterial UK therapy and 2 patients developed hemorrhagic transformation within 24 hours.

Discussion

The MELT Japan suggested beneficial effects of intra-arterial fibrinolysis in patients with acute stroke caused by MC artery occlusion of less than 6 hours' duration, although the result was negative with respect to the primary endpoint. The absolute increase of excellent neurological outcome, a secondary endpoint, was 19.3 % as mRS, which was better than other previous intra-venous rt-PA trials.^{2,15} Compared with the patients receiving conventional therapy, patients treated with intra-arterial UK infusion at a mean of 227 minutes from symptom onset were 86% more likely to achieve complete recovery at 90 days. The 19% absolute increase in favorable outcome with intra-arterial UK ($p=0.045$) indicates that one in 6 patients treated with intra-arterial UK will benefit.

The MELT Japan restricted patient selection to MC artery occlusion because of the poor natural history¹ and the most frequent location in patients with severe stroke of less than 6 hours' duration.¹⁶ In contrast to the National Institute of Neurological Disorders and Stroke (NINDS) rt-PA trial,² the MELT Japan used slight or no neurological disability (mRS score 0 to 2) as the primary outcome measure and complete recovery (mRS score 0 and 1) as the secondary outcome measure because of the anticipated baseline of high severity of stroke in patients with MC artery occlusion. It is debatable what is the optimal cut-off point of the functional outcome scale for dichotomization of outcomes in an acute stroke trial. The PROACT II trial defined mRS score of 2 or less (dichotomised for independency) as the favorable outcome, and showed the efficacy of intra-arterial fibrinolysis.¹³ The European Cooperative Acute Stroke Study II, defining mRS of 0 and 1 (dichotomised for complete recovery or minimal deficit) as the favorable outcome, failed to show the efficacy of intra-venous fibrinolysis, although an analysis of mRS score of 2 or less as the secondary end point demonstrated the efficacy of thrombolysis.¹⁵ On the contrary, the NINDS study demonstrated significant increase in the percentage of patients with mRS 0 and 1.²

Various reasons can be proposed for the failure to show efficacy as measured by mRS in this study. Firstly, the 38.6% of patients with mRS score 0 to 2 in the control group was relatively higher than the percentage expected in the present trial. The PROACT II trial reported that only 25% of the control group demonstrated mRS 2 or less. Administration of neuroprotective agents such as the free radical scavengers,¹⁷ which are widely used in Japan for improving functional outcome in patients with acute ischemic stroke, may also have affected this finding. Secondly, the number of enrolled patients was too small to recognize any difference between the UK and control groups at the primary endpoint in the present study. In the PROACT II study, 13 patients allocated to the proUK group did not receive fibrinolytic therapy and 5 control patients received fibrinolytic agent. If these 18 patients are removed from the intention-to-treat analysis, the difference is not significant. Thirdly, the CT criteria of the MELT Japan are different from other trials.^{2,3,13-15,18} The MELT Japan did not apply the one third rule of early ischemic CT sign which was widely used.^{2,3,13-15,18-20} Instead, patients were

included with no or subtle early ischemic signs only in the insular cortex, frontal and temporal opercula, or lenticular nuclei. Such an area may cover less than one third of the MC artery distribution area, so the patients in the MELT Japan might have had less severe stroke than in other randomized acute stroke trials. Thereby, the mean NIHSS score was smaller to those in other studies^{2,3,14,15}

To demonstrate the pharmacological effect of UK, the MELT Japan prohibited mechanical clot disruption except for the guidewire technique. The recanalization rate of 73.7% in the present study was better than that of 66% reported previously.¹³ The PROACT II study prohibited any mechanical procedure. Therefore, mechanical fibrinolysis may be one of the reasons for the present favorable recanalization rate. Advances in catheter technology, imaging techniques, mechanical clot removal, and more potent fibrinolytic agents should lead to faster and more complete recanalization and potentially even better patient outcomes.

The total ICH rates were consistent with those previously reported in patients with embolic stroke.²¹⁻²³ There was no significant difference in the rates of ICH throughout the trial period, which may reflect delayed recanalization in the control group with hemorrhagic transformation, whereas the higher early rate with UK reflected drug-induced recanalization and reperfusion hemorrhage.

The NINDS study supports the use of intravenous rt-PA within 3 hours of stroke, but limited data suggest that intravenous rt-PA may be relatively ineffective in the subset of patients with MC artery occlusion. An angiography-based trial demonstrated that successful recanalization with intravenous infusion of rt-PA was significantly less likely in complete MC artery occlusion than peripheral lesions.¹⁶ Baseline NIHSS score greater than 10 and hyperdense MC artery sign on CT (significant MC artery occlusion) both predict poor clinical outcome for patients treated with intravenous rt-PA at less than 3 hours from symptom onset.²⁴ Intra-arterial fibrinolysis may have greater potential in both the recanalization rate and the wider therapeutic window. Intra-arterial fibrinolysis may be feasible after intravenous rt-PA in patients with persistent MC artery occlusion.²⁵⁻²⁷

The therapeutic window in acute human ischemic stroke has been very controversial.^{28,29} Recent diffusion and perfusion magnetic resonance studies suggest that as many as two thirds of patients with acute MC artery distribution stroke have brain tissue at risk even 24 hours after stroke onset, but the clinical relevance of these observations is uncertain.³⁰ The MELT Japan demonstrated that the therapeutic window may extend to at least 6 hours for a significant number of patients with major stroke due to MC artery occlusion although it was clarified by the secondary endpoint.

Conclusion

Local intra-arterial UK fibrinolysis therapy for patients with acute MC artery occlusion at less than 6 hours after onset is safe and more likely to lead to complete recovery at 90 days. This should be reserved, however, because the trial was aborted prematurely and the primary end point did not reach

at the statistical significance. Further studies with a larger number of patients are needed to confirm these results.

Appendix: The MELT Japan Study Group

Steering Committee

A. Ogawa, Iwate Medical University School of Medicine; E. Mori, Tohoku University Graduate School of Medicine; K. Minematsu, National Cardiovascular Center; W. Taki, Mie University Graduate School of Medicine; A. Takahashi, Tohoku University Graduate School of Medicine; S. Nemoto, Jichi Medical University School of Medicine; S. Miyamoto, Kyoto University Graduate School of Medicine; M Sasaki, Iwate Medical University School of Medicine

Allocation Center

S. Nemoto, Jichi Medical University School of Medicine; T. Ueno; Teikyo University School of Medicine

Publicity Committee

S. Uchiyama, Tokyo Women's Medical University; S. Nemoto, Jichi Medical University School of Medicine; N. Sakai, Kobe City General Hospital; T. Ueno; Teikyo University School of Medicine

Protocol and Analysis Committee

E. Mori, Tohoku University Graduate School of Medicine; K. Minematsu, National Cardiovascular Center; Y. Yoneda; Hyogo Brain and Heart Center; T. Kiuchi, Tokyo University Graduate School of Medicine; S. Nemoto, Jichi Medical University School of Medicine; T. Ueno; Teikyo University School of Medicine

Imaging Technique Standardization Committee

W. Taki, Mie University Graduate School of Medicine; T. Terada, Wakayama Rosai Hospital; M. Maeda, Mie University Graduate School of Medicine; T. Ueda, Yokohama Stroke and Brain Center; T. Takada, St. Marianna University School of Medicine; T. Ueno; Teikyo University School of Medicine; J. Nakagawara, Nakamura Memorial Hospital

Fibrinolysis Technique Standardization Committee

A. Takahashi, Tohoku University Graduate School of Medicine; T. Hyogo, Nakamura Memorial Hospital; N. Sakai, Kobe City General Hospital; T. Takada, St. Marianna University School of Medicine; H. Matsuoka, National Hospital Organization Kyusyu Cardiovascular Center; T. Terada, Wakayama Rosai Hospital; M. Ezura, Tohoku University Graduate School of Medicine

Film Reading Committee

J. Nakagawara, Nakamura Memorial Hospital; Y. Okada, National Hospital Organization Kyushu

Medical Center; M. Sasaki, Iwate Medical University School of Medicine

Report Organization Committee

K. Minematsu, National Cardiovascular Center; T. Takada, St. Marianna University School of Medicine; A. Takahashi, Tohoku University Graduate School of Medicine; T. Hyogo, Nakamura Memorial Hospital; N. Sakai, Kobe City General Hospital; H. Matsuoka, National Hospital Organization Kyusyu Cardiovascular Center; T. Terada, Wakayama Rosai Hospital; M. Ezura, Tohoku University Graduate School of Medicine; K. Sato, Tokushima University School of Medicine

Protocol Surveillance Committee

N. Hashimoto, Kyoto University Graduate School of Medicine; S. Nagahiro, Tokushima University; M. Suzuki, Yamaguchi University; S. Miyamoto, Kyoto University Graduate School of Medicine; T. Hyogo, Nakamura Memorial Hospital; M. Ezura, Tohoku University Graduate School of Medicine; S. Nemoto, Jichi Medical University; S. Miyachi, Nagoya University Graduate School of Medicine; M. Sugita, Yamanashi University; T. Terada, Wakayama Rosai Hospital; N. Sakai, Kobe City General Hospital; K. Sugi, Okayama University; K. Sato, Tokushima University; A. Hyodo, Ryukyus University

Safety Monitoring Committee

T. Yamaguchi, National Cardiovascular Center; K. Yamada, Nagoya City University School of Medicine; S. Uchiyama, Tokyo Women's Medical University; S. Matsumoto, Hiroshima University School of Medicine

Secretariat

T. Inoue, K. Ogasawara, N. Mori, K. Hatayama, A. Sakurada, Iwate Medical University; H. Konno, Iwate Prefectural Hanamaki Kose Hospital; C. Sakai, Advanced Medical Center

MELT investigators (see supplement)

References

1. Mohr JP. Some clinical aspects of acute stroke. Excellence in clinical stroke award lecture. *Stroke*. 1997;28:1835-1839.
2. Tissue plasminogen activator for acute ischemic stroke. The National Institute of Neurological Disorders and Stroke rt-PA Stroke Study Group. *N Engl J Med*. 1995;333:1581-1587.
3. Yamaguchi T, Mori E, Minematsu K, Nakagawara J, Hashi K, Saito I, Shinohara Y. Alteplase at 0.6 mg/kg for acute ischemic stroke within 3 hours of onset: Japan Alteplase Clinical Trial (J-ACT). *Stroke*. 2006;37:1810-1815.
4. Reed SD, Cramer SC, Blough DK, Meyer K, Jarvik JG. Treatment with tissue plasminogen activator and inpatient mortality rates for patients with ischemic stroke treated in community hospitals. *Stroke*. 2001;32:1832-1840.
5. del Zoppo GJ, Pessin MS, Mori E, Hacke W. Thrombolytic intervention in acute thrombotic and embolic stroke. *Semin Neurol*. 1991;11:368-384.
6. Barr JD, Mathis JM, Wildenhain SL, Wechsler L, Jungreis CA, Horton JA. Acute stroke intervention with intraarterial urokinase infusion. *J Vasc Interv Radiol*. 1994;5:705-713.
7. Bendszus M, Urbach H, Ries F, Solymosi L. Outcome after local intra-arterial fibrinolysis compared with the natural course of patients with a dense middle cerebral artery on early CT. *Neuroradiology*. 1998;40:54-58.
8. del Zoppo GJ, Ferbert A, Otis S, Bruckmann H, Hacke W, Zyroff J, Harker LA, Zeumer H. Local intra-arterial fibrinolytic therapy in acute carotid territory stroke. A pilot study. *Stroke*. 1988;19:307-313.
9. Gonner F, Remonda L, Mattle H, Sturzenegger M, Ozdoba C, Lovblad KO, Baumgartner R, Bassetti C, Schroth G. Local intra-arterial thrombolysis in acute ischemic stroke. *Stroke*. 1998;29:1894-1900.
10. Mori E, Tabuchi M, Yoshida T, Yamadori A. Intracarotid urokinase with thromboembolic occlusion of the middle cerebral artery. *Stroke*. 1988;19:802-812.
11. Qureshi AI, Suri MF, Shatla AA, Ringer AJ, Fessler RD, Ali Z, Guterman LR, Hopkins LN. Intraarterial recombinant tissue plasminogen activator for ischemic stroke: An accelerating dosing regimen. *Neurosurgery*. 2000;47:473-476; discussion 477-479.
12. Zeumer H, Hacke W, Ringelstein EB. Local intraarterial thrombolysis in vertebrobasilar thromboembolic disease. *AJNR Am J Neuroradiol*. 1983;4:401-404.
13. Furlan A, Higashida R, Wechsler L, Gent M, Rowley H, Kase C, Pessin M, Ahuja A, Callahan F, Clark WM, Silver F, Rivera F. Intra-arterial prourokinase for acute ischemic stroke. The PROACT II study: a randomized controlled trial. Prolyse in Acute Cerebral Thromboembolism. *JAMA*. 1999;282:2003-2011.

14. Hacke W, Kaste M, Fieschi C, Toni D, Lesaffre E, von Kummer R, Boysen G, Bluhmki E, Hoxter G, Mahagne MH, Hennerice M. Intravenous thrombolysis with recombinant tissue plasminogen activator for acute hemispheric stroke. The European Cooperative Acute Stroke Study (ECASS). *JAMA*. 1995;274:1017-1025.
15. Hacke W, Kaste M, Fieschi C, von Kummer R, Davalos A, Meier D, Larrue V, Bluhmki E, Davis S, Donnan G, Schneider D, Diez-Tejedor E, Trouillas P. Randomised double-blind placebo-controlled trial of thrombolytic therapy with intravenous alteplase in acute ischaemic stroke (ECASS II). Second European-Australasian Acute Stroke Study Investigators. *Lancet*. 1998;352:1245-1251.
16. del Zoppo GJ, Poeck K, Pessin MS, Wolpert SM, Furlan AJ, Ferbert A, Alberts MJ, Zivin JA, Wechsler L, Busse O, Greenlee RJ, Brass L, Mohr JP, Feldmann E, Hacke W, Kase CS, Biller J, Gress D, Otis SM. Recombinant tissue plasminogen activator in acute thrombotic and embolic stroke. *Ann Neurol*. 1992;32:78-86.
17. Edaravone Acute Infarction Study Group. Effect of a novel free radical scavenger, edaravone (MCI-186), on acute brain infarction. Randomized, placebo-controlled, double-blind study at multicenters. *Cerebrovasc Dis*. 2003;15:222-229.
18. del Zoppo GJ, Higashida RT, Furlan AJ, Pessin MS, Rowley HA, Gent M. PROACT: a phase II randomized trial of recombinant pro-urokinase by direct arterial delivery in acute middle cerebral artery stroke. PROACT Investigators. Prolyse in Acute Cerebral Thromboembolism. *Stroke*. 1998;29:4-11.
19. Silver B, Demaerschalk B, Merino JG, Wong E, Tamayo A, Devasenapathy A, O'Callaghan C, Kertesz A, Young GB, Fox AJ, Spence JD, Hachinski V. Improved outcomes in stroke thrombolysis with pre-specified imaging criteria. *Can J Neurol Sci*. 2001;28:113-119.
20. Wardlaw JM, Mielke O. Early signs of brain infarction at CT: Observer reliability and outcome after thrombolytic treatment--systematic review. *Radiology*. 2005;235:444-453.
21. Okada Y, Yamaguchi T, Minematsu K, Miyashita T, Sawada T, Sadoshima S, Fujishima M, Omae T. Hemorrhagic transformation in cerebral embolism. *Stroke*. 1989;20:598-603.
22. Hornig CR, Dorndorf W, Agnoli AL. Hemorrhagic cerebral infarction--a prospective study. *Stroke*. 1986;17:179-185.
23. Lodder J, Krijne-Kubat B, Broekman J. Cerebral hemorrhagic infarction at autopsy: Cardiac embolic cause and the relationship to the cause of death. *Stroke*. 1986;17:626-629.
24. Tomsick T, Brott T, Barsan W, Broderick J, Haley EC, Spilker J, Khoury J. Prognostic value of the hyperdense middle cerebral artery sign and stroke scale score before ultraearly thrombolytic therapy. *AJNR Am J Neuroradiol*. 1996;17:79-85.
25. Pannell R, Black J, Gurewich V. Complementary modes of action of tissue-type plasminogen

- activator and pro-urokinase by which their synergistic effect on clot lysis may be explained. *J Clin Invest*. 1988;81:853-859.
26. Lewandowski CA, Frankel M, Tomsick TA, Broderick J, Frey J, Clark W, Starkman S, Grotta J, Spilker J, Khoury J, Brott T. Combined intravenous and intra-arterial r-TPA versus intra-arterial therapy of acute ischemic stroke: Emergency Management of Stroke (EMS) Bridging Trial. *Stroke*. 1999;30:2598-2605.
 27. Ueda T, Sakaki S, Nochide I, Kumon Y, Kohno K, Ohta S. Angioplasty after intra-arterial thrombolysis for acute occlusion of intracranial arteries. *Stroke*. 1998;29:2568-2574.
 28. Baron JC, von Kummer R, del Zoppo GJ. Treatment of acute ischemic stroke. Challenging the concept of a rigid and universal time window. *Stroke*. 1995;26:2219-2221.
 29. Fisher M, Garcia JH. Evolving stroke and the ischemic penumbra. *Neurology*. 1996;47:884-888.
 30. Fisher M, Prichard JW, Warach S. New magnetic resonance techniques for acute ischemic stroke. *JAMA*. 1995;274:908-911.

Table 1. Characteristics of Treated Patients

	UK	Control
n	57	57
Male sex	64.90%	64.90%
Age, y	66.9 ± 9.3	67.3 ± 8.5
NIHSS	14.7 ± 5.1	14.2 ± 4.2
Cardioembolic stroke	88%	83%
Time from onset to hospitalization, min	68 ± 46	79 ± 52
Time from onset to allocation, min	199 ± 61	206 ± 54
BP at hospitalization	154/88	145/82
violation of BP	14	6
Occlusion location		
proximal M1	14 (24.6%)	18 (31.6%)
distal M1	25 (43.9%)	24 (42.1%)
M2	18 (31.6%)	15 (26.3%)

UK, urokinase; NIHSS, National Institutes of Health Stroke Scale; BP, blood pressure. All values are mean ±SD or literal values.

Table 2. Clinical Outcomes at 90 days

Scores	UK (n=57)	Control (n=57)	P value
Modified Rankin Scale			
0 to 2	28 (49.1%)	22 (38.6%)	0.345
0 and 1	24 (42.1%)	13 (22.8%)	0.045
NIHSS 0 and 1	20 (35.1%)	8 (14.0%)	0.017
Barthel Index 95 or greater	28 (49.1%)	19 (33.3%)	0.087

Table 3. Mortality within 90 days

	UK (n=57)	Control (n=57)	P value
All mortality	3 (5.3%)	2 (3.5%)	0.647
intracerebral hemorrhage	1	0	
recurrence	1	0	
malignant tumor	1	0	
renal failure	0	1	
sepsis	0	1	

Table 4. Complications

	UK	Control
Intracerebral hemorrhage	5 (9%)*	1 (2%)
Brain edema	3 (5%)#	2 (4%)
Recurrence	4 (7%)	1 (2%)
Other	3 (5%)	5 (9%)

* including one patient caused by perforation by the guidewire.

including one patient followed by air embolism.

Figure legends

Fig.1

Distribution of patients for entry and randomization. UK indicates urokinase.

Fig.2

Representative computed tomography (CT) scans at baseline (left) and 24 hours after symptom onset (right) in a 72-year-old man with left hemiparesis allocated to the Control group. Baseline CT was performed 2 hours after symptom onset.

Left: Subtle early ischemic sign was seen in the right insular cortex (arrow) and lenticular nuclei (arrowhead).

Right: New infarctions are present in the insular cortex and lenticular nuclei.

Supplement

Secondary endpoint at 24 hours and 30 days

The UK group had a significantly better neurological outcome by mRS at 30 days ($p=0.009$), as NIHSS at 24 hours ($p=0.022$), and 30 days ($p=0.017$), and as Barthel Index at 30 days ($p=0.016$).

MELT Investigators (listed in order of number of patients allocated)

Kohnan Hospital: M. Ezura, Y. Matsumoto, R. Kondo, H. Nomura, A. Ohnuma, R. Mochizuki, S.

Sato, A. Kikuchi, T. Kinpara, K. Jin

Kobe City General Hospital: N. Sakai, T. Kuroiwa,

National Cardiovascular Center: K. Minematsu, T. Takada, J. Kasuya, C. Yokota

St.Mary's Hospital: Y. Saku, T. Kuwashiro, K. Annoura, H. Takaba, T. Imamura, T. Inoue, Y.

Fukushima, K. Fukuda, T. Yonekura, Y. Wakisaka, J. Jinnai, N. Santa

Kokura Memorial Hospital: I. Nakahara, M. Tanaka, T. Higashi, Y. Iwamuro, H. Shii

Nakamura Memorial Hospital: T. Hyogo, K. Hayase, J. Nakagawara, T. Kataoka, K. Sakoh

Sendai Medical Center: Y. Sakurai, S. Nishimura, I. Suzuki, J. Onodera

Yokohama Stroke and Brain Center: T. Ueda, R. Obama

Gifu University: S. Yoshimura, T. Iwama, N. Sakai, Y. Kaku, T. Inuzuka

Iwate Prefectural Ofunato Hospital: T. Yamanome, S. Ohmama

Yamagata Prefectural Central Hospital: N. Takeda, Y. Ibuchi, T. Kumagai, A. Inoue, Y. Yoneoka, H.

Shirai, T. Yamakawa, K. Kikuchi, Y. Jinbo

Iwate Medical University: A. Ogawa, T. Inoue, H. Konno, C. Sakai, M. Sasaki, M. Yamagata, R.

Kondo,

Yamaguchi University: M. Suzuki, S. Kato, H. Ishihara, H. Adachi, H. Nakayama, K. Negoro

Chiba Cardiovascular Center: J. Ono, S. Matsuda, S. Hirai, K. Honma

Hurukawa City Hospital: M. Ohba, M. Yoshida, H. Shimizu, J. Higuchi

Iwate Prefectural Central Hospital: H. Seki, T. Sugawara

Nagasaki University: N. Kitagawa, K. Hayashi, S. Shirabe, M. Motomura

Akita Red Cross Hospital: K. Nishimaki

International Medical Center Japan: Y. Uesaka, K. Hasuo, T. Hideyama, M. Ubano

Kanto Medical Center NTT EC: A. Morita, I. Shirouzu, C. Ochiai, T. Nishihara, M. Shoujima, K.

Arasaki

Kyoto Prefectural University of Medicine: K. Ohwada, K. Yoshikawa, M. Nawata, Y. Nagakane, Y.

Inoue

Kyoto University: K. Nakazawa, T. Sato, A. Sadatoh, T. Ohta, H. Tomimoto

Nippon Medical School Chiba Hokuso Hospital: S. Kominami

St.Luke's International Hospital: Y. Numaguchi, M. Oka, N. Kobayashi

Tokyo Metropolitan Police Hospital: H. Sato, S. Nemoto, T. Shirai

Wakayama Medical University: T. Itakura, T. Terada, H. Matsumoto, M. Tsuura, O. Masuo, T.

Tsumoto, K. Kubo, M. Ueno, H. Higashioka

Keio University: T. Kawase, S. Nogawa, Y. Mine, R. Ueda, S. Onozuka, T. Hayashi, H. Sato, T.

Denboh, S. Momoshima, M. Katayama, H. Takeda

Kinki University: F. Akai, N. Fukawa, A. Kimoto,

Kishiwada City Hospital: K. Yamashita, N. Matsuoka

Musashino Red Cross Hospital: S. Watabiki, H. Miake

Okayama University: K. Sugiu, K. Abe, T. Hayashi

Osaka Medical College: J. Deguchi, T. Kuroiwa, M. Yamada, H. Nakajima

Tokushima University Hospital: K. Sato, M. Nishimura

Ehime University School of Medicine: T. Ohnishi, S. Fukumoto Y. Mitsunaga

Kyorin University: Y. Konishi, I. Saito, A. Shimada, H. Yazaki, T. Ishiko

Nihon University: Y. Katayama, T. Kanoh, T. Hirayama

Osaka Koseinenkin Hospital: H. Ohtsuki, T. Fujita, H. Terakawa

Shakaihoken Kobe Central Hospital: K. Matsumoto, M. Shirato, E. Yoshino

Toho University Ohashi Medical University: S. Iwabuchi, T. Yokouchi, H. Nemoto

Aizawa Hospital: H. Nagashima, T. Koyama, K. Kitazawa, A. Watanabe, T. Iwashita, H. Hara

Hyogo Brain and Heart Center: T. Kawaguchi, Y. Shibata, Y. Yoneda, M. Iwakura, H. Komatsu, H.

Yamasaki, T. Uehara

Iwate Prefectural Kuji Hospital: K. Miura, T. Okuguchi

Mie University : W. Taki, F. Asakura, H. Sakaida, N. Toma, Y. Kokubo

National Hospital Organization Kyusyu Cardiovascular Center: H. Matsuoka, K. Uchimura,

Nishi-Kobe Medical Center: K. Fujita, S. Katayama, C. Yanagihara

Osaka City General Hospital: T. Yasui, M. Komiyama, Y. Matsusaka, H. Ikeda, K. Tokumoto

University of the Ryukyus: A. Hyodo, K. Isa

Saiseikai Utsunomiya Hospital: M. Nakatsukasa, H. Kagami, A. Imai

Fukushima Red Cross Hospital: Y. Watanabe

Kobe University: M. Nakamura, A. Fujita, Y. Nakamura, T. Kondo

Kumamoto University: Y. Kai, T. Todaka

Nagoya University: S. Miyachi

Nara Medical University: K. Kichikawa, T. Wada, H. Nakagawa, K. Fujimoto

Osaka City Unuversity: A. Nishio, M. Nishikawa, T. Tsutada

Tenri Hospital: F. Isaka, K. Higuchi, T. Suenaga

Tokyo Women's Medical University: T. Hori, T. Higa, F. Yamane, Y. Ono, K. Miyazaki

Yamagata University: T. Kayama, R. Kondo, Y. Kokubo

Fig. 1





Fig. 2



R