

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Table 1. Sensitivity and Specificity of Wavelet Analysis of RF IVUS Signals for Detecting Lipid-laden Plaque

Histology		Wavelet Coefficient	
		≥ 0.6	< 0.6
Lipid core (+)	In Vitro (n=29)	24/29(83%)	5/29(17 %)
	In Vivo (n=16)	13/16(81%)	3/16(19 %)
Lipid core (-)	In Vitro (n=56)	10/56(18%)	46/56(82%)
	In Vivo (n=13)	2/13(15%)	11/13(85%)

Sex Differences in Early Mortality of Patients Undergoing Primary Stenting for Acute Myocardial Infarction

Masami Kosuge, MD; Kazuo Kimura, MD; Sunao Kojima, MD*; Tomohiro Sakamoto, MD*; Masaharu Ishihara, MD**; Yujiro Asada, MD†; Chuwa Tei, MD††; Shunichi Miyazaki, MD‡; Masahiro Sonoda, MD‡‡; Kazufumi Tsuchihashi, MD§; Masakazu Yamagishi, MD‡; Yoshihiko Ikeda, MD§§; Mutsunori Shirai, MD***; Hisatoyo Hiraoka, MD†††; Takeshi Inoue, MD†††; Fumio Saito, MD§§§; Hisao Ogawa, MD*; on behalf of the Japanese Acute Coronary Syndrome Study (JACSS) Investigators

Background Limited information exists regarding the impact of gender on in-hospital outcome after primary stenting for acute myocardial infarction (AMI).

Methods and Results A total of 2,981 patients (790 women and 2,191 men) participated in the study who were admitted within 24 h after symptom onset and underwent emergency primary stenting for AMI. Compared with men, women were significantly older; had higher incidences of hypertension, diabetes mellitus, hyperlipidemia, Killip class ≥ 2 , and cardiogenic shock; had a higher blood glucose level and a lower serum creatinine level on admission. Other baseline characteristics, including the incidences of ST-segment elevation AMI, anterior infarction, 3-vessel disease, initial or final Thrombolysis in Myocardial Infarction (TIMI) flow grade did not significantly differ between the sexes. The in-hospital mortality rate was significantly higher in women than in men (9.4% vs 5.2%, $p < 0.001$). On multivariate analysis, age, Killip class, blood glucose level, serum creatinine level, and final TIMI grade were independent predictors of in-hospital death, but female gender was not (odds ratio 1.01, $p = 0.69$).

Conclusions Our findings suggest that in patients undergoing primary stenting for AMI, women have higher in-hospital mortality than men, but female gender itself is not independently associated with increased in-hospital mortality after adjustment for baseline differences. (Circ J 2006; 70: 217–221)

Key Words: Myocardial infarction; Stents; Survival; Women

A number of studies have addressed sex-related differences in outcomes in patients with acute myocardial infarction (AMI). A few studies have reported similar or lower mortality rates after AMI in women than in men^{1,4} but most have concluded that mortality is higher in women irrespective of reperfusion modality.^{5–12} The reasons for poorer outcomes in women remain unclear. Increased mortality in women might be

partially explained by their higher age at presentation and higher risk profiles. Some^{5–8} but not all^{9–12} studies have shown that female gender itself is independently associated with increased mortality after adjustment for baseline differences. One potential explanation for the persistence of increased mortality after risk adjustment is that women frequently receive less aggressive treatment for AMI than men.^{7,13,14} Primary balloon angioplasty, compared with thrombolytic therapy, has been shown to improve outcomes for women, but mortality remains high.^{2,10} Limited information exists regarding the impact of sex on outcomes after contemporary interventional techniques, such as stent implantation, for AMI. We therefore analyzed a database from a large, retrospective, multicenter observational study of patients with AMI who underwent emergency primary stenting to assess the outcomes of women compared with those of men.

Methods

Patients

The patients included in the current study were selected from those enrolled into the Japan Acute Coronary Syndrome Study,¹⁵ a retrospective, observational multicenter trial. Between January 2001 and December 2003, patients who were admitted to 35 participating hospitals in Japan within 48 h after the onset of AMI were studied. A diagnosis of AMI required at least 2 of the following characteristics: typical chest pain persisting for 30 min or longer,

(Received September 15, 2005; revised manuscript received December 8, 2005; accepted December 21, 2005)

Division of Cardiology, Yokohama City University Medical Center, Yokohama, *Department of Cardiovascular Medicine, Kumamoto University School of Medicine, Kumamoto, **Department of Cardiology, Hiroshima City Hospital, Hiroshima, †First Department of Pathology, Miyazaki University Medical School, Miyazaki, ††The First Department of Internal Medicine, Kagoshima University Faculty of Medicine, Kagoshima, ‡Division of Cardiology, Department of Internal Medicine, National Cardiovascular Center, Suita, ‡‡The Second Department of Cardiology, National Hospital Kyusyu Cardiovascular Center, Kagoshima, §Second Department of Internal Medicine, Sapporo Medical University School of Medicine, Sapporo, §§Department of Pathology, National Cardiovascular Center, Suita, ***Department of Microbiology, School of Medicine, Yamaguchi University, Ube, †††Department of Internal Medicine and Molecular Science, Graduate School of Medicine, Osaka University, Suita, ††††Division of Cardiology, Oita National Hospital, Oita and †††††Department of Cardiology, Nihon University Surugadai Hospital, Tokyo, Japan
Mailing address: Kazuo Kimura, MD, The Division of Cardiology, Yokohama City University Medical Center, 4-57 Urafune-cho, Minami-ku, Yokohama 232-0024, Japan. E-mail: c-kimura@urahp.yokohama-cu.ac.jp

Table 1 Baseline Characteristics

	Women (n=790)	Men (n=2,191)	p value
Age (years)	73±10	65±11	<0.001
Body surface area (m ²)	1.47±0.14	1.71±0.16	<0.001
Time from symptom onset to admission (h)	4.9±5.0	4.5±5.0	0.09
Killip ≥2 on admission	173 (22%)	331 (15%)	<0.001
Cardiogenic shock on admission	68 (9%)	142 (7%)	0.045
Previous infarction	78 (10%)	245 (11%)	0.31
Previous angina	308 (39%)	799 (37%)	0.21
Diabetes mellitus	273 (35%)	672 (31%)	0.044
Hyperlipidemia	293 (37%)	716 (33%)	0.025
Hypertension	506 (64%)	1,136 (52%)	<0.001
Smoking	131 (17%)	1,248 (57%)	<0.001
Blood glucose level on admission (mmol/L)	10.7±5.1	9.8±4.2	<0.001
Serum creatinine on admission (mg/dl)	0.8±0.7	1.0±0.9	<0.001
ST-segment elevation	688 (87%)	1,930 (88%)	0.46
Peak creatine kinase (IU/L)	2,829±2,676	3,418±3,387	<0.001

Data are presented as mean values ± SD or percentages of patients.

Table 2 Angiographic Findings

	Women (n=790)	Men (n=2,191)	p value
Number of diseased vessels			0.97
1	445 (57%)	1,246 (57%)	
2	240 (30%)	657 (30%)	
3	105 (13%)	288 (13%)	
Infarct-related artery			0.29
LAD	384 (49%)	1,024 (47%)	
RCA	283 (36%)	840 (38%)	
LCX	107 (13%)	287 (13%)	
LMT	16 (2%)	33 (2%)	
Bypass graft	0	7 (0.3%)	
TIMI flow grade 0 at initial CAG	512 (65%)	1,456 (67%)	0.40
Final TIMI flow grade ≥2	758 (96%)	2,090 (95%)	0.51
Final TIMI flow grade 3	714 (90%)	1,967 (87%)	0.63

Data are presented percentages of patients.

LAD, left anterior descending artery; RCA, right coronary artery; LCX, left circumflex coronary artery; LMT, left main trunk; TIMI, Thrombolysis in Myocardial Infarction.

ischemic electrocardiographic changes, and a peak creatine kinase level equivalent to more than twice the upper limit of normal. A total of 4,432 patients fulfilled the following criteria: 1 admission within 24 h after the onset of AMI; 2 coronary angiography performed immediately after admission; and 3 availability of a detailed clinical history. Reperfusion therapy was performed in 3,635 patients (82%). Of these 3,635 patients, we studied 2,981 (82%) who underwent stenting of the infarct-related artery. The study protocol was reviewed and approved by the ethical committee of each participating hospital.

Definitions

Diabetes mellitus was defined as a fasting glucose concentration of ≥ 7.0 mmol/L, a blood glucose concentration of ≥ 11.0 mmol/L on a 75-g, 2-h oral glucose tolerance test, or the use of anti-diabetic treatment. Hypertension was defined as a history of a systolic blood pressure of ≥ 140 mmHg, a diastolic pressure of ≥ 90 mmHg, or the use of anti-hypertensive treatment. Hyperlipidemia was defined as a fasting total cholesterol concentration of ≥ 220 mg/dl, a fasting triglyceride concentration of ≥ 150 mg/dl, or the use of anti-hyperlipidemic treatment. Preinfarction angina was defined as the presence of typical chest pain occurring at rest or during exercise and persisting for less than 30 min, within 24 h before the onset of AMI.

Coronary Angiography and Coronary Intervention

Coronary angiography was performed immediately after admission. The perfusion status of the infarct-related artery was assessed according to the Thrombolysis in Myocardial Infarction (TIMI) study classification.¹⁶ The recanalization method was left to the physicians' discretion. Stenting was done in all patients in whom the procedure was feasible. Final TIMI flow grade was assessed on the basis of the final angiograms obtained at admission.

Statistical Analysis

Data are expressed as means ± SD. Patients with and those without preinfarction angina were compared by using unpaired t-tests. Differences in prevalence were assessed by chi-square tests. A probability value of < 0.05 was considered to indicate a statistically significant difference. Multiple logistic regression analysis was used to examine determinants of in-hospital mortality. Analyzed variables included age, sex, hypertension, diabetes mellitus, prior infarction, preinfarction angina, body surface area, time to admission, Killip class, infarct location, blood glucose level and serum creatinine level on admission, occlusion status at the culprit lesion, number of diseased vessels, and final TIMI flow grade. Odds ratios and 95% confidence intervals were calculated. Data were analyzed with the use of SPSS software (Release 10, SPSS, Chicago, IL, USA).




Fig 1. Overall, in-hospital mortality was significantly higher in women (black bars) than in men (white bars). The rates of cardiac death and stroke were also significantly higher in women than in men.

Results

Baseline Characteristics

Of the 2,981 subjects, 2,191 (73%) were men and 790 (27%) were women. The baseline characteristics of the subjects are presented according to sex in Table 1. Women were more likely to be elderly and had a lower body surface area; higher incidences of hyperlipidemia, diabetes mellitus, hypertension, Killip class ≥ 2 , and cardiogenic shock; and a lower frequency of smoking. There was a trend toward a longer time from symptom onset to admission in women, but the difference with men did not reach statistical significance. The frequency of previous infarction, preinfarction angina, and ST-segment elevation did not differ between the sexes. On admission the blood glucose level was significantly higher and the peak creatine kinase level and serum creatinine level were significantly lower in women than in men.

Angiographic Findings

Angiographic findings of the patients are shown in Table 2. There were no significant differences between women and men in the number or distribution of diseased coronary vessels, including 3-vessel disease and left main coronary disease. The initial and final TIMI flow grades did not differ between the sexes.

In-Hospital Outcomes

Clinical outcomes in hospital are shown in Fig 1. In-hospital mortality was significantly higher in women than

Table 3 Multivariate Analysis of Factors Associated With In-Hospital Mortality in All Patients

Variable	Odds ratio (95%CI)	p value
Age	1.08 (1.04–1.12)	<0.001
Female	1.01 (0.47–2.04)	0.69
Hypertension	1.09 (0.42–1.29)	0.29
Diabetes mellitus	1.04 (0.54–2.02)	0.40
Prior infarction	1.81 (0.92–3.56)	0.09
Absence of preinfarction angina	1.55 (0.87–2.76)	0.14
Body surface area	0.68 (0.43–1.06)	0.09
Time to admission	1.03 (0.98–1.09)	0.23
Killip class ≥ 2	6.43 (3.67–11.3)	<0.001
Anterior infarction	1.56 (0.89–2.71)	0.11
Blood glucose level on admission	1.01 (1.01–1.30)	0.049
Serum creatinine level on admission	1.45 (1.24–1.69)	<0.001
TIMI flow grade 0 at initial CAG	1.21 (0.68–2.16)	0.51
Multivessel disease	1.31 (0.74–2.33)	0.30
Final TIMI flow grade	0.60 (0.41–0.85)	0.005

CI, confidence interval; CAG, coronary angiography; TIMI, Thrombolysis in Myocardial Infarction.

men. Of the patients who died in the hospital (74 women and 115 men), 51 women and 98 men died of cardiac causes. Women had a higher rate of stroke. Multivariate analysis showed that age, Killip class, blood glucose level or serum creatinine level on admission, and final TIMI grade were independent predictors of in-hospital death, whereas female gender was not (Table 3). Multivariate analysis according to sex showed that age, Killip class, blood glucose level or serum creatinine level on admission,

Table 4 Multivariate Analysis of Factors Associated With In-Hospital Mortality According to Sex

Variable	Women		Men	
	Odds ratio (95%CI)	P	Odds ratio (95%CI)	P
Age	1.11 (1.03–1.19)	0.004	1.07 (1.03–1.11)	0.001
Hypertension	1.09 (0.39–3.03)	0.86	0.62 (0.31–1.26)	0.19
Diabetes mellitus	1.03 (0.30–2.91)	0.91	1.10 (0.47–2.58)	0.82
Prior infarction	0.95 (0.22–4.19)	0.95	2.33 (1.06–5.16)	0.036
Absence of preinfarction angina	1.51 (0.24–1.83)	0.42	1.52 (0.74–3.17)	0.26
Body surface area	0.80 (0.06–13.3)	0.59	0.19 (0.01–2.50)	0.21
Time to admission	1.08 (1.01–1.17)	0.037	0.99 (0.91–1.07)	0.76
Killip class ≥ 2	5.90 (2.24–15.6)	<0.001	6.60 (3.24–13.4)	<0.001
Anterior infarction	1.74 (0.64–4.76)	0.28	1.68 (0.83–3.38)	0.15
Blood glucose level on admission	1.17 (1.01–20.6)	0.038	1.20 (1.05–23.7)	0.046
Serum creatinine level on admission	1.59 (1.15–11.3)	0.005	1.42 (1.17–1.71)	<0.001
TIMI flow grade 0 at initial CAG	0.95 (0.36–2.53)	0.91	1.36 (0.64–2.86)	0.42
Multivessel disease	1.05 (0.41–2.65)	0.94	1.51 (0.72–3.20)	0.28
Final TIMI flow grade	0.46 (0.23–0.90)	0.024	0.62 (0.40–0.97)	0.037

CI, confidence interval; CAG, coronary angiography; TIMI, Thrombolysis in Myocardial Infarction.

and final TIMI grade were independent predictors of in-hospital death in both women and men (Table 4).

Discussion

The present study showed that in patients undergoing primary stenting for AMI, women had a higher rate of in-hospital mortality than men did. However, multivariate analysis showed that female gender itself was not an independent predictor of in-hospital mortality.

Similar to previous studies,¹⁻¹¹ women were older than men by 8 years on average and had a lower body surface area, a higher Killip class, and higher incidences of hypertension, diabetes mellitus, and hyperlipidemia. On coronary angiography, the extent of underlying coronary atherosclerosis, evaluated on the basis of the number of diseased vessels, did not differ between women and men, consistent with the findings of previous investigations.^{2,5,10,12} The current study also found that baseline and post-procedural TIMI flow grades were similar in women and men. Several studies have similarly shown that the TIMI flow grade before and after primary angioplasty for AMI did not differ between women and men.^{2,10,17} In contrast, Lansky et al found that baseline and post-procedural TIMI grade 3 flows were better in women than in men.¹¹

Mortality was higher in women than in men, consistent with the results of most previous studies.⁵⁻¹² This poorer outcome in women was most likely related to the facts that women were older than men, had higher incidences of coronary risk factors such as hyperlipidemia, diabetes mellitus, and hypertension, as well as higher incidences of Killip class ≥ 2 , and cardiogenic shock. These and other factors such as a higher blood glucose level on admission might have negatively affected outcome. Several studies have shown that women have poorer outcomes than men;^{9-11,14} however, after adjustment for other baseline characteristics, female sex itself was not an independent risk factor for increased mortality. Those studies therefore concluded that a higher age or more adverse risk profiles in women contributed to the poorer outcomes. However, several studies reported that female gender itself was an independent predictor of increased mortality, even after adjusting for other baseline characteristics.⁵⁻⁸ These inconsistent results suggest that methodological, as well as biologic factors, must be considered when interpreting the impact of gender on survival after AMI.

Women have been reported to be significantly less likely to receive thrombolytic therapy, percutaneous transluminal coronary angioplasty (PTCA), and coronary artery bypass grafting.^{6,7,13} The less frequent use of reperfusion therapy in women might be related to their higher mortality. Furthermore, randomized studies have shown that percutaneous coronary interventions are more effective reperfusion strategies than intravenous thrombolysis.¹⁸ In the GUSTO II-B PTCA substudy, Tamis-Holland et al suggested that women derived a larger absolute benefit from primary PTCA than from thrombolytic therapy as compared with men.¹⁰ Stone et al reported that primary PTCA reduced the risk of intracranial bleeding and improved survival in women enrolled in the PAMI trial.⁹ In the present study, all patients underwent primary stenting, thereby eliminating possible differences in mortality caused by treatment bias.

Recent studies have investigated the influence of sex on outcome in patients receiving primary coronary intervention for AMI, but whether female gender itself is an inde-

pendent predictor of increased mortality after AMI remains controversial. Vakili et al reported that after correcting for age and baseline risk differences, women undergoing primary coronary intervention for a first AMI have a higher in-hospital mortality rate than men.⁵ Lansky et al have shown that female gender is not an independent determinant of death at 1 year and attributed the higher mortality rate in women after interventional treatment for AMI to differences in body size and clinical risk factors.¹¹ Mehilli et al showed that women with AMI who received percutaneous coronary intervention have outcomes similar to those of men, despite more adverse risk profiles.³ Differences in study design, entry criteria, and length of follow-up make it difficult to compare results in different studies. The present study included high-risk patients who are usually excluded from clinical trials, such as those who are elderly and have shock, as well as a high proportion of patients with diabetes mellitus (31%). In addition, we included the blood glucose level and serum creatinine level on admission in our risk analysis. These factors have been shown to be associated with an increased risk of death after AMI,¹⁹⁻²² but were not always included in risk analysis in previous studies assessing the impact of sex on outcomes after AMI.

Study Limitations

Several important limitations need to be considered when interpreting our results. First, this was a retrospective, observational, non-randomized study. However, our database was relatively large and included patients treated at hospitals of various sizes and settings, making it more representative of current practice patterns than previous single-site databases or randomized trials. Second, our data did not include information on treatment with aspirin, statin, β -blockers, angiotensin converting enzyme inhibitors and angiotensin II receptor antagonists or on the door-to-balloon time, all factors shown to influence mortality from AMI. Furthermore, angiographic data such as the number of stenosed lesions, lesion length, and reference diameter were not obtained. However, in the current study, body surface area, a surrogate for coronary vessel size, was lower in women than in men. Another limitation was that cardiac and non-cardiac causes of death were not fully examined. Further investigations are necessary to determine why women with AMI are more likely to have poorer outcomes compared with men. In addition, prospective studies are needed to verify the effects of gender on clinical outcomes after primary stenting for AMI.

Acknowledgments

The authors gratefully acknowledge that this study was supported by a Research Grant for Cardiovascular Disease (14C-4) from the Ministry of Health, Labour and Welfare.

References

1. Johansson S, Bergstrand R, Ulyenstam G, Vedin A, Wilhelmsson C, Wedel H, et al. Sex differences in preinfarction characteristics and long-term survival among patients with myocardial infarction. *Am J Epidemiol* 1984; **119**: 610-623.
2. Stone GW, Grines CL, Browne KF, Marco J, Rothbaum D, O'Keefe J, et al. Comparison of in-hospital outcome in men versus women treated by either thrombolytic therapy or primary coronary angioplasty for acute myocardial infarction. *Am J Cardiol* 1995; **75**: 987-992.
3. Mehilli J, Kastrati A, Dirschinger J, Pache J, Seyfarth M, Blasini R, et al. Sex-based analysis of outcome in patients with acute myocardial infarction treated predominantly with percutaneous coronary intervention. *JAMA* 2002; **287**: 210-215.

4. Merrilees MA, Scott PJ, Romo M. Five-year survival of 728 patients after myocardial infarction: A community study. *Br Heart J* 1980; 43: 176-183.
5. Vakili BA, Kaplan RC, Brown DL. Sex-based differences in early mortality of patients undergoing primary angioplasty for first acute myocardial infarction. *Circulation* 2001; 104: 3034-3038.
6. Kudenchuk PJ, Maynard C, Martin JS, Wirkus M, Weaver WD. Comparison of presentation, treatment, and outcome of acute myocardial infarction in men versus women (the Myocardial Infarction Triage and Intervention Registry). *Am J Cardiol* 1996; 78: 9-14.
7. Chandra NC, Ziegelstein RC, Rogers WJ, Tiefenbrunn AJ, Gore JM, French WJ, et al. Observations of the treatment of women in the United States with myocardial infarction: A report from the National Registry of Myocardial Infarction-I. *Arch Intern Med* 1998; 158: 981-988.
8. Marso SP, Gowda M, O'Keefe JH, Coen MM, McCallister BD, Giorgi LV, et al. Improving in-hospital mortality in the setting of an increasing risk profile among patients undergoing catheter-based reperfusion for an acute myocardial infarction without cardiogenic shock. *J Invasive Cardiol* 2003; 15: 711-716.
9. Dittrich H, Gilpin E, Nicod P, Cali G, Henning H, Ross J Jr. Acute myocardial infarction in women: Influence of gender on mortality and prognostic variables. *Am J Cardiol* 1988; 62: 1-7.
10. Tamis-Holland JE, Palazzo A, Stebbins AL, Slater JN, Boland J, Ellis SG, et al. Benefits of direct angioplasty for women and men with acute myocardial infarction: Results of the Global Use of Strategies to Open Occluded Arteries in Acute Coronary Syndromes Angioplasty (GUSTO II-B) Angioplasty Substudy. *Am Heart J* 2004; 147: 133-139.
11. Lansky AJ, Pietras C, Costa RA, Tsuchiya Y, Brodie BR, Cox DA, et al. Gender differences in outcomes after primary angioplasty versus primary stenting with and without abciximab for acute myocardial infarction: Results of the Controlled Abciximab and Device Investigation to Lower Late Angioplasty Complications (CADILLAC) trial. *Circulation* 2005; 111: 1611-1618.
12. Oe K, Shimizu M, Ino H, Yamaguchi M, Terai H, Hayashi K, et al. Effects of gender on the number of diseased vessels and clinical outcome in Japanese patients with acute coronary syndrome. *Circ J* 2002; 66: 435-440.
13. Yarzelski J, Col N, Pagley P, Savageau J, Gore J, Goldberg R. Gender differences and factors associated with the receipt of thrombolytic therapy in patients with acute myocardial infarction: A community-wide perspective. *Am Heart J* 1996; 131: 43-50.
14. Kostis JB, Wilson AC, O'Dowd K, Gregory P, Chelton S, Cosgrove NM, et al. Sex differences in the management and long-term outcome of acute myocardial infarction: A statewide study. MIDAS Study Group: Myocardial Infarction Data Acquisition System. *Circulation* 1994; 90: 1715-1730.
15. Kosuge M, Kimura K, Kojima S, Sakamoto T, Ishihara M, Asada Y, et al. Beneficial effect of preinfarction angina on in-hospital outcome is preserved in elderly patients undergoing coronary intervention for anterior acute myocardial infarction. *Circ J* 2005; 69: 630-635.
16. The TIMI study group. The Thrombolysis in Myocardial Infarction (TIMI) trial. *N Engl J Med* 1985; 312: 932-936.
17. Mehilli J, Ndrepepa G, Kastrati A, Nekolla SG, Markwardt C, Bollwein H, et al. Gender and myocardial salvage after reperfusion treatment in acute myocardial infarction. *J Am Coll Cardiol* 2005; 45: 828-831.
18. Weaver WD, Simes RJ, Betriu A, Grines CL, Zijlstra F, Garcia E, et al. Comparison of primary coronary angioplasty and intravenous thrombolytic therapy for acute myocardial infarction: A quantitative review. *JAMA* 1997; 278: 2093-2098.
19. Capes SE, Hunt D, Malmberg K, Gerstein HC. Stress hyperglycemia and increased risk after myocardial infarction in patients without diabetes: A systematic overview. *Lancet* 2000; 355: 773-778.
20. Wahab NN, Cowden EA, Pearce NJ, Gardner MJ, Merry H, Cox JL; ICONS Investigators. Is blood glucose an independent predictor of mortality in acute myocardial infarction in the thrombolytic era? *J Am Coll Cardiol* 2002; 40: 1748-1754.
21. Sadeghi HM, Stone GW, Grines CL, Mehran R, Dixon SR, Lansky AJ, et al. Impact of renal insufficiency in patients undergoing primary angioplasty for acute myocardial infarction. *Circulation* 2003; 108: 2769-2775.
22. Kosuge M, Kimura K, Kojima S, Sakamoto T, Matsui K, Ishihara M, et al. Effects of glucose abnormalities on in-hospital outcome after coronary intervention for acute myocardial infarction. *Circ J* 2005; 69: 375-379.

Determinants of rapid progression of aortic root dilatation and complications in Marfan syndrome^B

Aleksandar M. Lazarevic^a, Satoshi Nakatani^{a,T}, Yutaka Okita^b, Jelena Marinkovic^c,
Yutaka Takeda^a, Keiji Hirooka^a, Hiroshi Matsuo^a, Soichiro Kitamura^b,
Masakazu Yamagishi^a, Kunio Miyatake^a

^a*Department of Cardiology, National Cardiovascular Center, Suita, Osaka, Japan*

^b*Department of Cardiothoracic Surgery, National Cardiovascular Center, Suita, Osaka, Japan*

^c*Cardiovascular Research Center, Dedinje Cardiovascular Institute, Belgrade University Medical School, Belgrade, Serbia and Montenegro*

Received 3 August 2004; received in revised form 20 January 2005; accepted 28 January 2005

Available online 30 March 2005

Determinants of rapid progression of aortic root dilatation and complications in Marfan syndrome[☆]

Aleksandar M. Lazarevic^a, Satoshi Nakatani^{a,*}, Yutaka Okita^b, Jelena Marinkovic^c,
Yutaka Takeda^a, Keiji Hirooka^a, Hiroshi Matsuo^a, Soichiro Kitamura^b,
Masakazu Yamagishi^a, Kunio Miyatake^a

^aDepartment of Cardiology, National Cardiovascular Center, Suita, Osaka, Japan

^bDepartment of Cardiothoracic Surgery, National Cardiovascular Center, Suita, Osaka, Japan

^cCardiovascular Research Center, Dedinje Cardiovascular Institute, Belgrade University Medical School, Belgrade, Serbia and Montenegro

Received 3 August 2004; received in revised form 20 January 2005; accepted 28 January 2005

Available online 30 March 2005

Abstract

Background: Progressive aortic dilatation has prognostic significance in the Marfan syndrome.

Methods: To identify which patients were at high risk of rapid progression, we echocardiographically studied 43 patients (age 22 ± 14 years) with the mean follow-up period of 5.2 ± 3.2 years. Aortic diameters, left ventricular (LV) size, fractional shortening, and the severity of aortic and mitral regurgitation were assessed. Transmitral peak early and atrial flow velocities, their ratio and the deceleration time of peak early velocity were also obtained.

Results: Mean annual increases of aortic diameters were 0.4 ± 0.3 mm at the annulus, 1.5 ± 1.3 mm at the sinuses of Valsalva, 0.7 ± 0.6 mm at the supraaortic ridge and 0.4 ± 0.4 mm at the proximal ascending aorta. Patients were divided into 2 groups according to the aortic growth rate at the sinuses of Valsalva level: rapid (R, $>3\%$ per year, 15 patients) or slow (S, $\leq 3\%$ per year, 28 patients) progression groups. Measured variables did not show significant differences between the 2 groups except older age, higher blood pressure and more severe aortic regurgitation in group R. Multiple regression analysis identified prolonged deceleration time as the most important variable predicting aortic complications. Aortic dissection occurred more frequently in group R (7 patients, 47%) than in group S (0%, $P < 0.001$).

Conclusions: Marfan patients at older age, with higher blood pressure, and with significant aortic regurgitation were at high risk of progression of aortic dilatation, with the most remarkable increase at the sinuses of Valsalva. Prolonged deceleration time may relate to an increased risk for aortic complications.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Marfan syndrome; Echocardiography; Aortic dissection; LV function

1. Introduction

One of the most serious clinical manifestations of the Marfan syndrome is progressive aortic root dilatation which may lead to aortic dissection, rupture or regurgitation that are responsible for decreased life expectancy in these patients [1–4]. Although aortic root dilatation is observed in 60–80% of Marfan patients [5–7], aortic growth rate among individuals may vary [8] and factors predicting the rate of change of the aortic diameter are still not completely

[☆] Dr. Lazarevic was supported by a scholarship from the Ministry of Education, Science and Culture of Japan, Tokyo, Japan. This study was partly supported by a Research Grant from the Ministry of Health, Labor and Welfare, Japan.

* Corresponding author. Department of Cardiology National Cardiovascular Center 5-7-1 Fujishiro-dai, Suita Osaka 565-8565, Japan. Tel.: +81 6 6833 5012; fax: +81 6 6872 7486.

E-mail address: nakatas@hsp.ncvc.go.jp (S. Nakatani).

clarified. It would be clinically important to define which patients are at high risk of rapid progression.

Several factors have been reported to relate to aortic complications in the Marfan syndrome including aortic size, systolic blood pressure and aortic growth rate [8]. However, determinants of aortic growth rate remains elusive. Because the abnormality in extracellular connective tissue matrix in the Marfan syndrome [9] involves not only aorta but also heart muscle [10], left ventricular (LV) systolic and diastolic function may have some influence on aortic growth rate and complications. In the present study, we investigated determinants of rapid progression of aortic root dilatation and aortic complications with a special interest of the effect of LV function on them. Further, we clarified characteristics of patients who might have poor prognosis.

2. Materials and methods

2.1. Study population

We studied retrospectively 114 consecutive patients with the Marfan syndrome or relatives of patients, who underwent clinical and echocardiographic examinations. Inclusion criteria were: (1) positive strict, internationally established diagnostic criteria [11,12] for the Marfan syndrome, (2) echocardiographic follow-up ≥ 1 year (2-dimensional echocardiography and Doppler color flow mapping), and (3) no proximal aortic surgery before the initial examination. The final study group included 43 patients (age 22 ± 14 years [range 1–59 years], 23 males), with the mean follow-up period of 5.2 ± 3.2 years (range 1–10 years).

2.2. Echocardiography

Two-dimensional, M-mode and Doppler echocardiograms were obtained with a commercially available cardiac ultrasound system, using a 2.5-MHz transducer and stored on videotape and/or strip chart for later analysis. Aortic root measurements were made in 2-dimensional parasternal long-axis view at end-diastole at 4 levels: aortic annulus, sinuses of Valsalva, supraaortic ridge and proximal ascending aorta 1–2 cm above the supraaortic ridge according to the method of Roman et al. (Fig. 1) [13]. All measurements were made using the leading edge technique on up to 5 cycles and averaged. The echocardiographic evidence of aortic root dilatation was determined using Roman's nomograms based on age and body surface area [13]. Dimensions of the sinuses of Valsalva and supraaortic ridge were used to determine an aortic ratio as follows:

$$\text{Aortic ratio} = \frac{\text{measured diameter}}{\text{expected diameter}}$$




Fig. 1. Schematic illustration of the long-axis two-dimensional echocardiogram. Aortic root measurements were obtained at 4 levels, including; 1) the annulus, 2) sinuses of Valsalva, 3) supraaortic ridge and 4) proximal ascending aorta. AO=aorta; LA=left atrium; LV=left ventricle.

where the expected diameter was obtained using the regression equation derived from Roman's nomograms [13]. The mean annual rate of aortic root dilatation was determined by dividing the absolute change in aortic diameter between the final and initial echocardiograms by the duration of follow-up in years. We also assessed the annual change in the aortic ratio by dividing the change in aortic ratio between the final and initial echocardiograms by the time interval between them. The severity of aortic regurgitation and mitral regurgitation was graded semi-quantitatively (1–4) using color Doppler jet area criteria [14–16]. Echocardiographic evidence of mitral valve prolapse was evaluated using established echocardiographic criteria [17]. LV end-diastolic and end-systolic diameters, fractional shortening, and diastolic interventricular septum and LV posterior wall thicknesses were determined according to the recommendations of the American Society of Echocardiography [18]. Standard pulsed wave Doppler flow velocities measurements were obtained from the apical long-axis view, with the sample volume placed at the level of the mitral valve leaflet tips. The following Doppler indexes were measured: peak early (E) and atrial (A) transmitral filling velocities, E deceleration time and E/A ratio. A single investigator (A.M.L.) blinded to the clinical data performed the off-line analysis.

2.3. Statistical analysis

Results are reported as mean value \pm SD. Comparison of data between patients groups was performed by use of a two-tailed, unpaired Student's t test or Mann-Whitney U test. Mean annual increases at the aortic root levels were compared using analysis of variance. The χ^2 test (with continuity correction when applicable) was used to compare frequencies among the patients groups. Multivariate and univariate linear regression analysis was used to evaluate which variables such as blood pressure, heart rate, LV diameter indexes, fractional shortening, aortic regurgitation and LV diastolic filling indexes correlated best with the progression of aortic root dilatation. Independence of relation was tested using logistic regression. Significance was established at $P < 0.05$.

3. Results

3.1. Changes in aortic diameters and grouping of patients

Because of suboptimal images, measurements at the supraaortic ridge level and those at the ascending aorta level were not available in 5 and 11 patients, respectively. Aortic diameters increased with time (Fig. 2). Mean annual increases were 0.4 ± 0.3 mm at the level of annulus, 1.5 ± 1.3 mm at the level of the sinuses of Valsalva, 0.7 ± 0.6 mm at the supraaortic ridge and 0.4 ± 0.4 mm at the proximal ascending aorta. The most rapid change was seen at the sinuses of Valsalva diameter ($F=17$, $P<0.001$ vs. 3 other aortic diameters). The change in aortic ratio was also the highest at the level of the sinuses of Valsalva ($3 \pm 4\%$ per year). Because the annual changes in aortic root diameters were most prominent at the aortic sinus level, patients were divided into 2 groups according to the mean annual change in aortic ratio at the sinuses of Valsalva: rapid (R, $>3\%$ per year, 15 patients) or slow (S, $\leq 3\%$ per year, 28 patients) progression groups. During the follow-up period, 7 patients in group R had aortic dissection requiring aortic surgery and 2 patients died. One patient died 2 months after aortic root and valve replacement and another patient died 3 years after Bentall's operation from colon cancer. Aortic dissection [2 proximal (involving the ascending aorta) and 5 distal (all dissections that originate after the left subclavian artery)] occurred more frequently in group R (7 patients, 47%) than in group S (0%, $\chi^2=12.37$, $P<0.001$). Only 1 patient from group S underwent prophylactic aortic root and valve replacement due to ascending aortic aneurysm and severe aortic regurgitation.

3.2. Characteristics of the patients

Demographic and clinical data for all 43 patients at the time of the initial examination are described in Table 1. Patients in group R were older than those in group S ($P<0.01$). There were no significant differences in sex, heart rate and body surface area, but systolic ($P<0.02$) and diastolic ($P<0.05$) blood pressures were higher in group R.




Fig. 2. Mean annual increases of aortic root diameter at 4 levels in all study patients.

Table 1

Demographic and clinical data in all study patients and in the slow (S) and rapid (R) progression group at entry

	Total (n=43)	Group S (n=28)	Group R (n=15)
Age (years)	22 ± 14	18 ± 12	30 ± 15*
Male sex	23 (53%)	14 (50%)	9 (60%)
Body surface area (m ²)	1.56 ± 0.37	1.51 ± 0.4	1.67 ± 0.2
Heart rate (beats/min)	73 ± 8	72 ± 7	75 ± 9
Blood pressure (mm Hg)			
Systolic	117 ± 15	112 ± 13	124 ± 16†
Diastolic	67 ± 9	65 ± 8	70 ± 9‡
Pulse pressure	50 ± 12	48 ± 11	54 ± 13
Mitral valve prolapse	24 (56%)	17 (61%)	7 (47%)
Aortic root dilatation	40 (93%)	26 (93%)	14 (93%)
Duration of follow-up (years)	5.2 ± 3.2	6.3 ± 3.1	3.3 ± 2.5§

* $P<0.01$; † $P<0.02$; ‡ $P<0.05$; § $P<0.005$ compared with group S. Data presented are mean value ± SD or number (%) of patients.

Pulse pressure was comparable between the 2 groups. Aortic root dilatation was similarly seen in group R and in group S. The presence of mitral valve prolapse was comparable in both groups. Duration of follow-up was shorter in group R ($P<0.005$). There were no differences in the incidence of patients using β -adrenergic blocking drugs between the 2 groups.

3.3. Echocardiographic data (Table 2)

At the initial examination mean aortic sinus diameter ($P=0.001$) and aortic ratio ($P=0.01$) were larger in group R. LV diameters and fractional shortening were comparable between the 2 groups, but LV wall thickness was larger in group R ($P=0.01$ for interventricular septum and $P=0.02$ for posterior wall). Aortic regurgitation was more common ($P=0.001$) and more severe in group R ($P=0.02$). The prevalence ($P=0.349$) and severity of mitral regurgitation ($P=0.13$) was comparable in both groups (Table 2).

Echocardiographic data at the most recent follow-up or before surgery showed no significant differences in LV diameters, fractional shortening and severity of mitral regurgitation between groups R and S. However, aortic sinus diameter ($P<0.001$) and aortic ratio ($P<0.001$) were larger and aortic regurgitation was more severe in group R ($P<0.001$).

Pulsed Doppler indexes were obtained in 27 out of 43 patients (12 patients in group R and in 15 patients in group S) (Table 3). At the time of the initial pulsed Doppler examination, there were no differences in age (30 ± 14 vs. 27 ± 12 years), blood pressures (125 ± 28 vs. 117 ± 11 mm Hg for systolic pressure and 69 ± 9 vs. 68 ± 8 mm Hg for diastolic pressure), LV dimensions (29 ± 4 vs. 30 ± 4 mm/m² for end-diastolic diameter index and 18 ± 4 vs. 19 ± 3 mm/m² for end-systolic diameter index), fractional shortening ($37 \pm 5\%$ vs. $36 \pm 3\%$), LV wall thickness (8.4 ± 1.4 vs. 9.2 ± 1.6 mm for interventricular septum and 8.1 ± 1.2 vs. 8.8 ± 1.5 for posterior wall), and the severity of aortic

Table 2
Echocardiographic variables

	Total (n=43)		Group S (n=28)		Group R (n=15)	
	Initial	Follow-up	Initial	Follow-up	Initial	Follow-up
Aortic sinus diameter (mm)	36.8±8.7	43.2±11.5	33.7±7.2	39±7	42.7±8.4*	51±14 [†]
Aortic sinus ratio	1.36±0.23	1.49±0.32	1.30±0.20	1.37±0.2	1.48±0.24 [‡]	1.72±0.4 [†]
LV EDD (mm)	48±7	52±8	47±8	51±8	50±5	54±9
LV EDDI (mm/m ²)	32±8	32±6	33±9	31±6	31±6	33±7
LV ESD (mm)	31±5	34±7	30±6	33±6	32±4	35±8
LV ESDI (mm/m ²)	21±5	21±5	21±6	20±4	20±5	22±6
FS (%)	36±4	34±5	36±4	34±5	36±5	34±5
IVS (mm)	8.1±1.8	8.6±1.6	7.6±1.6	8.3±1.5	9.0±1.7 [‡]	9.2±1.7
PW (mm)	7.7±1.7	8.3±1.5	7.3±1.5	7.9±1.4	8.5±1.8 [§]	8.9±1.6
Aortic regurgitation						
Prevalence	16 (37%)	22 (51%)	5 (18%)	8 (29%)	11 (73%)*	14 (93%) [†]
Severity	0.5±0.8	0.9±1.2	0.3±0.8	0.4±0.9	0.9±0.7 [§]	1.9±1.1 [†]
Mitral regurgitation						
Prevalence	12 (28%)	19 (44%)	6 (21%)	11 (39%)	6 (40%)	8 (53%)
Severity	0.4±0.7	0.8±1.1	0.2±0.5	0.6±1.0	0.7±1.0	1.0±1.2

* $P=0.001$; [†] $P<0.001$; [‡] $P=0.01$; [§] $P=0.02$; ^{||} $P=0.04$ compared with group S. Data presented are mean value ± SD or number (%) of patients. EDD=end-diastolic diameter; EDDI=end-diastolic diameter index, ESD=end-systolic diameter; FS=fractional shortening; IVS=diastolic interventricular septum thickness; PW=diastolic LV posterior wall thickness.

regurgitation (0.7 ± 0.5 vs. 0.5 ± 1.0) and mitral regurgitation (0.7 ± 1.0 vs. 0.2 ± 0.5) between the 2 groups. However, patients in group R showed longer deceleration time ($P=0.005$), lower E ($P=0.015$) and lower E/A ($P=0.002$) compared to those in group S. A wave velocity did not differ significantly between the 2 groups. Follow-up data were available in 17 patients. At the most recent follow-up, deceleration time was longer in group R ($P=0.03$). Other Doppler indexes were comparable between the 2 groups.

3.4. Possible predictors of rapid progression of aortic root dilatation

To identify possible predictors of rapid aortic root dilatation, we performed multiple regression analyses. Aortic root dilatation at the sinuses of Valsalva level and that at the supraaortic ridge level were determined by diastolic pressure (t value=2.091, $P=0.043$ for the sinus level, t value=2.348, $P=0.024$ for the supraaortic ridge level). Changes of aortic ratios at the sinus and supraaortic ridge levels were influenced by diastolic pressure and aortic regurgitation (t value=2.340, $P=0.024$ and t value=2.163, $P=0.037$ for the aortic sinus, and t value=3.374, $P=0.002$ and t value=3.438, $P=0.002$ for the supraaortic ridge).

Fig. 3 shows t values of each parameter to predict aortic growth rate with overall multiple correlation coefficient.

Multivariate logistic regression analysis was used to identify the best predictor of rapid aortic dilatation (group R patients). In this analysis, the parameters which differed significantly between groups R and S at initial evaluation were included. They were aortic sinus diameter, aortic sinus ratio, systolic and diastolic blood pressures, degree of aortic regurgitation, deceleration time, E and E/A . The only independent predictor of rapid dilatation was E/A ratio ($P=0.0223$). We also assessed the independent predictors of aortic complications using the logistic regression with the same parameters. The only independent predictor of aortic complications was deceleration time ($P=0.0295$).

4. Discussion

4.1. Features of the patients with rapid progression of the aortic root dilatation

The present study showed that the Marfan patients at older age, with higher blood pressure, and with significant aortic regurgitation were at high risk of rapid progression of

Table 3
Doppler left ventricular diastolic filling variables.

	Total		Group S		Group R	
	Initial (n=27)	Follow-up (n=17)	Initial (n=15)	Follow-up (n=8)	Initial (n=12)	Follow-up (n=9)
DT	184±26	196±40	172±15	175±22	199±30*	216±43 [†]
E	62±15	57±14	68±13	63±13	54±14 [‡]	51±14
A	44±13	44±9	40±14	42±10	48±12	46±8
E/A	1.55±0.58	1.35±0.45	1.84±0.57	1.55±0.36	1.19±0.35 [§]	1.18±0.46

* $P=0.005$; [†] $P=0.03$; [‡] $P=0.015$; [§] $P=0.002$ compared with group S. Data presented are mean value ± SD or number of patients. A =atrial diastolic filling velocity; DT=deceleration time of the early diastolic filling velocity; E =early diastolic filling velocity.




Fig. 3. Impact of patients' characteristics on annual change of aortic root diameter and on change of the aortic ratio; t values of each parameter are shown in bar graphs. Black bars indicate $P < 0.05$ and white bars indicate $P > 0.05$. R shows multiple correlation coefficient. AR=severity of aortic regurgitation; DP=diastolic blood pressure; EDDI=end-diastolic diameter index; FS=left ventricular fractional shortening; HR=heart rate; Δ ridge=change in supraaortic ridge diameter per year; ridge rate=change in supraaortic ridge ratio; Δ SofV=change in aortic sinuses of Valsalva diameter per year; SofV rate=change in aortic sinus ratio.

aortic root dilatation and subsequent aortic complications. Despite significant initial differences between groups in the prevalence and severity of aortic regurgitation, we found that LV size and systolic function were comparable between groups with rapid and slow progression. However, interestingly, abnormal LV diastolic filling pattern, indicating impaired LV relaxation, was more common in patients with rapid progression. Note that in this analysis, factors influencing Doppler filling parameters such as age, blood pressure, LV size, wall thickness, and aortic and mitral regurgitation grades, were all comparable between patients with slow and rapid progression. Although there was a relatively small number of patients who had pulsed Doppler recordings, the features associated with impaired diastolic LV relaxation could predict more rapid aortic root dilatation and development of complications in select Marfan patients. Further, our findings could have a logical explanation when considering the potential abnormalities of the collagen matrix that may involve the myocardium in the Marfan syndrome [10]. The abnormalities of the extracellular connective tissue matrix in patients with the Marfan syndrome could involve heart muscle as well as aorta and heart valves, leading to ventricular diastolic dysfunction [10]. It has been speculated that defective microfibrils and elastic fibers in the cardiac cytoskeleton weaken the elastic restoring forces in the Marfan syndrome, and therefore LV relaxation is impaired [10].

4.2. Determinants of rapid aortic root dilatation

Since the Marfan patients with aortic root dilatation may have poor prognosis, it would be clinically important to predict which patients would progressively develop aortic

root dilatation. Therefore, we performed multivariate analysis considering clinical and echocardiographic parameters as independent variables and changes in aortic root diameters as the dependent variables. We found that aortic regurgitation grade, diastolic blood pressure and LV size had significant influence on the rate of aortic root dilatation. This finding is in accord with the previously published study regarding the poorer prognosis (decreased long-term survival) in patients with diastolic murmur or cardiomegaly on initial physical examination [4]. The presence of aortic regurgitation or dilated LV, or both should be a sign to proceed in an aggressive manner to further treatment.

4.3. Clinical implications

Our study confirms the importance of age, high blood pressure, larger initial aortic ratio, and presence and severity of aortic regurgitation, abnormal Doppler diastolic filling pattern at baseline as markers of high risk of rapid progression of aortic dilatation and subsequent aortic complications. This underscores the need for more frequent control and aggressive treatment in the Marfan patients with these features.

Diastolic dysfunction has been known to precede systolic dysfunction in some forms of heart failure. We found that even in patients with rapid progression of aortic dilatation (which could represent more severe disease expression) systolic function was preserved. It appears that in the Marfan syndrome, diastolic dysfunction precedes systolic dysfunction. It may be possible that our findings, indicating impaired relaxation, may show an overture to rapid progression of aortic dilatation and the risk for aortic complications in select Marfan patients.

4.4. Study limitations

In this retrospective study, some patients had suboptimal echocardiographic recordings for measurements of aortic diameters (at supraaortic ridge and ascending aorta level) and these measurements were not included in the analysis. Furthermore, although all study patients had color Doppler recordings, only 63% of all patients had pulsed Doppler recordings. Despite of the small number of patients, the difference in LV filling pattern between the 2 groups was very prominent and consistent.

Although transmitral flow velocity pattern provides a measure of diastolic filling, almost all of the indexes derived from the pattern are load dependent [19]. Again, note that in this study LV size, wall thickness, blood pressure, and the severity of aortic regurgitation and mitral regurgitation were comparable between the 2 patient groups in which pulsed Doppler examination was performed. Therefore, effects of loading conditions would be minor.

Since only 2 patients, both from the rapid progression group, received β -blocker therapy, effect of therapy could not be addressed in this study. To clarify the relationship between LV performance and progression of aortic root dilatation, a prospective study should be conducted with a larger number of patients.

4.5. Conclusions

Marfan patients at older age, with higher blood pressure, and with significant aortic regurgitation were at high risk of progression of aortic dilatation, with the most remarkable increase at the sinuses of Valsalva. LV systolic function appeared not to relate to the progression of aortic root dilatation. Prolonged deceleration time may relate to an increased risk for aortic complications.

References

- [1] Pyeritz RE. The Marfan syndrome. In: Royce PM, Steinmann B, editors. *Connective tissue and its heritable disorders: molecular, genetic, and medical aspects*. New York: Wiley-Liss; 1993. p. 437–68.
- [2] McKusick VA. The cardiovascular aspects of Marfan's syndrome: a heritable disorder of connective tissue. *Circulation* 1955;11:321–42.
- [3] Murdoch JL, Walker BA, Halpern BL, Kuzma JW, McKusick VA. Life expectancy and causes of death in the Marfan syndrome. *N Engl J Med* 1972;286:804–8.
- [4] Marsalese DL, Moodie DS, Vacante M, et al. Marfan's syndrome: natural history and long-term follow-up of cardiovascular involvement. *J Am Coll Cardiol* 1989;14:422–8.
- [5] Brown OR, DeMots H, Kloster FE, Roberts A, Menashe VD, Beals RK. Aortic root dilatation and mitral valve prolapse in Marfan's syndrome: an echocardiographic study. *Circulation* 1975;52:651–7.
- [6] Come PC, Fortuin NJ, White RI, McKusick VA. Echocardiographic assessment of cardiovascular abnormalities in the Marfan syndrome: comparison with clinical findings and with roentgenographic estimation of aortic root size. *Am J Med* 1983;74:465–74.
- [7] Bruno L, Tredici S, Mangiavacchi M, Colombo V, Mazzotta GF, Sirtori CR. Cardiac, skeletal, and ocular abnormalities in patients with the Marfan's syndrome and their relatives: comparison with the cardiac abnormalities in patients with kyphoscoliosis. *Br Heart J* 1984; 51:220–30.
- [8] Roman MJ, Rosen SE, Kramer-Fox R, Devereux RB. Prognostic significance of the pattern of aortic root dilatation in the Marfan syndrome. *J Am Coll Cardiol* 1993;22:1470–6.
- [9] Hollister DW, Goodfrey M, Sakai LY, Pyeritz RE. Immuno-histologic abnormalities of the microfibrillar-fiber system in the Marfan syndrome. *N Engl J Med* 1990;323:152–9.
- [10] Savolainen A, Nisula L, Keto P, et al. Left ventricular function in children with the Marfan syndrome. *Eur Heart J* 1994;15:625–30.
- [11] Pyeritz RE, McKusick VA. The Marfan syndrome: diagnosis and management. *N Engl J Med* 1979;300:772–7.
- [12] De Paepe A, Devereux RB, Dietz HC, Hennekam RCM, Pyeritz RE. Revised diagnostic criteria for the Marfan syndrome. *Am J Med Genet* 1996;62:417–26.
- [13] Roman MJ, Devereux RB, Kramer-Fox R, O'Loughlin J. Two-dimensional echocardiographic aortic root dimensions in normal children and adults. *Am J Cardiol* 1989;64:507–12.
- [14] Perry GJ, Helmcke F, Nanda NC, Byard C, Soto B. Evaluation of aortic insufficiency by Doppler color flow mapping. *J Am Coll Cardiol* 1987;9:952–9.
- [15] Miyatake K, Izumi S, Okamoto M, et al. Semiquantitative grading of severity of mitral regurgitation by real-time two-dimensional Doppler flow imaging technique. *J Am Coll Cardiol* 1986;7:82–8.
- [16] Helmcke F, Nanda NC, Hsiung MC, et al. Color Doppler assessment of mitral regurgitation with orthogonal planes. *Circulation* 1987;75: 175–83.
- [17] Levine RA, Stathogiannis E, Newell JB, Harrigan P, Weyman AG. Reconsideration of echocardiographic standards for mitral valve prolapse: lack of association between leaflet displacement isolated to the apical 4-chamber view and independent echocardiographic evidence of abnormality. *J Am Coll Cardiol* 1988;11:1010–9.
- [18] Sahn DJ, DeMaria AN, Kissio J, Weyman A. Recommendations regarding quantitation in M-mode echocardiography: results of a survey of echocardiographic measurements. *Circulation* 1978;58: 1072–83.
- [19] Thomas JD, Weyman AE. Echo-Doppler evaluation of left ventricular diastolic function: physics and physiology. *Circulation* 1991; 84:977–90.

Altered expression balance of matrix metalloproteinases and their inhibitors in human carotid plaque disruption: Results of quantitative tissue analysis using real-time RT-PCR method

Takeo Higashikata^a, Masakazu Yamagishi^{a,*}, Toshio Higashi^b, Izumi Nagata^b, Koji Iihara^b,
Susumu Miyamoto^b, Hatsue Ishibashi-Ueda^c, Noritoshi Nagaya^d,
Takashi Iwase^d, Hitonobu Tomoike^a, Aiji Sakamoto^e

^a Division of Cardiovascular Medicine and Bioscience, National Cardiovascular Center and Research Institute,
5-7-1 Fujishiro-dai, Suita, Osaka 565-8565, Japan

^b Division of Neurosurgery, National Cardiovascular Center, Suita, Osaka, Japan

^c Division of Pathology, National Cardiovascular Center, Suita, Osaka, Japan

^d Division of Regenerative Medicine and Tissue Engineering, National Cardiovascular Center and Research Institute, Suita, Osaka, Japan

^e Division of Biotechnology and Bioscience, National Cardiovascular Center and Research Institute, Suita, Osaka, Japan

Received 28 October 2004; received in revised form 17 May 2005; accepted 27 May 2005

Available online 21 July 2005

Abstract

Background: The balance between degradation and synthesis of extracellular matrix determines its content in atherosclerotic tissue. To examine the role of expression balance of matrix metalloproteinases (MMPs) to their inhibitors, tissue inhibitors of metalloproteinases (TIMPs) and tissue factor pathway inhibitor-2 (TFPI-2) in the development and disruption of atherosclerotic plaque, these gene expressions in human carotid plaque were quantitatively determined by real-time reverse transcription (RT)-polymerase chain reaction (PCR) method.

Methods: Total RNA for cDNA synthesis was extracted from tissues in 24 patients with carotid endarterectomy. The amounts of cDNAs for MMP-1, -2, -3 and -9, TFPI-2 and TIMP-1, -2 and -3 were determined by real-time RT-PCR method, and normalized with glutaraldehyde 3-dehydrogenase.

Results: In plaques, the expression MMP-1 (1.53 ± 0.25 , mean \pm S.E.M.), MMP-3 (1.99 ± 0.59) and MMP-9 (2.00 ± 0.51) was augmented compared to those in the adjacent control regions (0.60 ± 0.16 , 0.46 ± 0.18 and 0.58 ± 0.21 , respectively, $p < 0.05$). The expression of TFPI-2 was lower in plaques (0.32 ± 0.08) than in controls (0.94 ± 0.23 , $p < 0.01$). Although the expression of TIMP-1 was higher in plaques (1.28 ± 0.23) than in controls (0.81 ± 0.10 , $p < 0.05$), the indices of MMP-1/TIMP-1, MMP-3/TIMP-3 and MMP-9/TIMP-1 were still significantly higher in plaques. Interestingly, MMP-9 and the resulting MMP-9/TIMP-1 balance in plaques with disruption were significantly higher (3.36 ± 1.52 and 1.66 ± 0.12 , $n = 11$) than those in non-disrupted plaques (1.11 ± 0.52 and 0.76 ± 0.12 , $n = 13$, $p < 0.05$).

Conclusion: With the decreased expression of TFPI-2, upregulation of MMPs in atherosclerotic plaque was disproportional to that of TIMPs, suggesting that imbalanced degradation and synthesis of extracellular matrix persists in advanced lesions, particularly in plaques with disruption.

© 2005 Elsevier Ireland Ltd. All rights reserved.

Keywords: Atherosclerosis; Extracellular matrix; Matrix metalloproteinases; Tissue inhibitor of metalloproteinases; Tissue factor pathway inhibitor-2; Plaque disruption

1. Introduction

Disruption of atherosclerotic plaque during its development can expose the thrombogenic core to luminal blood, frequently resulting in ischemic cardiac events and stroke

* Corresponding author. Tel.: +81 6 6833 5012; fax: +81 6 6833 9865.
E-mail address: myamagi@hsp.ncvc.go.jp (M. Yamagishi).

[1,2]. During this process, structural changes in extracellular matrix (ECM) were shown to play a crucial role in plaque development and disruption [3]. The structural integrity of plaques seems to depend on a balance between synthesis and degradation of the ECM which is mainly regulated by proteinases such as matrix metalloproteinases (MMPs) including interstitial collagenase or MMP-1, gelatinase A or MMP-2, stromelysin 1 or MMP-3, gelatinase B or MMP-9 [4,5].

The activities of MMPs are controlled on multiple levels; transcription and translation of their inactive precursors (zymogens), post-translational activation of zymogens by proteolysis and interactions with tissue inhibitors of metalloproteinases (TIMPs) [6] and/or tissue factor pathway inhibitor-2 (TFPI-2) [7]. Indeed, TIMPs-1, -2, -3 and TFPI-2 are expressed in atherosclerotic lesions [7–9], and these inhibitors bind to and inactivate most of the MMPs [7,10]. Thus, the expression balance of MMP to TIMP and TFPI-2 is considered to regulate the net degeneration of ECM, thus contributing to maintaining plaque stability [7,11,12]. However, few systematic data exist regarding quantitative evaluation of the expression of MMPs and their inhibitors in human atherosclerotic plaques, probably because of technical difficulties in simultaneous determination of multiple gene expression in small tissue samples obtained in clinical settings. In the present study, we used real-time reverse transcription (RT)-polymerase chain reaction (PCR) and analyzed gene expression levels of MMPs, TIMPs and TFPI-2 in human carotid plaque and an adjacent control region. We also compared expression and function of MMPs between histologically disrupted and non-disrupted plaques.

2. Subjects and methods

2.1. Subjects

The protocol of this study was approved by the institutional committee for ethical review. Written informed consent was obtained from all 24 patients who underwent carotid endarterectomy for severe stenosis of the extracranial carotid artery (all male with mean age of 68 ± 2

years). All patients presented clinical symptoms of cerebral ischemic attack related to carotid stenosis. Seven patients had a history of recent ischemic attack within 1 month prior to endarterectomy. The prevalence of risk factors for atherosclerosis was as follows: hypertension (systolic pressure >160 mmHg) in 20, hyperlipidemia (total cholesterol >220 mg/dl) in 22, smoking in 15 and diabetes mellitus (fasting blood glucose >110 mg/dl) in 10 patients. High sensitive (hs) CRP level (normal range <3 mg/l) just before surgery was 2.45 ± 0.43 mg/l (Table 1).

2.2. Tissue sampling

Samples of the plaque region were obtained immediately after endarterectomy. Endarterectomy was extended in a caudal direction to include a sample of minimally affected common carotid artery proximal to the plaque but in continuity with the plaque to act as a paired control. Under these conditions, the stenotic segment and adjacent areas were dissected undisturbedly as a single specimen, preserving circumferential integrity as much as possible. Also special care was taken not to damage luminal surface and plaque interior. After removing a part of the tissue for histological examination, all samples were immediately frozen in liquid nitrogen and stored at -80°C until extraction of mRNA.

Procedures for RNA preparation and cDNA synthesis were already described elsewhere in detail [13]. Briefly, the samples were homogenized in 1.0 ml ISOGENTM reagent (Nippon Gene, Tokyo, Japan), thoroughly mixed with 0.2 ml chloroform and centrifuged at $15,000 \times g$ for 15 min at 4°C . The aqueous supernatant was transferred into a micro test tube, mixed with 0.6 ml isopropanol and centrifuged at $15,000 \times g$ for 15 min at 4°C . The precipitated total RNA was rinsed with 70% ethanol, air-dried and then resuspended in RNase-free water. Then, all the total RNA was treated with DNase FreeTM reagent (Ambion, Austin, TX) for 60 min, and then reverse-transcribed with Superscript IITM (Invitrogen, Carlsbad, CA) at 37°C for 60 min using random primers (TaKaRa, Tokyo, Japan). The integrity of each cDNA mixture was checked by amplification of glutaraldehyde 3-phosphate dehydrogenase (GAPDH) with *ExTaq* (TaKaRa), using the primer set 5'-ACCACAGTCCATGCCATCAC-3'/5'-TCCACCACCCTGTTGCTGTA-3'.

Table 1
Patient Characteristics

	All patients (n=24)	With disruption (n=11)	Without disruption (n=13)	p-Value
Age	68 ± 2	66 ± 3	69 ± 2	NS
Male sex	24	11	13	NS
Hypertension	20	8	12	NS
Diabetes	10	5	5	NS
HbA1c (%)	6.5 ± 0.4	7.0 ± 0.8	6.2 ± 0.4	NS
Hyperlipidemia	22	9	13	NS
LDL (mg/dl)	132 ± 6	140 ± 10	128 ± 7	NS
Smoking	15	5	10	NS
hs-CRP (mg/l)	2.45 ± 0.43	2.68 ± 0.49	2.12 ± 0.81	NS