

ulated with PBS, intracerebrally and intramuscularly. The inoculated mice were observed daily for neurological conditions and were killed at 2, 3, 4, 5, 7 and 9 days postinoculation (PI)—three to 6 mice per day. All experiments were performed in level-2 biosafety laboratories according to Committees on Biosafety and Animal Handling and Ethical Regulation of the National Institute of Infectious Diseases, Japan. Animal care, breeding, virus inoculation and observation were performed in accordance with the guidelines of Committees.

Necropsy and preparation of tissue sections: Each mouse was perfused transcardially with 10–15 ml of PBS followed by freshly prepared 4% paraformaldehyde in 0.1 M PBS, pH 7.4. Brains and muscles were removed and postfixed in 4% paraformaldehyde at room temperature for less than 24 hr. The spinal samples were decalcified in K-CX (Fujisawa, Japan) solution after fixation. Coronal sections of the brains at the positions of the olfactory bulb, forebrain, mamillary body and pons and transverse sections of the spinal cords at C2 to C4, T3 to T5, L2 to L4 and S1 to S3 were prepared. Complete series of paraffin sections about 3 μ m thick were cut and mounted on glass slides. Serial sections were used for hematoxylin and eosin (HE) staining, immunohistochemistry and TUNEL assays.

Immunohistochemistry: Paraffin sections were mounted on silane-coated slides, deparaffinized in xylene and rehydrated in alcohol, then treated with 0.025% trypsin at 37°C for 30 min. After blocking of endogenous peroxidase activity with 0.3% H₂O₂ in methanol for 30 min, sections were treated with 5% normal goat serum to block nonspecific reactions. For detection of rabies virus antigens in the tissues, all sections were stained using the streptavidin-biotin-peroxidase complex (LSAB) method using rabbit anti-CVS-11 strain nucleoprotein at a 1:2,000 dilution as previously described [13]. Tissues from uninfected mice and substitution of normal rabbit serum for the primary antibody were used as controls.

TUNEL assay: In order to compare the DNA strand breaks in tissue sections from infected mice, the terminal deoxynucleotidyl transferase mediated dUTP-biotin nick end labeling (TUNEL) method (R & D Systems, Minneapolis, MN) was used. Paraffin sections from infected mice sacrificed at 3, 5, 7 and 9 days PI were prepared according to the manufacturer's instructions. Selected slides were double labeled with the TUNEL method followed by immunostaining for rabies virus antigen without counterstaining.

RESULTS

Clinical signs: The mice became hunched and hypokinetic 4 to 5 days after intracerebral inoculation and subsequently became moribund at 8 days PI. No mice showed paralysis before their death at 10 to 11 days PI. In intramuscularly inoculated mice, paralysis was found at 4 days PI. Paralysis was initially mild with decreased hind limb movement, but as the disease worsened, the mice became severely paralyzed at 7 days PI and progressed to quadriparalysis at 9

days PI. All infected mice died between 11 to 13 days PI.

Intracerebral inoculation

Histology: No histological changes were observed in any area, including the hippocampus, up to 3 days PI. At 4 days PI, the nuclear pyknosis and fragmentation, cytoplasmic shrinkage were first seen in the pyramidal neurons (CA3, Fig. 1a) of the hippocampus. These findings became severe and most of the pyramidal neurons were destroyed at 7 days PI (Fig. 1b). Lesser changes were observed in multiple other areas, including the thalamus, brain stem and cerebellum at 5 days PI. Throughout the experiments, no histological changes were observed in the neurons of the dentate gyrus of the hippocampus, spinal neurons, spinal ganglia neurons and muscles.

Immunohistochemistry: The localization of viral antigens in the cerebral cortex, hippocampus, thalamus, brain stem, cerebellum, spinal cord and muscles is summarized in Table 1. Immunohistochemical examination revealed viral antigens in the right cerebral cortex and hippocampus (CA3) at 2 days PI. In the thalamus, brain stem, spinal cord (cervical, thoracic, lumbar, sacral), spinal dorsal ganglia, small numbers of virus positive neurons were first detected at 4 days PI. Viral antigens were not found in the glial cells in the spinal cord and brain. All of the neurons in the pyramidal layer of hippocampus were positive at 4 days PI (Fig. 1c) and more significant at 7 days PI (Fig. 1d), but neurons in the dentate gyrus was negative. In the cerebellum, antigen positive reactions were found in the Purkinje cells and neurons in the cerebellar nuclei at 5 days PI. At 5 days PI, a small number of positive cells were observed in the muscle spindles around the thoracic vertebrae.

TUNEL assays: TUNEL staining of the brain and spinal cord sections showed prominent staining in topographic regions with apoptotic morphologic changes and viral antigen positive cells. TUNEL staining of neurons in the pyramidal layer of the hippocampus was found from 4 days PI (Fig. 1e) and gradually increased as disease progressed. Staining was most prominent in the pyramidal neurons (CA3) at 7 days PI (Fig. 1f). Despite strong immunostaining for the viral antigen in cerebellar Purkinje cells, TUNEL staining was markedly less. No TUNEL staining was found in the dentate gyrus of the hippocampus, spinal cord, spinal ganglia, muscle spindle and muscle fibers.

Intramuscular inoculation

Histology: In the spinal cord and dorsal root ganglia, chromatolysis, Negri body-like inclusions and necrotic gangliocytes with proliferation of microglial cells, lymphocytes and macrophages were found from 4 days PI, and the numbers of necrotic neurons increased as the paralysis progressed at 7 days PI (Figs. 2a and 2b). In the cerebral cortex, thalamus, brain stem and cerebellum (Fig. 2c), similar histological changes were observed in infected neurons at the terminal stages. In the muscle fibers, at the site of inoculation, hyaline degeneration was found at 2 days PI and mild inflammatory cell infiltrates composed of neutrophils, lymphocytes and macrophages were observed around the small vessels. At 5 days PI, regeneration figures such as nuclear




Fig. 1. Hippocampus (CA3). Histology, immunohistochemistry and TUNEL assay of after intracerebral inoculation of CVS-11 at 4 days PI (Figs. 1a, c, e) and at 7 days PI (Figs. 1b, d, f). Eosinophilic cytoplasm and nuclear pyknosis (arrows, 1a), viral antigens (1c) and TUNEL positive cells (1e) was observed in pyramidal neurons. Nuclear fragmentation, pyknosis and vacuolation (arrows, 1b) and viral antigens (1d) and TUNEL positive cells (1f) are more significantly observed at 7 days. Bar=30 μ m.

Table 1. Distribution of CVS-11 virus antigens in mice infected intracerebrally

Days post-inoculation	2	3	4	5	7	9
Organ						
Brain						
Cerebral cortex	1/3*	3/3	3/3	3/3	3/3	3/3
Hippocampus	1/3	2/3	3/3	3/3	3/3	3/3
Thalamus	0/3	0/3	2/3	3/3	3/3	3/3
Brain stem	0/3	0/3	1/3	3/3	3/3	3/3
Purkinje cells	0/3	0/3	0/3	3/3	3/3	3/3
Cerebellar nuclei	0/3	0/3	0/3	3/3	3/3	3/3
Spinal cord [#]						
C	0/3	0/3	3/3	3/3	3/3	3/3
T	0/3	0/3	3/3	3/3	3/3	3/3
L	0/3	0/3	2/3	2/3	3/3	3/3
S	0/3	0/3	2/3	2/3	3/3	3/3
Muscle spindles [†]	0/3	0/3	0/3	1/3	2/3	3/3

*: Number of mice with positive staining for CVS-11 virus antigen/number of total mice used in experiment. [#]C: cervical, T: thoracic, L: lumbar, S: sacral cord. [†]: Muscle spindles around the infected cervical spinal cord.

chains and proliferating myoblasts were observed (Fig. 3a). No histological changes were found in any area of the hippocampus or muscle spindles throughout the experiments.

Immunohistochemistry: The localization of viral antigens in the cerebral cortex, hippocampus, thalamus, brain stem, cerebellum, spinal cord and muscles are summarized in Table 2. Antigen was initially observed in muscles at the site of inoculation in 2 of 3 infected mice at 2 days PI (Fig. 3b). Rarely, antigen positive reactions were also found in the peripheral nerves in the muscles. At 4 days PI, virus positive neurons were observed in the spinal dorsal root ganglia (Fig. 3c), spinal cord (Fig. 3d) and muscle spindles (Fig. 3e) around the sacral spina prior to the cerebrum, thalamus and cerebellum. At 5 days PI, virus positive neurons were detected in the cerebral cortex, thalamus, brain stem and cerebellum. The earliest hippocampal involvement was observed in the CA3 region at 7 days PI.

TUNEL assay: TUNEL staining was prominent in the external granular layer and scattered cells in the internal granular cell layer of the cerebellum. However, despite strong immunostaining for rabies virus antigen in cerebellar Purkinje cells, TUNEL staining was markedly less (Fig. 3f). TUNEL staining was not observed in the spinal motor and/or sensory neurons and spinal dorsal root ganglia gangliocytes. Some of the inflammatory cells and glial cells in the spinal cord and brain showed TUNEL staining.

DISCUSSION

In this study, clinicopathological appearances differed distinctly between intracerebrally and intramuscularly mice infected with the CVS-11 strain. After intracerebral inoculation, hippocampus and cerebral cortex neurons were the initial target, and apoptosis was induced before inflammatory cell infiltration. In addition, a good correlation between the distributions of apoptotic cells and viral antigen positive cells was observed. These findings are supported

by the data reported by Jackson *et al.* [16, 17] and Theerasurakarn and Ubol [29].

After hind limb inoculation, a more natural route of viral entry, viral antigen was first detected, and persisted until the terminal stages, in the muscle at the site of inoculation and was then detected in the spinal dorsal root ganglia, spinal cord and muscle spindles. These findings indicate that the virus reached the CNS in a retrograde direction through peripheral sensory nerves following replication in the muscles, and subsequently spread in an anterograde direction to distant muscle spindles. Limb paralysis appeared at the same time as necrotic changes and virus antigens were detected at 4 days PI. However, virus infected spinal motor and/or sensory neurons and spinal ganglion gangliocytes were TUNEL negative despite strong immunostaining. These results are partially in accordance with those of the previous report by Guignoniet *et al.* [10], in which rat embryonic spinal motor neurons and spinal cord of rat neonates were more resistant to cytolysis and apoptosis than hippocampus neurons after CVS strain infection.

In this study, infiltration of inflammatory cells into the muscles at the site of inoculation, the spinal dorsal root ganglia and the spinal cord were observed in early stage of infection. In addition, paralytic signs were distinct in intramuscularly infected mice in spite of less cerebral destruction. Therefore, it was suggested that activation of local immune responses in peripheral muscle and spinal cord, and the dysfunction of infected spinal neurons were precedence of paralytic manifestations. The pathogenesis of rabies includes encephalitis and paralysis [23]. However, little explanation has been provided that would account for these diverse manifestations of infection with a common agent [12]. Immunosuppressed and athymic mice develop encephalitic rabies. This encephalitis occurs concomitantly with increased virus replication and destruction of neurons in the absence of detectable immune mediators [14, 32]. Paralytic rabies is found in immunocompetent mice only.




Fig. 2. Intramuscularly infected C57BL/6J mouse. Spinal dorsal ganglia, spinal cord and cerebellum at 7 days PI. Degenerated gangliocytes (arrows, 2a), necrotic spinal neurons with microgliosis (arrowheads, 2b) and necrotic Purkinje cells (arrowheads, 2c) were observed. Bar=30 μ m (Figs. 2a and 2c), Bar=50 μ m (Fig. 2b).

Passive transfer of immune sera and activated lymphocytes to immunosuppressed mice results in paralysis and early death [14]. In contrast to mice, paralytic rabies in humans

Table 2. Distribution of CVS-11 virus antigens in mice infected intramuscularly

Days postinoculation	2	3	4	5	7	9
Organ						
Brain						
Cerebral cortex	0/3*	0/3	0/3	1/3	6/6	3/3
Hippocampus	0/3	0/3	0/3	0/3	3/6	2/3
Thalamus	0/3	0/3	0/3	1/3	6/6	3/3
Brain stem	0/3	0/3	0/3	2/3	6/6	3/3
Purkinje cells	0/3	0/3	0/3	1/3	6/6	3/3
Cerebellar nuclei	0/3	0/3	0/3	1/3	6/6	3/3
Spinal cord [#]						
C	0/3	0/3	2/3	3/3	6/6	3/3
T	0/3	0/3	2/3	3/3	6/6	3/3
L	0/3	0/3	2/3	3/3	6/6	3/3
S	0/3	0/3	2/3	3/3	6/6	3/3
Other						
Muscle spindles [†]	0/3	0/3	1/3	3/3	6/6	3/3
Muscle [‡]	2/3	3/3	3/3	3/3	6/6	3/3

*: Number of mice with positive staining for CVS-11 virus antigen/number of total mice used in experiment. [#]C: cervical, T: thoracic, L: lumbar, S: sacral cord. [†]: Muscle spindles around the infected sacral spinal cord. [‡]: Right triceps surae muscle.

has been found with defective immune reactions, while encephalitic rabies cases have an active immune response to rabies. This raises questions as to whether immune responses are significant in rabies pathogenesis and whether the pathogenesis process differs in mice and human.

Muscle plays an important role in the pathogenesis of rabies. However, it remains to be determined whether viral replication in muscle is essential for further virus spread. In several experiments, local replication in myocytes after inoculation into muscle seemed to precede the appearance of viral material in the nervous system [4, 5, 11, 24, 25]. In contrast, Coulon *et al.* [8] and Shankar *et al.* [26] did not find any evidence of replication in the muscle, suggesting that the virus could penetrate directly into the nerve endings. The present study supported the former case because viral antigen was initially detected in muscles. Baer *et al.* [2] and Charlton *et al.* [6] reported long incubation periods in muscle of rabid animals. In the present study, viral antigens were persistently detected in muscle at the inoculation site and did not spread directly from one muscle fiber to another.

In this study, viral antigens were detected in spindle muscles at an early stage of infection when infected mice showed paralysis. The virus that initially or subsequently replicated in the spinal cord or spinal ganglia had directly descended to the spindles via sensory or/and motor nerve fibers. Muscle spindles are skeletal muscle mechanoreceptors that mediate the stretch reflex and provide axial and limb position information to the CNS. Spindles consist of encapsulated muscle fibers (intrafusal fibers) that are innervated by specialized sensory and motor neurons. However, we do not know the direct effects of viral infection in spindles on cellular function or paralysis as no evidence that suggested morphological changes was observed until the terminal stage of infection. Therefore, further studies are




Fig. 3. Intramuscularly infected C57BL/6J mouse. At 5 days PI, hyaline degeneration, nuclear chain indicating muscular regeneration and perivascular infiltration (3a) are observed. Numerous viral antigen spots in the cytoplasm of muscle (arrows, 3b) and weak positive reaction (arrowhead, 3b) are seen in the peripheral nerve system. Viral antigens were observed in the cytoplasm of dorsal root ganglia gangliocytes (arrowhead, 3c). By double staining, viral positive neurons (arrowheads, 3d) and TUNEL positive glial cells (arrow, 3d) are seen. Viral antigens are also seen in the cytoplasm of muscle spindle (arrowhead, 3e) and peripheral nerve system (arrow, 3e). TUNEL staining was observed not only in the external granular cells (arrowheads, 3e) but also Purkinje cells (arrows, 3f). Bar=30 μ m (Figs. 3a, b, c and f), Bar=20 μ m (Figs. 3d and 3e).

required to address the effects of muscle spindle involvement in the clinical magnifications of rabies.

ACKNOWLEDGMENTS. This study was partly supported by a Research on Emerging and Reemerging Infectious Diseases grant from the Ministry of Health, Labor and Welfare of Japan, and a Grant for Scientific Research (Park CH: 2850) from the School of Veterinary Medicine and Animal Sciences of Kitasato University, Japan.

REFERENCES

1. Astic, L., Saucier, D., Coulon, P., Lafay, F. and Flamand, A. 1993. The CVS strain of rabies virus as transneuronal tracer in the olfactory system of mice. *Brain Res.* **619**: 146–156.
2. Baer, G. M. and Cleary, W. F. 1972. A model in mice for the pathogenesis and treatment of rabies. *J. Infect. Dis.* **125**: 520–527.
3. Burrage, T. G., Tignor, G. H. and Smith, A. L. 1985. Rabies virus binding at neuromuscular junctions. *Virus Res.* **2**: 273–289.
4. Charlton, K. M. and Casey, G. A. 1979. Experimental rabies in skunks: immunofluorescence light and electron microscopic studies. *Lab. Invest.* **41**: 36–44.
5. Charlton, K. M. and Casey, G. A. 1981. Experimental rabies in skunks: persistence of virus in denervated muscle at the inoculation site. *Can. J. Comp. Med.* **45**: 357–362.
6. Charlton, K. M., Nadin-Davis, S., Casey, G. A. and Wandeler, A. I. 1997. The long incubation period in rabies: delayed progression of infection in muscle at the site of exposure. *Acta Neuropathol.* **94**: 73–77.
7. Clem, R. J. and Miller, L. K. 1993. Apoptosis reduces both the *in vitro* replication and the *in vivo* infectivity of a baculovirus. *J. Virol.* **67**: 3730–3738.
8. Coulon, P., Derbin, C., Kucera, P., Lafay, F., Prehaud, C. and Flamand, A. 1989. Invasion of the peripheral nervous systems of adult mice by the CVS strain of rabies virus and its avirulent derivative AvO1. *J. Virol.* **63**: 3550–3554.
9. Gillet, J. P., Derer, P. and Tsiang, H. 1986. Axonal transport of rabies virus in the central nervous system of the rat. *J. Neuropathol. Exp. Neurol.* **45**: 619–634.
10. Guigoni, C. and Coulon, P. 2002. Rabies virus is not cytolytic for rat spinal motoneurons *in vitro*. *J. Neurovirol.* **8**: 306–317.
11. Harrison, A. K. and Murphy, F. A. 1978. Lyssavirus infection of muscle spindles and motor end plates in striated muscle of hamsters. *Arch. Virol.* **57**: 167–175.
12. Hemachudha, T., Wacharapluesadee, S., Lumlerdaecha, B., Orciari, L. A., Rupprecht, C. E., La-Ongpant, M., Juntrakul, S. and Denduangboripant, J. 2003. Sequence analysis of rabies virus in humans exhibiting encephalitic or paralytic rabies. *J. Infect. Dis.* **188**: 960–966.
13. Inoue, S., Sato, Y., Hasegawa, H., Noguchi, A., Yamada, A., Kurata, T. and Iwasaki, T. 2003. Cross-reactive antigenicity of nucleoproteins of lyssaviruses recognized by a monospecific antirabies virus nucleoprotein antiserum on paraffin sections of formalin-fixed tissues. *Pathol. Int.* **53**: 525–533.
14. Iwasaki, Y., Gerhard, W. and Clark, H. F. 1977. Role of host immune response in the development of either encephalitic or paralytic disease after experimental rabies infection in mice. *Infect. Immun.* **18**: 220–225.
15. Jackson, A. C. 1991. Biological basis of rabies virus neurovirulence in mice: comparative pathogenesis study using the immunoperoxidase technique. *J. Virol.* **65**: 537–540.
16. Jackson, A. C. and Park, H. 1998. Apoptotic cell death in experimental rabies in suckling mice. *Acta Neuropathol.* **95**: 159–164.
17. Jackson, A. C. and Rossiter, J. P. 1997. Apoptosis plays an important role in experimental rabies virus infection. *J. Virol.* **71**: 5603–5607.
18. Johnson, R. T. 1965. Experimental rabies. Studies of cellular vulnerability and pathogenesis using fluorescent antibody staining. *J. Neuropathol. Exp. Neurol.* **24**: 662–674.
19. Kucera, P., Dolivo, M., Coulon, P. and Flamand, A. 1985. Pathways of the early propagation of virulent and avirulent rabies strains from the eye to the brain. *J. Virol.* **55**: 158–162.
20. Lewis, P., Fu, Y. and Lentz, T. L. 2000. Rabies virus entry at the neuromuscular junction in nerve–muscle cocultures. *Muscle Nerve* **23**: 720–730.
21. Majno, G. and Joris, I. 1995. Apoptosis, oncosis, and necrosis. An overview of cell death. *Am. J. Pathol.* **146**: 3–15.
22. Morimoto, K., Hooper, D. C., Spitsin, S., Koprowski, H. and Dietzschold, B. 1999. Pathogenicity of different rabies virus variants inversely correlates with apoptosis and rabies virus glycoprotein expression in infected primary neuron cultures. *J. Virol.* **73**: 510–518.
23. Murphy, F. A. 1977. Rabies pathogenesis. *Arch. Virol.* **54**: 279–297.
24. Murphy, F. A. and Bauer, S. P. 1974. Early street rabies virus infection in striated muscle and later progression to the central nervous system. *Intervirology* **3**: 256–268.
25. Murphy, F. A., Bauer, S. P., Harrison, A. K. and Winn, W. C. 1973. Comparative pathogenesis of rabies and rabies-like viruses. Viral infection and transit from inoculation site to the central nervous system. *Lab. Invest.* **28**: 361–376.
26. Shankar, V., Dietzschold, B. and Koprowski, H. 1991. Direct entry of rabies virus into the central nervous system without prior local replication. *J. Virol.* **65**: 2736–2738.
27. Smith, J. S., Yager, P. A. and Baer, G. M. 1996. A rapid tissue culture test for determining rabies-neutralizing antibody. pp.371–373. *In: Laboratory Techniques in Rabies*, 4th ed. (Meslin, F. X., Kaplan, M. M. and Koprowski, H. eds.), Geneva: World Health Organization.
28. Tang, Y., Rampin, O., Giuliano, F. and Ugolini, G. 1999. Spinal and brain circuits to motoneurons of the bulbospongiosus muscle: retrograde transneuronal tracing with rabies virus. *J. Comp. Neurol.* **414**: 167–192.
29. Theerasurakarn, S. and Ubol, S. 1998. Apoptosis induction in brain during the fixed strain of rabies virus infection correlates with onset and severity of illness. *J. Neurovirol.* **4**: 407–414.
30. Ugolini, G. 1995. Specificity of rabies virus as a transneuronal tracer of motor networks: transfer from hypoglossal motoneurons to connected second-order and higher order central nervous system cell groups. *J. Comp. Neurol.* **356**: 457–480.
31. Watson, H. D., Tignor, G. H. and Smith, A. L. 1981. Entry of rabies virus into the peripheral nerves of mice. *J. Gen. Virol.* **56**: 372–382.
32. Weiland, F., Cox, J. H., Meyer, S., Dahme, E. and Reddehase, M. J. 1992. Rabies virus neuritic paralysis: immunopathogenesis of nonfatal paralytic rabies. *J. Virol.* **66**: 5096–5099.

New Criteria for Immunofluorescence Assay for Q Fever Diagnosis in Japan

A. Setiyono,^{1,3} M. Ogawa,^{1*} Y. Cai,¹ S. Shiga,¹ T. Kishimoto,¹ and I. Kurane²

Laboratory of Rickettsia and Chlamydia,¹ and Department of Virology I,² National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-Ku, Tokyo 162-8640, Japan, and Laboratory of Pathology, Department of Parasitology and Pathology, Faculty of Veterinary Medicine, Bogor Agricultural University, Jl. Agatis, Kampus IPB Darmaga Bogor 16680, Indonesia³

Received 26 October 2004/Returned for modification 24 January 2005/Accepted 9 August 2005

A study was made to evaluate the cutoff value of indirect immunofluorescent-antibody (IFA) test for Q fever diagnosis in Japan. We used 346 sera, including 16 from confirmed Q fever cases, 304 from Japanese pneumonia patients, and 26 from negative cases. Thirteen sera from the confirmed Q fever cases with an immunoglobulin M (IgM) titer of $\geq 1:128$ and/or IgG titer of $\geq 1:256$ by the IFA test were positive by both enzyme-linked immunosorbent assay (ELISA) and Western blotting assay (WBA), whereas 298 sera from pneumonia patients and 26 negative sera with an IgM titer of $\leq 1:16$ and an IgG titer of $\leq 1:32$ by the IFA test were negative by both ELISA and WBA. In the proposed “equivocal area,” with an IgM titer of $\geq 1:32$ and $\leq 1:64$ and/or an IgG titer of $\geq 1:64$ and $\leq 1:128$, we found 9 sera, 3 from confirmed Q fever cases and 6 from Japanese pneumonia patients, by the IFA test. Three sera from the confirmed Q fever cases and one of the sera from pneumonia patients were IgM and/or IgG positive by both ELISA and WBA. These results suggest that a single cutoff value for the IFA test may cause false-positive and false-negative results. In conclusion, this study showed that an “equivocal area” should be used for the IFA test rather than a single cutoff value and that sera in the equivocal area should be tested by additional serological assays for confirmation.

Q fever is a zoonosis caused by *Coxiella burnetii*, an obligate intracellular rickettsial organism. The clinical manifestations of Q fever are readily divided into acute and chronic forms. The most common clinical presentation of acute Q fever in human is an influenza-like illness, often accompanied by pneumonia. The chronic Q fever form, particularly endocarditis, may appear several years after the primary episode (1, 2, 6, 21). Because the clinical presentation of the infection is not specific, serological confirmation is required for the diagnosis of Q fever. In Japan, the currently used serological method is the indirect immunofluorescent-antibody (IFA) test, and more recently, some attempts have been made to use other methods, such as the enzyme-linked immunosorbent assay (ELISA).

The unique characteristic of *C. burnetii* is its antigenic phase variation (17). Virulent phase I can be isolated from natural infection of humans or from laboratory infections of animals. Phase II develops during serial passage in an immunologically incompetent host, such as cell cultures or fertilized eggs (1). Serologically, anti-phase I antibodies are normally found at high levels only during the chronic form of the disease, whereas specific anti-phase II antibodies predominate primarily in acute Q fever (14).

The IFA test has previously been used to detect immunoglobulin M (IgM) antibodies in the sera of Q fever patients within the first 2 weeks of illness (8, 12). The estimation of anti-*C. burnetii* IgM antibody using the IFA test in a single serum sample has been proven useful in confirming acute in-

fection in humans (8). The IFA method is more sensitive and specific than the complement fixation test; however, it is less sensitive than the ELISA (15). Therefore, the validation of the immunofluorescent threshold value for Q fever serology should be important for the establishment of the diagnosis.

Sodium dodecyl sulfate-polyacrylamide gel electrophoresis, followed by immunoblotting assay, has been used to identify the biologically important antigen in a complex mixture of proteins (9). The outer membrane-associated protein of *C. burnetii* is believed to be the antigenic target for the detection of antibody in clinical serum samples. This protein has been well characterized, with a molecular mass of approximately 27 kDa (18), and its usefulness as an immunodiagnostic reagent had also been evaluated (23).

In a preliminary study, we found some unsure results in the Q fever diagnosis by our IFA test and then compared them with ELISA results (data not published). Several samples reacted to phase II *C. burnetii* antigen in the IFA test but were negative by ELISA. On the other hand, a few of these sera were positive by ELISA but nonreactive in the IFA test. This phenomenon led us to reevaluate the criteria for the IFA test for the diagnosis of Q fever.

MATERIALS AND METHODS

Serum samples. A total of 346 human serum samples were included in this study. The specimens consisted of 16 serum samples from patients diagnosed with Q fever in past examinations by the IFA test, ELISA, and complement fixation test, which also served as a positive control, including three imported cases (13) and others kindly provided by Werner Slenczka (Institut für Virologie, Marburg, Germany), Barrie P. Marmion (Institute of Medical and Veterinary Science, Adelaide, Australia), and E. Kovacova (Institute of Virology, Slovak Academy of Sciences, Slovak Republic), 150 paired and 4 single sera from Japanese pneumonia patients in Okayama Prefecture with unknown fever and no clinical information regarding the microbes in those pneumonia cases, and 26

* Corresponding author. Mailing address: Laboratory of Rickettsia and Chlamydia, Department of Virology I, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-Ku, Tokyo 162-8640, Japan. Phone: 81-3-5285-1111, ext. 2563. Fax: 81-3-5285-1208. E-mail: ogawam@nih.go.jp.




FIG. 1. Determination of IgM and IgG titers to phase II *C. burnetii* in 346 serum samples by immunofluorescence assay. *, acute-phase sera from confirmed Q fever patients; ●, chronic- and convalescent-phase sera from confirmed Q fever patients; ▲, sera from Japanese pneumonia patients; ■, negative sera.

negative sera from healthy donors, including 1 serum sample from Slovakia and 25 sera from Japan.

***C. burnetii* antigen.** A total of 10^7 *C. burnetii* phase II strain Nine Mile (ATCC VR615) bacteria with a high passage number were purified by differential centrifugation and formalin treatment as described previously (20), with minor modifications. Briefly, after 3 to 5 days of inoculation of *C. burnetii* to Vero cell lines, the medium was discarded and the cells were collected with a cell scraper. The cells were then resuspended by Dounce homogenization in 0.02% formalin-phosphate-buffered saline (PBS) (pH 7.2) 20 times. The cell solution was centrifuged at $1,300 \times g$ for 5 min, and the supernatant was collected. Subsequently, the collected supernatant was filtered through a 5.0- μ m-pore-size filter (Millipore Corp., Bedford, MA) to remove soluble cell culture debris and centrifuged at $13,000 \times g$ at 4°C for 15 min. The supernatant was then discarded, and the pellets were washed with PBS twice. The concentration of *C. burnetii* antigen was measured with a Bio-Rad protein assay kit (Bio-Rad Lab., Hercules, CA). After microscopic examination of impression smears checked by the IFA test, the partially purified *C. burnetii* was pooled and stored at -80°C until use.

IFA test. The IFA test was performed by using prepared *C. burnetii* antigen with twofold dilutions of serum from 1:16 to 1:2,048 in PBS. In brief, 4 μ l of antigen was dotted in triplicate onto a clean 15-well multitest slide (ICN Bio-medicals, Inc., Aurora, Ohio) and allowed to air dry. Once dry, the slides were fixed in acetone for 15 min at room temperature. The diluted serum samples were then overlaid onto the antigen spots and incubated at 37°C in a humidified chamber for 1 h. After one wash with distilled water, two washes with 0.05%

PBS-Tween, and one more wash with distilled water, 8 μ l of fluorescein isothiocyanate-labeled anti-human IgM or IgG (Biosource, Camarillo, CA) diluted 1:200 in 0.001% Evans blue solution was added to each antigen spot and the slides were incubated for 1 h at 37°C. Finally, they were washed as before, dried in air, and mounted in 50% glycerol-PBS (pH 8.6). The slides were examined with a 40 \times objective ($\times 400$ magnification) using a fluorescence microscope (Axioskop 2 plus; Zeiss) equipped for visualizing brilliant green staining of *C. burnetii* microorganisms.

ELISA. A commercial ELISA kit (PanBio Co Ltd., Windsor, Queensland, Australia) was used as recommended by the manufacturer. The bound conjugates were detected by using tetramethyl benzidine as a substrate, and the color change was assessed in a microplate reader at a test wavelength of 450 nm. The ELISA index can be obtained by calculating the ratio of the cutoff absorbance to the sample absorbance and multiplying by 10. The serum samples were considered positive if they had an index of more than 11, equivocal if the index was between 9 and 11, and negative if the index was less than 9.

WBA. The outer membrane complex of *C. burnetii* was extracted by trichloroacetic acid and the temperature treatment method as described elsewhere (22). Sodium dodecyl sulfate-polyacrylamide gel electrophoresis was carried out with a 12% polyacrylamide gel as a separating gel (12), and a Western blotting assay (WBA) was performed using prepared outer membrane complex with 1:400 dilutions of the sera. The assay result was considered positive if the sera recognized the approximately 27-kDa protein of *C. burnetii*.

TABLE 1. Grouping of results

Group no.	Serum origin	No. of samples with result:			Total no. of:	
		Q fever	Negative	Equivocal	Serum samples	Patients
1	Confirmed Q fever patients	13		3	16	16
2	Japanese pneumonia patients		149 pairs	1 pairs, 4 single	304	150
3	Negative samples		26		26	4
Total					346	196

TABLE 2. Results of IFA test, ELISA, and Western blotting assay

Subtype	Test	Assay result ^a			
		Chromio- and convalescent-phase sera (4 samples)	Acute-phase sera (9 samples)	Negative	
				149 pairs of sera	26 samples
IgM	IFA	—	+	—	—
	ELISA	—	+	—	—
	WBA	—	+	—	—
IgG	IFA	+	+	—	—
	ELISA	+	+	—	—
	WBA	+	+	—	—

^a +, positive; —, negative. Equivocal results were obtained for three samples from confirmed Q fever cases and six samples (one paired and four single sera) from Japanese pneumonia patients. See Table 3 for details regarding equivocal results.

RESULTS

Serological analysis. Three hundred forty-six human sera, including 16 sera from confirmed Q fever cases, 304 sera from pneumonia patients, and 26 negative sera, were tested by the IFA test (Fig. 1), ELISA, and WBA (Table 1). Thirteen sera with an IgM titer of $\geq 1:128$ and/or an IgG titer of $\geq 1:256$ by the IFA test were positive by both ELISA and WBA, whereas 302 sera with an IgM titer of $\leq 1:16$ and an IgG titer of $\leq 1:32$ by the IFA test were negative by both ELISA and WBA. In the case of an IgM titer of $\geq 1:32$ and $\leq 1:64$ and/or an IgG titer of $\geq 1:64$ and $\leq 1:128$ by the IFA test, 4 sera were IgM and/or IgG positive by both ELISA and WBA, whereas 2 sera were negative by ELISA but positive by WBA. In this study, we labeled this area the “equivocal area” and divided the results of the IFA test into 3 groups, as shown in Table 1. The detailed results of the 3 groups are discussed in the next section.

Groups. Group I included 13 sera from confirmed Q fever cases. Four of these sera were IgG positive by both ELISA and WBA, indicating convalescent or chronic cases, whereas 9 of these sera were IgM and IgG positive by both ELISA and WBA, suggesting acute or subacute cases (Table 2). Group II included 298 sera from pneumonia patients and 26 confirmed negative sera that were negative by ELISA and WBA. Group III, which was proposed as the “equivocal area” in this study, included 3 sera from confirmed Q fever cases and 6 sera from pneumonia patients. The 3 confirmed Q fever sera and one serum from a pneumonia patient were IgM and/or IgG positive by both ELISA and WBA (Table 3). However, 2 of the sera from pneumonia patients were negative by ELISA but positive by WBA, and the other 3 sera were negative by both ELISA and WBA.

DISCUSSION

We evaluated new criteria for the IFA test in the diagnosis of Q fever in Japan. We used 346 human sera, including 16 from confirmed Q fever cases, 304 from Japanese pneumonia patients, and 26 that were negative. Thirteen sera from the confirmed Q fever cases were positive by the IFA test, ELISA, and WBA, whereas 298 sera from pneumonia patients and 26 negative sera were negative by the IFA test, ELISA, and WBA. In the proposed “equivocal area,” we found 9 sera with various

results by the IFA test, ELISA, and WBA for each serum sample.

The high cutoff value should emphasize the predictive value of a positive result with a high probability. In our temporary criteria, the combination of a phase II IgM titer of $\geq 1:128$ and/or a phase II IgG titer of $\geq 1:256$ gave a positive result for 13 sera from confirmed Q fever patients. Under these conditions, the diagnosis can be made even with only a single serum sample. Additionally, the serum samples recognized an approximately 27-kDa protein of *C. burnetii* by WBA, which other workers suggested as an immunodominant component in certain acute cases of the disease (18, 19), and were positive by ELISA. These criteria can be considered more reliable than those of a recent study that defined the high sensitivity of the IFA test at a cutoff titer of 1:400 (16).

The low cutoff value for either IgM or IgG should give a high predictive value of a negative result; thus, diagnosis cannot be made below this titer with a high probability. We may consider phase II IgM and phase II IgG titers of $\leq 1:16$ and $\leq 1:32$, respectively, as the low cutoff values (Fig. 1). Most samples under the low cutoff values, including 149 paired and 26 negative-control sera, were negative and also did not recognize the

TABLE 3. Confirmation of equivocal IFA test results with ELISA and Western blot assay

Subtype	Test	Result for sample no. with serum origin:							
		Confirmed Q fever patients			Japanese pneumonia patients				
		1	2	3	1 ^a	2	3	4 ^b	5 ^b
IgM	IFA ^c	32	<16	<16	32	32	32	64	64
	ELISA ^d	+	—	—	+	—	—	—	—
	WBA ^e	+	—	—	+	—	—	+	+
IgG	IFA	128	128	128	<16	<16	<16	128	128
	ELISA	+	+	+	—	—	—	—	—
	WBA	+	+	+	—	—	—	—	—

^a Probable acute case.

^b Suspected acute case.

^c The numbers show titers as twofold dilutions.

^d +, positive, if the index was more than 11 units; —, negative, if the index was less than 9 units.

^e +, positive, if sera recognized the approximately 27-kDa protein; —, negative, if sera did not recognize the protein.

TABLE 4. Confirmation of equivocal ELISA results with IFA test and Western blotting assay

Subtype	Test	Result for indicated sample no. with serum origin:																	
		Confirmed Q fever patients				Japanese clinical specimen													
		1	2	3	4	1 ^a	2	3	4	5	6	7	8	9	10	11	12	13	14
IgM	ELISA ^b	15.1	13.8	8.3	8.8	12.8	14	12.5	3.4	7.3	7.9	10.4	4.2	8.3	13.5	11.1	11.2	15.7	5.1
	IFA ^c	32	+	<16	<16	32	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16
	WB ^d	+	+	-	-	+	-	-	-	-	-	-	-	-	-	-	-	-	-
IgG	ELISA	14.7	14	15.1	14.7	9.7	4.3	5.2	15.8	13	14.8	11.9	19.4	14.1	4.1	2.7	4.5	4.6	12.2
	IFA	128	+	128	+	<16	32	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16	<16
	WB	+	+	+	+	-	-	-	-	-	-	-	-	-	-	-	-	-	-

^a Probable acute case.^b The numbers indicate the ELISA index.^c The numbers indicate the IFA titer of the diluted sera (see details in the text).^d +, positive; -, negative in the recognition of the approximately 27-kDa protein.

specific protein by WBA, although a few of them had a significant index by ELISA (Table 4). This is in accordance with a previous study showing that ELISA is suitable for use as a screening assay for Q fever diagnosis, with the IFA test used to confirm negative results (3, 4). We may explain this difference by the fact that a nonspecific reaction by ELISA may still occur due to cellular debris in the antigen preparation from culture. Although a recent study showed that LightCycler nested PCR can also be applied as a secondary tool in the diagnostic strategy for the early diagnosis of acute Q fever (5), the result of this study showed a good correlation between IFA titers and ELISA index values. A higher IFA titer correlated with a higher index in the ELISA result. Based on this clarification, confirmation with another serological test might not be required for samples categorized as negative by the IFA test.

We proposed the titer equivocal area and found 9 sera, including 3 from confirmed Q fever cases and 6 from Japanese pneumonia patients in the area, by the IFA test. Three sera from the confirmed Q fever cases and one serum from a pneumonia patient were positive by ELISA and WBA. However, 2 sera from pneumonia patients were negative by ELISA but positive by WBA, which left 3 sera negative by both ELISA and WBA. These results suggest that a single cutoff value for the IFA test may cause false-positive and false-negative results. In addition, only the IgM titer was positive, and the titer was very low in the positive case of pneumonia. This result suggests that serological assay with paired sera should be done for confirmation.

The results presented here illustrate the new criteria for the IFA test for Q fever. We recommended that an "equivocal area" should be used for the IFA test, rather than a single cutoff value and that sera in the equivocal area should be tested by additional serological assays to eliminate false-positive and false-negative results.

ACKNOWLEDGMENTS

We thank all the physicians who kindly provided the clinical samples for this study and Werner Stenczka, Barrie P. Marmion, and Elena Kovacova for kindly providing the control human sera.

This work was financially supported by The Japan Human Sciences Foundation.

REFERENCES

- Baca, O. G., and D. Paretsky. 1983. Q fever and *Coxiella burnetii*: a model for host-parasite interactions. *Microbiol. Rev.* 47:127-149.

- Brouqui, P., H. Tissot Dupont, M. Drancourt, Y. Berlan, J. Etienne, C. Lepout, F. Goldstein, P. Massip, M. Micoind, A. Bertrand, and D. Raoult. 1993. Chronic Q fever: 92 cases from France including 27 cases without endocarditis. *Arch. Intern. Med.* 153:642-648.
- Field, P. R., J. L. Mitchell, A. Santiago, D. J. Dickeson, S.-W. Chan, D. W. T. Ho, A. M. Murphy, A. J. Cuzzubbo, and P. L. Devine. 2000. Comparison of a commercial enzyme-linked immunosorbent assay with immunofluorescence and complement fixation tests for detection of *Coxiella burnetii* (Q fever) immunoglobulin M. *J. Clin. Microbiol.* 38:1645-1647.
- Field, P. R., A. Santiago, S.-W. Chan, D. B. Patel, D. Dickeson, J. L. Mitchell, P. L. Devine, and A. M. Murphy. 2002. Evaluation of a novel commercial enzyme-linked immunosorbent assay detecting *Coxiella burnetii*-specific immunoglobulin G for Q fever prevaccination screening and diagnosis. *J. Clin. Microbiol.* 40:3526-3529.
- Fournier, P. E., and D. Raoult. 2003. Comparison of PCR and serology assays for early diagnosis of acute Q fever. *J. Clin. Microbiol.* 41:5094-5098.
- Huldane, E. V., T. J. Marrie, R. S. Faulkner, S. H. Lee, J. H. Cooper, D. D. MacPherson, and T. J. Montague. 1983. Endocarditis due to Q fever in Nova Scotia: experience with five patients in 1981-1982. *J. Infect. Dis.* 148:978-985.
- Hendrix, L. R., L. P. Mallavia, and J. E. Samuel. 1993. Cloning and sequencing of *Coxiella burnetii* outer membrane protein gene *com1*. *Infect. Immun.* 61:470-477.
- Hunt, J. G., P. R. Field, and A. M. Murphy. 1983. Immunoglobulin responses to *Coxiella burnetii* (Q fever): single-serum diagnosis of acute infection, using an immunofluorescence technique. *Infect. Immun.* 39:977-981.
- Kenneth, O., and K. M. Yamada. 1977. Direct detection of antigens in sodium dodecyl sulfate-polyacrylamide gels. *Anal. Biochem.* 78:483-490.
- Kovacova, E., J. Gallo, S. Schramek, J. Kazar, and R. Brezina. 1987. *Coxiella burnetii* antigens for detection of Q fever antibodies by ELISA in human sera. *Acta Virol.* 31:254-259.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685.
- Murphy, A. M., and L. Magro. 1980. IgM globulin response in Q fever (*Coxiella burnetii*) infections. *Pathology* 12:391-396.
- Ogawa, M., T. Kawamoto, A. Kawamoto, T. Yamashita, Y. Uchida, K. Kato, A. Setiyono, S. Shiga, and T. Kishimoto. 2003. Time course of the levels of antibodies to *Coxiella burnetii* and detection of *C. burnetii*-DNA in three imported cases of acute Q fever. *Kansenshogaku Zasshi* 77:127-132. (In Japanese.)
- Peacock, M. G., R. N. Philip, J. C. Williams, and R. S. Faulkner. 1983. Serological evaluation of Q fever in humans: enhanced phase I titers of immunoglobulins G and A are diagnostic for Q fever endocarditis. *Infect. Immun.* 41:1089-1098.
- Peter, O., G. Dupuis, W. Burgdorfer, and M. Peacock. 1985. Evaluation of complement fixation and indirect immunofluorescence tests in the early diagnosis of primary Q fever. *Eur. J. Clin. Microbiol.* 4:394-396.
- Rollain, J. M., C. Lecam, and D. Raoult. 2003. Simplified serological diagnosis of endocarditis due to *Coxiella burnetii* and *Bartonella*. *Clin. Diagn. Lab. Immunol.* 10:1147-1148.
- Stocker, M. G. P., and P. Fiset. 1956. Phase variation of the Nine Mile and other strains of *Rickettsia burnetii*. *Can. J. Microbiol.* 2:310-321.
- To, H., A. Hotta, G. Q. Zhang, S. V. Nguyen, M. Ogawa, T. Yamaguchi, H. Fukushi, K. Amano, and K. Hirai. 1997. Antigenic characteristics of polypeptides of *Coxiella burnetii* isolates. *Microbiol. Immunol.* 42:81-85.
- To, H., A. Hotta, T. Yamaguchi, H. Fukushi, and K. Hirai. 1997. Antigenic characteristics of the lipopolysaccharides of *Coxiella burnetii* isolates. *J. Vet. Med. Sci.* 60:267-270.
- To, H., K. K. Htwe, N. Yamasaki, G. Q. Zhang, M. Ogawa, T. Yamaguchi, H.

- Fukushi, and K. Hirai. 1995. Isolation of *Coxiella burnetii* from dairy cattle and ticks, and some characteristics of the isolates in Japan. *Microbiol. Immunol.* 39:663-671.
21. Turk, W. P., G. Howitt, L. A. Turnberg, H. Fox, M. Longson, M. B. Matthews, and R. DasGupta. 1976. Chronic Q fever. *Q. J. Med.* 45:193-217.
22. Vavrekova, M., F. Ciampor, and M. Lukacova. 1992. Electron-microscopic study of the effect of various extractants on the morphology of *Coxiella burnetii*. *Folia Microbiol.* 37:87-92.
23. Zhang, G. Q., A. Hotta, T. Ho, T. Yamaguchi, H. Fukushi, and K. Hirai. 1998. Evaluation of a recombinant 27-kDa outer membrane protein of *Coxiella burnetii* as an immunodiagnostic reagent. *Microbiol. Immunol.* 42:423-428.