

That is, the SVR rates were still high for the aged patients of the non-1H group, but lower for the aged patients than the young patients in the 1H group. There are two possible reasons for this. First, the number of patients with no reduction of both drugs was significantly fewer for the patients aged 60–64 years and <60 years than for the patients aged ≥ 65 years, and the older patients tended to require ribavirin reduction or discontinuance (Fig. 3). Second, the liver fibrosis score tended to be higher in aged patients than in young patients, although the significant difference was not seen in this study (Table 1). These factors can decrease the SVR rates in aged patients in the 1H group, from which it is difficult to eliminate the virus, although the aged patients in the non-1H group whose viruses are easily eliminated were not affected. The results on ITT analysis account for the conclusion of the indication for IFN and ribavirin combination therapy of 24 weeks for aged patients; the patients of the 1H group do not have good application whose SVR is approximately 10%. On the other hand, patients of the non-1H group should be given the combination therapy because of the higher SVR rates of about 65%.

Better efficacy of treatments using new drugs, such as peg-IFN and ribavirin combination therapy or NS3/4 protease inhibitor, is greatly anticipated.

Acknowledgments

Other institutions and participants in the Osaka Liver Disease Study Group (Digestive Disease Study Group of Osaka Renaissance) are: National Hospital Organization Osaka National Hospital, Y. Izumi; Osaka Rousai Hospital, H. Aketa and K. Noda; Osaka Kouseinenkin Hospital, M. Kurokawa and T. Akasaka; Kansai Rousai Hospital, M. Yamamoto; Osaka General Medical Center, T. Inoue; National Hospital Organization Osaka Minami Medical Center, H. Hikita and M. Shigekawa; Osaka Police Hospital, J. Kondo; Kaizuka City Hospital, O. Nishiyama; and Osaka University Graduate School of Medicine, S. Shinzaki, M. Miyazaki, H. Miyatake, I. Itose, S. Egawa and T. Nishida.

This work was supported by a Grant-in-Aid for Research on Hepatitis and BSE from the Ministry of Health Labour and Welfare of Japan, and Scientific Research from the Ministry of Education, Science, and Culture of Japan.

References

- [1] Lauer G, Walker BD. Hepatitis C virus infection. *N Engl J Med* 2001;345:41–52.
- [2] Poynard T, Bedossa P, Opolon P, et al. Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet* 1997;349:825–32.
- [3] Hamada H, Yatsushashi H, Yano K, et al. Impact of aging on the development of hepatocellular carcinoma in patients with posttransfusion chronic hepatitis C. *Cancer* 2002;95(2):331–9.
- [4] Tanaka H, Tsukuma H. Characteristics of Japanese patients with liver cancer—epidemiological study based on comparison between male and female patients. *Hepatology Res* 2002;24:S11–20.
- [5] Yoshizawa H. Trends of hepatitis virus carriers. *Hepatology Res* 2002;24:S28–39.
- [6] Shiratori Y, Imazeki F, Moriyama M, et al. Histologic improvement of fibrosis in patients with hepatitis C who have sustained response to interferon therapy. *Ann Intern Med* 2000;132:517–24.
- [7] Hiramatsu N, Hayashi N, Kasahara A, et al. Improvement of liver fibrosis in chronic hepatitis C patients treated with natural interferon alpha. *J Hepatol* 1995;22:135–42.
- [8] Yoshida H, Shiratori Y, Moriyama M, et al. Interferon therapy reduces the risk for hepatocellular carcinoma: National surveillance program of chronic and noncirrhotic patients with chronic hepatitis C in Japan. *Ann Intern Med* 1999;131:174–81.
- [9] Kasahara A, Hayashi N, Mochizuki K, et al. Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998;27:1394–402.
- [10] Ikeda K, Saitoh S, Arase Y, et al. Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C: A long-term observation study of 1,643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* 1999;19:1124–30.
- [11] Kasahara A, Tanaka H, Okanoue T, et al. Interferon treatment improves survival in chronic hepatitis C patients showing biochemical as well as virological responses by preventing liver-related death. *J Viral Hepat* 2004;11(2):148–56.
- [12] Yoshida H, Arakawa Y, Sata M, et al. Interferon therapy prolonged life expectancy among chronic hepatitis C patients. *Gastroenterology* 2002;123:4483–91.
- [13] Imai Y, Kasahara A, Tanaka H, et al. Interferon therapy for aged patients with chronic hepatitis C: improved survival in patients exhibiting a biochemical response. *J Gastroenterol* 2004;39:1069–77.
- [14] McHutchison JG, Gordon SC, Schiff ER, et al. Interferon alpha 2b alone or combination with ribavirin as initial treatment for chronic hepatitis C. *N Engl J Med* 1998;229:1485–92.
- [15] Poynard T, Marcellin P, Lee SS, et al. Randomised trial of interferon alpha2b plus ribavirin for 48 weeks or 24 weeks versus interferon alpha2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. *Lancet* 1998;352:1426–32.
- [16] Hiramatsu N, Kasahara A, Nakanishi F, et al. The significance of interferon and ribavirin combination therapy followed by interferon monotherapy for patients with chronic hepatitis C in Japan. *Hepatol Res* 2004;29(3):142–7.
- [17] Bodenheimer HC, Lindsay KL, Davis GL, et al. Tolerance and efficacy of oral ribavirin treatment of chronic hepatitis C: A multicenter trial. *Hepatology* 1997;26:437–77.
- [18] Okanoue T, Sakamoto S, Itoh Y, et al. Side effects of high-dose interferon therapy for chronic hepatitis C. *J Hepatol* 1996;25:283–91.
- [19] Yuki N, Hayashi N, Kasahara A, et al. Pretreatment viral load and response to prolonged interferon- α course for chronic hepatitis C. *J Hepatol* 1995;22:457–63.
- [20] Lau YN, Davis G, Kniffen J, et al. Significance of serum hepatitis C virus RNA levels in chronic hepatitis C. *Lancet* 1993;341:1501–4.
- [21] Shiratori Y, Kato N, Yokosuka O, et al. Predictors of the efficacy of interferon therapy in chronic hepatitis C virus infection. *Gastroenterology* 1997;113:558–66.
- [22] Tanaka T, Tsukiyama-Kohara K, Yamaguchi K, et al. Significance of specific antibody assay for genotyping of hepatitis C virus. *Hepatology* 1994;19:1347–53.
- [23] Lindahl K, Schwarcz R, Bruchfeld A, et al. Evidence that plasma concentration rather than dose per kilogram body weight predicts ribavirin-induced anemia. *J Viral Hepatitis* 2004;11:84–7.
- [24] Maeda Y, Kiribayashi Y, Moriya T, et al. Dosage adjustment of ribavirin based on renal function in Japanese patients with chronic hepatitis C. *Ther Drug Monit* 2004;26:9–15.

Early decline of hemoglobin correlates with progression of ribavirin-induced hemolytic anemia during interferon plus ribavirin combination therapy in patients with chronic hepatitis C

TSUGIKO OZE¹, NAOKI HIRAMATSU¹, NAO KURASHIGE¹, NATSUKO TSUDA¹, TAKAYUKI YAKUSHIJI¹, TATSUYA KANTO¹, TETSUO TAKEHARA¹, AKINORI KASAHARA¹, MICHIO KATO², HARUMASA YOSHIHARA³, KAZUHIRO KATAYAMA⁴, SHINJI KUBOTA⁵, TAIZO HIJIOKA⁶, KAZUNOBU ISHIBASHI⁷, MASAHIDE OSHITA⁸, HIDEKI HAGIWARA⁹, YOSHIMICHI HARUNA¹⁰, EIJI MITA¹¹, SHINJI TAMURA¹, and NORIO HAYASHI¹

¹Department of Gastroenterology and Hepatology, Osaka University Graduate School of Medicine, 2-2 Yamadaoka, Suita 565-0871, Japan

²National Hospital Organization Osaka National Hospital, Osaka, Japan

³Osaka Rousai Hospital, Sakai, Japan

⁴Osaka Kouseinenkin Hospital, Osaka, Japan

⁵Kansai Rousai Hospital, Amagasaki, Japan

⁶National Hospital Organization Osaka Minami Medical Center, Kawachinagano, Japan

⁷Kaizuka City Hospital, Kaizuka, Japan

⁸Osaka Police Hospital, Osaka, Japan

⁹Higashiosaka City Central Hospital, Higashiosaka, Japan

¹⁰Osaka General Medical Center, Osaka, Japan

¹¹Saiseikai Senri Hospital, Suita, Japan

Background. The aim of this study was to examine the factors correlated with the progression of ribavirin-induced hemolytic anemia in patients with chronic hepatitis C treated by interferon and ribavirin combination therapy. **Methods.** This study was conducted on 505 patients by the Osaka Liver Disease Study Group. A decline of hemoglobin (Hb) concentration by 2 g/dl at the end of 2 weeks from the start of the treatment (“2 by 2” standard) was adopted as a predictive factor for progression to severe anemia. The ribavirin apparent clearance (CL/F) was also examined. **Results.** Of 482 patients whose Hb value was more than 12 g/dl before the treatment, 68 patients (14%) had to discontinue ribavirin owing to severe anemia. Patients in the “2 by 2”-positive group (Hb decline over 2 g/dl) and the group with lower CL/F were significantly more likely to discontinue ribavirin owing to severe anemia. Discontinuation was more common among patients aged 60 years or older than for those under 60 years old (21% vs. 9%, $P < 0.001$). Among patients aged 60 years or older, only the “2 by 2” standard was significantly associated with the discontinuance of ribavirin owing to severe anemia in a multivariate analysis (odds ratio, 4.18; $P < 0.001$). **Conclusions.** The “2 by 2” standard of Hb decline can be used to identify patients likely to develop severe anemia. The early reduction of ribavirin can help prevent progression to severe anemia, thus allowing ribavirin therapy to be completed even in older patients.

Key words: chronic hepatitis C, interferon and ribavirin combination therapy, progression of anemia, “2 by 2” standard

Introduction

Hepatitis C virus (HCV) is estimated to infect up to 170 million people worldwide,¹ and two million people in Japan. Long persistence of HCV infection can lead to progression of liver fibrosis, causing liver cirrhosis and ultimately hepatocellular carcinoma.^{2,3} Past studies have made clear that interferon (IFN) therapy is effective for eliminating HCV,^{4,5} but the sustained viral response (SVR) rate of IFN monotherapy is not sufficient. The addition of the nucleoside analog ribavirin to IFN in the treatment of patients with chronic hepatitis C can significantly improve the SVR rate, and combination therapy with IFN or pegylated-IFN (Peg-IFN) has been recommended as a standard regimen worldwide.^{6–10} However, additional side effects of ribavirin have been reported, such as hemolytic anemia, which have not been found with IFN monotherapy, leading to discontinuance of the treatment.^{11–14}

In previous studies, the discontinuance rate of IFN and ribavirin combination treatment due to severe side effects has been reported to be 6%–13%.^{6,7} Ribavirin-induced hemolytic anemia has been suggested to depend on a high plasma concentration of ribavirin.¹⁵ The ribavirin apparent clearance (CL/F), which reflects the plasma concentration of ribavirin at 4 weeks after the start of combination therapy, has been used as a

Received: March 30, 2006 / Accepted: June 12, 2006

Reprint requests to: N. Hiramatsu

predictive factor for ribavirin-induced hemolytic anemia before the start of treatment.¹⁶⁻¹⁸ Furthermore, in the manufacturer's drug information for ribavirin,¹⁹ a dose reduction is recommended when hemoglobin (Hb) levels decrease to less than 10 g/dl, and discontinuance of ribavirin is recommended when Hb levels fall to less than 8.5 g/dl during combination therapy with IFN and ribavirin. However, according to this guideline, not a few patients are forced to discontinue ribavirin because the dose reduction to avoid severe anemia does not occur in time.

What is needed is a convenient guideline for avoiding ribavirin discontinuance due to severe anemia. In this study, we evaluated the correlation of Hb decline at 2 weeks after the start of combination therapy with the discontinuance of treatment due to progression of ribavirin-induced hemolytic anemia. We also assessed the utility of an early decline of Hb in comparison with the CL/F standard for predicting the progression to severe anemia.

Patients and methods

Patients

The current study was conducted at Osaka University Hospital and other institutions participating in the Osaka Liver Disease Study Group. The 505 patients with chronic hepatitis C included in this study were treated with a combination of interferon- α -2b and ribavirin between January 2001 and December 2005. All patients were anti-hepatitis C virus antibody positive, had HCV RNA detectable in their serum by the polymerase chain reaction method, and had elevated serum alanine transaminase (ALT) (above the upper limit of normal) within the 6 months prior to treatment.

Excluded from this study were patients who were positive for hepatitis B surface antigen or anti-human immunodeficiency virus antibody or those with other forms of liver disease (alcoholic liver disease, hepatotoxic drugs, autoimmune hepatitis). Twenty-three patients whose Hb was under 12 g/dl before the treatment were also excluded because the aim of this study was to analyze the progression of anemia; patients with a low Hb level before treatment are known to have a tendency toward progression of anemia. The remaining 482 patients were followed in this study.

The baseline clinical features of the 482 patients are shown in Table 1. Their mean age was 55.2 ± 10.9 years, and 66% were men. Among the patients, 347 had HCV RNA with genotype 1 and high viral loads (1H group) and 130 had HCV RNA with genotype 2 or low viral loads (non-1H group). The mean ALT level was 100 ± 74 IU/l. In this study, a high viral load was defined as a serum HCV-RNA level of more than 10^6 equivalents/ml by branched DNA assay or more than 10^5 copies/ml serum by Amplicor-HCV monitor assay.

Treatment schedule

Of the 482 patients treated with a combination of interferon- α -2b and ribavirin, 273 were IFN naïve and 209 were undergoing retreatment. All patients were scheduled to receive interferon- α -2b (Intron-A, Schering-Plough, Kenilworth, NJ, USA) at a dose of 6 ($n = 371$) or 10 ($n = 111$) MU intramuscularly every day for the first 2 weeks and three times a week thereafter. Ribavirin (Rebetol; Schering-Plough) was given orally twice a day for a total dose of 800 mg ($n = 261$), 600 mg ($n = 215$), or 400 mg ($n = 6$) per day. The IFN dose was decreased from 10 to 6 MU or from 6 to 3 MU when the

Table 1. Baseline characteristics of patients

Number	482	
Age (y.o)	55.2 ± 10.9	(21-75)
Sex (male/female)	320/162	
Body weight (kg)	62.3 ± 9.9	(35-94)
HCV serotype (1/2/unknown)	364/111/7	
(1H/non-1H/unknown)	347/130/5	
Fibrosis (0/1/3/4/unknown)	19/192/202/13/56	
WBC (/mm ³)	5184 ± 1531	(2100-13200)
RBC ($\times 10^4$ /mm ³)	449 ± 42	(329-617)
Hb (g/dl)	14.4 ± 1.2	(12.0-19.2)
Plt ($\times 10^4$ /mm ³)	15.4 ± 5.4	(4.4-36.1)
ALT (IU/l)	100 ± 74	(17-736)
Serum creatinine (mg/dl)	0.8 ± 0.2	(0.3-1.7)
Ribavirin dosage/body weight (mg/kg)	11.4 ± 1.5	(4.6-17.8)

Data are shown as means \pm SD

HCV, hepatitis C virus; 1H group, patients with genotype 1 and high viral load; non-1H group, patients not in the 1H group; Fibrosis, Knodell's histological score (category4); WBC, white blood cells; RBC, red blood cells; Hb, hemoglobin; Plt, platelets; ALT, alanine aminotransferase

white blood cell (WBC) count was below 1500/mm³, the neutrocyte count below 750/mm³, or the platelet (Plt) count below 5×10^4 /mm³. IFN was discontinued when the WBC count was below 1000/mm³, the neutrocyte count below 500/mm³, or the Plt count below 2.5×10^4 /mm³. The ribavirin dose of 200mg was reduced when the Hb concentration decreased to less than 10g/dl, and the ribavirin was discontinued when the Hb concentration decreased to less than 8.5g/dl, in accordance with the manufacturer's drug information for ribavirin.¹⁹ Ferric medicine or erythropoietin to prevent anemia was not administered. Ribavirin was scheduled to be administered for 24 weeks for all patients, and IFN for 24 weeks for 307 patients and for 48 weeks for 175 patients.

Patients with persistently undetectable HCV RNA 6 months after completion of treatment were considered to have achieved SVR.

Blood tests

All patients were examined for serum HCV-RNA level and underwent hematological and biochemical tests just before therapy, at the end of week 2, and every 4 weeks thereafter during treatment. When treatment was completed, the patients were assessed every 4 weeks until 24 weeks after the end of treatment.

Total ribavirin clearance

Using the method of Kamar et al.,¹⁷ CL/F at the start of the treatment was calculated as follows:

$$\text{CL/F (l/h)} = 32.3 \times \text{BW} \times (1 - 0.0094 \times \text{Age}) \\ \times (1 - 0.42 \times \text{Sex})/\text{Scr},$$

where BW = body weight; sex = 0 for male and 1 for female; and Scr = serum creatinine.

Definition of "severe anemia" leading to discontinuance of ribavirin

In this study, "discontinuance of ribavirin due to severe anemia" was defined as follows: discontinuance of ribavirin due to a decrease of Hb to less than 8.5g/dl or clinical symptoms of anemia associated with a decrease of Hb of more than 3g/dl from the start of combination therapy.

Liver histology

Hepatic fibrosis was assessed by Knodell's histological score (category 4).²⁰ Fibrosis stage was evaluated on a scale from 0 to 4: 0 = no fibrosis; 1 = fibrosis portal expansion; 3 = bridging fibrosis (portal-portal or portal-central linkage); 4 = cirrhosis.

Statistical analysis

Age, body weight, ribavirin dosage/body weight, WBC count, red blood cell (RBC) count, Hb concentration, Plt, serum ALT levels, and Scr are expressed as means \pm SD. The SVR rate was evaluated using an intention-to-treat (ITT) analysis. The differences in proportions were tested by the χ -squared test. For univariate and multivariate analyses, a logistic regression analysis was used to predict ribavirin-induced severe anemia. A value of $P < 0.05$ (two-tailed) was considered to indicate significance.

Results

Efficacy of the combination therapy with dose reduction or discontinuance of ribavirin

The relationship between dose reduction or discontinuance of ribavirin and the SVR rate on ITT analysis is shown in Fig. 1. The SVR rate was 20% (71/347) for all 1H patients and 72% (93/130) for all non-1H patients. Among the 1H patients, SVR was achieved for 24% (45/189) without dose reduction of ribavirin and for 26% (20/76) with dose reduction. Significantly lower SVR rates were observed for patients who had to discontinue ribavirin treatment owing to adverse effects (7%, 6/82) in comparison with those with ($P < 0.01$) or without ($P < 0.01$) dose reduction. In the non-1H group, similar SVR rates were found with dose reduction of ribavirin [SVR rate without dose reduction, 83% (58/70), vs. SVR rate with dose reduction, 82% (23/28)], and the SVR rate of patients who had to discontinue ribavirin owing to adverse effects was significantly lower (38%, 12/32) than that for those with ($P < 0.001$) or without ($P < 0.0001$) dose reduction.

The same tendency was observed even in the 307 patients treated with IFN for 24 weeks. Among the 1H patients treated for 24 weeks, SVR was achieved for 19% (17/91) without dose reduction of ribavirin, 15% (6/41) with dose reduction, and 3% (2/75) with discontinuance. There were significant differences between the patients with discontinuance and those without ($P < 0.01$) or with ($P < 0.05$) dose reduction. Among the non-1H patients treated for 24 weeks, SVR rates were 85% (39/46) for the patients without dose reduction of ribavirin, 85% (17/20) for those with dose reduction, and 33% (10/30) for those with discontinuance. Significantly lower SVR rates were observed for patients who had to discontinue ribavirin than for those with ($P = 0.05$) or without ($P < 0.05$) dose reduction.




Fig. 1. Efficacy of combination therapy with dose reduction or discontinuance of ribavirin (intention-to-treat analysis). *1H group*, patients with genotype 1 and high viral load; *non-1H group*, patients not in the 1H group; SVR, sustained viral response. □ all patients; ▨ patients without dose reduction of ribavirin; ▩ patients with dose reduction of ribavirin; ■ patients with discontinuance of ribavirin. *, $P < 0.01$; **, $P < 0.0001$; ***, $P < 0.001$

Table 2. Rate of the ribavirin reduction or discontinuance due to adverse effects with different levels of CL/F

	No reduction	Dose reduction	Discontinuance	
			All cases	Cases due to severe anemia
20 ≤ CL/F (n = 45)	94% (42/45)	2% (1/45)	4% (2/45)	0% (0/45)
15 ≤ CL/F < 20 (n = 100)	66% (66/100)	19% (19/100)	15% (15/100)	6% (6/100)
10 ≤ CL/F < 15 (n = 179)	54% (96/179)	24% (42/179)	23% (41/179)	14% (25/179)
CL/F < 10 (n = 158)	37% (58/158)	28% (44/158)	35% (56/158)	23% (37/158)

Frequency of and reasons for dose reduction or discontinuance of ribavirin during combination therapy

We examined the rate of discontinuance of therapy due to adverse effects up to the end of 24 weeks, because all cases of discontinuation occurred before the end of 24 weeks. Of the 482 patients, 401 patients completed 24 weeks of therapy, and 81 patients (17%) had to discontinue both IFN and ribavirin before the end of the 24 weeks. Of the 401 patients undergoing 24 weeks of therapy, the entire treatment schedule without reduction or discontinuance of either drug was completed by 262 patients (54%). The ribavirin dose was decreased for 106 patients (22%) and was stopped without discontinuance of IFN for 33 patients (7%). Overall, 114 patients (24%) discontinued ribavirin treatment. The reasons for dose reduction or discontinuance of ribavirin were anemia, general fatigue, digestive disorder, eczema, neutropenia, thrombocytopenia, or psychological disorder. Among the patients discontinuing

ribavirin, the major reasons were anemia (14%), general fatigue (2%), or digestive disorder (2%).

CL/F and dose reduction or discontinuance of ribavirin

CL/F calculated for all patients was 4.6–32.5 l/h. The mean CL/F was 13.0 l/h, and the median was 11.9 l/h. At the start of treatment, CL/F was less than 10 l/h for 33% (158/482) of patients, 10–15 l/h for 37% (179/482), 15–20 l/h for 21% (100/482), and more 20 l/h for 9% (45/486).

Table 2 shows the rates of dose reduction or discontinuance of ribavirin in relation to different levels of CL/F. The rate of discontinuance of ribavirin among all patients was 4% (2/45) for patients with CL/F ≥ 20, 15% (15/100) for those with 15 ≤ CL/F < 20, 23% (41/179) for those with 10 ≤ CL/F < 15, and 35% (56/158) for those with CL/F < 10. The rate of discontinuance of ribavirin due to severe anemia was 14% (68/482) among all pa-

tients. There was no discontinuance of ribavirin due to severe anemia among patients with $CL/F \geq 20$, but the rate of discontinuance was 6% (6/100) among those with $15 \leq CL/F < 20$, 14% (25/179) among those with $10 \leq CL/F < 15$, and 23% (37/158) among those with $CL/F < 10$. The rate of continuance of ribavirin without dose reduction decreased in proportion to the decline of CL/F . In this study, we adopted two categories of CL/F , below 15l/h ($CL/F < 15$) and below 10l/h ($CL/F < 10$), to assess CL/F as a factor for predicting anemia progression.

We also analyzed the predictive factor of anemia progression according to patient age, because CL/F varies widely with patient age and tends to be lower among older patients. Among patients under 60 years old ($n = 288$), 17% (48/288) had CL/F under 10l/h, 38% (109/288) had CL/F 10–15l/h, 30% (86/288) had CL/F 15–20l/h, and 16% (45/288) had CL/F over 20l/h. On the other hand, among those 60 years old or older ($n = 194$), 57% (110/194) had CL/F under 10l/h, 36% (70/194) had CL/F 10–15l/h, 7% (14/194) had CL/F 15–20l/h, and none had CL/F over 20l/h. Thus, the majority (93%) of the patients 60 years old or older had a low CL/F (<15), whereas only 55% of those under 60 years old had $CL/F < 15$.

Early decline of Hb and progression of anemia during combination therapy

Figure 2 shows the decline of Hb from the start of combination therapy. We conducted this analysis for the 433 patients: those who did not need a dose reduction of ribavirin ($n = 262$), those who needed a dose reduction owing to a decrease of Hb to less than 10 g/dl ($n = 103$), and those who discontinued ribavirin due to "severe anemia" ($n = 68$). We excluded 49 patients from this analysis: 46 patients stopped combination therapy

for reasons other than anemia, such as general fatigue or digestive disorder, and the other three patients were not responding to antiviral treatment and stopped therapy before 24 weeks without a dose reduction of ribavirin. Following the initiation of combination therapy, Hb concentration decreased rapidly until the end of the 4th week. At the end of 2 weeks, Hb had decreased by 0.9 ± 1.2 g/dl among the patients without dose reduction of ribavirin, by 1.8 ± 1.3 g/dl among those with dose reduction, and by 2.3 ± 1.4 g/dl among those who discontinued ribavirin. At the end of 4 weeks, Hb had decreased by 2.1 ± 1.5 g/dl among the patients without dose reduction of ribavirin, by 3.2 ± 1.5 g/dl among those with dose reduction, and by 3.9 ± 1.5 g/dl among those discontinuing ribavirin.

ΔHb [$\Delta Hb = (\text{Hb value just before treatment}) - (\text{Hb value during treatment})$] both at the end of 2 weeks and at the end of 4 weeks were significantly larger among the patients discontinuing ribavirin than among those without dose reduction of ribavirin ($P < 0.0001$, $P < 0.0001$, respectively). In this study, we adopted the category of ΔHb at the end of 2 weeks because it allowed the progression of anemia to be estimated at an earlier phase of treatment than did ΔHb at the end of 4 weeks.

To establish the cutoff value of ΔHb at the end of 2 weeks, we used two categories of ΔHb : a decrease in Hb concentration at 2 weeks to 2 g/dl below the baseline ($\Delta Hb2.0$) or to 1.5 g/dl below the baseline ($\Delta Hb1.5$). We conducted this analysis for 480 patients, because two patients stopped combination therapy before 2 weeks for reasons other than anemia. With the $\Delta Hb2.0$ standard, the rate of discontinuance of ribavirin due to severe anemia was 10% (32/338) in the $\Delta Hb < 2.0$ group and 25% (36/142) in the $\Delta Hb \geq 2.0$ group, with the difference being significant ($P < 0.0001$) (Table 3). With the $\Delta Hb1.5$ standard, the rate of discontinuance of ribavirin due to severe anemia was significantly higher




Fig. 2. Decline of hemoglobin according to dose reduction or discontinuance of ribavirin. *Significantly different from patients with dose reduction ($P < 0.0001$) and patients with discontinuance ($P < 0.0001$); **significantly different from patients with discontinuance ($P < 0.02$); †significantly different from patients with dose reduction ($P < 0.0001$) and patients with discontinuance ($P < 0.0001$); ††significantly different from patients with discontinuance ($P < 0.01$)

Table 3. Rate of the ribavirin reduction or discontinuance due to adverse effects with rate of anemia progression

	No reduction	Dose reduction	Discontinuance	
			All cases	Cases due to severe anemia
$\Delta\text{Hb} \geq 2.0$ ($n = 142$)	37% (53/142)	29% (41/142)	34% (48/142)	25%* (36/142)
$\Delta\text{Hb} < 2.0$ ($n = 338$)	61% (209/338)	19% (65/338)	20% (64/338)	10% (32/338)

* $P < 0.0001$




Fig. 3. Receiver-operating characteristic curve for ΔHb at the end of 2 weeks for discontinuance of ribavirin due to severe anemia

in the $\Delta\text{Hb} \geq 1.5$ group than in the $\Delta\text{Hb} < 1.5$ group (8%, 22/279 vs. 23%, 46/201; $P < 0.0001$). Figure 3 shows the receiver-operating characteristic curve using ΔHb at the end of 2 weeks for the discontinuance of ribavirin due to severe anemia. Between the $\Delta\text{Hb}2.0$ and $\Delta\text{Hb}1.5$ standards, no significant difference was found in sensitivity (53%, 36/68, vs. 68%, 46/68; NS). On the other hand, the false positive rate was significantly lower with the $\Delta\text{Hb}2.0$ standard than with the $\Delta\text{Hb}1.5$ standard (26%, 93/360, vs. 38%, 136/360; $P < 0.001$), and accuracy was significantly higher with the $\Delta\text{Hb}2.0$ standard than with the $\Delta\text{Hb}1.5$ standard (71%, 303/428, vs. 63%, 270/428; $P = 0.02$). Therefore, we adopted $\Delta\text{Hb}2.0$ at the end of 2 weeks (the “2 by 2” standard) as a predictive factor for discontinuance of ribavirin due to severe anemia because of the higher specificity rate of $\Delta\text{Hb}2.0$ (lower false positive rate).

Logistic regression analysis for discontinuance of ribavirin in combination therapy

We assessed the factors correlated with the discontinuance of ribavirin due to severe anemia by logistic regression analysis. The following factors were evaluated: age, sex, body weight, ribavirin dosage/body weight, IFN dosage, Scr, Hb value at the start of the therapy, CL/F category, and early decline of Hb (“2 by 2” standard). Older age, lower body weight, lower Hb at the start of the therapy, lower CL/F (CL/F < 10 or CL/F < 15), and “2 by 2”-positive (the patients whose Hb had decreased by more than 2 g/dl at 2 weeks from the start of the treatment) were factors significantly associated with discontinuance of ribavirin due to severe anemia by univariate logistic regression analysis (Table 4). Next, we assessed the factors correlated with the discontinuance of ribavirin due to severe anemia by multivariate logistic regression analysis. Among the factors selected as significant by the univariate analysis, we omitted age and body weight from the multivariate analysis because they were included as parameters in the numerical formula for CL/F. Therefore, we evaluated the Hb value at the start of therapy, the CL/F category, and the “2 by 2” category by multivariate analysis. The CL/F borderline values of 10l/h and 15l/h were evaluated separately. In the multivariate logistic regression analysis, lower Hb at the start of therapy, lower CL/F (CL/F < 10 or CL/F < 15), and “2 by 2”-positive were significantly associated with discontinuance of ribavirin due to severe anemia (Table 5).

Useful predictive factors for discontinuance of ribavirin among older patients

Among the 288 patients under 60 years old, 50 (17%) had discontinued ribavirin by the end of 24 weeks for various reasons, including anemia, general fatigue, digestive disorder, and psychological disorders. Among the 194 patients aged 60 years and older, 64 (33%) had discontinued ribavirin, with severe anemia accounting for approximately 65% (41/64). More than twice as many patients aged 60 years and older discontinued ribavirin treatment compared with younger patients;

Table 4. Univariate analysis for the discontinuance of ribavirin due to severe anemia

Factor	Category	Odds ratio	95% CI	P value
Age			1.045–1.117	<0.0001
Sex	Male/Female	1/1.18	0.663–2.029	0.56
Body weight			0.928–0.981	<0.001
Serum creatinine			0.551–9.492	0.25
Ribavirin/Body weight			0.945–1.357	0.18
IFN dosage	6MU/10MU	1/1.03	0.557–1.893	0.93
Hb			0.480–0.780	<0.0001
CL/F	≥15/<15	1/5.56	0.076–0.427	0.0001
	≥10/<10	1/3.14	0.187–0.540	<0.0001
"2 by 2"	Negative/Positive	1/3.23	0.182–0.527	<0.0001

CI, confidence interval; IFN, interferon; CL/F, apparent clearance; "2 by 2", ΔHb2.0 at the end of 2 weeks; "2 by 2"-positive means ΔHb ≥ 2.0; "2 by 2"-negative means ΔHb < 2.0

Table 5. Multivariate analysis for the discontinuance of ribavirin due to severe anemia

Factor	Category	Odds ratio	95% CI	P value
Hb			0.446–0.785	0.0003
CL/F	≥15/<15	1/3.18	0.126–0.786	0.01
"2 by 2"	Negative/Positive	1/4.35	0.127–0.419	<0.0001
Hb			0.440–0.784	0.0003
CL/F	≥10/<10	1/1.98	0.278–0.923	0.03
"2 by 2"	Negative/Positive	1/4.63	0.119–0.393	<0.0001

this difference was significant (21%, 41/194, vs. 9%, 27/288; $P = 0.0003$) (Table 6).

We assessed the analysis for discontinuance of ribavirin due to severe anemia among the patients aged 60 years or older. Older age, lower CL/F (CL/F < 10), and "2 by 2"-positive were factors significantly associated with discontinuance of ribavirin due to severe anemia by univariate logistic regression analysis (Table 7A). Next, we assessed the factors correlated with the discontinuance of ribavirin due to severe anemia by multivariate logistic regression analysis. Among the three factors selected as significant by univariate analysis, we omitted the factor of age from the multivariate analysis as it was included as a parameter in the numerical formula for CL/F. In the multivariate logistic regression analysis of the CL/F category (CL/F < 10) and the "2 by 2" category, the latter was the only significant factor associated with the discontinuance of ribavirin due to severe anemia (Table 7B). Using the "2 by 2" standard, the rate of discontinuance of ribavirin due to severe anemia was 14% (18/133) in the "2 by 2"-negative (the patients whose Hb decreased by less than 2 g/dl from the start of treatment) group and 38% (23/60) in the "2 by 2"-positive group, with the difference being significant ($P < 0.0001$) (Table 8).

We next compared the sensitivity, specificity, and accuracy of the CL/F category with those of the "2 by 2" category as predictive factors for discontinuance of

Table 6. Major causes of discontinuance of ribavirin

	Age < 60	Age ≥ 60
Severe anemia	27 (9%)	41 (21%)*
General fatigue	7	3
Digestive disorders	5	3
Neutropenia	1	1
Thrombocytopenia	2	4
Eruption with itching	2	4
Psychological disorders	3	3
Others	3	5
Total	50/288 (17%)	64/194 (33%)

* $P < 0.001$

ribavirin due to severe anemia among patients aged 60 years or older. Table 9 shows the comparison between the CL/F < 15 category and the "2 by 2" category (Table 9A) and that between the CL/F < 10 category and the "2 by 2" category (Table 9B). Although sensitivity was higher for the lower CL/F category [CL/F < 15, 100% (41/41); CL/F < 10, 71% (29/41)] than for the "2 by 2" category (56%, 23/41), specificity and accuracy were significantly higher for the "2 by 2" category than for the CL/F category [specificity: "2 by 2," 77% (96/125) vs. CL/F < 15, 7% (9/125), $P < 0.0001$; "2 by 2" vs. CL/F < 10, 47% (59/125), $P < 0.0001$; accuracy: "2 by 2," 72% (119/166) vs. CL/F < 15, 30% (50/166), $P < 0.0001$; "2 by 2" vs. CL/F < 10, 53% (88/166), $P < 0.001$].

Table 7. Univariate and multivariate analysis for the discontinuance of ribavirin due to severe anemia among the patients aged 60 years and older

A. Univariate analysis

Factor	Category	Odds ratio	95% CI	P value
Age			1.007–1.250	0.04
Sex	Male/Female	1/1.67	0.280–1.286	0.19
Body weight			0.947–1.021	0.37
Serum creatinine			0.865–33.586	0.07
Ribavirin/Body weight			0.775–1.205	0.76
IFN dosage	6 MU/10 MU	1/1.92	0.803–4.579	0.14
Hb			0.537–1.106	0.16
CL/F	≥15/<15	—	—	0.97
	≥10/<10	1/2.16	0.217–0.989	0.047
“2 by 2”	Negative/Positive	1/4.24	0.112–0.497	0.0001

B. Multivariate analysis

Factor	Category	Odds ratio	95% CI	P value
CL/F	≥10/<10	1/2.12	0.213–1.042	0.063
“2 by 2”	Negative/Positive	1/4.18	0.112–0.507	0.0002

Table 8. Rate of the ribavirin reduction or discontinuance due to adverse effects with the rate of anemia progression among the patients aged 60 years and older

	No reduction	Dose reduction	Discontinuance	
			All cases	Cases due to severe anemia
ΔHb ≥ 2.0 (“2 by 2”-positive) (n = 60)	27% (16/60)	23% (14/60)	50% (30/60)	38%* (23/60)
ΔHb < 2.0 (“2 by 2”-negative) (n = 133)	46% (61/133)	29% (39/133)	25% (33/133)	14% (18/133)

*P < 0.0001

Table 9. Comparison of “2 by 2” standard and CL/F standard for the discontinuance of ribavirin due to severe anemia among the patients aged 60 years and older

A.

	“2 by 2”-positive	CL/F < 15	P value
Sensitivity	56% (23/41)	100% (41/41)	<0.0001
Specificity	77% (96/125)	7% (9/125)	<0.0001
Accuracy	72% (119/166)	30% (50/166)	<0.0001

B.

	“2 by 2”-positive	CL/F < 10	P value
Sensitivity	56% (23/41)	71% (29/41)	0.17
Specificity	77% (96/125)	47% (59/125)	<0.0001
Accuracy	72% (119/166)	53% (88/166)	<0.001

Discussion

Ribavirin, developed in 1972, is a synthetic nucleic acid analog, which has antiviral activity in vitro against a wide variety of RNA and DNA viruses. Combination

therapy of ribavirin with IFN or Peg-IFN led to remarkable progress in antiviral therapy for chronic hepatitis C. To raise the SVR rate for such combination therapy, it is very important to predict the discontinuance of the therapy due to an adverse effect and prevent it. In this study, we observed the incidence of hemolytic anemia, the major side effect of ribavirin. The factors correlated with the progression of anemia were analyzed to avert the need to discontinue ribavirin treatment of patients with chronic hepatitis C receiving combination therapy.

Several studies in the United States and European countries have reported that higher ribavirin dosage or a higher plasma concentration of ribavirin increases the SVR rate.^{21,22} However, a higher ribavirin dose or higher plasma concentration of ribavirin entails the risk of having to discontinue ribavirin treatment. In Japan, analysis of the relationship between the SVR rate and a dose reduction or discontinuance of ribavirin, has shown that reducing the dose of ribavirin does not affect the SVR rate. In the present study, the SVR rate of the patients discontinuing ribavirin was also shown to be significantly lower than the patients who did not discontinue it

in both the 1H group and the non-1H group ($P < 0.01$ and $P < 0.01$, respectively). The SVR rate was almost the same between patients without a dose reduction of ribavirin and those with a dose reduction in both groups (1H, 24% vs. 26%; non-1H, 83% vs. 83%). Therefore, averting ribavirin discontinuance, even if its dose must be reduced, can lead to improvement of the SVR rate. This means that it is important to identify patients prone to develop severe anemia leading to ribavirin discontinuance while they are still in the early phase of treatment, and to consider ribavirin dose reduction before anemia progression.

CL/F relating to the plasma concentration of ribavirin at the end of 4 weeks after initiation of the combination therapy has been used as a predictive factor for the progression of anemia.¹⁶⁻¹⁸ In this study, the patients with a lower CL/F value, which is thought to be correlated with a high plasma concentration of ribavirin, showed a higher rate of discontinuance of ribavirin due to severe anemia than those with a higher CL/F value. This indicates that prediction of anemia progression using the CL/F is useful before the initiation of combination therapy. We analyzed predictive factors for discontinuance of ribavirin due to severe anemia using two CL/F categories, $CL/F < 10$ and $CL/F < 15$, taking into account that the mean CL/F was 13.01/h and the median was 11.91/h, and compared the usefulness of those categories with that of the "2 by 2" standard.

We focused on the early decline of the Hb concentration after the initiation of combination therapy. Monitoring of the Hb decline allowed clear assignment of the patients into three groups: patients without dose reduction of ribavirin, those with dose reduction, and those who discontinued ribavirin. At the end of 2 weeks, a significant relationship was already observed among the three groups. Therefore, we examined the relationship between the beginning of a progression to severe anemia and the decrease in the Hb concentration at the end of 2 weeks (ΔHb). Since a standard value of ΔHb for dose reduction of ribavirin must be established, we compared $\Delta Hb 2.0$ with $\Delta Hb 1.5$, and found that the specificity and accuracy of $\Delta Hb 2.0$ as a predictive factor for the discontinuance of ribavirin due to severe anemia was higher than those of $\Delta Hb 1.5$. We therefore adopted $\Delta Hb 2.0$ at the end of 2 weeks from the start of treatment (the "2 by 2" standard) as the predictive factor for discontinuance of ribavirin due to severe anemia, because an early reduction of ribavirin should be limited to those patients with a higher specificity rate for the progression of anemia. Furthermore, $\Delta Hb 2.0$ is easier to calculate.

In the multivariate logistic regression analysis, both the CL/F category and the "2 by 2" category were useful for all patients as independent predictive factors for discontinuing ribavirin due to severe anemia (Table 5).

Patients with lower CL/F ($CL/F < 10$ or $CL/F < 15$) and those who were "2 by 2" positive were significantly associated with the discontinuance of ribavirin due to severe anemia. Thus, the CL/F standard should be used as a predictive factor before combination therapy is begun, and the "2 by 2" standard should be used during the combination therapy. We also assessed which would be the more useful predictive factor for discontinuance of ribavirin due to severe anemia among older patients. Multivariate analysis showed that only the "2 by 2" standard was significantly related to the discontinuance of ribavirin due to severe anemia among older patients (Table 7B). Moreover, the "2 by 2" standard showed higher specificity (77%) and accuracy (72%) for the discontinuance of ribavirin due to severe anemia among older patients than either CL/F value (Table 9). The ribavirin dose of 200 mg should be reduced for aged patients whose Hb decreases over 2 g/dl from the start of combination therapy in order to avoid having to discontinue ribavirin administration altogether.

Hemolytic anemia has been reported to be induced by ribavirin administration, depending on the plasma ribavirin concentration¹⁵ and the fragile membrane of RBC in which ribavirin accumulates.²³ Furthermore, the plasma clearance of ribavirin has been reported to depend on renal function.^{24,25} The anemia associated with IFN and ribavirin therapy is a "mixed anemia," in which both hemolysis and bone marrow suppression occur simultaneously. In this study, many patients, especially older ones, had to discontinue ribavirin due to severe anemia, as previously reported.²⁶ A major reason for this was thought to be the tendency of the plasma concentration of ribavirin to rise due to lower renal function and impaired hematogenous function as the anemia progressed. In predicting the discontinuance of ribavirin due to severe anemia using the CL/F category, the lower CL/F implies that older patients and patients with low renal function are high-risk groups. However, CL/F does not account for the fragile membrane of RBC or the hematogenous function. Therefore, the CL/F standard cannot be a good marker for individual patients, because CL/F does not reflect in vivo phenomena triggered by ribavirin. CL/F is related simply to the plasma concentration of ribavirin at the end of 4 weeks after the initiation of combination therapy. On the other hand, the "2 by 2" standard can be useful as a predictive factor of ribavirin discontinuance forces by severe anemia for all patients, including older patients. It indicates that the "2 by 2" standard reflects plural factors, such as the occurrence of hemolysis and hematogenous functions. We suggest that the "2 by 2" standard is more useful than the CL/F category as a predictive factor for discontinuance of ribavirin due to severe anemia, especially among older patients.

In conclusion, it is important to monitor the early decline of the Hb concentration after initiation of combination therapy and to reduce the dose of ribavirin at the end of 2 weeks based on the magnitude of the Hb decline. An early reduction of ribavirin before progression to severe anemia can reduce the number of patients who are destined to discontinue ribavirin therapy. This should help improve the patients' quality of life by preventing the progression to severe anemia. Further prospective study is necessary to evaluate the antiviral outcome by ITT analysis using early reduction of ribavirin based on the "2 by 2" standard.

Acknowledgments. Other institutions and participants in the Osaka Liver Disease Study Group (Digestive Disease Study Group of Osaka Renaissance) were the National Hospital Organization Osaka National Hospital, Y. Izumi; Osaka Rousai Hospital, K. Noda and M. Satoh; Osaka Kouseinenkin Hospital, M. Kurokawa; Kansai Rousai Hospital, M. Yamamoto; Osaka General Medical Center, T. Inoue; National Hospital Organization Osaka Minami Medical Center, Y. Inoue and M. Shigekawa; Osaka Police Hospital, J. Kondo; Kaizuka City Hospital, O. Nishiyama; and Osaka University Graduate School of Medicine, S. Shinzaki, I. Itose, S. Egawa, and T. Nishida.

This work was supported by a Grants-in-Aid for Research on Hepatitis and BSE from Health, Labour and Welfare Ministry of Japan, and for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Lauer GM, Walker BD. Hepatitis C virus infection. *N Engl J Med* 2001;345:41–52.
- Poynard T, Bedossa P, Opolon P. Natural history of liver fibrosis progression in patients with chronic hepatitis C. The OBSVIRC, METAVIR, CLINIVIR, and DOSVIRC groups. *Lancet* 1997; 349:825–32.
- Hamada H, Yatsuhashi H, Yano K, Daikoku M, Arisawa K, Inoue O, et al. Impact of aging on the development of hepatocellular carcinoma in patients with posttransfusion chronic hepatitis C. *Cancer* 2002;95:331–9.
- Hiramatsu N, Hayashi N, Kasahara A, Hagiwara H, Takehara T, Haruna Y, et al. Improvement of liver fibrosis in chronic hepatitis C patients treated with natural interferon alpha. *J Hepatol* 1995; 22:135–42.
- Kasahara A, Hayashi N, Mochizuki K, Takayanagi M, Yoshioka K, Kakumu S, et al. Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. Osaka Liver Disease Study Group. *Hepatology* 1998; 27:1394–402.
- Poynard T, Marcellin P, Lee SS, Niederau C, Minuk GS, Ideo G, et al. Randomised trial of interferon alpha2b plus ribavirin for 48 weeks or for 24 weeks versus interferon alpha2b plus placebo for 48 weeks for treatment of chronic infection with hepatitis C virus. International Hepatitis Interventional Therapy Group (IHIT). *Lancet* 1998;352:1426–32.
- McHutchison JG, Gordon SC, Schiff ER, Shiffman ML, Lee WM, Rustgi VK, et al. Interferon alpha-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. Hepatitis Interventional Therapy Group. *N Engl J Med* 1998;339: 1485–92.
- Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358: 958–65.
- Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL Jr, et al. Peginterferon alpha-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–82.
- Hiramatsu N, Kasahara A, Nakanishi F, Toyama T, Tsujii M, Tsuji S, et al. The significance of interferon and ribavirin combination therapy followed by interferon monotherapy for patients with chronic hepatitis C in Japan. *Hepatol Res* 2004;29:142–7.
- Bodenheimer HC Jr, Lindsay KL, Davis GL, Lewis JH, Thung SN, Seeff LB. Tolerance and efficacy of oral ribavirin treatment of chronic hepatitis C: a multicenter trial. *Hepatology* 1997;26:473–7.
- De Franceschi L, Fattovich G, Turrini F, Ayi K, Brugnara C, Manzato F, et al. Hemolytic anemia induced by ribavirin therapy in patients with chronic hepatitis C virus infection: role of membrane oxidative damage. *Hepatology* 2000;31:997–1004.
- Van Vlierbergh H, Delanghe JR, De Vos M, Leroux-Roel G. Factors influencing ribavirin-induced hemolysis. *J Hepatol* 2001;34:911–6.
- Tappero G, Ballare M, Farina M, Negro F. Severe anemia following combined alpha-interferon/ribavirin therapy of chronic hepatitis C. *J Hepatol* 1998;29:1033–4.
- Lindahl K, Schvarcz R, Bruchfeld A, Stahle L. Evidence that plasma concentration rather than dose per kilogram body weight predicts ribavirin-induced anaemia. *J Viral Hepat* 2004;11: 84–7.
- Jen JF, Glue P, Gupta S, Zambas D, Hajian G. Population pharmacokinetic and pharmacodynamic analysis of ribavirin in patients with chronic hepatitis C. *Ther Drug Monit* 2000;22: 555–65.
- Kamar N, Chatelut E, Manolis E, Lafont T, Izopet J, Rostaing L. Ribavirin pharmacokinetics in renal and liver transplant patients: evidence that it depends on renal function. *Am J Kidney Dis* 2004;43:140–6.
- Karino Y, Kato T, Arakawa T, Matsumoto S, Kuwata Y, Araike J, et al. Total clearance (CL/F) of ribavirin is the factor most influencing the incidence of hemolytic anemia during IFN plus ribavirin therapy. *Hepatology* 2004;40(Suppl 1): 358.
- Rebetron® Combination Therapy containing Rebetol® (ribavirin, USP) Capsules and Intron® A (interferon alpha-2b, recombinant) Injection prescribing information. Schering Corporation, Kenilworth, NJ. January 2001.
- Knodell RG, Ishak KG, Black WC, Chen TS, Craig R, Kaplowitz N, et al. Formulation and application of a numerical scoring system for assessing histological activity in asymptomatic chronic active hepatitis. *Hepatology* 1981;1:431–5.
- Hadziyannis SJ, Sette H Jr, Morgan TR, Balan V, Diago M, Marcellin P, et al. Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140: 346–55.
- Lindahl K, Stahle L, Bruchfeld A, Schvarcz R. High-dose ribavirin in combination with standard dose peginterferon for treatment of patients with chronic hepatitis C. *Hepatology* 2005; 41:275–9.
- Grattagliano I, Russmann S, Palmieri VO, Juni P, Bihl F, Portincasa P, et al. Low membrane protein sulfhydryls but not G6PD deficiency predict ribavirin-induced hemolysis in hepatitis C. *Hepatology* 2004;39:1248–55.
- Bruchfeld A, Lindahl K, Schvarcz R, Stahle L. Dosage of ribavirin in patients with hepatitis C should be based on renal function: a population pharmacokinetic analysis. *Ther Drug Monit* 2002;24: 701–8.

25. Maeda Y, Kiribayashi Y, Moriya T, Maruhashi A, Omoda K, Funakoshi S, et al. Dosage adjustment of ribavirin based on renal function in Japanese patients with chronic hepatitis C. *Ther Drug Monit* 2004;26:9-15.
26. Hiramatsu N, Oze T, Tsuda N, Kurashige N, Koga K, Toyama T, et al. Should aged patients with chronic hepatitis C be treated with interferon and ribavirin combination therapy? *Hepatol Res* 2006;35:185-9.

Case report

Squamous cell carcinoma as a rare entity of primary liver tumor with grave prognosis

Nobukazu Yuki^{a,*}, Yasuki Hijikata^a, Michio Kato^a,
Kunimitsu Kawahara^b, Kenichi Wakasa^c

^a Department of Gastroenterology, Osaka National Hospital, Hoenzaka 2-1-14, Chuo-ku, Osaka 540-0006, Japan

^b Department of Pathology, Osaka Prefectural Medical Center for Respiratory and Allergic Diseases, Habikino 583-8588, Japan

^c Department of Pathology, Osaka City University Hospital, Osaka 545-8586, Japan

Received 15 February 2006; received in revised form 2 August 2006; accepted 3 August 2006
Available online 15 September 2006

Abstract

The development of primary squamous cell carcinoma (SCC) of the liver has only rarely been reported in association with pre-existing hepatic cysts and biliary tract diseases. This report describes an unusual case of SCC originating in a cirrhotic liver. A 63-year-old male alcoholic was incidentally found to have a 6-cm liver tumor which showed mixed echogenic by sonography and a low-density area with rim enhancement by computed tomography. Tumor biopsy led to a diagnosis of SCC with a sarcomatoid change. The tumor showed fatal rapid growth accompanied by abdominal pain. Transcatheter arterial embolization and chemotherapy were not effective, and the tumor increased to 13 cm in diameter over an 8-month period. A post-mortem search revealed no alternative primary tumor site other than liver. Review of the literature shows that abdominal pain is a chief symptom of primary liver SCC. Only three of the nine patients treated with hepatic resection survived without recurrence during 8 months to 4 years of follow-up. We propose that SCC with a grave prognosis should be considered as a rare entity of primary liver tumor even in cirrhotic patients.

© 2006 Elsevier Ireland Ltd. All rights reserved.

Keywords: Liver cirrhosis; Liver tumor; Squamous cell carcinoma

1. Introduction

Squamous cell carcinoma (SCC) rarely arises from the liver. However, when it does occur, the prognosis can be worse than that of hepatocellular carcinoma, the most common primary liver tumor. Here, we describe an unusual case of SCC primarily arising from a cirrhotic liver. In this paper, we combine our experience with data obtained from the literature, and try to summarize this rare entity of primary liver tumor based on its clinical presentation, diagnostic findings and results of subsequent therapy.

2. Case report

2.1. Clinical features

The entire 8-month clinical course is shown in Fig. 1. A 63-year-old Japanese male with alcohol abuse (>80 g/day over 40 years) underwent a health checkup at a local hospital. He was asymptomatic, but abdominal sonography disclosed a cirrhotic liver with a 6-cm mixed echogenic tumor in segment 6. He had no confounding etiology of liver disease other than alcohol. He was negative for hepatitis B surface antigen and hepatitis C virus antibody, had no history of administration of hepatotoxic drugs and showed no evidence of autoimmune liver disease. Computed tomography confirmed the tumor as a low-density area showing rim enhancement after contrast injection. A needle biopsy demonstrated the possibility

* Corresponding author. Tel.: +81 6 6942 1331; fax: +81 6 6943 6467.
E-mail address: yuki@onh.go.jp (N. Yuki).




Fig. 1. Clinical course of a case with primary squamous cell carcinoma of the liver. The tumor showed no response to transcatheter arterial embolization (TAE) and chemotherapy resulting in fatal rapid growth.

of the tumor being SCC. A thorough search for a primary tumor was undertaken including chest radiography, computed tomography of the lungs, upper gastrointestinal endoscopy, colonoscopy, urological examination, magnetic resonance imaging of the brain and ear, nose and throat screening. No other tumor source was found. Hepatic angiography showed a 6-cm hypovascular mass in the lower part of the right liver lobe (Fig. 2). Epirubicin hydrochloride (EPI) 50 mg, fluorouracil (5-FU) 250 mg and mitomycin C (MMC) 4 mg were infused followed by transcatheter arterial embolization (TAE). Thereafter, the tumor was further treated by continuous transarterial infusion of cisplatin (CDDP) 100 mg over 90 min.

Five months after the initial diagnosis, the patient was referred to Osaka National Hospital, complaining of abdominal distension and vague right upper quadrant pain. He had an intermittent fever of up to 38.4 °C. Physical examination was significant for marked ascites and right upper quadrant tenderness. Abdominal sonography showed an 8-cm mixed echogenic liver tumor in segment 6 (Fig. 3). By computed tomography, the tumor appeared again as a low-density area showing rim enhancement (Fig. 4). Aspi-



Fig. 2. Hepatic angiography disclosed a 6-cm hypovascular mass in the lower part of the right liver lobe (arrows).




Fig. 3. On admission to our institution, abdominal sonography showed marked ascites and a cirrhotic liver with an 8-cm mixed echogenic tumor in segment 6.




Fig. 4. By computed tomography, the liver tumor appeared as a low-density area showing rim enhancement after contrast injection.

rated ascites was clear in appearance and appeared transudate in nature. Cytology revealed no tumor cells, and bacteriology was negative. Laboratory tests showed anemia and hypoproteinemia, with a hemoglobin level of 9.8 g/dL; total proteins, 5.6 g/dL; albumin, 2.5 g/dL; serum bilirubin, 2.3 mg/dL; aspartate aminotransferase, 32 IU/L; alanine aminotransferase, 13 IU/L; alkaline phosphatase, 577 IU/L (normal 109–344); γ -glutamyl transpeptidase, 137 IU/L. As for tumor markers, alpha-fetoprotein, PIVKA-II, carcinoembryonic antigen and CA19-9 were normal. Elevation was observed for SCC-related antigen (3.2 ng/mL [<1.5]) and cytokeratin fragment 21.1 (CYFRA21.1) (8.4 ng/mL [<2.8]). After admission, the liver tumor grew to 13 cm in diameter, and a 3-cm new tumor appeared in segment 3. In parallel with rapid growth of the tumor, the ascites increased further, and marked jaundice appeared. The patient was managed conservatively for right upper quadrant pain. His condition progressively deteriorated, and he died 8 months after the initial diagnosis.

2.2. Pathologic features

The histology of the tumor biopsy showed sarcomatoid spindle cell component and multiple foci of squamous cells with nuclear atypia and large eosinophilic keratinizing cytoplasm, which were compatible with SCC lesions (Fig. 5). Areas of transition from SCC to the sarcomatoid tissue were visible. SCC cells and spindle-shaped cells were both positive for monoclonal antibody MNF116 directed to human cytokeratins 5, 6, 8 and 17 and specific for epithelial lesions (Dako Cytomation Co., Ltd., Kyoto, Japan). Neither SCC cells nor spindle cells showed reactivity to anti-vimentin monoclonal antibody Vim3B4 specific for mesenchymal cells and anti-human hepatocyte monoclonal antibody OCH1E5 (Dako Cytomation Co., Ltd.). Thus, the diagnosis reached was liver SCC with a sarcomatoid change.

At autopsy, the liver appeared cirrhotic with a finely nodular surface. Cut surfaces of the liver revealed a large whitish mass arising primarily from the right liver lobe extending into the left lobe and the right kidney, the largest of which measured 14 cm in diameter (Fig. 6). Tumor thrombi were observed in the portal veins adjacent to the main tumor. No cystic changes were present in the liver, and the hepatic and bile ducts were normal. Lymph node metastasis was seen at the hepatic hilus, and tumor deposits due to direct spread were present in the intestinal serosa. In an attempt to demonstrate a possible alternative primary site of the tumor other than liver, we undertook a careful examination of the whole body, but no tumor was found elsewhere. Microscopically, the main liver tumor showed appearances similar to those seen in the previous biopsy sections. The vast majority (90%) had sarcomatoid features with keratinizing SCC lesions embedded in it. Thorough examination identified, albeit extremely infrequently, an SCC island which contained a few carcinoma cells with Alcian blue-positive mucin-like substance (Fig. 7). However, glandular formation was not evident. The




Fig. 5. Histology of the tumor biopsy revealed foci of squamous cell carcinoma showing keratinization embedded in the sarcomatoid change (hematoxylin-eosin, original magnification $\times 20$ (A) and $\times 100$ (B)).

lesion was very limited in that this type of carcinoma cells was not found in adjacent sections examined by Alcian blue, mucicarmine and PAS stain. The tumor cells were all negative for cytokeratins 7 and 19, known as markers of the bile duct epithelium. The same histological features with the primary




Fig. 6. A cut surface of the autopsied liver. A large whitish tumor arising primarily from the right liver lobe expanded into the left lobe and the right kidney.

Table 1
Summary of reported cases of primary squamous cell carcinoma of the liver

Case no.	Authors	Age (year)/sex	Underlying liver disease	Symptoms	Methods of diagnosis	Treatment	Outcome
1	Bloustein and Silverberg [1]	30/M	SBNHC	Abdominal pain	Exploratory laparotomy	Hepatic resection + chemotherapy and radiotherapy	Died 4 months post-operatively
2	Song et al. [2]	43/M	Hepatolithiasis	Abdominal pain, Jaundice	Exploratory laparotomy	Hepatic resection	Died 6 months post-operatively
3	Gresham and Rue [3]	78/M	SBNHC	Abdominal pain	Post-mortem examination	Symptomatic treatment	Died
4	Lynch et al. [4]	63/M	SBNHC	Abdominal pain, Jaundice	Exploratory laparotomy	Cyst drainage	Died 6 months after admission
5	Arase et al. [5]	63/M	Liver cirrhosis	Abdominal pain	Post-mortem examination	Transarterial infusion of mitomycin C	Died
6	Clements et al. [6]	73/M	Biliary strictures	Jaundice	Post-mortem examination	Symptomatic treatment	Died
7	Roediger and Dymock [7]	51/F	No	Abdominal pain	Exploratory laparotomy	Hepatic resection	Died post-operatively
8	Nieweg et al. [8]	62/F	SBNHC	Abdominal pain	Exploratory laparotomy	Symptomatic treatment	Died 5 months post-operatively
9	Pliskin et al. [9]	82/F	Multiple congenital cysts	No particular symptom	Percutaneous biopsy	Symptomatic treatment	Died
10	Banbury et al. [10]	59/F	SBNHC	Abdominal pain	Exploratory laparotomy	Hepatic resection	No recurrence (over 16mo)
11	Lombardo et al. [11]	59/F	Epidermoid cyst	Abdominal pain	Exploratory laparotomy	Hepatic resection	No recurrence (over 8mo)
12	Weimann et al. [12]	74/F	SBNHC	Abdominal pain	Exploratory laparotomy	Hepatic resection	No recurrence (over 4 years)
13	Shinagawa et al. [13]	56/F	Cystically dilated biliary tract	Jaundice	Bile cytology	Hepatic resection	Died 8 months post-operatively
14	Monteagudo et al. [14]	71/F	SBNHC	Abdominal pain	Percutaneous biopsy	Cystostomy	Died 1 month post-operatively
15	Doctor et al. [15]	68/F	No	Fever	Percutaneous biopsy	Hepatic resection + chemotherapy with cisplatin and 5-fluorouracil	Unknown
16	Vick et al. [16]	51/M	Ciliated foregut cyst	Abdominal pain	Exploratory laparotomy	Hepatic resection	Died 2 month post-operatively

SBNHC, solitary benign nonparasitic hepatic cyst.




Fig. 7. Thorough post-mortem histological examination revealed that an island of squamous cell carcinoma, albeit very rarely, contained a few carcinoma cells with mucin-like substance (Alcian blue stain, original magnification $\times 80$).




Fig. 8. Histology of the background liver showed the features of cirrhosis (hematoxylin-eosin, original magnification $\times 10$).

tumor were seen for the new liver tumor in segment 3, lymph node metastasis and tumor deposits in the intestinal serosa. Foci of SCC were embedded in the sarcomatoid change. The background liver showed the histological features of cirrhosis. Regenerative nodules of various sizes were surrounded by fibrous septa. Necroinflammation and fatty change were not evident (Fig. 8).

3. Discussion

The low prevalence of primary SCC of the liver is reflected by the small number of reported cases. A review of literature disclosed 16 cases [1–16] (Table 1) of 30–82-year-old subjects (median 63 years), seven males and nine females. The tumor had arisen from congenital hepatic cysts in 10 (63%) cases and from diseased biliary tracts in three (19%) cases. Solitary and multiple benign nonparasitic hepatic cyst(s), the most common congenital hepatic cysts originate from von Meyenburg's complexes (congenital rests of biliary epithelium)

and are lined with simple cuboidal or columnar (bile duct-type) epithelium. SCC is generally considered to result from secondary squamous metaplasia of such biliary epithelium due to chronic inflammation and subsequent neoplastic transformation. The primary liver SCC described in this report was unique because it arose from a cirrhotic liver of alcoholic origin and was not associated with pre-existing cystic and biliary tract diseases. In the literature, there are only three such cases. One case was of liver cirrhosis of uncertain etiology [5], while the other two had no documented liver disease [7,15]. The carcinogenesis process in this situation remains unclear. In our patient, thorough post-mortem examination of the tumor demonstrated an SCC focus containing a few carcinoma cells with Alcian blue-positive mucin-like substance. This observation, albeit extremely infrequent, raises a possibility that the SCC may have resulted from squamous metaplasia of an adenocarcinoma presumably originating from the biliary epithelium.

The liver tumor showed a sarcomatoid change and fatal rapid growth accompanied by right upper quadrant abdominal pain. In previous reports, such histological change has not been documented. However, the tumor size at presentation was large ranging between 5 and 23 cm (median 9 cm), and abdominal pain was the most common clinical manifestation. Thus, primary liver SCC seems to grow rapidly even in the absence of a sarcomatoid change. In our case, the liver tumor appeared as a mixed echogenic tumor and a low-density area showing rim enhancement by computed tomography, which agrees with the limited data available in the literature. Although changes in SCC-specific tumor markers have not been well documented, SCC-related antigen and CYFRA21.1 were elevated in this case. Early diagnosis is often difficult for primary liver SCC, but these observations may help toward a differential diagnosis. Neither chemotherapy nor radiation therapy have been shown to be effective. This was also the case with our patient. Hepatic resection seems the only promising therapy. According to the literature, five patients died within 8 months post-operatively, but three patients survived without evidence of recurrence during the 8 months to 4 years of follow-up.

We have described an extremely rare case with SCC primarily arising from a cirrhotic liver. Lessons from previous reports and the present case suggest that the prognosis of primary liver SCC is grave. This tumor displays the following characteristics: (1) frequent but not consistent association with cystic and biliary tract diseases, (2) frequent abdominal pain, (3) mixed echogenic and hypovascular nature and (4) possible elevation of SCC-specific tumor markers. Under such conditions, the occurrence of SCC should be kept in mind as a rare entity of primary liver tumor.

References

- [1] Bloustein PA, Silverberg SG. Squamous cell carcinoma originating in a hepatic cyst. *Cancer* 1976;38:2002–5.

- [2] Song E, Kew MC, Grieve T, Isaacson C, Myburgh JA. Primary squamous cell carcinoma of the liver occurring in association with hepatolithiasis. *Cancer* 1984;53:542–6.
- [3] Gresham GA, Rue LW. Squamous cell carcinoma of the liver. *Hum Pathol* 1985;16:413–6.
- [4] Lynch MJ, McLeod MK, Weatherbee L, Gilsdorf JR, Guice KS, Eckhauser FE. Squamous cell cancer of the liver arising from a solitary benign nonparasitic hepatic cyst. *Am J Gastroenterol* 1988;83:426–31.
- [5] Arase Y, Endo Y, Hara M, Kumada H, Ikeda K, Yoshida A. Hepatic squamous cell carcinoma with hypercalcemia in liver cirrhosis. *Acta Pathol Jpn* 1988;38:643–50.
- [6] Clements D, Newman P, Etherington R, Lawrie BW, Rhodes J. Squamous carcinoma in the liver. *Gut* 1990;31:1333–4.
- [7] Roediger WEW, Dymock RB. Primary squamous carcinoma of the liver: clinical and histopathological features. *Aust NZ J Surg* 1991;61:720–2.
- [8] Nieweg O, Slooff MJH, Grond J. A case of primary squamous cell carcinoma of the liver arising in a solitary cyst. *HPB Surg* 1992;5:203–8.
- [9] Pliskin A, Cualing H, Stenger RJ. Primary squamous cell carcinoma originating in congenital cysts of the liver. *Arch Pathol Lab Med* 1992;116:105–7.
- [10] Banbury J, Conlon KC, Ghossein R, Brennan MF. Primary squamous cell carcinoma within a solitary nonparasitic hepatic cyst. *J Surg Oncol* 1994;57:210–2.
- [11] Lombardo FP, Hertford DE, Tan LK, Kazam E, Ramirez de Arellano E. Epidermoid cyst of the liver complicated by microscopic squamous cell carcinoma: CT, ultrasound, and pathology. *J Comput Assist Tomogr* 1995;19:131–4.
- [12] Weimann A, Klempnauer J, Gebel M, et al. Squamous cell carcinoma of the liver originating from a solitary non-parasitic cyst: case report and review of the literature. *HPB Surg* 1996;10:45–9.
- [13] Shinagawa T, Tadokoro M, Takagi M, et al. Primary squamous cell carcinoma of the liver: a case report. *Acta Cytol* 1996;40:339–45.
- [14] Monteagudo M, Vidal G, Moreno M, et al. Squamous cell carcinoma and infection in a solitary hepatic cyst. *Eur J Gastroenterol Hepatol* 1998;10:1051–3.
- [15] Doctor N, Dafnios N, Jones A, Davidson BR. Primary squamous carcinoma of liver: presentation as liver abscess. *Indian J Gastroenterol* 1998;17:28–9.
- [16] Vick DJ, Goodman ZD, Ishak KG. Squamous cell carcinoma arising in a ciliated hepatic foregut cyst. *Arch Pathol Lab Med* 1999;123:1115–7.

Serum HBV-RNA is a Predictor of Early Emergence of YMDD Mutant in Patients Treated with Lamivudine

Tsuyoshi Hatakeyama^{1,2}, Chiemi Noguchi^{1,2}, Nobuhiko Hiraga^{1,2}, Nami Mori^{1,2},
5 Masataka Tsuge^{1,2}, Michio Imamura^{1,2}, Shoichi Takahashi^{1,2}, Yoshiiku Kawakami^{1,2},
Yoshifumi Fujimoto^{2,3}, Hidenori Ochi^{2,3}, Hiromi Abe^{1,2,3}, Toshiro Maekawa³, Hiroiku
Kawakami⁴, Hiromi Yatsuji^{1,2}, Yasuyuki Aisaka⁵, Hiroshi Kohno⁵, Shiomi Aimitsu⁵,
Kazuaki Chayama^{1,2,3}*

10 ¹Department of Medicine and Molecular science, Division of Frontier Medical
Science, Programs for Biomedical Research, Graduate School of Biomedical Sciences,
Hiroshima University, 1-2-3, Kasumi, Minami-ku, Hiroshima-shi, 734-8551, JAPAN

²Liver Research Project Center, Hiroshima University, Hiroshima, Japan

³Laboratory for Liver Disease, SNP Research Center, The Institute of Physical and
15 Chemical Research (RIKEN), Yokohama 230-0045, Japan

⁴Kawakami Clinic, 4-24, Hacchoubori, Naka-ku, Hiroshima-shi, 730-0013, Japan

⁵Department of Hepatology, Hiroshima Red Cross Hospital and Atomic Bomb
Survivors Hospital, 1-9-6, Senda-machi, Naka-ku, Hiroshima-shi, 730-8619, Japan

20 *Short title: HBV-RNA and LAM resistance*

Key words: ETV, TMA-HPA, nucleotide substitution, RT, resistance

Corresponding author:

Prof. Kazuaki Chayama, MD

25 Department of Medical and Molecular Science,
Division of Frontier Medical Science, Programs for Biomedical Research,
Graduate school of Biomedical Science, Hiroshima University,
1-2-3 Kasumi, Minami-ku, Hiroshima 734-8551, Japan
Tel: +81-82-257-5190
30 Fax: +81-82-255-6220
E-mail: chayama@hiroshima-u.ac.jp

Abbreviations:

ALT: alanine transaminase
35 cccDNA: covalently closed circular DNA
ETV: entecavir
HBV: hepatitis B virus
HIV: human immunodeficiency virus
LAM: lamivudine
40 PCR: polymerase chain reaction
RT: reverse transcription
TMA-HPA: transcription mediated amplification and hybridization protection assay
YMDD: tyrosine-methionine-aspartate-aspartate