

Fig. 6. RT-PCR analysis of IFN- β , interferon regulatory factors (IRF1, IRF3, and IRF7), type I IFN receptors (IFNAR1 and IFNAR2c), transcriptional factors (STAT1 and STAT2), and GAPDH in PH5CH8 cells infected with pCXbsr retrovirus encoding Core, NS5A, or NS5B. The pCXbsr retrovirus was used as a control infection (Ctr). At 48 h postinfection, total RNA was extracted and subjected to RT-PCR analysis using the primer set for IRF3 (312 bp), IFN- β (202 bp), IRF7 (346 bp), IRF1 (246 bp), IFNAR1 (256 bp), IFNAR2c (385 bp), STAT1 (300 bp), STAT2 (379 bp), or GAPDH (334 bp). The PCR products were detected by staining with ethidium bromide after 3% agarose gel electrophoresis. Lane (-), PH5CH8 cells without retrovirus infection.

suppressed by NS3-4A, and the synergistic effect of Core and NS5B was not invalidated by NS3-4A.

Since Core and NS5B equivalently activate the 2'-5'-OAS and IFN- β gene promoters and a synthetic ISRE promoter, it is of interest that IFN- β was induced by NS5B, but not by Core. Furthermore, the synergistic effect of Core on the activation of these gene promoters by NS5B is also interesting. Recently, we found that NS5B, but not Core, delayed the cell cycle progression through the S phase in PH5CH8 cells, and we considered that this phenomenon may have been caused by IFN- β induced by NS5B through the activation of TLR3 (Naka et al., submitted). In the present study, we observed that Core enhanced NS5B's induction of IFN- β mRNA and that Core might indirectly interact with NS5B. Therefore, our findings suggest not only that Core's activation of gene promoters through ISRE is synergistically enhanced by NS5B, but also that Core positively modifies the activation of the TLR3 signaling pathway by NS5B, although Core does not activate this pathway directly.

In the course of studies on the mechanism of Core's activation of gene promoters containing ISRE, we previously showed that Core had no effects on the gene expression levels or phosphorylation status of the major components involved in the JAK-STAT signaling pathway, such as STAT1 and STAT2 [16]. Recently, Imanaka et al. [35] reported that IFN- α -induced STAT-1 phosphorylation and the expression of antiviral genes were inhibited in the suppressor of cytokine signaling (SOCS)-1-expressing cells. Since SOCS-1 is referred to as the STAT-induced STAT

inhibitor or the JAK-binding protein, we speculated that Core may be able to suppress the SOCS-1 expression in PH5CH8 cells, resulting in activation of the JAK-STAT signaling pathway. However, RT-PCR analysis revealed that the level of SOCS-1 mRNA was not changed regardless of the expression of Core in PH5CH8 cells. The other possibility is that Core may modify the affinity of IFN-stimulated gene factor 3 (ISGF3) to the ISRE sequence, since it has recently been reported that Core expression is associated with increased ISGF3-binding activity to the ISRE sequence [36]. Further analysis will be necessary to clarify whether or not the same phenomenon is observed in PH5CH8 cells.

Our observations that NS3-4A drastically suppresses NS5B's activation of the IFN system, especially induction of IFN- β , are consistent with the recent findings [18] that NS3-4A effectively blocks the phosphorylation of IRF3 (a key molecule in innate immunity) that normally occurs in response to virus infection. Recent studies [19,37,38] have indicated that NS3-4A inhibits both the TLR3 signaling pathway (TRIF-dependent pathway) and RIG-I signaling pathway (TRIF-independent pathway). The present study showed that NS3-4A in a serine protease activity-dependent manner inhibited the TLR3 signaling pathway in PH5CH8 cells, based on our recent finding that the induction of IFN- β by NS5B was mediated through the TLR3 but not the RIG-I signaling pathway (Naka et al., submitted). It is likely that TRIF, which was identified recently [19] as a target molecule of cellular components upstream of IRF3, is cleaved by NS3-4A in PH5CH8 cells. However, the induction of IFN- β in PH5CH8 cells co-expressing Core and NS5B was only partially suppressed by NS3-4A, whereas the induction of IFN- β by NS5B only was completely suppressed by NS3-4A. As one of the biological implications of this phenomenon, we speculate that HCV proteins contribute to the maintenance of a low steady state of virus by controlling the expression level of IFN- β in the infected cells, thereby enabling HCV to escape from the host immuno-surveillance system, and facilitating persistent viral infection.

To evaluate this hypothesis, it is important to clarify whether or not the activation of the IFN system by Core and/or NS5B or suppression of the IFN system by NS3-4A occur during the HCV life cycle. Although four kinds of genome-length HCV RNA-replicating cells [39–42] and a reproducible HCV proliferation system in cell culture [43] have been established to date using HuH-7 cells, these HuH-7-based cells would not be suitable to prove our hypothesis, because the TLR3 and/or RIG-I signaling pathway does not function in these cells [38,44]. Therefore, a new HCV RNA-replicating or HCV proliferation cell system needs to be developed using other hepatocyte cell lines possessing intact TLR3 and RIG-I signaling pathways, such as PH5CH8 cells [20]. We are currently conducting a trial to establish genome-length HCV RNA-replicating PH5CH8 cells.




Fig. 7. NS3-4A did not suppress the synergistic effect of Core and NS5B on IFN-β gene expression. (A) Effect of NS3-4A on the activation of the *IFN-β* gene promoter by Core, by NS5B, and by the synergistic effect of Core and NS5B. The dual luciferase reporter assay was performed as described in Fig. 1A. The expression vector of the NS3-4A/S1165A mutant lacking serine protease activity was also used. (B) Real-time LightCycler PCR analysis of IFN-β in PH5CH8 cells stably expressing HCV protein(s). PH5CH8 cells were infected with retrovirus pCXbsr encoding various HCV proteins, and PH5CH8 cells stably expressing HCV protein(s) were obtained as described in Materials and methods. The pCXbsr retrovirus was used as a control infection (Ctr). Total RNA was extracted from the cells and was subjected to real-time LightCycler PCR analysis using the primer set for IFN-β (202 bp). The experiments were performed in at least triplicate. To correct the differences in RNA quality and quantity between the samples, data were normalized using the ratio of IFN-β mRNA concentration to that of GAPDH. The relative level of IFN-β mRNA calculated, when the level of IFN-β mRNA of PH5CH8 cells expressing NS5B alone was assigned to be 1.0, is presented here. (C) Western blot analysis of HCV proteins. Production of Core, NS3, and NS5B in PH5CH8 cells indicated in (B) was analyzed by immunoblotting using anti-Core, anti-NS3, and anti-NS5B antibodies, respectively. The PH5CH8 cells transfected with pCXbsr plasmid were used as control (Ctr). β-Actin was used as a control for the amount of protein loaded per lane.

Acknowledgments

We thank T. Nakamura for his helpful assistance with the experiments. This work was supported by Grants-in-

Aid for the third-term comprehensive 10-year strategy for cancer control, and for research on hepatitis from the Ministry of Health, Labor, and Welfare of Japan, and by the Program for Promotion of Fundamental Studies in Health

Sciences of the National Institute of Biomedical Innovation (NIBIO).

References

- [1] Q.L. Choo, G. Kuo, A.J. Weiner, L.R. Overby, D.W. Bradley, M. Houghton, Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome, *Science* 244 (1998) 359–362.
- [2] G. Kuo, Q.L. Choo, H.J. Alter, G.L. Gitnick, A.G. Redeker, R.H. Purcell, T. Miyamura, J.L. Dienstag, M.J. Alter, C.E. Stevens, G.E. Tegtmeier, F. Bonino, W.S. Colombo, W.S. Lee, C. Kuo, K. Berger, J.R. Shuster, L.R. Overby, D.W. Bradley, M. Houghton, An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis, *Science* 244 (1998) 362–364.
- [3] S. Ohkoshi, H. Kojima, H. Tawaraya, T. Miyajima, T. Kamimura, H. Asakura, A. Satoh, S. Hirose, M. Hijikata, N. Kato, K. Shimotohno, Prevalence of antibody against non-A, non-B hepatitis virus in Japanese patients with hepatocellular carcinoma, *Jpn. J. Cancer Res.* 81 (1990) 550–553.
- [4] I. Saito, T. Miyamura, A. Ohbayashi, H. Harada, T. Katayama, S. Kikuchi, Y. Watanabe, S. Koi, M. Onji, Y. Ohta, Q.L. Choo, M. Houghton, G. Kuo, Hepatitis C virus infection is associated with the development of hepatocellular carcinoma, *Proc. Natl. Acad. Sci. USA* 87 (1990) 6547–6549.
- [5] D.L. Thomas, Hepatitis C epidemiology, *Curr. Top. Microbiol. Immunol.* 242 (2000) 25–41.
- [6] N. Kato, M. Hijikata, Y. Ootsuyama, M. Nakagawa, S. Ohkoshi, T. Sugimura, K. Shimotohno, Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis, *Proc. Natl. Acad. Sci. USA* 87 (1990) 9524–9528.
- [7] T. Tanaka, N. Kato, M.-J. Cho, K. Shimotohno, A novel sequence found at the 3' terminus of hepatitis C virus genome, *Biochem. Biophys. Res. Commun.* 215 (1995) 744–749.
- [8] M. Hijikata, N. Kato, Y. Ootsuyama, M. Nakagawa, K. Shimotohno, Gene mapping of the putative structural region of the hepatitis C virus genome by in vitro processing analysis, *Proc. Natl. Acad. Sci. USA* 88 (1991) 5547–5551.
- [9] M. Hijikata, H. Mizushima, Y. Tanji, Y. Komada, Y. Hirowatari, T. Akagi, N. Kato, K. Kimura, K. Shimotohno, Proteolytic processing and membrane association of putative nonstructural proteins of hepatitis C virus, *Proc. Natl. Acad. Sci. USA* 90 (1993) 10773–10777.
- [10] N. Kato, Molecular virology of hepatitis C virus, *Acta Med. Okayama* 55 (2001) 133–159.
- [11] G.R. Stark, I.M. Kerr, B.R.G. Williams, R.H. Silverman, R.D. Schreiber, How cells respond to interferons? *Annu. Rev. Biochem.* 67 (1998) 227–264.
- [12] C.E. Samuel, Antiviral actions of interferons, *Clin. Microbiol. Rev.* 14 (2001) 778–809.
- [13] J.F. Lau, C.M. Horvath, Mechanisms of Type I interferon cell signaling and STAT mediated transcriptional responses, *Mt. Sinai J. Med.* 69 (2002) 156–168.
- [14] M.G. Katze, Y. He, M. Gale Jr., Viruses and interferon: a fight for supremacy, *Nat. Rev. Immunol.* 2 (2002) 675–687.
- [15] A. Naganuma, A. Nozaki, T. Tanaka, K. Sugiyama, H. Takagi, M. Mori, K. Shimotohno, N. Kato, Activation of the interferon-inducible 2'-5'-oligoadenylate synthetase gene by hepatitis C virus core protein, *J. Virol.* 74 (2000) 8744–8750.
- [16] H. Dansako, A. Naganuma, T. Nakamura, F. Ikeda, A. Nozaki, N. Kato, Differential activation of interferon-inducible genes by hepatitis C virus core protein mediated by the interferon stimulated response element, *Virus Res.* 97 (2003) 17–30.
- [17] N. Kato, K. Shimotohno, Systems to culture hepatitis C virus, *Curr. Top. Microbiol. Immunol.* 242 (2000) 261–278.
- [18] E. Foy, K. Li, C. Wang, R. Sumpter Jr., M. Ikeda, S.M. Lemon, M. Gale Jr., Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease, *Science* 300 (2003) 1145–1148.
- [19] K. Li, E. Foy, J.C. Ferreon, M. Nakamura, A.C.M. Ferreon, M. Ikeda, S.C. Ray, M. Gale Jr., S.M. Lemon, Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2992–2997.
- [20] K. Li, Z. Chen, N. Kato, M. Gale Jr., S.M. Lemon, Distinct poly-I:C and virus-activated signaling pathways leading to interferon- β production in hepatocytes, *J. Biol. Chem.* 280 (2005) 16739–16747.
- [21] M. Ikeda, K. Sugiyama, T. Mizutani, T. Tanaka, K. Tanaka, H. Sekihara, K. Shimotohno, N. Kato, Human hepatocyte clonal cell lines that support persistent replication of hepatitis C virus, *Virus Res.* 56 (1998) 157–167.
- [22] M. Noguchi, S. Hirohashi, Cell lines from non-neoplastic liver and hepatocellular carcinoma tissue from a single patient, *In Vitro Cell Dev. Biol. Anim.* 32 (1996) 135–137.
- [23] K. Sugiyama, N. Kato, T. Mizutani, M. Ikeda, T. Tanaka, K. Shimotohno, Genetic analysis of the hepatitis C virus (HCV) genome from HCV-infected human T cells, *J. Gen. Virol.* 78 (1997) 329–336.
- [24] T. Akagi, T. Shishido, K. Murata, H. Hanafusa, v-Crk activates the phosphoinositide 3-kinase/AKT pathway in transformation, *Proc. Natl. Acad. Sci. USA* 97 (2000) 7290–7295.
- [25] P. Benech, M. Vigneron, D. Peretz, M. Revel, J. Chebath, Interferon-responsive regulatory elements in the promoter of the human 2',5'-oligo(A) synthetase gene, *Mol. Cell. Biol.* 7 (1987) 4498–4504.
- [26] N. Kato, K. Sugiyama, K. Namba, H. Dansako, T. Nakamura, M. Takami, K. Naka, A. Nozaki, K. Shimotohno, Establishment of a hepatitis C virus subgenomic replicon derived from human hepatocytes infected in vitro, *Biochem. Biophys. Res. Commun.* 306 (2003) 756–766.
- [27] K. Abe, M. Ikeda, H. Dansako, K. Naka, K. Shimotohno, N. Kato, cDNA microarray analysis to compare HCV subgenomic replicon cells with their cured cells, *Virus Res.* 107 (2005) 73–81.
- [28] A. Naganuma, H. Dansako, T. Nakamura, A. Nozaki, N. Kato, Promotion of microsatellite instability by hepatitis C virus core protein in human non-neoplastic hepatocyte cells, *Cancer Res.* 64 (2004) 1307–1314.
- [29] S. Bressanelli, L. Tomei, F.A. Rey, R. De Francesco, Structural analysis of the hepatitis C virus RNA polymerase in complex with ribonucleotides, *J. Virol.* 76 (2002) 3482–3492.
- [30] T. Yamashita, S. Kaneko, Y. Shiota, W. Qin, T. Nomura, K. Kobayashi, S. Murakami, RNA-dependent RNA polymerase activity of the soluble recombinant hepatitis C virus NS5B protein truncated at the C-terminal region, *J. Biol. Chem.* 273 (1998) 15479–15486.
- [31] V.J. Leveque, R.B. Johnson, S. Parsons, J. Ren, C. Xie, F. Zhang, Q.M. Wang, Identification of a C-terminal regulatory motif in hepatitis C virus RNA-dependent RNA polymerase: structural and biochemical analysis, *J. Virol.* 77 (2003) 9020–9028.
- [32] M. Uchida, N. Hino, T. Yamanaka, H. Fukushima, T. Imanishi, Y. Uchiyama, T. Kodama, T. Doi, Hepatitis C virus core protein binds to a C-terminal region of NS5B RNA polymerase, *Hepatol. Res.* 22 (2002) 297–306.
- [33] H. Kishine, K. Sugiyama, M. Hijikata, N. Kato, H. Takahashi, T. Noshi, Y. Nio, M. Hosaka, Y. Miyinari, K. Shimotohno, Subgenomic replicon derived from a cell line infected with the hepatitis C virus, *Biochem. Biophys. Res. Commun.* 293 (2002) 993–999.
- [34] M. Hijikata, H. Mizushima, T. Akagi, S. Mori, N. Kakiuchi, N. Kato, T. Tanaka, K. Kimura, K. Shimotohno, Two distinct proteinase activities required for the processing of a putative nonstructural precursor protein of hepatitis C virus, *J. Virol.* 67 (1993) 4665–4675.
- [35] K. Imanaka, S. Tamura, K. Fukui, N. Ito, S. Kiso, Y. Imai, T. Naka, T. Kishimoto, S. Kawata, Y. Shinomura, The Kansai Viral Hepatitis Research Group, Enhanced expression of suppressor of cytokine signaling-1 in the liver of chronic hepatitis C: possible involvement in resistance to interferon therapy, *J. Viral Hepat.* 12 (2005) 130–138.
- [36] K. Miller, S. McArdle, M.J. Gale Jr., D.A. Geller, B. Tenover, J. Hiscott, D.R. Gretsch, S.J. Polyak, Effects of the hepatitis C virus core protein on innate cellular defense pathways, *J. Interferon Cytokine Res.* 24 (2004) 391–402.

- [37] E. Foy, K. Li, R. Sumpter Jr., Y.-M. Loo, C.L. Johnson, C. Wang, P.M. Fish, M. Yoneyama, T. Fujita, S.M. Lemon, M. Gale Jr., Control of antiviral defenses through hepatitis C virus disruption of retinoic acid-inducible gene-I signaling, *Proc. Natl. Acad. Sci. USA* 102 (2005) 2986–2991.
- [38] R. Sumpter Jr., Y.-M. Loo, E. Foy, K. Li, M. Yoneyama, T. Fujita, S.M. Lemon, M. Gale Jr., Regulating intracellular antiviral defense and permissiveness to hepatitis C virus RNA replication through a cellular RNA helicase, RIG-I, *J. Virol.* 79 (2005) 2689–2699.
- [39] K.J. Blight, J.A. McKeating, J. Marcotrigiano, C.M. Rice, Efficient replication of hepatitis C virus genotype 1a RNAs in cell culture, *J. Virol.* 77 (2003) 3181–3190.
- [40] M. Ikeda, M. Yi, K. Li, S.M. Lemon, Selectable subgenomic and genome-length dicistronic RNAs derived from an infectious molecular clone of the HCV-N strain of hepatitis C virus replicate efficiently in cultured Huh7 cells, *J. Virol.* 76 (2002) 2997–3006.
- [41] T. Pietschmann, V. Lohmann, A. Kaul, N. Krieger, G. Rinck, G. Rutter, D. Strand, R. Bartenschlager, Persistent and transient replication of full-length hepatitis C virus genomes in cell culture, *J. Virol.* 76 (2002) 4008–4021.
- [42] M. Ikeda, K. Abe, H. Dansako, T. Nakamura, K. Naka, N. Kato, Efficient replication of a full-length hepatitis C virus genome, strain O, in cell culture, and development of a luciferase reporter system, *Biochem. Biophys. Res. Commun.* 329 (2005) 1350–1359.
- [43] T. Wakita, T. Pietschmann, T. Kato, T. Date, M. Miyamoto, Z. Zhao, K. Murthy, A. Habermann, H.G. Krausslich, M. Mizokami, R. Bartenschlager, T.J. Liang, Production of infectious hepatitis C virus in tissue culture from a cloned viral genome, *Nat. Med.* 11 (2005) 791–796.
- [44] R.E. Lanford, B. Guerra, H. Lee, D.R. Averett, B. Pfeiffer, D. Chavez, L. Notvall, C. Bigger, Antiviral effect and virus–host interactions in response to alpha interferon, gamma interferon, poly(I)–poly(C), tumor necrosis factor alpha, and ribavirin in hepatitis C virus subgenomic replicons, *J. Virol.* 77 (2003) 1092–1104.

Distinct Poly(I-C) and Virus-activated Signaling Pathways Leading to Interferon- β Production in Hepatocytes*

Received for publication, December 16, 2004, and in revised form, February 28, 2005
Published, JBC Papers in Press, February 28, 2005, DOI 10.1074/jbc.M414139200

Kui Li^{‡§}, Zihong Chen[‡], Nobuyuki Kato[¶], Michael Gale, Jr.^{||}, and Stanley M. Lemon^{‡**}

From the [‡]Department of Microbiology and Immunology, Institute for Human Infections and Immunity, University of Texas Medical Branch, Galveston, Texas 77555-1019, the [¶]Department of Molecular Biology, Okayama University Graduate School of Medicine and Dentistry, Okayama 700-8558, Japan, and the ^{||}Department of Microbiology, University of Texas Southwestern Medical Center, Dallas, Texas 75390-9048

Innate cellular antiviral defenses are likely to influence the outcome of infections by many human viruses, including hepatitis B and C viruses, agents that frequently establish persistent infection leading to chronic hepatitis, cirrhosis, and liver cancer. However, little is known of the pathways by which hepatocytes, the cell type within which these hepatitis agents replicate, sense infection, and initiate protective responses. We show that cultured hepatoma cells, including Huh7 cells, do not activate the interferon (IFN)- β promoter in response to extracellular poly(I-C). In contrast, the addition of poly(I-C) to culture media activates the IFN- β promoter and results in robust expression of IFN-stimulated genes (ISG) in PH5CH8 cells, which are derived from non-neoplastic hepatocytes transformed with large T antigen. Small interfering RNA knockdown of TLR3 or its adaptor, Toll-interleukin-1 receptor domain-containing adaptor inducing IFN- β (TRIF), blocked extracellular poly(I-C) signaling in PH5CH8 cells, whereas poly(I-C) responsiveness could be conferred on Huh7 hepatoma cells by ectopic expression of Toll-like receptor 3 (TLR3). In contrast to poly(I-C), both cell types signal the presence of Sendai virus infection through a TLR3-independent intracellular pathway requiring expression of retinoic acid-inducible gene I (RIG-I), a putative cellular RNA helicase. Silencing of RIG-I expression impaired only the response to Sendai virus and not extracellular poly(I-C). We conclude that hepatocytes contain two distinct antiviral signaling pathways leading to expression of type I IFNs, one dependent upon TLR3 and the other dependent on RIG-I, with little cross-talk between these pathways.

The innate immune system represents the first line of defense against viral infections in mammalian cells. It senses invading viral pathogens and initiates signaling pathways

leading to the induction of protective cellular genes, including type I interferons (IFN- α and IFN- β)¹ and proinflammatory cytokines that directly limit viral replication and also help to shape subsequent adaptive immune responses (1, 2). Recognition of conserved molecular structures that are expressed by large groups of pathogens (pathogen-associated molecular patterns, PAMPs) is carried out by specific pattern recognition receptor (PRR) molecules (3, 4). The Toll-like receptors (TLRs) are a class of PRRs that have been shown to detect infection by many types of pathogens, including viruses (5). TLR3 is engaged specifically by double-stranded (ds) RNA that is present either in viral genomes or generated during viral replication, and is involved in the cellular recognition of RNA viruses and induction of type I IFN responses (6). However, several recent studies indicate that viral infection can also activate host responses through TLR3-independent pathways (7–9). Mouse TLR7 and human TLR8, which are expressed within endosomal membranes, detect GU-rich viral single-stranded RNA (7, 10, 11). Whereas TLR3 signaling requires the adaptor protein, Toll-IL1 receptor (TIR) domain-containing adaptor inducing IFN- β (TRIF/TICAM-1) (12–15), TLR7/8 engagement utilizes the MyD88 adaptor protein and requires endosomal acidification for activation of IFN responses (7, 10). Finally, virally encoded proteins may engage TLR2 or TLR4, activating MyD88-dependent pathways leading to expression of inflammatory cytokines and contributing to both viral clearance as well as pathogenesis (16–18).

The TLRs are not the only class of PAMP receptors that contribute to the recognition of virus infection. Yoneyama and colleagues (19) have demonstrated recently that IFN- β production is induced in response to Newcastle disease virus infection through a pathway that is independent of TLR3 but requires the retinoic acid-inducible gene I (RIG-I, or DDX58), a cellular RNA helicase with homology to caspase-recruitment domain (CARD) proteins.

The ability of these pathways to induce the expression of type I IFNs and subsequently a wide array of IFN-stimulated genes (ISGs) is likely to influence the outcome of infection by many human viruses. Such responses may have direct antiviral effects within the infected cell (2). Type I IFNs also exert

* This work was supported in part by National Institutes of Health Grants R21-DA018054 (to K. L.), U19-AI40035 (to S. M. L.), U01-AI48235 (to M. G.), and in part by Gastrointestinal Research Interdisciplinary Program Seed Money (to K. L.). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

§ John Mitchell Hemophilia of Georgia Liver Scholar of the American Liver Foundation. To whom correspondence may be addressed: 4.168A Medical Research Bldg., University of Texas Medical Branch, 301 University Blvd., Galveston, TX 77555-1019. Tel.: 409-772-4928; Fax: 409-772-5065; E-mail: kuli@utmb.edu.

** To whom correspondence may be addressed: Institute for Human Infections and Immunity, The University of Texas Medical Branch at Galveston, 301 University Blvd., Galveston, TX 77555-0428. Tel.: 409-747-6500; Fax: 409-747-6514; E-mail: smlemon@utmb.edu.

¹ The abbreviations used are: IFN, interferon; dsRNA, double-stranded RNA; HCV, hepatitis C virus; IRF-3, interferon regulatory factor 3; ISG, interferon-stimulated gene; MDA5, melanoma differentiation associated gene-5; MyD88, myeloid differentiation factor 88; RIG-I, retinoic acid-inducible gene I; SenV, Sendai virus; siRNA, small interfering RNA; TLR, Toll-like receptor; TRIF, Toll-IL1 receptor domain-containing adaptor inducing IFN- β ; PRR, pattern recognition receptor; PAMP, pathogen-associated molecular patterns; CARD, caspase-recruitment domain; HBV, hepatitis B virus; pAb, polyclonal antibody; RT, reverse transcription; HAU, hemagglutinin unit(s); GAPDH, glyceraldehyde-3-phosphate dehydrogenase.

critically important immunoregulatory effects, including stimulation of natural killer cell cytotoxicity, activation of $\gamma\delta$ T cells, and stimulation of immunoglobulin synthesis. Cytokines and chemokines induced by the activation of these signaling pathways also contribute to the maturation of dendritic cells and influence the priming of T_H1 and cytotoxic T cells (20). These responses thus may play critical roles in shaping subsequent adaptive T cell responses that are required for the ultimate elimination of viruses (21, 22).

Although the liver is a particularly important site of persistent viral infections in humans, very little is known about how these signaling pathways function specifically in hepatocytes. Despite this, there is strong, albeit indirect evidence, that type I IFN responses are important in the pathogenesis of chronic viral hepatitis. Both hepatitis B virus (HBV) and hepatitis C virus (HCV) cause persistent infections involving the hepatocyte, and both have evolved mechanisms to disrupt the induction of type I IFNs. Although not well understood, the core protein of HBV inhibits the transcription of IFN- β (23, 24). Similarly, the NS3/4A protease of HCV blocks Sendai virus (SenV)-induced activation of IFN regulatory factor 3 (IRF-3), a cellular transcription factor that plays a critical role in the expression of IFN- β (25). IRF-3 blockade is dependent upon the protease activity of NS3/4A, suggesting that NS3/4A proteolytically targets one or more cellular proteins residing within the signaling pathways leading to IRF-3 activation and type I IFN production (25). These and other data (26, 27) suggest that a more detailed understanding of the mechanisms by which virus infections trigger IRF-3 activation and IFN production in hepatocytes would be helpful in unraveling the pathogenesis of persistent HBV and HCV infections, and might possibly lead to the design of novel therapeutic interventions. Here, we describe efforts to better define the antiviral signaling pathways that are active in cultured hepatocyte-derived cell lines and that thus may be triggered by HBV or HCV infection *in vivo*.

MATERIALS AND METHODS

Cell Culture and Reagents—Murine macrophage RAW264.7, human hepatoma Huh7, Huh7.5 (kindly provided by C. M. Rice via Apath), HepG2, Hep3B cells, and PH5CH8, a simian virus 40 (SV40) large T antigen-immortalized non-neoplastic human hepatocyte cell line (28), were maintained in Dulbecco's modified Eagle's medium supplemented with 10% heat-inactivated fetal bovine serum, 100 units/ml penicillin G and streptomycin in a humidified 37 °C, 5% CO₂ incubator. Poly(I-C) was purchased from Sigma. Bafilomycin A1 was from Calbiochem. Recombinant human interleukin-1 β was from Raybiotech. R-848 was kindly provided by K. A. Fitzgerald.

Plasmids—pIFN- β -luc, pCMV1FlagTLR3, pCMV1FlagTLR3DN, and pEF-Bos Flag-RIG-I were generous gifts of Rongtuan Lin, Ruslan Medzhitov, and Takashi Fujita, respectively. The ISG54 ISRE-Luc and pNF- κ B-Luc plasmids were from Stratagene. Sequence encoding a dominant negative form of MyD88, which lacks the amino-terminal death domain (29), was amplified from cDNA transcribed from PH5CH8 cell RNA, and cloned into pCDNA3.1 V5-His TOPO (Invitrogen). pPRDII-luc has been described previously (30). pCMV- β -galactosidase (Clontech) was used to normalize transfection efficiencies. Cells were transfected with plasmid DNAs using TransIT-LT1 (Mirus) according to the manufacturer's instructions.

Sendai Virus Infection—Cells were infected with 100 hemagglutinin units/ml of SenV (Charles River Laboratory) and harvested 16 h later for luciferase/ β -galactosidase reporter assays or immunoblot analysis as previously described (25).

Poly(I-C) Treatment—Poly(I-C) was added directly to the medium at 50 μ g/ml (M-pIC), or complexed with Lipofectin for transfection (T-pIC). Cells were assayed for poly(I-C) induced responses 6 h after exposure by either route.

Reporter Gene Assay—Cells (5×10^4 cells per well in 24-well plates) were transfected with reporter plasmids (100 ng), pCMV- β -galactosidase (100 ng), and the indicated amounts of an expression vector. Twenty-four hours later, cells were mock treated or treated with 20 ng/ml interleukin-1 β , or challenged with poly(I-C) or Sendai virus, then subsequently lysed and assayed for luciferase and β -galactosidase activities as indicated. For

TABLE I
Gene-specific primers for semiquantitative RT-PCR

Gene name		Primer sequence ^a	Product size bp
IFN- β	Forward	gattcatctagactgctggtgg	186
	Reverse	cttcaggtaatgcagaatcc	
TLR3	Forward	tcactgtcctattctcctt	157
	Reverse	gacctctccattctctggc	
TRIF	Forward	ccagatgcaacctccactgg	339
	Reverse	ctgttccgatgatgatcc	
ISG56	Forward	tagccaacatgtctccacagac	396
	Reverse	tctctaccactggttctatgc	
RIG-I	Forward	cagtatattcaggctgag	389
	Reverse	ggccagttttctctgtc	
MDA5	Forward	agtttggcagaaggaagtgc	480
	Reverse	ggagttttcaaggatttgagc	

^a GAPDH primers were purchased from Clontech.

comparisons, luciferase activity was normalized to β -galactosidase activity. Data are expressed as mean relative luciferase activity \pm S.D. for one representative experiment carried out in triplicate, typically from a minimum of three separate experiments. The -fold induction of promoter activity was calculated by dividing the relative luciferase activity of stimulated cells with that of mock-treated cells.

Immunoblot Analysis—Cellular extracts were subjected to immunoblot analysis as described (25). Briefly, protein samples were separated by SDS-PAGE, and transferred to Hybond enhanced chemiluminescent (ECL) nitrocellulose membranes (Amersham Biosciences). The membranes were blocked with 3% nonfat milk in phosphate-buffered saline and processed for immunodetection using the following monoclonal or polyclonal (pAb) antibodies: anti-FLAG M2 and anti-actin monoclonal antibodies (Sigma), anti-TLR3 monoclonal antibody (Imgenex), rabbit anti-MyD88 pAb (Santa Cruz), rabbit anti-IRF-3 pAb (kindly provided by Michael David), rabbit anti-ISG15 pAb (kindly provided by Arthur Haas), rabbit anti-ISG56 pAb (kindly provided by Ganes Sen), rabbit anti-MxA and anti-Sendai virus pAbs (kindly provided by Ilkka Julkunen), peroxidase-conjugated secondary donkey anti-rabbit (Jackson ImmunoResearch), and sheep anti-mouse (Amersham Biosciences) pAbs. Protein bands were visualized using ECL Plus Western blotting detection reagents (Amersham Biosciences), followed by exposure to Kodak Bio-Max film.

RNA Interference—Transfections of siRNAs targeting TLR3, TRIF, or RIG-I were carried out using Oligofectamine (Invitrogen) with a final concentration of siRNA of 80 nM according to the manufacturer's instructions. The target sequences of siRNA used in this study were: TLR3, GGTATAGCCAGCTAACTAG; TRIF (13), GACCAGACGC-CACTCCAAC; RIG-I, GGAAGAGGTGCAGTATATT; and MDA5, GGTGAAGGAGCAGATTTCAG. siRNAs were purchased from Dharmacon and Ambion. The scrambled negative control siRNA was from Ambion.

RT-PCR—Total cellular RNA was extracted with TRIzol Reagent (Invitrogen), treated with DNase I to remove genomic DNA contamination, and reverse-transcribed using Advantage RT-4PCR kit (Clontech). The resulting cDNA was subjected to PCR using primers (Table I) specific for IFN- β , TLR3, TRIF, ISG56, RIG-I, MDA5, β -actin, and GAPDH, respectively. The quantity of the cDNA template included in these reactions and the number of amplification cycles were optimized to ensure that reactions were stopped during the linear phase of product amplification, permitting semiquantitative comparisons of mRNA abundance between different RNA preparations. To exclude the possibility of contaminating DNA, control reactions were performed in parallel in the absence of reverse transcriptase. PCR products were visualized by agarose gel electrophoresis.

RESULTS

Cultured Hepatocyte Cell Lines Differ in Their Ability to Activate IFN- β Transcription in Response to dsRNA or Virus Infection—The induction of type I IFNs represents an early protective response to many viral infections in mammalian cells. IFN- β induction represents the immediate response of cells to viral infection, and precedes the transcription of most IFN- α species, the induction of which depends on autocrine/paracrine feedback of IFN- β and activation of IRF-7 (31). We thus focused on characterizing the IFN- β response. The molecular basis for induction of IFN- β expression has been exten-

sively studied and shown to be induced by dsRNA or other products of virus infection through coordinate activation of transcription factors IRF-3, NF- κ B, and ATF-2/c-Jun (2). Most of these studies were conducted in human embryonic kidney 293 or epithelial cells, and relatively little is known about these events in hepatocytes, which constitute ~80% of the liver cell population (32), and are the primary cells within which both HBV and HCV replicate. Because primary differentiated hepatocytes are difficult to maintain in culture, we studied 3 different continuous cell lines derived from human hepatocellular carcinomas: Huh7 cells, which appear to be particularly permissive for HCV RNA replication (33, 34); HepG2 cells and Hep3B cells, the latter of which contains integrated HBV DNA and express the HBV envelope protein, HBsAg (35). Huh7 cells are not available from any standard repository, but are carried in many different laboratories and different laboratory variants may show significant variation in morphology and perhaps other characteristics. We thus studied 3 different Huh7 sublines: Huh7 SL, cells carried in this laboratory for many years; Huh7 2-3c, cells derived from Huh7 SL that were selected for their ability to support replication of genome-length HCV RNA and subsequently "cured" of the replicating RNA with IFN- α 2b treatment (36), and Huh7 MG cells, Huh7 cells carried in the Gale laboratory at the University of Texas Southwestern Medical Center. We also studied the PH5CH8 cell line, which is a clonal variant of a non-neoplastic hepatocyte cell line, PH5CH, immortalized with the simian virus 40 large T antigen (28, 37, 38).

We characterized virus-induced activation of the IFN- β promoter in these various hepatocyte-derived cell lines after transient transfection with a reporter plasmid expressing luciferase under control of the IFN- β promoter. Although the PH5CH8 cell line has been reported to support replication of HCV (28), no cultured cell has been reported to be fully for HCV replication. Thus, to challenge these cells, we exposed them to the dsRNA analog, poly(I-C), or infected them with SenV. Interestingly, when poly(I-C) was added to the culture medium, there was no induction of IFN- β promoter activity in the Huh7, HepG2, or Hep3B hepatoma cells (M-pIC, Fig. 1A, upper panel). In contrast, the PH5CH8 cells demonstrated a 6-fold up-regulation of IFN- β promoter activity upon exposure to poly(I-C) (Fig. 1A, upper panel). Consistent with the reporter data, M-pIC treatment significantly up-regulated the expression of ISG15 and ISG56, which are responsive to either IRF-3 or IFN, in PH5CH8 cells, but not in hepatoma cell lines Huh7, HepG2, or Hep3B (Fig. 1B). To mimic intracellular dsRNA generated during viral replication, we also transfected poly(I-C) into cells using a liposome-mediated procedure. This resulted in much more potent stimulation of the IFN- β promoter in PH5CH8 cells, leading to a 17-fold increase in activity over basal promoter levels (T-pIC, Fig. 1A, middle panel). It also resulted in significant activation of the IFN- β promoter in HepG2 (8-fold induction) and Hep3B (4-fold induction) cells. However, under the same conditions, there was little if any activation of the promoter in any of the Huh7 cell sublines. These promoter assay results were confirmed by RT-PCR analysis of endogenous IFN- β mRNA synthesis in cells transfected with increasing concentrations of poly(I-C) (Fig. 1C). Whereas transfection of as little as 1 μ g/ml poly(I-C) induced IFN- β transcription in HepG2, Hep3B, and PH5CH8 cells, there was no detectable IFN- β mRNA in Huh7 SL cells transfected with up to 100 μ g/ml poly(I-C).

A similar pattern of IFN- β induction was observed following infection of these different cell lines with SenV. The greatest induction of IFN- β promoter activity occurred in PH5CH8 cells




FIG. 1. Activation of IFN- β transcription in various hepatocyte cell lines by dsRNA and virus. A, cells grown in 24-well plates were cotransfected with IFN- β Luc and pCMV- β -galactosidase for 24 h before stimulation with 50 μ g/ml poly(I-C) directly added to culture medium for 6 h (M-pIC, upper panel), 5 μ g of poly(I-C) transfected with Lipofectin for 6 h (T-pIC, middle panel), or SenV 100 HAU/ml for 16 h (lower panel). -Fold induction of IFN- β promoter was calculated by dividing the relative luciferase activity of stimulated cells with that of mock-treated cells. B, immunoblot analysis of ISG15 and ISG56 expression in various hepatocyte cell lines either mock-treated or treated with 50 μ g/ml M-pIC for 12 h. Actin was included as a loading control. C, cells were mock-treated (Lipofectin) or transfected with differing concentrations of poly(I-C) for 6 h before total RNA isolation. IFN- β mRNA was detected by semiquantitative RT-PCR.

(96-fold increase over basal promoter activity), followed by HepG2 (35-fold), and Hep3B (10-fold) (Fig. 1A, lower panel). However, there were significant differences in the magnitude of the IFN- β promoter activation induced by SenV infection in the different Huh7 cell lines. SenV infection induced IFN- β promoter activity by ~10-fold in the Huh7 MG cells, but only 1.5-fold in Huh7 SL cells and 3-fold in the cured Huh7 2-3c cells (Fig. 1A, lower panel).

We conclude from these results that cultured hepatoma cells generally have impaired poly(I-C)- and virus-activated IFN responses, compared with the PH5CH8 cells that were established from non-neoplastic hepatocytes. These latter cells retain robust IFN- β responses to extracellular and intracellular poly(I-C) as well as SenV infection. They are likely to more closely resemble normal hepatocytes *in vivo*, and thus may represent a superior cell model for investigation of antiviral responses in hepatocytes. The considerable variation we observed in the IFN response in different Huh7 cell lines mandates caution in comparing studies carried out with these cells in different laboratories, an important point because Huh7 cells are widely used for cell culture studies with HCV.

TLR3 and TRIF Are Not Required for SenV Activation of the IFN- β Promoter in PH5CH8 Cells—TLR3 is the only known TLR that recognizes viral dsRNA and it is expressed both on the cell surface and within intracellular vesicles (39, 40). We were thus interested in determining whether the virus-induced IFN response involves TLR3 in PH5CH8 cells. To investigate this, we transfected the cells with a synthetic siRNA duplex targeting human TLR3, or a scrambled negative-control siRNA, then challenged the cells by adding poly(I-C) to the media or infecting them with SenV prior to measuring IFN- β




FIG. 2. Extracellular dsRNA, but not SenV, activates IFN response via TLR3 and TRIF in PH5CH8 hepatocytes. *A*, TLR3 expression is required for activation of IFN- β promoter by M-pIC but not by SenV. PH5CH8 cells grown in 24-well plates were transfected with control (*Ctrl*) or TLR3 siRNA, and pIFN- β Luc plus pCMV- β -galactosidase. 48 h later, cells were mock-treated or stimulated with 50 μ g/ml M-pIC for 6 h or infected with SenV (100 HAU/ml) for 16 h. *B*, semiquantitative RT-PCR detection of TLR3 and GAPDH mRNAs in control/TLR3 siRNA-transfected cells. *C*, TRIF expression is required for activation of IFN- β promoter by M-pIC but not by SenV. PH5CH8 cells were transfected with *Ctrl* or TRIF siRNA and reporter plasmids and treated similarly as in *A*. *D*, semiquantitative RT-PCR detection of TRIF, ISG56, and GAPDH mRNAs in cells under conditions as described for panel *C*.

promoter activity (Fig. 2). PH5CH8 cells express a readily detectable abundance of TLR3 mRNA, which was reduced to almost undetectable levels upon transfection of TLR3-specific siRNA (Fig. 2*B*). The induction of IFN- β promoter activity by extracellular poly(I-C) was substantially reduced by this siRNA knockdown of TLR3 expression (Fig. 2*A, left panel*). In contrast, there was no appreciable effect on SenV-induced IFN- β promoter activation (Fig. 2*A, right panel*).

TRIF is an essential adaptor protein that links TLR3 to downstream kinases responsible for IRF-3 activation and IFN- β production (13–15). It may well play additional roles in intracellular signaling events, given its relatively large size (712 amino acids) compared with other TLR adaptors. Mice that are deficient for TRIF function have an impaired response to murine cytomegalovirus infection (12). The liver contains a higher abundance of TRIF than any other organ (14), suggesting that it may play a particularly important role in intrahepatic signaling. To determine whether TRIF is required to mount a type I IFN response against SenV in hepatocytes, we carried out a TRIF knockdown experiment in PH5CH8 cells. As shown in Fig. 2*D*, semiquantitative RT-PCR demonstrated that transfection of the TRIF-specific siRNA efficiently reduced TRIF mRNA abundance. As expected from the TLR3 knockdown (Fig. 2*A*), TRIF knockdown also significantly inhibited the induction of IFN- β promoter activity in response to extracellular poly(I-C) (Fig. 2*C, left panel*). It

also significantly reduced extracellular poly(I-C) induction of ISG56 mRNA transcription (Fig. 2*D*, compare lanes 2 versus 4). In contrast, there was no effect on either IFN- β promoter activity (Fig. 2*C, right panel*) or ISG56 mRNA transcription (Fig. 2*D*, compare lanes 6 versus 8) triggered by SenV infection. Similar results were obtained in an ISG54 ISRE promoter assay (data not shown). We conclude from these data that exposure to extracellular poly(I-C) triggers activation of the IFN- β promoter in PH5CH8 cells through a TLR3-TRIF dependent pathway, whereas SenV induction of IFN- β transcription is TLR3- and TRIF-independent.

SenV Induction of IFN- β in Hepatocytes Does Not Utilize MyD88 nor Require Endosomal Acidification—In addition to TLR3, recent data indicate that TLR7, TLR8, and TLR9 may sense viral components and lead to type I IFN production. Murine TLR7 and human TLR8 recognize viral ssRNA, whereas TLR9 senses unmethylated CpG DNA present in the murine cytomegalovirus genome (41). A common feature of these three TLRs is that they all localize within endosomes and signal through MyD88 and IRF-7 (42, 43). In addition, the responses induced through these TLRs require intact endocytic pathways and are thus sensitive to endosomal acidification inhibitors, such as chloroquine and bafilomycin A1 (7, 10). Because SenV induction of IFN- β occurs via a TLR3-TRIF independent pathway in PH5CH8 cells, we considered the possibility that TLR7/8 may initiate this response. To determine whether MyD88 is required for SenV activation in the PH5CH8 cells, as would be the case were it mediated by TLR7, TLR8, or TLR9, we measured promoter activation in PH5CH8 cells transfected with a vector expressing a dominant-negative MyD88 mutant that lacks the amino-terminal death domain that is required for interaction with IRF-7 (42) (MyD88DN). Expression of MyD88DN significantly inhibited the activation of the NF- κ B-dependent PRDII promoter (44) by interleukin-1 β , which is known to signal through MyD88 (29). However, we found that the activity of both PRDII and IFN- β promoters was induced to similar levels by SenV with or without MyD88DN co-expression (Fig. 3*A*). Consistent with these reporter data, SenV-induced hyperphosphorylation of IRF-3 was not affected by overexpression of MyD88DN (Fig. 3*B*). This was also true for SenV-induced expression of ISG56, which is responsive to either IRF-3 or IFN (45), and MxA, which responds only to IFN (46).

To determine whether SenV activation of the IFN- β promoter requires endosomal acidification, PH5CH8 cells were pretreated with the endosomal inhibitor bafilomycin A1 prior to infection with SenV. Although bafilomycin A1 treatment completely ablated NF- κ B activation in RAW264.7 cells induced by a TLR7/8 ligand, R-848 (Fig. 3*C*), SenV-stimulated IFN- β and PRDII promoter activity was minimally reduced in cells treated with the compound (Fig. 3*D*). Moreover, immunoblot analyses indicated that there were no differences in SenV-induced hyperphosphorylation of IRF-3, or ISG15 and MxA expression in these cells in the presence or absence of bafilomycin A1 (Fig. 3*E*). Bafilomycin A1 did not alter the expression of SenV proteins.

These data suggest that SenV induction of IFN responses is MyD88-independent in PH5CH8 cells, and does not require active endocytic pathways. Both lines of evidence argue strongly against the involvement of TLR7/8. In addition, because MyD88 is thought to be used by all TLRs other than TLR3 (and TLR4 for signaling to IRF-3), these data, taken in the context of the absence of any inhibition of the SenV response by TRIF knockdown (Fig. 2, *C* and *D*), suggest that the recognition of SenV by PH5CH8 cells is not dependent upon any known TLR.




FIG. 3. SenV activation of type I IFN response in hepatocytes is MyD88-independent and does not require endosomal acidification. A, PH5CH8 cells in 24-well plates were cotransfected with the indicated reporter plasmids (100 ng), pCMV- β -galactosidase (100 ng), and 600 ng of a dominant-negative form of MyD88 (*MyD88DN*) or empty vector (V) for 24 h before mock-infected or infected with SenV (100 HAU/ml) for 16 h (*middle and right panels*) or mock/treated with 20 ng/ml interleukin-1 β for 8 h (*left panel*). B, immunoblot detection of IRF-3, MxA, ISG56, SenV, and MyD88 in PH5CH8 cells using the conditions as described for the *middle and right panels* of A. C, NF- κ B promoter activity in RAW264.7 cells mock-treated or treated with bafilomycin A1 and then mock-treated with R-848. Cells transfected with pNF- κ B Luc and pCMV- β -galactosidase for 24 h were pretreated with medium alone (M), or 100 nM bafilomycin A1 (*Bf-A1*) for 1 h and then mock-treated with 1 μ M R-848 for 6 h in the presence of M/Bf-A1. D, SenV-induced IFN- β and PRDII promoter activity in PH5CH8 cells treated with medium alone or Bf-A1. E, immunoblot detection of IRF-3, ISG15, MxA, and SenV in PH5CH8 cells as described for the conditions of *panel D*.

SenV Activates Type I IFN Responses through a RIG-I-dependent Pathway in Hepatocytes—Studies with knock-out mice have indicated that both virus infection and dsRNA can trigger type I IFN responses via TLR3-independent mechanisms (7, 8). Recently, Yoneyama and colleagues (19) demonstrated that RIG-I, a putative DExD/H box RNA helicase containing an N-terminal sequence with CARD-like homology domains, is essential for IFN- β production induced in response to infection with Newcastle disease virus. To determine whether SenV induces IFN- β promoter activity in PH5CH8 cells through a RIG-I-dependent pathway, we utilized RNA interference to knockdown RIG-I expression prior to virus challenge. Transfection of a RIG-I-specific siRNA, but not a scrambled control siRNA, reproducibly caused a ~50% reduction in SenV-induced activation of both the IFN- β (Fig. 4A, *right panel*) and PRDII promoters (Fig. 4B, *right panel*). Semiquantitative RT-PCR confirmed that the transfection of RIG-I siRNA efficiently knocked down the basal expression of RIG-I (Fig. 4C, *lanes 2 and 4* in the *left panel* and *lanes 7 and 11*, in *right panel*), and also significantly blunted the up-regulation of this IFN-induced




FIG. 4. SenV, but not TLR3 engagement, activates type I IFN response through RIG-I in hepatocytes. PH5CH8 cells grown in 24-well plates were transfected with control (*Ctrl*) or RIG-I siRNA, and pIFN- β Luc (A) or PRDII Luc (B), and pCMV- β -galactosidase. 48 h later, cells were mock-treated or stimulated with 50 μ g/ml M-pIC for 6 h or infected with SenV (100 HAU/ml) for 16 h. C, semiquantitative RT-PCR detection of RIG-I, ISG56, and GAPDH mRNAs in PH5CH8 cells under conditions of A and B. The *left panel* shows efficient knockdown of the basal expression of RIG-I in PH5CH8 cells by RIG-I siRNA determined by increasing cycles of RT-PCR. D, immunoblot analysis of ISG15 and SenV protein expression in PH5CH8 cells under conditions of *panels A and B*. Actin was included as a loading control.

protein upon SenV infection (Fig. 4C, *right panel*, compare *lanes 9 and 10* with *lanes 11 and 12*). SenV-induced transcription of ISG56 mRNA was also significantly reduced in cells transfected with RIG-I siRNA (Fig. 4C, *lanes 9–12*). Therefore, RIG-I is an essential component in the TLR-independent pathway by which PH5CH8 cells sense SenV infection and initiate a type I IFN response.

In contrast to these results, PH5CH8 cells transfected with the RIG-I siRNA responded normally when poly(I-C) were added to the media, in terms of activation of both the IFN- β (Fig. 4A, *left panel*) and NF- κ B-dependent PRDII (Fig. 4B, *left panel*) promoters. In contrast to SenV-induced signaling, the poly(I-C) induction of RIG-I transcription was not dramatically reduced by transfection of the RIG-I siRNA (Fig. 4C, compare *lanes 5 and 6* with *lanes 7 and 8*). Similarly, there was no inhibition of poly(I-C) induction of ISG56 transcription (Fig. 4C, *lanes 5–8*). We also confirmed the RT-PCR results by immunoblot analysis of ISG15 expression under these conditions (Fig. 4D). These data indicate that TLR3- and RIG-I-mediated signaling function independently of each other in PH5CH8




FIG. 5. Poly(I-C), as well as SenV activation of the IFN- β promoter is partially MDA5-dependent in hepatocytes. *A*, SenV (*left panel*) and M-pIC (*right panel*)-induced IFN- β promoter activity in PH5CH8 cells transfected with the indicated siRNAs. *B*, immunoblot detection of ISG56 and SenV proteins in PH5CH8 cells transfected with Ctrl (*lanes 1, 2, and 5, 6*) or MDA5 siRNA (*lanes 3, 4, and 7, 8*). Where indicated, cells were treated with 50 μ g/ml M-pIC for 8 h or infected with 100 HAU/ml SenV for 16 h. In *lane 9*, cells were treated with 500 units/ml of IFN- α 2b as a control for ISG56 expression. *C*, semiquantitative RT-PCR detection of MDA5 and GAPDH mRNAs in control (Ctrl)/MDA5 siRNA-transfected PH5CH8 cells.

cells, similar to what has been reported previously for epithelial HeLa cells (19).

Poly(I-C) Activation of the IFN- β Promoter Is Partially MDA5-dependent—Because siRNA knockdown of RIG-I did not ablate but only partially reduced the IFN- β promoter response to SenV infection in PH5CH8 cells (Fig. 4A, *right panel*), we carried out similar experiments to determine whether the human melanoma differentiation associated gene-5 product (MDA5) participates in this signaling pathway, possibly in a redundant role with respect to RIG-I. MDA5, another DExD/H box RNA helicase, is an IFN-inducible protein that shares a subdomain architecture and considerable sequence homology with RIG-I (47). It has been associated with the induction of apoptosis, but not clearly identified as playing a role in activation of IRF-3 similar to that of RIG-I. siRNA-mediated knockdown of MDA5 expression had only a minor, but reproducible, suppressive effect on SenV induction of the IFN- β promoter. The magnitude of the reduction in promoter activity was substantially less than that observed with RIG-I knockdown (Fig. 5A, *left panel*), but it was associated nonetheless within a modest reduction in SenV-induced expression of ISG56 (Fig. 5B, compare *lanes 6 versus 8*). The suppressive effect of MDA5 knockdown was not additive with RIG-I knockdown, as the degree of suppression of promoter activity was not increased when both RIG-I and MDA5 siRNAs were cotransfected into PH5CH8 cells. The MDA5 knockdown had no effect on SenV protein expression (Fig. 5B).

Interestingly, MDA5 knockdown caused \sim 30% reduction in the induction of IFN- β promoter activity by extracellular poly(I-C) (Fig. 5A, *right panel*). This effect was less than that observed with TLR3 or TRIF knockdown, yet nonetheless reproducible and significant. Consistent with this, MDA5 knockdown also caused a moderate reduction in poly(I-C)-induced





FIG. 6. The defect of TLR3 signaling in Huh7 cells is because of insufficient expression of TLR3. *A*, semiquantitative RT-PCR detection of TLR3, TRIF, and β -actin mRNAs in Huh7 and PH5CH8 cells. *B*, ectopic expression of FLAG-tagged TLR3 restored the induction of ISG15, ISG56, and MxA expression by M-pIC (50 μ g/ml) determined by immunoblot analysis. A nonspecific band detected by the TLR3 antibody serves as a loading control. In *lane 5*, cells were treated with 500 units/ml of IFN- α 2b as a positive control for ISG expression.

ISG56 expression (Fig. 5B, compare *lanes 2 and 4*). We conclude from these results that MDA5, like RIG-I, may contribute to viral activation of the IFN- β promoter but, unlike RIG-I, plays a greater role in poly(I-C) activation of the promoter than in SenV-induced responses. Whether this occurs in association with, or independent of, the TLR3-TRIF pathway remains to be determined.

TLR3 Overexpression Reconstitutes Poly(I-C)-induced ISG Expression in Huh7 Cells—As shown in Fig. 1, all three Huh7 cell sublines studied were defective in poly(I-C) signaling. This lack of poly(I-C) responsiveness may be explained by the fact that Huh7 cells express a negligible abundance of TLR3 mRNA, in contrast to PH5CH8 cells that demonstrate robust TLR3 expression (Fig. 6A). In contrast, TRIF mRNA abundance was approximately equal in these cell lines. Overexpression of FLAG-tagged TLR3 effectively restored the ability of Huh7 cells to respond to poly(I-C) when added to the culture medium (50 μ g/ml), as indicated by expression of ISG15, ISG56, and MxA (Fig. 6B, compare *lanes 2 and 4*). Thus, Huh7 cells are normally deficient in their ability to respond to externally applied poly(I-C) because of a lack of sufficient expression of TLR3.

TLR3- and RIG-I-mediated Signaling Function Independently in Huh7.5 Cells—TLR3 signaling has been reported not to be adversely affected by overexpression of a dominant-negative form of RIG-I in HeLa cells (19). Consistent with this, the siRNA knockdown experiments shown in Figs. 2 and 4 suggest that these pathways function largely independently of each other in PH5CH8 cells. To confirm this in another hepatocyte-derived cell line, we utilized a Huh7 subline, Huh7.5, which is highly permissive for replication of HCV RNA replicons (48) and which has recently been shown to be defective for RIG-I

Fig. 7. TLR3 and RIG-I signaling pathways function independently in Huh7.5 cells. *A*, the defect of SenV-induced ISG15 expression can be restored by overexpression of RIG-I, but not TLR3 in Huh7.5 cells. *B*, the defect of extracellular dsRNA-induced expression of ISG15 and ISG56 can be restored by overexpression of TLR3, but not a mutant TLR3 with the TIR domain deleted (Δ TIR) in Huh7.5 cells. *C*, ectopic expression of TLR3, but not RIG-I, restored the response to M-pIC in Huh7.5 cells. All panels shown were immunoblot analysis. Where indicated, cells were treated with 50 μ g/ml M-pIC for 12 h, or 100 HAU/ml SenV for 16 h, or 500 units/ml of IFN- α 2b. Please note that the TLR3 antibody detects transfected FLAG-RIG-I for unknown reasons. The superintense bands of RIG-I detected by anti-FLAG antibody is likely because of the 2 \times FLAG tag *versus* 1 \times FLAG tag in TLR3.



signaling because of a point mutation within its CARD-like homology domain (49). Thus, in contrast to PH5CH8 cells in which both TLR3 and RIG-I pathways are intact, neither pathway is functional in Huh7.5 cells. By reconstituting expression of either TLR3 or a functional RIG-I molecule in Huh7.5 cells, we were thus able to assess the role played by each signaling pathway independently in recognition of viral infection while determining the extent to which these pathways act independently of each other.

We first confirmed previous observations (49) that SenV-induced signaling through RIG-I is deficient in Huh7.5 cells, by demonstrating that SenV-induced expression of ISG15 was dependent upon ectopic expression of RIG-I in these cells (Fig. 7A). In contrast, ectopic expression of TLR3 did not rescue SenV-induced ISG15 expression in Huh7.5 cells, suggesting that the products of SenV infection do not engage TLR3, or that optimal TLR3 signaling requires functional RIG-I. The latter is unlikely, however, as RIG-I knockdown had no effect on IFN induction by extracellular poly(I-C) in PH5CH8 cells (Fig. 4). Moreover, ectopic expression of TLR3 alone was able to restore responsiveness to extracellular poly(I-C) in Huh7.5 cells (Fig. 7B, lanes 8 *versus* 10), to a degree similar to that in normal Huh7 cells (Fig. 7C, lanes 15 *versus* 21). The rescue of poly(I-C) responsiveness was specific for TLR3, as overexpression of a TIR domain-deleted TLR3 mutant failed to restore expression of ISG15 and ISG56 in response to poly(I-C) (Fig. 7B, lanes 10 *versus* 12), nor did RIG-I (Fig. 7C, lanes 15 *versus* 17). Dual ectopic expression of TLR3 and RIG-I only slightly increased the ISG15 induction in response to poly(I-C) (Fig. 7C, lanes 18 and 19). Taken together, the data suggest that there is little if any cross-talk between the TLR3- and RIG-I pathways in hepatocytes.

DISCUSSION

We have shown here that cultured hepatoma cells generally have impaired poly(I-C) and viral-activated IFN responses,

compared with PH5CH8 cells that are derived from normal hepatocytes (Fig. 1). These data are thus in agreement with a previous report (50) showing that hepatoma cells have impaired antiviral responses. HepG2 and Hep3B cells are deficient in signaling in response to extracellular poly(I-C), and appear to have diminished responses to either intracellularly delivered poly(I-C) or SenV infection. As described by others (51, 52), we found that Huh7 cells also failed to respond to poly(I-C), irregardless of whether it was added to the culture medium or introduced into cells via transfection. In addition, only a weak response was observed in some Huh7 cell sublines (Huh7 2–3c and MG) after infection with SenV, a more potent IFN inducer. Poly(I-C) signaling could be restored to Huh7 cells by ectopic expression of TLR3 (Fig. 6), indicating that the lesion in Huh7 cells is at the level of the PRR molecule, and that the downstream pathway involving TRIF is intact. In contrast, the non-neoplastic PH5CH8 cells retain robust IFN- β responses to extracellular and intracellular poly(I-C) as well as SenV infection, and thus may be more representative of the antiviral signaling pathways present within hepatocytes *in vivo*.

dsRNA is commonly expressed during the replication of most RNA viruses, and has long been considered a candidate viral PAMP. Treatment of mammalian cells with the synthetic dsRNA analog, poly(I-C), induces type I IFN production through activation of transcription factors IRF-3 and NF- κ B (53). TLR3 is well characterized as a PRR that is engaged specifically by dsRNA in many cell types. We have shown here that it is expressed in non-neoplastic hepatocytes, and that it plays an important role in these cells in activation of the IFN- β promoter following exposure to extracellular poly(I-C) (Fig. 2A). Furthermore, we have shown that poly(I-C) signaling through TLR3 leading to the induction of ISG expression is dependent upon the adaptor protein TRIF in these cells, as might be expected (Fig. 2C). Thus, the data presented demonstrate clearly that the TLR3-TRIF pathway is functional in the non-neoplastic hepatocyte-derived PH5CH8 cells, although it is

generally absent in cultured cells derived from hepatocellular carcinomas (Fig. 1).

Importantly, neither RNA interference directed silencing of TLR3 nor TRIF resulted in any impairment of IFN- β promoter activation or ISG expression in SenV-infected PH5CH8 cells (Fig. 2). This indicates that dsRNA, or possibly another PAMP produced during SenV replication, triggers activation of the IFN- β promoter in these hepatocytes through a distinctly different signaling pathway. This is consistent with several recent reports that dsRNA and/or virus infection can initiate antiviral signaling through TLR3-independent pathways (7, 8). SenV could potentially activate signaling through recognition of viral ssRNA by endosomally located TLR7/8, as reported for murine plasmacytoid dendritic cells and B lymphocytes (7). However, neither endosomal acidification inhibitors nor expression of a dominant-negative MyD88 mutant affected the response in SenV-infected PH5CH8 cells (Fig. 3). Thus, SenV appears to activate a TLR-independent intracellular signaling pathway in hepatocytes that is distinct from the TLR3-TRIF pathway activated by extracellular poly(I-C).

Consistent with a previous study carried out in mouse fibroblasts and human epithelial cells (19), we found that RIG-I is an essential component of this TLR-independent antiviral signaling pathway in cells derived from non-neoplastic hepatocytes (Fig. 4). Similarly, we have recently shown that SenV activation of IRF-3 is dependent upon RIG-I expression in hepatoma cells as well (49). RIG-I is a cytoplasmic RNA helicase that contains tandem motifs near its N terminus with limited homology to CARD domains and a downstream DExD/H-box helicase domain. It putatively binds viral dsRNA within its helicase domain, resulting in activation of IRF-3 and NF- κ B through signaling involving the N-terminal CARD-like homology domains (19). MDA5 is a closely related DExD/H box helicase with similar domain architecture, and has been associated with apoptotic signaling in terminally differentiating melanoma cells (47). Its expression is induced by IFN- α as well as infection with human immunodeficiency virus (47, 54). We found that it contributes to the induction of IFN signaling by extracellular poly(I-C) (Fig. 5), but has only a minor, yet clearly discernible, contribution to SenV induction of IFN- β promoter activity and downstream ISG expression. Whether it functions in a fashion similar to that proposed for RIG-I, as a PRR molecule binding dsRNA through its helicase domain, seems likely but remains to be demonstrated. It is also not known whether RIG-I and MDA5 are evolved to preferentially recognize dsRNAs derived from viruses. The relationship of MDA5 to the TLR3-TRIF pathway also remains to be defined.

Although HCV replicon RNAs expressing selectable markers are capable of replicating in some alternative cell types (55), Huh7 cells have proven to be nearly unique in their ability to support the autonomous replication of these viral RNAs (34). In addition, we previously demonstrated that Huh7 cells are uniquely permissive for self-amplification of subgenomic replicons derived from hepatitis A virus, another positive-strand RNA virus (56). Huh7.5, a Huh7 subline that is highly permissive for HCV RNA replication (48), has a lethal mutation in the RIG-I CARD-like homology domain that renders it unresponsive to structured HCV RNA or SenV induced signaling (49). Whereas this accounts for the highly permissive phenotype of Huh7.5 cells, it does not fully explain why parental Huh7 cells are already more permissive than other cell types. Although relatively little is known regarding the cellular factors associated with permissiveness for HCV (34, 57–59), it is intriguing to speculate that the enhanced permissiveness for positive-strand virus RNA replication in normal Huh7 cells may relate to the absence of significant TLR3-mediated antiviral re-

sponses in these cells. It remains to be determined, however, whether HCV RNA replication results in specific engagement of TLR3 and activation of the downstream signaling pathway, and whether the TLR3 signaling cascade acts to limit HCV replication. Recent data indicate that expression of the HCV NS3/4A protease inhibits poly(I-C)-induced, TLR3-dependent signaling by directing the proteolytic cleavage of TRIF in osteosarcoma cells as well as in HeLa cells supporting replication of subgenomic HCV RNA replicons (26). However, for reasons that remain unclear, the ectopic expression of NS3/4A does not appear to block TLR3 signaling in PH5CH8 cells.²

In summary, hepatocytes contain two distinct antiviral signaling pathways leading to expression of type I IFNs, one dependent upon TLR3 and the other on RIG-I, and with little evidence of significant cross-talk between them. However, although the RIG-I and TLR3 pathways function independently in hepatocytes (Figs. 4 and 7), it is very likely that during viral infection *in vivo* both pathways are activated in a coordinate manner to enhance innate immune responses. Activation of the RIG-I pathway by the intracellular presentation of dsRNA or another viral PAMP produced by replicating viruses may contribute to the initial induction of type I IFNs. Subsequently, viral dsRNA released by lysis of cells in later stages of the infection may engage TLR3 and induce TRIF-dependent signaling, resulting in further amplification of the antiviral response (53). Importantly, type I IFNs induced through either pathway would have a positive feedback on signaling through both pathways, as both RIG-I and TLR3 are ISGs (19, 60).

Acknowledgments—We thank Drs. Ilkka Julkunen, Michael David, Arthur Haas, Rongtuan Lin, Ganes Sen, Ruslan Medzhitov, Katherine Fitzgerald, and Takashi Fujita for providing reagents.

REFERENCES

- Janssens, S., and Beyaert, R. (2003) *Clin. Microbiol. Rev.* **16**, 637–646
- Sen, G. C. (2001) *Annu. Rev. Microbiol.* **55**, 255–281
- Boehme, K. W., and Compton, T. (2004) *J. Virol.* **78**, 7867–7873
- Gordon, S. (2002) *Cell* **111**, 927–930
- Akira, S., and Hemmi, H. (2003) *Immunol. Lett.* **85**, 85–95
- Alexopoulou, L., Holt, A. C., Medzhitov, R., and Flavell, R. A. (2001) *Nature* **413**, 732–738
- Lund, J. M., Alexopoulou, L., Sato, A., Karow, M., Adams, N. C., Gale, N. W., Iwasaki, A., and Flavell, R. A. (2004) *Proc. Natl. Acad. Sci. U. S. A.* **101**, 5598–5603
- Honda, K., Sakaguchi, S., Nakajima, C., Watanabe, A., Yanai, H., Matsumoto, M., Ohteki, T., Kaisho, T., Takaoka, A., Akira, S., Seya, T., and Taniguchi, T. (2003) *Proc. Natl. Acad. Sci. U. S. A.* **100**, 10872–10877
- Edelmann, K. H., Richardson-Burns, S., Alexopoulou, L., Tyler, K. L., Flavell, R. A., and Oldstone, M. B. (2004) *Virology* **322**, 231–238
- Diebold, S. S., Kaisho, T., Hemmi, H., Akira, S., and Reis e Sousa, C. (2004) *Science* **303**, 1529–1531
- Heil, F., Hemmi, H., Hochrein, H., Ampenberger, F., Kirschning, C., Akira, S., Lipford, G., Wagner, H., and Bauer, S. (2004) *Science* **303**, 1526–1529
- Hoebe, K., Du, X., Georgel, P., Janssen, E., Tabet, K., Kim, S. O., Goode, J., Lin, P., Mann, N., Mudd, S., Crozat, K., Sovath, S., Han, J., and Beutler, B. (2003) *Nature* **424**, 743–748
- Oshiumi, H., Matsumoto, M., Funami, K., Akazawa, T., and Seya, T. (2003) *Nat. Immunol.* **4**, 161–167
- Yamamoto, M., Sato, S., Mori, K., Hoshino, K., Takeuchi, O., Takeda, K., and Akira, S. (2002) *J. Immunol.* **169**, 6668–6672
- Yamamoto, M., Sato, S., Hemmi, H., Hoshino, K., Kaisho, T., Sanjo, H., Takeuchi, O., Sugiyama, M., Okabe, M., Takeda, K., and Akira, S. (2003) *Science* **301**, 640–643
- Kurt-Jones, E. A., Popova, L., Kwinn, L., Haynes, L. M., Jones, L. P., Tripp, R. A., Walsh, E. E., Freeman, M. W., Golenbock, D. T., Anderson, L. J., and Finberg, R. W. (2000) *Nat. Immunol.* **1**, 398–401
- Haynes, L. M., Moore, D. D., Kurt-Jones, E. A., Finberg, R. W., Anderson, L. J., and Tripp, R. A. (2001) *J. Virol.* **75**, 10730–10737
- Dolganiuc, A., Oak, S., Kodys, K., Golenbock, D. T., Finberg, R. W., Kurt-Jones, E., and Szabo, G. (2004) *Gastroenterology* **127**, 1513–1524
- Yoneyama, M., Kikuchi, M., Natsukawa, T., Shinobu, N., Imaizumi, T., Miyagishi, M., Taira, K., Akira, S., and Fujita, T. (2004) *Nat. Immunol.* **5**, 730–737
- Beignon, A. S., Skoberne, M., and Bhardwaj, N. (2003) *Nat. Immunol.* **4**, 939–941
- Iwasaki, A., and Medzhitov, R. (2004) *Nat. Immunol.* **5**, 987–995
- Hoebe, K., Janssen, E., and Beutler, B. (2004) *Nat. Immunol.* **5**, 971–974
- Twu, J. S., Lee, C. H., Lin, P. M., and Schloemer, R. H. (1988) *Proc. Natl. Acad.*

² K. Li, unpublished data.

- Sci. U. S. A.* **85**, 252–256
24. Whitten, T. M., Quets, A. T., and Schloemer, R. H. (1991) *J. Virol.* **65**, 4699–4704
 25. Foy, E., Li, K., Wang, C., Sumpster, R., Jr., Ikeda, M., Lemon, S. M., and Gale, M., Jr. (2003) *Science* **300**, 1145–1148
 26. Li, K., Foy, E., Ferreon, J. C., Nakamura, M., Ferreon, A. C., Ikeda, M., Ray, S. C., Gale, M., Lemon, S. M. (2005) *Proc. Natl. Acad. Sci. U. S. A.* **102**, 2992–2997
 27. Foy, E., Li, K., Sumpster, R., Loo, Y.-M., Johnson, C., Wang, C., Fish, P., Yoneyama, M., Fujita, T., Lemon, S. M., and Gale, M., Jr. (2005) *Proc. Natl. Acad. Sci. U. S. A.* **102**, 2986–2991
 28. Ikeda, M., Sugiyama, K., Mizutani, T., Tanaka, T., Tanaka, K., Sekihara, H., Shimotohno, K., and Kato, N. (1998) *Virus Res.* **56**, 157–167
 29. Muzio, M., Ni, J., Feng, P., and Dixit, V. M. (1997) *Science* **278**, 1612–1615
 30. Fredericksen, B., Akkaraju, G. R., Foy, E., Wang, C., Pflugheber, J., Chen, Z. J., and Gale, M., Jr. (2002) *Viral Immunol.* **15**, 29–40
 31. Levy, D. E., Marie, I., and Prakashi, A. (2003) *Curr. Opin. Immunol.* **15**, 52–58
 32. Kmiec, Z. (2001) *Adv. Anat. Embryol. Cell Biol.* **161**, III–XIII, 1–151
 33. Nakabayashi, H., Taketa, K., Miyano, K., Yamane, T., and Sato, J. (1982) *Cancer Res.* **42**, 3858–3863
 34. Lohmann, V., Korner, F., Koch, J., Herian, U., Theilmann, L., and Bartenschlager, R. (1999) *Science* **285**, 110–113
 35. Knowles, B. B., Howe, C. C., and Aden, D. P. (1980) *Science* **209**, 497–499
 36. Scholle, F., Li, K., Bodola, F., Ikeda, M., Luxon, B. A., and Lemon, S. M. (2004) *J. Virol.* **78**, 1513–1524
 37. Noguchi, M., and Hirohashi, S. (1996) *In Vitro Cell Dev. Biol. Anim.* **32**, 135–137
 38. Kato, N., Ikeda, M., Mizutani, T., Sugiyama, K., Noguchi, M., Hirohashi, S., and Shimotohno, K. (1996) *Jpn. J. Cancer Res.* **87**, 787–792
 39. Matsumoto, M., Funami, K., Oshiumi, H., and Seya, T. (2004) *Microbiol. Immunol.* **48**, 147–154
 40. Matsumoto, M., Funami, K., Tanabe, M., Oshiumi, H., Shingai, M., Seto, Y., Yamamoto, A., and Seya, T. (2003) *J. Immunol.* **171**, 3154–3162
 41. Tabeta, K., Georgel, P., Janssen, E., Du, X., Hoebe, K., Crozat, K., Mudd, S., Shamel, L., Sovath, S., Goode, J., Alexopoulou, L., Flavell, R. A., and Beutler, B. (2004) *Proc. Natl. Acad. Sci. U. S. A.* **101**, 3516–3521
 42. Honda, K., Yanai, H., Mizutani, T., Negishi, H., Shimada, N., Suzuki, N., Ohba, Y., Takaoka, A., Yeh, W. C., and Taniguchi, T. (2004) *Proc. Natl. Acad. Sci. U. S. A.* **101**, 15416–15421
 43. Kawai, T., Sato, S., Ishii, K. J., Coban, C., Hemmi, H., Yamamoto, M., Terai, K., Matsuda, M., Inoue, J., Uematsu, S., Takeuchi, O., and Akira, S. (2004) *Nat. Immunol.* **5**, 1061–1068
 44. Lenardo, M. J., Fan, C. M., Maniatis, T., and Baltimore, D. (1989) *Cell* **57**, 287–294
 45. Nakaya, T., Sato, M., Hata, N., Asagiri, M., Suemori, H., Noguchi, S., Tanaka, N., and Taniguchi, T. (2001) *Biochem. Biophys. Res. Commun.* **283**, 1150–1156
 46. Ronni, T., Matikainen, S., Lehtonen, A., Palvimo, J., Dellis, J., Van Eylen, F., Goetschy, J. F., Horisberger, M., Content, J., and Julkunen, I. (1998) *J. Interferon Cytokine Res.* **18**, 773–781
 47. Kang, D. C., Gopalkrishnan, R. V., Lin, L., Randolph, A., Valerie, K., Pestka, S., and Fisher, P. B. (2004) *Oncogene* **23**, 1789–1800
 48. Blight, K. J., McKeating, J. A., and Rice, C. M. (2002) *J. Virol.* **76**, 13001–13014
 49. Sumpster, R., Jr., Loo, Y. M., Foy, E., Li, K., Yoneyama, M., Fujita, T., Lemon, S. M., and Gale, M. J., Jr. (2005) *J. Virol.* **79**, 2689–2699
 50. Keskinen, P., Nyqvist, M., Sareneva, T., Pirhonen, J., Melen, K., and Julkunen, I. (1999) *Virology* **263**, 364–375
 51. Lanford, R. E., Guerra, B., Lee, H., Averett, D. R., Pfeiffer, B., Chavez, D., Notvall, L., and Bigger, C. (2003) *J. Virol.* **77**, 1092–1104
 52. McCormick, C. J., Challinor, L., Macdonald, A., Rowlands, D. J., and Harris, M. (2004) *J. Gen. Virol.* **85**, 429–439
 53. Levy, D. E., and Marie, I. J. (2004) *Nat. Immunol.* **5**, 699–701
 54. Cocude, C., Truong, M. J., Billaut-Mulot, O., Delsart, V., Darcissac, E., Capron, A., Mouton, Y., and Bahr, G. M. (2003) *J. Gen. Virol.* **84**, 3215–3225
 55. Zhu, Q., Guo, J. T., and Seeger, C. (2003) *J. Virol.* **77**, 9204–9210
 56. Yi, M., and Lemon, S. M. (2002) *J. Virol.* **76**, 1171–1180
 57. Bartenschlager, R., Kaul, A., and Sparacio, S. (2003) *Antiviral Res.* **60**, 91–102
 58. Bartenschlager, R., and Lohmann, V. (2001) *Antiviral Res.* **52**, 1–17
 59. Blight, K. J., Kolykhalov, A. A., and Rice, C. M. (2000) *Science* **290**, 1972–1974
 60. Miettinen, M., Sareneva, T., Julkunen, I., and Matikainen, S. (2001) *Genes Immun.* **2**, 349–355
 61. Deleted in proof
 62. Deleted in proof

Short
CommunicationInterferon resistance of hepatitis C virus
replicon-harboured cells is caused by functional
disruption of type I interferon receptorsKazuhito Naka,¹ Kazunori Takemoto,¹ Ken-ichi Abe,¹ Hiromichi Dansako,¹
Masanori Ikeda,¹ Kunitada Shimotohno² and Nobuyuki Kato¹

Correspondence

Nobuyuki Kato

nkato@md.okayama-u.ac.jp

¹Department of Molecular Biology, Okayama University Graduate School of Medicine, Dentistry
and Pharmaceutical Sciences, 2-5-1 Shikata-cho, Okayama 700-8558, Japan²Department of Viral Oncology, Institute for Virus Research, Kyoto University, 53 Kawara-cho
Shogo-in, Sakyo-ku, Kyoto 606-8507, Japan

Received 21 April 2005

Accepted 5 July 2005

Hepatitis C virus (HCV) replicon-harboured cell lines possessing interferon (IFN)-resistant phenotypes have recently been established. These were divided into two classes: partially IFN resistant and highly IFN resistant. Here, the viral and cellular factors contributing to the IFN resistance of HCV replicon-harboured cells were evaluated. The results revealed that cellular factors rather than viral factors contributed to a highly IFN-resistant phenotype. The possibility of genetic abnormality of the factors involved in IFN signalling was investigated. As a result, nonsense mutations and deletions in type I IFN receptor genes (IFNAR1 and IFNAR2c) were found in replicon-harboured cells showing a highly IFN-resistant phenotype, but rarely appeared in cells showing a partially IFN-resistant phenotype. Furthermore, similar genetic alterations were also found in IFN-resistant phenotype, replicon-harboured cell lines obtained additionally by IFN- β treatment. Moreover, it was shown that ectopic expression of wild-type IFNAR1 in IFN-resistant phenotype, replicon-harboured cells possessing the IFNAR1 mutant restored type I IFN signalling.

Persistent infection by hepatitis C virus (HCV) is a major cause of chronic hepatitis (Choo *et al.*, 1989; Kuo *et al.*, 1989), which can progress to liver cirrhosis and hepatocellular carcinoma (Saito *et al.*, 1990). Since at least 170 million people are currently infected with HCV worldwide, this infection constitutes a global health problem (Thomas, 2000). HCV is an enveloped RNA virus belonging to the family *Flaviviridae*, the genome of which consists of a positive-stranded RNA encoding an approximately 3000 aa polyprotein precursor (Kato *et al.*, 1990). This precursor protein is processed by the host and viral proteases to generate at least 10 proteins in the following order: core, envelope 1 (E1), E2, p7, non-structural protein 2 (NS2), NS3, NS4A, NS4B, NS5A and NS5B (Kato, 2001).

Since 1998, combined treatment with interferon (IFN)- α and ribavirin has been standard clinical therapy for patients with chronic hepatitis C; however, the effectiveness of IFN is limited to approximately 50% (Hadziyannis *et al.*, 2004). This clinical result suggests that HCV directly or indirectly attenuates the antiviral actions of IFN (Pawlotsky, 2000).

Although an HCV replicon system carrying autonomously replicating HCV subgenomic RNA containing the NS3–NS5B regions (Lohmann *et al.*, 1999) was considered to be

useful in studies on the mechanism(s) of IFN resistance of HCV, all HCV replicons established to date have been highly sensitive to IFN- α , - β and - γ (Frese *et al.*, 2001, 2002; Kato *et al.*, 2003). This seems to contradict the fact that half of the patients with chronic hepatitis C are resistant to current IFN therapy. Therefore, we assumed that prolonged IFN treatment might change HCV replicons from an IFN-sensitive phenotype to an IFN-resistant phenotype.

Based on this assumption, we recently established nine HCV replicon cell lines possessing two IFN-resistant phenotypes: a partially resistant phenotype (α R series: 1 α R, 3 α R, 4 α R, 5 α R and α Rmix) and a highly resistant phenotype (β R series: 1 β R, 3 β R, 4 β R and 5 β R) obtained by IFN- α and - β treatment, respectively (Namba *et al.*, 2004). Genetic analysis of these replicons found one common amino acid substitution (Q1737H) in the NS4B region and several additional amino acid substitutions (such as M2174V and T2242N) in the NS5A region of the β R series (Namba *et al.*, 2004). To examine which viral and cellular factors contribute to the IFN resistance of HCV replicons, we evaluated the IFN sensitivity of replicon-harboured cells (6M.m/6Mc, 1 β R.m/6Mc and 4 β R.m/6Mc) established by transfection of total RNAs isolated from an IFN-sensitive clone (6M) and from highly IFN-resistant clones (1 β R and 4 β R) into cured 6Mc cells, from which 50–1 replicons (Kishine *et al.*, 2002) had been eliminated by IFN- γ treatment (500 IU ml⁻¹ for

Supplementary material is available in JGV Online.




Fig. 1. IFN sensitivities of various HCV replicon-harboring cells. (a) 6M.m/6Mc, 1βR.m/6Mc and 4βR.m/6Mc cells obtained as G418-resistant mixed colonies were treated with IFN-α (400 IU ml⁻¹) for 3 weeks in the presence of G418 (300 μg ml⁻¹). 6M, 1βR and 4βR cells were also used for control experiments. G418-resistant colonies were stained with Coomassie brilliant blue as described previously (Naganuma *et al.*, 2004). (b) IFN sensitivities of 6MK1/6Mc, 1βR1/6Mc and 4βR2/6Mc cells were examined as described in (a) except that they were treated with IFN-β (200 IU ml⁻¹).

3 weeks). Although many colonies were found to have survived in 1βR and 4βR cells after IFN-α treatment, only a few colonies survived in 1βR.m/6Mc and 4βR.m/6Mc cells or in 6M and 6M.m/6Mc cells (Fig. 1a). Similar results were obtained when these replicon-harboring cells were treated with IFN-β (400 IU ml⁻¹) (data not shown), although two colonies (named 6βR and 7βR, described below) derived from 6M.m/6Mc and 4βR.m/6Mc cells, respectively, proliferated as highly IFN-resistant clones.

We further examined the IFN sensitivity of replicon-harboring cells (6MK1/6Mc, 1βR1/6Mc and 4βR2/6Mc) established by the transfection of *in vitro*-synthesized replicon RNAs (6MK1, 1βR1 and 4βR2 obtained from 6M, 1βR and 4βR cells, respectively) (Namba *et al.*, 2004; Kato *et al.*, 2005) into 6Mc cells. The results revealed that few or no colonies survived in 1βR1/6Mc and 4βR2/6Mc cells, as in

6M and 6MK1/6Mc cells, whereas many colonies survived in 1βR and 4βR cells (Fig. 1b). These results suggested that cellular factors rather than viral factors contributed to the highly IFN-resistant phenotype of HCV replicon-harboring cells. However, the present results obtained under a high concentration of IFN do not necessarily rule out a possible role for HCV mutations in conferring low degrees of IFN resistance, because effects of HCV mutations on IFN activity are presumably weaker than those of the cellular factors.

To obtain evidence in favour of the idea that alterations in cellular factor(s) are involved in the emergence of an IFN-resistant phenotype, we attempted to prepare cured cells from the replicon-harboring cells possessing a highly IFN-resistant phenotype. Since phosphorylation of signal transducer and activator of transcription 1 (STAT1) occurred in the 1βR and 4βR cells treated with IFN-γ (Fig. 2a), these replicon-harboring cells were treated with IFN-γ (500 IU ml⁻¹) for 3 weeks, and cured 1βRc and 4βRc cells were obtained. Western blot (Fig. 2b) and RT-PCR (data not shown) analyses showed that no replicons were detected in either type of cured cells. Analysis of a luciferase reporter assay indicated that the complete defect of the IFN-α/β signalling was not restored in the cured 1βRc and 4βRc cells (Fig. 2c).

To clarify whether or not the signalling defect in these replicon-harboring cells was restricted to type I IFN, we examined the phosphorylation status of STAT3 in 6M, 1βR and 4βR cells treated with interleukin-6 (IL6). Since it has been reported that STAT3 is also activated by IFN-α treatment (Pfeffer *et al.*, 1997), the phosphorylation status of STAT3 in these replicon-harboring cells after IFN-α treatment was also examined. Our results revealed that STAT3 was not phosphorylated in 1βR and 4βR cells treated with IFN-α, while phosphorylation of STAT3 was observed in 6M, 1βR and 4βR cells treated with IL6 and in 6M cells treated with IFN-α (Fig. 2d), indicating that only type I IFN signalling was defective in 1βR and 4βR cells. These results suggested that the initial reaction following the addition of IFN-α/β was defective in replicon-harboring cells possessing a highly IFN-resistant phenotype.

Following up this suggestion, we examined the genetic status of tyrosine kinase 2 (TYK2) and Janus kinase 1 (JAK1). Sequence analysis of TYK2 and JAK1 cDNAs obtained from 1βR and 4βR cells was performed after cloning into the pCXbsr vector (Akagi *et al.*, 2000), as described previously (Nozaki *et al.*, 2003). However, the results showed no mutations in these cDNAs (data not shown). We next focused on type I IFN receptors (IFNAR1 and IFNAR2c). Our results showed that the mRNA levels of the two receptors were almost equal among all examined replicon-harboring cells including 6Mc cells (see Supplementary Fig. S1a, available in JGV Online). More than three independent clones of each cDNA (1708 bp for IFNAR1 and 1582 bp for IFNAR2c) were sequenced as described above. Table 1 shows a summary of sequence analysis of IFNAR1




Fig. 2. Cellular factors rather than HCV replicons contribute to defects in type I IFN signalling. (a) Western blot analysis of STAT1 in 6M, 1βR and 4βR cells treated with IFN-α or -γ. The replicon-harboring cells were treated with or without IFN-α or -γ (500 IU ml⁻¹ each) for 30 min, and Western blot analysis for STAT1 and its phosphorylation status (Tyr-701) was then performed as previously described (Hijikata *et al.*, 1993). β-Actin was used as a control for the amount of protein loaded per lane. (b) Western blot analysis of NS5B. Anti-NS5B antibody was used for immunoblotting. β-Actin was used as described in (a). (c) Analysis of IFN signal transduction. Dual luciferase assays using pSRE(V2)-Luci (Dansako *et al.*, 2003) were performed as previously described (Naganuma *et al.*, 2000). Cells were treated with IFN-α or -β (500 IU ml⁻¹ each) for 6 h. (d) The defect in the signalling pathway in 1βR and 4βR cells is restricted to type I IFN. Replicon-harboring cells were left untreated or treated with IFN-α (500 IU ml⁻¹) or IL6 (100 ng ml⁻¹) for 30 min. Western blot analysis for STAT3 and its phosphorylation status (Tyr-705) was performed as described in (a).

and IFNAR2c mRNAs. Surprisingly, we found that nt 475 of IFNAR2c mRNA from 1βR cells had a U substituted for G in 13/13 clones, resulting in a nonsense mutation at codon 159 from glutamic acid (GAG) to the termination codon UAG (see Supplementary Fig. S1b). Furthermore, nt 319 of IFNAR1 mRNA from 4βR cells was also found to have a U substituted for G in 12/12 clones, resulting in a nonsense mutation at codon 107 from glutamic acid (GAA) to the termination codon UAA (see Supplementary Fig. S1c). However, interestingly, IFNAR1 and IFNAR2c mRNAs obtained from 1αR and 4αR cells, which were counterparts of 1βR and 4βR cells, respectively (see Supplementary Fig. S2, available in JGV Online), did not possess the nonsense mutations found in the mRNAs from 1βR and 4βR cells. In addition, several kinds of deletion and another nonsense mutation (lysine to a termination codon at codon 458) were found in approximately half of IFNAR1 cDNA clones obtained from 3βR and

5βR cells. In contrast to the finding of frequent mutations and deletions in IFNAR mRNAs from the βR series, such genetic abnormalities in IFN receptors were quite rare in the αR series (Table 1).

To evaluate the possibility that genetic mutants might pre-exist in the cloned replicon-harboring cells (clones 1, 3, 4 and 5 shown in supplementary Fig. S2) or that genetic mutants had appeared during the IFN-β treatment of the cloned replicon-harboring cells, we repeated IFN-β treatment of cloned 1, 3, 4 and 5 cells following the method described previously (Namba *et al.*, 2004). The result was almost identical to that obtained previously (Namba *et al.*, 2004), indicating the good reproducibility of the experiment with IFN-β treatment (data not shown). In the present study, each of the three colonies showing resistance to IFN-β was isolated and proliferated (see Supplementary Fig. S2) and we then performed sequence analysis of IFNAR mRNAs

Table 1. Genetic alterations of type I IFN receptors in HCV replicon-harboring cell lines possessing IFN-resistant phenotypes

The determined nucleotide sequences were compared with those of 6M and 6Mc cells, which confirmed that the deduced amino acid sequences were identical to the human IFNAR1 (GenBank accession no. NM_000629) and IFNAR2c (GenBank accession no. L41942) sequences. NM, Not mutated.

Cell line	IFNAR1 mRNA			IFNAR2c mRNA		
	Nucleotide change and position	Effect on protein	Mutation frequency*	Nucleotide change and position	Effect on protein	Mutation frequency*
6M	NM	None	0/3	NM	None	0/3
1 β R	NM	None	0/3	G to U at nt 475	E to stop at codon 159	13/13
3 β R	Deletion of 5 nt at nt 376	Truncated (135 aa)	2/11	NM	None	0/3
	A to U at nt 1372	K to stop at codon 458	3/11			
4 β R	G to U at nt 319	E to stop at codon 107	12/12	NM	None	0/3
5 β R	Deletion of 176 nt at nt 201	Truncated (78 aa)	4/10	NM	None	0/3
	Deletion of 79 nt at nt 201	Truncated (67 aa)	2/10			
1 α R	NM	None	0/3	NM	None	0/3
3 α R	NM	None	0/3	NM	None	0/3
4 α R	Deletion of 5 nt at nt 376	Truncated (135 aa)	1/3	NM	None	0/3
5 α R	Deletion of 79 nt at nt 201	Truncated (67 aa)	1/3	NM	None	0/3
α Rmix	NM	None	0/3	NM	None	0/3
6Mc	NM	None	0/3	NM	None	0/3

*Number of mutated or truncated clones/number of examined clones.

as described above. The results revealed that the nonsense mutations or deletions identified at this time (see Supplementary Table S1, available in JGV Online) were quite different from those obtained from the β R series (Table 1). Therefore, it is unlikely that the identified IFNAR mutants pre-existed in cloned 1, 3, 4 and 5 cells when these cells were obtained as colonies surviving IFN- α treatment.

To examine whether or not additional HCV replicon cell lines possessing the IFN-resistant phenotype could be obtained from HCV replicon-harboring cells other than the parental replicon-harboring cells used for the isolation of the α R and β R series, 6M.m/6Mc, 1 β R.m/6Mc, 4 β R.m/6Mc and 50-1 replicon-harboring cells were treated with IFN- β (see Supplementary Fig. S2). Finally, we obtained four replicon-harboring cell lines (6 β R obtained from the 4 β R.m/6Mc cells, 7 β R obtained from the 6M.m/6Mc cells, and 8 β R and 9 β R obtained from the 50-1 cells) showing resistance to IFN- β . These results indicated that HCV replicon-harboring cells showing the IFN-resistant phenotype were obtained from HCV replicon-harboring cells established immediately. By sequence analysis of IFNAR1 and IFNAR2c cDNAs as described above, the E107stop nonsense mutation in IFNAR1 cDNA, which was the same mutation found in the 4 β R cells, was found again in the 8 β R and 9 β R cells, while no IFNAR mutations were detected in the 6 β R and 7 β R cells (see Supplementary Table S1). The observation that IFNAR mutations occurred preferentially after IFN- β treatment is interesting. Since a variety of

mutations and deletions in the IFN receptors were obtained from the cloned replicon-harboring cells surviving after IFN- β treatment, such genetic alterations might occur accidentally in order to impair the antiviral states caused after IFN- β treatment. Thereafter, only replicon-harboring cells possessing the IFNAR mutants might be able to proliferate in the presence of G418, resulting in the β R series.

To clarify whether or not the IFNAR mutations found in the β R series were determinants for the IFN sensitivity of HCV replicons, we prepared 4 β R cells (possessing the IFNAR1 mutant) stably expressing wild-type IFNAR1 and examined its IFN sensitivity. Analysis of a luciferase reporter assay (see Supplementary Fig. S3a, available in JGV Online) clearly showed that IFN signalling in 4 β R cells was restored by the expression of wild-type IFNAR1 in comparison with those of 4 β R cells expressing the IFNAR1 mutant (see Supplementary Fig. S3b). The quantitative RT-PCR analysis of replicon RNA in the cells treated with IFN- β clearly showed that the level of 4 β R replicon in cells expressing wild-type IFNAR1 was drastically decreased after IFN- β treatment, as was the level of 6M replicon in cells expressing wild-type IFNAR1 (see Supplementary Fig. S3c). In summary, we demonstrated that the IFNAR mutation found in 4 β R cells was a major determinant for a strongly IFN-resistant phenotype of 4 β R cells, suggesting that IFNAR mutations, which lead to the impairment of IFN signalling, convert HCV replicon-harboring cells from an IFN-sensitive phenotype to a highly IFN-resistant phenotype.

IFNAR1 and IFNAR2c belonging to the class II cytokine receptor superfamily are structurally conserved transmembrane receptors located on the cell surface (see Supplementary Fig. S4a, available in JGV Online). However, since both the IFNAR1 E107stop mutant and the IFNAR2c E159stop mutant found in 4 β R and 1 β R cells, respectively, were N-terminally truncated and probably soluble forms, these truncated proteins may not be functional as IFN receptors or may act as dominant-negative inhibitors, and will lead to the interception of IFN signalling (see Supplementary Fig. S4b). Thus, the cause of the IFN-resistant phenotype of 1 β R or 4 β R cells appeared to be the functional disruption of IFNAR. The present results suggest that the downstream JAK/STAT pathway is intact, at least in 4 β R cells.

Although for the most part we could clarify the mechanism underlying a highly IFN-resistant phenotype of HCV replicon-harboring cells, at least in the case of 4 β R cells, the mechanism underlying a partially IFN-resistant phenotype remains unclear, because IFNAR mutations were rare in the α R series. Since the expression levels of IFNAR, TYK2 and JAK1 were not decreased in the α R series, a functional deficiency of other cellular factor(s) involved in the IFN signalling may contribute to the acquisition of IFN resistance. Alternatively, certain HCV mutation(s) may account for the partially IFN-resistant phenotype of the α R series.

Since Machida *et al.* (2004a, b) recently reported that the frequency of genetic mutation was enhanced by HCV replication in *in vitro*-infected B cells and that the HCV core and NS3 were involved in the induction of a mutator phenotype mediated through the activation of inducible nitric oxide synthase, we cannot exclude the possibility that persistent HCV replication induces some irreversible genetic mutations. To clarify whether or not HCV acts as a mutagen for cellular factors, further study using an HCV RNA replication system (Ikeda *et al.*, 2002, 2005; Naka *et al.*, 2005) will also be necessary.

The HCV replicon-harboring cells including 1 β R and 4 β R, in which IFN signalling is impaired, used or obtained in the present study may be useful for future studies, not only of the mechanism(s) underlying the IFN resistance of the replicons but also of the functional characterization of IFN receptors. Furthermore, these replicon cells may also be useful for screening novel anti-HCV reagents that act by mechanisms unrelated to IFN signalling.

Acknowledgements

This work was supported by grants-in-aid for the third-term comprehensive 10-year strategy for cancer control, for research on hepatitis from the Ministry of Health, Labour and Welfare of Japan, and by the programme for promotion of fundamental studies in Health Sciences of the Pharmaceuticals and Medical Devices Agency (PMDA). We would like to thank T. Nakamura, H. Tawara, T. Maeta, A. Morishita and Y. Edamatsu for their helpful assistance in experiments.

References

- Akagi, T., Shishido, T., Murata, K. & Hanafusa, H. (2000). v-Crk activates the phosphoinositide 3-kinase/AKT pathway in transformation. *Proc Natl Acad Sci U S A* **97**, 7290–7295.
- Choo, Q. L., Kuo, G., Weiner, A. J., Overby, L. R., Bradley, D. W. & Houghton, M. (1989). Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* **244**, 359–362.
- Dansako, H., Naganuma, A., Nakamura, T., Ikeda, F., Nozaki, A. & Kato, N. (2003). Differential activation of interferon-inducible genes by hepatitis C virus core protein mediated by the interferon stimulated response element. *Virus Res* **97**, 17–30.
- Frese, M., Pietschmann, T., Moradpour, D., Haller, O. & Bartenschlager, R. (2001). Interferon- α inhibits hepatitis C virus subgenomic RNA replication by an MxA-independent pathway. *J Gen Virol* **82**, 723–733.
- Frese, M., Schwarzle, V., Barth, K., Krieger, N., Lohmann, V., Mihm, S., Haller, O. & Bartenschlager, R. (2002). Interferon- γ inhibits replication of subgenomic and genomic hepatitis C virus RNAs. *Hepatology* **35**, 694–703.
- Hadziyannis, S. J., Sette, H., Jr, Morgan, T. R. & 11 other authors (2004). Peginterferon- α 2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* **140**, 346–355.
- Hijikata, M., Mizushima, H., Tanji, Y., Komoda, Y., Hirowatari, Y., Akagi, T., Kato, N., Kimura, K. & Shimotohno, K. (1993). Proteolytic processing and membrane association of putative nonstructural proteins of hepatitis C virus. *Proc Natl Acad Sci U S A* **90**, 10773–10777.
- Ikeda, M., Yi, M., Li, K. & Lemon, S. M. (2002). Selectable subgenomic and genome-length dicistronic RNAs derived from an infectious molecular clone of the HCV-N strain of hepatitis C virus replicate efficiently in cultured Huh7 cells. *J Virol* **76**, 2997–3006.
- Ikeda, M., Abe, K., Dansako, H., Nakamura, T., Naka, K. & Kato, N. (2005). Efficient replication of a full-length hepatitis C virus genome, strain O, in cell culture, and development of a luciferase reporter system. *Biochem Biophys Res Commun* **329**, 1350–1359.
- Kato, N. (2001). Molecular virology of hepatitis C virus. *Acta Med Okayama* **55**, 133–159.
- Kato, N., Hijikata, M., Ootsuyama, Y., Nakagawa, M., Ohkoshi, S., Sugimura, T. & Shimotohno, K. (1990). Molecular cloning of the human hepatitis C virus genome from Japanese patients with non-A, non-B hepatitis. *Proc Natl Acad Sci U S A* **87**, 9524–9528.
- Kato, N., Sugiyama, K., Namba, K., Dansako, H., Nakamura, T., Takami, M., Naka, K., Nozaki, A. & Shimotohno, K. (2003). Establishment of a hepatitis C virus subgenomic replicon derived from human hepatocytes infected *in vitro*. *Biochem Biophys Res Commun* **306**, 756–766.
- Kato, N., Nakamura, T., Dansako, H., Namba, K., Abe, K., Nozaki, A., Naka, K., Ikeda, M. & Shimotohno, K. (2005). Genetic variation and dynamics of hepatitis C virus replicons in long-term cell culture. *J Gen Virol* **86**, 645–656.
- Kishine, H., Sugiyama, K., Hijikata, M. & 7 other authors (2002). Subgenomic replicon derived from a cell line infected with the hepatitis C virus. *Biochem Biophys Res Commun* **293**, 993–999.
- Kuo, G., Choo, Q. L., Alter, H. J. & 17 other authors (1989). An assay for circulating antibodies to a major etiologic virus of human non-A, non-B hepatitis. *Science* **244**, 362–364.
- Lohmann, V., Korner, F., Koch, J., Herian, U., Theilmann, L. & Bartenschlager, R. (1999). Replication of subgenomic hepatitis C virus RNAs in a hepatoma cell line. *Science* **285**, 110–113.

- Machida, K., Cheng, K. T.-N., Sung, V. M.-H., Lee, K. J., Levine, A. M. & Lai, M. M. (2004a). Hepatitis C virus infection activates the immunologic (type II) isoform of nitric oxide synthase and thereby enhances DNA damage and mutations of cellular genes. *J Virol* **78**, 8835–8843.
- Machida, K., Cheng, K. T.-N., Sung, V. M.-H., Shimodaira, S., Lindsay, K. L., Levine, A. M., Lai, M.-Y. & Lai, M. M. (2004b). Hepatitis C virus induces a mutator phenotype: enhanced mutations of immunoglobulin and protooncogenes. *Proc Natl Acad Sci U S A* **101**, 4262–4267.
- Naganuma, A., Nozaki, A., Tanaka, T., Sugiyama, K., Takagi, H., Mori, M., Shimotohno, K. & Kato, N. (2000). Activation of the interferon-inducible 2'-5'-oligoadenylate synthetase gene by hepatitis C virus core protein. *J Virol* **74**, 8744–8750.
- Naganuma, A., Dansako, H., Nakamura, T., Nozaki, A. & Kato, N. (2004). Promotion of microsatellite instability by hepatitis C virus core protein in human non-neoplastic hepatocyte cells. *Cancer Res* **64**, 1307–1314.
- Naka, K., Ikeda, M., Abe, K., Dansako, H. & Kato, N. (2005). Mizoribine inhibits hepatitis C virus RNA replication: effect of combination with interferon- α . *Biochem Biophys Res Commun* **330**, 871–879.
- Namba, K., Naka, K., Dansako, H., Nozaki, A., Ikeda, M., Shiratori, Y., Shimotohno, K. & Kato, N. (2004). Establishment of hepatitis C virus replicon cell lines possessing interferon-resistant phenotype. *Biochem Biophys Res Commun* **323**, 299–309.
- Nozaki, A., Ikeda, M., Naganuma, A., Nakamura, T., Inudoh, M., Tanaka, K. & Kato, N. (2003). Identification of a lactoferrin-derived peptide possessing binding activity to hepatitis C virus E2 envelope protein. *J Biol Chem* **278**, 10162–10173.
- Pawlotsky, J. M. (2000). Hepatitis C virus resistance to antiviral therapy. *Hepatology* **32**, 889–896.
- Pfeffer, L. M., Mullersman, J. E., Pfeffer, S. R., Murti, A., Shi, W. & Yang, C. H. (1997). STAT3 as an adapter to couple phosphatidylinositol 3-kinase to the IFNAR1 chain of the type I interferon receptor. *Science* **276**, 1418–1420.
- Saito, I., Miyamura, T., Ohbayashi, A. & 10 other authors (1990). Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci U S A* **87**, 6547–6549.
- Thomas, D. L. (2000). Hepatitis C epidemiology. *Curr Top Microbiol Immunol* **242**, 25–41.



Mizoribine inhibits hepatitis C virus RNA replication: Effect of combination with interferon- α

Kazuhito Naka, Masanori Ikeda, Ken-ichi Abe, Hiromichi Dansako, Nobuyuki Kato*

Department of Molecular Biology, Okayama University Graduate School of Medicine and Dentistry, 2-5-1 Shikata-cho, Okayama 700-8558, Japan

Received 28 February 2005

Available online 18 March 2005

Abstract

Interferon (IFN)- α monotherapy, as well as the more effective combination therapy of IFN- α and ribavirin, are currently used for patients with chronic hepatitis C caused by hepatitis C virus (HCV) infection, although the mechanisms of the antiviral effects of these reagents on HCV remain ambiguous, and side effects such as anemia due to the administration of ribavirin present a problem for patients who are advanced in years. Using a recently developed reporter assay system in which genome-length dicistronic HCV RNA encoding *Renilla* luciferase gene was found to replicate efficiently, we found that mizoribine, an imidazole nucleoside, inhibited HCV RNA replication. The anti-HCV activity of mizoribine (IC₅₀: approximately 100 μ M) was similar to that of ribavirin. Using this genome-length HCV RNA replication monitor system, we were the first to demonstrate that the combination of IFN- α and ribavirin exhibited more effective anti-HCV activity than the use of IFN- α alone. Moreover, we found that the anti-HCV activity of mizoribine in co-treatment with IFN- α was at least equivalent to that of ribavirin. This effect was apparent in the presence of at least 5 μ M mizoribine. Since mizoribine is currently used in several clinical applications and has not been associated with severe side effects, mizoribine is considered to be of potential use as a new anti-HCV reagent in combination with IFN- α .

© 2005 Elsevier Inc. All rights reserved.

Keywords: Hepatitis C virus; HuH-7; HCV RNA replication system; Ribavirin; Mizoribine; Interferon

Hepatitis C virus (HCV) is the major causative agent of chronic hepatitis (CH) [1,2], which progresses to liver cirrhosis and hepatocellular carcinoma [3,4]. Since at least 170 million people are currently infected with HCV worldwide, this infection is a global health problem [5]. HCV is an enveloped RNA virus belonging to *Flaviviridae*, the genome of which consists of a positive-stranded 9.6-kilobase (kb) RNA encoding an approximately 3000 amino acid polyprotein precursor [6,7]. This precursor protein is cleaved by the host and viral proteinases to generate at least 10 proteins in the following order: core, envelope 1 (E1), E2, p7, nonstructural protein 2 (NS2), NS3, NS4A, NS4B, NS5A, and NS5B [8–10].

To date, interferon (IFN)- α (majority) and IFN- β (minority) are used as effective anti-HCV reagents in clinical therapy for patients with CH C; however, the effectiveness of IFN is limited to about 30% of the reported cases [11]. In 1998, combined treatment with IFN- α and ribavirin, a nucleoside analogue, has been shown to be more effective (although the effectiveness remains at less than 50%) than treatment with IFN alone [12,13]; nonetheless, it has been shown that ribavirin alone does not induce a decrease in HCV levels in patients with CH C. Furthermore, it has been reported that the combination of pegylated IFN with ribavirin led to significant improvements in terms of a sustained virological response, when compared to standard IFN and ribavirin combination therapy [14]. However, a sustained virological response is still not induced in approximately half of patients treated with these reagents.

* Corresponding author. Fax: +81 86 235 7392.

E-mail address: nkato@md.okayama-u.ac.jp (N. Kato).