

References

- [1] Alter MJ, Kruszon-Moran D, Nainan OV, et al. The prevalence of hepatitis C virus infection in the United States 1998 through 1994. *N Engl J Med* 1999;341:556–62.
- [2] Wong JB, McQuillan GM, McHutchison JG, Poynard T. Estimating future hepatitis C morbidity, mortality, and costs in the United States. *Am J Public Health* 2000;90:1562–9.
- [3] Diepolder HM, Zachoval R, Hoffmann R, et al. Possible mechanism involving T-lymphocyte response to non-structural protein 3 in viral clearance in acute hepatitis C virus infection. *Lancet* 1995;346:1006–7.
- [4] Missale G, Bertoni R, Lamonaca V, et al. Different clinical behaviors of acute hepatitis C virus infection are associated with different vigor of the anti-viral cell-mediated immune response. *J Clin Invest* 1996;98:706–14.
- [5] Watanabe H, Saito T, Shinzawa H, et al. Spontaneous elimination of serum hepatitis C virus (HCV) RNA in chronic HCV carriers: a population-based cohort study. *J Med Virol* 2003;71:56–61.
- [6] Fujisawa T, Komatsu H, Inui Y, et al. Spontaneous remission of chronic hepatitis C in children. *Eur J Pediatr* 1997;156:773–6.
- [7] Hattori Y, Orito E, Ohno T, et al. Loss of hepatitis C virus RNA after parturition in female patients with chronic HCV infection. *J Med Virol* 2003;71:205–11.
- [8] Fukuizumi S, Sata M, Suzuki H, Kumashiro R, Tanikawa K. Natural disappearance of serum HCV RNA: prospective study in a hyperendemic area. *Hepatol Res* 1997;9:144–51.
- [9] Yokosuka O, Kojima H, Imazeki F, et al. Spontaneous negatization of serum hepatitis C virus RNA is a rare event in type C chronic liver diseases: analysis of HCV RNA in 320 patients who were followed for more than 3 years. *J Hepatol* 1999;31:394–9.
- [10] Yoshikawa M, Morimoto Y, Shiroy A, Yoshiji H, Kuriyama S, Fukui H. Spontaneous elimination of serum HCV-RNA after total gastrectomy for early gastric cancer in a patient with chronic hepatitis C. *Am J Gastroenterol* 2001;96:922–3.
- [11] Kashiwakuma T, Hasegawa A, Kajita T, et al. Detection of hepatitis C virus specific core protein in serum of patients by a sensitive fluorescence enzyme immunoassay (FEIA). *J Immunol Methods* 1996;190:78–9.
- [12] Tanaka T, Lau JYN, Mizokami M, et al. Simple fluorescent EIA for detection and quantification of hepatitis C viremia. *J Hepatol* 1995;23:742–5.
- [13] Hayashi K, Hasuike S, Kusumoto K, et al. Usefulness of a new immuno-radiometric assay to detect hepatitis C core antigen in a community-based population. *J Viral Hepat* 2005;12:106–10.
- [14] Tanaka E, Kiyosawa K, Matsumoto A, et al. Serum levels of hepatitis C virus core protein in patients with chronic hepatitis C treated with interferon alfa. *Hepatology* 1996;23:1330–3.
- [15] Ono E, Shiratori Y, Okudaira T, et al. Platelet counts as predictors of degree of fibrosis in chronic hepatitis C virus infection. *Am J Gastroenterol* 2001;96:3142–6.
- [16] Murawaki Y, Ikuta Y, Koda M, Yamada S, Kawasaki H. Comparison of serum 7S fragment of type IV collagen and serum central triple-helix of type IV collagen for assessment of liver fibrosis in patients with chronic viral liver disease. *J Hepatol* 1996;24:148–54.
- [17] Ohkoshi S, Tawaraya H, Kuwana H, et al. A retrospective study of hepatitis C virus carriers in a local endemic town in Japan. A possible presence of asymptomatic carrier. *Dig Dis Sci* 1995;40:465–71.
- [18] Tanaka E, Kiyosawa K, Matsushima T, et al. Epidemiology of genotypes of hepatitis C virus in Japanese patients with type C chronic liver diseases: a multi-institution analysis. *J Gastroenterol Hepatol* 1995;10:538–45.
- [19] Mizoguchi N, Mizokami M, Orito E, Shibata H, Shibata H. Serologically defined genotypes of hepatitis C virus among Japanese patients with chronic hepatitis C. *J Virol Methods* 1996;58:71–9.
- [20] Inoue G, Horiike N, Michitaka K, Onji M. Hepatitis C virus clearance is prominent in women in an endemic area. *J Gastroenterol Hepatol* 2000;15:1054–8.