



の、経口抗ウイルス薬の発癌抑制効果を明らかにしたとしている。


Jang JWら⁵⁾は、肝動脈化学療法を施行中のB型肝炎ウイルス関連肝癌に対してラミブジン投与の無作為化比較試験を行った。彼らが本来行った治験の目的は、化学療法中に起こるB型肝炎ウイルスの再増殖を事前に抑え込むことにより有害事象を予防することにあった。73例の連続肝癌症例に対する肝癌治療としてエピルピシン50mg/m²＋シスプラチン60mg/m²＋リピオドール動注を毎月行い、これらの症例をラミブジン治療群(100mg内服)と無治療群に無作為に割り付けした。観察期間内に無治療群では11例(29.7%)がB型肝炎ウイルスの再増殖をきたしたのに対し、ラミブジン治療群の再燃は1例(2.8%)のみで、予防的ラミブジン治療により肝炎ウイルス再増殖が有意に抑制できた(p=0.002)。さらに、無治療群では肝炎全般(p=0.021)、高度の肝炎(p=0.035)ともラミブジン治療群より高率であった。多変量解析では

Chemolipiodolization治療を行ったB型肝炎関連肝癌において、HBV DNA量が10⁴コピー/mL以上であることが唯一のウイルス再増殖に寄与する要因であった。本研究では、ラミブジン投与により肝炎抑制・生存率延長に役立つとしているが、肝癌の無再発生存率、全生存率に関しては言及されていない。発癌に直結するHBV DNAの要因を抑制していることから、長期経過観察が可能であれば(肝癌の短期再発例が多くなれば)再発を含めた予後に関しても期待できるものと思われる。

文献

1. B型肝炎関連肝癌根治切除後のインターフェロンαは再発を遅らせ全体生存率を改善する:無作為化比較試験
Sun HC et al: Postoperative interferon α treatment postponed recurrence and improved overall survival in patients after curative resection of HBV-related hepatocellular carcinoma: a randomized clinical trial. *J Cancer Res Clin Oncol* 132(7): 458-465, 2006
2. B型肝炎関連肝癌発癌とその予防
Ikeda K et al: Hepatitis B virus-related hepatocellular carcinogenesis and its prevention. *Intervirol* 48(1): 29-38, 2005
3. 慢性B型肝炎進行例に対するラミブジンの効果
Liaw YF et al: Cirrhosis Asian Lamivudine Multicentre Study Group: Lamivudine for patients with chronic hepatitis B and advanced liver disease. *N Engl J Med* 351(15): 1521-1531, 2004
4. 慢性B型肝炎におけるラミブジンの肝癌発癌抑制効果。2795例の多施設レトロスペクティブ研究
Matsumoto A et al: Efficacy of Lamivudine for preventing hepatocellular carcinoma in chronic hepatitis B: A multicenter retrospective study of 2795 patients. *Hepatology Research* 32(3): 173-184, 2005
5. 経動脈的chemo-lipiodolization施行B型肝炎関連肝癌症例における予防的ラミブジン投与の無作為化比較試験
Jang JW et al: A randomized controlled study of preemptive lamivudine in patients receiving transarterial chemo-lipiodolization. *Hepatology* 43(2): 233-240, 2006

図1 インターフェロンによるB型肝炎ウイルス関連肝癌の再発予防



文献1をもとに改変引用

Anticarcinogenic Impact of Interferon on Patients with Chronic Hepatitis C: A Large-Scale Long-Term Study in a Single Center

Kenji Ikeda Yasuji Arase Satoshi Saitoh Masahiro Kobayashi
Takashi Someya Tetsuya Hosaka Hitomi Sezaki Norio Akuta
Fumitaka Suzuki Yoshiyuki Suzuki Hiromitsu Kumada

Department of Gastroenterology, Toranomon Hospital, and Okinaka Memorial Institute for Medical Research, Tokyo, Japan

Key Words

Cirrhosis · Fibrosis · Hepatitis C virus · Hepatocellular carcinoma · Interferon

Abstract

Background: The anticarcinogenic capacity of interferon (IFN) was assessed in a cohort of Japanese patients with chronic hepatitis C en masse. **Patients and Methods:** The rate of hepatocarcinogenesis was analyzed in 2,166 patients with chronic hepatitis C, of whom 1,654 had received IFN therapy while 512 had not. **Results:** Crude rates of hepatocarcinogenesis in treated and untreated patients were 2.6 and 4.6% at the end of the 5th year, 5.8 and 12.7% at the 10th year and 13.9 and 23.9% at the 15th year (after completion of IFN therapy for those treated) ($p < 0.001$). IFN decreased the hazard ratio of carcinogenesis to 0.42 ($p < 0.001$) in multivariate analysis with adjustments for significant covariates including fibrotic stage, γ -glutamyl transpeptidase level, gender, platelet count and age. Among the 1,654 patients treated with IFN, 606 (36.6%) achieved persistent loss of hepatitis C virus (HCV) RNA and an additional 266 (16.1%) gained normal levels of alanine aminotransferase without loss of HCV RNA for 6 months or longer after the completion of IFN therapy. Cumulative rates of hepatocarcinogenesis in sustained virological responders and biochemical responders were 1.4 and 2.0% at the end of the 5th year,

1.9 and 3.6% at the 10th year and 1.9 and 7.5% at the 15th year, respectively. The hazard ratio of sustained virological response was 0.10 ($p < 0.001$), and that of biochemical response was 0.12 ($p < 0.001$). Normalization of aminotransferase levels after IFN therapy without loss of serum HCV RNA decreased hepatocarcinogenesis. **Conclusion:** IFN significantly decreased the rate of hepatocarcinogenesis in patients with chronic hepatitis C as a whole in Japan, even in those who fail to clear HCV RNA from serum.

Copyright © 2006 S. Karger AG, Basel

Introduction

In most developed countries, hepatitis B virus (HBV) and hepatitis C virus (HCV) infections account for the great majority of hepatocellular carcinoma (HCC), with incidence rates dependent on the regional prevalence of these hepatitis viruses. HCV-associated HCC typically develops through a sequence of events that progress from chronic inflammation through fibrosis and cirrhosis accompanying dysplasia and ultimately to HCC. In our previous cohort study on Japanese patients with HCV-related cirrhosis [1], cumulative rates of developing HCC at 5, 10 and 15 years were 21.5, 53.2 and 75.2%, respectively. According to our observations of untreated patients with chronic hepatitis C [2], rates of hepatocarcino-

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 2006 S. Karger AG, Basel
0300-5526/06/0492-0082\$23.50/0

Accessible online at:
www.karger.com/int

Kenji Ikeda, MD
Department of Gastroenterology, Toranomon Hospital
Toranomon 2-2-2, Minato-ku
Tokyo 105-8470 (Japan)
Tel. +81 44 877 5111, Fax +81 44 860 1623. E-Mail ikedakenji@tora.email.ne.jp

genesis at 5, 10 and 15 years were estimated to be 4.8, 13.6 and 26.0%, respectively. The life expectancy of patients with HCV-related cirrhosis is largely influenced by the development of HCC in the clinical course. As the efficacy of radically curative therapies for HCC remains limited at best, and since a severe organ shortage does not provide with sufficient chances for liver transplantation, the prevention of HCC in patients with chronic liver disease is of great importance at the present.

Interferon (IFN) is effective in eliminating HCV and reducing serum levels of alanine aminotransferase (ALT) in some patients with chronic hepatitis C [3–6]. Reduced incidence of HCC in HCV-associated cirrhosis by IFN has been reported by many investigators including ourselves [7–14]; only a few studies have failed to find its benefit [15, 16]. However, many published studies had shortcomings in the study design, in terms of pooling patients who received IFN in diverse regimens, relatively short periods of follow-up despite a long incubation period of HCC, large numbers of dropouts and retrospective studies with historical controls. Moreover, almost all studies evaluated the activity of IFN to prevent HCC by comparing responders and nonresponders to the treatment. Due to difficulties in studying patients with chronic hepatitis C, a number of nonrandomized studies examined the effect of IFN on the incidence of hepatocarcinogenesis [17–20]. With invariable limitations in study design and interpretation of the results, these studies have disclosed useful information as regards the treatment of patients with chronic HCV infection.

In order to evaluate whether IFN can reduce the rate of carcinogenesis in patients with chronic hepatitis C, we compared 1,654 patients with IFN therapy with 512 patients without treatment in a single clinical center, who were adjusted for background features by the multivariate analysis. Therefore, the principal aims of our study were to show the role of IFN in preventing HCC in chronic hepatitis type C en masse and to establish the extent to which IFN decreases the rate of carcinogenesis as a sequel to chronic hepatitis C in a society.

Patients and Methods

Study Population

A total of 2,166 patients with chronic hepatitis were examined, whose initial sera tested negative for hepatitis B surface antigen by radioimmunoassay (Ausria, Dainabot, Tokyo, Japan) and positive for anti-HCV by the second-generation enzyme-linked immunosorbent assay (Dainabot); anti-HCV was tested in sera that had been stored frozen at -80°C . They included 1,421 men and 745

women aged 14–78 with a median of 50 years. They were all diagnosed with chronic hepatitis by liver biopsy with or without peritoneoscopy between 1970 and 2000 at the Department of Gastroenterology in Toranomon Hospital, Tokyo, Japan. Patients who had possibly developed HCC already at the time of diagnosis of hepatitis were strictly excluded from the study. In order to exclusively investigate hepatocarcinogenesis in HCV-related cirrhosis, patients coinfecting with HBV were excluded.

Among the 2,166 patients with HCV-related hepatitis, 1,654 (76.4%) received IFN therapy, mostly since 1987 when IFN was available in Japan; new antivirals or anticarcinogenic treatments of viral cirrhosis, except for IFN, were not introduced in 1987 or thereafter in Japan. The remaining 512 patients did not receive IFN or any other antiviral therapies. This is a retrospective cohort study with historical controls composed of patients before 1987 and those who refused or could not receive IFN for various reasons since 1987.

Background and Laboratory Findings

Table 1 shows demographic profiles and laboratory data for the 1,654 patients treated with IFN and the 512 without receiving IFN since they were diagnosed with chronic hepatitis. There were more males, with a median age 3 years lower in treated than in nontreated patients. There were 299 treated patients (18.1%) with a history of alcohol intake ≥ 500 ml until the diagnosis of chronic hepatitis (corresponding to daily consumption of 3,000 ml of beer or 300 ml of whiskey for 20 years) and 113 (22.1%) untreated patients ($p < 0.001$). Because IFN was introduced to our hospital in 1987, the observation period was significantly shorter in the treated than in untreated patients (median 10.4 vs. 12.3 years; $p < 0.0001$).

Although all patients tested positive for HCV RNA during their clinical courses, tests for the concentration of HCV RNA in the initial serum was possible in 1,863 (86.5%) patients. HCV genotypes were analyzed by the serological typing method with a commercial kit (Kokusai Diagnostic Corporation, Kobe, Japan) in which the serological group 1 represented genotypes 1a and 1b, and group 2 stood for 2a and 2b genotypes. HCV in the serological group 2 was significantly more frequent in patients with IFN treatment than in those without. Concentration of HCV RNA was determined in the initial sera from 1,873 (86.5%) patients by the competitive polymerase chain reaction (PCR) method with the HCV probe assay (Chiron Corp., Emeryville, Calif., USA) or by PCR with Amplicor HCV Monitor kits (Roche Diagnostics Japan Co., Tokyo, Japan). High concentration of HCV ($\geq 10^6$ copies/ml by the competitive PCR or $\geq 10^6$ equivalents/ml by the HCV probe assay) was significantly more frequent in untreated than in treated patients ($p < 0.0001$). The stage of hepatic fibrosis was not different between the two groups.

Interferon Treatment and Judgment of the Effect

A total of 1,654 patients underwent IFN therapy in one or more treatment courses: 1,358 patients (82.1%) received IFN once, 240 patients (14.5%) twice, and the remaining 56 patients (3.4%) three times or more. Initial treatment was performed with natural or recombinant IFN- α ($n = 1,238$), natural IFN- β ($n = 386$) or both ($n = 30$). Regimens of IFN were variable: 926 (56.0%) patients received IFN 6–9 million units (MU) daily for 8 weeks, followed by 2 or 3 times per week for 16 weeks; 329 (20.0%) received IFN 6–9 MU daily for 2–4 weeks, followed by 3 times per week for 20–22 weeks; 185 (11.2%) underwent a short-course therapy with IFN

Table 1. Patient profiles and laboratory data at the diagnosis of chronic hepatitis

Factors	Interferon therapy		p value
	yes (n = 1,654)	no (n = 512)	
Male	1,110 (67.1%)	311 (60.7%)	0.024
Age, years	50 (16–72)	53 (21–78)	<0.001
History of transfusion	607 (36.7%)	229 (44.7%)	0.001
Family member with liver disease	426 (25.8%)	140 (27.3%)	0.47
Alcohol intake \geq 500 kg	299 (18.1%)	113 (22.1%)	0.044
Observation period, year	10.4 (0.1–33.6)	12.3 (0.1–33.6)	<0.001
Laboratory data			
ALT, IU/l	63 (4–1,266)	67 (4–704)	0.098
AST, IU/l	106 (9–1,660)	96 (12–832)	0.0001
γ -GTP, IU/ml	62 (6–1,118)	70 (3–850)	0.39
Platelet counts, $\times 1,000/\text{mm}^3$	169 (27–433)	165 (35–560)	0.091
ICG R ₁₅ , %	14 (1–90)	16 (1–95)	0.003
AFP, ng/ml	4 (1–90)	5 (1–1,180)	0.42
HCV serological group			
Group 1, genotypes 1a/1b	1,021 (66.1%)	259 (81.4%)	<0.0001
Group 2, genotypes 2a/2b	488 (31.6%)	48 (15.1%)	
Undetermined	36 (2.3%)	11 (3.5%)	
HCV RNA concentration			
High ^a	937 (58.4%)	191 (71.3%)	<0.0001
Low ^b	668 (41.6%)	77 (28.7%)	
Histological stage of hepatitis			
F1, slight fibrosis	1,029 (62.2%)	298 (58.2%)	0.10
F2/F3, moderate/severe fibrosis	625 (37.8%)	214 (41.6%)	

AST = Aspartate aminotransferase; AFP = α -fetoprotein; ICG R₁₅ = retention of indocyanine green at 15 min.

^a HCV RNA concentration $\geq 10^6$ copies/ml by the competitive PCR or $\geq 10^6$ equivalents/ml by the HCV probe assay.

^b HCV RNA concentrations less than high concentrations.

daily for 4–8 weeks; 128 (7.7%) were administered with intermittent IFN 3 times per week for 24 weeks; 72 (4.4%) had a prolonged course of IFN for 8–36 months; 8 (0.5%) received IFN- β 6 MU daily for 6–18 months, and the remaining 6 (0.4%) were given IFN- α combined with IFN- β for 4 months. The median dose of 624 MU was administered during the median period of 24 weeks. IFN for 24 weeks or longer was given to 83.2% of the patients. IFN therapy was usually initiated within a few months after the diagnosis of chronic hepatitis, and all patients were started on it within 12 months. The median interval between liver biopsy and initiation of IFN was 9 days.

Almost all the patients given IFN showed varied degrees of fever, chills, myalgia, headache and general malaise after the first injection. Most patients developed leukocytopenia and thrombocytopenia in various degrees. A significant thrombocytopenia $\leq 40,000/\text{mm}^3$ required a reduction of the IFN dose in 39 patients. IFN therapy was discontinued due to psychosis in 35 patients and ophthalmological symptoms in 12 patients. None of the patients developed decompensated liver disease with ascites, encephalopathy, jaundice or variceal bleeding. Although only 88 (5.3%) patients could not continue injection with IFN, studies for carcinogenesis were analyzed on the intention-to-treat basis.

The efficacy of IFN was judged by the clearance of HCV RNA from serum and ALT levels 12 months after the completion of treatment. Sustained virological response (SVR) was defined as persistent disappearance of HCV RNA after therapy, biochemical response (BR) as normal ALT levels without elimination of HCV RNA for at least 6 months after therapy, and no response (NR) as persistently elevated or transiently normalized ALT levels without loss of HCV RNA lasting for less than 6 months.

Follow-Up of Patients and Diagnosis of HCC

Patients were followed up monthly after diagnosis of chronic hepatitis in our outpatient clinic and monitored for hematological, biochemical and virological parameters. With their admission, during and after the treatment with IFN, weekly or biweekly follow-up was performed in almost all patients who received IFN. Imaging diagnosis was made once or twice per year in the majority of patients with ultrasonography or computed tomography. Angiography was performed only when HCC was highly suspected on imaging by ultrasonography or computed tomography.

When angiography pictured a characteristic hypervascular nodule specific for HCC in patients, histological confirmation was not required in the majority of them. Microscopic examinations of liv-




Fig. 1. Crude rates of hepatocarcinogenesis in patients treated with IFN and those untreated. The carcinogenesis rate was significantly lower in treated than in untreated patients (log-rank test, $p < 0.0001$).

er tissues obtained by a fine-needle biopsy were performed in 14 patients whose angiogram could not portray a typical image of HCC. There were 89 patients in whom HCC was confirmed histologically on liver specimens obtained at surgery or autopsy. Detection of serological tumor markers and increase with time were also taken into account in the diagnosis of HCC.

There were 223 (10.3%) patients lost to follow-up, including 164 (9.9%) treated and 59 (11.5%) untreated. Rates of annual dropouts in treated and untreated patients were 0.95 and 0.93%, respectively. In 9 patients, the response to IFN was judged by information on aminotransferase levels determined in other clinics and by persistent HCV RNA, as well as aminotransferase levels at 6 months after the completion of therapy in an additional 3 patients. Therefore, the response to IFN could be judged in all patients including the 12 who were lost to our follow-up early. Since the eventual outcome with respect to the development of HCC was not confirmed in these patients, their data were censored in statistical analyses [21]. Deaths unrelated to liver disease were censored and withdrawn from the analysis. The date of the last follow-up in this study was May 1, 2004, and the median observation period of studied patients was 10.7 years, with a range of 0.1–33.6 years.

Statistical Analysis

Nonparametric Mann-Whitney U test and χ^2 test were used for analysis of background characteristics of patients. The rate of HCC development was calculated by the Kaplan-Meier method [22]; it was based on the duration between diagnosis of chronic hepatitis by liver biopsy and detection of HCC. Differences in slopes of carcinogenesis curves were evaluated by the log-rank test. To gain a robust statistical power for the anticarcinogenic activity of IFN, observation of treated patients was initiated at the commencement of IFN therapy, in lieu of the diagnosis of chronic hepatitis. Independent factors associated with the development of HCC were studied using the stepwise Cox regression analysis [23]. The follow-

ing 18 variables were analyzed for potential covariates in hepatocarcinogenesis at the time when hepatitis was diagnosed: age, sex, total alcohol intake, family history of liver disease, history of blood transfusion, stage of hepatic fibrosis, aspartic aminotransferase, ALT, albumin, bilirubin, globulin, γ -glutamyl transpeptidase (γ -GTP), platelet count, retention of indocyanine green at 15 min, serological grouping of HCV, HCV RNA level and IFN treatment.

Although continuous variables without conversion of data were evaluated in multivariate analyses, several variables were transformed into categorical data consisting of two or three ordinal numbers in calculating hazard ratios. All factors found to be marginally associated with hepatocarcinogenesis with p values < 0.15 were tested by the multivariate Cox proportional hazard model. All analyses of data were performed with the computer program SPSS version 11 [24], and a p value < 0.05 was considered significant.

Results

Response to IFN

Response to IFN was judged 12 months after the completion of therapy by both HCV RNA and serial ALT readings. Among the 1,654 patients with IFN treatment, SVR (elimination of HCV RNA) was achieved by 606 (36.6%), BR (ALT normalized for at least 6 months without clearance of HCV RNA from serum) in 266 (16.1%) and NR (elevated or transiently decreased ALT levels without loss of serum HCV RNA) in 782 (47.3%).

Crude Rates of Hepatocarcinogenesis

During the median observation period of 10.7 years, HCC developed in 199 of the 2,166 (9.2%) patients, including 96 of the 1,654 (5.8%) patients treated with IFN and 103 of the 512 (20.1%) patients without IFN (fig. 1). Among the 199 patients with HCC, 140 (70.4%) imaged a typical hypervascular stain on angiography and dynamic computed tomography, while 59 failed to exhibit tumor stains on angiography. HCC in these 59 patients was confirmed histologically on liver specimens obtained at surgery or by fine-needle biopsy.

Crude rates of hepatocarcinogenesis in patients treated with IFN and those untreated were 1.3 and 1.8% at the end of the 3rd year (after the completion of therapy), 2.6 and 4.6% at the end of the 5th year, 5.8 and 12.7% at the 10th year and 13.9 and 23.9% at the 15th year, respectively (fig. 1). The carcinogenesis rate was significantly lower in patients treated with IFN than in untreated patients (log-rank test, $p < 0.0001$).

Impact of IFN on Hepatocarcinogenesis

During the observation period, HCC developed in 96 of the 1,654 (5.8%) patients treated with IFN, including

11 patients (1.8%) with SVR, 10 (3.8%) with BR and 75 (9.6%) with NR to IFN. Rates of hepatocarcinogenesis in patients with SVR, BR and NR were 0.7, 0.8 and 2.0% at the end of the 3rd year, 1.4, 2.0 and 3.8% at the 5th year, 1.6, 2.9 and 6.5% at the 7th year, 1.9, 3.6 and 9.6% at the 10th year and 1.9, 7.5 and 27.6% at the end of 15th year (fig. 2). Hepatocarcinogenesis was significantly less frequent in patients with SVR or BR than in patients with NR and those untreated (log-rank test, $p < 0.0001$).

Factors Influencing Hepatocarcinogenesis

Univariate analysis identified 9 factors significantly associated with carcinogenesis. They were fibrotic stage ($p < 0.001$), age ($p < 0.001$), α -fetoprotein ($p < 0.001$), aspartic aminotransferase ($p = 0.001$), retention of indocyanine green at 15 min ($p = 0.002$), total alcohol intake ($p = 0.002$), γ -GTP ($p = 0.005$) and HCV serotype ($p = 0.045$). IFN therapy ($p = 0.064$), histological activity of hepatitis ($p = 0.069$) and ALT ($p = 0.70$) were marginally associated with carcinogenesis.

In order to prove the role of IFN on carcinogenesis in patients with chronic hepatitis type C en masse, multivariate analysis was performed by non-time-dependent proportional hazard analysis. Fibrotic stage, γ -GTP, gender, IFN therapy, platelet count and age independently influenced the development of HCC in the cohort (table 2). Advanced liver fibrosis in F2/F3 stages imposed a higher risk for carcinogenesis with a hazard ratio of 8.68, 95% confidence interval (CI) 5.08–14.81, compared with the F1 stage. Similarly, higher γ -GTP levels (hazard ratio 2.64), male sex (2.38), low platelet count (2.22) and older age (1.90) posed higher carcinogenesis risks. After adjusting background clinical biases between treated and untreated patients for the 5 significant covariates identified in the multivariate analysis, IFN therapy significantly decreased the hepatocarcinogenesis rate in the entire patients with chronic hepatitis C with a hazard ratio of 0.42 (95% CI 0.29–0.61) in comparison with untreated patients.

Based on the multivariate analysis, curves of carcinogenesis rates were theoretically illustrated in treated and untreated patients with the average histological stage, average γ -GTP value, average ratio of male to female, average platelet count and average age (fig. 3).

Hazard of Hepatocarcinogenesis Stratified by the Response to IFN

Since the carcinogenesis rate in patients with SVR or BR was significantly lower than that of patients with NR or untreated patients by the product limit method, a mul-




Fig. 2. Rates of hepatocarcinogenesis in patients with SVR, BR and NR to IFN. The rate in patients with NR (persistently elevated ALT or transiently normalized ALT for less than 6 months) was significantly higher than that in patients with SVR or BR.

Table 2. Factors associated with hepatocarcinogenesis in patients with chronic hepatitis C^a

Factors	HR	95% CI	p value
Fibrosis stage			
F1	1		
F2–F3	8.68	(5.08–14.81)	<0.001
γ -GTP, IU/ml			
<50	1		
≥ 50	2.64	(1.58–4.42)	<0.001
Gender			
Women	1		
Men	2.38	(1.56–3.70)	<0.001
IFN therapy			
No	1		
Yes	0.42	(0.29–0.61)	<0.001
Platelet count, $\times 10^3/\text{mm}^3$			
≥ 100	1		
<100	2.22	(1.47–3.44)	<0.001
Age, years			
<50	1		
≥ 50	1.90	(1.27–2.85)	0.002

HR = Hazard ratio.

^a Evaluated by the Cox proportional hazard analysis.




Fig. 3. Theoretical curves of hepatocarcinogenesis in patients treated with IFN and those untreated who have the average histological stage, average γ -GTP value, average ratio of male to female, average platelet count and average age. They are based on the analysis of 1,654 patients treated with IFN and 512 untreated patients.

Table 3. Factors associated with hepatocarcinogenesis in patients with chronic hepatitis C who had distinct responses to IFN therapy^a

Factors	HR	95% CI	p value
Fibrosis stage			
F1	1		
F2-F3	9.90	(4.19-23.40)	<0.001
Gender			
Women	1		
Men	3.44	(1.89-6.25)	<0.001
γ -GTP, IU/ml			
<50	1		
\geq 50	2.68	(1.30-5.54)	0.008
Age, years			
<50	1		
\geq 50	2.56	(1.50-4.38)	0.001
AFP, ng/ml			
<20	1		
\geq 20	2.32	(1.34-4.02)	0.003
Platelet count, $\times 10^3/\text{mm}^3$			
\geq 100	1		
<100	2.09	(1.14-3.75)	0.013
Response to IFN			
Without IFN	1		
NR	0.57	(0.13-2.56)	0.46
BR	0.12	(0.04-0.35)	<0.001
SVR	0.10	(0.03-0.30)	<0.001

HR = Hazard ratio; AFP = α -fetoprotein.

^a Evaluated by the Cox proportional hazard analysis.

tivariate analysis was performed taking into account the response to IFN. Hazard ratios of patients with SVR and BR to IFN therapy were 0.10 (95% CI 0.03-0.30, $p < 0.001$) and 0.12 (95% CI 0.04-0.35, $p < 0.001$), respectively, in comparison with that of untreated patients, when the other 5 factors served as significant covariates (table 3). The hazard ratio of NR at 0.57 (95% CI 0.13-2.56) was less than 1, but fell short of making a significant difference against untreated patients.

Mortality and Causes of Death

During the observation period, 116 of the 2,166 (5.4%) patients died, including 52 of the 1,654 (3.1%) subjects treated with IFN and 64 of the 512 (12.5%) subjects without IFN. Estimated survival rates in the treated and untreated patients were 99.3 and 98.3% at 5 years, 97.8 and 96.0% at 10 years and 93.8 and 86.9% at 15 years, respectively. The survival rate of treated patients was significantly higher than that of untreated patients (log-rank test, $p < 0.0001$).

Discussion

Based on our epidemiological data obtained by long-term observations of patients with chronic hepatitis [2] and patients with cirrhosis [1], the life expectancy of patients with HCV-related chronic liver disease heavily depends on the development of HCC. The possibility of eventually developing HCC in patients with HCV infection and cirrhosis is staggeringly high at 75% [1]. Theoretically, the treatment of chronic HCV infection with IFN can prevent the development of HCC. From the ethical point of view, a prospective randomized trial with control untreated patients is not to be allowed at present when IFN has become the standard radical therapy for chronic hepatitis C; everyone can receive IFN, as expenses are being covered for by the medical insurance in Japan. Another difficulty involves the informed consent in prospective randomized studies. It requires at least 5 years in order that IFN can decrease the incidence of carcinogenesis in chronic hepatitis C, with a statistical difference in the carcinogenesis rate between treated and 'untreated' patients. Since any randomized studies are considered extremely difficult in the future, we attempted to carry out this retrospective study by the multivariate analysis with statistical adjustments for possible covariates.

In the product limit analysis, IFN significantly decreased the crude rate of hepatocarcinogenesis in the

entire cohort of 2,166 patients with chronic hepatitis C. Since there were some background differences between treated and untreated patients, we tried to correct for biases including stage of fibrosis, γ -GTP value, sex, platelet count and age, which significantly affect the carcinogenesis rate. Demographic, histological and biochemical factors having been adjusted, IFN is proven to bring about a significant decrease in the hazard of carcinogenesis in patients with chronic hepatitis C en masse (hazard ratio 0.42, $p < 0.001$ by the non-time-dependent model). Taking into consideration that a significant number of patients without IFN had received anti-inflammatory medicines, which might have contributed to suppression of hepatocarcinogenesis, the actual anticarcinogenic activity of IFN may be higher than the observed. Having published results of a similar study on a cohort of 1,643 patients with a median observation period of 5.4 years in 1999 [18], we could not establish the anticarcinogenic activity of IFN because of a low risk of carcinogenesis in untreated patients (1.2% per year). Nevertheless, we expected a significant statistical difference if we could extend the median observation period to longer than 7 or 10 years in our studied patients. This has been realized in the present study, in which 2,166 patients with and without IFN therapy were observed for a median of more than 10 years. As far as we are aware, it represents the first study that has demonstrated preventive effects of IFN on the carcinogenesis rate in a large cohort of patients in a single center, in correlation with distinct responses to it, such as SVR, BR and NR.

Treatment of patients with chronic HCV infection using IFN- α and ribavirin has led to sustained loss of serum HCV RNA in 40–50% of recipients with HCV genotype 1 and 75–80% with HCV genotype 2 or 3. However, to date, the combination therapy with IFN- α and ribavirin has not been evaluated for its impact on the risk of developing HCC. Monotherapy with IFN- α achieves sustained clearance of serum HCV RNA in only 20–30% of patients; the impact of IFN- α on the development of HCC has been evaluated only in patients who had received IFN- α without ribavirin [17–20, 25–27].

Multivariate analysis definitively demonstrated that IFN lessens the carcinogenesis risk in the patients whose ALT levels decreased after therapy. Furthermore, the anticarcinogenic capacity of IFN was demonstrated not only in the patients with persistent aminotransferase normalization, but also in those with transient normalization of ALT for at least 6 or 12 months. Many authors have already described that the activity of IFN to suppress the

development of HCC in patients with HCV RNA clearance (SVR) is similar to that in patients with ALT normalization in the absence of eliminating HCV RNA (BR) [18, 25–27]. Based on these compelling lines of evidence, the anticarcinogenic activity of IFN is ascribed to the suppression of inflammatory and regenerative processes in hepatocytes. Moreno and Muriel [28] reported that IFN reverts liver fibrosis, and therefore, control of the necro-inflammatory process can suppress the growth of HCC. Tarao et al. [29] reported that high aminotransferase levels increase the rate of HCC recurrence in patients with cirrhosis. Our results stand in favor of the view that the carcinogenic process in patients with chronic hepatitis C would be enhanced by fluctuating as well as persistently elevated levels of aminotransferases. It does seem that IFN exerts suppressive effects on HCC through reduction or complete remission of inflammatory activity. Recently, a few authors reported that even transient disappearance of HCV RNA during IFN therapy contributed to a low carcinogenesis rate in the clinical course of hepatitis [17, 27]. The significance of transient HCV in decreasing hepatocarcinogenesis should be further explored and confirmed by multicenter clinical studies with rigorous virological assessments.

HCC developed in a few patients with SVR 5 years after the HCV infection had been terminated by IFN, along with normalized ALT levels. These patients would have developed minute HCC in their livers already while receiving IFN which escaped the detection by imaging modalities or screening for serological tumor markers. This would indicate the limitation of IFN in preventing HCC. IFN will not be able to suppress HCC once it has developed, even when it succeeds in eliminating HCV and suppressing necroinflammatory processes in the liver.

With many difficulties in vaccine development, the recent progress in treatment of chronic HCV infection, from IFN monotherapy to combination therapy with ribavirin, is very auspicious. SVR and BR can be achieved in up to 56% of patients with combined IFN and ribavirin [30]. There is evidence that a sustained virological response can lead to decrease in fibrosis and even reversal of cirrhosis [31]. Because HCV-associated HCC occurs almost exclusively in patients with cirrhosis, successful treatment for SVR in patients without cirrhosis is likely to prevent future development of HCC [32]. However, once cirrhosis has been established, a preventive benefit of IFN monotherapy is restricted to the patients who can achieve SVR or BR. In their meta-analysis of 3 randomized and 11 nonrandomized controlled trials, Camma et

al. [33] have reported a low but statistically significant preventive effect.

In conclusion, IFN significantly decreases the rate of hepatocarcinogenesis in patients with chronic hepatitis C, irrespective of the response to it.

Acknowledgement

This study was supported in part by a research grant from the Ministry of Health, Labour and Welfare in Japan.

References

- Ikeda K, Saitoh S, Koida I, Arase Y, Tsubota A, Chayama K, Kumada H, Kawanishi M: A multivariate analysis of risk factors for hepatocellular carcinogenesis: A prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology* 1993;18:47-53.
- Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Koida I, Arase Y, Fukuda M, Chayama K, Murashima N, Kumada H: Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: A prospective observation of 2,215 patients. *J Hepatol* 1998;28:930-938.
- Davis GL, Balart LA, Schiff ER, Lindsay K, Bodenheimer HC Jr, Perrillo RP, Carey W, Jacobson IM, Payne J, Dienstag JL, van Thiel DH, Tamburro C, Lefkowitz J, Albrecht J, Meschivitz C, Ortego T: Treatment of chronic hepatitis C with recombinant interferon alfa. A multicenter randomized, controlled trial. *N Engl J Med* 1989;321:1501-1506.
- Di Bisceglie AM, Martin P, Kassianides C, Lisker-Melman M, Murray L, Waggoner J, Goodman Z, Banks SM, Hoofnagle JH: Recombinant interferon alfa therapy for chronic hepatitis C. A randomized, double-blind, placebo-controlled trial. *N Engl J Med* 1989;321:1506-1510.
- Causse X, Godinot H, Chevallier M, Chossegros P, Zoulim F, Ouzan D, Heyraud JP, Fontanges T, Albrecht J, Meschivitz C, Trepo C: Comparison of 1 or 3 MU of interferon alfa-2b and placebo in patients with chronic non-A, non-B hepatitis. *Gastroenterology* 1991;101:497-502.
- Chayama K, Saitoh S, Arase Y, Ikeda K, Matsumoto T, Sakai Y, Kobayashi M, Unakami M, Morinaga T, Kumada H: Effect of interferon administration on serum hepatitis C virus RNA in patients with chronic hepatitis C. *Hepatology* 1991;13:1040-1043.
- Nishiguchi S, Kuroki T, Nakatani S, Morimoto H, Takeda T, Nakajima S, Shiomi S, Seki S, Kobayashi K, Otani S: Randomised trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 1995;346:1051-1055.
- Mazzella G, Accogli E, Sottili S, Festi D, Orsini M, Salzetta A, Novelli V, Cipolla A, Fabbri C, Pezzoli A, Roda E: Alpha interferon treatment may prevent hepatocellular carcinoma in HCV-related liver cirrhosis. *J Hepatol* 1996;24:141-147.
- Schalm SW, Fattovich G, Brouwer JT: Therapy of hepatitis C: Patients with cirrhosis. *Hepatology* 1997;26:S128-S132.
- Benvegno L, Chemello L, Noventa F, Fattovich G, Pontisso P, Alberti A: Retrospective analysis of the effect of interferon therapy on the clinical outcome of patients with viral cirrhosis. *Cancer* 1998;83:901-909.
- Niederer C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hurter D, Nawrocki M, Kruska L, Hensel F, Petry W, Haussinger D: Prognosis of chronic hepatitis C: Results of a large, prospective cohort study. *Hepatology* 1998;28:1687-1695.
- International Interferon-alpha Hepatocellular Carcinoma Study Group: Effect of interferon-alpha on progression of cirrhosis to hepatocellular carcinoma: A retrospective cohort study. *Lancet* 1998;351:1535-1539.
- Hu KQ, Tong MJ: The long-term outcomes of patients with compensated hepatitis C virus-related cirrhosis and history of parenteral exposure in the United States. *Hepatology* 1999;29:1311-1316.
- Ikeda K, Saitoh S, Kobayashi M, Suzuki Y, Suzuki F, Tsubota A, Arase Y, Murashima N, Chayama K, Kumada H: Long-term interferon therapy for 1 year or longer reduces the hepatocellular carcinogenesis rate in patients with liver cirrhosis caused by hepatitis C virus: A pilot study. *J Gastroenterol Hepatol* 2001;16:406-415.
- Fattovich G, Giustina G, Degos F, Tremolada F, Diodati G, Almasio P, Nevens F, Solinas A, Mura D, Brouwer JT, Thomas H, Njapoum C, Casarin C, Bonetti P, Fuschi P, Basho J, Tocco A, Bhalla A, Galassini R, Noventa F, Schalm SW, Realdi G: Morbidity and mortality in compensated cirrhosis type C: A retrospective follow-up study of 384 patients. *Gastroenterology* 1997;112:463-472.
- Valla DC, Chevallier M, Marcellin P, Payen JL, Trepo C, Fonck M, Bourliere M, Boucher E, Miguet JP, Parlier D, Lemonnier C, Opolon P: Treatment of hepatitis C virus-related cirrhosis: A randomized, controlled trial of interferon alfa-2b versus no treatment. *Hepatology* 1999;29:1870-1875.
- Kasahara A, Hayashi N, Mochizuki K, Takayanagi M, Yoshioka K, Kakumu S, Iijima A, Urushihara A, Kiyosawa K, Okuda M, Hino K, Okita K: Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998;27:1394-1402.
- Ikeda K, Saitoh S, Arase Y, Chayama K, Suzuki Y, Kobayashi M, Tsubota A, Nakamura I, Murashima N, Kumada H, Kawanishi M: Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C: A long-term observation study of 1,643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* 1999;29:1124-1130.
- Shindo M, Ken A, Okuno T: Varying incidence of cirrhosis and hepatocellular carcinoma in patients with chronic hepatitis C responding differently to interferon therapy. *Cancer* 1999;85:1943-1950.
- Yoshida H, Shiratori Y, Moriyama M, Arakawa Y, Ide T, Sata M, Inoue O, Yano M, Tanaka M, Fujiyama S, Nishiguchi S, Kuroki T, Imazeki F, Yokosuka O, Kinoyama S, Yamada G, Omata M: Interferon therapy reduces the risk for hepatocellular carcinoma: National surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy. *Ann Intern Med* 1999;131:174-181.
- Harrington DP, Fleming TR: A class of rank test procedures for censored survival data. *Biometrika* 1982;69:553-566.
- Kaplan EL, Meier P: Nonparametric estimation for incomplete observation. *J Am Stat Assoc* 1958;53:457-481.
- Cox DR: Regression models and life tables. *J R Stat Soc* 1972;34:248-275.
- SPSS Incorporation: SPSS for Windows Version 11.0 Manual. Chicago, SPSS Inc., 2001.
- Kasahara A, Hayashi N, Mochizuki K, Hiramatsu N, Sasaki Y, Kakumu S, Kiyosawa K, Okita K: Clinical characteristics of patients with chronic hepatitis C showing biochemical remission, without hepatitis C virus eradication, as a result of interferon therapy. The Osaka Liver Disease Study Group. *J Viral Hepat* 2000;7:343-351.
- Yabuuchi I, Imai Y, Kawata S, Tamura S, Noda S, Inada M, Maeda Y, Shirai Y, Fukuzaki T, Kaji I, Ishikawa H, Matsuda Y, Nishikawa M, Seki K, Matsuzawa Y: Long-term responders without eradication of hepatitis C virus after interferon therapy: Characterization of clinical profiles and incidence of hepatocellular carcinoma. *Liver* 2000;20:290-295.

- 27 Okanoue T, Itoh Y, Minami M, Sakamoto S, Yasui K, Sakamoto M, Nishioji K, Murakami Y, Kashima K: Interferon therapy lowers the rate of progression to hepatocellular carcinoma in chronic hepatitis C but not significantly in an advanced stage: A retrospective study in 1,148 patients. *J Hepatol* 1999;30:653-659.
- 28 Moreno MG, Muriel P: Remission of liver fibrosis by interferon-alpha 2b. *Biochem Pharmacol* 1995;50:515-520.
- 29 Tarao K, Takemiya S, Tamai S, Sugimasa Y, Ohkawa S, Akaike M, Tanabe H, Shimizu A, Yoshida M, Kakita A: Relationship between the recurrence of hepatocellular carcinoma (HCC) and serum alanine aminotransferase levels in hepatectomized patients with hepatitis C virus-associated cirrhosis and HCC. *Cancer* 1997;79:688-694.
- 30 Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncales FL Jr, Haussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Lin A, Hoffman J, Yu J: Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- 31 Shiratori Y, Imazeki F, Moriyama M, Yano M, Arakawa Y, Yokosuka O, Kuroki T, Nishiguchi S, Sata M, Yamada G, Fujiyama S, Yoshida H, Omata M: Histologic improvement of fibrosis in patients with hepatitis C who have sustained response to interferon therapy. *Ann Intern Med* 2000;132:517-524.
- 32 Di Bisceglie AM, Carithers RL Jr, Gores GJ: Hepatocellular carcinoma. *Hepatology* 1998;28:1161-1165.
- 33 Camma C, Giunta M, Andreone P, Craxi A: Interferon and prevention of hepatocellular carcinoma in viral cirrhosis: An evidence-based approach. *J Hepatol* 2001;34:593-602.



Springer *Reprint*

**A Long-Term Glycyrrhizin Injection Therapy
Reduces Hepatocellular Carcinogenesis Rate in
Patients with Interferon-Resistant Active Chronic
Hepatitis C: A Cohort Study of 1249 Patients**

Kenji Ikeda, Yasuji Arase, Masahiro Kobayashi, Satoshi Saitoh,
Takashi Someya, Tetsuya Hosaka, Hitomi Sezaki, Norio Akuta,
Yoshiyuki Suzuki, Fumitaka Suzuki, Hiromitsu Kumada

Digestive Diseases and Sciences 2006;51:603-609

A Long-Term Glycyrrhizin Injection Therapy Reduces Hepatocellular Carcinogenesis Rate in Patients with Interferon-Resistant Active Chronic Hepatitis C: A Cohort Study of 1249 Patients

KENJI IKEDA, YASUJI ARASE, MASAHIRO KOBAYASHI, SATOSHI SAITOH, TAKASHI SOMEYA, TETSUYA HOSAKA, HITOMI SEZAKI, NORIO AKUTA, YOSHIYUKI SUZUKI, FUMITAKA SUZUKI, and HIROMITSU KUMADA

To elucidate the influence of a glycyrrhizin therapy on hepatocarcinogenesis rate in interferon (IFN)-resistant hepatitis C, we retrospectively analyzed 1249 patients with chronic hepatitis with or without cirrhosis. Among 346 patients with high alanine transaminase value (twice or more of upper limit of normal), 244 patients received intravenous glycyrrhizin injection and 102 patients did not, after judgment of IFN resistance. Crude carcinogenesis rates in the treated and untreated group were 13.3%, 26.0% at the 5th year, and 21.5% and 35.5% at the 10th year, respectively ($P = .0210$). Proportional hazard analysis using time-dependent covariates disclosed that glycyrrhizin treatment significantly decreased the hepatocarcinogenesis rate (hazard ratio 0.49, 95% confidence interval 0.27–0.86, $P = .014$) after adjusting the background features with significant covariates. Glycyrrhizin injection therapy significantly decreased the incidence of hepatocellular carcinoma in patients with IFN-resistant active chronic hepatitis C, whose average aminotransferase value was twice or more of upper limit of normal after interferon.

KEY WORDS: chronic hepatitis; hepatitis C virus; glycyrrhizin; hepatocellular carcinogenesis; cancer prevention.

Until recently, hepatitis C virus (HCV) has been reported to be a causative agent of hepatocellular carcinoma (HCC) aside from hepatitis B virus (1–5). In our cohort studies of Japanese patients with HCV-related cirrhosis (5), the cumulative appearance rates of HCC at the 5, 10, and 15 years were 21.5%, 53.2%, and 75.2%, respectively.

The carcinogenesis rate was higher in those patients with cirrhosis caused by HCV than in those with hepatitis B virus-related cirrhosis.

Interferon (IFN) is effective in eliminating HCV in some patients with chronic hepatitis C (6–8) and cirrhosis (9–11), and in reducing hepatocellular carcinogenesis rate through suppression of necro-inflammatory process and reduction of serum alanine transaminase (ALT). Kasahara *et al.* (6) reported that sustained normal ALT value after IFN therapy was significantly associated with a decreased hepatocellular carcinogenesis rate in patients with chronic hepatitis C. Our data (7) also demonstrated an anticarcinogenic activity of IFN in patients who attained normal ALT

Manuscript received February 25, 2005; accepted July 12, 2005.

From the Department of Gastroenterology, Toranomon Hospital, Tokyo, Japan, and Okinaka Memorial Institute for Medical Research, Tokyo, Japan.

Address for reprint requests: Kenji Ikeda, MD, Department of Gastroenterology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo, 105-8470, Japan; ikedakenji@tora.email.ne.jp

level after the therapy compared with IFN-treated patients without normalization of ALT.

Oka *et al.* (12) reported in a randomized controlled trial that a kind of medicinal herb, *Sho-saiko-to*, significantly decreased hepatic carcinogenesis rate in patients with HBsAg-negative cirrhosis. Terao *et al.* (13) showed that HCC appearance rate was significantly higher in HCV-related cirrhotic patients with a high ALT value of 80 IU/mL or more than that of those with lower ALT value (<80 IU/mL), and also suggested that treatment of cirrhosis and prevention of HCC should be directed to suppress the necro-inflammation of HCV-related hepatitis. A glycyrrhizin-containing product, Stronger Neo-Minophagen C (SNMC; Minophagen Pharmaceutical Co. Ltd., Tokyo, Japan), is widely used in Japan for suppression of hepatitis activity and for prevention of disease progression in patients with hepatitis B virus and HCV-induced chronic hepatitis. Glycyrrhizin has been reported to suppress hepatic inflammation with an effect to improve the elevated ALT levels and histologic findings of the liver (14–17). We reported its favorable effect on hepatocellular carcinogenesis in those patients with chronic hepatitis C who received glycyrrhizin for more than 10 years (18).

To elucidate whether glycyrrhizin suppress the carcinogenesis rate in patients with IFN-resistant chronic hepatitis C, we retrospectively assessed a cohort of 1249 patients without sustained virologic response (SVR) after IFN therapy.

PATIENTS AND METHODS

Study Population. A total of 1249 consecutive Japanese patients with chronic hepatitis C with or without cirrhosis were examined, who did not show an SVR of HCV-RNA under IFN therapy. Sera of the patients showed positive anti-HCV (second-generation anti-HCV kit, enzyme-linked immunosorbent assay, Dainabot, Tokyo, Japan), positive HCV-RNA (nested PCR), and negative hepatitis B surface antigen (HBsAg; radioimmunoassay, Dainabot). Anti-HCV and HCV-RNA were assayed using stored frozen sera at -80°C . There were 778 men and 471 women aged 18–81 years, with a median age of 53 years in the study. They were diagnosed as having liver cirrhosis by peritoneoscopy, liver biopsy, or both between 1987–2002.

All the patients had a history of receiving once or more times of IFN therapy: 1179 patients underwent IFN monotherapy only and the other 70 patients had received an IFN plus ribavirin combination therapy before the entry of this study. A total of 347 patients showed a normal ALT for at least 6 months after cessation of IFN (biochemical responders), and the other 902 patients abnormal ALT at 6 months after the end of IFN therapy. A retrospective cohort study was performed using these 1249 consecutive patients with chronic hepatitis or cirrhosis who failed to show SVR.

Glycyrrhizin Treatment. Glycyrrhizin therapy was performed using intravenous injection of SNMC. The preparation contains 0.2% (4 mg) glycyrrhizinic acid as the main active con-

stituent, 2% (40 mg) glycine, and 0.1% (2 mg) L-cysteine in 20-mL ampoules.

Of 1249 patients with IFN-resistant chronic liver disease, 453 patients underwent glycyrrhizin injection therapy and the remaining 796 patients did not receive the therapy until the end of observation. The purpose of the introduction of the glycyrrhizin injection therapy was to suppress elevated ALT and to prevent disease progression in all the patients. Of the 453 patients, 129 (28.5%) received a daily dose of 40–60 mL of SNMC (80–120 mg as glycyrrhizin) and 324 (71.5%) received 80–100 mL (160–200 mg as glycyrrhizin). A total of 110 patients received the treatment for less than 2 years and 107 patients continued the therapy for 2–4 years, 132 patients for 4–6 years, and the remaining 104 patients for 6 years or longer. When the treatment was regarded as effective from the viewpoint of ALT levels, treatment was usually continued for a period as long as possible. As a result, a median daily dose of 100 mL of SNMC was administered 3 times a week during a median period of 4.3 years (range, 0.1–14.5 years) in the treated group.

Two (0.44%) of 453 treated patients were withdrawn from the glycyrrhizin injection therapy because of side effects: 1 because of hypertension and 1 from skin rash.

Background and Laboratory Data of Patients With and Without Glycyrrhizin Therapy. Table 1 summarizes the profiles and data of the patients at the time of diagnosis of chronic hepatitis with or without cirrhosis. The male/female ratio was not different between the 2 groups. Median age was older by 2 years in the treated group than in the untreated group ($P < .001$). Results of histologic staging of liver disease were classified according to Desmet *et al.* (19). F1stage hepatitis was found significantly more often in the untreated group than in the glycyrrhizin group ($P < .001$, χ^2 test). Both AST and ALT median levels were significantly higher in the treated group than in the untreated group ($P < .001$). HCV subtype was analyzed by the immunoserologic typing method with a commercial kit (Kokusai Diagnostic Corporation, Kobe, Japan): serologic group 1 indicated genotypes 1a and 1b, and group 2 included 2a and 2b subtypes. The rate of HCV serologic group 1 was significantly higher in the glycyrrhizin group than in the untreated group ($P = .032$).

Follow Up. Follow-up of the patients was made monthly after the judgment of IFN-resistance by monitoring hematologic, biochemical, and virologic data. Imaging diagnosis with ultrasonography (US) and/or computed tomography (CT) was made 3 or more times per year in a majority of patients with cirrhosis and once a year in patients without cirrhosis. Angiographic study was performed only when HCC was strongly suspected on US or CT.

When angiography revealed a characteristic hypervascular nodule suggesting a specific finding for HCC, no histologic examination was made in a majority of these patients. An increasing trend of tumor markers was also taken into account in establishment of the diagnosis of HCC. Microscopic examination through a fine needle biopsy was also performed in patients whose angiogram did not show a typical image of HCC.

The number of cases lost to follow-up was 121 (9.7%): 28 patients (6.2%) in the glycyrrhizin group and 93 (11.7%) in the untreated group. Because the outcomes regarding appearance of HCC were not identified in these patients, they were dealt as censored data in the following statistics (20). Death unrelated to HCC was also classified as withdrawal and regarded as a censored case. The median observation period of the total number of patients was 5.7 years with a range of 0.1–16.1 years. Because

GLYCYRRHIZIN FOR CHRONIC HEPATITIS

TABLE 1. PATIENT PROFILES AND LABORATORY DATA AT TIME OF JUDGMENT OF IFN RESISTANCE

	Glycyrrhizin Group (n = 453)	Untreated Group (n = 796)	P
Demographics			
Gender (M/F)	283/170	495/301	.92
Age (y)*	54 (25-81)	52 (18-77)	<.001
Observation period (y)*	8.3 (0.1-16.1)	5.1 (0.1-13.1)	<.001
Liver histology			
F1	146 (32.7%)	502 (64.0%)	<.001
F2	193 (43.3%)	192 (24.5%)	
F3	38 (8.5%)	52 (6.6%)	
F4	69 (15.5%)	38 (4.8%)	
Laboratory data*			
Aspartic transaminase (IU/L)*	81 (19-446)	54 (11-355)	<.001
ALT (IU/L)*	122 (12-630)	83 (10-822)	<.001
HCV serologic group 1 (1a or 1b)	360 (80.2%)	582 (73.7%)	.032
Group 2 (2a or 2b)	73 (16.3%)	165 (20.9%)	
Others	16 (3.6%)	43 (5.4%)	

*Expressed as median (range).

many patients receiving glycyrrhizin therapy migrated from the untreated group to the treated group, observation period of the untreated group was significantly shorter than that of the treated group (see Table 1). The date of the last follow-up for this study was September 1, 2003.

Statistical Analysis. Nonparametric procedures were employed for the analysis of background characteristics of the patients, including Mann-Whitney *U*-test and χ^2 method. HCC appearance rates were calculated from the time period between the judgment of IFN ineffectiveness and appearance of HCC in each group, using Kaplan-Meier technique (20). The differences in carcinogenesis curves were tested using the log-rank test. Independent factors associated with the appearance rate of HCC were studied using time-dependent Cox regression analysis (21). An interaction term of IFN treatment and "waiting time" to the therapy was introduced in the analysis as a time-dependent covariate. The independence of treatment factor from "waiting time" was also confirmed by log-minus-log plot of proportional hazard model. Several variables were transformed into categorical data consisting of 2-3 simple ordinal numbers to estimate each hazard ratio. All factors found to be at least marginally as-

sociated with liver carcinogenesis ($P < .15$) were tested by the multivariate Cox proportional hazard model. A *P*-value of less than .05 was considered to be significant. All data analysis was performed using the computer program SPSS version 11 (22).

RESULTS

Initial Aminotransferase and Carcinogenesis Rates

Patients with and without glycyrrhizin therapy were classified into 6 categories according to average ALT value during the first year after cessation of IFN therapy: group 1, normal ALT; group 2, <1.5 times of upper limit of normal (ULN); group 3, 1.5-2 times ULN; group 4, 2-3 times ULN; group 5, 3-4 times of ULN; and group 6, >4 times ULN. Hepatocellular carcinogenesis rates were 2.5%, 5.0%, 8.1%, 11.8%, 12.0%, and 12.7% at the end of 5 years and 6.6%, 7.2%, 19.6%, 15.1%, 21.0%, and 39.3% at 10 years, respectively (Figure 1). There was a significant statistical difference among the 6 subgroups (log-rank test, $P < .0001$). The higher the average ALT, the higher the carcinogenesis rate was.

Influence of Glycyrrhizin on Carcinogenesis in Patients With High Aminotransferase

Glycyrrhizin therapy was usually performed in patients with a high ALT value and high hepatitis activity. In this retrospective study, average ALT values were significantly different between the treated and untreated groups: group 1, normal average ALT was found in 38 among patients with glycyrrhizin therapy and in 188 among patients without therapy; in group 2, ALT <1.5 times of ULN was found in 42 and 331; in group 3, 1.5-2 times ULN in 84 and 138; in group 4, 2-3 times ULN in 143 and 92; in group 5, 3-4 times in 53 and 29; and in group 6, ALT

TABLE 2. INDEPENDENT RISK FACTORS AFFECTING HEPATOCELLULAR CARCINOGENESIS

Factors	Category	Risk Ratio (95% CI)	P
Fibrotic stage	F1	1	
	F2-3	2.94 (1.20-7.21)	.018
	F4 (cirrhosis)	9.21 (3.73-22.8)	<.001
Gender	1: Female	1	
	2: Male	2.80 (1.35-5.81)	.006
Glycyrrhizin injection (SNMC)*	1: No	1	
	2: Yes	0.49 (0.27-0.86)	.014

Time-dependent Cox proportional hazard analysis. *SNMC, Stronger Neo-Minophagen C (herbal medicine containing glycyrrhizin).




Fig 1. Carcinogenesis rates according to initial ALT values classified into six groups: (1) normal ALT, (2) <1.5 times ULN, (3) 1.5–2 times ULN, (4) 2–3 times ULN, (5) 3–4 times ULN, and (6) >4 times of ULN. The higher the average ALT, the higher the carcinogenesis rate was.

>4 times ULN in 93 of the glycyrrhizin group and 18 of the untreated group. The rate of a high ALT value of twice or more of ULN in the glycyrrhizin treated group (64.2%, 289/453) was significantly higher than that of the untreated group (16.2%, 129/796).

Of the 418 patients with a high average ALT in both groups, 68 patients showed a normal ALT value for at least 6 months just after IFN therapy (biochemical response). Because biochemical response with normal ALT for a certain period after IFN was likely to affect carcinogenesis rates in those patients, biochemical responders were excluded in the following analyses about the influence of glycyrrhizin on carcinogenesis: after all, 244 patients with glycyrrhizin therapy and the 102 patients without therapy were assessed.

Cumulative hepatocellular carcinogenesis rates were calculated in these 346 patients with a high average ALT values, excluding biochemical responders from both groups. Carcinogenesis rates in the glycyrrhizin group and the untreated group were 6.5% and 13.3% at the end of year 3, 13.3% and 26.0% at the end of year 5, 17.7% and 28.3% at the end of year 7, and 21.5% and 35.5% at year 10, respectively (Figure 2). In the stratified and selected patient group, the carcinogenesis rate of glycyrrhizin-treated group was significantly lower than that of the untreated group (log-rank test, $P = .0210$).

Carcinogenesis Rates According to Hepatitis Staging

Crude carcinogenesis rates were compared between the groups, according to each hepatitis stage. In F1 stage chronic hepatitis, hepatocellular carcinogenesis rates in the glycyrrhizin group ($n = 82$) and the untreated group ($n = 32$) were 1.4% and 4.2% at year 5 and 7.0% and 12.1% at 10 years, respectively (Figure 3A). In F2–3 stage chronic hepatitis, hepatocellular carcinogenesis rates in




Fig 2. Carcinogenesis rates in patients with high average ALT values of twice or more of ULN, excluding those patients with biochemical responders who continued a normal ALT value at least 6 months just after IFN therapy. The carcinogenesis rate of glycyrrhizin-treated group was significantly lower than that of the untreated group (log-rank test, $P = .0210$).

the glycyrrhizin group ($n = 121$) and the untreated group ($n = 53$) were 14.8% and 28.4% at the end of year 5, and 21.5% and 38.6% at year 10, respectively (Figure 3B). In patients with F4 stage chronic hepatitis (cirrhosis), hepatocellular carcinogenesis rates in the glycyrrhizin group ($n = 38$) and the untreated group ($n = 15$) were 35.2% and 58.0% at the end of year 5, and 57.2% and 58.0% at year 10, respectively (Figure 3C).

In each fibrotic stage of hepatitis, carcinogenesis rates were lower in the glycyrrhizin group than in the untreated group, but statistical significance was not obtained owing to shortage of patient number in these stratified groups.

Aminotransferase Activity Before and After Glycyrrhizin Therapy

ALT values in the patients with glycyrrhizin treatment were serially assessed in those patients who began the therapy after they had shown a high average ALT value (Figure 4). Median value of ALT at the beginning of the glycyrrhizin therapy was 150 IU/L (25th percentile 120, 75th percentile 221), 72 IU/L at month 3, 70 IU/L at month 6, and 64 IU/L (25th percentile 48, 75th percentile 93) at month 12, respectively. ALT value significantly decreased after the initiation of glycyrrhizin injection therapy.

Factors Affecting Carcinogenesis Rates in Active Hepatitis and Cirrhosis

In the selected patients with active hepatitis with an average ALT value of twice ULN or higher, multivariate analysis was performed to explore associating factors with carcinogenesis, using time-dependent Cox proportional hazard model. Time between the judgment of IFN




Fig 3. Carcinogenesis rates according to hepatitis staging: (a) F1 stage hepatitis, (b) F2–F3 stage hepatitis, and (c) F4 or cirrhotic stage. In each fibrotic stage of hepatitis, carcinogenesis rates were lower in the glycyrrhizin group than in the untreated group.

ineffectiveness and initiation of glycyrrhizin therapy was set as a time-dependent variable to clarify the significance of glycyrrhizin therapy in the clinical course of HCV-related chronic liver diseases. Patients with biochemical response with a normal ALT value sustained for at least 6 months after IFN therapy were also excluded from the analysis.

In multivariate analysis, following 3 factors influenced the carcinogenesis: fibrotic staging, gender ($P = .006$), and glycyrrhizin therapy ($P = .014$). When a hazard of F1 stage fibrosis for carcinogenesis was set as 1 in the model, hazard ratio of F2–F3 stage fibrosis was calculated as 2.94 ($P = .018$), and that of F4 (cirrhosis) was estimated as 9.21 ($P < .001$). Similarly, the hazard ratio for carcinogenesis of male gender was 2.80, and use of glycyrrhizin independently decreased the carcinogenesis rate in patients with active chronic hepatitis after IFN therapy. Following factors did not affect the HCC appearance rate

significantly: age, association of diabetes mellitus, serologic grouping of HCV, HCV-RNA concentration, AST, ALT at the time before IFN therapy, and bilirubin.

DISCUSSION

IFN is effective in patients with chronic liver disease caused by HCV, from the viewpoints of anti-inflammatory effect and cancer prevention (6–11). Although the carcinogenesis rate is noticeably reduced when aminotransferase becomes normal with or without HCV-RNA eradication (6–8) after the therapy, the rate of normalization of ALT after IFN therapy is approximately half of patients with high viral load and group 1 HCV-subtype.

This retrospective study was undertaken to evaluate whether long-term glycyrrhizin injection therapy could decrease hepatocellular carcinogenesis rate in patients with IFN-resistant HCV-related chronic hepatitis and cirrhosis. Because it requires at least 5 years to show a statistical difference in carcinogenesis rate from hepatitis or cirrhosis between glycyrrhizin-treated and “untreated” groups, a prospective randomized trial using untreated control patients is difficult from both ethical and medical viewpoints in Japan, where glycyrrhizin injection therapy is covered by standard medical insurance and is already regarded as a usual choice of therapy as a salvaging procedure for IFN-ineffective patients. We, therefore, attempted to carry out this retrospective cohort study to prove an anticarcinogenic activity of glycyrrhizin, with a statistical adjustment using possible covariates explored in multivariate analysis.




Fig 4. Aminotransferase activity before and after glycyrrhizin therapy. ALT value significantly decreased after the initiation of glycyrrhizin injection therapy.

Because glycyrrhizin injection therapy was chiefly performed for patients with a high ALT value and because cancer prevention was meaningful in just those patients with a high carcinogenesis risk with high hepatitis activity, we analyzed the role of a long-term glycyrrhizin injection therapy in the patients with a high ALT value. The treated group consisted of significantly more numbers of patients with a high ALT value of twice or more of ULN. When carcinogenesis rates were assessed only in those patients with a high ALT value of twice or more ULN excluding biochemical responders, the rate of the treated group became significantly higher than that of the untreated group ($P = .021$). The cancer preventive effect of glycyrrhizin in IFN-resistant patients was also confirmed by time-dependent Cox proportional analysis that adjusted the background features of the retrospective cohort (hazard ratio = 0.49, $P = .014$). We previously reported a study focused on the anticarcinogenic action of glycyrrhizin for patients with chronic hepatitis C, but the pilot study only demonstrated that 10 years or longer treatment with glycyrrhizin ($n = 84$) could suppress the carcinogenesis rate (18). Current study dealing with a large cohort ($n = 1249$) showed that glycyrrhizin injection therapy significantly decreased carcinogenesis rate irrespective of the length of treatment when comparison was made in a selected patient cohort with high hepatitis activity.

Although a statistically significant difference was not shown for a lack of sufficient patient number in subgroups of chronic hepatitis and cirrhosis, this study also demonstrated that glycyrrhizin was effective not only in chronic hepatitis but also in cirrhosis. Considering that liver cirrhosis generally shows a resistance to IFN treatment, our current study demonstrated encouraging results from the viewpoint of HCC prevention. When IFN therapy was attempted in 7 patients with decompensated cirrhosis by Nevens *et al.* (23), complications sometimes occurred in these patients, including variceal bleeding, aggravation of ascites or encephalopathy, development of pneumonia, and recurrence of spontaneous bacterial peritonitis or gastric ulcer bleeding. Because patients with cirrhosis usually showed lower platelet and leukocyte counts than those with chronic hepatitis and because cirrhotic patients tended to show deterioration with a large dose of IFN, glycyrrhizin therapy proved to be a useful alternative of therapy. Intermittent long-term glycyrrhizin therapy was well tolerated with withdrawal of only 2 patients (0.44%).

Because carcinogenesis is not a single-step event but a complex, multistep process, the exact mechanism of the glycyrrhizin activity in suppression of liver carcinogenesis remains unknown. One of the principal roles of long-term administration of glycyrrhizin in decreasing the carcinogenesis rate is considered to be anti-inflammatory,

which blocks the active carcinogenic process of continuous hepatic necro-inflammation and cell damage. In the treated group, median ALT values markedly decreased after initiation of the glycyrrhizin injection, suggesting that pathologic process of hepatocyte necrosis or apoptosis was significantly suppressed by glycyrrhizinic acid. The importance of the action of amino acids, glycine and cysteine contained in SNMC has not been completely explained, but they have been demonstrated to suppress increased aldosterone levels that are induced by glycyrrhizinic acid. Tarao *et al.* (24) reported that high aminotransferase level resulted in an increase of an HCC recurrence rate in patients with HCC. From the viewpoint of these anti-inflammatory activities, SNMC may be considered to only postpone the time of HCC appearance in the clinical course of cirrhosis. Because the entire process of hepatocellular carcinogenesis from the initial transformation of a hepatocyte to a detectable growth of cancer is considered to take at least several years, the influence of glycyrrhizin on the carcinogenesis rate will not be evaluated in a short period. Although several reports suggested a relationship of anti-hepatitis B core antibody or hepatitis B surface antibody with carcinogenesis (25–27), we could not show the association because of insufficient available data.

Because current data were obtained from a retrospective cohort analysis, dose of glycyrrhizin per time, times of injection per week, and duration of therapy varied in each patient in the treated group. To elucidate the cancer preventive effect of glycyrrhizin therapy in active HCV-related liver disease, we should further stratify the treated patients or perform much more detailed statistical procedures. Future studies should, therefore, aim at defining the basic oncogenic mechanisms and roles of long-term administration of glycyrrhizin in carcinogenesis in patients with cirrhosis caused by HCV.

In conclusion, a long-term intermittent glycyrrhizin therapy for a few years or more successfully reduced hepatocellular carcinogenesis in patients with HCV-related chronic liver disease. A randomized control study with a larger number of cases, with or without glycyrrhizin therapy, is expected to confirm the effectiveness of this therapy.

REFERENCES

1. Bruix J, Calvet X, Costa J. *et al.*: Prevalence of antibodies to hepatitis C virus in Spanish patients with hepatocellular carcinoma and hepatic cirrhosis. *Lancet* 2:1004–1006, 1989
2. Colombo M, Kuo G, Choo QL, *et al.*: Prevalence of antibodies to hepatitis C virus in Italian patients with hepatocellular carcinoma. *Lancet* 2:1006–1008, 1989
3. Hasan F, Jeffers LJ, Medina MD, *et al.*: Hepatitis C-associated hepatocellular carcinoma. *Hepatology* 12:589–591, 1990

GLYCYRRHIZIN FOR CHRONIC HEPATITIS

4. Kew MC, Houghton M, Choo QL, *et al.*: Hepatitis C virus antibodies in southern African blacks with hepatocellular carcinoma. *Lancet* 335:873-874, 1990
5. Ikeda K, Saitoh S, Koida I, *et al.*: A multivariate analysis of risk factors for hepatocellular carcinogenesis—a prospective observation of 795 cases with viral and alcoholic cirrhosis. *Hepatology* 18:47-53, 1993
6. Kasahara A, Hayashi N, Mochizuki K, *et al.*: Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. Osaka Liver Disease Study Group. *Hepatology* 27:1394-1402, 1998
7. Ikeda K, Saitoh S, Arase Y, *et al.*: Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C—a long-term observation study of 1643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* 29:1124-1130, 1999
8. Yabu K, Kiyosawa K, Mori H, *et al.*: Serum collagen type IV for the assessment of fibrosis and resistance to interferon therapy in chronic hepatitis C. *Scand J Gastroenterol* 29:474-479, 1994
9. Nishiguchi S, Kuroki T, Nakatani S, *et al.*: Randomized trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 346:1051-1055, 1995
10. Mazzella G, Accogli E, Sottili S, *et al.*: Alpha interferon treatment may prevent hepatocellular carcinoma in HCV-related liver cirrhosis. *J Hepatol* 24:141-147, 1996
11. Benvegna L, Chemello L, Noventa F, *et al.*: A. Retrospective analysis of the effect of interferon therapy on the clinical outcome of patients with viral cirrhosis. *Cancer* 83:901-909, 1998
12. Oka H, Yamamoto S, Kuroki T, *et al.*: Prospective study of chemoprevention of hepatocellular carcinoma with Sho-saiko-to (TJ-9). *Cancer* 76:743-749, 1995
13. Tarao K, Rino Y, Ohkawa S, *et al.*: Association between high serum alanine aminotransferase levels and more rapid development and higher rate of incidence of hepatocellular carcinoma in patients with hepatitis C virus-associated cirrhosis. *Cancer* 86:589-595, 1999
14. Fujisawa K, Watanabe Y, Kimura K: Therapeutic approach to chronic active hepatitis with glycyrrhizin. *Asian Med J* 23:745-756, 1980
15. Suzuki H, Ohta Y, Takino T, *et al.*: Effects of glycyrrhizin on biochemical tests in patients with chronic hepatitis. Double blind trial. *Asian Med J* 26:423-438, 1983
16. Wildhirt E: Experience in Germany with glycyrrhizinic acid for the treatment of chronic viral hepatitis. In: *Viral Hepatitis and Liver Disease* 658-661, 1994, Springer Verlag, Germany
17. Rossum TGJ van, Vulto AG, Hop WCJ, *et al.*: Intravenous glycyrrhizin for the treatment of chronic hepatitis C: a double-blind, randomized, placebo-controlled phase III trial. *J Gastroenterol Hepatol* 14:1093-1099, 1999
18. Arase Y, Ikeda K, Murashima N, *et al.*: The long term efficacy of glycyrrhizin in chronic hepatitis C patients. *Cancer* 79:1494-1500, 1997
19. Desmet VJ, Gerber M, Hoofnagle JH, *et al.*: Classification of chronic hepatitis: diagnosis, grading, and staging. *Hepatology* 19:1513-1520, 1994
20. Kaplan EL, Meier P: Nonparametric estimation for incomplete observation. *J Am Stat Assoc* 53:457-481, 1958
21. Cox DR: Regression models and life tables. *J R Stat Soc* 34:248-275, 1972
22. SPSS Inc. SPSS for Windows version 11.0 manual. Chicago, SPSS Inc., 2001
23. Nevens F, Goubau P, Van Eyken P, *et al.*: Treatment of decompensated viral hepatitis B-induced cirrhosis with low doses of interferon alpha. *Liver* 13:15-19, 1993
24. Tarao K, Takemiya S, Tamai S, *et al.*: Relationship between the recurrence of hepatocellular carcinoma (HCC) and serum alanine aminotransferase levels in hepatectomized patients with hepatitis C virus-associated cirrhosis and HCC. *Cancer*, 79:688-694, 1997
25. Gaeta GB, Rapicetta M, Sardaro C, *et al.*: High prevalence of co-occurrence of anti-HCV and anti-HBc antibodies in chronic hepatitis patients from southern Italy. *Ital J Gastroenterol* 22:350-351, 1990
26. Yu MC, Yuan JM, Ross RK, *et al.*: Presence of antibodies to hepatitis B surface antigen is associated with an excess risk for hepatocellular carcinoma among non-Asians in Los Angeles county, California. *Hepatology* 25:226-228, 1997
27. Koike K, Kobayashi M, Gondo M, *et al.*: Hepatitis B virus DNA is frequently found in liver biopsy samples from hepatitis C virus-infected chronic hepatitis patients. *J Med Virol* 54:249-255, 1998



Prediction model of hepatocarcinogenesis for patients with hepatitis C virus-related cirrhosis. Validation with internal and external cohorts

Kenji Ikeda^{1,*}, Yasuji Arase¹, Satoshi Saitoh¹, Masahiro Kobayashi¹, Takashi Someya¹, Tetsuya Hosaka¹, Norio Akuta¹, Yoshiyuki Suzuki¹, Fumitaka Suzuki¹, Hitomi Sezaki¹, Hiromitsu Kumada¹, Akihisa Tanaka², Hideharu Harada²

¹Department of Gastroenterology, Okinaka Memorial Institute for Medical Research, Toranomon Hospital, Toranomon 2-2-2, Minato-ku, Tokyo 105-8470, Japan

²Department of Gastroenterology, Tokyo National Hospital, National Hospital Organization, Tokyo, Japan

See Editorial, pages 1013–1016

Background/Aims: To estimate hepatocarcinogenesis rates in patients with hepatitis C virus (HCV)-related cirrhosis, an accurate prediction table was created.

Methods: A total of 183 patients between 1974 and 1990 were assessed for carcinogenesis rate and risk factors. Predicted carcinogenesis rates were validated using a cohort from the same hospital between 1991 and 2003 ($n=302$) and an external cohort from Tokyo National Hospital between 1975 and 2002 ($n=205$).

Results: The carcinogenesis rates in the primary cohort were 28.9% at the 5th year and 54.0% at the 10th year. A proportional hazard model identified alpha-fetoprotein (≥ 20 ng/ml, hazard ratio 2.30, 95% confidence interval 1.55–3.42), age (≥ 55 years, 2.02, 95% CI 1.32–3.08), gender (male, 1.58, 95% CI 1.05–2.38), and platelet count ($< 100,000$ counts/mm³, 1.54, 95% CI 1.04–2.28) as independently associated with carcinogenesis. When carcinogenesis rates were simulated in 16 conditions according to four binary variables, the 5th- and 10th-year rates varied from 9 to 64%, and 21–93%, respectively. Actual carcinogenesis rates in the internal and external validation cohorts were similar to those of the simulated curves.

Conclusions: Simulated carcinogenesis rates were applicable to patients with HCV-related cirrhosis. Since, hepatocarcinogenesis rates markedly varied among patients depending on background features, we should consider stratifying them for cancer screening and cancer prevention programs.

© 2006 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

Keywords: Cirrhosis; Hepatocellular carcinoma; Carcinogenesis; Hepatitis C virus; Simulation; Proportional hazard model; Validation; Prediction

1. Introduction

There is increasing evidence that chronic hepatitis C virus (HCV) infection is closely associated with the occurrence of hepatocellular carcinoma (HCC) [1–4]. The

incidence of patients with HCV-related HCC has increased recently in several parts of the world [5–9]. In Japan, blood transfusion and parenteral drug use became prevalent in 1960s, and patients with HCV-related cirrhosis gradually increased around 1980s. Since, an effective and truly curative therapy for a large and advanced HCC still remains limited at best, evaluation and assessment of carcinogenesis in chronic liver disease and detection at an early stage of HCC are of great importance. Reports of HCC development rates in HCV-cirrhosis differ [10–13], probably due to

Received 5 February 2005; received in revised form 31 January 2006; accepted 20 February 2006; available online 13 March 2006

* Corresponding author. Tel.: +81 44 877 5111; fax: +81 44 860 1623.

E-mail address: ikedakenji@tora.email.ne.jp (K. Ikeda).

differences of patient characteristics in varied study populations. The lack of reliable data as to the natural history of cirrhosis makes it difficult to evaluate the exact role and cost-effectiveness of interferon therapy.

Platelet count has been used to predict hepatocarcinogenesis [10,13,14], but its usefulness for distinguishing the HCC appearance rate is based on discrimination between chronic hepatitis and cirrhosis [15–18]. Predicting carcinogenesis solely on the basis of platelet count is less valuable in a cohort of patients with cirrhosis, because the liver disease has already advanced to a certain stage with a uniformly low platelet count. When a cohort of patients with HCV-related cirrhosis is analyzed by platelet count, it is usually not possible to discriminate between a super-high-risk group for carcinogenesis and a relatively low-risk group. The availability of a general model that can accurately predict the HCC development rate in HCV-related disease based on readily available data would be helpful in planning the treatment of these patients. Moreover, such a model could be used for the selection and stratification of patients for clinical trials.

In this study, we tried to develop a prediction model for hepatocarcinogenesis rate, using a large cohort with a long observation period. This model was also validated with two independent patient cohorts for generalization and clinical application.

2. Patients and methods

2.1. Study population

Among 457 consecutive patients diagnosed with liver cirrhosis between 1974 and 1990 at Toranomon Hospital, Tokyo, 258 patients had positive anti-HCV antibody (second-generation anti-HCV, enzyme-linked immunosorbent assay, Dainabot, Japan), positive HCV-RNA, and negative hepatitis B surface antigen (HBsAg, radioimmunoassay, Dainabot, Tokyo, Japan). Among them, 75 patients met either of the following exclusion criteria: (1) possible association with HCC, (2) association of hemochromatosis, autoimmune liver disease, primary biliary cirrhosis, alpha-1-anitrypsin deficiency, or Wilson disease, (3) daily drinking habit of 75 g or more, (4) alpha-fetoprotein (AFP) of 400 ng/ml or higher, (5) advanced and decompensated stage of cirrhosis with encephalopathy and refractory ascites, or (6) a short follow-up period of 6 months or less. We excluded those patients with Child–Pugh [19] stage C, because of substantial difference in carcinogenesis [20,21]. Consequently, 183 patients were retrospectively analyzed for HCC appearance rate.

2.2. Background and laboratory data

Table 1 summarizes the profiles and data of the 183 patients at the time of diagnosis. The group consisted of 92 men and 91 women aged from 28 to 80 (median, 55 years). The diagnosis of cirrhosis was made by peritoneoscopy, biopsy or both in 118 patients, and by clinical symptoms with ultrasonographic findings in 55 patients. When the ultrasonography (US) showed a typical irregular-surfaced liver with coarse internal architecture in addition to overt ascites or esophageal varices demonstrated by fiberoptic examination, we regarded the disease as cirrhosis. Although 12.7% of patients (23/181) showed normal aminotransferases at the time of the diagnosis of cirrhosis, all of those patients had been followed up as having chronic hepatitis with fluctuated aminotransferases.

Table 1
Patient profiles and laboratory data at the time of diagnosis of cirrhosis (primary cohort of Toranomon Hospital between 1974 and 1990, *n* = 183)

	Median (range)	Valid data
Demography and backgrounds		
Total number	183	
Sex (M/F)	92/91	
Age, median (range)	55 (28–80)	
Diagnostic method		
Peritoneoscopy and/or biopsy	118 (64.5%)	
Clinical (ultrasonography plus varices or ascites)	65 (35.5%)	
History of blood transfusion	82 (44.8%)	
Diabetes mellitus	23 (12.6%)	
Previous medical history of chronic hepatitis	34 (18.6%)	
Interferon therapy during observation	24 (12.0%)	
Refractory ascites and/or encephalopathy	0	
Hepatitis B surface antigen, positive	0 (100%)	
Anti-hepatitis C virus, positive	183 (100%)	
Hepatitis C virus RNA, positive	183 (100%)	
Child–Pugh score A	136 (74.3%)	
Child–Pugh score B	47 (25.7%)	
Observation period (year) median (range)	10.5 (0.5–26.0)	
Laboratory data		
Albumin (normal, 3.9–5.1 g/dl)	3.9 (2.5–5.1)	183
Bilirubin (normal, 0.3–1.1 mg/dl)	1.1 (0.4–4.4)	183
Aspartic transaminase (normal, ≤ 38 IU/L ^a)	69 (17–372)	181
Alanine transaminase (normal, ≤ 50 IU/L ^a)	56 (9–282)	181
Platelet (normal, $149\text{--}315 \times 1000^3/\text{mm}^3$)	95 (33–213)	183
ICG R15 ^b (normal, $\leq 10\%$)	27 (6–81)	173
Prothrombin time (normal, $\geq 70\%$)	79 (54–100)	183
Gamma-globulin (normal, < 1.5 g/dl)	1.9 (1.0–3.5)	174
Alpha-fetoprotein (normal, < 5 mg/L)	16.5 (3–256)	166
HCV genotype^c		
1b	107 (69.9%)	153
2a/2b	39 (25.5%)	
Combined/others	7 (4.6%)	
Not examined	30	

^a Numbers of normal aspartic and alanine transaminases were 25 (13.8%) and 69 (38.1%), respectively. Both transaminases were normal at the time of the diagnosis of cirrhosis in 23 patients (12.7%).

^b ICG R15: indocyanine green retention rate at 15 min.

^c HCV genotyping was classified according to Simmonds et al. [22].

HCV-RNA measurement and HCV genotyping [22] are analyzed with nested polymerase chain reaction using initial sera stored at -80°C .

2.3. Follow-up of patients and diagnosis of hepatocellular carcinoma

Patients were followed-up monthly following the diagnosis of cirrhosis by monitoring hematological and biochemical data. Diagnostic imaging by US was taken approximately once a year in each patient. After 1987, imaging procedures with US or computerized tomography (CT) were performed twice or more per year in the majority of patients for early detection of HCC. HCC was diagnosed by typical hypervascular characteristics on angiography. When combined use of imagings could not demonstrate a typical image of HCC (13/107, 12.1%), a fine needle biopsy was obtained for microscopic examination.

Twenty-four patients (13.1%) received interferon during the follow-up period. Since the therapy could affect the natural clinical course of viral hepatitis, they were treated as censored at the time of the initiation of interferon in the analysis. Sixteen (8.7%) cases were lost to follow-up, and median observation period was 10.5 years (range, 7.0–14.9). Those patients lost to follow-up were treated as censored data in the following statistics.