

Firstly, the HBs antigen was prematurely terminated in the mutant strain reported by Yeh et al. (34). In this regard, a similar amino acid substitution of the B domain of the polymerase FLLA motif in woodchuck hepatitis virus (WHV) treated with lamivudine was reported (15, 28). The HBs antigen in these WHV mutant strains also had premature stop codons. These findings suggest that the mutant strains of HBV and WHV cannot replicate and spread by themselves because of the lack of HBs antigen. Such strains are thought to replicate by using in vivo-supplied HBs antigen from wild-type strains as helper antigens. In contrast, the novel strain identified in this study had no premature termination of the HBs gene. The in vitro study suggested that the strain had a replication ability similar to that of the wild type. Furthermore, we also showed that the strain infected and reached a high viral load in human hepatocyte-chimeric mice. Although the inoculum contained only a small amount of wild-type strain (one of 12 clones), all clones obtained from mouse serum were mutant strains (rtA181T). Considering these results and the fact that the index patient showed high viral titers after breakthrough (more than 7.6 log copies/ml), this mutant strain can spread and replicate by itself and has strong replicative ability.

Secondly, the substitutions identified in this study appeared with nucleotide and amino acid substitutions in the spacer region of the polymerase (S331C). There are only a few studies that reported the function of the spacer domain (19–21, 28), leaving the biological significance of this region unknown. The substitution in the spacer region reappeared with the A181T mutation in the RT domain in the index patient after the patient restarted lamivudine therapy. Although our study showed no significant contribution of this mutation to drug resistance (Fig. 3 and 4; Table 1), the significance of the mutation in this region (fingers in the HBV polymerase homology model [8]) should further be investigated.

Recently, the amino acid substitutions rtA181T and rtA181V were reported to emerge with resistance against adefovir (11, 32). Tillmann et al. (29) reported one case in which the virus developed the rtA181T mutation during famciclovir breakthrough. The A556T mutation of WHV, analogous to the rtA181T mutation of HBV, has been reported to be associated with lamivudine resistance (15, 28). These results indicate that the amino acid substitutions at position 181 may associate with resistance against many nucleoside analogues, including lamivudine, famciclovir, and adefovir. Although our in vitro study indicated that the rtA181T mutant had no resistance against adefovir and the animal study showed that combination therapy with lamivudine and adefovir effectively reduced the virus load in woodchucks (15), such combination therapy did not produce sufficient suppression of HBV in the index patient (Fig. 1A). The amino acid substitution at position 181 has to be further analyzed with regard to resistance to anti-HBV drugs.

The rtA181T mutation detection system using RFLP PCR developed in this study is a useful tool, as we were able to distinguish the wild type from all mutants with nucleotide substitutions in a given region. The system also enabled us to monitor the fluctuation of the wild-type/mutant ratio during therapy against HBV (Fig. 1 and 6). The incidence of rtA181T mutants with an intact YMDD motif is rare in Japanese patients with chronic HBV infection treated with lamivudine. Interestingly, 4 of the 39 (10%) patients who developed lamivudine breakthrough and were positive for YMDD mutants were found to have small amounts of rtA181T mutant strains. Different from the previous report (34), the mutants did not take over another strain and were not preceded by exacerbation. We have to monitor these patients carefully for further population change of mutants and for exacerbation of hepatitis.

A recent study reported that the prevalence of genotype A HBV infection is increasing in Japan and that the incidence of disease chronicity is higher than for other genotypes (26). It is thus expected that an increasing number of the sexually active population will receive nucleoside analogue therapy against HBV and multiple mutant strains can potentially emerge and spread along with long-term treatment. There is an increasing possibility of emergence of novel mutants resistant to multiple anti-HBV drugs. The importance and significance of the rtA181 mutations, including the novel mutant strain identified in this study, should be investigated further to develop more useful treatment strategies.

#### ACKNOWLEDGMENTS

This work was carried out at the Research Center for Molecular Medicine, Faculty of Medicine, Hiroshima University. We thank Hiromi Ishino, Asako Kozono, Kana Kunihiro, Rie Akiyama, Yoshiko Seo, Yoshiko Nakata, Eiko Miyoshi and Kiyomi Toyota for their excellent technical assistance.

This work was supported in part by grants-in-aid for scientific research and development from the Ministry of Education, Sports, Culture, and Technology and the Ministry of Health, Labor and Welfare.

#### REFERENCES

- Allen, M. I., M. Deslauriers, C. W. Andrews, G. A. Tipples, K. A. Walters, D. L. Tyrrell, N. Brown, L. D. Condearey, et al. 1998. Identification and characterization of mutations in hepatitis B virus resistant to lamivudine. *Hepatology* 27:1670–1677.
- Angus, P., R. Vaughan, S. Xiong, H. Yang, W. Delaney, C. Gibbs, C. Brosgart, D. Colledge, R. Edwards, A. Ayres, A. Bartholomeusz, and S. Locarnini. 2003. Resistance to adefovir dipivoxil therapy associated with the selection of a novel mutation in the HBV polymerase. *Gastroenterology* 125:292–297.
- Bozdagi, A. M., O. Uzunalioglu, A. R. Turkyilmaz, N. Aslan, O. Sezgin, T. Sahin, G. Bozdagi, K. Cinar, S. B. Pai, R. Pai, H. Bozkaya, S. Karayalcin, C. Yurdaydin, and R. F. Schinazi. 2003. YSDD: a novel mutation in HBV DNA polymerase confers clinical resistance to lamivudine. *J. Viral Hepat.* 10:256–265.
- Bruix, J., and J. M. Llovet. 2003. Hepatitis B virus and hepatocellular carcinoma. *J. Hepatol.* 39(Suppl. 1):S59–S63.
- Chang, T. T., R. G. Gish, S. J. Hadziyannis, J. Cianciara, M. Rizzetto, E. R. Schiff, G. Pastore, B. R. Bacon, T. Poynard, S. Joshi, K. S. Kleszczewski, A. Thiry, R. E. Rose, R. J. Colonna, and R. G. Hindes. 2005. A dose-ranging study of the efficacy and tolerability of entecavir in lamivudine-refractory chronic hepatitis B patients. *Gastroenterology* 129:1198–1209.
- Chayama, K., Y. Suzuki, M. Kobayashi, M. Kobayashi, A. Tsubota, M. Hashimoto, Y. Miyano, H. Koike, M. Kobayashi, I. Koida, Y. Arase, S. Saitoh, N. Murashima, K. Ikeda, and H. Kumada. 1998. Emergence and takeover of YMDD motif mutant hepatitis B virus during long-term lamivudine therapy and re-takeover by wild type after cessation of therapy. *Hepatology* 27:1711–1716.
- Chien, R. N., C. T. Yeh, P. N. Wang, M. C. Kuo, S. Y. Hsieh, L. Y. Shih, and Y. F. Liaw. 2004. Acute leukaemia in chronic hepatitis B patients with lamivudine therapy. *Int. J. Clin. Pract.* 58:1088–1091.
- Das, K., X. Xiong, H. Yang, C. E. Westland, C. S. Gibbs, S. G. Sarafianos, and E. Arnold. 2001. Molecular modeling and biochemical characterization reveal the mechanism of hepatitis B virus polymerase resistance to lamivudine (3TC) and emtricitabine (FTC). *J. Virol.* 75:4771–4779.
- Delaney, W. E., IV, H. Yang, C. E. Westland, K. Das, E. Arnold, C. S. Gibbs, M. D. Miller, and S. Xiong. 2003. The hepatitis B virus polymerase mutation rtV173L is selected during lamivudine therapy and enhances viral replication in vitro. *J. Virol.* 77:11833–11841.
- Doong, S. L., C. H. Tsai, R. F. Schinazi, D. C. Liotta, and Y. C. Cheng. 1991. Inhibition of the replication of hepatitis B virus in vitro by 2',3'-dideoxy-3'-thiacytidine and related analogues. *Proc. Natl. Acad. Sci. USA* 88:8495–8499.

11. Fung, S. K., P. Andreone, S. H. Han, K. Rajender Reddy, A. Regev, E. B. Keeffe, M. Hussain, C. Cursaro, P. Richtmyer, J. A. Marrero, and A. S. Lok. 2005. Adefovir-resistant hepatitis B can be associated with viral rebound and hepatic decompensation. *J. Hepatol.* 43:937-943.
12. Ganem, D., and A. M. Prince. 2004. Hepatitis B virus infection—natural history and clinical consequences. *N. Engl. J. Med.* 350:1118-1129.
13. Gunther, S., G. Sommer, F. Von Breunig, A. Iwanska, T. Kalinina, M. Sterneck, and H. Will. 1998. Amplification of full-length hepatitis B virus genomes from samples from patients with low levels of viremia: frequency and functional consequences of PCR-introduced mutations. *J. Clin. Microbiol.* 36:531-538.
14. Hadziyannis, S. J., N. C. Tassopoulos, E. J. Heathcote, T. T. Chang, G. Kitis, M. Rizzetto, P. Marcellin, S. G. Lim, Z. Goodman, J. Ma, S. Arterburn, S. Xiong, G. Currie, and C. L. Brosgart. 2005. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B. *N. Engl. J. Med.* 352:2673-2681.
15. Jacob, J. R., B. E. Korba, P. J. Cote, I. Toshkov, W. E. t. Delaney, J. L. Gerin, and B. C. Tennant. 2004. Suppression of lamivudine-resistant B-domain mutants by adefovir dipivoxil in the woodchuck hepatitis virus model. *Antiviral Res.* 63:115-121.
16. Lai, C. L., R. N. Chien, N. W. Leung, T. T. Chang, R. Guan, D. I. Tai, K. Y. Ng, P. C. Wu, J. C. Dent, J. Barber, S. L. Stephenson, D. F. Gray, et al. 1998. A one-year trial of lamivudine for chronic hepatitis B. *N. Engl. J. Med.* 339:61-68.
17. Lai, C. L., M. Rosmawati, J. Lao, H. Van Vlierberghe, F. H. Anderson, N. Thomas, and D. Dehertogh. 2002. Entecavir is superior to lamivudine in reducing hepatitis B virus DNA in patients with chronic hepatitis B infection. *Gastroenterology* 123:1831-1838.
18. Lai, C. L., J. Dienstag, E. Schiff, N. W. Leung, M. Atkins, C. Hunt, N. Brown, M. Woessner, R. Boehme, and L. Condreay. 2003. Prevalence and clinical correlates of YMDD variants during lamivudine therapy for patients with chronic hepatitis B. *Clin. Infect. Dis.* 36:687-696.
19. Lanford, R. E., L. Notvall, H. Lee, and B. Beames. 1997. Transcomplementation of nucleotide priming and reverse transcription between independently expressed TP and RT domains of the hepatitis B virus reverse transcriptase. *J. Virol.* 71:2996-3004.
20. Lanford, R. E., Y. H. Kim, H. Lee, L. Notvall, and B. Beames. 1999. Mapping of the hepatitis B virus reverse transcriptase TP and RT domains by transcomplementation for nucleotide priming and by protein-protein interaction. *J. Virol.* 73:1885-1893.
21. Lin, X., Z. H. Yuan, L. Wu, J. P. Ding, and Y. M. Wen. 2001. A single amino acid in the reverse transcriptase domain of hepatitis B virus affects virus replication efficiency. *J. Virol.* 75:11827-11833.
22. Nevens, F., J. Main, P. Honkoop, D. L. Tyrrell, J. Barber, M. T. Sullivan, J. Fevery, R. A. De Man, and H. C. Thomas. 1997. Lamivudine therapy for chronic hepatitis B: a six-month randomized dose-ranging study. *Gastroenterology* 113:1258-1263.
23. Ono, S. K., N. Kato, Y. Shiratori, J. Kato, T. Goto, R. F. Schinazi, F. J. Carrilho, and M. Omata. 2001. The polymerase L528M mutation cooperates with nucleotide binding-site mutations, increasing hepatitis B virus replication and drug resistance. *J. Clin. Investig.* 107:449-455.
24. Pai, S. B., A. M. Bozdayi, R. B. Pai, T. Beker, M. Sarioglu, A. R. Turkyilmaz, J. Grier, C. Yurdaydin, and R. F. Schinazi. 2005. Emergence of a novel mutation in the FLLA region of hepatitis B virus during lamivudine therapy. *Antimicrob. Agents Chemother.* 49:2618-2624.
25. Suzuki, Y., H. Kumada, K. Ikeda, K. Chayama, Y. Arase, S. Saitoh, A. Tsubota, M. Kobayashi, M. Koike, N. Ogawa, and K. Tanikawa. 1999. Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J. Hepatol.* 30:743-748.
26. Suzuki, Y., M. Kobayashi, K. Ikeda, F. Suzuki, Y. Arase, N. Akuta, T. Hosaka, S. Saitoh, M. Kobayashi, T. Someya, M. Matsuda, J. Sato, S. Watabiki, Y. Miyakawa, and H. Kumada. 2005. Persistence of acute infection with hepatitis B virus genotype A and treatment in Japan. *J. Med. Virol.* 76:33-39.
27. Tateno, C., Y. Yoshizane, N. Saito, M. Kataoka, R. Utoh, C. Yamasaki, A. Tachibana, Y. Soeno, K. Asahina, H. Hino, T. Asahara, T. Yokoi, T. Furukawa, and K. Yoshizato. 2004. Near completely humanized liver in mice shows human-type metabolic responses to drugs. *Am. J. Pathol.* 165:901-912.
28. Tatti, K. M., B. E. Korba, H. L. Stang, S. Peek, J. L. Gerin, B. C. Tennant, and R. F. Schinazi. 2002. Mutations in the conserved woodchuck hepatitis virus polymerase FLLA and YMDD regions conferring resistance to lamivudine. *Antiviral Res.* 55:141-150.
29. Tillmann, H. L., C. Trautwein, T. Bock, K. H. Boker, E. Jackel, M. Glowienka, K. Oldhafer, I. Bruns, J. Gauthier, L. D. Condreay, H. R. Raab, and M. P. Manns. 1999. Mutational pattern of hepatitis B virus on sequential therapy with famciclovir and lamivudine in patients with hepatitis B virus reinfection occurring under HBIg immunoglobulin after liver transplantation. *Hepatology* 30:244-256.
30. Tsubota, A., Y. Arase, S. Saitoh, M. Kobayashi, Y. Suzuki, F. Suzuki, K. Chayama, N. Murashima, K. Ikeda, M. Kobayashi, and H. Kumada. 2001. Lamivudine therapy for spontaneously occurring severe acute exacerbation in chronic hepatitis B virus infection: a preliminary study. *Am. J. Gastroenterol.* 96:557-562.
31. Tsuge, M., N. Hiraga, H. Takaishi, C. Noguchi, H. Oga, M. Imamura, S. Takahashi, E. Iwao, Y. Fujimoto, H. Ochi, K. Chayama, C. Tateno, and K. Yoshizato. 2005. Infection of human hepatocyte chimeric mouse with genetically engineered hepatitis B virus. *Hepatology* 42:1046-1054.
32. Villeneuve, J. P., D. Durantel, S. Durantel, C. Westland, S. Xiong, C. L. Brosgart, C. S. Gibbs, P. Parvaz, B. Werle, C. Trepo, and F. Zoulim. 2003. Selection of a hepatitis B virus strain resistant to adefovir in a liver transplantation patient. *J. Hepatol.* 39:1085-1089.
33. Wright, T. L., and J. Y. Lau. 1993. Clinical aspects of hepatitis B virus infection. *Lancet* 342:1340-1344.
34. Yeh, C. T., R. N. Chien, C. M. Chu, and Y. F. Liaw. 2000. Clearance of the original hepatitis B virus YMDD-motif mutants with emergence of distinct lamivudine-resistant mutants during prolonged lamivudine therapy. *Hepatology* 31:1318-1326.



CLINICAL RESEARCH STUDY

AJM Theme Issue: Infectious Disease

## Long-term Outcome of HBV Carriers with Negative HBe Antigen and Normal Aminotransferase

Kenji Ikeda, MD, PhD,<sup>a,b</sup> Yasuji Arase, MD,<sup>a,b</sup> Satoshi Saitoh, MD,<sup>a,b</sup> Masahiro Kobayashi, MD,<sup>a,b</sup> Takashi Someya, MD,<sup>a,b</sup> Tetsuya Hosaka, MD,<sup>a,b</sup> Hitomi Sezaki, MD,<sup>a,b</sup> Norio Akuta, MD, PhD,<sup>a,b</sup> Yoshiyuki Suzuki, MD, PhD,<sup>a,b</sup> Fumitaka Suzuki, MD, PhD,<sup>a,b</sup> Hiromitsu Kumada, MD, PhD<sup>a,b</sup>

<sup>a</sup>Department of Hepatology, Toranomon Hospital, Tokyo, Japan, and <sup>b</sup>Okinaka Memorial Institute for Medical Research, Tokyo, Japan.

### ABSTRACT

**PURPOSES:** To elucidate the incidence of hepatitis activation and hepatocellular carcinogenesis in patients with negative HBe antigen and normal aminotransferase, long-term observation was performed in a retrospective cohort.

**METHODS:** Among 116 consecutive patients with normal aminotransferase and negative HBe antigen at the time of liver biopsy, sequential frozen sera for initial 5 years were available for 95 patients. Hepatitis B virus (HBV) DNA assay (sensitivity >400 copy/mL) and aminotransferase were annually examined in the initial 5 years after biopsy.

**RESULTS:** Liver biopsy showed minimal hepatitis (F0) in 9, F1 in 53, F2 in 21, F3 in 6, and F4 or cirrhosis in 6. Initial HBV DNA concentration was low (less than 10<sup>4</sup> copies/mL) in 33, intermediate (10<sup>4</sup>-10<sup>6</sup>) in 53, and high (10<sup>6</sup> or more) in 9. Hepatitis activation rates with twice as high as normal aminotransferase at the end of the third year were 12.1% in the low DNA group, 43.4% in the intermediate group, and 66.7% in the high DNA group. Initial HBV DNA values were significantly associated with future increase in aminotransferase ( $P < .0001$ ). Initial DNA, history of aminotransferase elevation, and histological staging independently affected future hepatitis activation. Carcinogenesis rate in patients with and without high DNA of 10<sup>6</sup> copies/mL during the initial 3 years were 6.9% and 0%, respectively, at the end of the 5th year, and 11.5% and 1.8%, respectively, at the 10th year, ( $P = .021$ ).

**CONCLUSION:** Advanced stages of hepatitis were sometimes found in HBe antigen-negative, aminotransferase-normal HBV carriers. Serial HBV DNA assessment in early 3 years predicted future hepatitis activation and carcinogenesis. © 2006 Elsevier Inc. All rights reserved.

**KEYWORDS:** Hepatitis B virus; DNA; HBe-antigen; Carcinogenesis; Prognosis; Chronic hepatitis

HBe antigen loss and appearance of anti-HBe antibody are usually associated with normalization of aminotransferase in patients with chronic hepatitis,<sup>1-3</sup> and these changes have long been considered as a reliable clinical indicator of inactive and innocent state of chronic hepatitis B. However, the level of aminotransferase sometimes fluctuates under the state of negative HBe antigen in chronic hepatitis B, and there is occasional histological evidence of disease progression.<sup>4-6</sup> Because hepatitis B virus (HBV) DNA is usually

detected in the sera of these patients, mutant HBV has been confirmed to cause the hepatitis activity.<sup>7,8</sup> Moreover, hepatocellular carcinoma (HCC) sometimes occurs in patients with mild hepatitis and normal aminotransferase levels.<sup>9,10</sup> It is an important problem from the viewpoint of early detection of HCC that general practitioners and even hepatologists are sometimes unaware of those patients with negative HBe antigen and normal aminotransferase as high-risk subjects for HCC.

There is little information about how many patients with both normal alanine aminotransferase (ALT) and negative HBe antigen have active or advanced histologically confirmed hepatitis,<sup>11,12</sup> and how many patients in

Requests for reprints should be addressed to Kenji Ikeda, MD, PhD, Department of Hepatology, Toranomon Hospital, 2-2-2 Toranomon, Minato-ku, Tokyo, 105-8470, Japan.

E-mail address: ikedakenji@tora.email.ne.jp

whom ALT level increases over long-term follow-up. Recently, for evaluation of the activity of chronic hepatitis B, simple and sensitive HBV DNA assays using polymerase chain reaction (PCR) have replaced serologic examination of HBe antigen-antibody. The high sensitivity of DNA assays has contributed to our understanding of the pathogenetic role of the virus and provided useful information about hepatitis activity in individual patients.<sup>12-18</sup> An arbitrary serum DNA level of  $10^5$  copies/mL was proposed by the National Institute of Health Workshop for the differentiation of chronic hepatitis B from an inactive carrier state (HBe antigen-negative, persistently normal ALT).<sup>19</sup> Martinot-Peignoux et al<sup>20</sup> also reported that a cutoff DNA value of  $10^5$  copies/mL effectively discriminated inactive hepatitis in a follow-up study of 1 to 6 years. On the other hand, Manesis et al<sup>21</sup> suggested that a cutoff DNA level of  $3 \times 10^4$  copies/mL was more appropriate for differentiating inactive carriers from patients with HBeAg-negative chronic hepatitis B. Furthermore, Chu et al<sup>22</sup> concluded that a single cutoff DNA value could not differentiate inactive carriers from patients with HBeAg-negative chronic hepatitis after retrospective analysis of 165 Chinese patients. Previous studies indicated that the above data still did not provide sufficient information as to the longitudinal outcome of hepatitis activation and carcinogenesis, considering the fluctuating course of HBeAg-negative chronic hepatitis.<sup>19,22</sup> Here we report the results of our long-term study (median follow-up of 10.2 years) in patients with negative HBe antigen and normal alanine aminotransferase (ALT).

The purposes of this retrospective study were to elucidate the incidence of advanced liver disease in patients with negative HBe antigen and normal ALT value, and to estimate the future activation of hepatitis and carcinogenesis rate in those patients, using life-table method with multivariate analysis. We also analyzed whether the initial few years of biochemical and virological observation (annual measurements of ALT and HBV DNA concentration) could predict future outcome of the patients.

## PATIENTS AND METHODS

### Patients

Among 1048 patients who were diagnosed with HBV-related chronic hepatitis by peritoneoscopy and/or biopsy from 1974 to 2002 in our hospital, 116 patients were negative HBe antigen and had normal aminotransferase at the time of liver biopsy. Of these, 95 (81.9%) were thoroughly observed for 5 years or longer, and their data were available for analysis of HBV DNA concentration.

The initial sera of all these 95 patients were positive for HBs antigen, negative HBe antigen, and negative anti-HCV antibody. ALT concentration was also normal for 2 months or longer before admission to the hospital for examination of liver biopsy. The enrolled patients included 75 men and 20 women aged 21 to 67 years, with a median of 39 years. None of the patients received antiviral therapy before liver biopsy. All the clinical practices were performed in accordance with the ethical committee of the hospital and with general agreement of domestic medical insurance.

Table 1 shows demographic and laboratory data of the 95 patients. Family history of liver disease or HBV infection has been closely investigated and assessed to a relative in the fourth degree using each family line. Because all the patients had been negative for HBe antigen and normal ALT since the initial visit to our hospital, the time of seroconversion from HBe antigen to HBe antibody was not clear in most of the patients. HBV subtyping showed that genotype C was the most prevalent type (74.7%) in the cohort, the rate of which was slightly lower than that of Japanese patients with chronic hepatitis type B.<sup>23,24</sup> Two authors (K.I. and H.K.) assessed the pathology and translated the biopsy findings to the classification of Desmet et al.<sup>25</sup>

## CLINICAL SIGNIFICANCE

- Advanced stages of hepatitis should not be overlooked in patients with HBe antigen-negative, aminotransferase-normal HBV carriers.
- Serial DNA assessment in the early few years predicted future hepatitis activation and the risk of carcinogenesis.

**Table 1** Demography and Laboratory Data of 95 Patients with Positive HBs Antigen who Showed Normal Aminotransferase and Negative HBe Antigen at the Time of Biopsy

Data	
<b>Demographic data</b>	
Men:Women	75:20
Age (years - median, range)	39 (18-67)
HBe antigen positive	0 (0%)
HBV genotype A	3 (3.2%)
B	18 (18.9%)
C	71 (74.7%)
Others/undetectable	3 (3.3%)
Anti-HCV antibody positive	0 (0%)
Past alcohol intake of 500 kg or more	15 (15.8%)
History of blood transfusion	4 (4.2%)
<b>Laboratory data*</b>	
Aspartic aminotransferase (normal 6-38 IU)	26 (10-37)
Alanine aminotransferase (normal 6-50 IU)	28 (6-50)
Platelet count (normal $149-315 \times 10^3/\text{mm}^3$ )	185 (86-311)
ICG <sub>R15</sub> ** (normal, $\leq 10\%$ )	11 (3-33)
Albumin (normal, 3.9-5.2 g/dL)	4.3 (3.6-5.4)
Bilirubin (normal, $\leq 1.1$ mg/dL)	0.9 (0.2-1.9)
Prothrombin time (normal, $\geq 70\%$ )	97.5 (67-111)

\*Data are median (range).

\*\*ICG<sub>R15</sub>: Indocyanine green retention rate at 15 minutes.

## Follow-up Studies

After liver biopsy, follow-up examinations were conducted on a monthly to tri-monthly basis by monitoring biochemical and virological data. Although 27 (28.4%) patients started to receive interferon (IFN) therapy because of ALT elevation, none underwent lamivudine therapy during the course of the study. "Activation of hepatitis" or "flare-up of ALT" was defined as an increase of ALT value of twice or more of the upper limit of normal.

Imaging studies were performed annually in each patient and included computed tomography (CT) or ultrasound (US) after 1985. At least 2 imaging procedures were performed in those patients with cirrhosis. HCC was diagnosed by typical hypervascular characteristics on angiography in addition to certain features of CT and US. Pathological confirmation of surgically resected specimens or autopsy was made in 3 (60.0%) of 5 patients who developed HCC during the study.

A total of 17 patients were lost to follow-up after 5 years. The observation period ranged from 5.1 to 24.7 years, with a median of 10.2 years.

## Assays of HBV Markers

Serum HBsAg was measured by radioimmunoassay (Dainabot, Tokyo, Japan), and HBeAg and anti-HBe were determined by enzyme-linked immunosorbent assay (ELISA, Institute of Immunology) using commercially available kits. Anti-HCV antibody was analyzed using second-generation anti-HCV kits (enzyme-linked immunosorbent assay, Dainabot). HBV DNA was assayed using frozen sera stored at  $-80^{\circ}\text{C}$  and measured with Amplicor HBV Monitor kits (Roche Diagnostics Japan Co., Tokyo). The effective measurement of the concentration ranged from  $10^{2.6}$  (400) copies/mL to  $10^{7.6}$  (40 000 000) copies/mL. All measurements of HBV DNA were performed at the same time using the same assay kits. Quantitative HBV DNA was conducted at the time of liver biopsy, at the end of 6 months, and at the end of the 1st, 2nd, 3rd, and 5th years.

## Statistical Analysis

The chi-square test, Fisher's exact test, and Kruskal-Wallis test were used to analyze the relationship between HBV markers and the clinical course. Cumulative flare-up rate of hepatitis and carcinogenesis rate were calculated with the Kaplan-Meier method,<sup>26</sup> and differences in the rates were analyzed by the log-rank test. Multivariate Cox proportional hazard model<sup>27</sup> was adopted to analyze the prediction of future hepatitis activation and carcinogenesis. A *P*-value of less than .05 with 2-tailed test was considered significant. Data analysis was performed using the SPSS software version 11.<sup>28</sup>

## RESULTS

### Histological Findings

According to the classification of Desmet et al,<sup>25</sup> liver biopsy showed minimal hepatitis (F0) in 9 (9.5%), F1 in 53 (55.8%), F2 in 21 (22.1%), F3 in 6 (6.3%), and F4 or cirrhosis in 6 (6.3%). Occult cirrhosis with normal ALT and negative HBe antigen was found in 6.3%. Inflammatory activity was minimal (A0) in 18 (18.9%), A1 in 72 (75.8%), and A2 in 5 (5.3%).

### Reasons for Liver Biopsy

The reasons for liver biopsy in spite of "stable hepatitis" included a history of ALT elevation in the past (group A,  $n = 41$ ), patient's apprehension based on family history of advanced liver diseases (group B,  $n = 25$ ), possible advanced liver disease on ultrasonography (US) or liver function tests (group C,  $n = 8$ ), or simple desire for thorough examination including biopsy (group D,  $n = 21$ ). We also assessed the relationship between background features and reasons for undergoing liver biopsy (Table 2).

Advanced liver disease of F3 or F4 was present in 4 of 41 (9.8%) patients in group A, 4 of 25 (16.0%) in group B, 3 of 8 (37.5%) in group C, and 1 of 18 (5.6%) in group D. The median platelet count was the lowest in group C, which suggested that the group included more numbers of patients with advanced stages of the disease. The median HBV DNA concentration at the beginning was the highest in group A. HBV genotype A was exclusively found in group D and the incidence of genotype B was higher in groups A and D.

### Initial and Serial Assays of HBV DNA Concentration

The median DNA concentration at the time of liver biopsy was  $10^{4.4}$  copies/mL, with 25th percentile of  $10^{3.2}$ , and 75th percentile of  $10^{5.2}$  (range,  $<10^{2.6}$  to  $>10^{7.6}$ ).

The relationship between the initial HBV DNA values and serial measurements of DNA during the observation were assessed, classifying the initial DNA concentration into 4 groups: low DNA group ( $<10^4$  copies/mL), DNA concentration of  $10^4$ - $10^{4.9}$  copies/mL, DNA concentration of  $10^5$ - $10^{5.9}$  copies/mL, and high DNA of  $\geq 10^6$  copies (Figure 1). Among 33 patients with initial HBV DNA concentration of  $<10^4$  copies/mL, 29 (87.9%) showed a continuously low DNA of less than  $10^6$  copies/mL, and 4 (12.1%) showed an intermittent rise of the DNA to  $>10^6$  copies/mL. Among 34 patients with initial HBV DNA of  $10^4$  to  $10^{4.9}$  copies/mL, 25 (73.5%) showed persistently low DNA, and 9 (26.5%) an intermittent rise of the DNA. Of 18 patients with initial DNA of  $10^5$  to  $10^{5.9}$  copies/mL, 10 (55.5%) showed persistently low DNA course, and 8 (44.4%) an intermittent elevation of the DNA. Among 10 patients with initial DNA of  $\geq 10^6$  copies/mL, an intermittent rise of the DNA was found in 9 patients (90.0%) and persistently high DNA in one (10.0%). The initial HBV DNA concentration was significantly associated with the subsequent serial HBV DNA course ( $P < .0001$ ).

**Table 2** Demographic, Histological, Biochemical and Virological Findings According to the Reasons for Examination Including Liver Biopsy

	Reasons for Liver Biopsy			
	Group A	Group B	Group C	Group D
Number of patients	41	25	8	21
Male/Female	36/5	15/10	5/3	19/2
Age (years - median, range)	39 (18-61)	38 (30-67)	41 (30-57)	41 (21-58)
Family history of liver disease or HBV infection	18 (37.5%)	38 (100%)	5 (62.5%)	6 (28.6%)
Liver histology				
F0 (minimal)	1	5	0	3
F1	21	15	2	15
F2	15	1	3	2
F3	2	2	1	1
F4 (cirrhosis)	2	2	2	0
AST (IU/L) (median, range)	27 (17-37)	22 (10-37)	24 (16-28)	21 (14-34)
ALT (IU/L)	34 (18-50)	20 (6-48)	21 (12-48)	20 (8-50)
Platelet ( $\times 1000/\text{mm}^3$ )	169 (94-274)	185 (96-311)	156 (86-216)	196 (128-296)
ICG <sub>15</sub> (%)	12 (6-24)	9.5 (3-33)	9 (7-13)	12 (5-72)
Albumin (g/dL)	4.3 (3.6-5.1)	4.4 (3.9-5.4)	4.15 (3.9-5.4)	4.4 (3.7-4.9)
Bilirubin (mg/dL)	1.0 (0.2-1.5)	1.1 (0.4-1.8)	0.75 (0.3-1.1)	0.8 (0.3-1.9)
Prothrombin (%)	89 (67-111)	100 (70-108)	100 (76-101)	100 (73-101)
HBV DNA (Log <sub>10</sub> copy/ml) (median, 25-75 percentiles)	4.6 (4.2-5.3)	4.2 (3.1-5.4)	3.45 (<2.6-4.8)	3.9 (<2.6-4.8)
HBV genotype				
A	0	0	0	3
B	11	0	0	7
C	29	25	7	10
Others/undidentified	1	0	1	1

Group A: History of ALT elevation in the past.

Group B: Advanced liver disease in family members.

Group C: Possible advanced disease on ultrasonography or blood examination.

Group D: Simple desire for thorough examination

### Activation of Hepatitis and Its Prediction

Twenty-five patients showed twice as high as normal ALT concentration, and the other 27 patients showed activation of hepatitis with ALT elevation followed by IFN therapy. Consequently, a total of 52 patients (54.7%) showed an abnormal ALT concentration during a median observation period of 10.2 years. The cumulative rate of hepatitis activation was 34.7% at the end of the 3rd year, 45.4% at the 5th year, and 55.7% at the 10th year.

The incidence of hepatitis activation was assessed in the following 4 groups according to the initial HBV DNA concentration:  $<10^4$  copies/mL ( $n = 33$ ),  $10^4$  to  $10^{4.9}$  ( $n = 34$ ),  $10^5$  to  $10^{5.9}$  ( $n = 18$ ), and  $\geq 10^6$  ( $n = 10$ ). Cumulative hepatitis flare-up rates in each group were 12.0%, 41.2%, 44.4%, and 70.0%, respectively, at the end of the 3rd year; 12.0%, 61.8%, 55.5%, and 80.0%, respectively, at the 5th year; and 18.6%, 70.3%, 80.0%, and 80.0%, respectively, at the 10th year (Figure 2). The initial DNA load correlated significantly with future hepatitis activation rate ( $P < .0001$ ). Average DNA concentration during the initial 5 years was closely associated with hepatitis activation for the period: 52 patients with ALT elevation had a high average DNA value of  $10^{3.3}$  (median, range:  $<10^{2.6}$ - $10^{5.3}$ ), and 43 patients without ALT elevation showed  $10^{4.8}$  (median, range:  $10^{2.9}$ - $>10^{7.6}$ ).

The hepatitis activation rates in groups A, B, C, and D were 70.7%, 36.0%, 25.0%, and 14.3%, respectively, at the end of the 5th year; and 81.7%, 45.1%, 25.0%, and 26.7%, respectively, at the end of the 10th year (Figure 3). The ALT flare-up rate in patients of group A was the highest ( $P < .0001$ ).

Multivariate proportional hazard analysis disclosed that initial HBV DNA concentration, a past history of ALT elevation, and histological staging were significantly associated with reactivation of hepatitis in the follow-up (Table 3). Patients with HBV DNA of  $10^4$  to  $10^{5.9}$  copy/mL had a high hazard ratio of 5.73, and those with  $10^6$  copy/mL or higher showed a hazard ratio of 10.43, compared with those with lower DNA concentration  $<10^4$ . Past history of ALT elevation also was associated with hepatitis activation with a hazard ratio of 3.62. Histological staging of F2 or higher also showed a high risk for carcinogenesis with a hazard ratio of 3.06.

### Relationship between HBV DNA in First 2 Years and Hepatitis Activation

The prediction of hepatitis activation was analyzed using HBV DNA level in the first 2 years (DNA measurements at the time of observation, 6th, 12, and 24th month). Using various cutoff values of DNA, patients were classified into




		Patterns of serial of HBV DNA for 5 years		
Initial HBV DNA (copies/ml)		lower than $10^6$ persistently	$10^6$ or higher intermittently	$10^6$ or higher persistently
				
< $10^4$	(N=33)	29 (87.9%)	4 (12.1%)	0
$10^4$ - $10^{4.9}$	(N=34)	25 (73.5%)	9 (26.5%)	0
$10^5$ - $10^{5.9}$	(N=18)	10 (55.6%)	8 (44.4%)	0
$\geq 10^6$	(N=10)	0	9 (90%)	1 (10%)

Figure 1 Relationship between initial HBV DNA values and serial DNA during the initial 5 years.

low DNA group (persistently low) and high DNA group (a high value at least once during the observation period).

When patients were divided using a cutoff value of  $10^6$  copies/mL (Figure 4a), hepatitis activation rates in low DNA group ( $n=69$ ) and high DNA group ( $n=26$ ) were 27.5% and 53.8%, respectively, at the end of the 3rd year; 37.7% and 65.4%, respectively, at the 5th year; and 46.0% and 79.2%, respectively, at the 10th year. The hepatitis activation rate in the high DNA group was significantly higher than that of the low DNA group ( $P=.0014$ ). If DNA

concentration remained  $<10^6$  copies/mL for 2 years, 53.6% (37/69) of the patients did not show flare-up of ALT during the observation period.

Next, when patients were divided using a cutoff value of  $10^5$  copies/mL (Figure 4b), hepatitis activation rates in the low DNA group ( $n=47$ ) and high DNA group ( $n=48$ ) were 14.9% and 54.2%, respectively, at the end of the 3rd year; 23.4% and 66.7%, respectively, at the 5th year; and 30.8% and 79.4%, respectively, at the 10th year. Hepatitis activation rate in the high DNA group was significantly





Figure 2 Cumulative hepatitis activation rates in the 4 patient groups according to initial HBV DNA values:  $\text{Log}_{10}$  HBV DNA of  $<4.0$ ,  $4.0$ - $4.9$ ,  $5.0$ - $5.9$ , and  $\geq 6$ .





**Figure 3** Cumulative hepatitis activation rates in subgroup A, B, C, and D: Patients in group A had a history of aminotransferase elevation in the past, those in group B had apprehension of their own diseases based on a significant family history, those in group C were suspected of having an advanced liver disease on ultrasonography or liver function tests, and those in group D had a simple desire for thorough examination with biopsy.

higher than that of the low DNA group ( $P < .0001$ ). If DNA concentration remained  $< 10^5$  copies/mL for 2 years, 70.2% (33/47) of the patients did not show flare-up of ALT during the observation period.

Similarly, when patients were divided using a cutoff value of  $10^4$  copies/mL (Figure 4c), hepatitis activation rates in the low DNA group ( $n = 24$ ) and high DNA group ( $n = 71$ ) were 4.2% and 45.1%, respectively, at the end of the 3rd year; 4.2% and 59.1%, respectively, at the 5th year; and 8.3% and 72.1%, respectively, at the 10th year. Hepatitis activation rate in the high DNA group was significantly higher than that of the low DNA group ( $P < .0001$ ). If DNA concentration remained  $< 10^4$  copies/mL for 2 years, 91.7%

(22/24) of the patients did not show flare-up of ALT during the observation period.

A receiver-operating characteristic (ROC) curve was generated to assess the best cutoff value of HBV DNA for the prediction of future hepatitis activation (Figure 5). Cutoff value of  $10^{6.0}$  or higher predicted 86.0% of future hepatitis activation, but persistently lower value than  $10^{6.0}$  predicted only 36.5% of stable disease. On the contrary, cutoff value of  $10^{4.0}$  or higher predicted only 51.2% of future ALT elevation, but constantly lower value than  $10^{4.0}$  predicted 96.2% of stable hepatitis.

### Hepatocellular Carcinogenesis and Its Prediction


During a median observation period of 10.2 years, 5 patients (5.3%) developed HCC. Among the 5 patients who developed HCC, one had low HBV DNA concentration of  $\leq 10^4$  copies/mL, 3 had intermediate value of  $10^4$  to  $10^6$ , and one had high DNA concentration of  $\geq 10^6$  copies/mL. Among 90 patients who did not develop HCC, 32 had low DNA, 50 had intermediate viral load, and 8 had high concentration ( $P = .62$ ). The cumulative carcinogenesis rates in patients with and without high DNA concentration of  $\geq 10^6$  copies/mL during the initial 3 years were 6.9% and 0%, respectively, at the end of the 5th year, and 11.5% and 1.8%, respectively, at the 10th year (Figure 6).

Those patients with high DNA concentration of  $\geq 10^6$  copies/mL at least once during the initial 3 years had a significantly higher carcinogenesis rate than those without DNA elevation ( $P = .021$ ). Average DNA concentration during the initial 5 years also was associated with hepatitis

**Table 3** Factors Associated with Future Hepatitis Activation in Patients with Positive HBs Antigen who Showed Normal Aminotransferase and Negative HBe Antigen at the Time of Biopsy

Factor	Category	Hazard Ratio	(95% CI)	P Value
Initial HBV DNA	0: $-10^{3.9}$	1		
	1: $10^4$ - $10^{5.9}$	5.73	(2.36-13.9)	<.0001
	2: $10^{6.0}$ -	10.43	(3.44-31.60)	<.0001
History of ALT elevation	0: no	1		
	1: yes	3.62	(1.97-6.67)	<.0001
Histological staging	0: F0-F1	1		
	1: F2-F4	3.06	(1.73-5.42)	<.0001





**Figure 6** Cumulative hepatocellular carcinogenesis rates in patients with and without high HBV DNA concentration during the initial 3 years.

hepatitis activity often fluctuates with significant changes of DNA concentration in the patients, even sensitive DNA assays demonstrate an obvious limitation from a single time measurement.<sup>22</sup> Our retrospective study, therefore, analyzed the chronological changes in DNA and biochemistry in a patient cohort over a period of more than 5 years (median, 10.2 years).

We found all patients with negative HBeAg and normal ALT had a detectable value of serum HBV DNA by PCR assay, and that the DNA value correlated well with high ALT concentration and the incidence of future ALT elevation, in agreement with other reports.<sup>19-22,32-35</sup> A few studies<sup>19,20,36</sup> reported that a cutoff value of  $10^5$  copies/mL was practical in discriminating chronic hepatitis from asymptomatic carrier state, whereas Manesis et al<sup>21</sup> reported that a cutoff DNA level of  $3 \times 10^4$  copies/mL was more appropriate for the same purpose. In forecasting future ALT elevation in a narrower patient cohort with negative HBe antigen and normal ALT concentrations, however, a cutoff value of  $10^4$  copies/mL effectively predicted reactivation of hepatitis. The efficacy of the cutoff value of  $10^4$  copies/mL was true in using both the initial DNA value at a single-time assay and multiple assays during 2-year follow-up. From the longitudinal data, only 8.3% of patients with persistently low DNA of  $< 10^4$  copies/mL in the initial 2 years showed ALT elevation during a median of 10.2 years of observation.

The fate of patients regarding future DNA elevation and ALT fluctuation also depends on the reason for undergoing invasive diagnostic procedures including liver biopsy. Patients of group A (patients with history of ALT elevation) showed the highest initial DNA concentration and future ALT elevation rate. Those of group B (patients with family history of advanced disease) showed the second highest value of initial DNA and high incidence of ALT elevation.

Although we analyzed more than 80% of consecutive patients in our hospital, the background features of patients should be taken into account in the study of HBV-related disease, when a tertiary medical center performs a retrospective cohort study using a selected patient group.

Among the stable patients with negative HBe antigen and normal ALT, HCC occurred in 5 patients (5.3%) during a median of 10.2 years. As was reported previously,<sup>33,37,38</sup> hepatocarcinogenesis was significantly associated with persistently high HBV DNA. In this cohort study, patients with DNA of  $\geq 10^6$  copies/mL had a significantly higher carcinogenesis rate than those with low DNA. The cutoff concentration of DNA of  $10^6$  copies/mL was consistent with other cohorts with cirrhosis<sup>33,37,38</sup> in the prediction of carcinogenesis. The reason for the higher cutoff value for carcinogenesis than that for hepatitis reactivation might be explained by differences in the pathogenetic processes. Although hepatocarcinogenesis in these patients may well be associated with DNA concentration severity of the liver disease, or merely persistence of high ALT concentration, general practitioners or physicians should be aware of the risk of HCC development in these stable patients.

The precise role of HBV in oncogenesis is still unknown, but increasing evidence indicates that the virus plays an important role in the development of HCC. Because the disease activity and carcinogenic potency can change significantly during the course of HBV-related liver disease, a longitudinal analysis is desirable for the study of carcinogenesis from HBV-related cirrhosis. Future studies should be aimed, therefore, at defining the basic oncogenic mechanisms and roles of serum DNA concentration in carcinogenesis in patients with HBV infection.

In conclusion, advanced stages of hepatitis should not be overlooked in patients with HBe antigen-negative, aminotransferase-normal HBV carriers. Serial DNA assessment

in the early few years predicted future hepatitis activation and the risk of carcinogenesis.

## ACKNOWLEDGMENT

This study was supported in part by a research grant from the Ministry of Health, Labour and Welfare, Japan.


## References

1. Realdi G, Alberti M, Rugge M, et al. Seroconversion from hepatitis B e antigen to anti-HBe in chronic hepatitis B virus infection. *Gastroenterology*. 1980;79:195-199.
2. Hoofnagle JH, Dusheiko GM, Seeff LB, Jones EA, Waggoner JG, Bales ZB. Seroconversion from hepatitis B e antigen to antibody in chronic type B hepatitis. *Ann Intern Med*. 1981;94:744-748.
3. Liaw YF, Chu CM, Su IJ, Huang MJ, Lin DY, Chang-Chien CS. Clinical and histological events preceding hepatitis B e antigen seroconversion in chronic type B hepatitis. *Gastroenterology*. 1983;84:216-219.
4. Bonino F, Rosina F, Rizzetto M, et al. Chronic hepatitis in HBs Ag carriers with serum HBV-DNA and anti-HBe. *Gastroenterology*. 1986;90:1268-1273.
5. Fattovich G, Rugge M, Brollo L, et al. Clinical, virologic and histologic outcome following seroconversion from HBeAg to anti HBe in chronic hepatitis type B. *Hepatology*. 1986;6:167-172.
6. Fattovich G, Brollo L, Alberti A, Pontisso P, Giustina G, Realdi G. Long-term follow up of anti HBe-positive chronic active hepatitis B. *Hepatology*. 1988;8:1651-1654.
7. Carman WF, Jacyna MR, Hadziyannis S, et al. Mutation preventing formation of hepatitis B e antigen in chronic HBV infection. *Lancet*. 1989;ii:588-591.
8. Brown JL, Carman WF, Thomas HC. The hepatitis B virus. *Baillieres Clin Gastroenterol*. 1990;4:721-747.
9. Omata M. Treatment of chronic hepatitis B infection (Editorial). *N Engl J Med*. 1998;339:114-115.
10. Hsu YS, Chien RN, Yeh CT, et al. Long-term outcome after spontaneous HBeAg seroconversion in patients with chronic hepatitis B. *Hepatology*. 2002;35:1522-1527.
11. Ruiz-Moreno M, Otero M, Millan A, et al. Clinical and histological outcome after hepatitis B e antigen to antibody seroconversion in children with chronic hepatitis B. *Hepatology*. 1999;29:572-575.
12. Lindh M, Horal P, Dhillon AP, Norrkans G. Hepatitis B virus DNA levels, precore mutations, genotypes and histological activity in chronic hepatitis B. *J Viral Hepatol*. 2000;7:258-267.
13. Niituma H, Ishii M, Miura M, Kobayashi K, Toyota T. Low level hepatitis B viremia detected by polymerase chain reaction accompanies the absence of HBe antigenemia and hepatitis B virus carriers. *Am J Gastroenterol*. 1997;92:119-123.
14. Ikeda K, Saitoh S, Suzuki Y, et al. Relationship of hepatocellular carcinogenesis with precore mutant virus and serum hepatitis B virus DNA concentration—a longitudinal analysis of patients with cirrhosis. *Hepatol Res*. 1998;10:142-155.
15. Gerken G, Gomes J, Lampertico P, et al. Clinical evaluation and applications of the Amplicor HBV Monitor™ test, a quantitative HBV DNA PCR assay. *J Virol Methods*. 1998;74:155-165.
16. Fujiwara K, Yokosuka O, Ehata T, et al. The two different states of hepatitis B virus DNA in asymptomatic carriers: HBe-antigen-positive versus anti-HBe-positive asymptomatic carriers. *Dig Dis Sci*. 1998;43:368-376.
17. Loeb KR, Jerome KR, Goddard J, Huang ML, Cent A, Corey L. High-throughput quantitative analysis of hepatitis B virus DNA in serum using the TaqMan fluorogenic detection system. *Hepatology*. 2000;32:626-629.
18. Kessler HH, Preininger S, Stelzl E, et al. Identification of different states of hepatitis B virus infection with a quantitative PCR assay. *Clin Diagn Lab Immunol*. 2000;7:298-300.
19. Lok AS, Heathcote EJ, Hoofnagle JH. Management of hepatitis B 2000, summary of a workshop. *Gastroenterology*. 2001;120:1828-1853.
20. Martinot-Peignoux M, Boyer N, Colombat M, et al. Serum HBV DNA levels and liver histology in inactive HBsAg carriers. *J Hepatol*. 2002;36:543-546.
21. Manesis EK, Papatheodoridis GV, Hadziyannis SJ. Serum HBV DNA levels in inactive HBsAg carriers. *Gastroenterology*. 2002;122:2092-2093.
22. Chu CJ, Hussain M, Lok ASF. Quantitative serum HBV DNA levels during different stages of chronic hepatitis B infection. *Hepatology*. 2002;36:1408-1415.
23. Orito E, Ichida T, Sakugawa H, et al. Geographic distribution of hepatitis B virus (HBV) genotype in patients with chronic HBV infection in Japan. *Hepatology*. 2001;34:590-594.
24. Kobayashi M, Arase Y, Ikeda K, et al. Clinical characteristics of patients infected with hepatitis B virus genotypes A, B, and C. *J Gastroenterol*. 2002;37:35-39.
25. Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Sheuer PJ. Classification of chronic hepatitis: diagnosis, grading, and staging. *Hepatology*. 1994;19:1513-1520.
26. Kaplan EL, Meier P. Nonparametric estimation for incomplete observation. *J Am Stat Assoc*. 1958;53:457-481.
27. Cox DR. Regression models and life tables. *J R Stat Soc*. 1972;34:248-275.
28. SPSS Inc. *SPSS for Windows Version 11.0 Manual*. Chicago, IL: SPSS Inc.; 2001.
29. Ranki M, Schatzl HM, Zachoval R, Uusi-Oukari M, Lehtvaara P. Quantification of hepatitis B virus DNA over a wide range from serum studying viral replicative activity in response to treatment and in recurrent infection. *Hepatology*. 1995;21:1492-1499.
30. Noborg U, Gusdal A, Pisa EK, Hedrum A, Lindh M. Automated quantitative analysis of hepatitis B virus DNA by using the Cobas Amplicor HBV Monitor test. *J Clin Microbiol*. 1999;37:2793-2797.
31. Pawlowsky JM, Bastie A, Hezode C, et al. Routine detection and quantification of hepatitis B virus DNA in clinical laboratories: performance of three commercial assays. *J Virol Methods*. 2000;85:11-21.
32. Mangia A, Chung YH, Hoofnagle JH, Birkenmeyer L, Mushahwar I, Di Bisceglie AM. Pathogenesis of chronic liver disease in patients with chronic hepatitis B virus infection without serum HBeAg. *Dig Dis Sci*. 1996;41:2447-2452.
33. Ikeda K, Saitoh S, Suzuki Y, et al. Interferon decreases hepatocellular carcinogenesis in patients with cirrhosis caused by hepatitis B virus—a pilot study. *Cancer*. 1998;82:827-835.
34. Sung JJ, Chan HL, Wong ML, et al. Relationship of clinical and virological factors with hepatitis activity in hepatitis B e antigen-negative chronic hepatitis B virus-infected patients. *J Viral Hepatol*. 2002;9:229-234.
35. Chan HL, Tsang SW, Liew CT, et al. Viral genotype and hepatitis B virus DNA levels are correlated with histological liver damage in HBeAg-negative chronic hepatitis B virus infection. *Am J Gastroenterol*. 2002;97:406-412.
36. Sakugawa H, Nakasone H, Nakayoshi T, et al. Correlation between serum aminotransferase activity and virus load among patients with chronic liver disease type B. *Hepatol Res*. 2001;21:159-168.
37. Ikeda K, Arase Y, Kobayashi M, et al. Consistently low hepatitis B virus DNA saves patients from hepatocellular carcinogenesis in HBV-related cirrhosis. *Intervirology*. 2003;43:96-104.
38. Ikeda K, Kobayashi M, Saitoh S, et al. Significance of hepatitis B-virus DNA clearance and its early prediction in hepatocellular carcinogenesis in patients with cirrhosis undergoing interferon therapy—a long-term follow-up of a pilot study. *J Gastroenterol Hepatol*. 2005;20:95-102.

# Anticarcinogenic Impact of Interferon on Patients with Chronic Hepatitis C: A Large-Scale Long-Term Study in a Single Center

Kenji Ikeda Yasuji Arase Satoshi Saitoh Masahiro Kobayashi  
Takashi Someya Tetsuya Hosaka Hitomi Sezaki Norio Akuta  
Fumitaka Suzuki Yoshiyuki Suzuki Hiromitsu Kumada

S. Karger  
Medical and Scientific Publishers  
Basel · Freiburg · Paris ·  
London · New York ·  
Bangalore · Bangkok ·  
Singapore · Tokyo · Sydney



Printed in Switzerland on acid-free paper by Reinhardt Druck, Basel

## Anticarcinogenic Impact of Interferon on Patients with Chronic Hepatitis C: A Large-Scale Long-Term Study in a Single Center

Kenji Ikeda Yasuji Arase Satoshi Saitoh Masahiro Kobayashi  
Takashi Someya Tetsuya Hosaka Hitomi Sezaki Norio Akuta  
Fumitaka Suzuki Yoshiyuki Suzuki Hiromitsu Kumada

Department of Gastroenterology, Toranomon Hospital, and Okinaka Memorial Institute for Medical Research, Tokyo, Japan

### Key Words

Cirrhosis · Fibrosis · Hepatitis C virus · Hepatocellular carcinoma · Interferon

### Abstract

**Background:** The anticarcinogenic capacity of interferon (IFN) was assessed in a cohort of Japanese patients with chronic hepatitis C en masse. **Patients and Methods:** The rate of hepatocarcinogenesis was analyzed in 2,166 patients with chronic hepatitis C, of whom 1,654 had received IFN therapy while 512 had not. **Results:** Crude rates of hepatocarcinogenesis in treated and untreated patients were 2.6 and 4.6% at the end of the 5th year, 5.8 and 12.7% at the 10th year and 13.9 and 23.9% at the 15th year (after completion of IFN therapy for those treated) ( $p < 0.001$ ). IFN decreased the hazard ratio of carcinogenesis to 0.42 ( $p < 0.001$ ) in multivariate analysis with adjustments for significant covariates including fibrotic stage,  $\gamma$ -glutamyl transpeptidase level, gender, platelet count and age. Among the 1,654 patients treated with IFN, 606 (36.6%) achieved persistent loss of hepatitis C virus (HCV) RNA and an additional 266 (16.1%) gained normal levels of alanine aminotransferase without loss of HCV RNA for 6 months or longer after the completion of IFN therapy. Cumulative rates of hepatocarcinogenesis in sustained virological responders and biochemical responders were 1.4 and 2.0% at the end of the 5th year,

1.9 and 3.6% at the 10th year and 1.9 and 7.5% at the 15th year, respectively. The hazard ratio of sustained virological response was 0.10 ( $p < 0.001$ ), and that of biochemical response was 0.12 ( $p < 0.001$ ). Normalization of aminotransferase levels after IFN therapy without loss of serum HCV RNA decreased hepatocarcinogenesis. **Conclusion:** IFN significantly decreased the rate of hepatocarcinogenesis in patients with chronic hepatitis C as a whole in Japan, even in those who fail to clear HCV RNA from serum.

Copyright © 2006 S. Karger AG, Basel

### Introduction

In most developed countries, hepatitis B virus (HBV) and hepatitis C virus (HCV) infections account for the great majority of hepatocellular carcinoma (HCC), with incidence rates dependent on the regional prevalence of these hepatitis viruses. HCV-associated HCC typically develops through a sequence of events that progress from chronic inflammation through fibrosis and cirrhosis accompanying dysplasia and ultimately to HCC. In our previous cohort study on Japanese patients with HCV-related cirrhosis [1], cumulative rates of developing HCC at 5, 10 and 15 years were 21.5, 53.2 and 75.2%, respectively. According to our observations of untreated patients with chronic hepatitis C [2], rates of hepatocarcino-

### KARGER

Fax +41 61 306 12 34  
E-Mail karger@karger.ch  
www.karger.com

© 2006 S. Karger AG, Basel  
0300-5526/06/0492-0082\$23.50/0

Accessible online at:  
www.karger.com/int

Kenji Ikeda, MD  
Department of Gastroenterology, Toranomon Hospital  
Toranomon 2-2-2, Minato-ku  
Tokyo 105-8470 (Japan)  
Tel. +81 44 877 5111, Fax +81 44 860 1623, E-Mail ikedakenji@tora.email.ne.jp

genesis at 5, 10 and 15 years were estimated to be 4.8, 13.6 and 26.0%, respectively. The life expectancy of patients with HCV-related cirrhosis is largely influenced by the development of HCC in the clinical course. As the efficacy of radically curative therapies for HCC remains limited at best, and since a severe organ shortage does not provide with sufficient chances for liver transplantation, the prevention of HCC in patients with chronic liver disease is of great importance at the present.

Interferon (IFN) is effective in eliminating HCV and reducing serum levels of alanine aminotransferase (ALT) in some patients with chronic hepatitis C [3–6]. Reduced incidence of HCC in HCV-associated cirrhosis by IFN has been reported by many investigators including ourselves [7–14]; only a few studies have failed to find its benefit [15, 16]. However, many published studies had shortcomings in the study design, in terms of pooling patients who received IFN in diverse regimens, relatively short periods of follow-up despite a long incubation period of HCC, large numbers of dropouts and retrospective studies with historical controls. Moreover, almost all studies evaluated the activity of IFN to prevent HCC by comparing responders and nonresponders to the treatment. Due to difficulties in studying patients with chronic hepatitis C, a number of nonrandomized studies examined the effect of IFN on the incidence of hepatocarcinogenesis [17–20]. With invariable limitations in study design and interpretation of the results, these studies have disclosed useful information as regards the treatment of patients with chronic HCV infection.

In order to evaluate whether IFN can reduce the rate of carcinogenesis in patients with chronic hepatitis C, we compared 1,654 patients with IFN therapy with 512 patients without treatment in a single clinical center, who were adjusted for background features by the multivariate analysis. Therefore, the principal aims of our study were to show the role of IFN in preventing HCC in chronic hepatitis type C en masse and to establish the extent to which IFN decreases the rate of carcinogenesis as a sequel to chronic hepatitis C in a society.

## Patients and Methods

### Study Population

A total of 2,166 patients with chronic hepatitis were examined, whose initial sera tested negative for hepatitis B surface antigen by radioimmunoassay (Ausria, Dainabot, Tokyo, Japan) and positive for anti-HCV by the second-generation enzyme-linked immunosorbent assay (Dainabot); anti-HCV was tested in sera that had been stored frozen at  $-80^{\circ}\text{C}$ . They included 1,421 men and 745

women aged 14–78 with a median of 50 years. They were all diagnosed with chronic hepatitis by liver biopsy with or without peritoneoscopy between 1970 and 2000 at the Department of Gastroenterology in Toranomon Hospital, Tokyo, Japan. Patients who had possibly developed HCC already at the time of diagnosis of hepatitis were strictly excluded from the study. In order to exclusively investigate hepatocarcinogenesis in HCV-related cirrhosis, patients coinfecting with HBV were excluded.

Among the 2,166 patients with HCV-related hepatitis, 1,654 (76.4%) received IFN therapy, mostly since 1987 when IFN was available in Japan; new antivirals or anticarcinogenic treatments of viral cirrhosis, except for IFN, were not introduced in 1987 or thereafter in Japan. The remaining 512 patients did not receive IFN or any other antiviral therapies. This is a retrospective cohort study with historical controls composed of patients before 1987 and those who refused or could not receive IFN for various reasons since 1987.

### Background and Laboratory Findings

Table 1 shows demographic profiles and laboratory data for the 1,654 patients treated with IFN and the 512 without receiving IFN since they were diagnosed with chronic hepatitis. There were more males, with a median age 3 years lower in treated than in nontreated patients. There were 299 treated patients (18.1%) with a history of alcohol intake  $\geq 500$  kg until the diagnosis of chronic hepatitis (corresponding to daily consumption of 3,000 ml of beer or 300 ml of whiskey for 20 years) and 113 (22.1%) untreated patients ( $p < 0.001$ ). Because IFN was introduced to our hospital in 1987, the observation period was significantly shorter in the treated than in untreated patients (median 10.4 vs. 12.3 years;  $p < 0.0001$ ).

Although all patients tested positive for HCV RNA during their clinical courses, tests for the concentration of HCV RNA in the initial serum was possible in 1,863 (86.5%) patients. HCV genotypes were analyzed by the serological typing method with a commercial kit (Kokusai Diagnostic Corporation, Kobe, Japan) in which the serological group 1 represented genotypes 1a and 1b, and group 2 stood for 2a and 2b genotypes. HCV in the serological group 2 was significantly more frequent in patients with IFN treatment than in those without. Concentration of HCV RNA was determined in the initial sera from 1,873 (86.5%) patients by the competitive polymerase chain reaction (PCR) method with the HCV probe assay (Chiron Corp., Emeryville, Calif., USA) or by PCR with Amplicor HCV Monitor kits (Roche Diagnostics Japan Co., Tokyo, Japan). High concentration of HCV ( $\geq 10^6$  copies/ml by the competitive PCR or  $\geq 10^6$  equivalents/ml by the HCV probe assay) was significantly more frequent in untreated than in treated patients ( $p < 0.0001$ ). The stage of hepatic fibrosis was not different between the two groups.

### Interferon Treatment and Judgment of the Effect

A total of 1,654 patients underwent IFN therapy in one or more treatment courses: 1,358 patients (82.1%) received IFN once, 240 patients (14.5%) twice, and the remaining 56 patients (3.4%) three times or more. Initial treatment was performed with natural or recombinant IFN- $\alpha$  ( $n = 1,238$ ), natural IFN- $\beta$  ( $n = 386$ ) or both ( $n = 30$ ). Regimens of IFN were variable: 926 (56.0%) patients received IFN 6–9 million units (MU) daily for 8 weeks, followed by 2 or 3 times per week for 16 weeks; 329 (20.0%) received IFN 6–9 MU daily for 2–4 weeks, followed by 3 times per week for 20–22 weeks; 185 (11.2%) underwent a short-course therapy with IFN

**Table 1.** Patient profiles and laboratory data at the diagnosis of chronic hepatitis

Factors	Interferon therapy		p value
	yes (n = 1,654)	no (n = 512)	
Male	1,110 (67.1%)	311 (60.7%)	0.024
Age, years	50 (16–72)	53 (21–78)	<0.001
History of transfusion	607 (36.7%)	229 (44.7%)	0.001
Family member with liver disease	426 (25.8%)	140 (27.3%)	0.47
Alcohol intake $\geq$ 500 kg	299 (18.1%)	113 (22.1%)	0.044
Observation period, year	10.4 (0.1–33.6)	12.3 (0.1–33.6)	<0.001
Laboratory data			
ALT, IU/l	63 (4–1,266)	67 (4–704)	0.098
AST, IU/l	106 (9–1,660)	96 (12–832)	0.0001
$\gamma$ -GTP, IU/ml	62 (6–1,118)	70 (3–850)	0.39
Platelet counts, $\times$ 1,000/mm <sup>3</sup>	169 (27–433)	165 (35–560)	0.091
ICG R <sub>15</sub> , %	14 (1–90)	16 (1–95)	0.003
AFP, ng/ml	4 (1–90)	5 (1–1,180)	0.42
HCV serological group			
Group 1, genotypes 1a/1b	1,021 (66.1%)	259 (81.4%)	<0.0001
Group 2, genotypes 2a/2b	488 (31.6%)	48 (15.1%)	
Undetermined	36 (2.3%)	11 (3.5%)	
HCV RNA concentration			
High <sup>a</sup>	937 (58.4%)	191 (71.3%)	<0.0001
Low <sup>b</sup>	668 (41.6%)	77 (28.7%)	
Histological stage of hepatitis			
F1, slight fibrosis	1,029 (62.2%)	298 (58.2%)	0.10
F2/F3, moderate/severe fibrosis	625 (37.8%)	214 (41.6%)	

AST = Aspartate aminotransferase; AFP =  $\alpha$ -fetoprotein; ICG R<sub>15</sub> = retention of indocyanine green at 15 min.

<sup>a</sup> HCV RNA concentration  $\geq$  10<sup>6</sup> copies/ml by the competitive PCR or  $\geq$  10<sup>6</sup> equivalents/ml by the HCV probe assay.

<sup>b</sup> HCV RNA concentrations less than high concentrations.

daily for 4–8 weeks; 128 (7.7%) were administered with intermittent IFN 3 times per week for 24 weeks; 72 (4.4%) had a prolonged course of IFN for 8–36 months; 8 (0.5%) received IFN- $\beta$  6 MU daily for 6–18 months, and the remaining 6 (0.4%) were given IFN- $\alpha$  combined with IFN- $\beta$  for 4 months. The median dose of 624 MU was administered during the median period of 24 weeks. IFN for 24 weeks or longer was given to 83.2% of the patients. IFN therapy was usually initiated within a few months after the diagnosis of chronic hepatitis, and all patients were started on it within 12 months. The median interval between liver biopsy and initiation of IFN was 9 days.


Almost all the patients given IFN showed varied degrees of fever, chills, myalgia, headache and general malaise after the first injection. Most patients developed leukocytopenia and thrombocytopenia in various degrees. A significant thrombocytopenia  $\leq$  40,000/mm<sup>3</sup> required a reduction of the IFN dose in 39 patients. IFN therapy was discontinued due to psychosis in 35 patients and ophthalmological symptoms in 12 patients. None of the patients developed decompensated liver disease with ascites, encephalopathy, jaundice or variceal bleeding. Although only 88 (5.3%) patients could not continue injection with IFN, studies for carcinogenesis were analyzed on the intention-to-treat basis.

The efficacy of IFN was judged by the clearance of HCV RNA from serum and ALT levels 12 months after the completion of treatment. Sustained virological response (SVR) was defined as persistent disappearance of HCV RNA after therapy, biochemical response (BR) as normal ALT levels without elimination of HCV RNA for at least 6 months after therapy, and no response (NR) as persistently elevated or transiently normalized ALT levels without loss of HCV RNA lasting for less than 6 months.

#### *Follow-Up of Patients and Diagnosis of HCC*

Patients were followed up monthly after diagnosis of chronic hepatitis in our outpatient clinic and monitored for hematological, biochemical and virological parameters. With their admission, during and after the treatment with IFN, weekly or biweekly follow-up was performed in almost all patients who received IFN. Imaging diagnosis was made once or twice per year in the majority of patients with ultrasonography or computed tomography. Angiography was performed only when HCC was highly suspected on imaging by ultrasonography or computed tomography.

When angiography pictured a characteristic hypervascular nodule specific for HCC in patients, histological confirmation was not required in the majority of them. Microscopic examinations of liv-



**Fig. 1.** Crude rates of hepatocarcinogenesis in patients treated with IFN and those untreated. The carcinogenesis rate was significantly lower in treated than in untreated patients (log-rank test,  $p < 0.0001$ ).

er tissues obtained by a fine-needle biopsy were performed in 14 patients whose angiogram could not portray a typical image of HCC. There were 89 patients in whom HCC was confirmed histologically on liver specimens obtained at surgery or autopsy. Detection of serological tumor markers and increase with time were also taken into account in the diagnosis of HCC.

There were 223 (10.3%) patients lost to follow-up, including 164 (9.9%) treated and 59 (11.5%) untreated. Rates of annual dropouts in treated and untreated patients were 0.95 and 0.93%, respectively. In 9 patients, the response to IFN was judged by information on aminotransferase levels determined in other clinics and by persistent HCV RNA, as well as aminotransferase levels at 6 months after the completion of therapy in an additional 3 patients. Therefore, the response to IFN could be judged in all patients including the 12 who were lost to our follow-up early. Since the eventual outcome with respect to the development of HCC was not confirmed in these patients, their data were censored in statistical analyses [21]. Deaths unrelated to liver disease were censored and withdrawn from the analysis. The date of the last follow-up in this study was May 1, 2004, and the median observation period of studied patients was 10.7 years, with a range of 0.1–33.6 years.

#### Statistical Analysis

Nonparametric Mann-Whitney U test and  $\chi^2$  test were used for analysis of background characteristics of patients. The rate of HCC development was calculated by the Kaplan-Meier method [22]; it was based on the duration between diagnosis of chronic hepatitis by liver biopsy and detection of HCC. Differences in slopes of carcinogenesis curves were evaluated by the log-rank test. To gain a robust statistical power for the anticarcinogenic activity of IFN, observation of treated patients was initiated at the commencement of IFN therapy, in lieu of the diagnosis of chronic hepatitis. Independent factors associated with the development of HCC were studied using the stepwise Cox regression analysis [23]. The follow-

ing 18 variables were analyzed for potential covariates in hepatocarcinogenesis at the time when hepatitis was diagnosed: age, sex, total alcohol intake, family history of liver disease, history of blood transfusion, stage of hepatic fibrosis, aspartic aminotransferase, ALT, albumin, bilirubin, globulin,  $\gamma$ -glutamyl transpeptidase ( $\gamma$ -GTP), platelet count, retention of indocyanine green at 15 min, serological grouping of HCV, HCV RNA level and IFN treatment.

Although continuous variables without conversion of data were evaluated in multivariate analyses, several variables were transformed into categorical data consisting of two or three ordinal numbers in calculating hazard ratios. All factors found to be marginally associated with hepatocarcinogenesis with  $p$  values  $< 0.15$  were tested by the multivariate Cox proportional hazard model. All analyses of data were performed with the computer program SPSS version 11 [24], and a  $p$  value  $< 0.05$  was considered significant.

## Results

### Response to IFN

Response to IFN was judged 12 months after the completion of therapy by both HCV RNA and serial ALT readings. Among the 1,654 patients with IFN treatment, SVR (elimination of HCV RNA) was achieved by 606 (36.6%), BR (ALT normalized for at least 6 months without clearance of HCV RNA from serum) in 266 (16.1%) and NR (elevated or transiently decreased ALT levels without loss of serum HCV RNA) in 782 (47.3%).

### Crude Rates of Hepatocarcinogenesis

During the median observation period of 10.7 years, HCC developed in 199 of the 2,166 (9.2%) patients, including 96 of the 1,654 (5.8%) patients treated with IFN and 103 of the 512 (20.1%) patients without IFN (fig. 1). Among the 199 patients with HCC, 140 (70.4%) imaged a typical hypervascular stain on angiography and dynamic computed tomography, while 59 failed to exhibit tumor stains on angiography. HCC in these 59 patients was confirmed histologically on liver specimens obtained at surgery or by fine-needle biopsy.

Crude rates of hepatocarcinogenesis in patients treated with IFN and those untreated were 1.3 and 1.8% at the end of the 3rd year (after the completion of therapy), 2.6 and 4.6% at the end of the 5th year, 5.8 and 12.7% at the 10th year and 13.9 and 23.9% at the 15th year, respectively (fig. 1). The carcinogenesis rate was significantly lower in patients treated with IFN than in untreated patients (log-rank test,  $p < 0.0001$ ).

### Impact of IFN on Hepatocarcinogenesis

During the observation period, HCC developed in 96 of the 1,654 (5.8%) patients treated with IFN, including



11 patients (1.8%) with SVR, 10 (3.8%) with BR and 75 (9.6%) with NR to IFN. Rates of hepatocarcinogenesis in patients with SVR, BR and NR were 0.7, 0.8 and 2.0% at the end of the 3rd year, 1.4, 2.0 and 3.8% at the 5th year, 1.6, 2.9 and 6.5% at the 7th year, 1.9, 3.6 and 9.6% at the 10th year and 1.9, 7.5 and 27.6% at the end of 15th year (fig. 2). Hepatocarcinogenesis was significantly less frequent in patients with SVR or BR than in patients with NR and those untreated (log-rank test,  $p < 0.0001$ ).

#### Factors Influencing Hepatocarcinogenesis


Univariate analysis identified 9 factors significantly associated with carcinogenesis. They were fibrotic stage ( $p < 0.001$ ), age ( $p < 0.001$ ),  $\alpha$ -fetoprotein ( $p < 0.001$ ), aspartic aminotransferase ( $p = 0.001$ ), retention of indocyanine green at 15 min ( $p = 0.002$ ), total alcohol intake ( $p = 0.002$ ),  $\gamma$ -GTP ( $p = 0.005$ ) and HCV serotype ( $p = 0.045$ ). IFN therapy ( $p = 0.064$ ), histological activity of hepatitis ( $p = 0.069$ ) and ALT ( $p = 0.70$ ) were marginally associated with carcinogenesis.

In order to prove the role of IFN on carcinogenesis in patients with chronic hepatitis type C en masse, multivariate analysis was performed by non-time-dependent proportional hazard analysis. Fibrotic stage,  $\gamma$ -GTP, gender, IFN therapy, platelet count and age independently influenced the development of HCC in the cohort (table 2). Advanced liver fibrosis in F2/F3 stages imposed a higher risk for carcinogenesis with a hazard ratio of 8.68, 95% confidence interval (CI) 5.08–14.81, compared with the F1 stage. Similarly, higher  $\gamma$ -GTP levels (hazard ratio 2.64), male sex (2.38), low platelet count (2.22) and older age (1.90) posed higher carcinogenesis risks. After adjusting background clinical biases between treated and untreated patients for the 5 significant covariates identified in the multivariate analysis, IFN therapy significantly decreased the hepatocarcinogenesis rate in the entire patients with chronic hepatitis C with a hazard ratio of 0.42 (95% CI 0.29–0.61) in comparison with untreated patients.

Based on the multivariate analysis, curves of carcinogenesis rates were theoretically illustrated in treated and untreated patients with the average histological stage, average  $\gamma$ -GTP value, average ratio of male to female, average platelet count and average age (fig. 3).

#### Hazard of Hepatocarcinogenesis Stratified by the Response to IFN

Since the carcinogenesis rate in patients with SVR or BR was significantly lower than that of patients with NR or untreated patients by the product limit method, a mul-




**Fig. 2.** Rates of hepatocarcinogenesis in patients with SVR, BR and NR to IFN. The rate in patients with NR (persistently elevated ALT or transiently normalized ALT for less than 6 months) was significantly higher than that in patients with SVR or BR.

**Table 2.** Factors associated with hepatocarcinogenesis in patients with chronic hepatitis C<sup>a</sup>

Factors	HR	95% CI	p value
Fibrosis stage			
F1	1		
F2–F3	8.68	(5.08–14.81)	<0.001
$\gamma$ -GTP, IU/ml			
<50	1		
$\geq 50$	2.64	(1.58–4.42)	<0.001
Gender			
Women	1		
Men	2.38	(1.56–3.70)	<0.001
IFN therapy			
No	1		
Yes	0.42	(0.29–0.61)	<0.001
Platelet count, $\times 10^3/\text{mm}^3$			
$\geq 100$	1		
<100	2.22	(1.47–3.44)	<0.001
Age, years			
<50	1		
$\geq 50$	1.90	(1.27–2.85)	0.002

HR = Hazard ratio.

<sup>a</sup> Evaluated by the Cox proportional hazard analysis.



**Fig. 3.** Theoretical curves of hepatocarcinogenesis in patients treated with IFN and those untreated who have the average histological stage, average  $\gamma$ -GTP value, average ratio of male to female, average platelet count and average age. They are based on the analysis of 1,654 patients treated with IFN and 512 untreated patients.

**Table 3.** Factors associated with hepatocarcinogenesis in patients with chronic hepatitis C who had distinct responses to IFN therapy<sup>a</sup>

Factors	HR	95% CI	p value
Fibrosis stage			
F1	1		
F2–F3	9.90	(4.19–23.40)	<0.001
Gender			
Women	1		
Men	3.44	(1.89–6.25)	<0.001
$\gamma$ -GTP, IU/ml			
<50	1		
$\geq 50$	2.68	(1.30–5.54)	0.008
Age, years			
<50	1		
$\geq 50$	2.56	(1.50–4.38)	0.001
AFP, ng/ml			
<20	1		
$\geq 20$	2.32	(1.34–4.02)	0.003
Platelet count, $\times 10^3/\text{mm}^3$			
$\geq 100$	1		
<100	2.09	(1.14–3.75)	0.013
Response to IFN			
Without IFN	1		
NR	0.57	(0.13–2.56)	0.46
BR	0.12	(0.04–0.35)	<0.001
SVR	0.10	(0.03–0.30)	<0.001

HR = Hazard ratio; AFP =  $\alpha$ -fetoprotein.

<sup>a</sup> Evaluated by the Cox proportional hazard analysis.

tivariate analysis was performed taking into account the response to IFN. Hazard ratios of patients with SVR and BR to IFN therapy were 0.10 (95% CI 0.03–0.30,  $p < 0.001$ ) and 0.12 (95% CI 0.04–0.35,  $p < 0.001$ ), respectively, in comparison with that of untreated patients, when the other 5 factors served as significant covariates (table 3). The hazard ratio of NR at 0.57 (95% CI 0.13–2.56) was less than 1, but fell short of making a significant difference against untreated patients.

#### Mortality and Causes of Death

During the observation period, 116 of the 2,166 (5.4%) patients died, including 52 of the 1,654 (3.1%) subjects treated with IFN and 64 of the 512 (12.5%) subjects without IFN. Estimated survival rates in the treated and untreated patients were 99.3 and 98.3% at 5 years, 97.8 and 96.0% at 10 years and 93.8 and 86.9% at 15 years, respectively. The survival rate of treated patients was significantly higher than that of untreated patients (log-rank test,  $p < 0.0001$ ).

#### Discussion

Based on our epidemiological data obtained by long-term observations of patients with chronic hepatitis [2] and patients with cirrhosis [1], the life expectancy of patients with HCV-related chronic liver disease heavily depends on the development of HCC. The possibility of eventually developing HCC in patients with HCV infection and cirrhosis is staggeringly high at 75% [1]. Theoretically, the treatment of chronic HCV infection with IFN can prevent the development of HCC. From the ethical point of view, a prospective randomized trial with control untreated patients is not to be allowed at present when IFN has become the standard radical therapy for chronic hepatitis C; everyone can receive IFN, as expenses are being covered for by the medical insurance in Japan. Another difficulty involves the informed consent in prospective randomized studies. It requires at least 5 years in order that IFN can decrease the incidence of carcinogenesis in chronic hepatitis C, with a statistical difference in the carcinogenesis rate between treated and 'untreated' patients. Since any randomized studies are considered extremely difficult in the future, we attempted to carry out this retrospective study by the multivariate analysis with statistical adjustments for possible covariates.

In the product limit analysis, IFN significantly decreased the crude rate of hepatocarcinogenesis in the

entire cohort of 2,166 patients with chronic hepatitis C. Since there were some background differences between treated and untreated patients, we tried to correct for biases including stage of fibrosis,  $\gamma$ -GTP value, sex, platelet count and age, which significantly affect the carcinogenesis rate. Demographic, histological and biochemical factors having been adjusted, IFN is proven to bring about a significant decrease in the hazard of carcinogenesis in patients with chronic hepatitis C en masse (hazard ratio 0.42,  $p < 0.001$  by the non-time-dependent model). Taking into consideration that a significant number of patients without IFN had received anti-inflammatory medicines, which might have contributed to suppression of hepatocarcinogenesis, the actual anticarcinogenic activity of IFN may be higher than the observed. Having published results of a similar study on a cohort of 1,643 patients with a median observation period of 5.4 years in 1999 [18], we could not establish the anticarcinogenic activity of IFN because of a low risk of carcinogenesis in untreated patients (1.2% per year). Nevertheless, we expected a significant statistical difference if we could extend the median observation period to longer than 7 or 10 years in our studied patients. This has been realized in the present study, in which 2,166 patients with and without IFN therapy were observed for a median of more than 10 years. As far as we are aware, it represents the first study that has demonstrated preventive effects of IFN on the carcinogenesis rate in a large cohort of patients in a single center, in correlation with distinct responses to it, such as SVR, BR and NR.

Treatment of patients with chronic HCV infection using IFN- $\alpha$  and ribavirin has led to sustained loss of serum HCV RNA in 40–50% of recipients with HCV genotype 1 and 75–80% with HCV genotype 2 or 3. However, to date, the combination therapy with IFN- $\alpha$  and ribavirin has not been evaluated for its impact on the risk of developing HCC. Monotherapy with IFN- $\alpha$  achieves sustained clearance of serum HCV RNA in only 20–30% of patients; the impact of IFN- $\alpha$  on the development of HCC has been evaluated only in patients who had received IFN- $\alpha$  without ribavirin [17–20, 25–27].

Multivariate analysis definitively demonstrated that IFN lessens the carcinogenesis risk in the patients whose ALT levels decreased after therapy. Furthermore, the anticarcinogenic capacity of IFN was demonstrated not only in the patients with persistent aminotransferase normalization, but also in those with transient normalization of ALT for at least 6 or 12 months. Many authors have already described that the activity of IFN to suppress the

development of HCC in patients with HCV RNA clearance (SVR) is similar to that in patients with ALT normalization in the absence of eliminating HCV RNA (BR) [18, 25–27]. Based on these compelling lines of evidence, the anticarcinogenic activity of IFN is ascribed to the suppression of inflammatory and regenerative processes in hepatocytes. Moreno and Muriel [28] reported that IFN reverts liver fibrosis, and therefore, control of the necro-inflammatory process can suppress the growth of HCC. Tarao et al. [29] reported that high aminotransferase levels increase the rate of HCC recurrence in patients with cirrhosis. Our results stand in favor of the view that the carcinogenic process in patients with chronic hepatitis C would be enhanced by fluctuating as well as persistently elevated levels of aminotransferases. It does seem that IFN exerts suppressive effects on HCC through reduction or complete remission of inflammatory activity. Recently, a few authors reported that even transient disappearance of HCV RNA during IFN therapy contributed to a low carcinogenesis rate in the clinical course of hepatitis [17, 27]. The significance of transient HCV in decreasing hepatocarcinogenesis should be further explored and confirmed by multicenter clinical studies with rigorous virological assessments.

HCC developed in a few patients with SVR 5 years after the HCV infection had been terminated by IFN, along with normalized ALT levels. These patients would have developed minute HCC in their livers already while receiving IFN which escaped the detection by imaging modalities or screening for serological tumor markers. This would indicate the limitation of IFN in preventing HCC. IFN will not be able to suppress HCC once it has developed, even when it succeeds in eliminating HCV and suppressing necroinflammatory processes in the liver.

With many difficulties in vaccine development, the recent progress in treatment of chronic HCV infection, from IFN monotherapy to combination therapy with ribavirin, is very auspicious. SVR and BR can be achieved in up to 56% of patients with combined IFN and ribavirin [30]. There is evidence that a sustained virological response can lead to decrease in fibrosis and even reversal of cirrhosis [31]. Because HCV-associated HCC occurs almost exclusively in patients with cirrhosis, successful treatment for SVR in patients without cirrhosis is likely to prevent future development of HCC [32]. However, once cirrhosis has been established, a preventive benefit of IFN monotherapy is restricted to the patients who can achieve SVR or BR. In their meta-analysis of 3 randomized and 11 nonrandomized controlled trials, Camma et

al. [33] have reported a low but statistically significant preventive effect.

In conclusion, IFN significantly decreases the rate of hepatocarcinogenesis in patients with chronic hepatitis C, irrespective of the response to it.

## Acknowledgement

This study was supported in part by a research grant from the Ministry of Health, Labour and Welfare in Japan.

## References

- Ikeda K, Saitoh S, Koida I, Arase Y, Tsubota A, Chayama K, Kumada H, Kawanishi M: A multivariate analysis of risk factors for hepatocellular carcinogenesis: A prospective observation of 795 patients with viral and alcoholic cirrhosis. *Hepatology* 1993;18:47-53.
- Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Koida I, Arase Y, Fukuda M, Chayama K, Murashima N, Kumada H: Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: A prospective observation of 2,215 patients. *J Hepatol* 1998;28:930-938.
- Davis GL, Balart LA, Schiff ER, Lindsay K, Bodenheimer HC Jr, Perrillo RP, Carey W, Jacobson IM, Payne J, Dienstag JL, van Thiel DH, Tamburro C, Lefkowitz J, Albrecht J, Meschivitz C, Ortego T: Treatment of chronic hepatitis C with recombinant interferon alfa. A multicenter randomized, controlled trial. *N Engl J Med* 1989;321:1501-1506.
- Di Bisceglie AM, Martin P, Kassianides C, Lisker-Melman M, Murray L, Waggoner J, Goodman Z, Banks SM, Hoofnagle JH: Recombinant interferon alfa therapy for chronic hepatitis C. A randomized, double-blind, placebo-controlled trial. *N Engl J Med* 1989;321:1506-1510.
- Causse X, Godinot H, Chevallier M, Chossegros P, Zoulim F, Ouzan D, Heyraud JP, Fontanges T, Albrecht J, Meschivitz C, Trepo C: Comparison of 1 or 3 MU of interferon alfa-2b and placebo in patients with chronic non-A, non-B hepatitis. *Gastroenterology* 1991;101:497-502.
- Chayama K, Saitoh S, Arase Y, Ikeda K, Matsumoto T, Sakai Y, Kobayashi M, Unakami M, Morinaga T, Kumada H: Effect of interferon administration on serum hepatitis C virus RNA in patients with chronic hepatitis C. *Hepatology* 1991;13:1040-1043.
- Nishiguchi S, Kuroki T, Nakatani S, Morimoto H, Takeda T, Nakajima S, Shiomi S, Seki S, Kobayashi K, Otani S: Randomised trial of effects of interferon-alpha on incidence of hepatocellular carcinoma in chronic active hepatitis C with cirrhosis. *Lancet* 1995;346:1051-1055.
- Mazzella G, Accogli E, Sottili S, Festi D, Orsini M, Salzetta A, Novelli V, Cipolla A, Fabbri C, Pezzoli A, Roda E: Alpha interferon treatment may prevent hepatocellular carcinoma in HCV-related liver cirrhosis. *J Hepatol* 1996;24:141-147.
- Schalm SW, Fattovich G, Brouwer JT: Therapy of hepatitis C: Patients with cirrhosis. *Hepatology* 1997;26:S128-S132.
- Benvegna L, Chemello L, Noventa F, Fattovich G, Pontisso P, Alberti A: Retrospective analysis of the effect of interferon therapy on the clinical outcome of patients with viral cirrhosis. *Cancer* 1998;83:901-909.
- Niederer C, Lange S, Heintges T, Erhardt A, Buschkamp M, Hurter D, Nawrocki M, Kruska L, Hensel F, Petry W, Haussinger D: Prognosis of chronic hepatitis C: Results of a large, prospective cohort study. *Hepatology* 1998;28:1687-1695.
- International Interferon-alpha Hepatocellular Carcinoma Study Group: Effect of interferon-alpha on progression of cirrhosis to hepatocellular carcinoma: A retrospective cohort study. *Lancet* 1998;351:1535-1539.
- Hu KQ, Tong MJ: The long-term outcomes of patients with compensated hepatitis C virus-related cirrhosis and history of parenteral exposure in the United States. *Hepatology* 1999;29:1311-1316.
- Ikeda K, Saitoh S, Kobayashi M, Suzuki Y, Suzuki F, Tsubota A, Arase Y, Murashima N, Chayama K, Kumada H: Long-term interferon therapy for 1 year or longer reduces the hepatocellular carcinogenesis rate in patients with liver cirrhosis caused by hepatitis C virus: A pilot study. *J Gastroenterol Hepatol* 2001;16:406-415.
- Fattovich G, Giustina G, Degos F, Tremolada F, Diodati G, Almasio P, Nevens F, Solinas A, Mura D, Brouwer JT, Thomas H, Njapoum C, Casarin C, Bonetti P, Fuschi P, Basho J, Tocco A, Bhalla A, Galassini R, Noventa F, Schalm SW, Realdi G: Morbidity and mortality in compensated cirrhosis type C: A retrospective follow-up study of 384 patients. *Gastroenterology* 1997;112:463-472.
- Valla DC, Chevallier M, Marcellin P, Payen JL, Trepo C, Fonck M, Bourliere M, Boucher E, Miguet JP, Parlier D, Lemonnier C, Opolon P: Treatment of hepatitis C virus-related cirrhosis: A randomized, controlled trial of interferon alfa-2b versus no treatment. *Hepatology* 1999;29:1870-1875.
- Kasahara A, Hayashi N, Mochizuki K, Takayanagi M, Yoshioka K, Kakumu S, Iijima A, Urushihara A, Kiyosawa K, Okuda M, Hino K, Okita K: Risk factors for hepatocellular carcinoma and its incidence after interferon treatment in patients with chronic hepatitis C. *Hepatology* 1998;27:1394-1402.
- Ikeda K, Saitoh S, Arase Y, Chayama K, Suzuki Y, Kobayashi M, Tsubota A, Nakamura I, Murashima N, Kumada H, Kawanishi M: Effect of interferon therapy on hepatocellular carcinogenesis in patients with chronic hepatitis type C: A long-term observation study of 1,643 patients using statistical bias correction with proportional hazard analysis. *Hepatology* 1999;29:1124-1130.
- Shindo M, Ken A, Okuno T: Varying incidence of cirrhosis and hepatocellular carcinoma in patients with chronic hepatitis C responding differently to interferon therapy. *Cancer* 1999;85:1943-1950.
- Yoshida H, Shiratori Y, Moriyama M, Arakawa Y, Ide T, Sata M, Inoue O, Yano M, Tanaka M, Fujiyama S, Nishiguchi S, Kuroki T, Imazeki F, Yokosuka O, Kinoyama S, Yamada G, Omata M: Interferon therapy reduces the risk for hepatocellular carcinoma: National surveillance program of cirrhotic and noncirrhotic patients with chronic hepatitis C in Japan. IHIT Study Group. Inhibition of Hepatocarcinogenesis by Interferon Therapy. *Ann Intern Med* 1999;131:174-181.
- Harrington DP, Fleming TR: A class of rank test procedures for censored survival data. *Biometrics* 1982;69:553-566.
- Kaplan EL, Meier P: Nonparametric estimation for incomplete observation. *J Am Stat Assoc* 1958;53:457-481.
- Cox DR: Regression models and life tables. *J R Stat Soc* 1972;34:248-275.
- SPSS Incorporation: SPSS for Windows Version 11.0 Manual. Chicago, SPSS Inc., 2001.
- Kasahara A, Hayashi N, Mochizuki K, Hiramatsu N, Sasaki Y, Kakumu S, Kiyosawa K, Okita K: Clinical characteristics of patients with chronic hepatitis C showing biochemical remission, without hepatitis C virus eradication, as a result of interferon therapy. The Osaka Liver Disease Study Group. *J Viral Hepat* 2000;7:343-351.
- Yabuuchi I, Imai Y, Kawata S, Tamura S, Noda S, Inada M, Maeda Y, Shirai Y, Fukuzaki T, Kaji I, Ishikawa H, Matsuda Y, Nishikawa M, Seki K, Matsuzawa Y: Long-term responders without eradication of hepatitis C virus after interferon therapy: Characterization of clinical profiles and incidence of hepatocellular carcinoma. *Liver* 2000;20:290-295.

- 27 Okanoue T, Itoh Y, Minami M, Sakamoto S, Yasui K, Sakamoto M, Nishioji K, Murakami Y, Kashima K: Interferon therapy lowers the rate of progression to hepatocellular carcinoma in chronic hepatitis C but not significantly in an advanced stage: A retrospective study in 1,148 patients. *J Hepatol* 1999;30:653-659.
- 28 Moreno MG, Muriel P: Remission of liver fibrosis by interferon-alpha 2b. *Biochem Pharmacol* 1995;50:515-520.
- 29 Tarao K, Takemiya S, Tamai S, Sugimasa Y, Ohkawa S, Akaike M, Tanabe H, Shimizu A, Yoshida M, Kakita A: Relationship between the recurrence of hepatocellular carcinoma (HCC) and serum alanine aminotransferase levels in hepatectomized patients with hepatitis C virus-associated cirrhosis and HCC. *Cancer* 1997;79:688-694.
- 30 Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves FL Jr, Haussinger D, Diago M, Carosi G, Dhumeaux D, Craxi A, Lin A, Hoffman J, Yu J: Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975-982.
- 31 Shiratori Y, Imazeki F, Moriyama M, Yano M, Arakawa Y, Yokosuka O, Kuroki T, Nishiguchi S, Sata M, Yamada G, Fujiyama S, Yoshida H, Omata M: Histologic improvement of fibrosis in patients with hepatitis C who have sustained response to interferon therapy. *Ann Intern Med* 2000;132:517-524.
- 32 Di Bisceglie AM, Carithers RL Jr, Gores GJ: Hepatocellular carcinoma. *Hepatology* 1998;28:1161-1165.
- 33 Camma C, Giunta M, Andreone P, Craxi A: Interferon and prevention of hepatocellular carcinoma in viral cirrhosis: An evidence-based approach. *J Hepatol* 2001;34:593-602.