

activity in the liver [1–5], along with marked biochemical and histological improvements. However, HBV mutants resistant to lamivudine have appeared in some patients during their treatment with this drug [6–10]. An increase in virus load during therapy is seen, reflecting a viral breakthrough that often brings about biochemical flare-ups of liver enzymes. Such lamivudine-resistant mutants have a characteristic mutation converting the 552nd amino acid for methionine, in the YMDD motif of reverse transcriptase, to isoleucine (YIDD mutant) [11–13] or valine (YVDD mutant) [11, 13]. An additional amino acid substitution (L528M) is often identified along with the YVDD mutant. A mutant with both these mutations has been reported to have the strongest drug resistance [14–16].

A method has been developed by us for detecting lamivudine-resistant HBV mutants [8]. It is based on restriction fragment length polymorphism (RFLP) of HBV DNA sequences amplified by polymerase chain reaction (PCR), and can detect the emergence of such mutants a few months before virological breakthroughs. By means of this RFLP method, the takeover of lamivudine-resistant mutants by the wild-type HBV was found to occur after the withdrawal of lamivudine [8]. No lamivudine-resistant mutants were detected, however, in any patients before they were placed on lamivudine, which might be due to the sensitivity of this method, which is not high enough for detecting mutants in low concentrations. To further explore the possibility of there being drug-resistant mutants even before the commencement of therapy, we have developed a more sensitive method for detecting rare mutants amongst a vast majority of the wild-type HBV [17].

The method involved PCR with peptide nucleic acid (PNA) clamping [18–20], and could detect 10² copies of YMDD mutants in the presence of 10⁶ copies of the wild-type HBV [17]. By means of this PCR-PNA method, YMDD mutants were detected in 1 of the 62 (1.6%) serum samples from patients with acute or chronic HBV infection positive for antibody to hepatitis B e antigen (anti-HBe) in serum who had not received lamivudine [17]. In addition, many YMDD mutants were found that had not been reported before [17]. In the present study, the analysis for YMDD mutants was conducted in serum samples from hepatitis B e antigen (HBeAg)-positive patients, and many rare mutants were identified, some of which have not been documented previously.

Materials and Methods

Serum Samples

A total of 85 serum samples from 72 adult Japanese patients with HBV infection were studied, of whom 10 (14%) were positive for HBeAg and 62 (86%) for the corresponding antibody (anti-HBe) in serum. Lamivudine was given to 3 of the 10 (30%) HBeAg-positive patients and 19 of the 62 (31%) anti-HBe-positive patients. All serum samples were stored frozen at –80° until molecular biological examinations. All patients were negative for serum markers of hepatitis C virus (HCV) or human immunodeficiency virus type 1 infection.

Serological Markers of HBV Infection

Hepatitis B surface antigen (HBsAg) was determined by enzyme immunoassay (Roche Diagnostics, Basel, Switzerland), and HBeAg as well as anti-HBe by radioimmunoassay (Abbott Diagnostics, Abbott Park, Ill., USA). HBV DNA was determined by transcription-mediated amplification and hybridization-protection assay (Chugai Diagnostics, Tokyo, Japan), and the results were expressed as log genome equivalents (LGE) per milliliter. The lower detection limit of this assay is 3.7 LGE/ml, which is equivalent to 5,000 copies/ml. The activity of HBV DNA polymerase was determined by a radioassay (Perkin Elmer, Boston, Mass., USA) that has a lower detection limit of 30 cpm. The antibody against HCV (anti-HCV) was tested for by the third-generation enzyme immunoassay (Roche Diagnostics).

Detection of YMDD Mutants by PCR with PNA Clamping

HBV DNA was extracted from 100 µl of serum by SMITEST (Genome Science Laboratories, Tokyo, Japan) and was dissolved in 20 µl of H₂O. The first-round of PCR was performed with an outer primer set (PLF1: 5'-GGT ATG TTG CCC GTT TGT CC-3' and BR123: 5'-TTC CAA TTA CAT ATC CCA T-3') and a second-round PCR with an inner primer set (PLF2: 5'-CCT ATG GGA GTG GGC CTC AG-3' and PLR2: 5'-CCA ATT ACA TAT CCC ATG AAG TTA AGG GA-3'). The PNA used was an 18-mer (PNA 552: H₂N-CAC ATC ATC CAT ATA ACT-CON₂H) that exactly matches the original YMDD sequence (fig. 1). The PNA-mediated PCR clamping method was optimized by changing the annealing temperature of PNA from 68 to 75°, and the PNA concentration from 1.25 to 10 µM. The optimized PCR with PNA clamping was performed in a total volume of 25 µl, consisting of a reaction buffer [100 mM Tris-HCl (pH 8.3), 50 mM KCl and 15 mM MgCl₂], 0.2 mM each of dNTPs, 1 µl of the DNA solution, 12.5 pmol each of primers, 150 pmol PNA 552 and 1 U Taq DNA polymerase (Gene Taq, Wako Pure Chemicals, Tokyo) along with 0.2 µg of anti-Taq high (Toyobo, Osaka, Japan). The optimized amplification conditions included initial denaturation at 95° for 4 min and 25 cycles of amplification (denaturation at 95° for 45 s, PNA annealing at 73° for 2 min, annealing and extension of primer at 63° for 50 s), followed by the final extension at 63° for 7 min. YVDD and YIDD mutants were detected by primer sets, PLF2 and TMApalI: 5'-CAG ACT TGG CCC CCA ATA CCA CAT CGT GCA-3' as well as YNSsp I: 5'-TTT CCC CCA CTG TTT GGC TTTC AGT AAT AT-3' and BR109: 5'-AAG GGA GTA GCC CCA ACG TT-3', as described previously [8]. YVDD and YIDD mutants can be detected after they had been digested by restriction enzyme *Alw44I* (Toyobo, Osaka, Japan) and *SspI* (Takara Shuzo, Otsu, Japan), respectively (fig. 1). The error rate of the Taq DNA polymerase was estimated to be 1.76 × 10⁻⁵ per site when amplifying about 10² copies of plasmid under the same conditions as above and cloning and sequencing [17].

Fig. 1. The experimental design. PNA is an 18-mer oligonucleotide of antiparallel polarity, and exactly matches the YMDD sequence. The amplified HBV DNA sequences from YVDD mutants only become sensitive to digestion with *Alw44I* by a change from GAT to CAC triplet. Consequently, the amplified HBV DNA sequences from YIDD mutants alone become sensitive to digestion with *SspI* by a change from TAT to AAT triplet. Sequence positions are in accord with those of Ono et al. [23].





Fig. 2. Clinical and virological course of an anti-HBe-positive patient with chronic hepatitis B who possessed the YVDD mutant before the start of lamivudine therapy. The YVDD mutant was detected by PCR-PNA [17] in 2 of 4 serum samples obtained before lamivudine therapy. The patient responded to lamivudine completely. IFN = Interferon; V = YVDD mutant; M = wild-type with YMDD motif.



Cloning and Sequencing Rare YMDD Mutants

Products (1 μ l) of the second-round of PNA-PCR were subjected to PCR with primers PLF2 and BR109 for 35 cycles (94°, 1 min; 58°, 1 min; 72°, 1.5 min) after initial denaturation at 94° for 4 min and followed by the final extension at 72° for 7 min. Amplicons were purified by electrophoresis on 2% (wt/vol) agarose gel and cloned into pGEM-T Easy Vector (Promega, Madison, Wisc., USA) with the standard method, and then transformed into *Escherichia coli* JM 109 (Takara Shuzo, Otsu, Japan). Sequencing was performed in the ABI PLISMTM 310NT Genetic analyzer (Applied Biosystems, Tokyo, Japan) with Big Dye terminator version 3.0 Cycle Sequencing Ready Reaction Kit (Applied Biosystems). Independent clones from 10 patients, including 5 with HBeAg positive and 5 with anti-HBe, were sequenced for analysis.

Results

Detection of YMDD Mutants in a Patient before the Start of Lamivudine Therapy

Only 1 of 72 patients (1.4%) tested positive for YVDD mutants before they received lamivudine therapy. The detection of YMDD mutants was comparatively infrequent as in our previous study in which they were found in only 1 of the 62 (1.6%) patients with chronic hepatitis B with anti-HBe in serum [17]. It was much less, however, than the detection by others [21, 22] of YMDD mutants in patients before the commencement of lamivudine. The single patient in the present study was positive for anti-HBe and responded completely to lamivudine without any breakthroughs as illustrated in figure 2. When HBV DNA was extracted from serial frozen serum samples and analyzed, YVDD mutants were detected only twice; lami-

Fig. 3. Nucleotide and amino acid sequences of clones from a patient including the YMDD motif, a clone having a YMDD sequence. The sequence of clone 1 with YVDD mutation was the same as that reported with lamivudine-resistant YVDD mutants. The YIDD sequence in clone 2, however, was different from the reported lamivudine-resistant YIDD mutant in that it possessed TAT ATA in place of TAT and ATT. These substitutions induce a TAG stop codon in the overlapping envelope gene (shown by an asterisk). Other variants, such as YRDD and YMDN, were also found in this patient (data not shown).

| | PNA | | | | | | | | | | | | | |
|----------|--|--|--|--|--|--|--|--|--|--|--|--|--|-----|
| (V00867) | TGTTTGGCTTTCAGTTATATGGATGATGTGGTATTGGGGGCCAAG | | | | | | | | | | | | | pol |
| | C L A F S Y M D D V V L G G K | | | | | | | | | | | | | HBs |
| | V W L S V I W M M W Y W G P S | | | | | | | | | | | | | |
| Clone 1 | TGTTTGGCTTTCAGTTATATGGATGATGTGGTATTGGGGGCCGAAA | | | | | | | | | | | | | pol |
| | C L A F S Y V D D V V L G A K | | | | | | | | | | | | | HBs |
| | V W L S V M W M M W Y W G R N | | | | | | | | | | | | | |
| Clone 2 | TGTTTGGCTTTCAGTTATATAGATGATGTGATATTGGGGGCCGAAA | | | | | | | | | | | | | pol |
| | C L A F S Y I D D V I L G A K | | | | | | | | | | | | | HBs |
| | V W L S V I * M M * Y W G R N | | | | | | | | | | | | | |
| Clone 3 | TGTTTGGCTTTCAGTTATAGGATGATGTGGTATTGGGGGCCAAG | | | | | | | | | | | | | pol |
| | C L A F S Y R D D V V L G A K | | | | | | | | | | | | | HBs |
| | V W L S V I G M M W Y W G R S | | | | | | | | | | | | | |
| Clone 4 | TGTTTGGCTTTCAGTTATATGGATAATGTGGTATTGGGGGCCAAG | | | | | | | | | | | | | pol |
| | C L A F S Y M D N V V L G A K | | | | | | | | | | | | | HBs |
| | V W L S V I W R M W Y W G R S | | | | | | | | | | | | | |

vudine suppressed HBV very well in this patient. The patient was negative for HBV DNA by PCR during therapy, and he experienced no breakthroughs (fig. 2).

To confirm the presence of YVDD mutants, HBV DNA clones were propagated from his pretreatment sera and sequenced. Many unusual mutants with mutations in the YMDD motif were recovered. Only 1 of the 43 (2%) clones had the YVDD sequence. The wild-type sequence (YMDD) was detected in 6 of the 43 clones (14%). Notably, 26 of these 43 clones (61%) possessed a stop codon in the overlapping envelope (S) gene. None of the 16 clones that had been obtained by PCR without PNA clumping disclosed any nucleotide mutations or amino acid substitutions in the YMDD motif (table 1, fig. 3). Only a few mutations were detected in the sequences where PNA does not anneal.

Detection of YMDD Mutants in a Patient with Chronic Hepatitis B Who Rebounded after the Withdrawal of Lamivudine

Serial serum samples from 1 patient who rebounded after the withdrawal of lamivudine therapy were examined. The activity of HBV-DNA polymerase started to increase soon after the patient was taken off lamivudine (fig. 4). Both YIDD and YVDD mutants were detected with the RFLP method [8] while he was receiving lamivudine. However, the YIDD mutant was detected about 2 months earlier with PCR with PNA clamping than with the RFLP method. After the withdrawal of lamivudine, both YIDD and YVDD mutants became undetectable

Table 1. Amino acid sequences of YMDD variants in HBV DNA clones from a patient before lamivudine therapy tested with or without PNA

| Mutation | With PNA (n = 43) | Without PNA (n = 16) |
|----------|----------------------|-------------------------|
| YMDD | 6 (14) | 16 (100) |
| YVDD | 1(2) | 0 |
| YIDD | 13 (30) ^a | 0 |
| YRDD | 3 (7) | 0 |
| YMDN | 10 (23) ^b | 0 |
| YIND | 3 (7) ^a | 0 |
| YINN | 7 (16) ^a | 0 |

Figures in parentheses represent percentage.

^a Created a stop codon in the overlapping envelope (S) gene in the YMDD motif.

^b Included the three clones with a stop codon in the overlapping envelope gene.

with the RFLP method, but they were still detectable with PCR with PNA clamping immediately after the rebound. Neither of these mutants, however, was detectable by either detection method 3 years thereafter. These results indicate that PCR with PNA clamping is useful for monitoring the appearance and disappearance of YMDD mutants during and after lamivudine therapy.

Fig. 4. Clinical and virological course of an HBeAg-positive patient with chronic hepatitis B who developed YMDD mutants during lamivudine therapy. The results of detecting YMDD mutants with the RFLP method of Chayama et al. [8] without PNA clamping and the PCR with PNA clamping [17] are compared. SNMC = Stronger Neo-Minophagen C (Minophagen Pharmaceutical Co., Tokyo, Japan); M = wild type with YMDD motif; I = YIDD mutant; V = YVDD mutant.




Table 2. Amino acid sequences of YMDD variants in HBV DNA clones from nine patients who tested negative for them by PCR with PNA clamping before or without lamivudine therapy

| | Clone | YMDD | YIDD | YRDD | YLDD | YMDN | YMND | YMNN | YMED | YMHD | YIND | YINN |
|------------------------|-----------|----------------|---------------------|--------------|--------------|--------------|---------------------|---------------------|--------------|--------------|---------------------|---------------------|
| Anti-HBe | | | | | | | | | | | | |
| Patient 1 | 10 | 4 (40) | 2 (20) ^a | 0 | 0 | 0 | 3 (30) ^a | 0 | 0 | 0 | 0 | 1 (10) ^a |
| Patient 2 | 10 | 0 | 5 (50) ^a | 0 | 0 | 0 | 3 (30) ^a | 1 (10) ^a | 0 | 0 | 0 | 1 (10) ^a |
| Patient 3 ^b | 11 | 10 (91) | 0 | 0 | 0 | 1 (9) | 0 | 0 | 0 | 0 | 0 | 0 |
| Patient 4 | 10 | 10 (100) | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| HBeAg | | | | | | | | | | | | |
| Patient 5 | 11 | 9 (82) | 0 | 0 | 1 (9) | 0 | 0 | 0 | 1 (9) | 0 | 0 | 0 |
| Patient 6 | 11 | 3 (27) | 8 (72) ^a | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Patient 7 | 11 | 5 (45) | 0 | 1 (9) | 0 | 2 (18) | 2 (18) ^a | 0 | 0 | 1 (9) | 0 | 0 |
| Patient 8 | 10 | 2 (20) | 3 (30) ^a | 0 | 0 | 0 | 3 (30) ^a | 0 | 0 | 0 | 2 (20) ^a | 0 |
| Patient 9 ^c | 11 | 1 (9) | 3 (27) ^a | 0 | 3 (27) | 1 (9) | 0 | 1 (9) ^a | 0 | 0 | 1 (9) ^a | 1 (9) ^a |
| Total | 95 | 44 (47) | 21 (22) | 1 (1) | 4 (4) | 4 (4) | 11 (12) | 2 (2) | 1 (1) | 1 (1) | 3 (3) | 3 (3) |

Figures in parentheses represent percentage.

- ^a Created a stop codon in the overlapping envelope (S) gene.
^b Developed viral breakthrough after 21 months on lamivudine.
^c Developed viral breakthrough after 7 months on lamivudine.

Detection of YMDD Mutants in Patients with Chronic Hepatitis B

Although the YVDD mutant was detected in only 1 of 72 patients with PCR-PNA and restriction enzyme digestion analyses, we went on to analyze quasispecies of HBV in this region in 9 additional patients by cloning and sequencing. Multiple clones from 4 patients with anti-

HBe and 5 with HBeAg were examined for YMDD mutants. Only 1 patient, who tested positive for anti-HBe in serum, did not exhibit YMDD mutants (patient 4 in table 2). However, their YMDD clones had a unique nucleotide sequence (TAC ATG GAT GAT); the biological relevance of these substitutions is unknown.

YIDD sequences were detected in 5 patients, with the incidence of YIDD mutants in 60% of the 10 patients examined (including the patient positive for YVDD mutant). However, all these YIDD sequences were different from the typical lamivudine-resistant strain (TAT ATT GAT GAT) and they all possessed a stop codon in the overlapping envelope (S) gene (clone 2 in fig. 3). This suggests that these clones with YIDD mutation would represent defective that were unrelated to typical lamivudine resistance.

Discussion

Although lamivudine is effective in suppressing viral replication and lowering alanine aminotransferase levels in serum [1–5], a frequent emergence of YMDD mutants detracts enormously from its effects [6–10]. It is not known, however, if such mutants exist before the therapy. In an attempt to detect YMDD mutants, we previously developed a PCR-based RFLP method that can sensitively identify YVDD and YIDD mutants [8]. The RFLP method is not sensitive, however, in detecting a minor population of YMDD mutants that coexist with the vast majority of the wild-type HBV without mutations in the YMDD motif. Only 10^2 copies of YMDD mutants were detected when mixed with 10^4 copies of the wild-type HBV, even when wild-type sequences were digested by restriction enzyme in the products of the first-round PCR. Hence the sensitivity was not sufficient for detecting very small amounts of mutants that exist among overwhelming numbers of the wild-type virus. We developed, therefore, a more sensitive detection method of mutants by using PCR with PNA clamping [17].

Although the PCR-PNA method could detect YMDD mutants 10- to 100-fold more sensitively than the previous PCR-RFLP method [17], such mutants were not detected in sera from patients before the start of lamivudine, even though they later developed a viral breakthrough within 1 year on lamivudine (patient 9 in table 2). This suggests that it would be very difficult to detect YMDD mutants before the commencement of therapy by increasing the sensitivity of the methods. It does need to be pointed out that the patients who tested positive for YVDD mutant did not develop a breakthrough during at least 23 months and showed a good response to the therapy. This suggests that the YMDD mutant detected in our patient might have been an incompetent virus that would have a mutation in the other part of the genome not analyzed so far. Combined, these observations suggest that

increasing the sensitivity of the assay per se would be inappropriate for detecting mutants before therapy. Different approaches should be considered in selecting patients suitable for antiviral treatments.

Previous studies have found YMDD mutants in pre-treatment sera from the patients who tested positive for anti-HBe [17, 21, 22]. The detection goes along with the fact that HBV sequences with multiple substitutions exist in the patients who are mounting active immune responses against HBV with anti-HBe immune responses. In fact, 5 patients with HBeAg were found to harbor many YMDD mutants in the present study. This suggests that mutants would not be exclusive for the patients who are under immune pressure against HBV with anti-HBe in serum. Many mutants identified in this study and a previous study [17] underscore the production of a number of mutants during viral genome synthesis. Only a part of these mutants might be able to survive and produce their progeny and evolve under the immunological pressure. Further study should be directed to gain a full profile of rare quasispecies of HBV mutants, which would help understand the mechanism for the emergence of drug-resistant HBV mutants. Detection of rare quasispecies by PCR with PNA clamping will be a useful tool in monitoring for YMDD mutants arising before and after lamivudine therapy.

Acknowledgments

This work was carried out at the Research Center for Molecular Medicine, Faculty of Medicine, Hiroshima University. We thank Eiko Okutani, Chiemi Noguchi, Kiyomi Toyota for their excellent technical assistance, and Yutaka Fujii for helpful comments. This study was supported in part by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Health, Labor and Welfare.

References

- 1 Lai CL, Ching CK, Tung AK, Li E, Young J, Hill A, Wong BC, Dent J, Wu PC: Lamivudine is effective in suppressing hepatitis B virus DNA in Chinese hepatitis B surface antigen carriers: A placebo-controlled trial. *Hepatology* 1997;25:241-244.
- 2 Nevens F, Main J, Honkoop P, Tyrrell DL, Barber J, Sullivan MT, Fevery J, De Man RA, Thomas HC: Lamivudine therapy for chronic hepatitis B: A six-month randomized dose-ranging study. *Gastroenterology* 1997;113:1258-1263.
- 3 Dienstag JL, Schiff ER, Wright TL, Perrillo RP, Hann HW, Goodman Z, Crowther L, Condreay LD, Woessner M, Rubin M, Brown NA: Lamivudine as initial treatment for chronic hepatitis B in the United States. *N Engl J Med* 1999;341:1256-1263.
- 4 Suzuki Y, Kumada H, Ikeda K, Chayama K, Arase Y, Saitoh S, Tsubota A, Kobayashi M, Koike M, Ogawa N, Tanikawa K: Histological changes in liver biopsies after one year of lamivudine treatment in patients with chronic hepatitis B infection. *J Hepatol* 1999;30:743-748.
- 5 Liaw YF, Leung NW, Chang TT, Guan R, Tai DI, Ng KY, Chien RN, Dent J, Roman L, Edmundson S, Lai CL: Effects of extended lamivudine therapy in Asian patients with chronic hepatitis B. *Gastroenterology* 2000;119:172-180.
- 6 Honkoop P, Niesters HG, de Man RA, Osterhaus AD, Schalm SW: Lamivudine resistance in immunocompetent chronic hepatitis B. Incidence and patterns. *J Hepatol* 1997;26:1393-1395.
- 7 Allen MI, Deslauriers M, Andrews CW, Tipples GA, Walters KA, Tyrrell DL, Brown N, Condreay LD: Identification and characterization of mutations in hepatitis B virus resistant to lamivudine. *Hepatology* 1998;27:1670-1677.
- 8 Chayama K, Suzuki Y, Kobayashi M, Kobayashi M, Tsubota A, Hashimoto M, Miyano Y, Koike H, Kobayashi M, Koida I, Arase Y, Saitoh S, Murashima N, Ikeda K, Kumada H: Emergence and takeover of YMDD motif mutant hepatitis B virus during long-term lamivudine therapy and re-takeover by wild type after cessation of therapy. *Hepatology* 1998;27:1711-1716.
- 9 Liaw YF, Chien RN, Yeh CT, Tsai SL, Chu CM: Acute exacerbation and hepatitis B virus clearance after emergence of YMDD motif mutation during lamivudine therapy. *Hepatology* 1999;30:567-572.
- 10 Ono-Nita SK, Kato N, Shiratori Y, Lan KH, Yoshida H, Carrilho FJ, Omata M: Susceptibility of lamivudine-resistant hepatitis B virus to other reverse transcriptase inhibitors. *J Clin Invest* 1999;103:1635-1640.
- 11 Ling R, Mutimer D, Ahmed M, Boxall EH, Elias E, Dusheiko GM, Harrison TJ: Selection of mutations in the hepatitis B virus polymerase during therapy of transplant recipients with lamivudine. *Hepatology* 1996;24:711-713.
- 12 Tipples GA, Ma MM, Fischer KP, Bain VG, Kneteman NM, Tyrrell DL: Mutation in HBV RNA-dependent DNA polymerase confers resistance to lamivudine in vivo. *Hepatology* 1996;24:714-717.
- 13 Bartholomew MM, Jansen RW, Jeffers LJ, Reddy KR, Johnson LC, Bunzendahl H, Condreay LD, Tzakis AG, Schiff ER, Brown NA: Hepatitis-B-virus resistance to lamivudine given for recurrent infection after orthotopic liver transplantation. *Lancet* 1997;349:20-22.
- 14 Xiong X, Flores C, Yang H, Toole JJ, Gibbs CS: Mutations in hepatitis B DNA polymerase associated with resistance to lamivudine do not confer resistance to adefovir in vitro. *Hepatology* 1998;28:1669-1673.
- 15 Gutfreund KS, Williams M, George R, Bain VG, Ma MM, Yoshida EM, Villeneuve JP, Fischer KP, Tyrrel DL: Genotypic succession of mutations of the hepatitis B virus polymerase associated with lamivudine resistance. *J Hepatol* 2000;33:469-475.
- 16 Ono SK, Kato N, Shiratori Y, Kato J, Goto T, Schinazi RF, Carrilho FJ, Omata M: The polymerase L528M mutation cooperates with nucleotide binding-site mutations, increasing hepatitis B virus replication and drug resistance. *J Clin Invest* 2001;107:449-455.
- 17 Ohishi W, Shirakawa H, Kawakami Y, Kimura S, Kamiyasu M, Tazuma S, Nakanishi T, Chayama K: Identification of rare polymerase variants of hepatitis B virus using two-stage PCR with peptide nucleic acid clamping. *J Med Virol*, in press.
- 18 Nielsen PE, Egholm M, Berg RH, Buchardt O: Sequence-selective recognition of DNA by strand displacement with a thymine-substituted polyamide. *Science* 1991;254:1497-1500.
- 19 Egholm M, Buchardt O, Christensen L, Behrens C, Freier SM, Driver DA, Berg RH, Kim SK, Norden B, Nielsen PE: PNA hybridizes to complementary oligonucleotides obeying the Watson-Crick hydrogen-bonding rules. *Nature* 1993;365:566-568.
- 20 Zhong S, Nguyen NY, Eggerman TL: Detection of apolipoprotein B mRNA editing by peptide nucleic acid mediated PCR clamping. *Biochem Biophys Res Commun* 1999;259:311-313.
- 21 Kobayashi S, Ide T, Sata M: Detection of YMDD motif mutations in some lamivudine-untreated asymptomatic hepatitis B virus carriers. *J Hepatol* 2001;34:584-586.
- 22 Kirishima T, Okanoue T, Daimon Y, Itoh Y, Nakamura H, Morita A, Toyama T, Minami M: Detection of YMDD mutant using a novel sensitive method in chronic liver disease type B patients before and during lamivudine treatment. *J Hepatol* 2002;37:259-265.
- 23 Ono Y, Onda H, Sasada R, Igarashi K, Sugino Y, Nishioka K: The complete nucleotide sequences of the cloned hepatitis B virus DNA; subtype adr and adw. *Nucleic Acids Res* 1983;11:1747-1757.

Molecular evolutionary analyses implicate injection treatment for schistosomiasis in the initial hepatitis C epidemics in Japan

Yasuhito Tanaka¹, Kousuke Hanada², Etsuro Orito³, Yoshihiro Akahane⁴, Kazuaki Chayama⁵, Hiroshi Yoshizawa⁶, Michio Sata⁷, Nobuo Ohta⁸, Yuzo Miyakawa⁹, Takashi Gojobori², Masashi Mizokami^{1,*}

¹Department of Clinical Molecular Informative Medicine, Nagoya City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan

²Center for Information Biology, National Institutes of Genetics, Mishima, Japan

³Department of Internal Medicine and Molecular Science, Nagoya City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan

⁴First Department of Internal Medicine, Faculty of Medicine, University of Yamanashi, Yamanashi, Japan

⁵Department of Medicine and Molecular Science, Hiroshima University Graduate School of Biomedical Sciences, School of Medicine, Hiroshima, Japan

⁶Department of Infectious Disease and Control, Hiroshima University Graduate School of Biomedical Sciences, School of Medicine, Hiroshima, Japan

⁷Second Department of Internal Medicine, Kurume University School of Medicine, Fukuoka, Japan

⁸Department of Molecular Parasitology, Nagoya City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan

⁹Miyakawa Memorial Research Foundation, Tokyo, Japan

Background/Aims: The mortality due to hepatocellular carcinoma (HCC) has ranged widely in various areas of Japan since 30 years ago and the incidence was particularly high in once *Schistosoma japonicum* (*Sj*)-endemic areas. Our aim was to estimate the spread time of hepatitis C virus (HCV) infection in the past with possible relevance to a higher incidence of HCC in once *Sj*-endemic than *Sj*-nonendemic areas.

Methods: During 2001, 131 strains of HCV-1b were obtained from patients in three previously *Sj*-endemic areas, as well as *Sj*-nonendemic areas in Japan and a cross-sectional study was conducted on them with molecular evolutionary analyses.

Results: A phylogenetic tree reconstructed on HCV-1b sequences in the NS5B region disclosed 2 independent clusters for *Sj*-positive and -negative groups with a high bootstrap value. The estimated effective number of HCV-infections indicated a transition from quiescence to rapid exponential growth in the 1920s among patients with schistosomiasis, which is 20 years earlier than that among patients without schistosomiasis.

Conclusions: The estimated spread time in previously *Sj*-endemic areas in Japan coincides with injection treatment for *Sj* since 1921. A high incidence of HCC there would be attributed to a long duration of HCV infection since 1920s.

© 2004 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

Keywords: Hepatitis C virus; *Schistosoma japonicum*; Molecular evolutionary analysis; Hepatocellular carcinoma

1. Introduction

Recently, the molecular clock has been successfully applied to long-term serial serum samples containing hepatitis C virus (HCV) from the US and Japan and estimated the spread time of HCV in the 1930s in Japan, which is 30 years earlier than that in the US in the 1960s [1]. Insofar as a long duration of HCV infection is the most important factor for the development of hepatocellular

Received 17 June 2004; received in revised form 1 September 2004; accepted 6 September 2004; available online 22 October 2004

* Corresponding author. Tel.: +81 52 853 8292; fax: +81 52 842 0021.

E-mail address: mizokami@med.nagoya-cu.ac.jp (M. Mizokami).

Abbreviations HCV, hepatitis C virus; Anti-HCV, antibody to HCV; HCC, hepatocellular carcinoma; *Sj*, *Schistosoma japonicum*.

0168-8278/\$30.00 © 2004 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.
doi:10.1016/j.jhep.2004.09.023

carcinoma (HCC), it can be predicted that the incidence of HCC will increase in the US over the next 2–3 decades. Thus, a combination of classical epidemiological approaches and molecular evolutionary analyses would be particularly useful in the study of contagious diseases, typified by HCV infection.

The way how individuals contracted HCV infection has remained unclear in Japan. Recently, a Japanese report (Ministry of Health, Labour and Welfare: Distribution of age-adjusted mortality rate from liver cancer by prefecture between 1971 and 1975, Tokyo, 2001) indicated that the mortality due to HCC has already varied widely in various areas of Japan since 30 years ago; the incidence of HCC was much higher in Saga/Fukuoka, Hiroshima and Yamanashi Prefectures, which were once endemic for schistosomiasis japonica in the long past. Hence, a high incidence of HCC in the 1970s would be related to HCV transmitted by injection treatment for *Schistosoma japonicum* (*Sj*) conducted since 1921 in these areas. In fact, shared needles and syringes for intravenous injection treatment with antimony potassium tartrate or sodium antimony tartrate posed a significant risk for HCV transmission in endemic areas [2]. Indeed, the prevalence of antibody to HCV (anti-HCV) is high (36.5; 95% CI=28.1–44.9%) in patients with chronic schistosomiasis [2] and therefore, HCV infection is considered responsible for the development of HCC in patients with chronic schistosomiasis.

Since, once popular intravenous injection for schistosomiasis was a risk factor for HCV transmission, the spread time of HCV in the areas once endemic for *Sj* in Japan would deserve determination. In this study, molecular evolutionary analyses using principles of both population genetics and mathematical epidemiology [3] were applied to HCV-infected patients with and without a past history of chronic schistosomiasis in once *Sj*-endemic areas.

2. Materials and methods

2.1. Sample collection

In Japan during 2001, 181 random serum samples positive for anti-HCV were obtained from patients with chronic liver disease in widely separated areas previously endemic for *Sj*, including Kofu in Yamanashi ($n=75$), Katayama in Hiroshima ($n=50$) and Chikugo in Saga/Fukuoka Prefectures ($n=56$). Schistosomiasis was diagnosed by ultrasonographic (US) and/or computer tomographic (CT) modalities or serological examinations [4]. Two kinds of serological tests, which can detect past history of schistosomiasis, were available in this study. In brief, IgG antibodies binding to two different *schistosoma* antigens, *Sj* adult worm antigen and *Sj* egg antigen, were detected using an enzyme-linked immunosorbent assay (ELISA). As it is now accepted that ELISA titer of egg antigen-specific IgG is reliable for case-detection rather than IgG for adult worm antigen [4–6], the results based on the egg antigen-specific IgG were accepted in this study. Samples of more than 0.25 of optical density at 415 nm were determined to be positive, as previously confirmed [4–6]. The serum samples were tested for anti-HCV by Lumipulse II Ortho HCV (Ortho-Clinical Diagnostics K.K., Tokyo, Japan). As patients with *Sj* treatments were estimated to be old,

relatively older patients were selected in the *Sj*-endemic areas to match age factor that might influence duration of HCV infection or HCC incidence. For a cross-sectional study, 30 serum samples were obtained from patients infected with HCV in Aichi Prefecture where *Sj* has not been endemic. The age- and sex-matched patients were also selected from the *Sj*-nonendemic areas excluding influence of these factors on HCC incidence. The study protocol conformed to the 1975 Declaration of Helsinki and was approved by Ethic Committees of institutions. Every patient gave a written informed consent to participate in the virological research of HCV. Information of injection treatment for *Sj* was obtained by means of self-administrated questionnaires or structured interviews. None had been treated with interferon therapy for HCV infection. HCC incidence was estimated by historical information from patients themselves and/or medical records during 2001. HCC was diagnosed by liver biopsy or combination of imaging modalities such as US, enhanced CT and angiography.

2.2. Genotyping and sequencing

Nucleic acids were extracted using a SepaGean RV-R Nucleic acid extracting kit (Sanko Junyaku Co., Ltd., Tokyo, Japan) in accordance with the manufacturer's protocol. They were reverse-transcribed to cDNA using SuperScript II RNase H⁻ Reverse Transcriptase (Invitrogen Corp., Carlsbad, California, USA) and random hexamer primer (Takara Shuzo Co. Ltd, Tokyo, Japan) by the method described previously [7].

A sequence spanning 339 nucleotides (nt) in the NS5B region was amplified by polymerase chain reaction (PCR) with primers described previously [1]. PCR products were directly sequenced with Prism Big Dye (Applied Biosystems, Foster City, California, USA) in an ABI 3100 DNA automated sequencer. To reduce the number of artificial substitutions arising in PCR, PLATINUM Pfx DNA Polymerase (Invitrogen Corp.) with a very high fidelity was used. The sequences determined were utilized to confirm HCV genotypes and construct phylogenetic trees.

2.3. Test for clustering between *Sj*-positive and -negative groups

The phylogenetic tree was first constructed to examine the evolutionary history for *Sj*-positive and *Sj*-negative groups by the neighbor joining method [8]. Furthermore, to test whether either *Sj*-positive or *Sj*-negative group have evolved independently or not, we conducted an interior branch test for the neighbor-joining tree [9]. Thereafter, a *t*-test was conducted for the interior branch length and its standard error, which is computed using the bootstrap procedure.

2.4. Demographic model

A reconstructed tree was built on the NS5B sequence of 339 nt by a heuristic maximum-likelihood topology search with stepwise-addition and the nearest neighbor-interchange algorithms. Tree likelihood scores were calculated using HKY85 with the molecular clock enforced by PAUP version 4.0b8.

As estimates of the demographic history, a nonparametric function $N(t)$, known also as the skyline plot, was obtained by transforming coalescent intervals of an observed genealogy into a piecewise plot that represents an effective number of infections through time [3,10]. A parametric maximum-likelihood was estimated by several models with the computer software Genie v3.5 to build a statistical framework for inferring the demographic history of a population on phylogenies reconstructed on sampled DNA sequences [10]. This model assumes a continuous epidemic process in which the viral transmission parameters remain constant through time. Model fitting was evaluated by likelihood ratio tests of the parametric maximum-likelihood estimates [11,12].

2.5. Statistical method

Data for continuous variables were demonstrated as the mean \pm standard deviation. The Fishers' exact test, Chi square test with Yates' correction and one-way ANOVA followed by the Scheffe's multiple comparison test were used to evaluate differences in the mean age, sex ratio

and incidence of HCC between groups, respectively. Differences with *P* values less than 0.05 were considered significant.

3. Results

Of 181 anti-HCV positive samples, 113 were classified into HCV genotype 1b (HCV-1b), which is predominant in Japan. Fifty-two of 181 samples (29%) were negative for HCV RNA or incomplete for sequencing and the remaining 16 samples (9%) of genotype 2a were excluded in this study due to a minor population. Of the HCV-1b strains, 47 were recovered from patients in Yamanashi, 31 in Hiroshima and 35 in Saga/Fukuoka Prefectures. Along with 18 HCV-1b strains in Aichi Prefecture serving as controls, a cross-sectional study was conducted on them with molecular evolutionary analyses. The patients in areas previously endemic for *Sj* revealed a significantly higher prevalence of chronic schistosomiasis [24/47 (51%) in Yamanashi (Kofu area), 21/31 (68%) in Hiroshima (Katayama area) and 19/35 (54%) in Saga/Fukuoka (Chikugo area)] than that in Aichi Prefecture (0/18 [0%], $P < 0.0001$). There were no significant differences in the mean age or sex ratio among patients from these four areas (Fig. 1). Although the mean age of *Sj*-positive patients was just higher than that of *Sj*-negative patients in once *Sj*-endemic areas or matched-control patients in Aichi Prefecture, there were also no significant differences between these groups (Table 1).




Fig. 1. Geographic distribution of *Schistosoma japonicum* (*Sj*) and characteristics of patients infected with HCV. *Sj* (+) and *Sj* (-) denote, respectively, presence and absence of infection with *Sj* diagnosed by ultrasonographic and/or computer tomographic methods or serological examinations. Pie graphs include the age (mean \pm standard deviation) and sex ratio (male/female).

Table 1
Characteristics of patients with and without schistosomiasis

| | Schistosoma japonicum | | Controls (Aichi) (n = 18) |
|-------------------|-----------------------|----------------------|---------------------------------|
| | Positive (n = 64) | Negative (n = 49) | |
| Mean age | | | |
| Total | 69.9 \pm 7.7 | 67.4 \pm 8.7 | 66.5 \pm 9.2 |
| Yamanashi | 69.9 \pm 7.2 | 67.3 \pm 11.2 | |
| Hiroshima | 71.2 \pm 8.7 | 67.6 \pm 6.5 | |
| Saga/Fukuoka | 69.0 \pm 7.7 | 67.5 \pm 7.1 | |
| Sex (male/female) | | | |
| Total | 34/30 | 24/25 | 9/9 |
| Yamanashi | 13/11 | 11/12 | |
| Hiroshima | 10/11 | 5/5 | |
| Saga/Fukuoka | 11/8 | 8/8 | |
| Incidence of HCC | 25/55 (45%) | 11/48 (23%) | 3/18 (17%) |

The incidence of HCC in *Sj*-positive patients was significantly higher than that in *Sj*-negative patients ($P = 0.0226$) or controls ($P = 0.0488$).

Abbreviations: HCC, hepatocellular carcinoma.

For cross-sectional study on the viral population size between HCV-infected patients with and without a past history of schistosomiasis, a phylogenetic tree for HCV-1b strains in the *Sj*-positive and -negative patients was constructed with use of the maximum-likelihood method enforced by the molecular clock as introduced in our previous report [1] and an independent study by Pybus et al. [3]; a substitution rate of 5.3×10^{-4} per site per year [1,3] was assumed for HCV. The phylogenetic tree disclosed 2 independent clusters for *Sj*-positive and -negative groups, with a high bootstrap value (81%) by the interior branch testing (Fig. 2), which is comparative with past epidemiological backgrounds in Japan. From distinct evolutionary histories in the two populations, the effective number of HCV-1b infections through time, $N(t)$, were assessed by the skyline plot. The parameters for several models in Genie v3.5 [3,10] were also examined. Time t was then transformed to year using the same rate, assuming the collecting time (year 2001) as the present. Fig. 3 shows the skyline plots and population growth for *Sj*-positive and -negative patients, according to a specific demographic model in Genie v3.5 with three parameters, piecewise expansion growth model, that was evaluated by the likelihood ratio testing [11,12]. Molecular evolutionary results thus obtained supported our previous study in which the divergence time of the most recent common ancestor of HCV-1b in each area in Japan was estimated before 1850 [1]. Our estimates of the effective number of HCV-infections showed a transition from constant size to rapid exponential growth in the 1920s among patients with chronic schistosomiasis in endemic areas, which is 20 years earlier than that among patients without schistosomiasis in the 1940s. Information on HCC was available for 121 of the 131 patients with HCV-1b. Although they were relatively small in number, the incidence of HCC was significantly higher in *Sj*-positive than -negative patients ($P = 0.0226$) or controls ($P = 0.0488$) (Table 1).




Fig. 2. A phylogenetic tree constructed on NS5B sequences of HCV-1b strains in *Schistosoma japonicum* (*Sj*)-positive ($n=64$) and -negative ($n=67$) groups. The numbers in the tree indicate bootstrap reliability by the interior branch test. *Sj*+ indicates *Sj*-positive strains. YAM; Yamanashi, HI/HIS; Hiroshima, KU/KUS; Saga/Fukuoka, Aichi; control strains.

4. Discussion

The specific demographic model based on the neutral theory [3,11,12], which has a constant size in the past and changes to exponential growth until the present, is applied to investigate the Japanese endemic of HCV. By means of the molecular evolutionary analyses, the spread time of HCV in *Sj*-positive patients was estimated 20 years earlier than that in *Sj*-negative patients from three areas in Japan where *Sj* was previously endemic (Yamanashi, Hiroshima, Saga/Fukuoka Prefectures). The spread time of HCV much earlier in *Sj*-positive than -negative patients indicates that the previous intravenous injection treatment with antimony compounds (antimonyl potassium tartarate or antimony sodium tartarate) on patients with schistosomiasis since 1921 [2] would have been a significant risk factor for HCV transmission in endemic areas through re-used needles and syringes. Indeed, it might be possible that HCV transmission from *Sj*-positive patients to *Sj*-negative patients occurs in the once *Sj*-endemic areas, but we could not find such strains in this study. One of the reasons is that residents in the village around the river, where schistosomiasis had been the most prevalent, might have been isolated from those in the other areas of the same Prefecture in the past due to the endemic disease 'schistosomiasis'. Interestingly, most Japanese strains from *Sj*-nonendemic areas in the database clustered with the *Sj*-negative group of the present study. Hence, factors other than the injection treatment for *Sj*, such as intravenous stimulants popular during and after World War II [13] and medical treatments including transfusion with blood units from paid donors in the past, would have imposed the risk for HCV transmission in most areas in Japan [14]. In addition, there would have been opportunities for HCV transmission through inadequately sterilized needles and syringes in general practices, which have contributed to a large reservoir of chronic HCV infection in Japan during the 1950s [13]. Such inadequately sterilized medical injections were still common in the less-developed world in the 20th century. WHO estimates that unsafe injections result in 2.3–4.7 million new HCV infections worldwide every year [15].

Although the spread time of HCV in *Sj*-positive group was earlier than that in *Sj*-negative group, there was no significant difference of mean age between the 2 groups. Two possibilities are considered. One is a sampling bias; as patients with *Sj* treatments were estimated to be old, relatively older patients were selected in the *Sj*-endemic areas to match age factor that might influence duration of HCV infection or HCC incidence. Second, the ages that patients had been infected with HCV were different between the 2 groups; the treatments for *Sj* in Japan were mainly conducted among relatively younger people including school children after screening of *Sj* [4,16,17], while the




Fig. 3. The maximum-likelihood estimates of $N(t)$ on the effective number of infections with HCV genotype 1b in Japan for *Schistosoma japonicum* (*Sj*)-positive group (a) and *Sj*-negative group (b) separated in the phylogenetic tree (Fig. 2). The parametric model is indicated by the grey line and stepwise plots by the black line that represents corresponding nonparametric estimates of $N(t)$ (number as a function of time). Genetic distances are transformed into a time scale of year using estimates of the molecular clock in the NSSB region.

other risk factors such as blood transfusion were found in older people excluding at least children.

A disease possibly caused by schistosomal infection in Japan is documented in a book written some 300 years ago. In 1847, the clinical picture of this disease was precisely described by Yoshinao Fujii in the book 'Katayama-ki' that documented an endemic disease in Katayama area as Katayama's disease (equivalent to schistosomiasis). Water-borne epidemics of schistosomiasis prevailed in inhabitants around rivers (the tributaries of the Fuji river in Yamanashi, the Takaya river in Hiroshima and the Chikugo river in Saga/Fukuoka) in Japan, mediated by

small shellfish (Miyairi-kai) serving as the natural host. More than 200,000 individuals were estimated to have been infected with *Sj* in Yamanashi Prefecture alone during 1965 through 1990 [16] and approximately 1,000,000 patients in the entire Japan since 1920s [17]. To cope with these endemics, more than 10 million intravenous injections with antimony compounds had been given in Japan since 1921 [17]. Thus, Japan would have started ahead of any other countries, in terms of HCV spread in association with schistosomiasis, wherein intravenous drugs were invented. Although acute schistosomal infection has disappeared in Japan since long ago, there are still elderly people with

chronic schistosomiasis in previously endemic areas, some of whom are developing HCC [2,14]. Substantial transmission among regions is supported by the lack of regional clustering of HCV sequences in this study.

A similar situation is reported in the Nile delta in Egypt where schistosomiasis once prevailed mediated by small shellfish [18] and the national campaigns for injection treatment with antimony potassium tartarate (tartar emetic) from the 1961 until 1986 are suspected to have given rise to the highest endemicity of HCV in the world ever, involving >20% of the national population there [19]. The prevalence of anti-HCV is extremely high (>70%) in patients with schistosomiasis there [18,20,21]. Highly prevalent HCV infection in the general Egyptian population accounts for most HCC cases in Egypt [22]. A question may arise whether schistosomiasis alone is responsible for the development of HCC. Patients co-infected with HCV and *Schistosoma mansoni* (*Sm*) may have a high incidence of viral persistence, accelerated fibrosis and development of HCC [23,24]. A recent population-based study between two large populations with district histories of *Sm* and hepatitis C infections, however, failed to indicate any interaction between *Sm* infection and the prevalence or severity of hepatitis C [25]. Moreover, no significant histological differences were found between anti-HCV-positive Egyptian patients with and without schistosoma [26]. Hence, the long duration of persistent HCV infection would be a more important factor for the development of HCC than the pathogenicity of *Sm* itself.

Estimating the effective number of HCV infections has been very informative in looking back epidemic spreads of HCV infection in the United States [1] and Egypt [12,27]. In addition, it would also be useful in predicting the population size and extent of HCV infection. Studies to foresee future spreads of HCV would be required to cope with and prevent healthcare problems where de novo infections are increasing. The advantage of molecular evolutionary analyses, its ability to accurately estimate the dynamics of HCV based on a limited number of isolates in particular [3], will extend its application anywhere in the world where clinical sequelae of persistent HCV infection pose increasing burdens on the public health of nations.

In conclusion, the evolutionary analyses indicated that the estimated spread time in previously *Sj*-endemic areas in Japan coincides with injection treatment for *Sj* conducted since 1921. The high incidence of HCC in *Sj*-endemic areas is most likely attributed to long duration of HCV infection there transmitted through injection treatments.

Acknowledgements

We greatly appreciate Dr Oliver G. Pybus (Department of Zoology, University of Oxford, Oxford, UK) for his enlightening advice on molecular evolutionary analyses

using Genie v3.5. We thank Ms Yoshiko Kobayashi (Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan) for technical assistance and Dr Tatsuya Ide (Kurume University School of Medicine, Fukuoka, Japan) and Dr Hirotaka Kono (Hiroshima University Graduate School of Biomedical Sciences, School of Medicine, Hiroshima, Japan) for providing us with valuable materials. The work was supported by a grant-in-aid from the Ministry of Health, Labour and Welfare of Japan (H13-kanen-2), grants-in-aid for Young Scientists (A) from the Ministry of Education, Culture, Science and Sports of Japan (16689016), The Hori Information Science Promotion Foundation and Miyakawa Memorial Research Foundation.

References

- [1] Tanaka Y, Hanada K, Mizokami M, Yeo AE, Shih JW, Gojobori T, et al. Inaugural article: a comparison of the molecular clock of hepatitis C virus in the United States and Japan predicts that hepatocellular carcinoma incidence in the United States will increase over the next two decades. *Proc Natl Acad Sci USA* 2002;99:15584–15589.
- [2] Iida F, Iida R, Kamijo H, Takaso K, Miyazaki Y, Funabashi W, et al. Chronic Japanese schistosomiasis and hepatocellular carcinoma: ten years of follow-up in yamanashi prefecture, Japan. *Bull World Health Organ* 1999;77:573–581.
- [3] Pybus OG, Charleston MA, Gupta S, Rambaut A, Holmes EC, Harvey PH. The epidemic behavior of the hepatitis C virus. *Science* 2001;292:2323–2325.
- [4] Minai M, Hosaka Y, Ohta N. Historical view of schistosomiasis japonica in Japan: implementation and evaluation of disease-control strategies in yamanashi prefecture. *Parasitol Int* 2003;52:321–326.
- [5] Matsuda H, Tanaka H, Blas BL, Nosenas JS, Tokawa T, Ohsawa S. Evaluation of ELISA with ABTS, 2-2'-azino-di-(3-ethylbenzthiazoline sulfonic acid), as the substrate of peroxidase and its application to the diagnosis of schistosomiasis. *Jpn J Exp Med* 1984;54:131–138.
- [6] Matsumoto J, Kirinoki M, Kawai S, Chigusa Y, Ilagan EJ, Ducusin BE, et al. Prevalence of schistosomiasis japonica among schoolchildren and animal reservoirs in oriental mindoro, Philippines. *Jpn J Trop Med Hyg* 1999;27:175–180.
- [7] Ohno T, Mizokami M, Wu RR, Saleh MG, Ohba K, Orito E, et al. New hepatitis C virus (HCV) genotyping system that allows for identification of HCV genotypes 1a, 1b, 2a, 2b, 3a, 3b, 4, 5a and 6a. *J Clin Microbiol* 1997;35:201–207.
- [8] Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987;4:406–425.
- [9] Dopazo J. Estimating errors and confidence intervals for branch lengths in phylogenetic trees by a bootstrap approach. *J Mol Evol* 1994;38:300–304.
- [10] Pybus OG, Rambaut A. GENIE: estimating demographic history from molecular phylogenies. *Bioinformatics* 2002;18:1404–1405.
- [11] Lemey P, Pybus OG, Wang B, Saksena NK, Salemi M, Vandamme AM. Tracing the origin and history of the HIV-2 epidemic. *Proc Natl Acad Sci USA* 2003;100:6588–6592.
- [12] Pybus OG, Drummond AJ, Nakano T, Robertson BH, Rambaut A. The epidemiology and iatrogenic transmission of hepatitis C virus in Egypt: a Bayesian coalescent approach. *Mol Biol Evol* 2003;20:381–387.
- [13] Fukui S, Wada K, Iyo M. History and current use of methamphetamine in Japan. *Proceedings of Japan-US Scientific*

- Symposium '90 on Drug Dependence and Abuse; 7–8 September 1990. Tokyo, Japan: Cocaine and methamphetamine behavioral toxicology, clinical psychiatry and epidemiology; 1991 [p. 219–37].
- [14] Yoshizawa H. Hepatocellular carcinoma associated with hepatitis C virus infection in Japan: projection to other countries in the foreseeable future. *Oncology* 2002;62:8–17.
- [15] Drucker E, Alcabes PG, Marx PA. The injection century: massive unsterile injections and the emergence of human pathogens. *Lancet* 2001;358:1989–1992.
- [16] Kitani K, Iuchi M. Schistosomiasis japonica: a vanishing endemic in Japan. *J Gastroenterol Hepatol* 1990;5:160–172.
- [17] Okabe K. Epidemiology of schistosomiasis japonica. *Kurume Med J* 1975;38:11–55.
- [18] Kamel MA, Ghaffar YA, Wasef MA, Wright M, Clark LC, Miller FD. High HCV prevalence in Egyptian blood donors. *Lancet* 1992;340:427.
- [19] Cohen J. The scientific challenge of hepatitis C. *Science* 1999;285:26–30.
- [20] Hibbs RG, Corwin AL, Hassan NF, Kamel M, Darwish M, Edelman R, et al. The epidemiology of antibody to hepatitis C in Egypt. *J Infect Dis* 1993;168:789–790.
- [21] Frank C, Mohamed MK, Strickland GT, Lavanchy D, Arthur RR, Magder LS, et al. The role of parenteral antischistosomal therapy in the spread of hepatitis C virus in Egypt. *Lancet* 2000;355:887–891.
- [22] Hassan MM, Zaghoul AS, El-Serag HB, Soliman O, Patt YZ, Chappell CL, et al. The role of hepatitis C in hepatocellular carcinoma: a case control study among Egyptian patients. *J Clin Gastroenterol* 2001;33:123–126.
- [23] Angelico M, Renganathan E, Gandin C, Fathy M, Profili MC, Refai W, et al. Chronic liver disease in the Alexandria governorate, Egypt: contribution of schistosomiasis and hepatitis virus infections. *J Hepatol* 1997;26:236–243.
- [24] Kamal S, Madwar M, Bianchi L, Tawil AE, Fawzy R, Peters T, et al. Clinical, virological and histopathological features: long-term follow-up in patients with chronic hepatitis C co-infected with *S. mansoni*. *Liver* 2000;20:281–289.
- [25] Blanton RE, Salam EA, Kariuki HC, Magak P, Silva LK, Muchiri EM, et al. Population-based differences in *Schistosoma mansoni*- and hepatitis C-induced disease. *J Infect Dis* 2002;185:1644–1649.
- [26] Helal T, Danial M, Ahmed H. The relationship between hepatitis C virus and schistosomiasis: histopathologic evaluation of liver biopsy specimens. *Hum Pathol* 1998;29:743–749.
- [27] Tanaka Y, Agha S, Saady N, Kurbanov F, Orito E, Kato T, et al. Exponential spread of hepatitis C virus genotype 4a in Egypt. *J Mol Evol* 2004;58:191–195.

Influence of Age, Sex, and Degree of Liver Fibrosis on the Association Between Serum Alanine Aminotransferase Levels and Liver Inflammation in Patients with Chronic Hepatitis C

HIDENORI TOYODA, MD, TAKASHI KUMADA, MD, SEIKI KIRIYAMA, MD, YASUHIRO SONE, MD, MAKOTO TANIKAWA, MD, YASUHIRO HISANAGA, MD, KAZUHIKO HAYASHI, MD, TAKASHI HONDA, MD, and TELJI KUZUYA, MD

In 504 patients with chronic hepatitis C who underwent liver biopsy, the correlation between serum ALT levels and histologic liver inflammation was investigated, and the effects of age, sex, and degree of histologic liver fibrosis on this correlation were analyzed. Serum ALT levels were significantly higher in male than in female patients only when activity of hepatitis was mild. In contrast, ALT levels were significantly higher in younger (≤ 50 years old) than in older patients only when activity of liver inflammation was severe. Fibrosis was closely associated with activity of hepatitis and, also, serum ALT level. More importantly, a significant number of older patients and patients with severe hepatic fibrosis had severe hepatic inflammation even when their serum ALT level was not markedly elevated (≤ 70 IU/L). Age, sex, and degree of liver fibrosis independently influenced serum ALT levels in patients with chronic hepatitis C.

KEY WORDS: chronic hepatitis C; serum alanine aminotransferase; age; sex; fibrosis.

Aminotransferase levels are sensitive indicators of liver-cell injury (1). The serum alanine aminotransferase (ALT) level, in particular, is the most commonly used variable in assessments of liver disease; it is used as an indicator of the presence of chronic liver disease and the severity of liver damage. In patients with chronic viral hepatitis, the serum ALT level is sometimes considered in the decision to initiate antiviral therapy (2), although this is now controversial (3). Furthermore, ALT levels may be associated

with the risk of hepatocellular carcinoma in patients with cirrhosis (4). However, precise correlation between the serum ALT level and activity of hepatitis (liver inflammation) has not been sufficiently established, especially as it relates to the degree of liver fibrosis. In addition, the influences of patient age and sex on the correlation between serum ALT level and activity of hepatitis have not been well analyzed. Better interpretation of serum ALT levels in routine follow-up for patients with chronic hepatitis C would be important for physicians to make more appropriate clinical decisions. In the present study, therefore, we investigated the influence of patient age, sex, and degree of liver fibrosis on the correlation between serum ALT level and histologically evaluated activity of hepatitis in patients with chronic hepatitis C.

Manuscript received June 25, 2003; accepted October 28, 2003.

From the Department of Gastroenterology, Ogaki Municipal Hospital, Ogaki, Japan.

Address for reprint requests: Hidenori Toyoda, MD, Department of Gastroenterology, Ogaki Municipal Hospital, 4-86 Minaminokawa, Ogaki, Gifu, 503-8502, Japan; tkumada@he.mirai.ne.jp.

METHODS

Between 1990 and 1998, a total of 504 patients with chronic hepatitis C underwent ultrasonography-guided needle liver biopsy. These patients comprised 332 men and 172 women (mean age, 49.0 ± 10.4 years). None of them had received interferon therapy before liver biopsy. In all patients, infection with hepatitis C virus (HCV) had been confirmed by detection of anti-HCV antibody by second-generation assay (Dinabot; Tokyo) and of serum HCV RNA by reverse-transcription polymerase chain reaction (PCR) assay (5). PCR with genotype-specific primers on a core region (6) showed HCV genotype 1b in 304 patients, 2a in 152 patients, and 2b in 25 patients; the test showed mixed genotypes in 8 patients, and genotype was not determined in 15 patients.

Activity of hepatitis and liver fibrosis were evaluated by two independent liver pathology specialists who were blinded to the clinical data. Evaluation was according to the classification of Desmet *et al.* (7). Activity of hepatitis was graded as minimal (A0), mild (A1), moderate (A2), or severe (A3). The degree of fibrosis was scored as none (F0), mild (F1), moderate (F2), or severe (F3). Serum ALT level was measured by the routine method on the day of liver biopsy.

Data are shown as mean \pm SD. Differences in proportions were analyzed by chi-square test. Differences in mean quantitative values were analyzed by Student's *t*-test. Multivariate analysis was done using logistic regression (8); the variables analyzed were age at biopsy (≤ 50 vs > 50 years), sex, and degree of liver fibrosis (F0/F1 vs. F2/F3). Statistical analyses were performed with the SAS statistical package (SAS Institute, Cary, NC) (9). All *P* values were two-tailed, and values less than 0.05 were accepted as statistically significant.

RESULTS

Association Among Serum ALT Level, Activity of Hepatitis, and Degree of Liver Fibrosis. The mean serum ALT level for all patients was 91.7 ± 67.5 IU/L. Among the 504 patients, activity scores were A0 ($n = 42$), A1 ($n = 286$), A2 ($n = 165$), and A3 ($n = 11$). Fibrosis scores were F0 ($n = 92$), F1 ($n = 225$), F2 ($n = 138$), and F3 ($n = 49$). Serum ALT levels increased according to the degree of activity (Figure 1). Also, ALT levels increased in association with the progression of liver fibrosis (Figure 2).

Distribution of the activity of hepatitis is shown according to the degree of liver fibrosis in Table 1. The proportion of patients with high activity (A2 or A3) markedly increased in patients with a high degree of fibrosis (F2 or F3). Thus, activity of hepatitis and degree of liver fibrosis together were associated with an elevation in serum ALT.

Association Between Serum ALT Level and Patient Age and Sex. Serum ALT levels were analyzed according to age and sex of patients. Serum ALT levels were significantly higher in male patients than in female patients (males, 98.1 ± 69.9 IU/L, vs. females, 80.1 ± 63.1 IU/L; $P = 0.0048$). However, serum ALT levels were similar




Fig 1. Serum alanine aminotransferase (ALT) levels per activity of hepatitis. ALT levels increase in association with the grade of the activity. ALT levels are significantly higher in patients with A1 activity than in those with A0 activity ($P = 0.0475$) and in patients with A2 activity than in those with A1 activity ($P = 0.0003$).

between younger patients (≤ 50 years old; $n = 246$) and older patients (> 50 years old; $n = 258$) (younger, 95.8 ± 69.5 IU/L, vs. older, 88.2 ± 66.8 IU/L; $P = 0.2148$).

When patients were stratified by low activity of hepatitis (A0/A1) and high activity of hepatitis (A2/A3), we found that serum ALT levels did not differ by sex among patients with A2/A3 (males, 111.9 ± 70.4 IU/L, vs. females, 104.9 ± 72.2 IU/L; $P = 0.5362$); however, we found a marked difference between the sexes in patients with A0/A1 (males, 90.9 ± 68.8 IU/L, vs. females, 66.0 ± 52.8 IU/L; $P = 0.0010$) (Figure 3). In contrast, we found a significant difference in serum ALT levels between younger and older patients with A2/A3 (younger, 128.6 ± 75.7 IU/L, vs. older, 98.2 ± 65.8 IU/L; $P =$




Fig 2. Serum alanine aminotransferase (ALT) levels per degree of liver fibrosis. ALT levels increase in association with the degree of fibrosis. ALT levels are significantly higher in patients with F2 fibrosis than in those with F1 fibrosis ($P = 0.0097$).

SERUM ALT IN PATIENTS WITH CHRONIC HEPATITIS C

TABLE 1. CORRELATION BETWEEN THE ACTIVITY OF HEPATITIS AND THE DEGREE OF FIBROSIS IN PATIENTS WITH CHRONIC HEPATITIS C

| Activity of hepatitis | Degree of fibrosis | | | |
|-----------------------|--------------------|-----|----|----|
| | F0 | F1 | F2 | F3 |
| A0 | 30 | 12 | 0 | 0 |
| A1 | 60 | 171 | 52 | 3 |
| A2 | 2 | 42 | 80 | 41 |
| A3 | 0 | 0 | 6 | 5 |

Note. $P < 0.0001$ for A0 or A1/A2 or A3 vs. F0 or F1/F2 or F3.

0.0058) and no difference between younger and older patients with A0/A1 (younger, 84.0 ± 63.4 IU/L, vs. older, 80.7 ± 66.8 IU/L; $P = 0.6492$) (Figure 4). Thus, the difference in serum ALT by sex was observed only when the activity of hepatitis was low, whereas the difference by age was observed only when the activity of hepatitis was high.

Multivariate Analysis for Influence of Age, Sex, and Degree of Liver Fibrosis on Serum ALT Levels. Multivariate analysis showed that all three factors (age, sex, and degree of liver fibrosis) independently affected serum ALT levels (>70 IU/L) (degree of liver fibrosis, $P = 0.0043$; age, $P = 0.0038$; sex, $P = 0.0184$), in addition to activity of hepatitis ($P = 0.0366$). When patients were stratified by activity of hepatitis, a high degree of liver fibrosis (F2 or F3) ($P = 0.0229$) and male sex ($P = 0.0153$) independently increased serum ALT levels in patients with A0 or A1. In patients with A2 or A3, age greater than 50 years independently increased serum ALT levels ($P = 0.0002$).

Influence of Age, Sex, and Degree of Liver Fibrosis in Patients with Persistently Normal ALT Levels or with Persistently Mild ALT Elevation. In considering patients in whom serum ALT levels were persistently less than 70 IU/L for more than 6 months before liver biopsy,

which is less than twice the normal limit of ALT levels in our hospital, only 10.8% of patients with F0/F1 fibrosis had A2/A3 activity of hepatitis. In contrast, 67.1% of patients with F2/F3 fibrosis had A2/A3 activity of hepatitis ($P < 0.0001$; Table 2). There was no difference in the proportion of patients with moderate or severe activity of hepatitis (A2/A3) between men and women ($P = 0.9943$). However, 39.0% of patients >50 years old had A2/A3 activity of hepatitis, whereas only 11.8% of patients ≤ 50 years old had A2/A3 activity of hepatitis ($P < 0.0001$; Table 3). In addition, even in patients with persistently normal ALT levels (<35 IU/L), 6 of 13 patients (46.2%) with F2/F3 fibrosis had A2/A3 activity of hepatitis and 10 of 42 patients (23.8%) >50 years old had A2/A3 activity of hepatitis.

DISCUSSION

Initial evaluation of a patient with chronic HCV infection focuses on assessment of the activity and stage of liver disease, based on the histology of a liver biopsy specimen. Liver histology is frequently considered the gold standard for assessing the progression of chronic hepatitis in terms of the activity of hepatitis and fibrosis. However, it is usually difficult to monitor patients by serial histological evaluation because liver biopsy has potential complications.

Liver fibrosis is usually irreversible, except in patients in whom HCV has been eradicated by antiviral treatment, and it progresses slowly, over years. In contrast, the activity of hepatitis (i.e., liver inflammation) is reversible and is likely to fluctuate. Therefore, routine follow-up of patients with chronic hepatitis C focuses on the fluctuation in the activity of hepatitis, and patients are usually evaluated by laboratory tests.




Fig 3. Serum alanine aminotransferase (ALT) levels per sex. Left: Comparison among patients with A0/A1 activity. Right: Comparison among patients with A2/A3 activity. Male patients had higher serum ALT levels than female patients among patients with A0/A1 activity (males, 90.9 ± 68.8 IU/L, vs. females, 66.0 ± 52.8 IU/L; $P = 0.0010$).




Fig 4. Serum alanine aminotransferase (ALT) levels per age. Left: Comparison among patients with A0/A1 activity. Right: Comparison among patients with A2/A3 activity. Younger patients (≤ 50 years old) had higher serum ALT levels than older patients (> 50 year old) among patients with A2/A3 activity (younger, 128.6 ± 75.7 IU/L, vs. older, 98.2 ± 65.8 IU/L; $P = 0.0058$).

In the clinical setting, ALT is generally considered the most reliable indicator among laboratory measures of active liver injury. The highest concentration of ALT in the body is in the liver, and levels of this enzyme are therefore the most specific indicators of liver injury. ALT is reportedly released into the blood in increasing amounts when the liver cell membrane is damaged (1). An old study showed that elevations in serum aminotransferase levels are markers of liver cell injury (10), and another study showed a close correlation between elevated transaminases and histologic necroinflammation in biopsied specimens from patients with chronic liver disease (11, 12).

Most patients with low serum ALT levels show histologically low activity of hepatitis and low fibrosis (3, 13–16). Some previous studies, however, yielded a poor correlation between serum ALT levels and histological findings of liver-cell damage (13, 14, 17, 18). The volume of residual hepatocytes is reduced in patients with severe fibrosis because of the accumulation of fibrotic tissue caused by chronic liver disease; this reduction in the volume of intact hepatocytes may modify the correlation between liver inflammation and serum ALT levels. As a result, the elevation in serum ALT levels in association with the increase

in the activity of hepatitis can be suppressed in patients with severe fibrosis. In addition, serum ALT levels reportedly vary according to sex and body mass index (BMI) in healthy individuals (19, 20).

In the present study, we evaluated the influence of age, sex, and degree of liver fibrosis on the elevation in serum ALT in patients with chronic hepatitis C. We found that patients with a higher degree of liver fibrosis had higher serum ALT levels, mainly because patients with severe fibrosis had a higher activity of hepatitis than that of patients with mild fibrosis. More importantly, however, we found that a large proportion of patients with severe fibrosis also had high activity of hepatitis (A2 or A3), even those with persistently normal ALT levels or those with persistently mild elevation in serum ALT (less than twice the normal value for more than 6 months). This should be taken into consideration when physicians evaluate the activity of hepatitis on the basis of serum ALT levels.

The increase in serum ALT levels in association with the activity of hepatitis seems to differ between the sexes. Two previous studies reported lower ALT levels in healthy females than in healthy males (19, 20). Thus, there is a trend toward lower ALT levels in females without liver

TABLE 2. DISTRIBUTION OF THE ACTIVITY OF HEPATITIS ACCORDING TO THE DEGREE OF FIBROSIS IN PATIENTS WITH SERUM ALT LEVELS ≤ 70 IU/L

| Activity of hepatitis | Degree of fibrosis | |
|-----------------------|--------------------|-------|
| | F0/F1 | F2/F3 |
| A0/A1 | 157 | 23 |
| A2/A3 | 19 | 47 |

Note. $P < 0.0001$.

TABLE 3. DISTRIBUTION OF THE ACTIVITY OF HEPATITIS ACCORDING TO AGE IN PATIENTS WITH SERUM ALT LEVELS ≤ 70 IU/L

| Activity of hepatitis | Age of patients | |
|-----------------------|-----------------|-----------|
| | ≤ 50 yr | > 50 yr |
| A0/A1 | 97 | 83 |
| A2/A3 | 13 | 53 |

Note. $P < 0.0001$.

disease. This difference in serum ALT levels in healthy individuals seems to be maintained when the activity of hepatitis and fibrosis of the liver is mild. However, this difference disappears when the activity is severe.

Interestingly, in contrast, the age of patients influenced the increase in serum ALT levels when the activity of hepatitis became severe. This suggests a reduced elevation in ALT in response to liver injury in older patients. Liver morphology reportedly undergoes changes such as hepatocyte ploidy as a person ages (21), which may account for the difference in ALT elevation between younger and older patients observed in our study. In addition, 39.0% of older patients with a minor elevation in serum ALT (less than twice the normal value) had higher activity of hepatitis (A2 or A3). Thus, physicians should also consider patient age in the interpretation of serum ALT levels.

In the present study, we did not evaluate the influence of BMI because of a lack of data. BMI reportedly influences serum ALT levels in healthy individuals (19, 20). Therefore, the influence of BMI on serum ALT levels in patients with chronic hepatitis C will require further study.

In summary, age, sex, and degree of liver fibrosis independently influence serum ALT levels in patients with chronic hepatitis C. Physicians should consider these factors during routine follow-up of this population. Especially in older patients or patients with severe fibrosis, the activity of hepatitis can be severe even with mild ALT elevation. Serum ALT is still the most commonly used indicator of liver cell injury, and there is no current alternative method of monitoring fluctuation in liver inflammation that is more reliable than serum ALT. Therefore, further studies are needed to evaluate factors influencing the association between liver inflammation and serum ALT for a better understanding of the degree of liver injury.

REFERENCES

- Pratt DS, Kaplan MM: Evaluation of abnormal liver-enzyme results in asymptomatic patients. *N Engl J Med* 342:1266-1271, 2000
- Shakil AO, Conry-Cantilena C, Alter HJ, Hayashi P, Kleiner DE, Tedeschi V, Krawczynski K, Conjeevaram HS, Sallie R, Di Bisceglie AM: Volunteer blood donors with antibody to hepatitis C virus: clinical, biochemical, virologic, and histologic features. *Ann Intern Med* 123:330-337, 1995
- Bacon BR: Treatment of patients with hepatitis C and normal serum aminotransferase levels. *Hepatology* 36:S179-S184, 2002
- Tarao K, Rino Y, Ohkawa S, Shimizu A, Tamai S, Miyakawa K, Aoki H, Imada T, Shindo K, Okamoto N, Totsuka S: Association between high serum alanine aminotransferase levels and more rapid development and higher rate of incidence of hepatocellular carcinoma in patients with hepatitis C virus-associated cirrhosis. *Cancer* 86:589-595, 1999
- Okamoto H, Okada S, Sugiyama Y, Tanaka T, Sugai Y, Akahane Y, Machida A, Mishiro S, Yoshizawa H, Miyakawa Y, Mayumi M: Detection of hepatitis C virus RNA by a two-stage polymerase chain reaction with two pairs of primers deduced from the 5'-noncoding region. *Jpn J Exp Med* 60:215-222, 1990
- Okamoto H, Sugiyama Y, Okada S, Kurai K, Akahane Y, Sugai Y, Tanaka T, Sato K, Tsuda F, Miyakawa Y, Mayumi M: Typing hepatitis C virus by polymerase chain reaction with type-specific primers. An application to clinical surveys and tracing infection sources. *J Gen Virol* 73:673-679, 1992
- Desmet VJ, Gerber M, Hoofnagle JH, Manns M, Scheuer PJ: Classification of chronic hepatitis: diagnosis, grading and staging. *Hepatology* 19:1513-1520, 1994
- Cox DR: Regression models and life tables. *J Royal Stat Soc B* 34:187-220, 1972
- SAS Institute: SAS/STAT User's Guide: Basics, Version 6 (1st ed.). Cary, NC, SAS Institute, 1990
- Schmidt E, Schmidt FW, Wildhirt E: Aktivitäts-Bestimmungen von Enzymen des Energieliefernden Stoffwechsels in menschlichen Serum und in Leberpunkaten bei Leberkrankungen. *Klin Wschr* 36:280-287, 1958
- Zelman S, Wang CC: Transaminases in serum and liver correlated with liver cell necrosis in needle aspiration biopsies. *Am J Med Sci* 237:323-334, 1959
- Baier H, Hennemann G, Muhler E: Beitrag zur Aktivitätsbeurteilung chronischer Leberkrankungen durch Bestimmung von Serumenzymen. *Klin Wschr* 39:117-119, 1961
- Haber MM, West AB, Haber AD, Reuben A: Relationship of aminotransferase to liver histological status in chronic hepatitis C. *Am J Gastroenterol* 90:1250-1257, 1995
- McCormick SE, Goodman ZD, Maydonovitch CL, Sjogren MH: Evaluation of liver histology, ALT elevation, and HCV RNA titer in patients with chronic hepatitis C. *Am J Gastroenterol* 91:1516-1522, 1996
- Gordon SC, Fang JW, Silverman AL, McHutchison JG, Albrecht JK: The significance of baseline serum alanine aminotransferase on pretreatment disease characteristics and response to antiviral therapy in chronic hepatitis C. *Hepatology* 32:400-404, 2000
- Pradat P, Alberti A, Poynard T, Esteban JI, Weiland O, Marcellin P, Badalamenti S, Trepo C: Predictive value of ALT levels for histologic findings in chronic hepatitis C: a European collaborative study. *Hepatology* 36:973-977, 2002
- Zarski JP, Bohn B, Bastie A, Pawlowsky JM, Baud M, Bost-Bezeaux F, Tran van Nhieu J, Seigneurin JM, Buffet C, Dhumeaux D: Characteristics of patients with dual infection by hepatitis B and C viruses. *J Hepatol* 28:27-33, 1998
- Benhamou Y, Bochet M, Di Martino V, Charlotte F, Azria F, Coutellier A, Vidaud M, Bricaire F, Opolon P, Katlama C, Poynard T, for the MULTIVIRC Group: Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. *Hepatology* 30:1054-1058, 1999
- Piton A, Poynard T, Imbert-Bismut F, Khalil L, Delattre J, Pelissier E, Sansonetti N, Opolon P: Factors associated with serum alanine transaminase activity in healthy subject: consequence for the definition of normal values, for selection of blood donors, and for patients with chronic hepatitis C. *Hepatology* 27:1213-1219, 1998
- Prati D, Taioli E, Zanella A, Della Torre E, Butelli S, Del Vecchio E, Vianello L, Zanuso F, Mozzi F, Milani S, Conte D, Colombo M, Sirchia G: Updated definitions of healthy ranges for serum alanine aminotransferase levels. *Ann Intern Med* 137:1-9, 2002
- Kudryavtsev BN, Kudryavtseva MV, Sakuta GA, Stein GI: Human hepatocyte polyploidization kinetics in the course of life cycle. *Virchows Arch B Cell Pathol* 64:387-393, 1993

Prevalence and Clinical Implications of Occult Hepatitis B Viral Infection in Hemophilia Patients in Japan

Hidenori Toyoda,^{1,3*} Kazuhiko Hayashi,¹ Yoshiki Murakami,⁴ Takashi Honda,¹ Yoshiaki Katano,¹ Isao Nakano,¹ Hidemi Goto,¹ Takashi Kumada,³ and Junki Takamatsu²

¹Department of Gastroenterology, Nagoya University School of Medicine, Nagoya, Japan

²Department of Transfusion Medicine, Nagoya University School of Medicine, Nagoya, Japan

³Department of Gastroenterology, Ogaki Municipal Hospital, Ogaki, Japan

⁴Department of Gastroenterology, National Tsuruga Hospital, Tsuruga, Japan

The prevalence and clinical implications of occult hepatitis B virus (HBV) infection were investigated in the Japanese patients with hemophilia in whom a high prevalence of infection with transfusion-transmissible viruses has been reported. HBV DNA was detected in the sera of 22 of 43 (51.2%) patients with hemophilia who were negative for HBV surface antigen (HBs), indicating that these patients had occult HBV infection. No factor, including age, type or severity of hemophilia, presence of HBs or HBV core (HBc) antibody, or coinfection with hepatitis C virus (HCV) or human immunodeficiency virus (HIV) was associated with occult HBV infection, except for high anti-HBc titer and/or coinfection with HCV genotype 1 (1a or 1b). In general, occult HBV infection did not appear to have significant clinical implications. However, in patients in whom HBV was detected by PCR specific for the surface (S)-region, higher alanine aminotransferase levels were observed. The genotype of the occult HBV in the present study was exclusively the domestic type indigenous to Japan (genotype C), suggesting a different route of transmission for HBV in comparison to HCV and HIV in this population. *J. Med. Virol.* 73:195–199, 2004. © 2004 Wiley-Liss, Inc.

KEY WORDS: hemophilia; hepatitis B virus; occult infection; genotype; route of transmission

INTRODUCTION

Patients with hemophilia are at high risk of infection with parenterally transmissible viruses due to the frequent use of blood products. High prevalence of infection with hepatitis C virus (HCV) [Makris et al., 1990; Troisi et al., 1993], human immunodeficiency virus (HIV) [Tsuchie et al., 1985; Kroner et al., 1994],

and GB virus C (GBV-C) [Hanley et al., 1998; Toyoda et al., 1998] has been reported. A few studies have also been carried out on hepatitis B virus (HBV) infection in this population by serological evaluation [Kumar et al., 1993; Goedert et al., 2002].

Recently, several studies have reported occult HBV infection in subjects without HBV surface (HBs) antigen (HBsAg), and its clinical implications are suggested [Cacciola et al., 1999; Brechot et al., 2001; Torbenson and Thomas, 2002]. In addition to a history of frequent use of blood products, a large number of hemophilia patients are infected with HIV, which can result in dynamic changes in the immune status. Reactivation of HBV in association with changes in the immune status can occur and can cause liver damage, which sometimes results in liver failure [Xunrong et al., 2001]. Thus, hemophilia patients are a population in which the assessment of occult HBV infection is important, especially for those patients with HIV.

In the present study, we attempted to clarify the prevalence and clinical importance of occult HBV infection in the Japanese hemophilia patients without HBs antigen. As a result, occult HBV infection was found in around one-half of the patients.

PATIENTS AND METHODS

Patients

Among 44 patients with hemophilia who had been followed-up as outpatients at Nagoya University Hospital and who were admitted regularly to the hospital during 2002, 1 patient had HBs antigen and the other

*Correspondence to: Hidenori Toyoda, MD, PhD, Department of Gastroenterology, Ogaki Municipal Hospital, 4-86 Minaminkawa, Ogaki, Gifu, 503-8502, Japan.
E-mail: tkumada@he.mirai.ne.jp

Accepted 18 February 2004

DOI 10.1002/jmv.20075

Published online in Wiley InterScience
(www.interscience.wiley.com)

43 patients were negative for HBs antigen. These 43 patients were enrolled in this study. All were males with a mean age of 34.0 ± 12.1 -years-old. Thirty-four patients had hemophilia A, and the remaining 9 had hemophilia B. Thirty-seven patients had severe, 4 patients had moderate, and 2 patients had mild hemophilia. All 24 patients with HIV infection were receiving HAART therapy at the time of sampling serum. Eight of 38 patients with HCV infection had a history of interferon therapy, but no patients were treated with interferon at the time serum was sampled.

Evaluation for Coinfection With HIV and HCV

HIV infection was confirmed by anti-HIV1 antibody detected by particle agglutination (SERODIA-HIV, Fuji Rebio, Tokyo, Japan). Serum HIV RNA concentration was measured by the Amplicor HIV Monitor test (Roche Diagnostics K.K., Tokyo, Japan). The presence of HCV was confirmed by both an HCV antibody assay (2nd generation, Dinabot; Tokyo, Japan), and detection of HCV RNA by nested reverse transcription-polymerase chain reaction (RT-PCR) [Okamoto et al., 1990]. HCV genotypes, according to Simmonds et al. [1994] classification, were determined by RT-PCR with genotype-specific primers [Okamoto et al., 1996]. Serum HCV RNA concentrations were measured by Amplicor Monitor assay (Roche Diagnostics K.K., Tokyo).

Serological Tests for HBV Infection and Detection of HBV DNA

HBV serum markers (HBs antigen, HBs antibody, and HBV core [HBc] antibody) were examined by means of commercial immunoenzyme assays (Abbott Laboratories, North Chicago, IL). For detection of HBV DNA, extracted DNA was amplified by nested touchdown PCR [Don et al., 1991] with three independent primer sets specific for HBV surface (s)-(sense: 5'-CTCTTGTCCTCCAAATTTGTCCT-3' and antisense: 5'-CAGCAAAGCCAAAAGACCCAC-3' for the first PCR, and sense: 5'-AGGTA-TGTTGCCCGTTTGTCT-3' and antisense: 5'-GGGTTTAAATGTATACCCA-3' for the second PCR), core (c)-(sense: 5'-ACTGTTCAAGCCTCCAAGCT-3' for the first and second PCR, antisense: 5'-GGAATACTAACATTGAGATTCCCGAG-3' for the first PCR, and antisense: 5'-AGTGCGAATCCACACTC-3' for the second PCR), and X-regions (sense: 5'-TGCCAAGTGTGCTGACGC-3' for the first PCR, sense: 5'-CTGCCGATCCACTACTGCGGAAC-3' for the second PCR, and antisense: 5'-TTCCTGCAGTGAGACCACCGTGAACG-3' for the first and second PCR). HBV DNA was amplified from 100 ng of extracted DNA in a total volume of 50 μ l, in the presence of 10 pmol of each primer, 125 μ M dNTP, and 2.5 U Taq polymerase (Toyobo, Tokyo, Japan). PCR was performed in a DNA thermal cycler (Perkin-Elmer Cetus, Norwalk, CT). The PCR program consisted of 20 cycles of denaturation at 94°C for 30 sec, annealing at 65°C for 30 sec with a 0.5°C decrease per one cycle (55.5°C at final cycle), and extension at 70°C for 3 min with an initial denaturation at 94°C for 1 min, and a

subsequent 20 cycles of denaturation at 94°C for 30 sec, annealing at 55°C for 30 sec, and extension at 70°C for 3 min with a final extension at 70°C for 10 min. The same PCR program was used for both the first and second PCR amplifications. Amplified PCR products were analyzed by electrophoresis on 1.0% agarose gel and transferred to a Hybond-N+ nylon membrane (Amersham-Pharmacia, Buckinghamshire, UK). The amplified products were detected by hybridization with a specific probe based on the entire HBV sequences. This probe was generated with a DIG DNA Labeling and Detection kit (Roche Diagnostics, Mannheim, Germany). Results were considered to be valid only if identical results were obtained in at least two separate experiments.

The patients who were positive for both the S- and C-regions were considered to be positive for HBV. The patients who were positive for only one of the two regions examined were then referred to the result of PCR specific for the X-region, and HBV infection was confirmed according to this result.

Genotyping of HBV DNA

Genotyping of occult HBV was performed in seven patients in whom HBV DNA was detected by PCR specific for the S-region. For genotyping of HBV DNA, the PCR product from amplification of the S-region was sequenced directly, and phylogenetic analysis was performed with the neighbor-joining and bootstrap methods.

The entire protocol was approved by the hospital ethics committee and carried out in compliance with the Helsinki Declaration.

RESULTS

Serological Prevalence of Hepatitis B Viral Markers

Among the 43 patients with hemophilia but without HBs antigen, HBs antibody was detected in 27 patients (62.8%), and HBc antibody was detected in 37 patients (86.0%). There was no correlation between the type and/or severity of hemophilia and serological prevalence of HBV. There was no association between coinfections with HIV and/or HCV and HBV serological status.

Detection of HBV DNA in Serum

HBV DNA was detected in 22 of the 43 hemophilia patients (51.2%). HBV was detected by PCR specific for the S-region in 7 patients and by PCR specific for the C-region in 19 patients. In 4 patients HBV DNA was detected by both methods. In the 18 patients in whom HBV DNA was detected by only one of the PCR methods, additional PCR targeting the X-region was performed and all patients were positive for this region.

We compared the background, serological markers for HBV, and coinfection status between HBV DNA-positive and -negative patients (Table I). There were no significant differences in the background of patients, including age, and type or severity of hemophilia. Both