

70% of HCCs and that genes related to telomere synthesis, the Ras signaling pathway, and calcium signaling were recurrently affected.^{98,99} In particular, HBV integration into the *hTERT* gene was the first one found that was common to different HCCs. Two independent groups other than Brechot and colleagues have reported HBV integration into *hTERT* in HCCs and hepatoma cell lines,^{100,101} and one study demonstrated that expression of *hTERT* gene was *cis*-activated by HBV integration in vitro.¹⁰⁰ It is of note that viral integrations into *hTERT* were recurrently found in uterine cervical cancers with HPV infection,¹⁰¹ underlining the importance of viral integration and its insertional mutagenesis regardless of viral species or organ.

Recent reports revealed the second common integration site for HCC. Murakami et al. found HBV integration into the *MLL* gene in 1 of 68 cases,¹⁰² and Tamori et al. independently found it in 3 of 15 HCCs.¹⁰³ Taken together, HBV integration and the resulting insertional mutagenesis are not rare events, and they play a role in hepatocarcinogenesis possibly by producing fusion protein,^{104,105} by *cis*-activation of cellular genes, or by disrupting gene function.

Integration of HBV is not a late event during a course of chronic infection. One can identify HBV integration in chronic hepatitis tissues and even in tissues after acute self-limiting hepatitis.¹⁰⁶ We have analyzed host genes affected by HBV integration in chronic hepatitis tissues and identified candidate cellular genes related to cell growth and survival.¹⁰⁷

Pathogenesis of hepatitis B and natural history

Hepatitis B virus is not directly hepatotoxic. Many HBV carriers are asymptomatic and have no or minimal liver injury even with high viral replication. It has been demonstrated that host immune responses to viral antigens result in hepatocellular injury, and it is clear clarified that covalently closed circular DNA (cccDNA) plays an important role in maintaining chronic HBV infection.¹⁰⁸

Antiviral therapy

Recent studies have demonstrated that sustained viral suppression (<10⁵ copies/ml in serum HBV DNA) results in normal serum ALT and prevents progression to cirrhosis. Thus, the treatment goal is sustained viral suppression with antiviral therapy including interferon, lamivudine, adefovir dipivoxil, entecavir, and various combination therapies. We describe here antiviral treatment, including combination therapy, mainly focusing on the treatments available in Japan.

Interferon

Conventional interferon- α (IFN α), IFN β , and pegylated IFN α (PEG-IFN α) are available for treating chronic hepatitis B. IFN has many actions including antiviral and immunomodulatory effects.

Initially, IFN had been used for only 4 weeks in chronic hepatitis B patients in Japan, and its effects were limited, whereas 4–6 months of therapy was popular in many Western and Asian countries. With the latter regimen, HBeAg loss was achieved in approximately 33% of HBeAg-positive patients (three times that in controls),¹⁰⁹ and loss of HBsAg was noted in 7.8% (controls 1.8%) after IFN therapy.¹¹⁰ More than 12 months of therapy was more effective in HBeAg-positive patients with low serum HBV DNA levels.¹¹¹ The daily dosage of IFN was 3–10 MU thrice weekly.

A high seroconversion rate was noted in patients with high serum ALT levels, low serum HBV DNA levels, and moderate to severe hepatitis, whereas a lower response rate was observed in patients with lower baseline serum ALT levels (≤ 1.3 – 3.0 times the upper limit of normal),¹¹² high serum HBV DNA levels, and minimal inflammatory changes. Corticosteroid withdrawal therapy induced long-term clinical remission in chronic hepatitis B patients,¹¹³ and priming with a corticosteroid before IFN therapy resulted in a higher seroconversion rate.¹¹⁴ A long-term follow-up study demonstrated that the sustained virological response was 10%–15% within 4–6 months of treatment, 22% within 12 months, and 30% within 24 months.^{115–118} Furthermore, IFN-induced HBeAg seroconversion is durable and results in good overall survival and survival free of hepatic decompensation.^{119–121} It was reported that IFN therapy for cirrhotic patients significantly decreased the rate of HCC development, especially in patients with a high level of HBV DNA.¹²² HBsAg loss was seen in up to 10% of patients in Western countries but was rare in Asian patients.

Most HBV infections in Asian patients are acquired perinatally or during early childhood, and the distribution of their HBV genotype is quite different from that in Caucasian patients. More than 80% of HBV carriers in Asian countries including Japan naturally seroconvert by the age of 25–30 years. Thus, we must take into consideration the age and HBV genotype of those who are given antiviral therapy.

A pilot study suggested that IFN β is effective and safe for re-treating patients with chronic hepatitis B who had not responded to a previous IFN α cycle.¹²³

Recently, PEG-IFN α 2a^{19,124} and PEG-IFN α 2b¹²⁵ (long-acting forms of IFN α) have been used in both Asian and Caucasian patients but are still not available in Japan. Both 24-week and 52-week courses were found to be tolerable and produced a higher

seroconversion rate in Asian and Caucasian patients.^{19,124,125}

Lamivudine

Lamivudine, an oral nucleoside analogue, inhibits HBV replication. A daily dose of 100mg lamivudine markedly reduces the serum HBV DNA level. However, when short-term treatment is stopped, the serum HBV DNA levels generally return to pretreatment levels.^{8,126,127}

Lamivudine has usually been administered for 1 year. The first-year HBeAg seroconversion rate in 100mg lamivudine-treated patients with a pretreatment ALT level more than five times the upper limit of normal was 80%; there was no further increase in HBeAg seroconversion during a second year of therapy.¹²⁸ However, reappearance of HBeAg and hepatitis flares occurred at a high rate after seroconversion with lamivudine therapy for HBeAg-positive chronic hepatitis patients with high serum HBV DNA levels.^{129,130}

It is thought that prolonged therapy is needed in patients with low ALT levels or a long endogenous antiviral immune response.¹³¹ Three years of lamivudine therapy reduced necroinflammatory activity and reversed fibrosis formation in most patients, but the emergence of YMDD variants blunted the histological response.¹³² The response to lamivudine therapy in HBeAg-negative patients is similar to that of HBeAg-positive patients.^{133,134}

A lamivudine-resistant strain with altered YMDD motif of the polymerase gene (rtM204I and rtM204V with or without rtL180M) developed in 10%–20% after 1 year, 30%–40% after 3 years, and 50%–70% after 5 years, resulting in flare-up hepatitis due to resistant virus.^{129,135,136} A flare-up due to a YMDD mutant results in hepatic failure in some cirrhotic patients, who then require liver transplantation¹³⁷; however, most patients with flare-up hepatitis have a serum ALT level of <80IU/l.¹³⁶ It was reported that lamivudine initially selected wild-type virus from precore/core promoter mutants, but precore mutation reappeared during prolonged therapy.¹³⁸

Lamivudine resistance in HBV does not seem to depend on the HBV genotype, although it was significantly higher in the Ba ("a" means Asia) subgroup of HBV than in the Bj ("j" means Japan) subgroup.¹³⁹

A highly sensitive method to detect the YMDD motif mutant demonstrated that the mutant was noted in a few patients with HBeAb-positive chronic hepatitis B without previous administration of lamivudine.¹⁴⁰ YMDD motif mutants may be selected during continuing lamivudine therapy and elicit another hepatitis flare-up.^{19,140,141} Flare-up hepatitis develops when the serum YMDD motif mutant level is >10^{2.7} copies/ml.¹⁴²

Adefovir dipivoxil

Adefovir dipivoxil (Adefovir) is an acyclic nucleotide analogue, and it has been proven that adefovir is effective for both wild-type and lamivudine-resistant HBV strains. A daily dose of 10mg was recently approved for treatment of both HBeAg-positive and HBeAg-negative chronic hepatitis B patients.^{11,12,143,144} Histological improvement, HBV DNA suppression, ALT normalization, and HBsAg loss (1.6%–2.0% vs. 0%) was seen in HBeAg-positive and HBeAg-negative chronic hepatitis B patients.^{11,12} HBeAg loss and HBeAg seroconversion also increased compared to that in controls (12% vs. 6%).¹¹ There was no significant difference in the antiviral effect of adefovir among the various HBV genotypes.¹³ Combination therapy with lamivudine led to a stronger antiviral effect and achieved HBV DNA levels of <200 copies/ml, as seen by PCR. HBeAg seroconversion was 6%–8% after 1 year of combination therapy compared with 0%–2% with lamivudine monotherapy and 11% with adefovir monotherapy; at week 104, HBeAg seroconversion increased 12% on combination therapy.¹⁴⁴ A few patients showed a relapse after seroconversion when they stopped adefovir administration. Long-term adefovir therapy decreased the replicative form of HBV DNA (termed cccDNA) levels by a noncytolytic mechanism.¹⁴⁵

It has been reported that the rate of adefovir-resistant mutant appearance is low compared with that after lamivudine therapy. Recently, it was clarified that resistance mutations rtN236T and rtA181V were identified in 5.9% of HBeAg-negative chronic hepatitis B patients after 144 weeks.¹⁴⁶ Adefovir dipivoxil-resistant rtN236T mutant is susceptible to lamivudine and other nucleoside analogues, such as entecavir, emtricitabine, and telbivudine.^{147,148}

Disturbed renal function was reported with a daily dose of adefovir of 30mg but not with 10mg. Increased serum creatinine was reported in 2.5% when therapy was extended to 3 years, but it was reversible upon stopping therapy.¹⁴⁹

Entecavir

Entecavir became available for chronic hepatitis B patients in the United States in 2005 and might become available in Japan in 2006. A Phase III clinical trial demonstrated that entecavir was superior to lamivudine for reducing HBV DNA in both HBeAg-positive and HBeAg-negative patients.¹⁵⁰ Entecavir at daily doses of 1.0 and 0.5mg resulted in significantly greater reductions in the HBV DNA level and normalization of serum ALT levels than lamivudine 100mg daily after as little as 24 weeks of treatment.¹⁵¹ At 48 weeks, the

mean reductions in HBV DNA levels were 5.06, 4.46, and $2.86 \log_{10}$ copies/ml after entecavir 1.0, 0.5, and 0.1 mg, respectively, which is significantly higher than the $1.37 \log_{10}$ copies/ml achieved with lamivudine,¹⁵¹ and these amounts of entecavir were well tolerated. Entecavir 1 mg might be used for lamivudine-resistant mutants, and 0.5 mg may be suitable for the wild strain. In HBeAg-positive chronic hepatitis B patients, 96 weeks of treatment with entecavir 0.5 mg results in continued clinical benefit as measured by the reduction in serum HBV DNA (<300 copies/ml by PCR; 80% vs. 39%; $P < 0.0001$) and ALT levels and continued HBeAg seroconversion compared with lamivudine (31% vs. 25%, $P = \text{NS}$).¹⁵² No resistant mutants were noted during the 96 weeks of treatment. Entecavir demonstrated an overall safety profile comparable to that of lamivudine throughout the 96 weeks.¹⁵²

Combination therapy

Interferon and lamivudine

There have been many reports concerning the combination or sequential therapies with lamivudine and IFN;^{18,153–157} however, its efficacy is not certain and might be limited. We tried combination and sequential therapy with lamivudine and natural IFN α in genotype C patients according to the method by Serfaty et al.,¹⁸ but HBV DNA suppression, HBeAg negativity, and ALT normalization in our study were not comparable to their results (unpublished data). This discrepancy might be due to the differences in the distribution of HBV genotype (genotype A was prevalent in the study of Serfaty et al.) and in the mode of HBV transmission. Controlled studies demonstrated the efficacy of combination therapy for HBeAg-positive patients with high serum ALT levels^{153,154} but not for HBeAg-negative patients.^{156,158}

Pegylated interferon and lamivudine

As mentioned already, PEG-IFN might be more effective than conventional IFN α .^{19,124,125} Patients with HBeAg-negative chronic hepatitis B given PEG-IFN α 2a had significantly higher response rates (which were sustained 24 weeks after the cessation of therapy) than did patients given lamivudine. Addition of lamivudine to PEG-IFN α 2a did not improve the posttherapeutic response rate.^{19,159} In patients with HBeAg-positive chronic hepatitis, combination treatment of PEG-IFN α 2b (32 weeks) and lamivudine (52 weeks) produced a higher sustained virological response than did lamivudine monotherapy (52 weeks) after up to 3 years after treatment.¹⁶⁰ At the end of treatment, HBeAg loss occurred in 63% of patients in the combination group and in 28% of patients in the lamivudine group ($P = 0.0001$). The probabilities of

sustained response for combination treatment and lamivudine monotherapy were, respectively, 33% and 13% at week 24, 31% and 11% at week 52, and 29% and 9% at week 76 (log-rank test, $P = 0.0015$).

Other antiviral agents

Many promising nucleoside and nucleotide analogues for chronic hepatitis B are being evaluated in Phase I, II, and III studies. Unfortunately, these studies are not ongoing in Japan. Telbivudine suppresses wild-type HBV by $5\text{--}8 \log_{10}$ and was more potent than lamivudine in a Phase II study.¹⁶¹ Clevudine 30 mg/day for 24 weeks resulted in an HBV DNA reduction of $4.46 \log_{10}$ undetectable by PCR in 59%, HBeAg loss in 24%, and ALT normalization in 76%.¹⁶² Tenofovir disoproxil fumarate show strong suppression of HBV with YMDD motif mutants and has a good safety record.¹⁶³

References

- Okamoto H, Tsuda F, Sakugawa H, Sastrosoewignjo RI, Imai M, Miyakawa Y, et al. Typing hepatitis B virus by homology in nucleotide sequence: comparison of surface antigen subtypes. *J Gen Virol* 1998;69:2575–83.
- Norder H, Courouce AM, Magnius LO. Complete genotypes, phylogenetic relatedness, and structural proteins of six strains of the hepatitis B virus, four of which represent two new genotypes. *Virology* 1994;198:489–503.
- Stuyver L, De Grandt S, Van Geyt C, Zoulin F, Fried M, Schinazi RF, et al. A new genotype of hepatitis B virus: complete genome and phylogenetic relatedness. *J Gen Virol* 2000;81:67–74.
- Arauz-Ruiz P, Norder H, Robertson BH, Magnius LO. A new American genotype of hepatitis B virus revealed in Central America. *J Gen Virol* 2002;83:2059–73.
- Orito E, Ichida T, Sakugawa H, Sata M, Horiike N, Hino K, et al. Geographic distribution of hepatitis B virus (HBV) genotype in patients with chronic HBV infection in Japan. *Hepatology* 2001;34:590–4.
- Miyakawa Y, Mizokami M. Classifying hepatitis B. *Intervirology* 2003;46:329–38.
- Sugauchi F, Orito E, Ichida T, Kato H, Sakugawa H, Kakumu S, et al. Hepatitis B virus genotype B with or without recombination with genotype C over the precore region plus the core gene. *J Virol* 2002;76:5985–92.
- Lai CL, Chien RN, Leung NW, Chang TT, Guan R, Tai DI, et al. A one-year trial of lamivudine for chronic hepatitis B: Asian Hepatitis Lamivudine Study Group. *N Engl J Med* 1998;339:61–8.
- Dienstag JL, Schiff ER, Wright TL, Perrillo RP, Hann HW, Goodman Z, et al. Lamivudine as initial treatment for chronic hepatitis B in the United States. *N Engl J Med* 1999;341:1256–63.
- Chayama K, Suzuki Y, Kobayashi M, Kobayashi M, Tsubota A, Hashimoto M, et al. Emergence and takeover of YMDD motif mutant hepatitis B virus during long-term lamivudine therapy and takeover by wild type after cessation of therapy. *Hepatology* 1998;27:1711–6.
- Marcellin P, Chang TT, Lim SG, Tong MJ, Sievert W, Shiffman ML, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med* 2003;348:808–16.

12. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med* 2003;348:800-7.
13. Westland C, Delaney W 4th, Yang H, Chen SS, Marcellin P, Hadziyannis S, et al. Hepatitis B virus genotypes and virological response in 694 patients in phase III studies of adefovir dipivoxil. *Gastroenterology* 2003;125:107-16.
14. Balfour HHJ. Antiviral drugs. *N Engl J Med* 1999;340:1255-68.
15. Colonno RJ, Genovesi EV, Medina I, Lamb L, Durham SK, Huang ML, et al. Long-term entecavir treatment results in sustained antiviral efficacy and prolonged life span in woodchuck model of chronic hepatitis infection. *J Infect Dis* 2001;184:1236-45.
16. Perry CM, Jarvis B. Peginterferon-alpha 2a (40kDa): a review of its use in the management of chronic hepatitis C. *Drugs* 2001;61:2263-88.
17. Glue P, Fang JW, Rouzier-Panis R, Raffanel C, Sabo R, Gupta SK, et al. Pegylated interferon alfa-2b: pharmacokinetics, safety and preliminary efficacy data. *Clin Pharmacol Ther* 2000;68:556-67.
18. Serfaty L, Thabut D, Zoulim F, Andreani T, Chazouilleres O, Carbonell N, et al. Sequential treatment with lamivudine and interferon monotherapies in patients with chronic hepatitis B not responding to interferon alone: results of a pilot study. *Hepatology* 2001;34:573-7.
19. Marcellin P, Lau GK, Bonino F, Farci P, Hadziyannis S, Jin R, et al. Peginterferon alfa-2a alone, lamivudine alone, and the two in combination in patients with HBeAg-negative chronic hepatitis B. *N Engl J Med* 2004;351:1206-17.
20. Lau GK, Piratvisuth T, Luo KX, Marcellin P, Thongsawat S, Cooksley G, et al. Peginterferon alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2005;352:2682-95.
21. Orito E, Mizokami M. Hepatitis B virus genotypes and hepatocellular carcinoma. *Intervirology* 2003;46:408-12.
22. Sumi H, Yokosuka O, Seki N, Arai M, Imazeki F, Kurihara T, et al. Influence of hepatitis B virus genotypes on the progression of chronic type B liver disease. *Hepatology* 2003;37:19-26.
23. Orito E, Mizokami M, Sakugawa H, Michitaka K, Ishikawa K, Ichida T, et al. A case-control study for clinical and molecular biological differences between hepatitis B viruses of genotypes B and C. *Hepatology* 2001;33:218-23.
24. Usuda S, Okamoto H, Iwanari H, Baba K, Tsuda F, Miyakawa Y, et al. Serological detection of hepatitis B virus genotypes by ELISA with monoclonal antibodies to type-specific epitopes in the preS2-region product. *J Virol Methods* 1999;80:97-112.
25. Usuda S, Okamoto H, Tanaka T, Kidd-Ljunggren K, Holland PV, Miyakawa Y, et al. Differentiation of hepatitis B virus genotypes D and E by ELISA using monoclonal antibodies to epitopes on the preS2-region product. *J Virol Methods* 2001;87:81-9.
26. Sakugawa H, Nakasone H, Nakayoshi T, Orito E, Mizokami M, Yamashiro T, et al. Preponderance of hepatitis B virus genotype B contributes to a better prognosis of chronic HBV infection in Okinawa, Japan. *J Med Virol* 2002;67:484-9.
27. Ding X, Mizokami M, Yao G, Xu B, Orito E, Ueda R, et al. Hepatitis B virus genotype distribution among chronic hepatitis B virus carriers in Shanghai, China. *Intervirology* 2001;44:43-7.
28. Kao JH, Chen PJ, Lai MY, Chen DS. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology* 2000;118:554-9.
29. Chu CJ, Keeffe EB, Han SH, Perrillo RP, Min AD, Soldevilla-Pico C, et al. Hepatitis B virus genotypes in the United States: results of a nationwide study. *Gastroenterology* 2003;125:444-51.
30. Lindh M, Hannoun C, Dhillon AP, Norkrans G, Horal P. Core promoter mutations and genotypes in relation to viral replication and liver damage in East Asian hepatitis B virus carriers. *J Infect Dis* 1999;179:775-82.
31. Chu CJ, Hussain M, Lok AS. Hepatitis B virus genotype B is associated with earlier HBeAg seroconversion compared with hepatitis B virus genotype C. *Gastroenterology* 2002;122:1756-62.
32. Nakayoshi T, Maeshiro T, Nakayoshi T, Nakasone H, Sakugawa H, Kinjo F, et al. Difference in prognosis between patients infected with hepatitis B virus with genotype B and those with genotype C in the Okinawa Islands: a prospective study. *J Med Virol* 2003;70:350-4.
33. Kao JH, Chen PJ, Lai MY, Chen DS. Hepatitis B virus genotypes and spontaneous hepatitis B e antigen seroconversion in Taiwanese hepatitis B carriers. *J Med Virol* 2004;72:363-9.
34. Ni YH, Chang MH, Wang KJ, Hsu HY, Chen HL, Kao JH, et al. Clinical relevance of hepatitis B virus genotype in children with chronic infection and hepatocellular carcinoma. *Gastroenterology* 2004;127:1733-8.
35. Thakur V, Guptan RC, Kazim SN, Malhorta V, Sarin SK. Profile, spectrum and significance of HBV genotypes in chronic liver disease patients in the Indian subcontinent. *J Gastroenterol Hepatol* 2002;17:165-70.
36. Mayerat C, Mantegani A, Frei PC. Does hepatitis B virus (HBV) genotype influence the clinical outcome of HBV infection? *J Viral Hepatol* 1999;6:299-304.
37. Kikuchi K, Niituma H, Ishii M, Cervantes JG, Hong S, Ojima T, et al. Genoepidemiology and its relationship to clinical features in patients infected chronically with hepatitis B virus (HBV). *Hepatol Res* 2000;17:43-55.
38. Kobayashi M, Arase Y, Ikeda K, Tsubota A, Suzuki Y, Saitoh S, et al. Clinical characteristics of patients infected with hepatitis B virus genotypes A, B, and C. *J Gastroenterol* 2002;37:35-9.
39. Kobayashi M, Arase Y, Ikeda K, Tsubota A, Suzuki Y, Hosaka T, et al. Clinical features of hepatitis B virus genotype A in Japanese patients. *J Gastroenterol* 2003;38:656-62.
40. Kobayashi M, Suzuki F, Arase Y, Akuta N, Suzuki Y, Hosaka T, et al. Infection with hepatitis B virus genotype A in Tokyo, Japan during 1976 through 2001. *J Gastroenterol* 2004;39:844-50.
41. Carman WF, Jacyna MR, Hadziyannis S, Karayiannis P, McGarvey MJ, Makris A, et al. Mutation preventing formation of hepatitis B e antigen in patients with chronic hepatitis B infection. *Lancet* 1989;2:588-91.
42. Li JS, Tong SP, Wen YM, Vitvitski L, Zhang Q, Treppe C. Hepatitis B virus genotype A rarely circulates as an HBe-minus mutant: possible contribution of a single nucleotide in the precore region. *J Virol* 1993;67:5402-10.
43. Lok AS, Akarca U, Greene S. Mutations in the precore region of hepatitis B virus serve to enhance the stability of the secondary structure of the pre-genome encapsidation signal. *Proc Natl Acad Sci USA* 1994;91:4077-81.
44. Buckwold VE, Xu Z, Chen M, Yen TS, Ou JH. Effects of a naturally occurring mutation in the hepatitis B virus basal core promoter on precore gene expression and viral replication. *J Virol* 1996;70:5845-51.
45. Scaglioni PP, Melegari M, Wands JR. Biologic properties of hepatitis B viral genomes with mutations in the precore promoter and precore open reading frame. *Virology* 1997;233:374-81.
46. Yu X, Mertz JE. Distinct modes of regulation of transcription of hepatitis B virus by the nuclear receptors HNF4alpha and COUP-TF1. *J Virol* 2003;77:2489-99.
47. Gunther S, Piwon N, Will H. Wild-type levels of pregenomic RNA and replication but reduced pre-C RNA and e-antigen synthesis of hepatitis B virus with C(1653) → T, A(1762) → T and G(1764) → A mutations in the core promoter. *J Gen Virol* 1998;79:375-80.
48. Tang H, Raney AK, Mclachlan A. Replication of the wild type and a natural hepatitis B virus nucleocapsid promoter variant is differentially regulated by nuclear hormone receptors in cell culture. *J Virol* 2001;75:8937-48.

89. Tokino T, Matsubara K. Chromosomal sites for hepatitis B virus integration in human hepatocellular carcinoma. *J Virol* 1991; 65:6761-4.
90. Suzuki T, Shen H, Akagi K, Morse HC, Malley JD, Naiman DQ, et al. New genes involved in cancer identified by retroviral tagging. *Nat Genet* 2002;32:166-74.
91. Shen H, Suzuki T, Munroe DJ, Stewart C, Rasmussen L, Gilbert DJ, et al. Common sites of retroviral integration in mouse hematopoietic tumors identified by high-throughput, single nucleotide polymorphism-based mapping and bacterial artificial chromosome hybridization. *J Virol* 2003;77:1584-8.
92. Broussard DR, Lozano MM, Dudley JP. Rorgamma (Rorc) is a common integration site in type B leukemogenic virus-induced T-cell lymphomas. *J Virol* 2004;78:4943-6.
93. Cousens C, Bishop JV, Philbey AW, Gill CA, Palmarini M, Carlson JO, et al. Analysis of integration sites of Jaagsiekte sheep retrovirus in ovine pulmonary adenocarcinoma. *J Virol* 2004;78:8506-12.
94. Kustikova O, Fehse B, Modlich U, Yang M, Dullmann J, Kamino K, et al. Clonal dominance of hematopoietic stem cells triggered by retroviral gene marking. *Science* 2005;308:1171-4.
95. Collier LS, Carlson CM, Ravimohan S, Dupuy AJ, Largaespada DA. Cancer gene discovery in solid tumours using transposon-based somatic mutagenesis in the mouse. *Nature* 2005;436:272-6.
96. Hacein-Bey-Abina S, Von Kalle C, Schmidt M, McCormack MP, Wulffraat N, Leboulch P, et al. LMO2-associated clonal T cell proliferation in two patients after gene therapy for SCID-X1. *Science* 2003;302:415-9.
97. Minami M, Poussin K, Brechot C, Paterlini P. A novel PCR technique using Alu-specific primers to identify unknown flanking sequences from the human genome. *Genomics* 1995;29:403-8.
98. Gozuacik D, Murakami Y, Saigo K, Chami M, Mugnier C, Lagorce D, et al. Identification of human cancer-related genes by naturally occurring hepatitis B virus DNA tagging. *Oncogene* 2001;20:6233-40.
99. Paterlini-Brechot P, Saigo K, Murakami Y, Chami M, Gozuacik D, Mugnier C, et al. Hepatitis B virus-related insertional mutagenesis occurs frequently in human liver cancers and recurrently targets human telomerase gene. *Oncogene* 2003;22:3911-6.
100. Horikawa I, Barrett JC. cis-Activation of the human telomerase gene (hTERT) by the hepatitis B virus genome. *J Natl Cancer Inst* 2001;93:117-3.
101. Ferber MJ, Montoya DP, Yu C, Aderca I, McGee A, Thorland EC, et al. Integrations of the hepatitis B virus (HBV) and human papillomavirus (HPV) into the human telomerase reverse transcriptase (hTERT) gene in liver and cervical cancers. *Oncogene* 2003;22:3813-20.
102. Murakami Y, Saigo K, Takashima H, Minami M, Okanou T, Brechot C, et al. Large scaled analysis of hepatitis B virus (HBV) DNA integration in HBV related hepatocellular carcinomas. *Gut* 2005;54:1162-8.
103. Tamori A, Yamanishi Y, Kawashima S, Kanehisa M, Enomoto M, Tanaka H, et al. Alteration of gene expression in human hepatocellular carcinoma with integrated hepatitis B virus DNA. *Clin Cancer Res* 2005;11:5821-6.
104. Wang J, Zindy F, Chenivresse X, Lamas E, Henglein B, Brechot C. Modification of cyclin A expression by hepatitis B virus DNA integration in a hepatocellular carcinoma. *Oncogene* 1992;7:1653-6.
105. De The H, Marchio A, Tiollais P, Dejean A. A novel steroid thyroid hormone receptor-related gene inappropriately expressed in human hepatocellular carcinoma. *Nature* 1987;330:667-70.
106. Murakami Y, Minami M, Daimon Y, Okanou T. Hepatitis B virus DNA in liver, serum, and peripheral blood mononuclear cells after the clearance of serum hepatitis B virus surface antigen. *J Med Virol* 2004;72:203-14.
107. Minami M, Daimon Y, Mori K, Takashima H, Nakajima T, Itoh Y, et al. Hepatitis B virus-related insertional mutagenesis in chronic hepatitis B patients as an early drastic genetic change leading to hepatocarcinogenesis. *Oncogene* 2005;24:4340-8.
108. Yokosuka O, Omata M, Imazeki F, Okuda K, Summers J. Changes of hepatitis B virus DNA in liver and serum caused by recombinant leukocyte interferon treatment: analysis of intrahepatic replicative hepatitis B virus DNA. *Hepatology* 1985;5:728-34.
109. Guan R. Interferon monotherapy in chronic hepatitis B. *J Gastroenterol Hepatol* 2000;15(Suppl):E34-40.
110. Wong DK, Cheung AM, O'Rourke K, Naylor CD, Detsky AS, Heathcote J. Effect of alpha-interferon treatment in patients with hepatitis B e antigen-positive chronic hepatitis B: a meta analysis. *Ann Intern Med* 1999;119:312-23.
111. Janssen HL, Gerken G, Carreno V, Marcellin P, Naoumov NV, Craxi A, et al. Interferon alpha for chronic hepatitis B infection: increased efficacy of prolonged treatment: the European Concerted Action on Viral Hepatitis (EUROHEP). *Hepatology* 1999;30:238-43.
112. Liaw YF, Lin SM, Chen TJ, Chien RN, Sheen IS, Chu CM. Beneficial effect of prednisolone withdrawal followed by lymphoblastoid interferon on the treatment of chronic hepatitis B in Asians: a randomized controlled trial. *J Hepatol* 1994;20:175-80.
113. Akuta N, Suzuki F, Tsubota A, Arase Y, Suzuki Y, Someya T, et al. Long-term clinical remission induced by corticosteroid withdrawal therapy (CSWT) in patients with chronic hepatitis B infection: a randomized controlled trial-CSWT with and without follow-up interferon alpha therapy. *Dig Dis Sci* 2002;47:405-14.
114. Lin SM, Sheen IS, Chien RN, Chu CM, Liaw YF. Long-term beneficial effect of interferon therapy in patients with chronic hepatitis B virus infection. *Hepatology* 1999;29:971-5.
115. Manesis EK, Hadziyannis SJ. Interferon alpha treatment and retreatment of hepatitis B e antigen negative chronic hepatitis B mutants. *Gastroenterology* 2001;121:101-9.
116. Lampertico P, Del Ninno E, Manzin A, Donato MF, Rumi MG, Lunghi G, et al. A randomized controlled trial of a 24-month course of interferon alpha 2b in patients with chronic hepatitis B who had hepatitis B virus DNA without hepatitis B e antigen in serum. *Hepatology* 1997;26:1621-35.
117. Papatheododdis GV, Manesis E, Hadziyannis SJ. The long term outcome of interferon-alpha treated and untreated patients with HBeAg-negative chronic hepatitis B. *J Hepatol* 2001;34:306-13.
118. Brunetto MR, Oliveri F, Coco B, Leandro G, Colombatto P, Gorin JM, et al. The outcome of chronic anti-HBe positive chronic hepatitis B in alpha interferon treated and untreated patients: a long term cohort study. *J Hepatol* 2002;36:263-70.
119. Niederau C, Heintges T, Lange S, Goldmann G, Niederau CM, Mohr L, et al. Long-term follow-up of HBeAg-positive patients treated with interferon alpha for chronic hepatitis B. *N Engl J Med* 1996;334:1422-7.
120. Lin SM, Tai DI, Chien RN, Sheen IS, Chu CM, Liaw YF. Comparison of long-term effects of lymphoblastoid interferon alpha and recombinant interferon alpha-2a therapy in patients with chronic hepatitis B. *J Viral Hepatol* 2004;11:349-57.
121. Van Zonneveld M, Honkoop P, Hansen BE, Niesters HG, Murad SD, de Man RA, et al. Long-term follow-up of alpha-interferon treatment of patients with chronic hepatitis B. *Hepatology* 2004;39:804-10.
122. Ikeda K, Saitoh S, Suzuki Y, Kobayashi M, Tsubota A, Fukuda M, et al. Interferon decreases hepatocellular carcinogenesis in patients with cirrhosis caused by the hepatitis B virus. *Cancer* 1998;82:827-35.
123. Munoz R, Castellano G, Fernandez I, Alvarez MV, Manzano ML, Marcos MS, et al. A pilot study of beta-interferon for treatment of patients with chronic hepatitis B who failed to respond to alpha-interferon. *J Hepatol* 2002;37:655-9.
124. Cooksley WG, Piratvisuth T, Lee SD, Mahachai V, Chao YC, Tanwadee T, et al. Peginterferon alpha-2a (40kDa): an ad-

- vance in the treatment of hepatitis B e antigen-positive chronic hepatitis B. *J Viral Hepatol* 2003;10:298-305.
125. Janssen HL, van Zonneveld M, Senturk H, Zeuzem S, Akarca US, Cakaloglu Y, et al. Pegylated interferon alpha-2b alone or in combination with lamivudine for HBeAg-positive chronic hepatitis B: a randomized trial. *Lancet* 2005;365:123-9.
 126. Dienstag JL, Perrillo RP, Schiff ER, Bartholomew M, Vicary C, Rubin M. A preliminary trial of lamivudine for chronic hepatitis B infection. *N Engl J Med* 1995;333:1657-61.
 127. Nevens F, Main J, Honkoop P, Tyrrell DL, Barber J, Sullivan MT, et al. Lamivudine therapy for chronic hepatitis B: six month dose-ranging study. *Gastroenterology* 1997;113:1258-63.
 128. Liaw YF, Leung NW, Chang TT, Guan R, Tai DI, Ng KY, et al. Effects of extended lamivudine therapy in Asian patients with chronic hepatitis B. *Gastroenterology* 2000;119:172-80.
 129. Yao GB, Cui ZY, Wang BE, Yao JL, Zeng MD. A 3-year clinical trial of lamivudine in treatment of patients with chronic hepatitis B. *Hepatobiliary Pancreat Dis Int* 2004;3:188-93.
 130. Song BC, Suh DJ, Lee HC, Chung YH, Lee YS. Hepatitis B e antigen seroconversion after lamivudine therapy is not durable in patients with chronic hepatitis B in Korea. *Hepatology* 2000;32:803-6.
 131. Chien RN, Liaw YF, Atkins M. Pretherapy alanine aminotransferase level as a determinant for hepatitis B e antigen seroconversion during lamivudine therapy in patients with chronic hepatitis B. *Hepatology* 1999;30:770-4.
 132. Dienstag JL, Goldin RD, Heathcote EJ, Hann HW, Woessner M, Stephenson SL, et al. Histological outcome during long-term lamivudine therapy. *Gastroenterology* 2003;124:105-17.
 133. Tassopoulos NC, Volpes R, Pastore G, Heathcote J, Buti M, Goldin RD, et al. Efficacy of lamivudine in patients with hepatitis B e antigen-negative/hepatitis B virus DNA-positive (precore mutant) chronic hepatitis B. *Hepatology* 1999;29:889-96.
 134. Rizzetto M, Tassopoulos NC, Goldin RD, Esteban R, Santantonio T, Heathcote EJ, et al. Extended lamivudine treatment in patients with HBeAg-negative chronic hepatitis B. *J Hepatol* 2005;42:173-9.
 135. Chang TT, Lai CL, Chien RN, Guan R, Lim SG, Lee CM, et al. Four years of lamivudine treatment in Chinese patients with chronic hepatitis B. *J Gastroenterol Hepatol* 2004;19:1276-82.
 136. Lok AS, Lai CL, Leung N, Yao GB, Cui ZY, Schiff ER, et al. Long-term safety of lamivudine treatment in patients with chronic hepatitis B. *Gastroenterology* 2003;125:1714-22.
 137. Yuen MF, Kato T, Mizokami M, Chan AO, Yuen JC, Yuan HJ, et al. Clinical outcome and virologic profiles of severe hepatitis B exacerbation due to YMDD mutants. *J Hepatol* 2003;39:850-5.
 138. Suzuki F, Suzuki Y, Tsubota A, Akuta N, Someya T, Kobayashi M, et al. Mutations of polymerase, precore and core promoter gene in hepatitis B virus during 5-year lamivudine therapy. *J Hepatol* 2002;37:824-30.
 139. Akuta N, Suzuki F, Kobayashi M, Tsubota A, Suzuki Y, Hosaka T, et al. The influence of hepatitis B virus genotype on the development of lamivudine resistance during long-term treatment. *J Hepatol* 2003;38:315-21.
 140. Kirishima T, Okanoue T, Daimon Y, Itoh Y, Nakamura H, Morita A, et al. Detection of YMDD motif mutant using a novel sensitive method in chronic liver disease type B patients before and during lamivudine treatment. *J Hepatol* 2002;37:259-65.
 141. Yeh CT, Chien RN, Chu CM, Liaw YF. Clearance of the original hepatitis B virus YMDD-motif mutants with emergence of distinct lamivudine-resistant mutants during prolonged lamivudine therapy. *Hepatology* 2000;31:1318-26.
 142. Mori K, Minami M, Kirishima T, Kunimoto K, Okita M, Nakayama M, et al. Prediction of breakthrough hepatitis due to lamivudine-resistant hepatitis B virus by a sensitive semi-quantitative assay using peptide nucleic acids. *Intervirology* (in press).
 143. Peters MG, Hann HWH, Martin P, Heathcote EJ, Buggisch P, Rubin R, et al. Adefovir dipivoxil alone or in combination with lamivudine in patients with lamivudine-resistant chronic hepatitis B. *Gastroenterology* 2004;126:91-101.
 144. Perrillo R, Hann HW, Mutimer D, Willems B, Leung N, Lee WM, et al. Adefovir dipivoxil added to ongoing lamivudine in chronic hepatitis B with YMDD mutant hepatitis B virus. *Gastroenterology* 2004;126:81-90.
 145. Werle-Lapostolle B, Bowden S, Locarnini S, Wurstthorn K, Petersen J, Lau G, et al. Persistence of cccDNA during the natural history of chronic hepatitis B and decline during adefovir dipivoxil therapy. *Gastroenterology* 2004;126:1750-8.
 146. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, et al. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B. *N Engl J Med* 2005;352:2673-81.
 147. Angus P, Vaughan R, Xiong S, Yang H, Delaney W, Gibbs C, et al. Resistance to adefovir dipivoxil therapy associated with the selection of a novel mutation in the HBV polymerase. *Gastroenterology* 2003;125:292-7.
 148. Yang H, Qi X, Das K, Arnold E, Westland CE, Delaney IV WE, et al. In vitro characterization and molecular modeling analysis of a novel adefovir resistance mutation rtN236T in the HBV polymerase. *J Hepatol* 2004;40(Suppl 1):A114 (abstract 382).
 149. Hadziyannis S, Tassopoulos N, Chang TT, Heathcote J, Kitis G, Rizzetto M, et al. Three year study of adefovir dipivoxil (ADV) demonstrates sustained efficacy of in presumed precore mutant chronic hepatitis B patients in a long term safety and efficacy study. *J Hepatol* 2004;40(Suppl 1):17 (abstract 46).
 150. Lai CL, Rosmawati M, Lao J, Van Vlierberghe H, Anderson FH, Thomas N, et al. Entecavir is superior to lamivudine in reducing hepatitis B virus DNA in chronic hepatitis B infection. *Gastroenterology* 2002;123:1831-8.
 151. Chang TT, Gish RG, Hadziyannis SJ, Cianciara J, Rizzetto M, Schiff ER, et al. A dose-ranging study of the efficacy and tolerability of entecavir in lamivudine-refractory chronic hepatitis B patients. *Gastroenterology* 2005;129:1198-209.
 152. Gish RG, Chang TT, De Man RA, Gadano A, Sollano J, Han KH, et al. Entecavir results in substantial virologic and biochemical improvement and HBeAg seroconversion through 96 weeks of treatment in HBeAg(+) chronic hepatitis B patients (study ETV-022). *Hepatology* 2005;42(Suppl 1):267A (abstract 181).
 153. Schalm SW, Heathcote J, Cianciara J, Farrell G, Sherman M, Willems B, et al. Lamivudine and alpha interferon combination therapy of patients with chronic hepatitis B infection: a randomized trial. *Gut* 2000;46:562-8.
 154. Barbaro G, Zechini F, Pellicelli AM, Francavilla R, Scotto G, Bacca D, et al. Long-term efficacy of interferon alpha-2b and lamivudine in combination compared to lamivudine monotherapy in patients with chronic hepatitis B: an Italian multicenter, randomized trial. *J Hepatol* 1998;28:923-9.
 155. Tatulli I, Francavilla R, Rizzo GL, Vinciguerra V, Ierardi E, Amoroso A, et al. Lamivudine and alpha-interferon in combination long term for precore mutant chronic hepatitis B. *J Hepatol* 2001;35:805-10.
 156. Santantonio T, Niro GA, Sinisi E, Leandro G, Insalata M, Guastadisegni A, et al. Lamivudine/interferon combination therapy in anti-HBe positive chronic hepatitis B patients: a controlled pilot study. *J Hepatol* 2002;36:799-804.
 157. Schiff ER, Dienstag JL, Karayalcin S, Grimm IS, Perrillo RP, Husa P, et al. Lamivudine and 24 weeks lamivudine/interferon combination therapy for hepatitis B e antigen-positive chronic hepatitis B in interferon nonresponders. *J Hepatol* 2003;38:818-26.
 158. Akarca US, Ersoz G, Gunsar F, Karasu Z, Saritas E, Yuce G, et al. Interferon-lamivudine combination is no better than lamivudine alone in anti-HBe-positive chronic hepatitis B. *Antivir Ther* 2004;9:325-34.
 159. Sypsa VA, Mimidis K, Tassopoulos NC, Chrysagis D, Vassiliadis T, Moulakakis A, et al. A viral kinetic study using pegylated

- interferon alfa-2b and /or lamivudine in patients with chronic hepatitis B/HBeAg negative. *Hepatology* 2005;42:77-85.
160. Chan HL, Hui AY, Wong VW, Chim AM, Wong ML, Sung JJ. Long-term follow-up of peginterferon and lamivudine combination treatment in HBeAg-positive chronic hepatitis B. *Hepatology* 2005;41:1357-64.
 161. Lai CL, Leung N, Teo EK, Tong M, Wong F, Hann HW, et al. A 1-year trial of telbivudine, lamivudine, and combination in patients with hepatitis B e antigen-positive chronic hepatitis. *Gastroenterology* 2005;129:528-36.
 162. Lee HS, Chung YH, Lee KS, Byun KS, Paik SW, Han JY, et al. A 12-week clevudine therapy showed durable antiviral activity and normalization of alanine transaminase levels for 6 months after discontinuation of treatment in patients with chronic hepatitis B. *Hepatology* 2004;40(Suppl 1):652A (abstract 1130).
 163. Van Bommel F, Wunsche T, Mauss S, Reinke P, Bergk A, Schurmann D, et al. Comparison of adefovir and tenofovir in the treatment of lamivudine-resistant hepatitis B virus infection. *Hepatology* 2004;40:1421-5.

First phase viral kinetic parameters and prediction of response to interferon alpha-2b/ribavirin combination therapy in patients with chronic hepatitis C

Akiko Makiyama*, Yoshito Itoh, Kohichiroh Yasui, Kohjiroh Mori, Mika Okita, Mika Nakayama, Junko Yamaoka, Masahito Minami, Tomoki Nakajima, Takeshi Okanoue

Molecular Gastroenterology and Hepatology, Kyoto Prefectural University of Medicine, Graduate School of Medical Science, Kawaramachi-Hirokouji, Kamigyou-ku, Kyoto 602-8566, Japan

Received 17 January 2006; received in revised form 15 June 2006; accepted 3 July 2006
Available online 6 September 2006

Abstract

The aim of the present study was to assess parameters in early phase HCV dynamics for predicting the outcome of interferon (IFN)/ribavirin combination therapy in patients with chronic hepatitis C (CH-C). Sixty-five CH-C patients who received IFN alpha-2b/ribavirin combination therapy were enrolled. The serum levels of HCV RNA 0 h and 3 months after commencing therapy were serially quantified. HCV kinetic parameters such as quantity, ratio of decline, and half-life were analyzed. In genotype 1 patients, both the quantity and the ratio of decline of HCV RNA 24 h after the start of therapy were useful predictors of a poor response. No patients who had serum HCV RNA above 200 KIU/ml 24 h after the start of therapy achieved a sustained viral response (SVR). In genotype 2 patients, conversely, these two parameters were predictors of a sustained viral response. The efficacy of these parameters in predicting the outcome of therapy was comparable to that of the disappearance of HCV RNA from sera at 4 weeks. These results demonstrate that parameters of HCV kinetics 24 h after the start of therapy are useful for the early prediction of outcome in response to IFN alpha-2b/ribavirin combination therapy.
© 2006 Elsevier Ireland Ltd. All rights reserved.

Keywords: Chronic hepatitis C; HCV kinetics; Interferon; Ribavirin; Early prediction

1. Introduction

Interferon (IFN) is used to treat patients with chronic hepatitis C (CH-C) worldwide. However, most patients with genotype 1b and high hepatitis C virus (HCV) loads did not benefit from IFN monotherapy. In recent years, IFN/ribavirin (Rib) and peginterferon/Rib combination therapy are often used to treat patients with genotype 1, however, more than half of the patients did not succeed in clearing HCV RNA from the sera [1,2].

IFN therapy is expensive, requires a long period of treatment, and sometimes is accompanied by severe adverse

effects. Until recently, the most widely accepted factors for predicting the outcome of IFN or IFN/Rib combination therapy include HCV genotype, quantity of HCV RNA, mutations in the non-structural 5A (NS5A) region of HCV, histological staging, age, and duration of HCV infection before therapy [1–8]. The early identification of non-responders (NRs) soon after initiating the combination therapy is worthwhile because it provides an indication as to when it may be advisable to discontinue unnecessary therapy. Early identification of responders motivates patients to adhere to therapy.

In CH-C patients, a biphasic decline in the serum levels of HCV RNA after initiating IFN or IFN/Rib combination therapy is well known [9]. A third phase decline has also been recently reported [10]. The first phase decline occurs within 24 h of starting therapy, and it is thought to occur due

* Corresponding author. Tel.: +81 75 251 5519; fax: +81 75 251 0710.
E-mail address: akiko@koto.kpu-m.ac.jp (A. Makiyama).

to the IFN-mediated direct reduction of HCV RNA from the sera. The second phase decline is observed after 24 h and is thought to reflect the death of HCV-infected hepatocytes. After starting IFN or IFN/Rib combination therapy, the disappearance of HCV RNA at 2, 4, or 12 weeks is a useful predictor of responsiveness to treatment [11–23] and the second phase HCV decline is associated with sustained response [24].

Several recent reports have indicated that the first phase viral decline is also useful for early prediction of the response to IFN or IFN/Rib combination therapy [25,26], however, a general consensus has not been obtained. In the present study, we investigated HCV kinetics in Japanese patients treated with IFN/Rib combination therapy and we confirmed that first phase viral kinetics are useful for predicting treatment outcome.

2. Methods

2.1. Patients

Sixty-five patients with CH-C were enrolled in this study. All patients were admitted to and followed at the outpatient clinic of the University Hospital of Kyoto Prefectural University of Medicine between December 2001 and September 2003. They consisted of 38 men and 27 women, ranging from 27 to 70 years old [54.2 ± 9.7 (mean \pm S.D.)]. All patients were positive for anti-HCV antibody and serum HCV RNA and all had elevated serum alanine aminotransferase (ALT) levels for at least 6 months. They were also negative for hepatitis B virus surface antigen and human immunodeficiency virus. Patients who had co-existing liver diseases such as autoimmune hepatitis, primary biliary cirrhosis, or evidence for alcohol abuse were excluded from this study. Liver needle biopsy was performed prior to IFN therapy and the histological diagnoses were reached according to the classification of Desmet [27]. Sustained viral responders (SVRs) were defined as those who showed negative serum HCV RNA for 6 months after finishing the combination therapy. The rest of the patients were regarded as non-responders. Informed consent was obtained from all participants and the study protocol was approved by the ethical committee of the university.

2.2. Study protocol

Six MU of IFN alpha-2b (Intoron-A, Schering-Plough Corp., Kenilworth, NJ) was injected intramuscularly daily for 2 weeks, then switched to thrice weekly for 22 weeks in combination with Rib (Rebetol, Schering-Plough Corp., Kenilworth, NJ). Rib was given orally at a dose of either 800 mg (body weight \geq 60 kg) or 600 mg (body weight $<$ 60 kg) daily for 24 weeks. Blood samples were obtained at 0, 6, 12 and 24 h; at 2, 4, 7, and 14 days; and at 1, 3, 6, and 12 months after the initiation of combination therapy.

2.3. Quantification and determination of HCV RNA and genotyping

Serum HCV RNA levels were determined by use of the Amplicor GT HCV monitor (Roche Diagnostic Systems, Tokyo, Japan). The detection range of this assay was between 0.5–850 KIU/ml. When the serum HCV RNA level was below 0.5 KIU/ml, the existence of serum HCV RNA was determined by reverse transcription-nested polymerase chain reaction (RT-nested PCR) using the Amplicor HCV v2.0 (Roche Diagnostic Systems, Tokyo, Japan), which had a detection limit of 50 IU/ml. HCV genotypes 1 and 2 were determined by a serologic genotyping assay [28–30]. Genotypes 1 and 2 in this assay correspond to genotypes 1 (1a, 1b) and 2 (2a, 2b) respectively proposed by Simmonds et al. [31].

2.4. HCV kinetic parameters

We determined the half-lives of HCV RNA in the first and second phases using logarithmic approximate curves for the two phases. Samples containing HCV RNA greater than 850 KIU/ml or less than 0.5 KIU/ml were omitted from the curve and the upward curves were also omitted.

2.5. Statistical analysis

Positive predictive value (PPV) was calculated as the percentage of SVRs among patients who were predicted to have a sustained viral response (SVR). Negative predictive value (NPV) was calculated as the percentage of NRs among patients not meeting the criteria for prediction of a SVR. *p*-values were calculated by Fischer's exact probability test and Mann–Whitney *U*-test. Statistical significance was set at $p < 0.05$.

3. Results

Baseline characteristics of 65 chronic hepatitis C patients (49 genotype 1 patients and 16 genotype 2 patients) who received IFN/Rib combination therapy were shown in Table 1. As have been already known [3–6], SVRs had less advanced staging, lower HCV load, and increased ratio of genotype 2.

HCV dynamics during combination therapy are presented in four groups depending on the genotype and response to the therapy (Fig. 1A–D). In genotype 1 patients, the mean half-life of HCV RNA in the first phase was 5.5 ± 1.5 h in SVRs, which was significantly ($p = 0.0361$) shorter than that of NRs (9.8 ± 10.0 h). The mean half-life of HCV RNA in the second phase was 121.4 ± 104.2 h in SVRs, which was also significantly ($p = 0.0003$) shorter than that of NRs (470.0 ± 752.3 h). Although the half-life of HCV RNA in SVRs in both the first and second phase was significantly shorter than that of genotype 1 patients who were NRs, the

Table 1
Characteristic of chronic hepatitis C patients

	SVR (n = 23)	NR (n = 42)	p
Gender (male/female)	13/10	25/17	>0.9999
Age (years) ^a	55.2 ± 6.9	53.6 ± 11.0	0.9890
Weight (kg) ^a	61.4 ± 10.3	64.3 ± 10.3	0.2301
BMI ^a	23.0 ± 2.6	23.9 ± 2.6	0.1514
ALT (IU/ml) ^a	122.4 ± 79.9	122.4 ± 103.8	0.6706
AST (IU/ml) ^a	88.1 ± 42.9	98.2 ± 79.6	0.8584
PLT (×10 ⁴ /μl) ^a	16.3 ± 5.6	14.0 ± 5.0	0.1765
Fibrosis (stage)			
1	7	7	
2	10	15	0.0398
3	4	18	
Not available	2	2	
Viral load (KIU/ml) ^a	440.3 ± 308.3	669.6 ± 197.7	0.0066
Genotype (1/2)	10/13	39/3	<0.0001

ALT: alanine aminotransferase; AST: aspartate aminotransferase; BMI: body mass index; SVR: sustained viral responder (serum HCV RNA was negative after 6 months from the end of therapy); NR: non-responder (serum HCV RNA was positive after 6 months from the end of therapy).

^a Data are expressed as mean ± S.D. *p*-values were calculated by Fischer's exact probability test and Mann–Whitney *U*-test.

Table 2
Mean half-life of HCV dynamics during combination therapy

	SVR	NR	p
Genotype 1, n = 49			
First phase (hour)	n = 10 5.5 ± 1.5	n = 39 9.8 ± 10.0	0.0361
Second phase (hour)	121.4 ± 104.2	470.0 ± 752.3	0.0003
Genotype 2, n = 16			
First phase (hour)	n = 13 5.8 ± 4.2	n = 3 9.1 ± 1.4	0.0693
Second phase (hour)	110.9 ± 124.9	437.9 ± 320.6	0.0734

First phase is from before IFN therapy to 24 h after the start of combination therapy. Second phase is from 24 h to 3 months after the initiation of therapy. *p*-values were calculated by Mann–Whitney *U*-test.

half-life of HCV RNA in genotype 2 patients did not differ significantly between SVRs and NRs partly because of the small number of patients with genotype 2 (Table 2).

Among the 49 patients with genotype 1, 8 patients were negative for serum HCV RNA after 4 weeks of treatment, and a SVR was achieved in 5 of 8 patients. The PPV, as determined from the disappearance of serum HCV RNA after 4 weeks of treatment was 62.5% for SVRs, and the NPV was 90.2%. Six patients showed serum HCV RNA levels less than 5.0 KIU/ml after the first 24 h of therapy, and SVR was achieved in 4 patients. The PPV determined using HCV RNA levels less




Fig. 1. (A–D) Serum HCV RNA dynamics during the first 3 months of IFN alpha-2b and Rib combination therapy. (A) Sustained viral responders (SVRs) in genotype 1. (B) Non-responders (NR) in genotype 1. (C) Sustained viral responders (SVR) in genotype 2. (D) Non-responders (NR) in genotype 2.

Table 3
Predicting SVR and NR using HCV kinetics

Criteria for SVR prediction	PPV (%)	NPV (%)	Sensitivity (%)	Specificity (%)
Genotype 1 (<i>n</i> = 49)				
Negative for serum HCV RNA after 4 week of treatment	62.5 (5/8)	90.2 (37/41)	50.0 (5/10)	94.9 (36/39)
Viral levels at 24 h after the initiation of therapy <5.0 KIU/ml	66.7 (4/6)	86.0 (37/43)	40.0 (4/10)	94.9 (37/39)
The ratio of decrease in serum HCV RNA (24 h/0 h) >1.5 log	36.8 (7/19)	90.0 (27/30)	70.0 (7/10)	69.2 (27/39)
Genotype 2 (<i>n</i> = 16)				
Negative for serum HCV RNA after 4 week of treatment	100 (10/10)	50.0 (3/6)	76.9 (10/13)	100 (3/3)
Viral levels at 24 h after the initiation therapy <50 KIU/ml	100 (12/12)	75.0 (3/4)	92.3 (12/13)	100 (3/3)
The ratio of decrease in serum HCV RNA (24 h/0 h) >1 log	100 (10/10)	50.0 (3/6)	76.9 (10/13)	100 (3/3)

PPV: positive predictive value (% meeting the criteria for SVR prediction that were SVR). NPV: negative predictive value (% not meeting the criteria for SVR prediction that were NR).

than 5.0 KIU/ml after the first 24 h of the therapy was 66.7%, and the NPV was 86.0%. Decrease in the serum levels of HCV RNA (24 h/0 h) greater than 1.5 log were seen in 19 patients, and SVR was achieved in 7 patients. The PPV calculated based on the decrease in serum HCV RNA (24 h/0 h) was 36.8% in SVRs, and NPV was 90.0% (Table 3).

In genotype 2 patients, HCV dynamics did not differ significantly between SVRs and NRs at either phase. Among the 16 patients with genotype 2, 10 patients who had negative serum HCV RNA after 4 weeks of treatment all achieved SVR. The PPV calculated using negative serum HCV RNA after 4 weeks of treatment was 100% in SVRs, and the NPV was 50.0%. Twelve patients with serum HCV RNA levels lower than 50 KIU/ml after the first 24 h of treatment all achieved SVR. The PPV of SVRs, as determined using serum HCV RNA levels lower than 50 KIU/ml after the first 24 h of treatment, was 100%, and the NPV was 75.0%. Decrease in serum HCV RNA levels (24 h/0 h) greater than 1.0 log was seen in 10 patients and all of them achieved SVR. The PPV of SVRs, determined by a decrease in serum HCV RNA (24 h/0 h) of greater than 1.0 log was 100%, and the NPV was 50.0% (Table 3).

Because patients with genotype 1 and high HCV load showed a poor response to IFN/Rib combination therapy, we assessed the HCV load threshold in genotype 1 SVRs before or early after starting the combination therapy. Before therapy, the maximum serum HCV RNA level in SVRs was greater than 850 KIU/ml, which demonstrated that IFN/Rib combination therapy was effective in some patients with genotype 1 and very high HCV load. After initiating the combination therapy, the maximum serum HCV RNA levels in genotype 1 SVRs were 170 KIU/ml (24 h), 130 KIU/ml (2 d) and 140 KIU/ml (4 d) (Fig. 1A). Among genotype 1 patients, those who showed serum HCV RNA levels greater than 200 KIU/ml at 24 h after starting combination therapy did not achieve SVR (Fig. 1A).

4. Discussion

Biphasic decline in the serum levels of HCV RNA during IFN therapy for CH-C was reported by our group [9] and

Neumann et al. [32]. Subsequent studies showed the second phase decline in the serum HCV RNA level to be an excellent predictor of SVR in IFN mono-therapy [24] and IFN/Rib combination therapy. For CH-C patients as well as for their physicians, easy evaluation of the probability of SVR/non-response (NR) soon after starting therapy is desirable both from a financial and emotional standpoint.

In the present study, we showed that HCV viral load and viral decline within the first 24 h are good predictors of NR in genotype 1 patients. These parameters are also good predictors of SVR in genotype 2 patients receiving IFN/Rib combination therapy, and their predictive value was comparable to that of the disappearance of HCV RNA from the sera after the first 4 weeks of treatment (Table 3).

Our findings are consistent with previous reports by Jessner et al. [25,33] who demonstrated that NR to IFN/Rib combination therapy could be predicted when the HCV viral load declined by less than 70% of baseline levels after a single injection of 5 MU of IFN alpha-2b [25] and that the decline in viral load 24 h after a single test dose of standard IFN had also high predictive value for the outcome of peginterferon alpha-2a/Rib combination therapy [33]. Layden et al. [26] reported that only those patients with 24 h viral load less than 250 KIU/ml and 24 h viral declines of greater than 98% after 7.5–15 µg/ml consensus IFN achieve SVR [26]. The authors therefore advocated the use of the 250 KIU/ml at 24 h of the therapy threshold for achieving SVR. Karino et al. [34] reported that the decline in serum HCV RNA levels 24 h after the first injection of IFN was regulated by HCV genotypes and the extent of hepatic fibrosis [34]. In the present study, however, hepatic fibrosis had no relation to the 24 h viral load or viral decline (data not shown).

One of the most important effects of Rib on HCV dynamics is that it shortens the half-life of serum HCV RNA in the second phase of IFN/Rib combination therapy. It is therefore unlikely that Rib would have exerted a significant influence on the decline of serum HCV RNA after a single oral administration. There are at least two explanations for why HCV RNA levels after the first injection of IFN can predict the effectiveness of combination therapy. Firstly, serum HCV RNA levels may have to be below a certain threshold at the first phase of HCV kinetics during IFN/Rib combination therapy

to eradicate HCV in the second phase. Secondly, individual sensitivity to IFN may regulate the antiviral effect of IFN/Rib combination therapy.

Forns et al. [35] reported that “a viral load decrease of ≥ 2 log 10 at week 4 was the strongest predictor of virological response, when HCV-infected cirrhotic patients awaiting liver transplantation were treated with IFN/Rib combination therapy” [35]. Early prediction of the outcome at 24 h after starting IFN/Rib combination therapy may also be useful for avoiding unnecessary therapy prior to liver transplantation when IFN and Rib is administered to HCV-positive patients with liver cirrhosis or HCC whose white blood cells or platelet counts are reduced. It also motivates those patients who are likely to clear HCV before transplantation to adhere to the therapy. Further studies are required to clarify these issues.

We assessed the utility of evaluating the probability of SVR/NR soon after starting therapy and we found that those patients with serum HCV RNA levels greater than 200 KIU/ml at 24 h after starting combination therapy did not achieve SVR (Fig. 1A and C). This threshold of HCV RNA is recommended because patients with a low probability of SVR could be judged easily. Unexpectedly, patients with greater than 850 KIU/ml of HCV genotype 1b were verified to be capable of achieving SVR when the serum HCV RNA levels were lower than 200 KIU/ml at 24 h after the initiation of therapy.

In conclusion, quantity and the ratio of decline of HCV RNA after the first 24 h of the IFN/Rib combination therapy were a useful predictor of NR in genotype 1 patients and an accurate predictor of SVR in genotype 2 patients. No patients with serum HCV RNA greater than 200 KIU/ml 24 h after starting therapy achieved SVR. These results demonstrate the usefulness of HCV kinetic parameters after 24 h of therapy for the early prediction of the outcome of IFN alpha-2b/Rib combination therapy. In Japan, standard therapy for chronic hepatitis C patients with genotype 1 and high viral load had been the IFN/ribavirin combination therapy with around 20% of SVR rate until December 2004. With the advent of peginterferon/ribavirin combination therapy for 48 weeks, which is the standard therapy for these patients now in Japan, SVR rate has risen to approximately 50%. Now, the early prediction of the response to peginterferon/Rib combination therapy in patients with CH-C is under investigation.

References

- [1] Manns MP, McHutchison JG, Gordon SC, et al. Peginterferon alpha-2b plus ribavirin compared with interferon alpha-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomized trial. *Lancet* 2001;358:958–65.
- [2] Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon alpha-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–82.
- [3] Martinot-Peignoux M, Marcellin P, Pouteau M, et al. Pretreatment serum hepatitis C virus RNA levels and hepatitis C virus genotype are the main and independent prognostic factors of sustained response to interferon alfa therapy in chronic hepatitis C. *Hepatology* 1995;22:1050–6.
- [4] Tsubota A, Chayama K, Ikeda K, et al. Factors predictive of response to interferon-alpha therapy in hepatitis C virus infection. *Hepatology* 1994;19:1088–94.
- [5] Yamada G, Takatani M, Kishi F, et al. Efficacy of interferon alfa therapy in chronic hepatitis C patients depends primarily on hepatitis C virus RNA level. *Hepatology* 1995;22:1351–4.
- [6] Shiratori Y, Kato N, Yokosuka O, et al. Predictors of the efficacy of interferon therapy in chronic hepatitis C virus infection. Tokyo-Chiba Hepatitis Research Group. *Gastroenterology* 1997;113:558–66.
- [7] Enomoto N, Sakuma I, Asahina Y, et al. Mutations in nonstructural protein 5A gene and response to interferon in patients with chronic hepatitis C virus 1b infection. *N Engl J Med* 1996;334:77–81.
- [8] Murakami T, Enomoto N, Kurosaki M, et al. Mutations in nonstructural protein 5A gene and response to interferon in hepatitis C virus genotype 2 infection. *Hepatology* 1999;30:1045–53.
- [9] Yasui K, Okanoue T, Murakami Y, et al. Dynamics of hepatitis C viremia following interferon- α administration. *J Infect Dis* 1998;177:1475–9.
- [10] Herrmann E, Lee JH, Marinos G, Modi M, et al. Effect of ribavirin on hepatitis C viral kinetics in patients treated with pegylated interferon. *Hepatology* 2003;37:1351–8.
- [11] Castro FJ, Esteban JI, Juarez A, et al. Early detection of nonresponse to interferon plus ribavirin combination treatment of chronic hepatitis C. *J Viral Hepatol* 2002;9:202–7.
- [12] Brouwer JT, Hansen BE, Niesters HG, Schalm SW. Early prediction of response in interferon monotherapy and in interferon-ribavirin combination therapy for chronic hepatitis C: HCV RNA at 4 weeks versus ALT. *J Hepatol* 1999;30:192–8.
- [13] Di Marco V, Ferraro D, Almasio P, et al. Early viral clearance and sustained response in chronic hepatitis C: a controlled trial of interferon and ribavirin after high-dose interferon induction. *J Viral Hepatol* 2002;9:354–9.
- [14] McHutchison JG, Shad JA, Gordon SC, et al. Predicting response to initial therapy with interferon plus ribavirin in chronic hepatitis C using serum HCV RNA results during therapy. *J Viral Hepatol* 2001;8:414–20.
- [15] Castro FJ, Esteban JI, Sauleda S, et al. Utility of early testing for HCV viremia as predictive factor of sustained response during interferon or interferon plus ribavirin treatment. *J Hepatol* 2000;32:843–9.
- [16] Rosen HR, Ribeiro RR, Weiberger L, et al. Early hepatitis C viral kinetics correlate with long-term outcome in patients receiving high dose induction followed by combination interferon and ribavirin therapy. *J Hepatol* 2002;37:124–30.
- [17] McHutchison J, Blatt L, Sedghi-Vaziri A, Russell J, Schmid P, Conrad A. Is there an optimal time to measure quantitative HCV RNA to predict non-response following interferon treatment for chronic HCV infection? *J Hepatol* 1998;29:362–8.
- [18] Kakumu S, Aiyama T, Okumura A, Iwata K, Ishikawa T, Yoshioka K. Earlier loss of hepatitis C virus RNA in interferon therapy can predict a long-term response in chronic hepatitis C. *J Gastroenterol Hepatol* 1997;12:468–72.
- [19] Orito E, Mizokami M, Suzuki K, et al. Loss of serum HCV RNA at week 4 of interferon-alfa therapy is associated with more favorable long-term response in patients with chronic hepatitis C. *J Med Virol* 1995;46:109–15.
- [20] Knolle PA, Kremp S, Hohler T, Krummenauer F, Schirmacher P, Gerken G. Viral and host factors in the prediction of response to interferon-alpha therapy in chronic hepatitis C after long-term follow-up. *J Viral Hepatol* 1998;5:399–406.
- [21] Karino Y, Toyota J, Sugawara M, et al. Early loss of serum hepatitis C virus RNA can predict a sustained response to interferon therapy in patients with chronic hepatitis C. *Am J Gastroenterol* 1997;92:61–115.
- [22] Zeuzem S, Lee JH, Franke A, et al. Quantification of the initial decline of serum hepatitis C virus RNA and response to interferon alfa. *Hepatology* 1998;27:1149–56.

- [23] Gavier B, Martinez-Gonzalez MA, Reizu-Boj JI, et al. Viremia after one month of interferon therapy predicts treatment outcome in patients with chronic hepatitis C. *Gastroenterology* 1997;113:1647–53.
- [24] Layden TJ, Mika B, Wiley TE. Hepatitis C kinetics: mathematical modeling of viral response to therapy. *Semin Liver Dis* 2000;20:173–83.
- [25] Jessner W, Gschwantler M, Steindl-Munda P, et al. Primary interferon resistance and treatment response in chronic hepatitis C infection: a pilot study. *Lancet* 2001;358:1241–2.
- [26] Layden JE, Layden TJ, Reddy KR, Levy-Drummer RS, Poulakos J, Neumann AU. First phase viral kinetic parameters as predictors of treatment response and their influence on the second phase viral decline. *J Viral Hepat* 2002;9:340–5.
- [27] Desmet VJ. Histological classification of chronic hepatitis. *Acta Gastroenterol Belg* 1997;60:259–67.
- [28] Tsukiyama-Kohara K, Yamaguchi K, Maki N, et al. Antigenicities of Group 1 and 2 hepatitis C virus polypeptides: molecular basis of diagnosis. *Virology* 1993;192:430–7.
- [29] Tanaka T, Tsukiyama-Kohara K, Yamaguchi K, et al. Significance of specific antibody assay for genotyping of hepatitis C virus. *Hepatology* 1994;19:1347–53.
- [30] Kohara M, Tanaka T, Tsukiyama-Kohara K, et al. Hepatitis C virus genotypes 1 and 2 respond to interferon-alpha with different virologic kinetics. *J Infect Dis* 1995;172:934–8.
- [31] Simmonds P, Alberti A, Alter HJ, et al. A proposed system for the nomenclature of hepatitis C viral genotypes. *Hepatology* 1994;19:1321–4.
- [32] Neumann AU, Lam NP, Dahari H, et al. Hepatitis C viral dynamics in vivo and the antiviral efficacy of interferon-alpha therapy. *Science* 1998;280:103–7.
- [33] Jessner W, Stauber R, Hackl F, et al. Early viral kinetics on treatment with pegylated interferon-alpha-2a in chronic hepatitis C virus genotype 1 infection. *J Viral Hepat* 2003;10:37–42.
- [34] Karino Y, Toyota J, Sugawara M, et al. Hepatitis C virus genotypes and hepatic fibrosis regulate 24-h decline of serum hepatitis C virus RNA during interferon therapy in patients with chronic hepatitis C. *J Gastroenterol Hepatol* 2003;18:404–10.
- [35] Forn X, Garcia-Retortillo M, Serrano T, et al. Antiviral therapy of patients with decompensated cirrhosis to prevent recurrence of hepatitis C after liver transplantation. *J Hepatol* 2003;39:389–96.

Influence of Genotypes and Precore Mutations on Fulminant or Chronic Outcome of Acute Hepatitis B Virus Infection

Atsushi Ozasa,^{1,2} Yasuhito Tanaka,¹ Etsuro Orito,² Masaya Sugiyama,¹ Jong-Hon Kang,³ Shuhei Hige,⁴ Tomoyuki Kuramitsu,⁵ Kazuyuki Suzuki,⁶ Eiji Tanaka,⁷ Shunichi Okada,⁸ Hajime Tokita,⁹ Yasuhiro Asahina,¹⁰ Kazuaki Inoue,¹¹ Shinichi Kakumu,¹² Takeshi Okanoue,¹³ Yoshikazu Murawaki,¹⁴ Keisuke Hino,¹⁵ Morikazu Onji,¹⁶ Hiroshi Yatsuhashi,¹⁷ Hiroshi Sakugawa,¹⁸ Yuzo Miyakawa,¹⁹ Ryuzo Ueda,² and Masashi Mizokami¹

The outcome of acute hepatitis B virus (HBV) infection is variable, influenced by host and viral factors. From 1982 through 2004, 301 patients with acute HBV infection entered a multi-center cross-sectional study in Japan. Patients with fulminant hepatitis (n = 40) were older (44.7 ± 16.3 vs. 36.0 ± 14.3 years, $P < .0017$), less predominantly male (43% vs. 71%, $P = .0005$), less positive for hepatitis B e antigen (HBeAg) (23% vs. 60%, $P < .0001$), less infected with subgenotype Ae (0% vs. 13%, $P < .05$), and more frequently with Bj (30% vs. 4%, $P < .0001$) than those with acute self-limited hepatitis (n = 261). Precore (G1896A) and core-promoter (A1762T/G1764A) mutations were more frequent in patients with fulminant than acute self-limited hepatitis (53% vs. 9% and 50% vs. 17%, $P < .0001$ for both). HBV infection persisted in only three (1%) patients, and they represented 2 of the 23 infected with Ae and 1 of the 187 with the other subgenotypes (9% vs. 0.5%, $P = .032$); none of them received antiviral therapy. In multivariate analysis, age 34 years or older, Bj, HBeAg-negative, total bilirubin 10.0 mg/dL or greater, and G1896A mutation were independently associated with the fulminant outcome. In *in vitro* transfection experiments, the replication of Bj clone was markedly enhanced by introducing either G1896A or A1762T/G1764A mutation. **In conclusion**, persistence of HBV was rare (1%) and associated with Ae, whereas fulminant hepatitis was frequent (13%) and associated with Bj and lack of HBeAg as well as high replication due to precore mutation in patients with acute HBV infection. *Supplementary material for this article can be found on the HEPATOLOGY website (<http://interscience.wiley.com/jpages/0270-9139/suppmat/index.html>). (HEPATOLOGY 2006; 44:326-334.)*

Abbreviations: HBV, hepatitis B virus; HBeAg, hepatitis B e antigen; HBe, hepatitis B core antigen; HBsAg, hepatitis B surface antigen; ELA, enzyme immunoassay; PCR, polymerase chain reaction; RFLP, restriction fragment length polymorphism; ALT, alanine aminotransferase.

From the ¹Department of Clinical Molecular Informative Medicine and the ²Department of Internal Medicine and Molecular Science, Nagoya City University Graduate School of Medical Sciences, Nagoya, Japan; ³Teinekeijinkai Hospital, Sapporo, Japan; the ⁴Department of Internal Medicine, Hokkaido University Graduate School of Medicine, Sapporo, Japan; ⁵Akita City Hospital, Akita, Japan; the ⁶First Department of Internal Medicine, Iwate Medical University, Morioka, Japan; ⁷Shinshu University Graduate School of Medicine, Matsumoto, Japan; ⁸University of Yamanashi, Yamanashi, Japan; ⁹National Tokyo Hospital, Kiyose, Tokyo, Japan; ¹⁰Musashino Red Cross Hospital, Musashino, Tokyo, Japan; ¹¹Showa University Fujigaoka Hospital, Yokohama, Japan; ¹²Aichi Medical University School of Medicine, Aichi, Japan; ¹³Kyoto Prefectural University of Medicine, Kyoto, Japan; ¹⁴Tottori University, Tottori, Japan; ¹⁵Yamaguchi University School of Medicine, Yamaguchi, Japan; ¹⁶Ehime University School of Medicine, Matsuyama, Japan; ¹⁷National Hospital Organization Nagasaki Medical Center, Nagasaki, Japan; ¹⁸University of the Ryukyus, Okinawa, Japan; and ¹⁹Miyakawa Memorial Research Foundation, Tokyo, Japan.

Received February 8, 2006; accepted April 27, 2006.

Supported in part by a grant-in-aid from the Ministry of Health, Labour and Welfare of Japan (H16-kaken-3), Uehara Memorial Foundation, Toyoaki Foundation, and Miyakawa Memorial Research Foundation.

The nucleotide sequences of HBV DNA isolates used in this study have been deposited in the international DNA database under accession numbers AB249373-AB249636.

Address reprint requests to: Masashi Mizokami, M.D., Ph.D., Department of Clinical Molecular Informative Medicine, Nagoya, City University Graduate School of Medical Sciences, Kawasumi, Mizuho, Nagoya 467-8601, Japan. E-mail: mizokami@med.nagoya-cu.ac.jp; fax: (81) 52-842-0021.

Copyright © 2006 by the American Association for the Study of Liver Diseases.

Published online in Wiley InterScience (www.interscience.wiley.com).

DOI 10.1002/hep.21249

Potential conflict of interest: Nothing to report.

Approximately 3 billion people, one half of the world population, have been exposed to hepatitis B virus (HBV), of whom approximately 350 million are persistently infected with it.¹ Acute infection with HBV resolves in the great majority but can induce fulminant hepatitis or go on to become chronic. Host and viral factors may influence fulminant or chronic outcome of acute HBV infection, but they are not fully defined.

Eight genotypes have been detected by a sequence divergence greater than 8% in the entire HBV genome of approximately 3,200 nucleotides (nt), and designated by capital alphabet letters from A (HBV/A) to H in the order of documentation.²⁻⁵ They have distinct geographical distributions associated with severity of liver disease as well as response to antiviral therapies.⁶⁻⁸ Furthermore, subgenotypes have been reported for HBV/A, B, and C and named Aa/A1 (Asian/African type) and Ae/A2 (European type),⁹ Bj/B1 (Japanese type) and Ba/B2 (Asian type),¹⁰ as well as Cs/C1 (Southeast Asian type) and Ce/C2 (East Asian type).¹¹⁻¹³ Increasing lines of evidence indicate that subgenotypes of HBV/A and B influence the replication of HBV and bear clinical relevance.¹⁴⁻¹⁶ Furthermore, genotypes affect mutations in precore region and core promoter, thereby influencing the expression of hepatitis B e antigen (HBeAg).^{8,17}

During the 23 years from 1982 to 2004, a multi-center cross-sectional study was conducted throughout Japan on 301 patients with acute hepatitis B. We examined the influence of genotypes/subgenotypes on their fulminant or chronic outcome. Furthermore, the influence of G1896A or A1762T/G1764A on replication of HBV was evaluated in an *in vitro* replication model.

Patients and Methods

Patients With Acute Hepatitis B. During 1982 through 2004, 336 consecutive cases of acute hepatitis B were registered in 16 hospitals throughout Japan. These hospitals were from the following eight areas: Hokkaido (represented by J.-H. K. and S.H.), Tohoku (T.K. and K.S.), Kanto (H.T., Y.A. and K.I.), Koshin (E.T. and S.O), Tokai (A.O., Y.T., E.O., M.S., R.U., M.M., and S.K.), Kinki (T.O.), Honshu/Shikoku (Y.M., K.H., and M.O.), and Kyushu (H.Y. and H.S.). The diagnosis of acute hepatitis B was contingent on a sudden onset of clinical symptoms of hepatitis and detection of high-titered antibody to hepatitis B core antigen (anti-HBc) of IgM class in serum. Patients with initial high-titered anti-HBc ($\geq 90\%$ inhibition by a 1:200 diluted serum) were excluded; they were diagnosed as exacerbation of chronic hepatitis B. Patients with acute hepatitis A, hepatitis C, or human immunodeficiency virus co-infection, and drug-

or alcohol-induced acute hepatitis also were excluded; hepatitis D virus infection was not examined because of its extreme rarity in Japan.¹⁸ Most of them were followed for clinical outcomes until the disappearance of hepatitis B surface antigen (HBsAg) during 24 weeks or longer after the presentation. The criteria of fulminant hepatitis are based on the report by Trey et al.,¹⁹ with a slight modification in 1981 (Inuyama symposium, Aichi, Japan): coma of grade II or higher and prothrombin time less than 40% developing within 8 weeks after the onset. Serum samples were collected at the presentation and had been stored at -80°C . HBV genotypes, HBV DNA, and HBeAg were determined, and clinical outcomes of acute hepatitis were analyzed. The study protocol conformed to the 1975 Declaration of Helsinki, and was approved by the Ethics Committees of the institutions. Every patient gave an informed consent for this study.

Serological Markers of HBV Infection. HBsAg was determined by hemagglutination (MyCell; Institute of Immunology Co., Ltd., Tokyo, Japan) or enzyme immunoassay (EIA) (Axsym; Abbott Japan, Tokyo, Japan), and HBeAg by enzyme-linked immunosorbent assay (F-HBe; Kokusai Diagnostic, Kobe, Japan) or chemiluminescent EIA (Fujirebio Inc., Tokyo, Japan). Anti-HBc of IgM and IgG classes were determined by radioimmunoassay (Abbott Japan).

Genotypes and Subgenotypes of HBV. The six major HBV genotypes (A-F) were determined serologically by EIA using commercial kits (HBV GENOTYPE EIA; Institute of Immunology). The method depends on the combination of epitopes on preS2-region products detected by monoclonal antibodies, which is specific for each of them.²⁰ HBV/G was determined by a slight modification of the polymerase chain reaction (PCR) with specific primers.²¹

Subgenotypes of HBV/A designated Ae prevalent in Europe and Aa frequent in Africa as well as Asia,⁹ which corresponds to subgroup A' originally reported by Bowyer et al.,²² were determined by PCR restriction fragment length polymorphism (RFLP) involving nucleotide conversions in an immediate upstream of the precore region that are specific for each of them.^{16,23} HBV/Bj (Japanese type) lacking the recombination with C over the precore region and the core gene and Ba (Asian type) with the recombination were determined by its absence or presence on HBV DNA sequences, as well as RFLP based on specific nucleotide substitutions, after the methods described previously.^{15,24}

Subgenotypes of HBV/C, Cs (Southeast Asian type) found only in Southeast Asia, including Vietnam, Myanmar, Thailand, Laos, Bangladesh, Hong Kong, and Southern China, and Ce (East Asian type), found in Far

East Asia, including Japan, Korea, and Northern China, were determined by the PCR-RFLP method described previously.¹²

Quantification of HBV DNA and Sequencing. HBV DNA sequences spanning the S gene were determined by real-time detection PCR according to the method of Abe et al.,²⁵ with the detection limit of 100 copies/mL. HBV DNA sequences bearing core promoter, precore region, and the core gene were amplified by PCR with hemi-nested primers by the method described previously.¹⁰ Negative samples were tested by another more sensitive second-round PCR with HB7F and HBV1917R (5'-CTC CAC AGT AGC TCC AAA TTC TTT A-3'). Thereafter, PCR products were directly sequenced with Prism Big Dye (Applied Biosystems, Foster City, CA) in the ABI 3100 DNA automated sequencer.

Construction of Plasmid and Site-Directed Mutagenesis of HBV DNA. Serum samples were obtained from two patients infected with HBV/Bj and a patient with Ce. HBV DNA was extracted from 100 μ L serum using QIAamp DNA blood kit (QIAGEN, GmbH, Hilden, Germany). Four primer sets were designed to amplify two fragments covering the entire HBV genome. Amplified fragments were inserted into pGEM-T Easy Vector (Promega, Madison, WI) and cloned in DH5a competent cells (TOYOBO, Osaka, Japan). At least five clones of each fragment were sequenced and the consensus sequence determined. Among them, those containing the consensus sequence were identified and adopted as templates for further construction. Finally, 1.24-fold the HBV genome (nt 1413-3215/1-2185), just enough to transcribe oversized pregenome and precore mRNA, was constructed into pUC19 vector (Invitrogen Corp., Carlsbad, CA). For site-directed mutagenesis, the wild-type HBV was digested by *HindIII* and *EcoO65I* and ligated with the fragment carrying T1762/A1764 to produce 1.24-fold the genome carrying the core-promoter double mutation. Similarly, 1.24-fold the HBV genome with the precore stop-codon mutation (1896A) was generated. Further details are available online at: <http://interscience.wiley.com/jpages/0270-9139/suppmat/index.html>.

Cell Culture and DNA Transfection. For the standard replication assay, 10-cm-diameter dishes were seeded with 1×10^6 Huh7 cells each. After 16 hours of culture, cells were transfected with 5 μ g DNA construct using the FuGENE 6 transfection reagent (Roche Diagnostics, Indianapolis, IN) and harvested 3 days later. Transfection efficiency was measured by cotransfection with 1 μ g reporter plasmid expressing secreted alkaline phosphatase and estimating its enzymatic activity in the culture supernatant.

Southern Blot Hybridization. HBV DNA samples

from cells at day 3 in culture were separated on 1.2% (wt/vol) agarose gel, transferred to a positive-charged nylon membrane (Roche Diagnostics), and hybridized with full-length HBV DNA labeled with alkaline phosphatase. Detection was performed with CDP-star (Amersham Biosciences, Piscataway, NJ), and signals were analyzed in the LAS-1000 image analyzer (Fuji Photo Film, Tokyo, Japan).

Statistical Analysis. Categorical variables were compared between groups by the chi-squared test and non-categorical variables by the Mann-Whitney *U*-test. A *P* value less than .05 was considered significant. Multivariate analyses with logistic regression were used to determine independent factors for fulminant hepatitis. STATA Software (StataCorp LP, College Station, TX) version 8.0 was employed for analyses.

Results

Demographic and Clinical Differences in Patients Infected With Various HBV Genotypes/Subgenotypes.

Genotypes of HBV were not classifiable in 28 (8%), and sufficient clinical data were not available in 7 (2%) of the 336 patients with acute hepatitis B. Exclusive of these 35 patients, 301 (90%) were left for evaluation of HBV genotypes in reference to clinical outcome.

HBV genotypes/subgenotypes were Aa in 10 (3%), Ae in 33 (11%), Ba in 22 (7%), Bj in 22 (7%), Cs in 11 (4%), Ce in 192 (64%), D in 5 (2%), and G in 6 (2%); none of them were infected with F or H (Table 1). All six patients with HBV/G were co-infected with another genotype; Ae in two, Ba in two, and Ce in the remaining two. The mean age was lower in the patients with HBV/Ae than Ba ($P = .0001$), Aa ($P < .01$), Bj or Cs ($P < .05$ for each) and Ce than Ba ($P < .05$). Men predominated in HBV infections with foreign (Ae and Ba) compared with domestic genotypes (Bj and Ce) ($P < .05$).

HBeAg was detected in 79% of patients with HBV/Ae at a frequency much higher than that with Bj ($P < .005$), Ce ($P < .001$) or Ba ($P < .05$). HBeAg in four of the six (67%) patients with HBV/G was coded for by HBV of the other genotypes co-infecting them, because it has two stop codons and an insertion in the core gene that prohibit encoding HBeAg.²¹ HBV DNA levels as well as HBeAg-positive rates at the presentation were higher in HBV/Ae than Ce ($P < .005$) or Bj ($P < .05$) infection.

The peak alanine aminotransferase (ALT) level was higher in HBV/Bj than Ae infection ($P < .05$). Fulminant hepatitis was significantly more frequent in patients infected with HBV/Bj (55%) than the other genotypes ($P < .05$); it occurred in two of the five (40%) patients with HBV/D, also. In reflection of severe clinical course,

49. Shiraki K, Yoshihara N, Sakurai M, Eto T, Kawana T. Acute hepatitis B in infants born to carrier mother with the antibody to hepatitis B e antigen. *J Pediatr* 1980;97:768-70.
50. Terazawa S, Kojima M, Yamanaka T, Yotsumoto S, Okamoto H, Tsuda F, et al. Hepatitis B virus mutants with precore-region defects in two babies with fulminant hepatitis and their mothers positive for antibody to hepatitis B e antigen. *Pediatr Res* 1991;29:5-9.
51. Chen HL, Chang CJ, Kong MS, Huang FC, Lee HC, Lin CC, et al. Pediatric fulminant hepatic failure in endemic areas of hepatitis B infection: 15 years after universal hepatitis B vaccination. *Hepatology* 2004;39:58-63.
52. Fagan EA, Smith PM, Davison F, Williams R. Fulminant hepatitis B in successive female sexual partners of two anti-HBe-positive males. *Lancet* 1986;2:538-40.
53. Carman WF, Fagan EA, Hadziyannis S, Karayiannis P, Tassopoulos NC, Williams R, Thomas HC. Association of a precore genomic variant of hepatitis B virus with fulminant hepatitis. *Hepatology* 1991;14:219-22.
54. Sato S, Suzuki K, Akahane Y, Akamatsu K, Akiyama K, Yunomura K, et al. Hepatitis B virus strains with mutations in the core promoter in patients with fulminant hepatitis. *Ann Intern Med* 1995;122:241-8.
55. Kosaka Y, Takase K, Kojima M, Shimizu M, Inoue K, Yoshida M, et al. Fulminant hepatitis B: induction by hepatitis B virus mutants defective in the precore region and incapable of encoding e antigen. *Gastroenterology* 1991;100:1087-94.
56. Omata M, Ehata T, Yokosuka O, Hosoda K, Ohto M. Mutations in the precore region of hepatitis B virus DNA in patients with fulminant and severe hepatitis. *N Engl J Med* 1991;324:1699-704.
57. Liang TJ, Hasegawa K, Rimon N, Wands JR, Ben-Porath E. A hepatitis B virus mutant associated with an epidemic of fulminant hepatitis. *N Engl J Med* 1991;324:1705-9.
58. Liang TJ, Hasegawa K, Munoz SJ, Shapiro CN, Yoffe B, McMahon BJ, et al. Hepatitis B virus precore mutation and fulminant hepatitis in the United States: a polymerase chain reaction-based assay for the detection of specific mutation. *J Clin Invest* 1994;93:550-5.
59. Chen MT, Billaud JN, Sallberg M, Guidotti LG, Chisari FV, Jones J, et al. A function of the hepatitis B virus precore protein is to regulate the immune response to the core antigen. *Proc Natl Acad Sci USA* 2004;101:14913-8.
60. Milich D, Liang TJ. Exploring the biological basis of hepatitis B e antigen in hepatitis B virus infection. *Hepatology* 2003;38:1075-86.
61. Fang ZL, Ling R, Wang SS, Nong J, Huang CS, Harrison TJ. HBV core promoter mutations prevail in patients with hepatocellular carcinoma from Guangxi, China. *J Med Virol* 1998;56:18-24.
62. Baptista M, Kramvis A, Kew MC. High prevalence of 1762(T) 1764(A) mutations in the basic core promoter of hepatitis B virus isolated from black Africans with hepatocellular carcinoma compared with asymptomatic carriers. *Hepatology* 1999;29:946-53.
63. Hsia CC, Yuwen H, Tabor E. Hot-spot mutations in hepatitis B virus X gene in hepatocellular carcinoma. *Lancet* 1996;348:625-6.
64. Kao JH, Chen PJ, Lai MY, Chen DS. Basal core promoter mutations of hepatitis B virus increase the risk of hepatocellular carcinoma. *Gastroenterology* 2003;124:327-34.
65. Kuang SY, Jackson PE, Wang JB, Lu PX, Munoz A, Qian GS, et al. Specific mutations of hepatitis B virus in plasma predict liver cancer development. *Proc Natl Acad Sci USA* 2004;101:3575-80.
66. Yeh CT, Shen CH, Tai DI, Chu CM, Liaw YF. Identification and characterization of a prevalent hepatitis B virus X protein mutant in Taiwanese patients with hepatocellular carcinoma. *Oncogene* 2000;19:5213-20.
67. Carman WF, Zanetti AR, Karayiannis P, Waters J, Manzillo G, Tanzi E, et al. Vaccine-induced escape mutant of hepatitis B virus. *Lancet* 1990;336:325-9.
68. Gerken G, Kremsdorf D, Capel F, Petit MA, Dauguet C, Manns MP, et al. Hepatitis B defective virus with rearrangements in the preS gene during chronic HBV infection. *Virology* 1991;183:555-65.
69. Minami M, Okanoue T, Nakajima E, Yasui K, Kagawa K, Kashima K. Significance of pre-S region-defective hepatitis B virus that emerged during exacerbation of chronic type B hepatitis. *Hepatology* 1993;17:558-63.
70. Takayanagi M, Kakumu S, Ishikawa T, Higashi Y, Yoshioka K, Wakita T. Comparison of envelope and precore/core variants of hepatitis B virus (HBV) during chronic HBV infection. *Virology* 1993;196:138-45.
71. Nakajima E, Minami M, Ochiya T, Kagawa K, Okanoue T. PreS1 deleted variants of hepatitis B virus in patients with chronic hepatitis. *J Hepatol* 1994;20:329-35.
72. Melegari M, Bruno S, Wands JR. Properties of hepatitis B virus pre-S1 deletion mutants. *Virology* 1994;199:292-300.
73. Melegari M, Scaglioni PP, Wands JR. The small envelope protein is required for secretion of a naturally occurring hepatitis B virus mutant with pre-S1 deleted. *J Virol* 1997;71:5449-54.
74. Wang HC, Chang WT, Chang WW, Wu HC, Huang W, Lei HY, et al. Hepatitis B virus pre-S2 mutant upregulates cyclin A expression and induces nodular proliferation of hepatocytes. *Hepatology* 2005;41:761-70.
75. Hass M, Hannoun C, Kalinina T, Sommer G, Manegold C, Gunther S. Functional analysis of hepatitis B virus reactivating in hepatitis B surface antigen-negative individuals. *Hepatology* 2005;42:93-103.
76. Beasley RP, Hwang LY, Lin CC, Chien CS. Hepatocellular carcinoma and hepatitis B virus: a prospective study of 22 707 men in Taiwan. *Lancet* 1981;2:1129-33.
77. Yu MW, Chang HC, Liaw YF, Lin SM, Lee SD, Liu CJ, et al. Familial risk of hepatocellular carcinoma among chronic hepatitis B carriers and their relatives. *J Natl Cancer Inst* 2000;92:1159-64.
78. Arbutnot P, Kew M. Hepatitis B virus and hepatocellular carcinoma. *Int J Exp Pathol* 2001;82:77-100.
79. Brechot C. Pathogenesis of hepatitis B virus-related hepatocellular carcinoma: old and new paradigms. *Gastroenterology* 2004;127:S56-61.
80. Boucharad MJ, Schneider RJ. The enigmatic X gene of hepatitis B virus. *J Virol* 2004;78:12725-34.
81. Ganem D, Schneider RJ. Hepadnaviridae: the viruses and their replication. In: Knipe DM, Howley PM, editors. *Fields' virology*. Vol 2, 4th edn. Philadelphia: Lippincott Williams & Wilkins; 2001. p. 2923-69.
82. Kim CM, Koike K, Saito I, Miyamura T, Jay G. HBx gene of hepatitis B virus induces liver cancer in transgenic mice. *Nature* 1991;351:317-20.
83. Cha MY, Kim CM, Park YM, Ryu WS. Hepatitis B virus X protein is essential for the activation of Wnt/beta-catenin signaling in hepatoma cells. *Hepatology* 2004;39:1683-93.
84. Sirna H, Giannini C, Poussin K, Paterlini P, Kremsdorf D, Brechot C. Hepatitis B virus X mutants, present in hepatocellular carcinoma tissue abrogate both the antiproliferative and transactivation effects of HBx. *Oncogene* 1999;18:4848-59.
85. Tu H, Bonura C, Giannini C, Mouly H, Soussan P, Kew M, et al. Biological impact of natural COOH-terminal deletions of hepatitis B virus X protein in hepatocellular carcinoma tissues. *Cancer Res* 2001;61:7803-10.
86. Chisari FV, Klopchin K, Moriyama T, Pasquinelli C, Dunsford HA, Sell S, et al. Molecular pathogenesis of hepatocellular carcinoma in hepatitis B virus transgenic mice. *Cell* 1989;59:1145-56.
87. Dejean A, Bougueleret L, Grzeschik KH, Tiollais P. Hepatitis B virus DNA integration in a sequence homologous to v-erb-A and steroid receptor genes in a hepatocellular carcinoma. *Nature* 1986;322:70-2.
88. Wang J, Chenivresse X, Henglein B, Brechot C. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 1990;343:555-7.

Table 1. Clinical Characteristics of Patients Acutely Infected With HBV of Distinct Genotypes/Subgenotypes

Features	Genotypes/Subgenotypes							
	Aa (n = 10)	Ae (n = 33)	Ba (n = 22)	Bj (n = 22)	Cs (n = 11)	Ce (n = 192)	D ^a (n = 5)	G ^{a,b} (n = 6)
Age (years)	42.2 ± 13.1	31.2 ± 10.3 ^d	41.5 ± 10.7 ^e	43.5 ± 19.1	38.5 ± 11.1	36.3 ± 15.0	38.6 ± 20.8	42.7 ± 17.5
Men	8 (80%)	30 (91%) ^f	19 (86%) ^g	9 (41%)	7 (64%)	122 (64%)	2 (40%)	6 (100%)
HBeAg positive	7 (70%)	26 (79%) ^h	11 (50%)	8 (36%)	8 (73%)	101 (53%)	1 (20%)	4 (67%)
ALT (IU/L)	1875 ± 759	2070 ± 1113 ⁱ	2523 ± 1185	3472 ± 2720	2269 ± 995	2610 ± 1719	2559 ± 1672	2142 ± 722
Duration of elevated ALT (weeks) ^f	7.9 ± 5.8	9.5 ± 6.2	8.8 ± 3.7 ⁱ	6.0 ± 2.5	10.1 ± 7.5	7.7 ± 5.1	5.7 ± 2.1	9.8 ± 1.5
Total bilirubin (mg/dL)	14.1 ± 10.3	9.0 ± 7.2	9.3 ± 5.9	10.9 ± 9.0	11.0 ± 13.8	9.8 ± 10.7	8.2 ± 2.2	13.0 ± 7.8
HBV DNA (log copies/mL)								
Median	4.76	6.08 ^k	5.15	4.93	5.61	4.94	5.91	5.97
(range)	(2.90-8.08)	(2.00-8.46)	(2.00-8.19)	(2.00-8.44)	(2.00-8.50)	(2.00-9.06)	(2.00-8.37)	(3.35-7.11)
<2.00 (undetectable)	0 (0%)	1 (3%)	2 (9%)	3 (14%)	2 (18%)	28 (15%)	1 (20%)	0 (0%)
Medication with								
Lamivudine	1 (10%)	9 (27%)	2 (9%)	5 (23%)	2 (18%)	28 (15%)	4 (80%)	2 (33%)
Steroid	0	3 (9%)	0	5 (23%)	1 (9%)	16 (8%)	0	0

^aPatients with HBV genotype D or G were not included in the analysis.

^bAll patients with HBV genotype G were co-infected with HBV of another genotype; Ae in two, Ba in two, and Ce in two.

^cExclusive of the 16 patients who died of fulminant hepatitis, 3 receiving liver transplantation and 10 without clinical data available.

^dP = .0001, Ae vs. Ba. P < .01, Ae vs. Aa. P < .05, Ae vs. Bj or Cs.

^eP < .05, Ba vs. Ce.

^fP = .0001, Ae vs. Bj. P < .005, Ae vs. Ce.

^gP < .005, Ba vs. Bj. P < .05, Ba vs. Ce.

^hP < .005, Ae vs. Bj. P < .01, Ae vs. Ce. P < .05, Ae vs. Ba.

ⁱP < .05, Ae vs. Bj.

^jP < .01, Ba vs. Bj. P < .05, Ba vs. Ce.

^kP < .005, Ae vs. Ce. P < .05, Ae vs. Bj.

the peak ALT level tended to be high in patients with HBV/Bj.

Presumed infection routes of 301 patients were sexual transmission in 172 (57%), blood transfusion in 4 (1%), medical accidents in 17 (6%), and unknown in the remaining 108 (36%).

Clinical Outcome of Patients With Acute Hepatitis B.

Fulminant hepatitis developed in 40 (13%) patients. To cope with severe acute liver disease, lamivudine and steroid were administered to 53 (18%) and 25 (8%) patients, respectively. Fulminant hepatitis led to death in 16 (5%) patients, and three (1%) received liver transplantation. Exclusive of the 40 patients with fulminant hepatitis who received various treatments and five without clinical data, 256 (85%) were followed for the chronic outcome (Fig. 1). Serum ALT levels stayed elevated for longer than 24 weeks for the diagnosis of chronic hepatitis in eight (3%) of them. Among them, five had cleared HBsAg from serum until then, and therefore, their liver function abnormality was not attributed to persistent HBV infection. Table 2 summarizes persistence of HBV infection in the 256 patients with acute hepatitis; 253 (99%) lost serum HBsAg by 6 months. Hence, HBV infection evolved into chronicity in only 3 of the 256 (1%) patients, representing 2 of the 32 (6%) infected with HBV/Ae and 1 of the 21 (5%) with Ba. All of the three with chronic outcome had low-titered IgG anti-HBc at the presentation, and

two of them had been negative for HBsAg before the presentation. None of them had received lamivudine or steroid treatment during their acute phase of illness. Of the patients without antiviral therapy, chronic outcome was significantly more frequent in those infected with HBV/Ae than non-Ae genotypes (9% ³/₂₃ vs. 0.5% ¹/₁₈₇, P = .032).




Fig. 1. A flow diagram of 336 patients studied. Comparison was made between patients with fulminant and acute self-limited hepatitis (upper dotted area), and the chronicity was compared between patients with and without treatments (lower dotted area). *Of 301 patients, 37 were negative for HBV DNA, including 27 with acute and 10 with fulminant hepatitis.

Table 2. Persistence of HBV Infection in the Patients With Acute Hepatitis Who Did or Did Not Receive Lamivudine or Steroid

Treatment	Total	Genotypes/Subgenotypes							
		Aa (n = 8) ^a	Ae (n = 32) ^a	Ba (n = 21) ^a	Bj (n = 10) ^a	Cs (n = 10) ^a	Ce (n = 167) ^a	D (n = 3) ^a	G (n = 5) ^a
Total (n = 256)	3/256 (1.2%)	0	2/32 (6%) ^c	1/21 (5%)	0	0	0	0	0
Lamivudine (n = 36) ^b	0/36 (0%)	0/1 (0%)	0/9 (0%)	0/2 (0%)	0	0/1 (0%)	0/19 (0%)	0/2 (0%)	0/2 (0%)
Steroid (n = 16) ^b	0/16 (0%)	0	0/3 (0%)	0	0	0/1 (0%)	0/12 (0%)	0	0
Neither	3/210 (1.4%)	0/7 (0%)	2/23 (9%) ^c	1/19 (5%)	0/10 (0%)	0/8 (0%)	0/139 (0%)	0/1 (0%)	0/3 (0%)

^aExclusive of 40 patients with fulminant hepatitis and 5 without clinical data available.

^bSix patients received steroid along with lamivudine.

^c $P < .05$, Ae vs. non-Ae.

Comparison Between Patients With Fulminant and Acute Self-Limited Hepatitis. Table 3 compares demographic, clinical, and virological characteristics between the 40 patients with fulminant and the 261 with acute self-limited hepatitis for whom analysis was feasible. Patients with fulminant hepatitis were significantly older (44.7 ± 16.3 vs. 36.0 ± 14.3 years, $P = .0017$), less predominantly male (43% vs. 71%, $P = .0005$) and less often positive for HBeAg (23% vs. 60%, $P < .0001$) than those with acute hepatitis. Peak ALT and total bilirubin levels were higher for fulminant than acute hepatitis ($P < .0001$), reflecting severe hepatic lesions. Notably, the median HBV DNA level was lower in patients with fulminant than acute hepatitis (4.89 vs. 5.19 log copies/mL, $P = .0178$); the frequency of unde-

etectable HBV DNA at the presentation was higher in fulminant hepatitis (25% vs. 10%, $P = .0086$). Lamivudine or steroid was given significantly more often to patients with fulminant hepatitis.

There were marked differences in the distribution of genotypes between patients with fulminant and acute hepatitis. HBV/Ae was less frequent (0% vs. 13%, $P = .0121$), whereas Bj was more often (30% vs. 4%, $P < .0001$) in patients with fulminant than acute hepatitis. Although HBV/Ce tended to be less frequent in patients with fulminant than acute hepatitis (55% vs. 65%), the difference fell short of being significant.

Precore stop-codon mutation (G1896A) and core-promoter double mutation (A1762T/G1764A) were more

Table 3. Comparison Between Patients With Fulminant and Acute Self-Limited Hepatitis Who Were Infected With HBV

Features	Fulminant (n = 40)	Acute (n = 261)	P Value
Age (years)	44.7 ± 16.3	36.0 ± 14.3	.0017
Men	17 (43%)	186 (71%)	.0005
HBeAg positive	9 (23%)	157 (60%)	<.0001
ALT (IU/L)	4315 ± 2889	2284 ± 1221	<.0001
Total bilirubin (mg/dL)	20.5 ± 16.4	8.3 ± 7.3	<.0001
HBV DNA (log copies/mL)			
Median	4.89	5.19	.0178
(range)	(2.00-8.44)	(2.00-9.06)	
<2.00 (undetectable)	10 (25%)	27 (10%)	.0086
Treatment			
Lamivudine	16 (40%)	37 (14%)	.0003
Steroid	9 (23%)	16 (6%)	.0022
Genotypes/subgenotypes			
Aa	1 (2.5%)	9 (3%)	NS
Ae	0 (0%)	33 (13%)	.0121
Ba	1 (2.5%)	21 (8%)	NS
Bj	12 (30%)	10 (4%)	<.0001
Cs	1 (2.5%)	10 (4%)	NS
Ce	22 (55%)	170 (65%)	NS
D	2 (5%)	3 (1%)	NS
G	1 (2.5%)	5 (2%)	NS
Mutations ^a			
nt 1753 and/or nt1754 ^b	11/30 (37%)	28/234 (12%)	.0003
A1762T/G1764A	15/30 (50%)	39/234 (17%)	<.0001
G1896A	16/30 (53%)	21/234 (9%)	<.0001
G1899A	7/30 (23%)	8/234 (3%)	<.0001

^aExclusive of 37 patients in whom precore region and core-promoter could not be amplified by PCR.

^bT1753C/A/G and/or T1754C/A/G.

Table 4. Multivariate Analysis for Factors Independently Associated With Fulminant Hepatitis

Factors	Odds Ratio	95% Confidence Interval	P Value
Age (yr)			
<34 ^a	1		
≥34	3.472	1.094-11.023	.0347
Sex			
Male	1		
Female	2.272	0.780-6.613	.1323
HBeAg			
Positive	1		
Negative	3.344	1.065-10.506	.0387
ALT (IU/L)			
<2200 ^a	1		
≥2200	2.094	0.683-6.414	.1957
Total bilirubin (mg/dL)			
<10.0 ^a	1		
≥10.0	18.818	4.320-81.980	<.0001
HBVDNA (log copies/mL)			
<5.00 ^a	1		
≥5.00	1.042	0.367-2.961	.9383
Treatment			
Lamivudine (-)	1		
Lamivudine (+)	2.650	0.814-8.625	.1056
Steroid (-)	1		
Steroid (+)	2.515	0.668-9.472	.1728
Genotypes/Subgenotypes			
Non-Bj	1		
Bj	7.001	1.737-28.228	.0062
Mutations			
nt 1753 and/or 1754 ^b			
Absent	1		
Present	2.316	0.698-7.683	.1700
A1762T/G1764A			
Absent	1		
Present	1.013	0.295-3.478	.9841
G1896A			
Absent	1		
Present	4.157	1.265-13.657	.0189
G1899A			
Absent	1		
Present	2.525	0.534-11.949	.2427

^aMedian values.

^bT1753C/A/G or T1754C/A/G.

frequent in patients with fulminant than acute hepatitis (53% vs. 9% and 50% vs. 17%, respectively, $P < .0001$ for each). Likewise, mutations in core-promoter at nt 1753 or nt 1754, and G1899A mutation were more frequent in patients with fulminant than acute hepatitis ($P = .0003$ and $P < .0001$, respectively).

Factors Independently Associated With the Development of Fulminant Hepatitis. Various factors found in association with fulminant hepatitis were evaluated for the independence in multivariate analysis (Table 4). Age 34 years or older (odds ratio 3.47 [95% confidence interval 1.09-11.02], $P = .035$), HBV/Bj (7.00 [1.74-28.23], $P = .006$), HBeAg-negative (3.34 [1.07-10.51], $P = .039$), total bilirubin ≥ 10.0 mg/dL (18.82 [4.32-81.98], $P < .0001$) and G1896A (4.16 [1.27-13.66], $P = .019$)




Fig. 2. Frequencies of precore and core-promoter mutations compared between patients with fulminant and acute self-limited hepatitis who were infected with HBV/Bj (A) or Ce (B).

were independent risk factors for the development of fulminant hepatitis.

In view of the majority of Japanese patients who were infected with Bj or Ce, mutations in the precore region and core-promoter were compared between those with fulminant and acute self-limited hepatitis for each subgenotype (Fig. 2). G1896A and A1762T/G1764A were significantly more frequent in patients with fulminant than acute hepatitis infected with either HBV/Bj or Ce (56% vs. 0% and 67% vs. 0% for Bj or 44% vs. 11% and 56% vs. 22% for Ce, respectively, $P \leq .01$ for all). For the patients infected with HBV/Bj, in particular, precore and core-promoter mutations were highly frequent in those with fulminant hepatitis (56% and 67%, respectively), whereas they occurred in none of those with acute hepatitis. G1899A was equally frequent in both patients with fulminant and acute hepatitis infected with HBV/Bj; it was rarely seen in those with Ce. Mutations involving nt 1753 or nt 1754 tended to be more frequent in patients with fulminant than acute hepatitis.

Replication of the Wild-Type HBV as Well as Precore and Core-Promoter Mutants In Vitro. Full-length HBV DNA of the wild-type HBV/Bj from a patient with chronic hepatitis B was incorporated with G1896A or A1762T/G1764A mutation *in vitro*. Another plasmid was constructed with HBV/Bj_58 carrying G1896A from a fulminant patient. Figure 3 compares