

## Risk Factors and Impact of $\beta$ -D Glucan on Invasive Fungal Infection for the Living Donor Liver Transplant Recipients

NAOKI KAWAGISHI, KAZUSHIGE SATOH, YOSHITAKA ENOMOTO, YORIHICO AKAMATSU, SATOSHI SEKIGUCHI, KEISEI FUJIMORI and SUSUMU SATOMI

*Division of Advanced Surgical Science and Technology, Graduate School of Medicine, Tohoku University, Sendai, Japan*

KAWAGISHI, N., SATOH, K., ENOMOTO, Y., AKAMATSU, Y., SEKIGUCHI, S., FUJIMORI, K. and SATOMI, S. *Risk Factors and Impact of  $\beta$ -D Glucan on Invasive Fungal Infection for the Living Donor Liver Transplant Recipients.* Tohoku J. Exp. Med., 2006, 209 (3), 207-215 — Invasive fungal infection is a fatal complication in liver transplantation and it is very difficult to diagnose at the early stage. The aim of this study was to review our experience with invasive fungal infections in living donor liver transplantation (LDLT) and to analyze the risk factors and the impact of  $\beta$ -D glucan. From 1991 to 2005, 96 LDLTs were performed in our institution and we measured the serum level of  $\beta$ -D glucan in order to clarify the diagnosis. Invasive fungal infection was diagnosed based on clinical symptoms, culture, radiological evidence and  $\beta$ -D glucan. Active fungal infection was treated with fluconazole, amphotericin B, flucytosine and micafungin. Risk factors both pre- and post- LDLT were analyzed. *Candida albicans* was the most frequently isolated species (70%). The risk factors identified by univariate analysis include the following four conditions: acute blood purification (plasma exchange with or without continuous hemodiafiltration), hepatic vein complications, renal failure and respiratory failure. By logistic regression analysis, hepatic vein complications and respiratory failure were identified as independent risk factors. The risk factors for invasive fungal infection of LDLT in Japan have not been well analyzed and this report will provide valuable information for the prevention of the fungal infection. ——— invasive fungal infection; living donor liver transplantation;  $\beta$ -D glucan; risk factor; complication

© 2006 Tohoku University Medical Press

Fungal infection is a potentially fatal complication in patients subjected to liver transplantation. The incidence of fungal infection in liver transplantation is higher than in transplantation of other solid organs, ranging from 4 to 42% with a mortality of 25-69% (Patel et al. 1996; Singh et al. 1997, 2002; Rabkin et al. 2000; Paya 2001).

Many reports have been published about risk factors for invasive fungal infection in recipients of deceased donor liver grafts (Gladdy et al. 1999; Fortun et al. 2002); on the other hand, there have been no reports about the risk factors for living donor liver transplantation (LDLT). Most LDLTs are elective surgery, thus it is possible to treat pre-

---

Received March 13, 2006; revision accepted for publication April 13, 2006.

Correspondence : Naoki Kawagishi, M.D., Division of Advanced Surgical Science and Technology, Graduate School of Medicine, Tohoku University, 1-1 Seiryomachi, Aoba-ku, Sendai 980-8574, Japan.  
e-mail: kawan@mail.tains.tohoku.ac.jp

existing infections and prepare the patient by administering prophylactic agents. However, in LDLT the graft is smaller than that recovered from a deceased donor, and it involves technical difficulties such as the small diameter of vessels and biliary ducts. Therefore, it needs to have special consideration for the invasive fungal infection in LDLT (Suzuki et al. 2000). So it is important to know the risk factors in LDLT for the early management.

In the clinical field, it is very difficult to diagnose an invasive fungal infection in the early stage, because the pathogens cannot be detected until the infection has become quite severe (Aikawa et al. 2002). Besides, there are no significantly superior chemical makers for the diagnosis of invasive fungal infection. Therefore, there are no global standardized criteria in this regard (Avery 2001). In Japan serum  $\beta$ -D glucan is used for the diagnosis of fungal infection. Obayashi et al. (1995) reported that serum  $\beta$ -D glucan was a highly sensitive and specific test for invasive deep mycosis and fungal febrile episodes, and would substantially benefit immunocompromised patients. In the present study we investigated the specificity and sensitivity of  $\beta$ -D glucan for the detection of invasive fungal infection in LDLT recipients. This test, however, provides no information on cryptococcosis, because *C neoformans* produces less  $\beta$ -D glucan than do *candida* and *asperillus* (Obayashi et al. 1995). But we experienced its usefulness in case of *Pneumocystis carinii* pneumonia whose clinical symptoms are closely correlated with the level of  $\beta$ -D glucan.

In this study we review our experience with fungal infections in LDLT recipients and report the risk factors for invasive fungal infection. We also discuss the usefulness of the  $\beta$ -D glucan in the diagnosis and therapeutical evaluation of invasive fungal infection in LDLT recipients.

#### PATIENTS AND METHODS

##### *Characteristics of the patients*

From July 1991 to November 2005, 96 living related liver transplantations were performed in pediatric and adult patients with end-stage liver disease in the Division

of Advanced Surgical Science and Technology, Graduate School of Medicine, Tohoku University. This program was approved by the ethics committees in our hospital and the patients were given informed consent about the LDLT. Donors were selected among parents, brothers or sisters and spouses on the basis of liver function tests, serological markers of hepatitis, ABO blood group, graft/recipient size matching, lymphocyte cross-matching and human leukocyte (lymphocyte) antigen typing. All the recipients were transplanted the right lobe, left lobe or the left lateral segment of the liver. The primary diseases of the recipients were biliary atresia in 54, primary biliary cirrhosis (PBC) in 12, hepatocellular carcinoma in 6, cryptogenic liver cirrhosis in 4, metabolic disorders in 4, and other diseases in 16. These 96 recipients were 57 females and 39 males. The mean age was 18 years 8 month, ranging from 5 months to 63 years, and the mean weight was 31.2 kg, ranging from 5.3 kg to 89.0 kg at the time of transplantation.

##### *Anti-infection therapy and immunosuppressive therapy*

Several specimens, including pharyngeal swabs, nasal swabs, urine and feces were cultured before LDLT and if methicillin-resistant staphylococcus aureus (MRSA) or fungus were isolated, they were treated by vancomycin, mupirocin calcium and fluconazole. Routine preoperative bowel preparation was performed orally by fluconazole (3 mg/kg/day) and kanamycin (50 mg/kg/day) for 3 days before surgery and gargle by amphotericin B. For the children above 2 years old who had been performed splenectomy or had a plan of splenectomy at the time of transplantation, they had vaccination for *Streptococcus pneumoniae* and *Hemophilus influenzae* before transplantation. Fluconazole (5 mg/kg/day) or micafungin (50 mg/day) were given intravenously for fungal prophylaxis for the first 1 week after operation. From the second week after transplantation fluconazole was given orally at the dose of 3 mg/kg/day for another one or two weeks. Trimethoprim-sulfamethoxazole was given routinely for *Pneumocystis carinii*. Febrile episode were treated empirically with broad-spectrum antibiotic until culture results were known. Thereafter, bacterial infections were treated with antibiotics based on specific sensitivity patterns. Recipients with suspected fungal infection underwent a chest or abdominal computed tomography (CT) scan, when appropriate, and fundoscopy for *Candida* endophthalmitis, and suspected sites of infection were cultured. Fungal infections were identified in some ways,

they were treated with an initial dose of fluconazole (3-6 mg/kg/day), amphotericin B (0.25 mg/kg/day), micafungin (100-150 mg/day) or flucytosine (100 mg/day). Routine surveillance cultures after transplantation were obtained from pharyngeal swabs, nasal swabs, urine, central venous lines, abdominal drains, nasogastric tubes, endotracheal tubes and stool. Venous and urinary catheters were routinely changed when colonization of an indwelling tube was suspected. Serum level of *Candida*, *Aspergillus* and *Cryptococcus* antigens were also measured for the diagnosis. The culture medium used was fungus specific. *Candida* antigen was detected by latex agglutination and the positive value was more than two times higher compared to control. *Aspergillus* antigen was detected by latex agglutination and this is qualitative analysis. *Cryptococcus* antigen was detected by latex agglutination and the positive value was more than eight times higher compared to control.  $\beta$ -D glucan was determined by test kits (Wako, Inc., Tokyo), and the cut-off value was 11 pg/ml. The timing of sampling for these parameters was twice before transplantation and twice per week for one or two months after transplantation. The value was the maximal one.

The immunosuppressive regimen consisted of cyclosporine (CsA), tacrolimus (TAC), azathioprine (AZ), mycophenolate mofetil (MMF) and methylprednisolone (MP).

#### Definitions of invasive fungal infection

Fungal colonization was defined as the presence of a fungus in one or more surveillance superficial cultures in the absence of any clinical symptoms or signs of infection. Invasive fungal infections were divided into definite and probable infections according to the following criteria: definite invasive fungal infection was diagnosed by one of these criteria as follows: 1) presence of fungus in the blood, cerebrospinal fluid, ascites and biopsy specimens; 2) ophthalmologic evidence; 3) isolation of fungus from bronchoalveolar lavage with radiological evidence of pneumonitis. Probable invasive fungal infection was defined as isolation of a fungus from one superficial site and more than 11 pg/ml serum level of  $\beta$ -D glucan or one of these two criteria together with febrile symptom above 38°C which was not responded by antibiotics for more than one week or radiological evidence of pneumonitis or esophagitis.

#### Clinical data analysis

Clinical data were analyzed to determine the risk

factors that might contribute to the fungal infections. Preoperative factors such as gender, age at transplantation, body weight, serum total bilirubin and creatinine levels, number of pre-transplant abdominal surgery, pre-transplant plasmapheresis and pre-transplant fungal colonization were compiled. Intraoperative variables included duration of operation, blood loss, graft vs recipient ratio and the type of biliary reconstruction. Postoperative variables were collected until present time. This included the number of laparotomies and plasma exchange, total number of days in intensive care unit (ICU), total number of days intubated, vascular and biliary complications, number of viral and methicillin-resistant staphylococcus aureus (MRSA) infections, episodes of biopsy proven rejections, graft-recipient weight ratio. Respiratory failure was defined as requiring intubation for more than 2 weeks or re-intubation, renal failure was defined as continuous hemodiafiltration (CHDF), continuous hemofiltration (CHF) or hemodialysis (HD) was initiated.

Statistical analyses were performed using Stat View 4.5 (Abacus Concepts, Inc., Berkeley, CA, USA). Univariate statistical analysis for invasive fungal infection vs control group was performed by means of the Fisher's exact test for categorical variables and Mann-Whitney's U-test for continuous variables. *P* value less than 0.05 was considered statistically significant. The factors identified on univariate analysis to be associated with *p* value less than 0.20 were then entered into a stepwise logistic regression analysis to identify independent risk factors for invasive fungal infection.

## RESULTS

During the study period, mycoses were detected in 48 patients (50.0%) before LDLT, in 63 (65.6%) after LDLT, and in 38 (39.6%) patients during both periods (Table 1). Before LDLT *C. albicans* was isolated from 72.9% of the recipients followed by *C. glabrata* (18.8%), and others (8.3%). After LDLT, *C. albicans* was isolated from 71.4% of the recipients followed by *C. glabrata* (31.7%), yeast like fungus (15.9%), *Candida spp* (12.7%), *C. guilliermondii* (7.9%) and *C. tropicalis* (7.9%). The intestines and pharynx were the most frequent colonized sites followed by urine and stomach. Forty-eight recipients who had fungal colonization before LDLT had no symptoms of infection. Sera from 8

TABLE 1. Isolates by fungal species, site of colonization.

|                              | Pre-Tx<br>(n = 48) | Post-Tx<br>(n = 63) |
|------------------------------|--------------------|---------------------|
| <i>Species</i>               |                    |                     |
| <i>C. albicans</i>           | 72.9% (35/48)      | 71.4% (45/63)       |
| <i>C. glabrata</i>           | 18.8% (9/48)       | 31.7% (20/63)       |
| Yeast like fungus            | 4.1% (2/48)        | 15.9% (10/63)       |
| <i>Candida spp.</i>          | 10.4% (5/48)       | 12.7% (8/63)        |
| <i>C. guilliermondii</i>     | 0                  | 7.9% (5/63)         |
| <i>C. tropicalis</i>         | 0                  | 7.9% (5/63)         |
| <i>C. parapsilosis</i>       | 6.3% (3/48)        | 4.8% (3/63)         |
| <i>C. krusei</i>             | 4.1% (2/48)        | 4.8% (3/63)         |
| <i>Trichosporon beigelli</i> | 0                  | 3.2% (2/63)         |
| <i>Aspergillus spp.</i>      | 2.1% (1/48)        | 1.6% (1/63)         |
| <i>Cryptococcus luteolus</i> | 2.1% (1/48)        | 0                   |
| Others                       | 8.3% (4/48)        | 1.6% (1/63)         |
| <i>Site</i>                  |                    |                     |
| Stool                        | 64.6% (31/48)      | 71.4% (45/63)       |
| Pharynx                      | 70.8% (34/48)      | 63.5% (40/63)       |
| Urine                        | 18.8% (9/48)       | 28.6% (18/63)       |
| Gastric fluid                | 0                  | 14.3% (9/63)        |
| Abdominal drain              | 0                  | 12.7% (8/63)        |
| Bile juice                   | 0                  | 9.5% (6/63)         |
| Sputum                       | 8.3% (4/48)        | 7.9% (5/63)         |
| Nasal cavity                 | 4.1% (2/48)        | 7.9% (5/63)         |
| Jejunum                      | 0                  | 4.8% (3/63)         |
| Blood                        | 0                  | 1.6% (1/63)         |

patients were positive for *Candida* antigens before LDLT and from 18 after LDLT. *Aspergillus* antigen was not detected in serum from any patient in our institution and *Aspergillus spp.* was isolated from two recipients but it was not responsible for fungal infection. There was no patient with pneumonia or meningitis caused by *Aspergillus* species. *Cryptococcus luteolus* was isolated once before LDLT and serum *Cryptococcus* antigen was detected in one recipient after LDLT. None of the patients had severe infection caused by *Cryptococcus* species.

We experienced 2 definite and 6 probable cases of invasive fungal infections (Table 2). Most of the invasive fungal infections occurred

within 2 months after LDLT. Seven recipients were diagnosed as having pneumonia by chest x-ray or CT scan and one recipient was diagnosed sepsis based on the results of an arterial blood culture. *C. albicans* was isolated from sputa of 4 out of eight recipients with a body temperature of 38°C or more and radiological evidence of pneumonia. Five out of eight recipients with invasive fungal infections died because of graft failure or multiple organ failure. No fungus was isolated from two of 6 patients with probable fungal infections who had high serum levels of  $\beta$ -D glucan (86, 38 pg/ml), a body temperature of 38°C or more and radiological evidence of pneumonia. Two recipients who showed an improvement of

TABLE 2. Clinical spectrum of definite and probable invasive fungal infection ( $n = 8$ ).

|                                       | Patient No. | Diagnosis | Species                                  | $\beta$ -D glucan*<br>(pg/ml) | Outcome |
|---------------------------------------|-------------|-----------|--|-------------------------------|---------|
| Definite invasive<br>fungal infection | 68          | Pneumonia | <i>P. carinii</i>                        | 169                           | Alive   |
|                                       | 87          | Sepsis    | <i>C. albicans</i>                       | 29                            | Dead    |
|                                       |             | Pneumonia |  |                               |         |
| Probable invasive<br>fungal infection | 12          | Pneumonia | <i>C. albicans</i><br><i>C. glabrata</i> | NA                            | Dead    |
|                                       | 22          | Pneumonia | <i>Candida spp</i><br><i>C. glabrata</i> | 57                            | Dead    |
|                                       | 37          | Pneumonia | -  | 86                            | Dead    |
|                                       | 39          | Pneumonia | -  | 38                            | Alive   |
|                                       | 40          | Pneumonia | <i>C. albicans</i>                       | 39                            | Alive   |
|                                       | 48          | Pneumonia | <i>C. albicans</i><br><i>C. glabrata</i> | 122                           | Dead    |

\* Peak value. NA, not available.

their fungal pneumonia were treated with fluconazole and antibiotics. One recipient who was diagnosed as having *Pneumocystis carinii* pneumonia had a quite high serum level of  $\beta$ -D glucan (169 pg/ml) and his clinical symptoms correlated with the level of  $\beta$ -D glucan. He was found to be positive for *Pneumocystis carinii* in his sputa from bronchoalveolar lavage (BAL) by polymerase chain reaction (PCR) and x-ray findings revealed pneumonia; thus, he was treated with trimethoprim-sulfamethoxazole.

Twenty-six factors, including 7 pretransplantation, 4 intraoperative, and 15 posttransplantation variables, were examined by univariate analysis to determine the risk factors that predispose to invasive fungal infection (Table 3). We did not show  $\beta$ -D glucan as a factor, because it was not measured in the first 21 patients. But there was a significant difference between recipients with invasive fungal infections and those with non-invasive fungal infections ( $p = 0.001$ ). The statistically significant factors were the number of post-transplant episodes of acute blood purification

(plasma exchange with or without CHDF) ( $p = 0.031$ ), hepatic vein complications ( $p = 0.014$ ), renal failure ( $p = 0.004$ ) and respiratory failure ( $p = 0.005$ ), etc. Multivariate analysis showed an independent association of the development of invasive fungal infection with hepatic vein complications (OR, 26.1 [95% CI, 1.23-553.19]) and respiratory failure (OR, 52.3 [95% CI, 1.51-1,811.85]) (Table 4). Among the patients of invasive fungal infections patient no. 37 and 40 had hepatic vein complication, and no. 12, 22, 39, 40, 48, 87 had respiratory failure. Overall mortality was significantly higher in the group of recipients with invasive fungal infections than in the group of those without invasive fungal infections (62.5% [5 of 8] vs 15.9% [14 of 88];  $p = 0.0015$ ).

## DISCUSSION

Invasive fungal infection is one of the fatal complications and influences the survival of LDLT recipients. In this report we investigated the risk factors for invasive fungal infection and the usefulness of  $\beta$ -D glucan as an indicator of

TABLE 3. Univariate analysis of risk factors.

|   | Invasive fungal infection (n = 8) | No invasive fungal infection (n = 88) | <i>p</i> |
|---|-----------------------------------|---------------------------------------|----------|
| Pretransplantation variables                      |                                   |                                       |          |
| No. of male recipients                            | 2                                 | 37                                    | 0.466    |
| Mean age (yr)                                     | 28.3                              | 17.8                                  | 0.106    |
| Mean body weight (kg)                             | 46.1                              | 29.9                                  | 0.047    |
| Mean total bilirubin (mg/dl)                      | 15.8                              | 10.2                                  | 0.070    |
| Mean serum creatinine (mg/dl)                     | 0.6                               | 0.4                                   | 0.064    |
| Mean no. of acute blood purification              | 0.8                               | 0.6                                   | 0.404    |
| Mean no. of laparotomies                          | 0.5                               | 1.2                                   | 0.074    |
| No. of patients with pre-LDLT fungal colonization | 5                                 | 43                                    | 0.715    |
| Intraoperative variables                          |                                   |                                       |          |
| Mean operation time (min)                         | 882                               | 902                                   | 0.551    |
| Mean blood loss (ml)                              | 9,350                             | 9,705                                 | 0.223    |
| Mean graft/recipient ratio                        | 1.153                             | 1.906                                 | 0.015    |
| Biliary reconstruction (duct-to-duct)             | 0                                 | 13                                    | 0.593    |
| Posttransplantation variables                     |                                   |                                       |          |
| Mean no. of rejection                             | 1.3                               | 0.7                                   | 0.573    |
| Mean no. of laparotomies                          | 1.1                               | 0.8                                   | 0.446    |
| Mean no. of acute blood purification              | 16.8                              | 3.3                                   | 0.031    |
| Mean no. of days in ICU                           | 23.1                              | 17.5                                  | 0.989    |
| Mean no. of days intubated                        | 28.1                              | 14.2                                  | 0.011    |
| Biliary complication                              | 2                                 | 19                                    | 0.999    |
| Hepatic artery thrombosis                         | 1                                 | 7                                     | 0.515    |
| Portal vein complication                          | 0                                 | 6                                     | 0.999    |
| Hepatic vein complication                         | 2                                 | 5                                     | 0.014    |
| CMV infection                                     | 1                                 | 13                                    | 0.999    |
| EBV infection                                     | 0                                 | 10                                    | 0.595    |
| MRSA infection                                    | 3                                 | 21                                    | 0.408    |
| Renal failure                                     | 5                                 | 12                                    | 0.004    |
| Respiratory failure                               | 6                                 | 20                                    | 0.005    |
| Mean no. of days in hospital                      | 131.5                             | 78.6                                  | 0.448    |

TABLE 4. Multivariate analysis of risk factors for invasive fungal infection.

| Variable                   | Odds ratio (95% CI)  | <i>p</i> |
|----------------------------|----------------------|----------|
| Hepatic vein complications | 26.1 (1.23-553.19)   | 0.036    |
| Respiratory failure*       | 52.3 (1.51-1,811.85) | 0.029    |

\*More than 2 weeks intubation.  
CI, confidence interval.

this type of infection in LDLT recipients.

In our institution *Candida* spp. was the species most frequently isolated pre- and post-transplantation, and among them *C. albicans* accounted for over 70% of the isolates. This result derived from the high level of gastrointestinal colonization with *Candida* spp., manipulation of the gastrointestinal tract during surgery with increased translocation across the gut mucosal barrier, and impaired hepatic reticuloendothelial function (Matsumoto et al. 2004), as well as prolonged use of an indwelling catheter which promotes fungal colonization by providing a vascular portal of entry.

In deceased donor liver transplantation (DDLT) *Candida* spp. constitute the most frequent cause of invasive fungal infection followed by *Aspergillus* spp., and the same tendency is observed in LDLT. Among *Candida* spp., *C. albicans* was the most predominant followed by *C. glabrata*, *C. tropicalis*, *C. parapsilosis*, *C. krusei* (Gladdy et al. 1999; Kawagishi et al. 2004; Takakura et al. 2004). *Candida* overgrowth in the lumen of the gastrointestinal tract is secondary to changes in the bacterial flora. It has been clearly established that the presence of anaerobic bacteria in the gut conveys a protective effect as they neutralize the overgrowth of *Candida* (van der Waaij 1982). This has been the basis for the use of selective bowel decontamination regimens, the main goal of which is to maintain a healthy anaerobic flora (Arnouk et al. 1996; van der Waaij et al. 1982; Wiesner et al. 1988). Some selective bowel decontamination regimens have included oral amphotericin preparations to further enhance the elimination of *Candida* from the bowel. The clinical effectiveness of this technique in reducing systemic *Candida* infection, however, remains unknown. In our experience drug resistant *C. glabrata* might be increased in the invasive fungal infection through the intensive use of prophylactic agents with our protocol.

In our series there were 8 recipients with invasive fungal infections, including 6 probable cases, and of them 5 died. Other authors also reported that the development of invasive fungal infection after liver transplantation was associated

with a high mortality rate of up to 80% due to the difficulty in making an early diagnosis of the disease (Nieto-Rodriguez et al. 1996; Gladdy et al. 1999; Fortun et al. 2002). Particularly, invasive aspergillosis was quite fatal (Fortun et al. 2002). Diagnosis of invasive fungal infection by conventional superficial cultures and fungal isolation from tissues, bronchoalveolar lavage or cerebrospinal fluid from patients with clinical symptoms has some limitations. But recently, diagnostic testing in the field of medical mycology has improved allowing an earlier diagnosis of infection and the possibility to monitor the response to therapy in a noninvasive manner. An assay that measures 1-3  $\beta$ -D glucan, a panfungal cell wall component, as a potentially promising indicator of invasive fungal infection has been developed (Obayashi et al. 1995, 2004; Ostrosky-Zeichner et al. 2005; Pazos et al. 2005). Although some false positives may result from hemodialysis using certain cellulose membranes, gauzes and albumin products (Nakao et al. 1997; Sakai et al. 2000; Kato et al. 2001), we think that this method is useful for the early diagnosis of invasive fungal infection. In our experience the specificity of  $\beta$ -D glucan was 100%. Besides, it was very useful to monitor the results of therapy for *Pneumocystis carinii* pneumonia. However it should also be recognized that  $\beta$ -D glucan is not expected to detect cryptococcosis due to the low content of 1-3- $\beta$ -D-glucan in the cell wall of *Cryptococcus* species (Miyazaki et al. 1995).

The risk factors for invasive fungal infection in liver transplantation have been reported by many authors (Gladdy et al. 1999; Avery et al. 2001; Paya 2001; Fortun et al. 2002; Singh et al. 2002; Takakura et al. 2004). But these were identified mainly from their experiences with DDLT. Therefore, we think that risk factors in LDLT recipients should be discussed. In Japan, we have experienced more than 3,000 cases of LDLT so far (Japan Liver Transplantation Society, personal communication), fungal infection is one of the fatal complications (Suzuki et al. 2000). In DDLT, bile leakage, hepatic artery thrombosis, preoperative steroids, blood transfusion during the operation, respiratory failure, hyperglycemia,

retransplantation, and renal failure were reported as independent risk factors (Nieto-Rodriguez et al. 1996; Gladdy et al. 1999; Fortun et al. 2002). Some other series identified thrombocytopenia, bacterial and viral infections and pre-transplant fungal colonization as risk factors (Avery et al. 2001; Verma et al. 2005). In our series, there were no pre- nor intra-operative variables identified as risk factors (except body weight and graft/recipient ratio) by univariate analysis. This may derive from the fact that LDLT is usually set up under the condition of elective surgery and prophylaxis for fungal infection is provided before transplantation, contrary to what happens in case of DDLT. Moreover, donor-transmitted infection does not happen in LDLT. Hepatic vein complications and respiratory failure were identified as risk factors by multivariate analysis in our series, these results were comparable to the reported risk factors in DDLT. But we consider that hepatic vein complication is a characteristic risk factor for LDLT, particularly when the left lobe is used as the graft. Because in such cases once hepatic vein complications occur, liver damage becomes more serious unless proper therapy is provided. This kind of serious liver damage is an important risk for the development of invasive fungal infection.

In conclusion, invasive fungal infection is an important cause of death after LDLT. We identified two factors associated with the development of invasive fungal infection: hepatic vein complications and respiratory failure. For the prevention and treatment of invasive fungal infection in LDLT, it is important to notify these risk factors, and we need more studies on early non-invasive quantitative markers like  $\beta$ -D glucan.

## References

- Aikawa, N., Sumiyama, Y., Kusachi, S., Hirasawa, H., Oda, S. & Yamazaki, Y. (2002) Use of antifungal agents in febrile patients nonresponsive to antibacterial treatment: the current status in surgical and critical care patients in Japan. *J. Infect. Chemother.*, **8**, 237-241.
- Arnou, P.M., Carandang, G.C., Zabner, R. & Irwin, M.E. (1996) Randomized controlled trial of selective bowel decontamination for prevention of infections following liver transplantation. *Clin. Infect. Dis.*, **22**, 997-1003.
- Avery, R.K. (2001) Fungal infection in the organ transplant recipient. *Curr. Opin. Organ. Transplant.*, **6**, 284-289.
- Fortun, J., Martin-Davila, P., Moreno, S., De Vicente, E., Nuno, J., Candelas, A., Barcena, R. & Garcia, M. (2002) Risk factors for invasive aspergillosis in liver transplant recipients. *Liver Transpl.*, **8**, 1065-1070.
- Gladdy, R.A., Richardson, S.E., Davies, H.D. & Sperina, R.A. (1999) Candida infection in pediatric liver transplant recipients. *Liver Transpl. Surg.*, **5**, 16-24.
- Kato, A., Takita, T., Furuhashi, M., Takahashi, T., Maruyama, Y. & Hishida, A. (2001) Elevation of blood (1 $\rightarrow$ 3)-beta-D-glucan concentrations in hemodialysis patients. *Nephron*, **89**, 15-19.
- Kawagishi, N., Fujimori, K. & Satomi, S. (2004) Diagnosis and management of postoperative fungal infection in recipients of living donor liver transplantation. *J. Abdom. Emerg. Med.*, **24**, 57-65.
- Matsumoto, C.S., Fishbein, T.M. & Kaufman, S.S. (2004) Gastrointestinal infection in solid organ transplant recipients. *Curr. Opin. Organ. Transplant.*, **9**, 406-410.
- Miyazaki, T., Kohno, S., Mitsutake, K., Maesaki, S., Tanaka, K., Ishikawa, N. & Hara, K. (1995) Plasma (1 $\rightarrow$ 3)-beta-D-glucan and fungal antigenemia in patients with candidemia, aspergillosis, and cryptococcosis. *J. Clin. Microbiol.*, **33**, 3115-3118.
- Nakao, A., Yasui, M., Kawagoe, T., Tamura, H., Tanaka, S. & Takagi, H. (1997) False-positive endotoxemia derives from gauze glucan after hepatectomy for hepatocellular carcinoma with cirrhosis. *Hepato-gastroenterology*, **44**, 1413-1418.
- Nieto-Rodriguez, J.A., Kusne, S., Manez, R., Irish, W., Linden, P., Magnone, M., Wing, E.J., Fung, J.J. & Starzl, T.E. (1996) Factors associated with the development of candidemia and candidemia-related death among liver transplant recipients. *Ann. Surgery*, **223**, 70-76.
- Obayashi, T., Yoshida, M., Mori, T., Goto, H., Yasuoka, A., Iwasaki, H., Teshima, H., Kohno, S., Horiuchi, A. & Ito, A. (1995) Plasma (1 $\rightarrow$ 3)-beta-D-glucan measurement in diagnosis of invasive deep mycosis and fungal febrile episodes. *Lancet*, **345**, 17-20.
- Odabasi, Z., Mattiuzzi, G., Estey, E., Kantarjian, H., Saeki, F., Ridge, R.J., Ketchum, P.A., Finkelman, M.A., Rex, J.H. & Ostrosky-Zeichner, L. (2004) Beta-D-glucan as a diagnostic adjunct for invasive fungal infections: validation, cutoff development, and performance in patients with acute myelogenous leukemia and myelodysplastic syndrome. *Clin. Infect. Dis.*, **39**, 199-205.
- Ostrosky-Zeichner, L., Alexander, B.D., Kett, D.H., Vazquez, J., Pappas, P.G., Saeki, F., Ketchum, P.A., Wingard, J., Schiff, R., Tamura, H., Finkelman, M.A. & Rex, J.H. (2005) Multicenter clinical evaluation of the (1-3)beta-D-glucan assay as an aid to diagnosis of fungal infections in humans. *Clin. Infect. Dis.*, **41**, 654-659.
- Patel, R., Portela, D., Badley, A.D., Harmsen, W.S., Larson-Keller, J.J., Ilstrup, D.M., Keating, M.R., Wiesner, R.H., Krom, R.A. & Paya, C.V. (1996) Risk factors of invasive Candida and non-Candida fungal infections after liver transplantation. *Transplantation*, **62**, 926-934.
- Paya, C.V. (2001) Prevention of fungal and hepatitis virus infections in liver transplantation. *Clin. Infect. Dis.*, **33**, S47-S52.
- Pazos, C., Ponton, J. & Del Palacio, A. (2005) Contribution of (1 $\rightarrow$ 3)-beta-D-glucan chromogenic assay to diagnosis and therapeutic monitoring of invasive aspergillosis in neutropenic adult patients: a comparison with serial screening for circulating galactomannan. *J. Clin. Microbiol.*, **43**, 299-305.



- Rabkin, J.M., Orolloff, S.L., Corless, C.L., Benner, K.G., Flora, K.D., Rosen, H.R. & Olyaei, A.J. (2000) Association of fungal infection and increased mortality in liver transplant recipients. *Am. J. Surg.*, **179**, 426-430.
- Sakai, T., Ikegami, K., Yoshinaga, E., Uesugi-Hayakawa, R. & Wakizaka, A. (2000) Rapid, sensitive and simple detection of candida deep mycosis by amplification of 18S ribosomal RNA gene; comparison with assay of serum  $\beta$ -D-glucan level in clinical samples. *Tohoku J. Exp. Med.*, **190**, 119-128.
- Singh, N., Gayowski, T., Wagener, M.M., Doyle, H. & Marino, I.R. (1997) Invasive fungal infections in liver transplant recipients receiving tacrolimus as the primary immunosuppressive agent. *Clin. Infect. Dis.*, **24**, 179-184.
- Singh, N., Wagener, M.M., Marino, I.R. & Gayowski, T. (2002) Trends in invasive fungal infections in liver transplant recipients: correlation with evolution in transplantation practices. *Transplantation*, **73**, 63-67.
- Suzuki, T., Hashimoto, T., Nakamura, T., Kondo, S., Shimizu, Y., Nakamura, Y., Hayashi, S., Itoh, K. & Manabe, T. (2000) Fungal infection in living related liver transplantation patients. *Transplant. Proc.*, **32**, 2231-2232.
- Takakura, S., Fujihara, N., Saito, T., Kudo, T., Iinuma, Y. & Ichiyama, S. (2004) National surveillance of species distribution in blood isolates of *Candida* species in Japan and their susceptibility to six antifungal agents including voriconazole and micafungin. *J. Antimicrobial Chemotherapy*, **53**, 283-289.
- van der Waaij, D. (1982) Colonization resistance of the digestive tract: clinical consequences and implications. *J. Antimicrob. Chemother.*, **10**, 263-270.
- Verma, A., Wade, J.J., Cheeseman, P., Samaroo, B., Rela, M., Heaton, N.D., Mieli-Vergani, G. & Dhawan, A. (2005) Risk factors for fungal infection in paediatric liver transplant recipients. *Pediatr. Transplant.*, **9**, 220-225.
- Wiesner, R.H., Hermans, P.E., Rakela, J., Washington, J.A., Perkins, J.D., DiCecco, S. & Krom, R. (1988) Selective bowel decontamination to decrease gram-negative aerobic bacterial and *Candida* colonization and prevent infection after orthotopic liver transplantation. *Transplantation*, **45**, 570-574.

## Serum-derived hepatitis C virus infectivity in interferon regulatory factor-7-suppressed human primary hepatocytes

Hussein H. Aly<sup>1,2,3</sup>, Koichi Watashi<sup>2</sup>, Makoto Hijikata<sup>2</sup>, Hiroyasu Kaneko<sup>2</sup>, Yasutugu Takada<sup>1</sup>, Hiroto Egawa<sup>1</sup>, Shinji Uemoto<sup>1</sup>, Kunitada Shimotohno<sup>2,\*</sup>

<sup>1</sup>Graduate School of Medicine, Department of Transplant Surgery, Kyoto University Hospital, Kyoto, Japan

<sup>2</sup>Laboratory of Human Tumor Viruses, Institute of Virus Research, Kyoto University, Japan

<sup>3</sup>Hepatology Department, National Hepatology and Tropical Medicine Research Institute, Cairo, Egypt

See Editorial, pages 1–5

**Background/Aims:** The development of an efficient *in vitro* infection system for HCV is important in order to develop new anti-HCV strategy. Only Huh7 hepatocyte cell lines were shown to be infected with JFH-1 fulminant HCV-2a strain and its chimeras. Here we aimed to establish a primary hepatocyte cell line that could be infected by HCV particles from patients' sera.

**Methods:** We transduced primary human hepatocytes with human telomerase reverse transcriptase together with human papilloma virus 18/E6E7 (HPV18/E6E7) genes or simian virus large T gene (SV40 T) to immortalize cells. We also established the HPV18/E6E7-immortalized hepatocytes in which interferon regulatory factor-7 was inactivated. Finally we analyzed HCV infectivity in these cells.

**Results:** Even after prolonged culture HPV18/E6E7-immortalized hepatocytes exhibited hepatocyte functions and marker expression and were more prone to HCV infection than SV40 T-immortalized hepatocytes. The susceptibility of HPV18/E6E7-immortalized hepatocytes to HCV infection was further improved, in particular, by impairing signaling through interferon regulatory factor-7.

**Conclusions:** HPV18/E6E7-immortalized hepatocytes are useful for the analysis of HCV infection, anti-HCV innate immune response, and screening of antiviral agents with a variety of HCV strains.

© 2006 European Association for the Study of the Liver. Published by Elsevier B.V. All rights reserved.

**Keywords:** Immortalization; Primary hepatocytes; HCV infection; IRF-7; IRF-3; HPV18/E6E7; Innate immune response

### 1. Introduction

Infection with Hepatitis C virus (HCV) is a serious problem worldwide since 3% of the world's population is chronically infected [1]. Chronic HCV may lead to liver cirrhosis and hepatocellular carcinoma. Current stan-

dard therapy utilizes the combination of pegylated interferon- $\alpha$  and ribavirin, which results in a sustained response in only 30–60% of patients [2–5]. Many patients, however, do not qualify for or tolerate standard therapy [6]. Thus, it is important to develop an efficient *in vitro* infection system for HCV to facilitate the discovery of new anti-HCV strategies. Only Huh7 cell line is permissive for replication, infection and release of the fulminant hepatitis-derived HCV-2a (JFH-1) strain and its chimeric derivatives [7–9]. No other hepatocyte cell lines are able to support HCV replication efficiently.

Received 5 June 2006; received in revised form 24 July 2006; accepted 1 August 2006; available online 30 October 2006

\* Corresponding author. Tel.: +81 75 751 4000; fax: +81 75 751 3998.

E-mail address: kshimoto@virus.kyoto-u.ac.jp (K. Shimotohno).

Normal human hepatocytes are the ideal system in which to study HCV infectivity. When cultured in vitro, however, they proliferate poorly and divide only a few times [10]. Continuous proliferation could be achieved however by introducing oncogenes, such as Simian virus large tumor antigen (SV40 T) [11]. This often resulted in tumor development [12] together with numerical (aneuploidy) and structural (aberrations) chromosome abnormalities [13]. The human papilloma virus E6E7 genes (HPV/E6E7) immortalized multiple cell types that were phenotypically and functionally similar to the parental cells [14–20]. As yet, no human hepatocytes have been immortalized with HPV18/E6E7.

We established a human primary non-neoplastic hepatocyte cell line transduced with the HPV18/E6E7 that retained primary hepatocyte characteristics even after prolonged culture, and were more prone to HCV infection than those cells immortalized with SV40 T antigen. We further improved the susceptibility of HPV18/E6E7-immortalized hepatocytes to HCV infectivity by impairing interferon regulatory factor-7 (IRF-7) expression. These cells are useful to assay infectivity of HCV strains other than JFH-1, HCV replication, innate immune system engagement of HCV, and screening of anti-HCV agents. This infection system using non-neoplastic cells also suggested that IRF-7 plays an important role in eliminating HCV infection.

## 2. Materials and methods

### 2.1. Cell cultures

We obtained the approval of the Ethical Committee of Kyoto University for the use of human hepatocytes and sera obtained from HCV-positive patients. Informed consent was obtained from both the hepatocyte donor and HCV-positive patients. Primary hepatocytes (P.H.) were cultured as described [21]. HeLa, 293, Huh-7.5, and PH5CH8 cells were cultured as previously described [22]. For three-dimensional (3D) cultures, Mebiol Gel (Mebiol Inc.) was prepared according to the manufacturer's instructions.

### 2.2. Plasmids construction

The SV40 T, hTERT and HPV/E6E7 fragments from pAct-SVT, PCX4neo/hTERT, and pLXSN-E6E7 plasmids were inserted into pCSII-EF-RFA plasmid creating the pCSII-EF-SVT, pCSII-EF-hTERT, and pCSII-EF-E6E7 plasmids, respectively. The full-length IRF-3 and IRF-7 genes were cloned by RT-PCR using total RNA isolated from 293 cells as a template and were inserted into pcDNA3 vector. Dominant-negative forms of IRF-3 (DNIRF-3) and IRF-7 (DNIRF-7) were constructed by PCR amplification of the coding region for amino acid residues 108–427 of IRF-3 and 237–514 of IRF-7, respectively. The amplified IRF-3 fragment was cloned into pcDNA3 in frame with a FLAG epitope tag generating pcFLAG-DNIRF-3. The amplified IRF-7 fragment was cloned into pLXSH in frame with HA epitope tag generating pLXSH-HA-DNIRF-7. The pIFN $\beta$  promoter-luc and pIFN $\alpha$  promoter-luc plasmids were gifts from Dr. Taniguchi of the Tokyo University. The psiRNA-hIRF-3 and psiRNA-hIRF-7 plasmids were purchased from InvivoGen (USA).

### 2.3. Immunoblot analysis

Immunoblot analysis was performed as described previously [22]. We used anti-SV40 T (Santa Cruz), anti-HPV18/E7 (Santa Cruz), anti-tubulin (Sigma), anti-FLAG (Sigma), and anti-HA (Sigma) antibodies.

### 2.4. Transfection, small interfering RNA silencing and luciferase assays

Transfection of plasmid DNA was performed using Effectene transfection reagent (Qiagen) as recommended by the manufacturer. The pLXSH-HA-DNIRF-7 plasmid was transfected into the HuS-E/2 clone; transfectants were selected in 100  $\mu$ g/ml hygromycin B (Gibco). The psiRNA-hIRF-3 and psiRNA-hIRF-7 plasmids were separately transfected into HuS-E/2 cells followed by Zeocin (250  $\mu$ g/ml) selection. After two weeks of continuous selection, cells were infected with HCV. Luciferase assays were conducted as previously described [22]. The results are presented as relative light units (RLU) normalized to the total content of protein in the cell lysates.

### 2.5. Reverse transcriptase polymerase chain reaction (RT-PCR) and real-time RT-PCR

Using 250 ng of total RNA as a template, we performed RT-PCR with a one-step RNA PCR kit (Takara) according to the manufacturer's instructions. The primer sets and reaction conditions used are detailed in Table 1. To measure HCV-RNA titers by real-time RT-PCR, we collected RNA from infected wells. Five hundred nanograms of total cellular RNA was analyzed for the quantity of HCV-RNA as previously described [23].

### 2.6. HCV infection experiment

HCV infection experiment from serum was done as mentioned before [22]. HCV-infected-serums were titrated and  $1 \times 10^5$  HCV-RNA copies/ml were used for each infection experiment. Concentrated culture medium for HCV/JFH1-producing cells was prepared as previously described [7]. HCV titer in the concentrated medium was measured, adjusted and added to the cells as mentioned above.

### 2.7. Blocking of HCV infectivity by anti-CD81

Inhibition of HCV infectivity was performed by blocking CD81 as previously described [7].

## 3. Results

### 3.1. Establishment of immortalized primary human hepatocytes

Primary hepatocytes were isolated from liver tissue obtained from a 9-year-old male patient with Primary Hyperoxaluria who had undergone liver transplantation. Hepatocytes were left unmanipulated or transduced with CSII-EF-hTERT alone or in combination with CSII-EF-SVT or CSII-EF-E6E7 to enhance the efficiency of immortalization. After six weeks only cells transduced by the combination of hTERT and either LT or HPV18/E6E7 continued to proliferate. Initially appearing colonies with a growth advantage were picked up and expanded. SV40 T-immortalized cell clones were named HuS-T cells and given numbers from 1 to 7,

**Table 1**  
**Primer sequences and RT-PCR parameters**

| Genes            | Primer sequence 5'-3'   | PCR parameters <sup>a</sup>       |
|------------------|---|-----------------------------------|
| HGF              | F: AGGAGCCAGCCTGAATGATGA<br>R: CCCTCTGATGTCCCAAGATTAGC                      | 95, 56, 72<br>1 min, 45 s, 1 min  |
| TGF $\alpha$     | F: ATGGTCCCCTCGGCTGGA<br>R: GGCCTGCTTCTTCTGGCTGGCA                          | 95, 59, 72<br>45 s, 30 s, 1 min   |
| TGF $\beta$ 1    | F: GCCCTGGACACCAACTATTGCT<br>R: AGGCTCCAAATGTAGGGGCAGG                      | 95, 58, 72<br>45 s, 30 s, 1 min   |
| TGF $\beta$ 2    | F: GATTTCCATCTACAAGACCACGAGGGACTTGC<br>R: CAGCATCAGTTACATCGAAGGAGAGCCATTTCG | 95, 58, 72<br>45 s, 30 s, 1 min   |
| HGFR             | F: TGGTCCTTGGCGTCGTCCTC<br>R: CTCATCATCAGCGTTATCTTC                         | 95, 54, 72<br>30 s, 45 s, 1 min   |
| EGFR             | F: CTACCACCACTCTTTGAACTGGACCAAGG<br>R: TCTATGCTCTCACCCGTTCCAAGTATCG         | 95, 58, 72<br>45 s, 30 s, 1 min   |
| TGF $\beta$ 1R   | F: CGTGCTGACATCTATGCAAT<br>R: AGCTGCTCCATTGGCATAAC                          | 95 s, 54, 72<br>30 s, 45 s, 1 min |
| TGF $\beta$ 2R   | F: TGCACATCGTCTGTGGAC<br>R: GTCTCAAAGTCTCTGAAGTGTTTC                        | 95, 58, 72<br>45 s, 30 s, 1 min   |
| FGFR             | F: ATGTGGAGCTGGAAGTGCCTC<br>R: GGTGTTATCTGTTCTTTCTCC                        | 95, 54, 72<br>30 s, 45 s, 1 min   |
| IGF-1R           | F: ACCCGGAGTACTTCAGCGCT<br>R: CACAGAAGCTTCGTTGAGAA                          | 95, 54, 72<br>30 s, 45 s, 1 min   |
| HNF1 $\alpha$    | F: GTGTCTACAAGTGGTTTGCC<br>R: TGTAGACACTGTCACTAAGG                          | 95, 52, 72<br>45 s, 30 s, 1 min   |
| HNF1 $\beta$     | F: GAAACAATGAGATCACTTCCTCC<br>R: CTTTGTGCAATTGCCATGACTCC                    | 95, 52, 72<br>1 m, 45 s, 1 min    |
| HNF3 $\beta$     | F: CACCCTACGCCTTAACCAC<br>R: GGTAGTAGGAGGTATCTGCGG                          | 95, 56, 72<br>1 m, 45 s, 1 min    |
| HNF4             | F: CTGCTCGGAGCCACAAAGAGATCCATG<br>R: ATCATCTGCCACGTGATGCTCTGCA              | 95, 58, 72<br>45 s, 30 s, 1 min   |
| Albumin          | F: AGTTTGCAGAAGTTTCCAAGTTAGTG<br>R: AGGTCCGCCCTGTATCAG                      | 95, 55, 72<br>45 s, 30 s, 1 min   |
| Apolipoprotein-a | F:AGGCTCGGCATTTCTGGCAG<br>R: TATCCCAGAAGTCTGGGTC                            | 95, 55, 72<br>45 s, 30 s, 1 min   |
| HTF              | F: TCGCTACAGCCTTTGCAATG<br>R: TTGAGGGTACGGAGGAGTTCC                         | 95, 55, 72<br>45 s, 30 s, 1 min   |
| E-cadherin       | F: TCCATTTCTGGTCTACGCC<br>R: TTTGTCTACCGACTTCCAC                            | 95, 55, 72<br>45 s, 30 s, 1 min   |
| CYP 1B1          | F: CACCAAGGCTGAGACAGTGA<br>R: GCCAGGTAAACTCCAAGCAC                          | 94, 57, 72<br>30 s, 30 s, 1 min   |
| CYP 2C9          | F: GGACAGAGACGACAAGCACA<br>R: TGGTGGGGAGAAGGTCAAT                           | 94, 57, 72<br>30 s, 30 s, 1 min   |
| CYP 2B           | F: GGCACACAGCCAAGTTTACA<br>R: CCAGCAAAGAAGAGCGAGAG                          | 94, 57, 72<br>30 s, 30 s, 1 min   |
| CYP 3A4          | F: TGTGCCTGAGAACACCAGAG<br>R: GCAGAGGAGCCAAATCTACC                          | 94, 57, 72<br>30 s, 30 s, 1 min   |
| CYP 2E1          | F: CCGCAAGCATTTTGACTACA<br>R: GCTCCTTACCCTTTCAGAC                           | 94, 57, 72<br>30 s, 30 s, 1 min   |
| CYP 1A1          | F: AGGCTTTTACATCCCAAGG<br>R: GCAATGGTCTCACCGATACA                           | 94, 57, 72<br>30 s, 30 s, 1 min   |
| GAPDH            | F: CCATGGAGAAGGCTGGGG<br>R: CAAAGTTGTCTGGATGACC                             | 95, 8, 72<br>45 s, 30 s, 1 min    |

Table 1 (continued)

| Genes | Primer sequence 5'–3'                               | PCR parameters <sup>a</sup>     |
|-------|---|---------------------------------|
| CD81  | F: CTCAACTGTTGTGGCTCCAAC<br>R: CCAATGAGGTACAGCTTCCC | 95, 55, 72<br>45 s, 30 s, 1 min |
| TLR3  | F: GATCTGTCTCATAATGGCTTG<br>R: GACAGATCCGAATGCTTGTG | 95, 55, 72<br>45 s, 30 s, 1 min |
| TLR7  | F: CCAGACATCTCCCACGGTC<br>R: GGCAAAACAGTAGGGACGGC   | 95, 55, 72<br>45 s, 30 s, 1 min |
| TLR8  | F: CTGTGAGTTATGCGCGAAG<br>R: CGGGATTTCGGTCTGGTGC    | 95, 55, 72<br>45 s, 30 s, 1 min |
| Myd88 | F: GGTCTCCTCCACATCCTCCC<br>R: CCAGCTTGGTAAGCAGCTCG  | 95, 55, 72<br>45 s, 30 s, 1 min |
| IRF3  | F: GAACCCCAAAGCCACGGATC<br>R: CCTCCCGGAACATATGCAC   | 95, 55, 72<br>45 s, 30 s, 1 min |
| IRF7  | F: GTGCTGTTCGGAGAGTGGCTC<br>R: CAGCCCAGGCCTGAAGATG  | 95, 55, 72<br>45 s, 30 s, 1 min |

CYP, cytochrome P450; EGFR, epidermal growth factor receptor; F, forward primer; FGFR, fibroblast growth factor receptor; GAPDH, glyceraldehyde phosphate dehydrogenase; HGF, hepatocyte growth factor; HGFR, hepatocyte growth factor receptor; HNF, hepatocyte nuclear factor; HTF, human transferrin; IGF-1R, insulin-like growth factor-type I receptor; IRF, interferon regulatory factor; R, reverse primer; TGF, transforming growth factor; TGFR, transforming growth factor receptor; TLR, toll like receptor.

<sup>a</sup> Temperatures are tabulated in the first lane in degrees celsius and the corresponding times in the second lane. Performing one-step RT-PCR, reverse transcription was carried out at 42 °C for 20 min with a pre-PCR denaturation at 95 °C for 10 min.




Fig. 1. (A) Immunoblot detection of SV40 T and HPV E7 expression in HuS-T/2 and HuS-E/2 cells, respectively. 293-FT and HeLa cells were used as positive controls for SV40 T and HPV E7 expression, respectively. The specific bands representing the targets are indicated. Detection of tubulin expression in all cells served as an internal control. (B) Human Telomerase Reverse Transcriptase (hTERT) expression was examined by RT-PCR in freshly isolated hepatocytes and the HuS-E/2 and HuS-T/2 cell lines. GAPDH expression was used as an internal control. The hTERT-specific bands are shown. (C) Morphological characteristics of HuS-E/2 and HuS-T/2 cells after two (panels 1 and 2) and 30 (panels 3 and 4) weeks in culture. [This figure appears in colour on the web.]

while the HPV18/E6E7-immortalized clones were named HuS-E cells and given numbers from 1 to 4. Expression of SV40 T and HPV E7 proteins was detected in the appropriate cells by immunoblot analysis (Fig. 1A). In both immortalized cell lines, expression of hTERT-mRNA was enhanced in comparison to non-transduced, freshly isolated hepatocytes as determined by RT-PCR (Fig. 1B). HuS-E cells were larger in size and exhibited slower growth than HuS-T cells (Fig. 1C).

3.2. Characterization of HuS-E and HuS-T immortalized hepatocytes

The HuS-E/2 and HuS-T/2 clones demonstrated the highest expression of hepatocyte-specific markers and transcription factors by RT-PCR (data not shown); these cells were used as representative for each group in this study. To address if HuS-E/2 and HuS-T/2 maintained similar characteristics as primary hepatocytes, they were both cultured continuously for 30 weeks and the expression profiles of a variety of growth factors (Fig. 2A),

growth factor receptors (Fig. 2B), hepatocyte-specific nuclear factors (Fig. 2C), albumin, apolipoprotein-A1, transferrin (Fig. 2D), cytochrome p450 (CYP) genes (Fig. 2E), and GAPDH were compared with freshly isolated primary hepatocytes after isolation or two weeks of culture, Huh-7.5 cells, and 293 cells. After two weeks in culture, the expression of nearly all examined genes was similar between freshly isolated hepatocytes and the HuS-E/2 cell line. HuS-E/2 cells, however, exhibited higher expression of TGFβ2 (Fig. 2A), TGFβ2R, and HGFR (Fig. 2B) and lower expression of CYP 3A4 and 2C9 (Fig. 2E) in comparison to freshly isolated hepatocytes. Primary hepatocytes displayed reduced expression of TGFβ1 and TGFβ2 (Fig. 2A) and a loss of CYP1A1 expression (Fig. 2E) after two weeks of culture. HuS-E/2 cells exhibited higher expression of HGF (Fig. 2A), HGF receptor (Fig. 2B), HNF-4, (Fig. 2C), albumin, apolipoprotein-A1, HTF, and E-cadherin (Fig. 2D) in comparison to HuS-T/2 cells. Expression of CYP 3A4 (Fig. 2E) was lost from both HuS-T/2 and HuS-E/2 cells, while HuS-T/2 cells also lost the expression of HNF-1α (Fig. 2C), and CYPs 2B, 2E1 (Fig. 2E).




Fig. 2. Expression of the genes encoding growth factors (A), growth factor receptors (B), hepatocyte-specific nuclear factors (C), hepatocyte differentiation and functional markers (D), and CYP enzymes (E) in 293 cells, freshly isolated primary hepatocytes (P.H.), primary hepatocytes cultured for two weeks (Cult. P.H.), Huh-7.5 cells, and HuS-E/2 and HuS-T/2 cells cultured for two and 30 weeks were investigated by RT-PCR. The bands representing specific targets are indicated in the representative reactions.




Fig. 3. (A) The morphology of HuS-E/2 and HuS-T/2 cells in 3D culture. HuS-E/2 and HuS-T/2 cells were cultured in Mebiol Gel in 12-well plates at a concentration of  $5 \times 10^5$  cells/well. The microscopic characteristics of these cells after one week of 3D culture are shown. (B) The expressions of HNF4 $\alpha$ , HNF1 $\beta$ , albumin, and apo-A by HuS-E/2 and HuS-T/2 cells in both flat and 3D cultures are detailed. After one week of culture of HuS-E/2 and HuS-T/2 cells in flat and 3D cultures, the expressions of HNF4 $\alpha$ , HNF1 $\beta$ , albumin, and apo-A were measured by RT-PCR in 250 ng total RNA.

HuS-T/2 but not in HuS-E/2 cells showed a transformed-like character starting from the 13th week of culture. This was demonstrated by continuing proliferation after confluence, pile-up formations (Fig. 1C), and proliferating in serum-depleted condition. However, HuS-E/2 cells did not show any transformed-like characters even after 30 weeks of culture.

### 3.3. The characteristics of HuS-E and HuS-T immortalized hepatocytes in 3D culture

After one week in 3D culture, HuS-E/2 (Fig. 3A, panels 1, 2, and 3) cells adopted a donut-shaped structure with a central pore, while HuS-T/2 cells (Fig. 3A, panels 4, 5, and 6) displayed irregular mass formations (similar to the growth pattern of Huh-7.5 cells in 3D culture (data not shown)). In 3D culture, while the expression of HNF4, HNF1 $\beta$ , and albumin was enhanced in HuS-E/2, it was decreased in HuS-T/2 cells (Fig. 3B).

### 3.4. HCV infection to HuS-E/2

We further assessed the HCV infectivity of HuS-E- and HuS-T-derived clones by infection with HCV-1b-in-

fecting serum. Of the three HuS-E clones examined, HuS-E/2 clone demonstrated the highest infectability with HCV genotype 1b in comparison to Huh-7.5, PH5CH8 (Fig. 4A), and HuS-T cells (data not shown), which were excluded from further experiments.

### 3.5. Anti-CD81 blocked HCV infectivity

CD81 is involved in the entry of HCV pseudoparticles [24] and in vitro-synthesized JFH-1 [7]. To determine if authentic viral particles follow the same route of entry when infecting HuS-E/2 cells, we first examined the CD81 expression by RT-PCR. Both HuS-E/2 and HuS-T/2 cells expressed similar amounts of CD81 as freshly isolated hepatocytes and Huh-7.5 cells (Fig. 4B). Antibodies against CD81 reduced HCV infectivity of HuS-E/2 cells from the levels seen using a non-specific control antibody, confirming the importance of CD81 in HCV infectivity (Fig. 4C).

### 3.6. IFN $\alpha$ blocked HCV infectivity

We treated HuS-E/2 cells with HCV-containing serum. Cells were then cultured in fresh medium supplemented




Fig. 4. (A) Serum from an HCV patient was used to infect Huh-7.5 cells, PH5CH8 cells, and three HPV E6E7-immortalized clones (HuS-E/1-3) for 24 h. After washing three times in phosphate-buffered saline (PBS), cells were cultured in fresh medium. Cells were then harvested and lysed at the indicated time points. The quantity of HCV genome RNA per 500 ng total RNA was determined by real-time RT-PCR analysis. (B) HuS-E/2 and HuS-T/2 cells both expressed CD81. Expression of CD81 (upper panel) and GAPDH as an internal control (lower panel) in 293 cells, freshly isolated P.H., cultured P.H., and Huh-7.5, HuS-E/2, and HuS-T/2 cells was investigated by RT-PCR. (C) Anti-CD81 antibodies blocked HCV infectivity. HCV infection was performed as described in (A) with the addition of CD81-specific (black bar) or anti-tubulin antibodies (control, white bar). (D) IFN $\alpha$  inhibits HCV multiplication in HuS-E/2 cells infected with HCV-containing serum. HuS-E/2 cells were infected with HCV as described in (A). After washing three times with PBS, cells were cultured in fresh medium supplemented with (black bar) or without (white bar) 100 U/ml IFN $\alpha$ .

without or with 100 U/ml IFN $\alpha$ . The enhancement of the HCV-RNA genome titers on the fifth day (about 10-fold) was not observed in cells treated continuously with IFN $\alpha$  (Fig. 4D). This result suggests that IFN $\alpha$  inhibited HCV replication in infected HuS-E/2 cells.

### 3.7. The effect of blocking IRF-3 and IRF-7 signaling on HCV infectivity

Production of interferon-alpha (IFN $\alpha$ ) and interferon-beta (IFN $\beta$ ) limits viral replication and spread, providing one of the most effective innate antiviral responses [25]. Signaling through IRF-3 and IRF-7 plays important roles in the stimulation of IFN- $\alpha/\beta$  production [25]. To determine which molecules (IRF-3 or IRF-7) play an important role in modulation of the innate immune response against HCV infection in these cells, we first detected intrinsic expression of double-stranded RNA-stimulated Toll-like receptor (TLR) 3, the downstream effector IRF-3, single-stranded RNA-stimulated TLR7, and 8, and the downstream effectors MyD88 and IRF-7 by RT-PCR. TLR3 exhibited very low expression in freshly isolated hepatocytes, Huh-7.5, HuS-E/2, and HuS-T/2 cells, while TLR7, TLR8, MyD88, and IRF-7 were easily detectable in both freshly isolated and immortalized cell lines (Fig. 5A).

The abilities of DNIRF-3 and DNIRF-7 to inhibit IFN $\beta$  and IFN $\alpha$  production by HuS-E/2 cells infected with Sendai virus were confirmed using assays of IFN $\beta$  or IFN $\alpha$  promoter-driven luciferase reporters. DNIRF-3 exhibited strong inhibition of IFN $\beta$  production (Fig. 5B) and weaker inhibition of IFN $\alpha$  transcription (Fig. 5C), while DNIRF-7 strongly inhibited IFN $\alpha$  production (Fig. 5C) and only weakly inhibited IFN $\beta$  production (Fig. 5B).

We then assessed the inhibition of HCV infectivity by DNIRF-3 and DNIRF-7. Transient transfection with DNIRF-3, DNIRF-7, or an empty vector was performed prior to HCV infection. Using Effectene reagent, the efficiency of plasmid transfection into HuS-E/2 cells was approximately 70% (data not shown). While there was no significant effect of DNIRF-3 on HCV infectivity, DNIRF-7 demonstrated a marked increase in HCV titers on days 3 and 5 after infection in comparison to control cells (Fig. 5D). To confirm that the enhancement of HCV replication by DNIRF-7 is not mediated by the impairment of IRF-3 signaling by heterodimeric interactions between IRF-3 and DNIRF-7, we performed siRNA inhibition of IRF-3 and IRF-7. The reduction of IRF-3 and IRF-7 expression by siRNA was obvious by RT-PCR (Fig. 5E). siRNA-mediated suppression of either IRF-3 or IRF-7 inhibited IFN $\beta$  and IFN $\alpha$  production






Fig. 5. (A) We examined the expression of TLR3, TLR7, TLR8, MyD88, IRF-3, and IRF-7, as well as GAPDH as an internal control in freshly isolated primary hepatocytes and Huh-7.5, HuS-E/2, and HuS-T/2 cells was investigated by RT-PCR. (B and C) HuS-E/2 cells were cotransfected with pIFN $\beta$ -luc (B) or pIFN $\alpha$ -luc (C) with an expression plasmid encoding DNIRF-3, DNIRF-7, or the appropriate empty vector (pcDNA3 and PLXSH, respectively). Twenty-four hours later, cells were infected (black bar) with Sendai virus or mock-infected (white bar), then analyzed for luciferase activity after 12 h. (D) IRF-7, but not IRF-3, suppression enhanced HCV infectivity of HuS-E/2 cells. HuS-E/2 cells were transiently transfected with empty pcDNA3, DNIRF-3, empty pLXSH, or DNIRF-7 plasmids. Twenty-four hours later, serum from a patient with HCV was used to infect transfected cells for 24 h. After washing, cells were cultured in fresh medium. The cells were then harvested and lysed at the indicated time points. The quantity of HCV genome RNA per 500 ng total RNA was determined by real-time RT-PCR analysis. (E) IRF-3 and IRF-7 levels were suppressed by specific siRNAs. HuS-E/2 cells were harvested and assessed for the expression of IRF-3 and IRF-7 by RT-PCR. (F and G) HuS-E/2 cells were transfected with control psiRNA-hTLR2, psiRNA-hIRF-3, or psiRNA-hIRF-7, followed by selection in Zeocin at 250  $\mu$ g/ml. Two weeks later, cells were cotransfected with pIFN $\beta$ -luc (F) or pIFN $\alpha$ -luc (G). Twenty-four hours later, cells were infected (black bar) with Sendai virus or mock-infected (white bar), then analyzed for luciferase activity after 12 h. (H) Transfected cells were infected with serum from HCV patient; HCV infectivity was assessed as described above.

in HuS-E/2 cells infected with Sendai virus in patterns similar to the effects seen following DNIRF-3 and DNIRF-7 expression, respectively (Figs. 5F and G). Blockade of IRF-7 expression resulted in a significantly higher titer of HCV after infection, while IRF-3 down-regulation did not have any significant effect on HCV titers (Fig. 5H). The enhancement of IRF-7 silencing by siRNA improved the infectivity of HCV (data not shown). These results suggest that IRF-7 plays the major role in the innate immune response to HCV in HuS-E/2 cells.

### 3.8. Establishment of stable DNIRF-7 expressing clones derived from HuS-E/2 cells

Since DNIRF-7 enhanced HCV infectivity, we transduced the plasmid encoding DNIRF-7 and a hygromycin-B resistance gene, into HuS-E/2 cells. Following selection with hygromycin-B, we obtained the HuS-E7/DN22 and HuS-E7/DN24 clones. As detected by RT-PCR, both clones demonstrated similar expression levels

of albumin, apolipoprotein-A1, and HNF4 as the parental HuS-E/2 cells (Fig. 6A). The HuS-E7/DN24 clone exhibited stronger expression of DNIRF-7 than the HuS-E7/DN22 clone by immunoblotting (Fig. 6B). The induction of IFN $\alpha$  in HuS-E7/DN24 in response to infection with an RNA virus (Sendai virus) was low in comparison to the parental HuS-E/2 and HuS-E7/DN22 clones, as detected by IFN $\alpha$ -luciferase reporter assay (Fig. 6C). HuS-E7/DN24 also exhibited a higher HCV infectability in comparison to parental HuS-E/2 cells and the HuS-E7/DN22 clone (Fig. 6D).

### 3.9. Infection of HuS-E7/DN24 cells with different HCV genotypes

Huh7.5 and HuS-E7/DN24 cells were separately infected with serums derived from 3 different HCV-patients or by JFH-1 concentrated medium (HCV-2a). Two serums were infected by HCV-1b, while the third by HCV-2b. Inoculated virus titer was adjusted to be the same in all cases. Except for JFH-1, which efficiently




Fig. 6. (A) The pLXSH-HA-DNIRF-7 plasmid was transfected into HuS-E/2 cells, followed by selection in 100  $\mu$ g/ml Hygromycin B. Two clones, HuS-E7/DN22 (lane 2) and HuS-E7/DN24 (lane 3), were obtained. We investigated the expression of albumin, apo-A, HNF4, and GAPDH as an internal control in parental HuS-E/2, HuS-E7/DN22, and HuS-E7/DN24 hepatocytes cultured for two weeks by RT-PCR. (B) Expression of HA-tagged DNIRF-7 (upper panel) and tubulin (control, lower panel) was detected by immunoblotting analysis. HuS-E/2 cells transiently transfected with either empty pLXSH vector (lane 1) or pLXSH-HA-DNIRF-7 (lane 2) were used as negative and positive controls, respectively, after 48 h. (C) HuS-E/2, HuS-E7/DN24, and HuS-E7/DN22 cells were transfected with IFN $\alpha$ -luc. HuS-E/2 cells were also cotransfected with pLXSH. All of these cells were then infected (black bar) or with Sendai virus or mock-infected, then analyzed for luciferase activity after 12 h. (D) HuS-E7/DN24 cells exhibited high infectivity to HCV samples derived from patient serum. HuS-E/2 cells were transiently transfected with empty pLXSH. Twenty-four hours later, serum from a recurrently transplanted HCV patient was used to infect transfected cells and HuS-E7/DN22 and HuS-E7/DN24 cells for 24 h. After washing three times, cells were cultured in fresh medium. Cells were then harvested and lysed at the indicated time points.




Fig. 7. The infectability of Huh-7.5 and HuS-E7/DN24 cells to different HCV genotypes. Huh-7.5 (A) and HuS-E7/DN24 (B) cells were infected with same titer of JFH1 (HCV-2a), two different HCV-1b serums and one HCV-2b serum. After removing the infected medium, the cells were washed in PBS and recultured in fresh medium. Cells were harvested and lysed at the indicated time points. The quantity of HCV genome RNA per 500 ng RNA was detected by real-time RT-PCR analysis.

replicated in Huh7.5 cells (Fig. 7A), HuS-E7/DN24 cells showed a higher and reproducible infectability for the different HCV strains than Huh7.5 cells (Fig. 7B). Similar higher infectability of HuS-E7/DN24 cells was observed with HCV-4a genotype (unpublished data). These results suggest that the high infectability of Huh-7.5 with JFH-1 is specific among the combinations of HCV strains and cell lines; while HuS-E7/DN24 cells were generally permissive to HCV-infected serum independent of HCV strains.

#### 4. Discussion

This study demonstrates that ectopic expression of the HPV18/E6E7 genes in combination with hTERT could efficiently immortalize mature human hepatocytes, generating a cell line with stable expression of hepatocyte markers and functions for more than 30 weeks in culture. HuS-E/2 cells continuously exhibited higher expression of both HGF and HGFR than HuS-T/2 cells. This result suggests that HPV18/E6E7-immortalized hepatocytes maintain responsiveness to paracrine signals capable of inducing cell differentiation to a greater extent than SV40 T-immortalized hepatocytes. This conclusion is further supported by the increased expression of HNF4 in HuS-E/2 cells in comparison to HuS-T/2 cells. HNF4 is a major hepatocyte transcription factor, required for hepatocyte differentiation and liver-specific gene expression [26]. HNF4 drives hepatocytes differentiation by acting upstream in a transcription factor cascade that included HNF1 $\alpha$  [27]. HuS-E/2 cells continued to express HNF1 $\alpha$  throughout prolonged culture, while HuS-T/2 cells lost expression completely. Maintenance of hepatocellular functions was demonstrated by continuous and high expression of albumin, apolipoprotein-A, human transferrin, and E-cadherin by HuS-E/2 in comparison to HuS-T/2 cells. These differences became more pronounced in the late passages. In a similar manner, HuS-E/2 cells continued to express all of the examined CYP genes, with the exception of CYP 3A4, while HuS-T/2 cells lost expression of CYP 3A4, 1B, and 2E1 completely and displayed markedly lower expression of CYP 1B1 than HuS-E/2 cells. Thus, human hepatocytes immortalized by HPV E6/E7 transfection are phenotypically similar to primary hepatocytes, even during extended cultures.

Recently, it was reported that the JFH-1 strain and derived chimeras could only infect and propagate efficiently in Huh7.5.1 and Huh7.5 cells, both of which are subclones of Huh7 cells [7–9]. This limitation, however, may be specific to the JFH-1 strain, which may not accurately reflect the course of other HCV strains' infection. Thus, usage of HCV particles isolated from patient serum could be more useful to study authentic HCV infection. Using sera from HCV patients as a source

of infective virus, HPV18/E6E7-immortalized cell lines exhibited higher reproducible susceptibility to HCV infection than HuS-T, PH5CH8, and Huh-7.5 cell lines.

IRF3 and IRF7 play an important role in the activation of interferon signaling [28]. We suppressed the functions of IRF-3 or IRF-7 to assess their role in HCV infectivity. In fact, we observed significant increase of HCV replication in HuS-E/2 cells bearing dominant-negative IRF7 that impaired IFN signaling. The suppression of IRF-3, however, did not have any significant effect on HCV infectivity or replication in this cell line. This may result from the blockade of IRF-3 activation by an HCV NS3/4A serine protease [29] through at least two independent pathways that inhibit the TLR3-dependent and RIG-I-dependent signaling pathways [29–33]. Although HCV was shown to inhibit basal expression levels of IRF-7 at both mRNA and protein levels and it was shown that NS5A suppresses IRF-7-induced IFN $\alpha$  promoter activation [34]. Stimulation of TLR7 was shown to activate IRF-7 and induce suppression of HCV replicon levels in Huh-7 cells [35]. This suggests that the inhibition of IRF7 by HCV is not complete. Using IRF-7-deficient (IRF-7 $-/-$ ) mice, Honda [36] demonstrated that the transcription factor IRF-7 is essential for the induction of IFN $\alpha/\beta$  genes. We established a clone stably expressing DNIRF-7 (HuS-7E/DN24), which demonstrated higher infectivity with different HCV strains than the parental HuS-E/2 clone.

In summary, we have established a human hepatocyte-derived cell line that maintains the characteristic features of primary hepatocytes by transduction with HPV18/E6E7. This cell line is highly infectable by HCV, which suggests that these cells may be useful to characterize the molecular mechanisms involved with HCV infection and to develop novel HCV treatment modalities.

#### Acknowledgements

We thank Dr. Akagi at Osaka Bioscience Institute for providing hTERT expressing vector, Dr. Sakai at the Institute for Virus Research, Kyoto University, for providing HPV18/E6E7 expressing plasmid, and Dr. Taniguchi of the University of Tokyo for providing IFN $\beta$ , and IFN $\alpha$  promoters' reporter plasmids. This work was supported by Grants-in-Aid for cancer research and for the second-term comprehensive 10-year strategies for cancer control from the Ministry of Health, Labor and Welfare, by Grants-in-Aid for scientific research from the Ministry of Education, Culture, Sports, Science and Technology, by Grants-in-Aid for the research for the future program from the Japanese society for the Promotion of Science. Dr. Hussein H. Aly is a receiver of the Japanese *Gakushu Shorei* scholarship and was partly supported by Prof. Yassin A. El Ghaffar memorial scholarship for the Improvement of Liver research in Egypt.

## References

- [1] Wasley A, Alter MJ. Epidemiology of hepatitis C: geographic differences and temporal trends. *Semin Liver Dis* 2000;20:1–16.
- [2] Manns MP, McHutchison JG, Gordon SC, Rustgi VK, Shiffman M, Reindollar R, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet* 2001;358:958–965.
- [3] Fried MW, Shiffman ML, Reddy KR, Smith C, Marinos G, Goncalves Jr FL, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med* 2002;347:975–982.
- [4] Hadziyannis SJ, Sette Jr H, Morgan TR, Balan V, Diago M, Marcellin P, et al. Peginterferon-alpha2a and ribavirin combination therapy in chronic hepatitis C: a randomized study of treatment duration and ribavirin dose. *Ann Intern Med* 2004;140:346–355.
- [5] Muir AJ, Bornstein JD, Killenberg PG. Peginterferon alfa-2b and ribavirin for the treatment of chronic hepatitis C in blacks and non-Hispanic whites. *N Engl J Med* 2004;350:2265–2271.
- [6] Falck-Ytter Y, Kale H, Mullen KD, Sarbah SA, Sorescu L, McCullough AJ. Surprisingly small effect of antiviral treatment in patients with hepatitis C. *Ann Intern Med* 2002;136:288–292.
- [7] Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med* 2005;11:791–796.
- [8] Zhong J, Gastaminza P, Cheng G, Kapadia S, Kato T, Burton DR, et al. Robust hepatitis C virus infection in vitro. *Proc Natl Acad Sci USA* 2005;102:9294–9299.
- [9] Lindenbach BD, Evans MJ, Syder AJ, Wolk B, Tellinghuisen TL, Liu CC, et al. Complete replication of hepatitis C virus in cell culture. *Science* 2005;309:623–626.
- [10] Dellgado JP, Parouchev A, Allain JE, Pennarun G, Gauthier LR, Dutrillaux AM, et al. Long-term controlled immortalization of a primate hepatic progenitor cell line after Simian virus 40 T-Antigen gene transfer. *Oncogene* 2005;24:541–551.
- [11] Mizuguchi T, Mitaka T, Katsuramaki T, Hirata K. Hepatocyte transplantation for total liver repopulation. *J Hepatobiliary Pancreat Surg* 2005;12:378–385.
- [12] Isom HC, Tevethia MJ, Kreider JW. Tumorigenicity of simian virus 40-transformed rat hepatocytes. *Cancer Res* 1981;41:2126–2134.
- [13] Ray FA, Waltman MJ, Lehman JM, Little JB, Nickoloff JA, Kraemer PM. Identification of SV40 T-antigen mutants that alter T-antigen-induced chromosome damage in human fibroblasts. *Cytometry* 1998;31:242–250.
- [14] Chen WH, Lai WF, Deng WP, Yang WK, Lo WC, Wu CC, et al. Tissue engineered cartilage using human articular chondrocytes immortalized by HPV-16 E6 and E7 genes. *J Biomed Mater Res A* 2006;76:512–520.
- [15] Dimri G, Band H, Band V. Mammary epithelial cell transformation: insights from cell culture and mouse models. *Breast Cancer Res* 2005;7:171–179.
- [16] Harms W, Rothamel T, Miller K, Harste G, Grassmann M, Heim A. Characterization of human myocardial fibroblasts immortalized by HPV16 E6–E7 genes. *Exp Cell Res* 2001;268:252–261.
- [17] Shiga T, Shirasawa H, Shimizu K, Dezawa M, Masuda Y, Shimizu B. Normal human fibroblasts immortalized by introduction of human papillomavirus type 16 (HPV-16) E6–E7 genes. *Microbiol Immunol* 1997;41:313–319.
- [18] Akimov SS, Ramezani A, Hawley TS, Hawley RG. Bypass of senescence, immortalization, and transformation of human hematopoietic progenitor cells. *Stem Cells* 2005;23:1423–1433.
- [19] Hung SC, Yang DM, Chang CF, Lin RJ, Wang JS, Low-Tone Ho L, et al. Immortalization without neoplastic transformation of human mesenchymal stem cells by transduction with HPV16 E6E7 genes. *Int J Cancer* 2004;110:313–319.
- [20] Wang G, Johnson GA, Spencer TE, Bazer FW. Isolation, immortalization, and initial characterization of uterine cell lines: an in vitro model system for the porcine uterus. *In vitro Cell Dev Biol Anim* 2000;36:650–656.
- [21] Hino H, Tateno C, Sato H, Yamasaki C, Katayama S, Kohashi T, et al. A long-term culture of human hepatocytes which show a high growth potential and express their differentiated phenotypes. *Biochem Biophys Res Commun* 1999;256:184–191.
- [22] Watashi K, Hijikata M, Hosaka M, Yamaji M, Shimotohno K. Cyclosporin A suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* 2003;38:1282–1288.
- [23] Murata T, Ohshima T, Yamaji M, Hosaka M, Miyanari Y, Hijikata M, et al. Suppression of hepatitis C virus replication by TGF-beta. *Virology* 2005;331:407–417.
- [24] Dasgupta A, Hughey R, Lancin P, Larue L, Moghe PV. E-cadherin synergistically induces hepatospecific phenotype and maturation of embryonic stem cells in conjunction with hepatotrophic factors. *Biotechnol Bioeng* 2005;92:257–266.
- [25] Civas A, Island ML, Genin P, Morin P, Navarro S. Regulation of virus-induced interferon-A genes. *Biochimie* 2002;84:643–654.
- [26] Ishiyama T, Kano J, Minami Y, Iijima T, Morishita Y, Noguchi M. Expression of HNFs and C/EBP alpha is correlated with immunocytochemical differentiation of cell lines derived from human hepatocellular carcinomas, hepatoblastomas and immortalized hepatocytes. *Cancer Sci* 2003;94:757–763.
- [27] Wege H, Le HT, Chui MS, Liu L, Wu J, Giri R, et al. Telomerase reconstitution immortalizes human fetal hepatocytes without disrupting their differentiation potential. *Gastroenterology* 2003;124:432–444.
- [28] Mamane Y, Heylbroeck C, Genin P, Algarte M, Servant MJ, LePage C, et al. Interferon regulatory factors: the next generation. *Gene* 1999;237:1–14.
- [29] Foy E, Li K, Wang C, Sumpter Jr R, Ikeda M, Lemon SM, et al. Regulation of interferon regulatory factor-3 by the hepatitis C virus serine protease. *Science* 2003;300:1145–1148.
- [30] Sumpter Jr R, Loo YM, Foy E, Li K, Yoneyama M, Fujita T, et al. Regulating intracellular antiviral defense and permissiveness to hepatitis C virus RNA replication through a cellular RNA helicase, RIG-I. *J Virol* 2005;79:2689–2699.
- [31] Breiman A, Grandvaux N, Lin R, Ottone C, Akira S, Yoneyama M, et al. Inhibition of RIG-I-dependent signaling to the interferon pathway during hepatitis C virus expression and restoration of signaling by IKKepsilon. *J Virol* 2005;79:3969–3978.
- [32] Li K, Foy E, Ferreon JC, Nakamura M, Ferreon AC, Ikeda M, et al. Immune evasion by hepatitis C virus NS3/4A protease-mediated cleavage of the Toll-like receptor 3 adaptor protein TRIF. *Proc Natl Acad Sci USA* 2005;102:2992–2997.
- [33] Foy E, Li K, Sumpter Jr R, Loo YM, Johnson CL, Wang C, et al. Control of antiviral defenses through hepatitis C virus disruption of retinoic acid-inducible gene-I signaling. *Proc Natl Acad Sci USA* 2005;102:2986–2991.
- [34] Zhang T, Lin RT, Li Y, Douglas SD, Maxcey C, Ho C, et al. Hepatitis C virus inhibits intracellular interferon alpha expression in human hepatic cell lines. *Hepatology* 2005;42:819–827.
- [35] Lee J, Wu CC, Lee KJ, Chuang TH, Katakura K, Liu YT, et al. Activation of anti-hepatitis C virus responses via Toll-like receptor 7. *Proc Natl Acad Sci USA* 2006;103:1828–1833.
- [36] Honda K, Yanai H, Negishi H, Asagiri M, Sato M, Mizutani T, et al. IRF-7 is the master regulator of type-I interferon-dependent immune responses. *Nature* 2005;434:772–777.