

Fig. 3. Regeneration rates of liver weight (A) and PCNA expression rates (B, C) after PH were determined as described in Materials and Methods. No significant differences in the regeneration rates of liver weight and PCNA positive hepatocytes rates (B) were observed between the two groups. However, the PCNA positive SEC rates (C) significantly increased in the HBO group at 48 h ($P < 0.05$), in contrast, it significantly decreased at 72 h after PH ($P < 0.05$). Values are expressed as mean \pm SEM.




Fig. 4. PCNA staining of regenerating liver at 48 h after PH ($\times 400$). (A) Control group. (B) HBO group. H, hepatocyte; S, sinusoidal endothelial cell.




Fig. 5. The hepatic VEGF protein expression after PH by western blot analysis ($n = 4$). HBO treatment significantly increased the VEGF expression at 24 and 48 h after PH, respectively. Values are expressed as mean \pm SEM. * $P < 0.05$ versus control group at the same time point.

liver after PH by immunohistochemical staining. As shown Fig. 6, the majority of the VEGF-positive cells were hepatocytes in the both groups at any time points after PH. The VEGF-positive hepatocytes progressively increased in the periportal and midzonal areas during regeneration.

4. Discussion

In the present study, HBO treatment induces VEGF expression and reduces liver injury during liver regeneration after PH. Although HBO treatment did not significantly affect the hepatic ATP levels, the biliary function was significantly improved in the HBO group. No significant differences were observed in the restitution of the liver weight rates and the PCNA labeling index of hepatocytes, which was peaked at 48 h in the both groups. In contrast, the peak in that of SEC was at 48 h in the HBO group, accelerated compared with that in the control group.

Several studies have confirmed the angiogenic effects of HBO treatment on wound models [21]. Furthermore,




Fig. 6. Immunohistochemical staining for VEGF of regenerating livers at 48 h after PH (x400). (A) Control group. (B) HBO group.

HBO has been shown to induce several angiogenic factors, VEGF concentration in wound fluid [29], bFGF production by fibroblasts [19], and angiopoietin-2 expression in human umbilical vein endothelial cells [30]. A variety of growth factors have been reported to be associated with angiogenesis during liver regeneration [7], and recently, those three angiogenic factors also implicated in SEC proliferation [12,31]. Our data show that HBO treatment does not significantly affect the hepatocytes proliferation, however, the SEC proliferation is accelerated and significantly increases at 48 h after PH, as measured by the PCNA labeling index. To exposure the angiogenic effect of HBO treatment, we examined the expression of those angiogenic factors in the regenerating liver, and found that HBO treatment selectively induced VEGF during liver regeneration after PH and the early induction of VEGF might be a critical step toward HBO-induced SEC proliferation.

VEGF plays an important role in liver regeneration, and the effects of VEGF are mediated through two tyrosine kinase receptors, Flt-1 and KDR/Flk-1 [32,33]. The increased hepatocyte production of VEGF and expression of the receptors correlate with the endothelial proliferation after PH [12,34]. As shown in a study from Shimizu et al., we also found that the expression of Flt-1 and KDR/Flk-1 was observed along sinusoids by immunohistochemical staining, however, HBO treatment did not significantly affect the expression after PH (data not shown). Although hepatocytes do not typically express VEGF receptors, exogenous VEGF administration has been shown to promote the proliferative activity of hepatocytes and stimulate liver regeneration after PH [35]. Taniguchi et al suggests that VEGF has an influence on SEC proliferation after PH and the reconstruction of hepatic vascular architecture appears to promote the proliferation of hepatocytes [36]. Our results show that endogenous VEGF induced by HBO treatment during liver regeneration seems to promote SEC proliferation, however, it cannot affect the hepatocytes proliferation after PH. From the data presented, we suggest that endogenous VEGF increased by HBO treatment did not exert a paracrine effect on hepatocytes proliferation during regeneration.

The HBO treatment after PH significantly reduced liver injury and improved biliary function, as reflected by lower serum ALT, total bilirubin, and total bile acid levels during liver regeneration. It is possible that the beneficial effect of HBO treatment could be due to the increased oxygen delivery to the regenerating liver. During wound healing the presence of oxygen takes on additional importance due to the increased demand of reparative processes like cell proliferation and synthesis of collagen [37]. After PH, single cell wide plates of hepatocytes have grown to avascular clusters [6,7], and hepatocytes that are devoid of sinusoids also seem to be under hypoxic states until reconstruction of normal vascular architecture. In the present study, HBO treatment after PH appears to transiently elevate tissue oxygen levels in the regenerating liver and play a role to improve the hypoxia. Moreover, early upregulated VEGF in the liver also seems to have hepatoprotective effect during regeneration. VEGF has been shown to exert antiapoptotic activities and is highly protective for liver cells [16,38]. Yagi et al. have demonstrated a lower portal venous VEGF level has been associated with poor outcome after living donor liver transplantation [39].

To investigate whether HBO treatment could improve the survival after extended hepatectomy, we performed 95% PH, as a lethal hepatic failure model [40]. However, we found that HBO treatment did not significantly affect the survival rate after 95% PH, and in the both groups, 95% hepatectomized rats all died within 7 days (data not shown). Kamimukai et al. have shown that inhibition of hepatocyte apoptosis by transfection of the human bcl-2 gene did not lead to prolonged survival after 95% PH, and other factors such as a regeneration stimulator might be required to increase the survival rate [41]. Our data demonstrate that HBO treatment on regenerating liver cannot induce hepatocytes proliferation, and therefore, we assume that it did not maintain life after 95% PH.

Distinct from our studies, several studies reported that HBO treatment could affect hepatocytes proliferation during liver regeneration [23,24]. The differences between our findings and previous studies may be

related to the timing of HBO treatment performed on the regenerating liver. Our data have shown that HBO treatment can promote functional recovery after PH, however, further studies are necessary to determine the protocol of HBO treatment which can be most beneficial during liver regeneration.

The question of how HBO treatment influences VEGF expression after PH is not resolved by our data. Hypoxia has been reported to be one of the factors that stimulate VEGF production [42], in contrast, hyperoxia also reported to upregulate the expression [43]. Different oxygen tensions seem to activate different signaling pathways to stimulate VEGF expression for angiogenesis. Hyperoxia enhances reactive oxygen species (ROS) production, which potently induces VEGF expression via a hypoxia-inducible factor-1 independent pathway [44,45]. In addition, during wound healing ROS is required for prevention of wound infection, and specific levels of ROS may act as a signaling mediator that regulates a variety of cellular responses [45,46].

In summary, our studies demonstrate that HBO treatment after PH has beneficial effect on regenerating liver, increasing hepatic VEGF, accelerating SEC proliferation, reducing liver injury, and improving biliary function. Because its safety and simplicity in clinical usage have already been confirmed, HBO treatment may become therapeutic modality in situations in which massive hepatectomy is needed. This may happen in surgery for hepatocellular carcinoma or in living donor liver transplantation.

Acknowledgements

We thank Mizuki Ninomiya for technical help, and we are also indebted to Hiroshi Yagi for supplying the hyperbaric chamber.

References

- [1] Al-Hadeedi S, Chori TK, Wong J. Extended hepatectomy for hepatocellular carcinoma. *Br J Surg* 1990;77:1247–1250.
- [2] Matsumata T, Taketomi A, Kawahara N, Higashi H, Shirabe K, Takenaka K. Morbidity and mortality after hepatic resection in the modern era. *Hepatogastroenterology* 1995;42:456–460.
- [3] Taub R. Liver regeneration: from myth to mechanism. *Nat Rev* 2004;5:836–847.
- [4] Michalopoulos GK, DeFrances MC. Liver regeneration. *Science* 1997;276:60–66.
- [5] Widmann JJ, Fahimi HD. Proliferation of mononuclear phagocytes (Kupffer cells) and endothelial cells in regenerating rat liver. *Am J Pathol* 1975;80:349–366.
- [6] Wack KE, Ross MA, Zegarra V, Sysko LR, Watkins SC, Stolz DB. Sinusoidal ultrastructure evaluated during the revascularization of regenerating rat liver. *Hepatology* 2001;33:363–378.
- [7] Ross MA, Sander CM, Kleeb TB, Watkins SC, Stolz DB. Spatiotemporal expression of angiogenesis growth factor receptors during the revascularization of regenerating rat liver. *Hepatology* 2001;34:1135–1148.
- [8] Drixler TA, Vogten MJ, Ritchie ED, van Vroonhoven TJ, Gebbink MF, Voest EE, et al. Liver regeneration is an angiogenesis associated phenomenon. *Ann Surg* 2002;236:703–711.
- [9] LI J, Zhang YP, Kirsner RS. Angiogenesis in wound repair: angiogenic growth factors and extracellular matrix. *Microsc Res Tech* 2003;60:107–114.
- [10] Arai M, Mochida S, Ohno A, Ogata I, Obama H, Maruyama I, et al. Blood coagulation equilibrium in rat liver microcirculation as evaluated by endothelial cell thrombomodulin and macrophage tissue factor. *Thromb Res* 1995;80:113–123.
- [11] Mochida S, Ishikawa K, Inao M, Shibuya M, Fujiwara K. Increased expression of vascular endothelial growth factor and its receptors, flt-1 and KDR/flk-1, in regenerating rat liver. *Biochem Biophys Res Commun* 1996;226:176–179.
- [12] Sato T, El-Assal ON, Ono T, Yamanoi A, Dhar DK, Nagasue N. Sinusoidal endothelial cell proliferation and expression of angiopoietin/Tie family in regenerating rat liver. *J Hepatol* 2001;34:690–698.
- [13] Yamane A, Seetharam L, Yamaguchi S, Gotoh N, Takahashi T, Neufeld G, et al. A new communication system between hepatocytes and sinusoidal endothelial cells in liver through vascular endothelial growth factor and Flt tyrosine kinase receptor family (Flt-1 and KDR/Flk-1). *Oncogene* 1994;9:2683–2690.
- [14] Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, Dvorak HF. Tumor cells secrete a vascular permeability factor that promotes accumulation of ascitic fluid. *Science* 1983;219:983–985.
- [15] LeCounter J, Moritz DR, Li B, Phillips GL, Liang XH, Gerber HP, et al. Angiogenesis-independent endothelial protection of liver: role of VEGFR-1. *Science* 2003;299:890–893.
- [16] Beraza N, Marques JM, Martinez-Anso E, Iniguez M, Prieto J, Bustos M. Interplay among cardiotrophin-1, prostaglandins, and vascular endothelial growth factor in rat liver regeneration. *Hepatology* 2005;41:460–469.
- [17] Tibble PM, Edelsberg JS. Hyperbaric-oxygen therapy. *N Engl J Med* 1996;334:1642–1648.
- [18] Mazariegos GV, O'Toole K, Miele LA, Dvorchik I, Meza MP, Briassoulis G, et al. Hyperbaric oxygen therapy for hepatic artery thrombosis after liver transplantation in children. *Liver Transpl Surg* 1999;5:429–436.
- [19] Kang TS, Gorti GK, Quan SY, Ho M, Koch RJ, et al. Effect of hyperbaric oxygen on the growth factor profile of fibroblasts. *Arch Facial Plast Surg* 2004;6:31–35.
- [20] Yamashita M, Yamashita M. Hyperbaric oxygen treatment attenuates cytokine induction after massive hemorrhage. *Am J Endocrinol Metab* 2000;278:E811–E816.
- [21] Sheikh AY, Rollins MD, Hopf HW, Hunt TK. Hyperoxia improves microvascular perfusion in a murine wound model. *Wound Rep Reg* 2005;13:303–308.
- [22] Roedel-Wiedmann I, Bennett M, Kranke P. Systematic review of hyperbaric oxygen in the management of chronic wounds. *Br J Surg* 2005;92:24–32.
- [23] Uwagawa T, Unemura Y, Yamazaki Y. Hyperbaric oxygenation after portal vein embolization for regeneration of the predicted remnant liver. *J Surg Res* 2001;100:63–68.
- [24] Nagamine K, Kubota T, Togo S, Nagashima Y, Mori M, Shimada H. Beneficial effect of hyperbaric oxygen therapy on liver regeneration after 90% hepatectomy in rats. *Eur Surg Res* 2004;36:350–356.
- [25] Higgins GM, Anderson RM. Experimental pathology of liver: restoration of liver of the white rat following partial surgical removal. *Arch Pathol* 1934;12:186–202.
- [26] Selzner M, Clavian PA. Failure of regeneration of the steatotic rat liver: disruption at two different levels in the regeneration pathway. *Hepatology* 2000;31:35–42.
- [27] Yamaoka Y, Ohsawa T, Takasan H, Ozawa K. Energy requirement in regenerative and atrophic processes of the liver in man and other mammals. *Surg Gynecol Obstet* 1974;139:234–240.

- [28] Sheikh AY, Gibson JJ, Rollins MD, Hopf HW, Hussain Z, Hunt TK. Effect of hyperoxia on vascular endothelial growth factor levels in a wound model. *Arch Surg* 2000;135:1293–1297.
- [29] Lin S, Shyu KG, Lee CC, Wang BW, Chang CC, Liu YC, et al. Hyperbaric oxygen selectively induces angiopoietin-2 in human umbilical vein endothelial cells. *Biochem Biophys Res Commun* 2002;296:710–715.
- [30] Greene AK, Wiener S, Puder M, Yoshida A, Shi B, Perez-Atayde AR, et al. Endothelial-directed hepatic regeneration after partial hepatectomy. *Ann Surg* 2003;237:530–535.
- [31] De Vries C, Escobedo JA, Ueno H, Houck K, Ferrara N, Williams LT. The fms-like tyrosine kinase, a receptor for vascular endothelial growth factor. *Science* 1992;255:989–991.
- [32] Terman BI, Dougher-Vermazen M, Carrion ME, Dimitrov D, Armellino DC, Gospodarowicz D, et al. Identification of the KDR tyrosine kinase as a receptor for vascular endothelial cell growth factor. *Biochem Biophys Res Commun* 1992;187:1579–1586.
- [33] Shimizu H, Miyazaki M, Wakabayashi Y, Mitsuhashi N, Kato A, Ito H, et al. Vascular endothelial growth factor secreted by replicating hepatocytes induces sinusoidal endothelial cell proliferation during regeneration after partial hepatectomy in rats. *J Hepatol* 2001;34:683–689.
- [34] Assy N, Spira G, Paizi M, Shenkar L, Kraizer Y, Cohen T, et al. Effect of vascular endothelial growth factor on hepatic regenerative activity following partial hepatectomy in rats. *J Hepatol* 1999;30:911–915.
- [35] Taniguchi E, Sakisaka S, Matsuo K, Tanikawa K, Sata M. Expression and role of vascular endothelial growth factor in liver regeneration after partial hepatectomy in rats. *J Histochem Cytochem* 2001;49:121–130.
- [36] Hunt TK, Zederfeldt B, Goldstick TK. Oxygen and healing. *Am J Surg* 1969;118:521–525.
- [37] Morinaga T, Arai S, Takeda Y, Furuyama H, Mizumoto M, Mori A, et al. Protection by vascular endothelial growth factor against sinusoidal endothelial damage and apoptosis induced by cold preservation. *Transplantation* 2000;69:141–147.
- [38] Yagi S, Iida T, Taniguchi K, Hori T, Hamada T, Fujii K, et al. Impact of portal venous pressure on regeneration and graft damage after living-donor liver transplantation. *Liver Transpl* 2005;11:68–75.
- [39] Hasegawa S, Kubota T, Fukuyama N, Kurowawa H, Sekido H, Togo S, et al. Apoptosis of hepatocytes is a main cause of including lethal hepatic failure after excessive hepatectomy in rats. *Transplant Proc* 1999;31:558–559.
- [40] Kamimukai N, Togo S, Hasegawa S, Kubota T, Kurosawa H, Li XK, et al. Expression of bcl-2 family reduces apoptotic hepatocytes after excessive hepatectomy. *Eur Surg Res* 2001;33:8–15.
- [41] Sandner P, Wolf K, Bergmaier U, Gess B, Kurtz A. Induction of VEGF and VEGF receptor gene expression by hypoxia: divergent regulation in vivo and vitro. *Kidney Int* 1997;51:448–453.
- [42] Maniscalco WM, Watkins RH, Finkelstein JN, Campbell MH. Vascular endothelial growth factor mRNA increases in alveolar epithelial cells during recovery from oxygen injury. *Am J Respir Cell Mol Biol* 1995;13:377–386.
- [43] Tandara AA, Mustoe TA. Oxygen in wound healing—more than a nutrient. *World J Surg* 2004;28:294–300.
- [44] Sen CK, Khanna S, Babior BM, Hunt TK, Ellison EC, Roy S. Oxidant-induced vascular endothelial growth factor expression in human keratinocytes and cutaneous wound healing. *J Biol Chem* 2002;277:33284–33290.
- [45] Sen CK, Packer L. Antioxidant and redox regulation of gene transcription. *Fed Am Soc Exp Biol J* 1996;10:709–720.

特集「ウイルス肝炎の肝移植」

生体肝移植後ステロイドフリー免疫抑制法によるC型肝炎再発の制御

吉住朋晴^{1,2}, 武富紹信¹, 副島雄二¹, 内山秀昭¹, 池上 徹¹, 原田 昇¹,
萱島寛人¹, 山下洋市¹, 居村 暁², 森根裕二², 島田光生², 前原喜彦¹¹九州大学大学院消化器・総合外科, ²徳島大学大学院臓器病態外科

はじめに

非免疫抑制下のC型肝炎ウイルス(HCV)陽性患者では、感染後20年で約20%が肝硬変に進展し、そのうちの3%から6%が毎年非代償性肝硬変へ進行するとされている¹⁾。従来、HCV陽性レシピエントにおける肝移植後の患者生存率およびグラフト生着率は、その緩やかな経過の予測の下、HCV陰性例と比較して、遜色ないものとされていた²⁾。しかし、最近、HCV陽性例では、肝移植後早期に肝炎再発から肝硬変、さらに非代償性肝硬変へと進行し、患者生存率およびグラフト生着率が1990年代に比べ著しく低下していることが報告された³⁾。その中で、Berenguerらは近年の過剰な免疫抑制法が、早期の肝硬変への進展と成績の悪化をもたらしている可能性を示唆した。この移植後C型肝炎再発は現在の移植医療従事者が最も頭を悩ませている問題といっても過言ではない。

疫学的に、世界中のHCVキャリアー数は約1億7,500万人で、そのうちわが国におけるHCVキャリアー数は150万から200万人、約4,500人が毎年新規感染していると予想されている。つまり、HCVに起因する肝硬変は元来肝移植の対象として、頻度が高いと考えられる疾患であり、日本肝移植研究会の集計では、2003年までに施行された全成人間生体肝移植数に占めるHCV陽性症例の割合は、22%であった⁴⁾。さらに2004年1月から、ウイルス性肝硬変に対する生体肝移植が保険適応として認められ、HCV陽性レシピエントに対する生体肝移植施行症例数は今後ますます増加するものと思われる。

生体肝移植に用いられるグラフトは部分肝であり、

術後の経過でグラフトの再生が起こる。またわが国において、生体ドナーはレシピエントの近親者に限定されているため、ドナーとレシピエントのHLAがより近似している可能性がある。これらのため、生体肝移植術後のC型肝炎再発は、脳死肝移植に比べ、早い時期に起こり予後不良であるとする報告もあるが、脳死肝移植との間に差はないとする報告も多く、いまだ結論は出ていない。

本稿では、HCV陽性例に対する免疫抑制法の最近の報告をレビューするとともに、当施設におけるステロイドフリー免疫抑制法の経験について、概説する。

免疫抑制剤のHCVに与える影響

1. カルシニューリン阻害剤

HCVのレプリコン細胞を用いた実験において、シクロスポリンには濃度依存性に、シクロフィリンBを介するHCV増殖抑制効果があることが報告された^{5,6)}。臨床例では、シクロスポリン使用によりC型肝炎再発後の線維化が緩徐であるとするものや⁷⁾、肝移植後の再発例で、シクロスポリンはインターフェロンとの併用により抗ウイルス療法に対する相乗効果をもつことが報告された⁸⁾。しかし一方で、移植後5年生存率は、タクロリムス使用群がシクロスポリン使用群より有意に良好であったとする報告がみられ⁹⁾、最近米国でのprospective randomized trialにより、移植後のHCV RNA量やC型肝炎再発率はカルシニューリン阻害剤の種類で差を認めなかったことが報告された¹⁰⁾。

これら相反する報告から、どちらかのカルシニュー

リン阻害剤が有意に優れているとは、現時点では言えないと考えるのが妥当であろう。

2. ステロイド

Ganeらはメチルプレドニゾロンのbolus投与により、HCV RNA量は4~100倍に増加し、HCV RNA量の高値は、肝移植後C型肝炎の早期再発および再発肝炎の重症度と相関することを報告した¹¹⁾。メチルプレドニゾロンのbolus投与に続くHCV量の著増はfibrosing cholestatic hepatitis (FCH)とも関連するとされている¹²⁾。FCHは肝移植後5-10%の症例に起こる、移植約1カ月後の肝機能異常から始まり半年以内に肝不全に進行する特殊なC型肝炎再発パターンである¹³⁾。FCHはインターフェロン治療からの脱落が多く、高い死亡率のため、避けねばならぬ病態である。これらの報告による過剰なステロイドへの拒絶感のためか、rapid steroid taperingのプロトコールを多くの施設が取り入れた。しかし最近、プレドニゾロンを少量長期使用後、移植後6カ月以降に緩徐にtaperするプロトコールで重症再発が予防可能あるいは線維化進展が抑制可能との報告がされた^{14,15)}。一方、ステロイドフリーの免疫抑制に関する報告もみられ^{16,17)}、HCVの増殖速度を抑える¹⁰⁾あるいはグラフト生着率および患者生存率を改善する¹⁷⁾とされているが、C型肝炎再発自体への影響に関しては、いまだ有意な報告はされていない。

3. Mycophenolate Mofetil (MMF)

de novo系の核酸合成阻害剤である本薬は、EBウイルスや黄熱病ウイルスの複製阻害効果、抗ヘルペス薬の効果増強作用、in vitroでのHBV複製阻害効果などが知られており、HCVへの抗ウイルス作用の有無が注目されていたが¹⁸⁾、HCV陽性症例に対する効果に関しては議論が分かれている。HCV陽性症例に対して、タクロリムスとステロイドにMMFを追加すると、C型肝炎の再発には影響を及ぼさず、急性拒絶反応の頻度は低下し、さらに移植後4年生存率およびグラフト生着率は有意に良好であった¹⁹⁾、術後のALTは有意に低下し、肝炎や線維化の進行を認めなかった²⁰⁾というMMFの有用性を示唆する報告がみられる。一方、MMFを用いた免疫抑制導入は移植後2年以内の肝線維化の進行を早める危険因子である²¹⁾、あるいはC型肝炎再発例に対してazathioprineに代えてMMFを使用するとHCV RNA量が増加する¹⁸⁾といっ

たnegativeな報告もみられ、その効果に関する結論は出ていない。

4. Basiliximab

本薬はIL-2レセプターの α 鎖に結合し、Tリンパ球の増殖を抑制するモノクローナル抗体である。術中および術後4日目に使用することで、術後約30日にわたり、CD25陽性活性化T細胞の出現を抑制し、急性拒絶反応の発生率を低下させる。CalmusらはHCV陽性例に使用し肝移植6カ月後の肝炎再発率は48.4%で他の免疫抑制法と差を認めなかったと報告した²²⁾。その後既出のごとく、Basiliximabとの併用でステロイドフリー免疫抑制法が可能であることが報告された^{16,17)}。わが国からもMarubashiらが、生体肝移植症例においてもBasiliximabをタクロリムスおよびMMFと併用することで、急性拒絶反応を発生させることなく、ステロイドフリーの免疫抑制が可能であり、術後早期のHCV RNA量を抑制したと報告した²³⁾。

■ HCV陽性例における

■ 生体肝移植と脳死肝移植の相違

慢性的な臓器不足を背景に、欧米においても生体肝移植を施行する施設が増加している。HCV陽性症例に対する生体肝移植の長所として、待機期間が調整可能であるため移植前の抗ウイルス治療が可能であること、短い冷保存時間、比較的若いドナー、重度の脂肪肝のない良好なグラフト機能などが挙げられる。一方、欠点としては親族からの提供の場合にドナーとレシピエント間のHLAの相同性が高いこと、肝細胞の再生のためにHCVの増殖を促進する可能性があること、薬剤代謝の違いから免疫抑制剤が強く作用する可能性があること、高率な胆道系の合併症から持続する胆汁うっ滞を招くこと、などが挙げられる²⁴⁾。GaglioらのHCV陽性例における生体肝移植と脳死肝移植を比較した最初の報告では、生体肝移植後のC型肝炎再発は脳死肝移植に比べ早期に起こり、FCHなどのより激しい再発形式をとるとされた²⁵⁾。Schianoらは、生体ドナーでは脳死ドナーに比べ、再灌流直後の血中HCV RNAは速やかに低下するが、その後の増殖速度も速く、また移植後2年間のALTが有意に高値であったと報告した²⁶⁾。最近の報告では両者間の生存率・グラフト生着率あるいは組織学的再発率に差がないとするものも多いが、いずれも症例数が限られており、さらに各施設間におけるC型肝炎再発の定義の違いか

らか生体ドナーの善悪に関する結論は出ていない²⁴⁾。HCV陽性例に対する多数の生体肝移植症例数を持つが国から、多施設共同研究を行い、積極的に情報を発信していく必要があると思われる。

当施設における生体肝移植後ステロイドフリー免疫抑制法

九州大学第二外科で、2006年3月までに施行した成人間生体肝移植186例中、HCV陽性症例は73例(39.2%)であった。特に2002年以降は全成人症例のうち、約半数をHCV陽性例が占めている。図1に当科におけるHCV陽性症例に対する免疫抑制法の変遷を示す。当初はタクロリムスとステロイドを用いていたが、その後MMFおよびBasiliximabの使用、またステロイドの短期中止などを取り入れ、現在では、図1の3)に示す如く、BasiliximabとMMFにより導入し、カルシニューリン阻害剤を術後7日目より開始している。カルシニューリン阻害剤のトラフ値はタクロリムスでは10-15 ng/ml、サイクロスポリンでは200-250 ng/mlでコントロールしている。ステロイドは原則的に用いていない。自験例を免疫抑制導入法により3群に分け、3群間における術後HCV RNA量、急性

表1 各免疫抑制導入法の背景因子

因子	ステロイドフリー群 (n=14)	ステロイド短期/一回投与群 (n=33)	従来群 (n=26)	P値
Recipient 年齢	56.4	58.5	51.8	<0.05
Recipient 性別 (M/F)	10/4	20/13	18/8	N.S.
Donor 年齢	35.0	35.2	33.4	N.S.
Donor 性別 (M/F)	5/9	26/7	18/8	<0.05
術前HCV RNA (KU/ml)	554.6	420.3	162.8	<0.005
Virus Grouping 1型/2型	11/0	24/4	13/4	N.S.
Graft種類 (左葉/右葉)	7/7	22/11	17/9	N.S.
GV/SLV (%)	41.0	42.8	42.2	N.S.
術前MELD	11.0	12.3	13.9	N.S.
出血量	6240	5792	12352	0.13
手術時間(分)	852	788	893	0.10

拒絶反応の発生率、肝炎再発率、線維化進展率および生存率を検討した。表1に各群の背景因子を示す。ステロイドフリー群でレシピエント年齢が高く、ドナーは女性が多かった。また、術前HCV RNA量が有意に多かった。

急性拒絶反応の発生率はステロイドフリー群で14.3%と低い傾向にあった。このため、急性拒絶反応の治療として、ステロイドのbolus投与を要した症例も、14.3%に留まった(図2)。HCV RNA量の測定法が同一の2004年1月以降の症例で、術後1カ月のHCV RNA量を比較したところ、ステロイドフリー群

図1 九州大学におけるHCV陽性例に対する免疫抑制導入プロトコール




図2 各免疫抑制導入法の急性拒絶反応発生率と拒絶反応の治療にステロイド bolus 投与を施行した頻度




図3 各免疫抑制導入法と移植1カ月後のHCV RNA量





図4 術後HCV RNA量の推移



でHCV RNA量は低い傾向にあった(図3)。

次にステロイドフリー群とステロイド短期/一回投与群の術後6カ月間のHCV RNAの推移を示す(図4)。ステロイドフリー群に対して、短期/一回投与群では、術後早期から急速なHCV RNA量の増加を示し、測定上限である5,000 KIU/mlに達した。これは、rapid steroid taperingがHCVのウイルス動態に悪影響を及ぼすとの報告に矛盾しないと考えられる。ステロイド短期/一回投与群のうち2例では、FCH様の経過を呈し、そのうち1例は急速な肝不全の進行により術後7カ月で失った¹³⁾。McCaughanらが報告しているように術直後の急速なウイルスの増加は、FCHの発生

と関連するものと考えられた¹²⁾。ステロイドフリー群においても、多くの症例が術後2~3カ月で測定上限に達していたが、術直後の急速な増殖を避け得たために、FCHを呈した症例は経験しなかった。ウイルスの増殖が上限に達する前に予防的なインターフェロン治療を開始し、ウイルスの陰性化を目指すことが術後管理を行う上で重要と考えられる²⁷⁾。

当施設における移植後の組織学的累積再発率は術後6カ月で48.1%、5年では88.8%(図5)、F3以上の線維化進展率は、1年で13.2%、5年で65.9%であっ

図5 移植後C型肝炎の組織学的再発率




図6 移植後F3以上の肝線維化進展率




表2

組織学的線維化(F3, F4)進展の危険因子(多変量解析)

因子	ハザード比 (95% CI)	p 値
Recipient 女性 vs. 男性	3.81 (1.3-11.2)	0.015
Donor 年齢 ≥40才 vs. <40才	2.92 (1.0-8.1)	0.040

た(図6)。予防的治療を開始しない場合の再発はほぼ必発であるため、線維化の進展をいかに防ぐかが、現在の術後治療のポイントと考えられる。組織学的再発率および線維化進展率を各免疫抑制導入法で比較したが、有意な差は認めなかった。多変量解析による再発の危険因子はレシピエントの性別とドナーの年齢40歳以上のみであった。免疫抑制導入法あるいは免疫抑制剤の種類は危険因子とはならなかった(表2)。

まとめ

2004年1月の保険適応改正により、HCV陽性例は、今後しばらくはわが国における生体肝移植の最大の適応であり続けると考えられる。HCV陽性例の移植術後管理の困難さは、急性拒絶反応とC型肝炎再発の鑑別が組織像をもってしても、非常に難しいことにある。わが国において症例が集積される以前は、試行錯誤の中で術後管理が行われていたのが現状であった。自験例での検討から、われわれが現在行っているステロイドフリー免疫抑制法により、組織学的再発あるいは線維化進展の予防は有意には改善しなかった。しかし、ステロイドフリー免疫抑制導入法の最大の長所は、急性拒絶反応の頻度は低いままで、術直後におけるHCV RNA量の急速な増殖を抑制することが可能な点である。これにより、予後不良なFCHを起こすことなく、より早期に安定した状態で、術後の予防的インターフェロン治療を開始することが可能となった。現在のわれわれのプロトコルで術後管理を行う上で、急性拒絶とC型肝炎再発の鑑別に悩むことはほぼ皆無となった。

1980年代まで移植後の成績が不良であったB型肝炎では、HBIG・ラミブジンの使用により、背景に存在する肝炎までも肝移植により同時に根治させることが可能となり、現在では肝移植の最も良い適応とさえ

考えられる。現在、HCVが感染した細胞内におけるHCVの翻訳あるいはHCV RNAの複製を制御すると考えられる種々のHCVに対する治療薬が開発されているが²⁸⁾、HCVのグラフト肝への再感染は、再灌流時から始まる^{12,26)}。肝移植の分野から言えば、B型肝炎におけるHBIGの働き、つまり肝細胞へのHCVのbindingそのものを予防しうるモノクローナル抗体あるいは免疫グロブリンなどの開発が、より効率の良い移植後再感染およびC型肝炎再発予防を可能にするであろう²⁹⁾。最近、memory T細胞を利用したHCVワクチンのHCV感染に対する有効性が報告された³⁰⁾。肝移植術後の症例では、急性拒絶反応とのバランスという問題もあるが、術後のT細胞の特異的な制御が、今後の治療の主役となる可能性もある。いずれにしろ、HCVの分子構造解明に基づく治療法の早急な開発が望まれる。

最後に、HCV陽性例に対する移植後の管理に関しては、いまだコンセンサスが得られていないのが現状である。われわれが積極的に取り組んでいるステロイドフリー免疫抑制法は、そのものがHCVの治療法ではないが、安定した状態で早期に抗ウイルス治療を開始するための周術期管理法としては、大変有効な手段である。現在、厚生労働省班研究による症例の集積が進んでおり、今後さらに新たな知見が得られるものと考えられる。

文献

- 1) Fattovich G, Giustina G, Degos F, *et al.* Morbidity and mortality in compensated cirrhosis type C: a retrospective follow-up study of 384 patients. *Gastroenterology* 1997; 112: 463-472.
- 2) Gane EJ, Portmann BC, Naoumov NV, *et al.* Long-term outcome of hepatitis C infection after liver transplantation. *N Engl J Med* 1996; 334: 815-820.
- 3) Berenguer M, Prieto M, San Juan F, *et al.* Contribution of donor age to the recent decrease in patient survival among HCV-infected liver transplant recipients. *Hepatology* 2002; 36: 202-210.
- 4) 日本肝移植研究会・肝移植症例登録報告. 移植 2004; 39: 634-642.
- 5) Watashi K, Hijikata M, Hosaka M, *et al.* Cyclosporin A suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* 2003; 38: 1282-

- 1288.
- 6) Watashi K, Ishii N, Hijikata M, *et al.* Cyclophilin B is a functional regulator of hepatitis C virus RNA polymerase. *Mol Cell.* 2005; 19: 111-122.
 - 7) Pollard S. Calcineurin inhibition and disease recurrence in the hepatitis C virus-positive liver transplant recipient. *Liver Int.* 2004; 24: 402-406.
 - 8) Firpi RJ, Zhu H, Morelli G, *et al.* Cyclosporine suppresses hepatitis C virus in vitro and increases the chance of a sustained virological response after liver transplantation. *Liver Transpl* 2006; 12: 51-57.
 - 9) Wiesner RH. A long-term comparison of tacrolimus (FK 506) versus cyclosporine in liver transplantation: a report of the United States FK 506 Study Group. *Transplantation* 1998; 66: 493-499.
 - 10) Martin P, Busuttil RW, Goldstein RM, *et al.* Impact of tacrolimus versus cyclosporine in hepatitis C virus-infected liver transplant recipients on recurrent hepatitis: a prospective, randomized trial. *Liver Transpl* 2004; 10: 1258-1262.
 - 11) Gane EJ, Naoumov NV, Qian KP, *et al.* A longitudinal analysis of hepatitis C virus replication following liver transplantation. *Gastroenterology* 1996; 110: 167-177.
 - 12) McCaughan GW, Zekry A. Mechanisms of HCV reinfection and allograft damage after liver transplantation. *J Hepatol* 2004; 40: 368-374.
 - 13) 米村祐輔, 島田光生, 吉住朋晴, 他. C型肝硬変に対する生体肝移植後に急速な経過をたどり死亡した fibrosing cholestatic hepatitis の1例. *日本臨床外科学会雑誌* 2004; 65: 3257-3262.
 - 14) Samonakis DN, Triantos CK, Thalheimer U, *et al.* Immunosuppression and donor age with respect to severity of HCV recurrence after liver transplantation. *Liver Transpl* 2005; 11: 386-395.
 - 15) Berenguer M, Aguilera V, Prieto M, *et al.* Significant improvement in the outcome of HCV-infected transplant recipients by avoiding rapid steroid tapering and potent induction immunosuppression. *J Hepatol* 2006; 44: 717-722.
 - 16) Garcia-Retortillo M, Forns X, Feliu A, *et al.* Hepatitis C virus kinetics during and immediately after liver transplantation. *Hepatology* 2002; 35: 680-687.
 - 17) Filippini F, Callea F, Salizzoni M, *et al.* Double-blind comparison of hepatitis C histological recurrence Rate in HCV+Liver transplant recipients given basiliximab+steroids or basiliximab+placebo, in addition to cyclosporine and azathioprine. *Transplantation* 2004; 78: 1488-1495.
 - 18) Zekry A, Gleeson M, Guney S, *et al.* A prospective cross-over study comparing the effect of mycophenolate versus azathioprine on allograft function and viral load in liver transplant recipients with recurrent chronic HCV infection. *Liver Transpl* 2004; 10: 52-57.
 - 19) Wiesner RH, Shorr JS, Steffen BJ, *et al.* Mycophenolate mofetil combination therapy improves long-term outcomes after liver transplantation in patients with and without hepatitis C. *Liver Transpl* 2005; 11: 750-759.
 - 20) Bahra M, Neumann UI, Jacob D, *et al.* MMF and calcineurin taper in recurrent hepatitis C after liver transplantation: impact on histological course. *Am J Transplant* 2005; 5: 406-411.
 - 21) Berenguer M, Crippin J, Gish R, *et al.* A model to predict severe HCV-related disease following liver transplantation. *Hepatology* 2003; 38: 34-41.
 - 22) Calmus Y, Scheele JR, Gonzalez-Pinto I *et al.* Immunoprophylaxis with basiliximab, a chimeric anti-interleukin-2 receptor monoclonal antibody, in combination with azathioprine-containing triple therapy in liver transplant recipients. *Liver Transpl* 2002; 8: 123-131.
 - 23) Marubashi S, Dono K, Amano K, *et al.* Steroid-free living-donor liver transplantation in adults. *Transplantation* 2005; 80: 704-706.
 - 24) Berenguer M. Live donor liver transplantation for hepatitis C: new data, old story. *Liver Transpl* 2006; 12: 516-519.
 - 25) Gaglio PJ, Malireddy S, Levitt BS, *et al.* Increased risk of cholestatic hepatitis C in recipients of grafts from living versus cadaveric liver donors. *Liver Transpl* 2003; 9: 1028-1035.
 - 26) Schiano TD, Gutierrez JA, Walewski JL, *et al.* Accelerated hepatitis C virus kinetics but similar survival rates in recipients of liver grafts from living versus deceased donors. *Hepatology* 2005; 42: 1420-1428.

- 27) Sugawara Y, Makuuchi M, Matsui Y, *et al.* Preemptive therapy for hepatitis C virus after living-donor liver transplantation. *Transplantation* 2004; 78: 1308-1311.
- 28) Sakamoto H, Okamoto K, Aoki M, *et al.* Host sphingolipid biosynthesis as a target for hepatitis C virus therapy. *Nat Chem Biol* 2005; 1: 333-337.
- 29) Eren R, Landstein D, Terkieltaub D, *et al.* Preclinical evaluation of two neutralizing human monoclonal antibodies against hepatitis C virus (HCV): a potential treatment to prevent HCV reinfection in liver transplant patients. *J Virol*. 2006; 80: 2654-2664.
- 30) Folgosi A, Capone S, Ruggeri L, *et al.* A T-cell HCV vaccine eliciting effective immunity against heterologous virus challenge in chimpanzees. *Nat Med* 2006; 12: 190-197.

Living donor liver transplantation to patients with hepatitis C virus cirrhosis

Yasuhiko Sugawara, Masatoshi Makuuchi

Yasuhiko Sugawara, Masatoshi Makuuchi, Artificial Organ and Transplantation Division, Department of Surgery, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

Supported by Grant-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan

Correspondence to: Yasuhiko Sugawara, MD, Artificial Organ and Transplantation Division, Department of Surgery, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. yasusuga-ky@umin.ac.jp

Telephone: +81-3-38155411 Fax: +81-3-56843989

Received: 2005-09-12 Accepted: 2005-11-18

Abstract

Living donor liver transplantation (LDLT) is an alternative therapeutic option for patients with end-stage hepatitis C virus (HCV) cirrhosis because of the cadaveric organ shortage. HCV infection is now a leading indication for LDLT among adults worldwide, and there is a worse prognosis with HCV recurrence. The antiviral strategy after transplantation, however, is currently under debate. Recent updates on the clinical and therapeutic aspects of living donor liver transplantation for HCV are discussed in the present review.

© 2006 The WJG Press. All rights reserved.

Key words: Hepatitis C virus; Living donor liver transplantation; Interferon; Rivabirin

Sugawara Y, Makuuchi M. Living donor liver transplantation to patients with hepatitis C virus cirrhosis. *World J Gastroenterol* 2006; 12(28): 4461-4465

<http://www.wjgnet.com/1007-9327/12/4461.asp>

INTRODUCTION

The use of live donors for liver transplantation was initiated more than a decade ago as a solution to the cadaveric donor shortage for pediatric recipients^[1]. After the first successful case in an adult patient in 1994^[2], this procedure is now widely applied to adult recipients, especially in countries where the availability of brain-dead donors is severely restricted^[3] and also in the United States and European countries, due to a critical shortage

of cadaveric organs. Improved surgical techniques and the introduction of new immunosuppressive agents have enhanced the long-term results of living donor liver transplantation (LDLT), leading to an increased demand for liver transplantation that exceeds the number of potential donor organs. In initial experiences with adult LDLT in Japan, the most common indication was cholestatic liver disease, including primary biliary cirrhosis and primary sclerosing cholangitis in Japan. The number of LDLT patients indicated for hepatitis C virus (HCV) has recently increased rapidly.

A recent study^[4] of deceased donor liver transplantation (DDLT) reported that HCV infection was associated with a 23% increase in mortality and a 30% increase in the rate of graft failure. The poor results might be due to the recurrence of HCV disease in the graft^[5]. HCV-induced graft hepatitis and fibrosis/cirrhosis occur in 75% to 80% and 10% to 30% of recipients, respectively, at 5 years^[6,7]. Once liver cirrhosis is established, the cumulative probability of developing clinical decompensation is close to 50% after 1 year and survival after decompensation is extremely short^[8]. Cholestatic hepatitis occurs in approximately 10% of patients infected with HCV and leads to accelerated graft failure and death^[9]. One of the hottest debates is the possibility of increased severity of recurrent HCV in LDLT patients. The benefit of LDLT might be offset if the outcome of LDLT for HCV patients is worse than that of DDLT. In this review, we describe current trends and controversies in LDLT for patients with HCV. Our results for LDLT and HCV are also reported.

CURRENT STATUS OF LDLT

According to the Japan Liver Transplantation Society^[10], the number of adult patients (≥ 18 years old) is increasing annually, and has reached 300 in 2003. The most common indication for adults has been hepatocellular carcinoma ($n = 311$), followed by primary biliary cirrhosis ($n = 255$), and HCV-related cirrhosis without carcinoma ($n = 113$). The 1, 3, and 5 year survival rates of all the adult patients were 76%, 72%, and 69%, respectively. Those of HCV-positive patients were 76%, 73%, and 65%, respectively.

In the United States in 2000, there was a high level of enthusiasm for adult LDLT, with 49 centers performing at least one LDLT. Overall, in experienced centers, about a third of adults on the waiting list had a potential living donor and half of them had undergone LDLT; thus, LDLT might be applicable for up to 15% of individuals on the list^[11]. The enthusiasm was, however, quickly tempered

Table 1 Comparison between LDLT and DDLT for new HCV-IT[†] patients

Study	Author	Year	Institution	N		Dif ¹	Protocol biopsy	Findings
				LDLT	DDLT			
Gaglio ^[23]		2003	Colombia U.	23	45	Yes	No	Cholestatic hepatitis in 17% of LDLT and 0% of DDLT ($P = 0.001$). No significant difference in incidence of Rec.
Shiffman ^[26]		2003	Virginia Commonwealth U.	22	53	No	Yes	79% patient survival in LDLT and 91% in DDLT during 3 year (NS). No significant difference in inflammation score in liver specimen after 3 years
Russo ^[29]		2004	UNOS data	279	3955	No	No	87% 1-year patient survival in both.
Thuluvath ^[30]		2004	UNOS data	207	408	No	No	No significant difference in patient survival ($P = 0.6$).
Van Vlierberghe ^[32]		2004	Ghent U.	17	26	No	No	Rec in 35% of LDLT and 38% of DDLT during 1 year ($P = 0.1$)
Bozorgzadeh ^[34]		2004	Rochester U.	35	65	No	No	Rec in 77% of LDLT and 72% of DDLT during 1 year (NS), 89% patient survival in LDLT and 75% in DDLT during 39 mo (NS)

¹Difference in short-term outcomes or severity of virus recurrence between living and deceased donor liver transplantation. Abbreviations: Rec, Virus recurrence; U, University; NS, not significant; UNOS, United Network for Organ Sharing.

by the death of a donor in 2002 in the United States^[12]. Since 2001, the number of patients who have undergone LDLT has declined^[13]. Currently less than 5% of all adult liver recipients use living donors. By July 2005, 2734 LDLT cases had been performed. There were 1761 adult patients and HCV was the most common indication. HCV is the most common indication for LDLT^[14] and the number of HCV-positive patients is stable, approximately 100 per year between 2000 and 2002.

By the end of 2003, 1743 LDLT cases were recorded in the European Liver Transplantation Registry^[15]. According to the Transplant Procurement Management^[16], the number of LDLT peaked in 2003 and has gradually decreased over recent years. LDLT accounts for approximately 5% of the total liver transplants performed in Europe. Among the 806 LDLT cases from October 1991 to December 2001^[17], the overall 5-year graft survival rate was 75%, better for children than for adults (80% vs 66% at 3 years). Cirrhosis secondary to HCV infection is a leading indication for LDLT among adults in Europe^[18]. The number of LDLT patients is shown in the Table 1.

INDICATIONS

In areas with low deceased donor organ availability, the indications for LDLT are similar to those for DDLT. In contrast, in Western countries, LDLT is conducted in an attempt to alleviate the shortage of donor organs and to decrease the mortality among the patients awaiting transplants. That is, a balance needs to be achieved between the candidate's liver disease severity and the adequacy of a partial graft for transplantation. The candidate's liver disease should be advanced to the extent that transplantation is justified, but the liver disease cannot be so advanced that a partial graft will not provide adequate hepatic mass.

According to Russo's report^[19] a substantial proportion of patients were United Network for Organ Sharing (UNOS) status 3 at the time of LDLT (43%). The policy at their centers prior to the implementation of a model for end stage liver disease (MELD)-based allocation was not to proceed with LDLT in patients meeting UNOS status 2A criteria. Their patient survival rate was 57% with an average stay of 23 d in the intensive care unit. In

comparison, 1-year patient survival was 82% in DDLT recipients who were UNOS status 2A at the time of transplant^[20].

The waiting list mortality increases in patients with advanced liver disease and patients with a MELD score of 25 have a 20% 3-mo mortality^[21]. In general, it is uncommon to proceed with LDLT in patients with MELD scores above 25. Thus, depending on the region of the country and the average MELD score at the time of the transplant within the area served by the organ procurement organization, LDLT might offer patients transplantation before they die waiting for a deceased donor liver. The lower MELD score limit with LDLT is more controversial and varies from center to center. Russo^[19] commented that they do not proceed with LDLT in candidates with MELD scores under 11.

LDLT AS A RISK FACTOR FOR RECURRENCE OF HCV

One study from Barcelona^[22] reported that LDLT patients ($n = 22$) had younger donors, less graft steatosis, more frequent biliary complications, and earlier and more severe acute hepatitis compared with DDLT ($n = 95$) patients. A report from Colombia University^[23] indicates that cholestatic hepatitis or severe HCV recurrence occurs more frequently in LDLT. These reports indicate that more intensive antiviral therapy might be necessary for recipients of living donor grafts.

The possible causes of HCV recurrence include HLA matching between donor and recipient. Because cellular immune reactions restricted by both HLA class I and II antigens are involved in the recognition of HCV peptides^[24], HLA matching between donor and recipient could potentially increase damage to the graft from recurrent viral infections by facilitating host recognition of viral antigens^[5]. Recently, a beneficial effect of a complete HLA-DQ mismatch was reported in 14 patients after transplantation for HCV cirrhosis^[25]. Another possible cause might be related to liver regeneration^[26], although recent data^[27] did not support this hypothesis. *In vitro*, HCV internal ribosome entry site activity and replication are higher in actively dividing cells, and it is possible that

viral translation is enhanced by factors that stimulate the regeneration of hepatocytes. Moreover, there are experimental data suggesting that liver regeneration induces low density lipoprotein receptor expression, which might facilitate HCV entrance into the hepatocytes.

In contrast, comparable data between LDLT and DDLT for HCV was recently reported^[28]. Russo and colleagues^[29] compared patient and graft survival in recipients transplanted for chronic HCV who received a living donor organ ($n = 279$) and deceased donor organ ($n = 3955$) using the UNOS liver transplant database. One-year patient survival was 87% in both groups and 2-year patient survival was 83% and 81% in the living donor group and deceased donor group ($P = 0.68$), respectively. Similar results (DDLT, $n = 480$ vs LDLT, $n = 207$) were obtained from another analysis using the UNOS data base^[30]. Analyses from the Mayo Clinic^[31] and Gent University^[32] also demonstrated no negative impacts of LDLT on the results of liver transplantation for HCV-related cirrhosis.

These data should be interpreted with caution, however, because of the important clinical distinction between LDLT and DDLT recipients. At the time of transplantation, the LDLT group recipients are far less sick than their DDLT group counterparts^[33]. The LDLT ($n = 35$) and DDLT ($n = 65$) data from a single institution, Rochester University, were examined^[34]. Patient survival, graft survival, rate of HCV recurrence, severity of HCV recurrence, graft loss from HCV, and interval for HCV recurrence in DDLT and LDLT were similar. It remains unclear, however, whether LDLT is truly disadvantageous compared to DDLT for HCV-positive patients because the number of cases or follow-up duration is not yet sufficient.

According to the data from Russo^[29], from 1999 to 2000, the 1-year patient survival in the LDLT group increased from 69% to 90% ($P = 0.04$), and 1-year graft survival increased from 63% to 79% ($P = 0.16$). In contrast, in the DDLT group, 1-year patient and graft survival did not substantially change from 1999 to 2000. As a result, 1-year survival rates became similar between the LDLT and DDLT groups in 2000. The results indicated an experience effect and learning curve on outcomes after LDLT for HCV. Therefore, the initial reports indicating poorer results of LDLT might be due to technical problems from a lack of experience. Recent data indicating similar results between LDLT and DDLT might be due to the increased experience with LDLT. The multicenter adult to adult LDLT cohort study (A2ALL) might soon provide some answers to the questions about recurrent HCV after LDLT and DDLT^[35].

MANAGEMENT OF HCV

Therapy for recurrence in DDLT

If HCV recurs earlier and more severely after LDLT, a specific strategy for preventing the detrimental effects of HCV on living donor grafts must be developed. One strategy might be aggressive treatment for HCV. Treatment of recurrent HCV disease with interferon and ribavirin after DDLT is used in some centers^[36-38]. One standard regimen includes interferon-alpha2b (3 MU \times 3

per week) and ribavirin (1000 mg/d) for 6 mo. In a recent trial, polyethylene glycol-conjugated interferon therapy was used^[35,39-44], with a sustained viral response rate ranging from 13% to 47%.

Preemptive therapy for HCV after DDLT

Preemptive therapy in the early post-transplantation period with interferon either alone or in combination with ribavirin has been attempted in DDLT, although its effectiveness is controversial. In one study, HCV-positive recipients were randomized within 2 wk of transplantation to receive either interferon alone (3 MU \times 3 per wk, $n = 30$) or placebo ($n = 41$) for 1 year^[39]. Only 17 patients could complete 1 year of interferon therapy. Eight patients (27%) in the interferon group and 22 (54%) of the untreated patients had recurrent hepatitis ($P = 0.02$). Patient and graft survival at 2 years did not differ between the groups, however, and the rate of viral persistence was not affected by treatment.

In another controlled trial^[45], 24 recipients were randomized at 2 weeks post-transplantation to receive interferon (3 MU \times 3 per wk) or placebo for 6 mo. There were no differences in graft or patient survival. There were no differences between groups in the incidence of histological recurrence or its severity differed between groups. Recurrent HCV was delayed 408 d in treated patients versus 193 d in the control cohort.

In a case series by Mazzaferro^[46], 36 recipients were treated with interferon-alpha 2b (3 MU \times 3 per wk) and ribavirin (10 mg/kg per d). They started treatment at a median of 18 d after the operation and treatment continued for 11 mo. After a median follow-up of 52 mo, the 5-year patient survival was 88%. Serum HCV RNA clearance was obtained in 12 patients (33%). They did not require further antiviral treatment because of negative HCV RNA in serum and normal liver histology for a median of an additional 36 mo. The former two randomized trials on preemptive interferon monotherapy demonstrated minimal benefits of the drug. In contrast, Mazzaferro reported more encouraging results, although their protocol brings into question how long therapy is needed once embarking on a preemptive strategy.

Re-transplantation

The approach to retransplantation for recurrent HCV varies widely among the transplant centers of DDLT^[11]. The results after retransplantation for HCV (45% at 5 years) are poorer than that for other causes^[47] (56%, $P < 0.001$). The patients with recurrent HCV in the early timing and graft failure within the first year have poor outcomes after retransplantation. These individuals should be considered contraindicated for retransplantation. The experience of retransplantation for HCV in LDLT has not been well accumulated.

OUR EXPERIENCE

We performed preemptive therapy for LDLT patients with HCV infection^[40]. From 1996 to 2004, 67 patients underwent LDLT for HCV cirrhosis at the Tokyo University Hospital. The patients were 51 men and 16

women and their ages ranged from 23 to 63 years (median 55). The HCV genotype was 1b in 53 patients (79%). Forty-one patients (61%) had hepatocellular carcinoma. All the patients received the same immunosuppressive regimens with tacrolimus and methylprednisolone.

All the patients preemptively received antiviral therapy consisting of interferon α -2b and ribavirin, which was started approximately 1 mo after the operation. The therapy was continued for 12 mo after the first negative HCV RNA test. The standard regimen included interferon α -2b (3 MU \times 3 per wk) and ribavirin (800 mg/d) for 6 mo. The patients were then observed without the therapy for 6 mo. The therapy was continued for at least 12 mo even if the HCV RNA test remained positive.

Therapy was discontinued when there was significant leukopenia (< 1500 /mL), thrombocytopenia (< 50000 /mL) despite application of granulocyte colony stimulating factor (Gran[®], Sankyo, Co. Ltd., Tokyo, Japan), hemolytic anemia (hemoglobin < 8 g/L), renal dysfunction (serum creatinine > 20 mg/L), depressive psychological status, or general fatigue. The subjects were removed from the protocol if they did not continue the therapy for 12 mo due to adverse effects or could not start the therapy due to early death.

Blood counts and liver function tests were checked every 2 wk for the first month, and at 4 wk intervals thereafter. Serum samples were collected once a month for quantitative HCV RNA detection. Protocol liver biopsy was not performed. The log-rank test was used to compare the survival rate of the HCV-positive patients with the HCV-negative patients who underwent transplantation during the same period ($n = 168$).

A total of 28 patients were excluded from the analysis; 12 patients were removed from the protocol because of early death ($n = 9$) or because of drug cessation ($n = 3$). Another 16 patients are currently on the protocol and were therefore excluded from the analysis. Of the remaining 39 patients, 16 (16/39; 41%) obtained a sustained virologic response. The cumulative 5-year survival of the HCV-positive patients was 84%, comparable with that of patients negative for HCV ($n = 168$, 86%).

CONCLUSIONS

LDLT will remain an indispensable therapeutic tool for HCV related end stage liver disease and an alternative to DDLT. The association between LDLT and early HCV recurrence remains to be determined, although most of the recent papers suggest that live donor graft has no effect on short-term outcome or severity of virus recurrence. If living donor graft is associated with early HCV recurrence and consequently poorer graft survival, an aggressive antiviral protocol might improve the outcome of LDLT for HCV.

ACKNOWLEDGMENTS

This study was supported by a Grant-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan and Grants-in-aid

for Research on HIV/AIDS, a multicenter pilot clinical study to compare the safety and efficacy of a steroid free immunosuppression protocol with monoclonal anti-IL2R antibody in HCV positive living donor liver transplantation and Research on Measures for Intractable Diseases from the Ministry of Health, Labor and Welfare of Japan.

REFERENCES

- Strong RW, Lynch SV, Ong TH, Matsunami H, Koido Y, Balderson GA. Successful liver transplantation from a living donor to her son. *N Engl J Med* 1990; **322**: 1505-1507
- Hashikura Y, Makuuchi M, Kawasaki S, Matsunami H, Ikegami T, Nakazawa Y, Kiyosawa K, Ichida T. Successful living-related partial liver transplantation to an adult patient. *Lancet* 1994; **343**: 1233-1234
- Sugawara Y, Makuuchi M. Technical advances in living-related liver transplantation. *J Hepatobiliary Pancreat Surg* 1999; **6**: 245-253
- Forman LM, Lewis JD, Berlin JA, Feldman HI, Lucey MR. The association between hepatitis C infection and survival after orthotopic liver transplantation. *Gastroenterology* 2002; **122**: 889-896
- Berenguer M, Lopez-Labrador FX, Wright TL. Hepatitis C and liver transplantation. *J Hepatol* 2001; **35**: 666-678
- Gane EJ, Portmann BC, Naoumov NV, Smith HM, Underhill JA, Donaldson PT, Maertens G, Williams R. Long-term outcome of hepatitis C infection after liver transplantation. *N Engl J Med* 1996; **334**: 815-820
- Prieto M, Berenguer M, Rayon JM, Cordoba J, Arguello L, Carrasco D, Garcia-Herola A, Olaso V, De Juan M, Gobernado M, Mir J, Berenguer J. High incidence of allograft cirrhosis in hepatitis C virus genotype 1b infection following transplantation: relationship with rejection episodes. *Hepatology* 1999; **29**: 250-256
- Berenguer M, Prieto M, Rayon JM, Mora J, Pastor M, Ortiz V, Carrasco D, San Juan F, Burgueno MD, Mir J, Berenguer J. Natural history of clinically compensated hepatitis C virus-related graft cirrhosis after liver transplantation. *Hepatology* 2000; **32**: 852-858
- Dickson RC, Caldwell SH, Ishitani MB, Lau JY, Driscoll CJ, Stevenson WC, McCullough CS, Pruett TL. Clinical and histologic patterns of early graft failure due to recurrent hepatitis C in four patients after liver transplantation. *Transplantation* 1996; **61**: 701-705
- The Japanese Liver Transplantation Society. Liver Transplantation in Japan. Registry by the Japanese Liver Transplantation Society. *Jpn J Transplant* 2004; **38**: 401-408 (in Japanese).
- Brown RS. Hepatitis C and liver transplantation. *Nature* 2005; **436**: 973-978
- Miller C, Florman S, Kim-Schluger L, Lento P, De La Garza J, Wu J, Xie B, Zhang W, Bottone E, Zhang D, Schwartz M. Fulminant and fatal gas gangrene of the stomach in a healthy live liver donor. *Liver Transpl* 2004; **10**: 1315-1319
- Data from the United Network for Organ Sharing. <http://www.optn.org/latestData/rptData.asp>
- Strader DB, Wright T, Thomas DL, Seeff LB. Diagnosis, management, and treatment of hepatitis C. *Hepatology* 2004; **39**: 1147-1171
- Data from European Liver Transplant Registry. http://www.eltr.org/publi/index_rv.php3
- Data base from Transplant Procurement Management <http://www.tpm.org/presentation.htm>
- Adam R, McMaster P, O'Grady JG, Castaing D, Klempnauer JL, Jamieson N, Neuhaus P, Lerut J, Salizzoni M, Pollard S, Muhlbacher F, Rogiers X, Garcia Valdecasas JC, Berenguer J, Jaeck D, Moreno Gonzalez E. Evolution of liver transplantation in Europe: report of the European Liver Transplant Registry. *Liver Transpl* 2003; **9**: 1231-1243

- 18 Berenguer M. What determines the natural history of recurrent hepatitis C after liver transplantation? *J Hepatol* 2005; **42**: 448-456
- 19 Russo MW, Brown RS Jr. Adult living donor liver transplantation. *Am J Transplant* 2004; **4**: 458-465
- 20 Testa G, Malago M, Nadalin S, Hertl M, Lang H, Frilling A, Broelsch CE. Right-liver living donor transplantation for decompensated end-stage liver disease. *Liver Transpl* 2002; **8**: 340-346
- 21 Wiesner R, Edwards E, Freeman R, Harper A, Kim R, Kamath P, Kremers W, Lake J, Howard T, Merion RM, Wolfe RA, Krom R. Model for end-stage liver disease (MELD) and allocation of donor livers. *Gastroenterology* 2003; **124**: 91-96
- 22 Garcia-Retortillo M, Forns X, Llovet JM, Navasa M, Feliu A, Massaguier A, Bruguera M, Fuster J, Garcia-Valdecasas JC, Rimola A. Hepatitis C recurrence is more severe after living donor compared to cadaveric liver transplantation. *Hepatology* 2004; **40**: 699-707
- 23 Gaglio PJ, Malireddy S, Levitt BS, Lapointe-Rudow D, Lefkowitz J, Kinkhabwala M, Russo MW, Emond JC, Brown RS Jr. Increased risk of cholestatic hepatitis C in recipients of grafts from living versus cadaveric liver donors. *Liver Transpl* 2003; **9**: 1028-1035
- 24 Manez R, Mateo R, Tabasco J, Kusne S, Starzl TE, Duquesnoy RJ. The influence of HLA donor-recipient compatibility on the recurrence of HBV and HCV hepatitis after liver transplantation. *Transplantation* 1995; **59**: 640-642
- 25 Gretch D, Wile M, Gaur L. Donor-recipient match at the HLA-DQB locus is associated with recrudescence of chronic hepatitis following liver transplantation for end stage hepatitis C. *Hepatology* 1993; **18** Suppl: 108A
- 26 Olthoff KM. Hepatic regeneration in living donor liver transplantation. *Liver Transpl* 2003; **9**: S35-41
- 27 Humar A, Horn K, Kalis A, Glessing B, Payne WD, Lake J. Living donor and split-liver transplants in hepatitis C recipients: does liver regeneration increase the risk for recurrence? *Am J Transplant* 2005; **5**: 399-405
- 28 Shiffman ML, Stravitz RT, Contos MJ, Mills AS, Sterling RK, Luketic VA, Sanyal AJ, Cotterell A, Maluf D, Posner MP, Fisher RA. Histologic recurrence of chronic hepatitis C virus in patients after living donor and deceased donor liver transplantation. *Liver Transpl* 2004; **10**: 1248-1255
- 29 Russo MW, Galanko J, Beavers K, Fried MW, Shrestha R. Patient and graft survival in hepatitis C recipients after adult living donor liver transplantation in the United States. *Liver Transpl* 2004; **10**: 340-346
- 30 Thuluvath PJ, Yoo HY. Graft and patient survival after adult live donor liver transplantation compared to a matched cohort who received a deceased donor transplantation. *Liver Transpl* 2004; **10**: 1263-1268
- 31 Rodriguez-Luna H, Vargas HE, Sharma P, Ortiz J, De Petris G, Balan V, Byrne T, Moss A, Mulligan D, Rakela J, Douglas DD. Hepatitis C virus recurrence in living donor liver transplant recipients. *Dig Dis Sci* 2004; **49**: 38-41
- 32 Van Vlierberghe H, Troisi R, Colle I, Ricciardi S, Praet M, de Hemptinne B. Hepatitis C infection-related liver disease: patterns of recurrence and outcome in cadaveric and living-donor liver transplantation in adults. *Transplantation* 2004; **77**: 210-214
- 33 Forman LM, Trotter JF, Emond J. Living donor liver transplantation and hepatitis C. *Liver Transpl* 2004; **10**: 347-348
- 34 Bozorgzadeh A, Jain A, Ryan C, Ornt D, Zand M, Mantry P, Lansing K, Orloff M. Impact of hepatitis C viral infection in primary cadaveric liver allograft versus primary living-donor allograft in 100 consecutive liver transplant recipients receiving tacrolimus. *Transplantation* 2004; **77**: 1066-1070
- 35 Russo MW, Shrestha R. Is severe recurrent hepatitis C more common after adult living donor liver transplantation? *Hepatology* 2004; **40**: 524-526
- 36 Bizollon T, Ahmed SN, Radenne S, Chevallier M, Chevallier P, Parvaz P, Guichard S, Ducerf C, Baulieux J, Zoulim F, Trepo C. Long term histological improvement and clearance of intrahepatic hepatitis C virus RNA following sustained response to interferon-ribavirin combination therapy in liver transplanted patients with hepatitis C virus recurrence. *Gut* 2003; **52**: 283-287
- 37 Giostra E, Kullak-Ublick GA, Keller W, Fried R, Vanlemmens C, Kraehenbuhl S, Locher S, Egger HP, Clavien PA, Hadengue A, Mentha G, Morel P, Negro F. Ribavirin/interferon-alpha sequential treatment of recurrent hepatitis C after liver transplantation. *Transpl Int* 2004; **17**: 169-176
- 38 Abdelmalek MF, Firpi RJ, Soldevila-Pico C, Reed AI, Hemming AW, Liu C, Crawford JM, Davis GL, Nelson DR. Sustained viral response to interferon and ribavirin in liver transplant recipients with recurrent hepatitis C. *Liver Transpl* 2004; **10**: 199-207
- 39 Sheiner PA, Boros P, Klion FM, Thung SN, Schluger LK, Lau JY, Mor E, Bodian C, Guy SR, Schwartz ME, Emre S, Bodenheimer HC Jr, Miller CM. The efficacy of prophylactic interferon alfa-2b in preventing recurrent hepatitis C after liver transplantation. *Hepatology* 1998; **28**: 831-838
- 40 Abdelmalek MF, Firpi RJ, Soldevila-Pico C, Reed AI, Hemming AW, Liu C, Crawford JM, Davis GL, Nelson DR. Sustained viral response to interferon and ribavirin in liver transplant recipients with recurrent hepatitis C. *Liver Transpl* 2004; **10**: 199-207
- 41 Neff GW, Montalbano M, O'Brien CB, Nishida S, Safdar K, Bejarano PA, Khaled AS, Ruiz P, Slapak-Green G, Lee M, Nery J, De Medina M, Tzakis A, Schiff ER. Treatment of established recurrent hepatitis C in liver-transplant recipients with pegylated interferon-alfa-2b and ribavirin therapy. *Transplantation* 2004; **78**: 1303-1307
- 42 Castells L, Vargas V, Allende H, Bilbao I, Luis Lazaro J, Margarit C, Esteban R, Guardia J. Combined treatment with pegylated interferon (alpha-2b) and ribavirin in the acute phase of hepatitis C virus recurrence after liver transplantation. *J Hepatol* 2005; **43**: 53-59
- 43 Moreno Planas JM, Rubio Gonzalez E, Boullosa Grana E, Garrido Botella A, Barrios Peinado C, Lucena Poza JL, Jimenez Garrido M, Sanchez Turrion V, Cuervas-Mons Martinez V. Peginterferon and ribavirin in patients with HCV cirrhosis after liver transplantation. *Transplant Proc* 2005; **37**: 2207-2208
- 44 Yedibela S, Schuppan D, Muller V, Schellerer V, Tannapfel A, Hohenberger W, Meyer T. Successful treatment of hepatitis C reinfection with interferon-alpha2b and ribavirin after liver transplantation. *Liver Int* 2005; **25**: 717-722
- 45 Yedibela S, Schuppan D, Muller V, Schellerer V, Tannapfel A, Hohenberger W, Meyer T. Successful treatment of hepatitis C reinfection with interferon-alpha2b and ribavirin after liver transplantation. *Liver Int* 2005; **25**: 717-722
- 46 Mazzaferro V, Tagger A, Schiavo M, Regalia E, Pulvirenti A, Ribero ML, Coppa J, Romito R, Burgoa L, Zucchini N, Urbanek T, Bonino F. Prevention of recurrent hepatitis C after liver transplantation with early interferon and ribavirin treatment. *Transplant Proc* 2001; **33**: 1355-1357
- 47 Pelletier SJ, Schaubel DE, Punch JD, Wolfe RA, Port FK, Merion RM. Hepatitis C is a risk factor for death after liver retransplantation. *Liver Transpl* 2005; **11**: 434-440

S- Editor Guo SY L- Editor Alpini G E- Editor Ma N

the graft duodenal mucosa only showed mild infiltration with mononuclear cells. The postendoscopy course was uneventful. The pain subsided spontaneously on the next day. She was discharged home on the 21st postoperative day with both grafts functioning well.

In conclusion, double-balloon enteroscopy is feasible in recipients of pancreatic transplants with enteric drainage. This method makes possible to directly visualize duodenal graft and tissue biopsy. It might be employed to look for and possibly intervene at bleeding from the enteric anastomoses, and in special cases for endoscopic retrograde pancreatography of the graft. Its safety must be, however, carefully evaluated as threading on small bowel might potentially be damaging to anastomosed blood vessels supplying transplanted pancreas.

Marek Durlik

Department of Gastroenterological and Transplantation Surgery
Central Clinical Hospital of Ministry of Internal Affairs and Administration
Warsaw, Poland

Wojciech Kosmala
Janusz Milewski

Department of Internal Medicine and Gastroenterology
Central Clinical Hospital of Ministry of Internal Affairs and Administration
Warsaw, Poland

Agnieszka Serwacka

Department of Internal Medicine and Nephrology
Central Clinical Hospital of Ministry of Internal Affairs and Administration
Warsaw, Poland

Grażyna Rydzewska

Andrzej Rydzewski
Institute of Experimental and Clinical Medicine
Polish Academy of Sciences
Warsaw, Poland

Address correspondence to: Andrzej Rydzewski, M.D., Ph.D., Klinika Chorob Wewnętrznych i Nefrologii, Centralny Szpital Kliniczny MSWiA, Woloska 137, 02-507 Warszawa, Poland.
E-mail: Andrzej.Rydzewski@cskmswia.pl

Received 13 February 2006. Revision requested 15 March 2006.

Accepted 17 March 2006.

Copyright © 2006 by Lippincott Williams & Wilkins

ISSN 0041-1337/06/8204-578

DOI: 10.1097/01.tp.0000231834.20634.b2

REFERENCES

1. Laftavi MR, Gruessner AC, Bland BJ, et al. Diagnosis of pancreas rejection: cystoscopic transduodenal versus percutaneous computed tomography scan-guided biopsy. *Transplantation* 1998; 65: 528.
2. Lee BC, McGahan JP, Perez RV, et al. The role of percutaneous biopsy in detection of pancreatic transplant rejection. *Clin Transplant* 2000; 14: 493.
3. Kayler LK, Merion RM, Rudich SM, et al. Evaluation of pancreatic allograft dysfunction by laparoscopic biopsy. *Transplantation* 2002; 74: 1287.
4. Mark W, Hechenleitner P, Dietze O, et al. Duodenal histology for monitoring treatment of acute rejection in pancreaticoduodenal allografts in rats. *Transplantation* 2002; 73: 198.
5. Hawthorne WJ, Allen RD, Greenberg ML, et al. Simultaneous pancreas and kidney transplant rejection: separate or synchronous events? *Transplantation* 1997; 63: 352.
6. Boykin KN, Zibari GB. Management of complications of simultaneous kidney-pancreas transplantation with temporary venting jejunostomy. *Clin Transplant* 2003; 17 Suppl 9: 65.
7. Rudroff C, Scheele J. Pancreas transplantation: Eine neue chirurgische Methode mit endoskopisch zugänglicher enteraler Drainage. *Transplantationsmedizin* 1998; Suppl: 28.
8. Yamamoto H, Sekine Y, Sato Y, et al. Total enteroscopy with a nonsurgical steerable double-balloon method. *Gastrointest Endosc* 2001; 53: 216.

Cyclosporin A for Treatment of Hepatitis C Virus After Liver Transplantation

Cirrhosis secondary to hepatitis C virus (HCV) infection is a leading indication for liver transplantation. HCV recurrence, however, is nearly certain and might worsen patient/graft outcome. A recent paper (1) reported that HCV infection is associated with a 23% increase in mortality and a 30% increase in the graft failure rate. The goal of HCV management in the transplantation setting is to prevent graft loss due to recurrent HCV infection, raising questions about a possible role for immunosuppression regimens and antiviral therapy.

A previous report (2) indicated that

cyclosporin A (CyA) and interferon-alpha2b might effectively inhibit HCV replication in vitro. The antiviral effects of CyA for patients with chronic HCV (3) and those for HCV recurrence after transplantation (4), however, are controversial. We conducted a pilot study of the use of CyA, interferon, and ribavirin for preemptive therapy of HCV after liver transplantation.

Until October 2003, 41 HCV-positive patients underwent liver transplantation from living donors at the University of Tokyo Hospital. The immunosuppression regimens consisted of steroids and

tacrolimus (5). The targeted whole-blood tacrolimus level was 15 to 20 ng/ml during the seven days after living donor liver transplantation (LDLT), which was gradually tapered to 5 ng/ml six months after LDLT. Prednisolone was tapered to 0.05 mg/day/kg six months after LDLT but was not stopped. All of the HCV-positive patients preemptively received interferon-alpha2b and ribavirin therapy (6), which was started approximately one month after transplantation. HCV RNA level was measured by real-time-polymerase chain reaction (7) and Amplicor HCV (Roche Molecular Systems, Pleasanton, USA). Of the 41 patients, six died within two years; 14 obtained a viral response (<50 IU/ml by Amplicor HCV) at the end of the treatment period (one year) and the response was sustained for another six months without the antiviral therapy; 21 did not respond to the antiviral therapy. Of the 21 nonresponders, eight patients continued with the protocol, which called for a change from tacrolimus to CyA without changing the antiviral therapy. The targeted CyA trough level was 100 ng/ml.




FIGURE 1. Change of the viral titer in the eight patients.

All of the patients were genotype 1b. The serum HCV titer was measured once a month by reverse transcriptase-polymerase chain reaction. The Institutional Review Boards at University of Tokyo Hospital approved the protocol.

In five of the eight subjects (63%), the HCV titer was negative by Amplicor HCV within three months after the conversion and remained negative for another six months on CyA, interferon, and ribavirin (Figure 1). Liver and renal functions remained stable in all of the patients, and none of them had complications of acute cellular rejection after the conversion.

Our findings support the use of CyA in combination with interferon and ribavirin for the eradication of HCV in previous nonresponders. These findings suggest that a controlled study to confirm the benefit of CyA for preemptive treatment of HCV after liver transplantation is warranted.

Yasuhiko Sugawara
Junichi Kaneko
Masatoshi Makuuchi

Artificial Organ and Transplantation
Division
Department of Surgery
Graduate School of Medicine
University of Tokyo
Tokyo, Japan

This work was supported by a Grant-in-aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan and Grants-in-aid for Research on HIV/AIDS and Research on Measures for Intractable Diseases from the Ministry of Health, Labor and Welfare of Japan.

Address correspondence to: Yasuhiko Sugawara, MD, Artificial Organ and Transplantation Division, Department of Surgery, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan.

E-mail: yasusuga-tky@umin.ac.jp

Received 8 April 2006. Revision requested 12 April 2006.

Accepted 13 April 2006.

Copyright © 2006 by Lippincott Williams & Wilkins

ISSN 0041-1337/06/8204-579

DOI: 10.1097/01.tp.0000229397.81425.51

REFERENCES

1. Forman LM, Lewis JD, Berlin JA, et al. The association between hepatitis C infection

- and survival after orthotopic liver transplantation. *Gastroenterology* 2002; 122: 889.
2. Watashi K, Hijikata M, Hosaka M, et al. Cyclosporin A suppresses replication of hepatitis C virus genome in cultured hepatocytes. *Hepatology* 2003; 38: 1282.
3. Cotler SJ, Morrissey MJ, Wiley TE, et al. A pilot study of the combination of cyclosporin A and interferon alfacon-1 for the treatment of hepatitis C in previous nonresponder patients. *J Clin Gastroenterol* 2003; 36: 352.
4. Martin P, Busuttil RW, Goldstein RM, et al. Impact of tacrolimus versus cyclosporine in hepatitis C virus-infected liver transplant recipients on recurrent hepatitis: a prospective, randomized trial. *Liver Transpl* 2004; 10: 1258.
5. Sugawara Y, Makuuchi M, Kaneko J, et al. Correlation between optimal tacrolimus doses and the graft weight in living donor liver transplantation. *Clin Transplant* 2002; 16: 102.
6. Sugawara Y, Makuuchi M, Matsui Y, et al. Preemptive therapy for hepatitis C virus after living-donor liver transplantation. *Transplantation* 2004; 78: 1308.
7. Takeuchi T, Katsume A, Tanaka T, et al. Real-time detection system for quantification of hepatitis C virus genome. *Gastroenterology* 1999; 116: 636.

Does Using HTK Solution for Cold Perfusion of Cadaveric Kidneys Save Money?

Since 1987, University of Wisconsin (UW) solution has been the standard solution for cold perfusion and preservation of kidneys for transplant. In 2001, the U.S. Food and Drug Administration approved histidine-tryptophan-ketoglutarate (HTK) for cold perfusion and storage of renal allografts. Data has shown comparable outcomes between UW and HTK for preservation of renal, liver, and pancreatic allografts (1-3). Theoretical advantages for HTK compared to UW solution include: lower viscosity (which might lead to better microperfusion of the allograft), lower potassium concentrations, and easier handling properties, although the actual effects on clinical outcomes are not fully understood. On

the other hand, HTK is significantly less expensive than UW solution on a per liter basis. For this reason, HTK has become an attractive alternative to UW solution for some Organ Procurement Organizations. In reported clinical experience with HTK, larger volumes of solution have been used to achieve exsanguination of the recovered organs. The argument is frequently made that these larger volumes negated any significant cost savings associated with the cheaper HTK solution.

Gift of Life Michigan switched from UW to HTK solution in December 2003. This decision was based on an impression of clinical equipoise between UW and HTK solutions, with a projected materials cost savings of 47% per donor.

We compared the material costs of 77 consecutive kidney-pancreas donor procurements, in which the organs were used for transplant. The perfusion volumes include the aortic flush, the back-table flush of the renal allografts, and the storage volume of the renal grafts. As predicted, significantly larger volumes of fluid were used in the perfusion of the organs with HTK (Table 1). The cost of a liter of UW (UW solution \$282, filter \$10, and additives \$30) was \$322/L compared to \$148/L for HTK (no additives or filters needed). Costs of UW perfusion were significantly higher than HTK perfusion. The transition from UW to HTK saved \$548 (USD) per renal donor. This represents a 43% actual cost reduction, which is similar to the 47% projected reduction. In conclusion, the transition from UW to HTK has resulted in a significant cost savings for Gift of Life Michigan.

TABLE 1. UW vs. HTK flush solution for cold perfusion of kidneys for transplant

	UW	HTK	Difference
N	41	36	
Mean flush volume per donor (mL)	3516 ± 795	5575 ± 1506	2059 ^a
Cost per liter solution (US \$)	\$322	\$148	\$174
Cost per renal donor (US \$)	\$1288	\$740	\$548 ^a

^a P < 0.05.

Michael J. Englesbe
David Heidt
Randall Sung
Department of Surgery
Division of Transplantation
University of Michigan
Ann Arbor, MI

Identification of an Oligopeptide Binding to Hepatocellular Carcinoma

Akira Shimizu^a Fukuto Maruta^a Noriyuki Akita^a Shiro Miwa^a
Leonard W. Seymour^b David J. Kerr^b Alan L. Parker^c Shinichi Miyagawa^a

^aDepartment of Surgery, Shinshu University School of Medicine, Matsumoto, Japan; ^bDepartment of Clinical Pharmacology, Oxford University Radcliffe Infirmary, Oxford, and ^cCardiovascular and Medical Sciences, University of Glasgow, Glasgow, UK

Key Words

Biopanning · Hepatocellular carcinoma · Peptide · Phage display · Tumor targeting

Abstract

Objectives: We carried out identification of a small peptide binding to human hepatocellular carcinoma (HCC) cells with the aim of applying the peptide for future HCC-targeted therapy or imaging. **Methods:** The biopanning technique using phage peptide display libraries was performed on HCC cells in vitro, and a phage clone expressing the HCC-binding peptide motif was selected. The binding activity of the selected phage was evaluated by plaque infection assay and immunofluorescence on cell lines. In addition, the binding activity of the peptide-expressing phage was investigated using HCC specimens derived from patients who had undergone hepatectomy for HCC. **Results:** A heptapeptide, Thr-Thr-Pro-Arg-Asp-Ala-Tyr (TTPRDAY), was identified as a motif binding to HCC. TTPRDAY bound specifically to HCC cells in comparison with other cancer cells, and the binding to HCC cells was also confirmed by immunofluorescence. In addition, the synthesized TTPRDAY peptide showed binding activity and a non-mitogenic effect on HCC cells in vitro. TTPR-

DAY-presenting phage showed more significant binding to HCC cells derived from specimens obtained from actual patients than to non-cancerous liver tissue. **Conclusion:** The motif TTPRDAY, identified by the biopanning technique, shows significant binding to HCC cells.

Copyright © 2006 S. Karger AG, Basel

Introduction

Hepatocellular carcinoma (HCC) is one of the commonest malignancies worldwide [1], and the prognosis of affected patients remains poor [2–4]. In many patients with HCC, obtaining curativity by surgery or radiofrequency ablation of the cancer is difficult due to the high frequency of intrahepatic metastases or multicentric occurrence. Other therapeutic options for intrahepatic localized HCC are percutaneous ethanol injection, microwave therapy, cryotherapy and transarterial chemoembolization. However, the therapeutic efficacy of these treatments is limited. For example, the 5-year survival rate after transarterial chemoembolization for HCC is only 12.5% [5]. Patients with distant metastatic HCC are treated with systemic chemotherapy, but its reported

therapeutic efficacy is dismal; Chung et al. [6] reported that intra-arterial cisplatin infusion and systemic interferon- α yielded a response rate of 33% and a median survival time of 19 weeks for HCC patients with major portal vein thrombosis or distant metastases.

The chief obstacles to establishing effective therapies for HCC are the multiplicity of the lesions (multicentric occurrence of the tumor and intrahepatic or distant metastases) and severe liver dysfunction due to underlying liver cirrhosis. Therefore, establishment of a methodology that could target HCC efficiently would bring about a dramatic improvement in the efficacy of treatments for HCC. For example, HCC-selective ligands would enable targeted delivery of a variety of therapeutic agents, such as genes or cytotoxic drugs, and would also facilitate better imaging of multiple HCC lesions. Such agents are clearly appropriate for HCC patients with liver cirrhosis, because they could minimize the degree of drug entry into non-HCC cells and the resulting damage to liver function.

Candidate targeting agents have been studied by several groups in attempts to achieve tumor tropism. The ligands that have been evaluated include some antibodies, as well as fragments and single-chain Fv molecules and growth factors. Recently, however, this empirical approach to the identification of targeting ligands has been largely superseded by the use of library-based screening systems, which have been designed to allow iterative selection of high-affinity ligands by repeated screening and enrichment of living libraries.

Barry et al. [7] have introduced peptide-presenting phage libraries to enable selection of peptides binding to several different cell types. Since then, the biopanning technique using phage peptide display libraries has been shown to be a powerful tool for identifying specific ligands on target organs and tumors [8–10]. The peptide-presenting phage library used is based on a combinatorial library of random peptide heptamers fused to a minor coat protein (pIII) of phage M13 and contains about 2.8×10^9 different sequences. With successive rounds of biopanning, different peptides selective for the target cells can be identified.

In the present study, we used the phage panning technique to identify a peptide that would have the specific ability to bind to human HCC cells. In addition, in order to confirm the feasibility of applying the identified peptide to clinical practice, its HCC-binding activity was assessed on HCC samples derived from patients who had undergone hepatectomy for operable HCC.

Materials and Methods

Cell Lines

The cell line Huh-7 (human HCC) was obtained from the Health Science Research Resources Bank (HSRRB, Osaka, Japan), and the cell lines Hep3B (human HCC), HepG2 (human hepatoblastoma), DLD-1 (human colorectal carcinoma), AZ521 (human gastric cancer) and Saos2 (human osteosarcoma) were obtained from the Institute of Development, Aging and Cancer, Tohoku University (Sendai, Japan). Cells were maintained in Dulbecco's modified Eagle medium (DMEM, Sigma Chemical, Poole, UK) or RPMI-1640 medium (Sigma) supplemented with 10% fetal bovine serum, 100 U/ml penicillin and 100 mg/ml streptomycin at 37°C in 5% CO₂. The cells were passaged and expanded by trypsinization of the cell monolayers followed by replating every 4 days.

Selection of Cell-Targeting Peptides

The Ph.C7C M13 heptapeptide phage display library (New England Biolabs, Beverly, Mass., USA) was used for biopanning. Huh-7 cells were grown to confluence in a 6-well plate. They were acclimated to 4°C for 30 min and washed twice in phosphate-buffered saline (PBS, Dainippon, Osaka, Japan) containing 1% (w/v) bovine serum albumin (BSA, Sigma). The phage library of 2×10^{11} plaque-forming units (pfu)/well diluted in 1 ml DMEM containing 1% BSA was added to each well at 4°C. After 1 h of incubation with gentle agitation, medium containing unbound phage was removed, and the cells were washed 4 times in PBS containing 1% BSA. This was followed by addition of 1 ml/well 0.2 M glycine-HCl (pH 2.2) as a general buffer for nonspecific disruption of binding interactions for 5 min, and fluids were neutralized with 150 μ l 1 M Tris-HCl (pH 9.1).

The cells were scraped from the 6-well plate. Numbers of eluted phages were established by titrating on X-gal (5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside; Wako, Osaka, Japan) and IPTG (isopropyl- β -D-thiogalactopyranoside; Wako) agar plates containing *Escherichia coli* strain ER2738. Remaining phages were amplified by addition to a 20-ml early log phase culture of ER2738 for 5 h at 37°C with vigorous shaking (150 rpm). Amplified phages were isolated from the resulting culture using the manufacturer's recommended protocol, concentrated, titered and used for subsequent rounds of biopanning. In total, four rounds of biopanning were performed in triplicate.

Isolation and Sequencing of Phage DNA

After each round of biopanning, individual phage clones were isolated from each replicate and their total DNA was isolated according to the recommended protocol of the sequencing kit manufacturer (Biosystems, Perkin Elmer, Foster City, Calif., USA). The resulting DNA was used for sequencing analysis using the -96 primer together with a BigDye terminator cycle sequencing kit (Amersham Biosciences, Amersham, UK). The DNA sequences were determined using an automated ABI PRISM sequencer 3100 Genetic Analyzer (Applied Biosystems). Next, searches for human proteins mimicked by the selected peptide motifs were carried out using online databases through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/BLAST/>).