

Region 1		
Outer sense	nt 2222-2241	CTTACTTTTGGAGAGAAAC
Outer antisense	nt 2490-2509	GGACAGTAGAACAATAAAG
Inner sense	nt 2222-2241	CTTACTTTTGGAGAGAAAC
Inner antisense	nt 2478-2497	GAATAAAGCCCAGTAAAGTT
Region 2		
Outer sense	nt 2413-2434	CGGTCCGAG AAGATCTCAATC
Outer antisense	nt 2816-2835	GTTCUUAAGAATATGGTGAC
Inner sense	nt 2434-2452	CTCCGGAAATCTCAATGTTAG
Inner antisense	nt 2816-2835	GTTCUUAAGAATATGGTGAC
Region 3		
Outer sense	nt 2490-2509	CTTTATTTCTTACTGTACC
Outer antisense	nt 3121-3143	CGATTGGTGGAGGCAGGAGGAGG
Inner sense	nt 2637-2656	ATGCC TACTAGGTTTATCC
Inner antisense	nt 3121-3143	CGATTGGTGGAGGCAGGAGGAGG

each plate, from which recombinant plasmid DNA was purified using a commercially available kit (Plasmid midi kit, Qiagen, Valencia, CA). Nucleotide sequences were determined bidirectionally using the dye terminator cycle sequencing ready reaction kit (PE Applied Biosystems, CA) and the PCR primers. Sequencing was performed using an automated DNA sequencer (ABI 377; PE Applied Biosystems).

The determined amino acid sequences in the polymerase region are shown in Figure 3. No amino acid sequence changes were found at the start of lamivudine treatment. At the time of viral breakthrough, rtM204I with rtL80I became dominant. After the start of interferon treatment, rtM204I was replaced by rtM204V and rtL80V by rtL80V. At the final stage of the disease, mutation rtL180M appeared besides rtM204V and rtL80V.

## DISCUSSION

Lamivudine monotherapy is effective in suppressing HBV replication and ameliorating liver disease in chronic hepatitis B patients regardless of HBeAg positivity. A one-year study of HBeAg-positive chronic hepatitis B patients showed that 16% of these patients become seroconverted to anti-HBe and 72% of these patients showed normalization of their ALT levels<sup>29</sup>. Furthermore, treatment with lamivudine is associated with histologic improvement not only in terms of necroinflammatory score but also in terms of fibrosis score after long-term treatment<sup>29</sup>.

One advantage of lamivudine is that it can be used safely in patients with decompensated cirrhosis<sup>13,14</sup>. In contrast to IFN- $\alpha$ , lamivudine is well tolerated without any significant side effects even in patients with decompensated cirrhosis. Furthermore, lamivudine can improve liver function and survival prognosis.

However, the emergence of a drug-resistant mutant is a big problem in lamivudine treatment. A large-scale Asian study showed that lamivudine resistant HBV infection occurred in 23% of patients in year one and 65% of patients in year five. Hepatitis flares, which occurred more commonly in patients with lamivudine resistant mutations, occurred in 10% of patients in year one, and in 18% to



Figure 2 Histopathological findings of liver specimens showing irregularly-shaped parenchymal cells with massive necrosis (A) and scarce hepatocyte regeneration (B) surrounded by extensive fibrosis (A, HE  $\times$  20; B, HE  $\times$  80)

21% of patients in years two to five. Among patients with lamivudine resistant HBV infection, occurrence of hepatic decompensation increased significantly in patients with lamivudine resistant HBV infection for more than 4 years (from 0% to 6%)<sup>15,31</sup>. In this large-scale Asian study, liver-disease-related death occurred in two patients.

The prognosis of patients with lamivudine-resistant HBV infection, particularly those with advanced liver disease, may be determined by the timing and severity of breakthrough hepatitis. However, the viral factors that may influence the severity of this hepatitis remain to be clarified. A recent study indicated that patients with a normal ALT level even after the emergence of a YMDD motif mutant are characterized by HBeAg negativity during pretreatment, HBeAg loss during therapy, a longer duration from the commencement of therapy until the emergence of YMDD mutant, and lack of mixed-type YMDD mutants<sup>16,31</sup>. In contrast, patients with severely exacerbated hepatitis after the emergence of a YMDD mutant tend to have more substitutions in the reverse transcriptase (rt) region within the polymerase gene at the time of hepatitis exacerbation than those without hepatitis exacerbation<sup>16,31</sup>.

Our patient acquired amino acid mutations in the polymerase region one after the other. Amino acid changes in rtM204I appeared at the time of viral breakthrough. After the initial treatment with vidarabine, rtM204V substituted for rtM204I in one of the four clones. During the interferon treatment, rtM204V became predominant.

Another mutation observed in our patient was rtL80I/V. Ogata *et al.*<sup>27</sup> showed that rtL180M is accompanied with rtM204I in some patients with resistance to lamivudine. Because the mutation at aa position 80 was found at the same time as that at aa position 204 in our patient, it is not clear whether the mutation at aa position 80 affects the clinical course.




Figure 3 Comparison of amino acid sequences of HBV polymerase gene of isolates before lamivudine treatment (sample 1) and four sequential isolates (samples 2-5) during treatment. A HBV mutant with substitutions of isoleucine for leucine at residue 80 (rtL80I) in combination with isoleucine for methionine at residue 204 (rtM204I) was observed 12 mo after treatment (sample 3). After vidarabine treatment, another HBV mutant with substitutions of valine for leucine at residue 80 (rtL80V) and valine for methionine at residue 204 (rtM204V) was observed (sample 4). These mutations predominated in combination with methionine for leucine at residue 180 (rtL180M) after interferon treatment (sample 5). The published HBV DNA sequence of hepatitis B virus variant (genotype C, AB033550, Okamoto *et al.*<sup>25</sup>) was used for comparison.

At the final stage of the disease with deterioration of the condition of the patient, rtL180M, rtM204I/V and rtL80I/V became predominant. Natsuzaka *et al.*<sup>25</sup> showed that rtL180M and rtM204V are related to the exacerbation of hepatitis. Interestingly, a patient with a marked elevation in HBV DNA level in Ogata's report had rtL180M in addition to rtL80I and rtM204I. Therefore, the acquisitions of rtL80I/V and rtL180M in addition to rtM204I/V may be associated with a severe exacerbation of hepatitis. Large-scale studies are necessary to elucidate this hypothesis.

In our patient, vidarabine decreased HBV DNA levels and improved liver function. Although the long-term use of vidarabine is contraindicated because of its possible side effects including irreversible neurotoxicity<sup>22</sup>, its short-term use is effective for controlling active HBV infection and herpes simplex viral infection.

Vidarabine was replaced by interferon- $\alpha$  because adefovir dipivoxil was not available in 2002. Serum HBV DNA and bilirubin levels increased again, which led to a

fatal outcome, and HBV clones that have rtL180M became predominant, indicating that withdrawal of vidarabine and administration of interferon may be dangerous for the treatment of severe breakthrough hepatitis. Since interferon may potentially precipitate immunological flares and liver failure<sup>27</sup>, nucleotide analogues that are effective for lamivudine-resistant HBV such as adefovir dipivoxil<sup>16, 22</sup>, entecavir<sup>13, 28</sup> and tenofovir<sup>16, 27</sup>, should be used for the treatment of severe breakthrough hepatitis instead of vidarabine or interferon.

In conclusion, antiviral therapy should be considered in the treatment of patients with hepatic failure after breakthrough hepatitis caused by HBV mutants to lamivudine. The serial acquisition of amino acid mutations outside the YMDD motif in the polymerase region may be associated with severe hepatitis.

REFERENCES

1 Lok AS, McMahon BJ. Chronic hepatitis B: update of

- recommendations. *Hepatology* 2004; **39**: 857-861
- 2 Liaw YF, Leung N, Guan R, Lau GK, Merikan L, McCaughan G, Kane E, Kao JH, Omata M. Asian-Pacific consensus statement on the management of chronic hepatitis B: a 2005 update. *Liver Int* 2005; **25**: 472-489
  - 3 Villeneuve JP, Condroy LD, Willems B, Pomier-Layrargues G, Fenyves D, Bilodeau M, Leduc R, Peltekian K, Wong F, Margulies M, Heathcote EJ. Lamivudine treatment for decompensated cirrhosis resulting from chronic hepatitis B. *Hepatology* 2000; **31**: 207-210
  - 4 Yao FY, Bass NM. Lamivudine treatment in patients with severely decompensated cirrhosis due to replicating hepatitis B infection. *J Hepatol* 2000; **33**: 301-307
  - 5 Kapoor D, Gupta RC, Wakil SM, Kazim SN, Kaul R, Agarwal SR, Raisuddin S, Hasnain SE, Sarin SK. Beneficial effects of lamivudine in hepatitis B virus-related decompensated cirrhosis. *J Hepatol* 2000; **33**: 308-312
  - 6 Sponseller CA, Bacon BR, Di Bisceglie AM. Clinical improvement in patients with decompensated liver disease caused by hepatitis B after treatment with lamivudine. *Liver Transpl* 2000; **6**: 715-720
  - 7 Fontana RJ, Koofie EB, Carey W, Fried M, Reddy R, Kowdley KV, Soldevila-Pico C, McClure LA, Lok AS. Effect of lamivudine treatment on survival of 309 North American patients awaiting liver transplantation for chronic hepatitis B. *Liver Transpl* 2002; **8**: 433-439
  - 8 Yao FY, Terrault NA, Freise C, Maslow L, Bass NM. Lamivudine treatment is beneficial in patients with severely decompensated cirrhosis and actively replicating hepatitis B infection awaiting liver transplantation: a comparative study using a matched, untreated cohort. *Hepatology* 2001; **34**: 411-416
  - 9 Hann HW, Fontana RJ, Wright T, Everson G, Baker A, Schiff ER, Riely C, Anschuetz G, Gardner SD, Brown N, Griffiths D. A United States compassionate use study of lamivudine treatment in nontransplantation candidates with decompensated hepatitis B virus-related cirrhosis. *Liver Transpl* 2003; **9**: 49-56
  - 10 Manolakopoulos S, Karatapanis S, Elefsiniotis J, Mathou N, Vlachogiannakos I, Iliadou E, Kougioumtzan A, Economidou M, Triantos C, Tzourmakliotis D, Avgerinos A. Clinical course of lamivudine monotherapy in patients with decompensated cirrhosis due to HBeAg negative chronic HBV infection. *Am J Gastroenterol* 2004; **99**: 57-63
  - 11 Fattovich G, Broffo L, Giustina G, Noventa F, Pontisso P, Alberti A, Realdi G, Ruol A. Natural history and prognostic factors for chronic hepatitis type B. *Gut* 1991; **32**: 294-298
  - 12 Liaw YF, Tai DI, Chu CM, Chen TJ. The development of cirrhosis in patients with chronic type B hepatitis: a prospective study. *Hepatology* 1988; **8**: 493-496
  - 13 Liaw YF, Lin DY, Chen TJ, Chu CM. Natural course after the development of cirrhosis in patients with chronic type B hepatitis: a prospective study. *Liver* 1989; **9**: 235-241
  - 14 Fattovich G, Giustina G, Schalm SW, Hadziyannis S, Sanchez-Tapias J, Almasio P, Christensen E, Krogsgaard K, Degos F, Carneiro de Moura M. Occurrence of hepatocellular carcinoma and decompensation in western European patients with cirrhosis type B. The EUROHEP Study Group on Hepatitis B Virus and Cirrhosis. *Hepatology* 1995; **21**: 77-82
  - 15 Kim JW, Lee HS, Woo GH, Yoon JH, Jang JI, Chi JG, Kim CY. Fatal submassive hepatic necrosis associated with tyrosine-methionine-aspartate-aspartate-motif mutation of hepatitis B virus after long-term lamivudine therapy. *Clin Infect Dis* 2001; **33**: 403-405
  - 16 Wang JH, Lu SN, Lee CM, Lee JF, Chou YP. Fatal hepatic failure after emergence of the hepatitis B virus mutant during lamivudine therapy in a patient with liver cirrhosis. *Scand J Gastroenterol* 2002; **37**: 366-369
  - 17 Kagawa T, Watanabe N, Kanouda H, Takayama I, Shiba T, Kanai T, Kawazoe K, Takashimizu S, Kumaki N, Shimamura K, Matsuzaki S, Mine T. Fatal liver failure due to reactivation of lamivudine-resistant HBV mutant. *World J Gastroenterol* 2004; **10**: 1686-1687
  - 18 Kamisango K, Kamogawa C, Sumi M, Goto S, Hirao A, Gonzales F, Yasuda K, Iino S. Quantitative detection of hepatitis B virus by transcription-mediated amplification and hybridization protection assay. *J Clin Microbiol* 1999; **37**: 310-314
  - 19 Hoofnagle JH, Minuk GY, Dusheiko GM, Schafer DF, Johnson R, Straus S, Jones EA, Gerin JL, Ishak K. Adenine arabinoside 5'-monophosphate treatment of chronic type B hepatitis. *Hepatology* 1982; **2**: 784-788
  - 20 Weller IV, Lok AS, Mindel A, Karayiannis P, Galpin S, Monjardino J, Sherlock S, Thomas HC. Randomised controlled trial of adenine arabinoside 5'-monophosphate (ARA-A) in chronic hepatitis B virus infection. *Gut* 1985; **26**: 745-751
  - 21 Marcellin P, Ouzan D, Degos F, Brechot C, Metman EH, Degott C, Chevalier M, Berthelot P, Trepo C, Benhamou JP. Randomized controlled trial of adenine arabinoside 5'-monophosphate in chronic active hepatitis B: comparison of the efficacy in heterosexual and homosexual patients. *Hepatology* 1989; **10**: 328-331
  - 22 Hoofnagle JH, Hanson RG, Minuk GY, Pappas SC, Schafer DF, Dusheiko GM, Straus SE, Popper H, Jones EA. Randomized controlled trial of adenine arabinoside monophosphate for chronic type B hepatitis. *Gastroenterology* 1984; **86**: 150-157
  - 23 Lai CL, Chien RN, Leung NW, Chang TT, Guan R, Tai DL, Ng KY, Wu PC, Dent K, Barber J, Stephenson SL, Gray DF. A one-year trial of lamivudine for chronic hepatitis B. Asia Hepatitis Lamivudine Study Group. *N Engl J Med* 1998; **339**: 61-68
  - 24 Dienstag JL, Goldin RD, Heathcote EJ, Hann HW, Woessner M, Stephenson SL, Gardner S, Gray DF, Schiff ER. Histological outcome during long-term lamivudine therapy. *Gastroenterology* 2003; **124**: 105-117
  - 25 Lok AS, Lai CL, Leung N, Yao GB, Cui ZY, Schiff ER, Dienstag JL, Heathcote EJ, Little NR, Griffiths DA, Gardner SD, Castiglia M. Long-term safety of lamivudine treatment in patients with chronic hepatitis B. *Gastroenterology* 2003; **125**: 1714-1722
  - 26 Suzuki F, Akuta N, Suzuki Y, Sezaki H, Arase Y, Hosaka T, Someya T, Kobayashi M, Saitoh S, Ikeda K, Kobayashi M, Matsuda M, Satoh J, Watahiki S, Kumada H. Clinical and virological features of non-breakthrough and severe exacerbation due to lamivudine-resistant hepatitis B virus mutants. *J Med Virol* 2006; **78**: 341-352
  - 27 Ogata N, Fujii K, Takigawa S, Nomoto M, Ichida T, Asakura H. Novel patterns of amino acid mutations in the hepatitis B virus polymerase in association with resistance to lamivudine therapy in Japanese patients with chronic hepatitis B. *J Med Virol* 1999; **59**: 270-276
  - 28 Natsuizaka M, Hige S, Ono Y, Ogawa K, Nakanishi M, Chuma M, Yoshida S, Asaka M. Long-term follow-up of chronic hepatitis B after the emergence of mutations in the hepatitis B virus polymerase region. *J Viral Hepat* 2005; **12**: 154-159
  - 29 Janssen HL, Brouwer JT, Nevens F, Sanchez-Tapias JM, Craxi A, Hadziyannis S. Fatal hepatic decompensation associated with interferon alfa. European concerted action on viral hepatitis (Eurohep). *BMJ* 1993; **306**: 107-108
  - 30 Perrillo R, Schiff E, Yoshida E, Stalter A, Hirsch K, Wright T, Gutfreund K, Lamy P, Murray A. Adefovir dipivoxil for the treatment of lamivudine-resistant hepatitis B mutants. *Hepatology* 2000; **32**: 129-134
  - 31 Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, Chang TT, Kitis G, Rizzetto M, Marcellin P, Lim SG, Goodman Z, Wulfsohn MS, Xiong S, Fry J, Brosgart CL. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med* 2003; **348**: 800-807
  - 32 Marcellin P, Chang TT, Lim SG, Tong MJ, Sievert W, Shiffman ML, Jeffers L, Goodman Z, Wulfsohn MS, Xiong S, Fry J, Brosgart CL. Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *N Engl J Med* 2003; **348**: 808-816
  - 33 Chang TT, Gish RG, Hadziyannis SJ, Cianciara J, Rizzetto M, Schiff ER, Pastore G, Bacon BR, Poynard T, Ioshi S, Kleszczewski KS, Thiry A, Rose RE, Colonna RJ, Hines RC. A dose-ranging study of the efficacy and tolerability of entecavir

- in Lamivudine-refractory chronic hepatitis B patients. *Gastroenterology* 2005; **129**: 1198-1209
- 34 Chang TT, Gish RG, de Man R, Gadano A, Sollano J, Chao YC, Lok AS, Han KH, Goodman Z, Zhu J, Cross A, DeHertogh D, Wilber R, Colonna R, Apelian D. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med* 2006; **354**: 1001-1010
- 35 Lai CL, Shouval D, Lok AS, Chang TT, Cheinquer H, Goodman Z, DeHertogh D, Wilber R, Zink RC, Cross A, Colonna R, Fernandes L. Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N Engl J Med* 2006; **354**: 1011-1020
- 36 van Bommel F, Wunsche T, Mauss S, Reinke P, Bergk A, Schurmann D, Wiedenmann B, Berg T. Comparison of adefovir and tenofovir in the treatment of lamivudine-resistant hepatitis B virus infection. *Hepatology* 2004; **40**: 1421-1425
- 37 van der Eijk AA, Hansen BE, Niesters HG, Janssen HL, van de Ende M, Schalm SW, de Man RA. Viral dynamics during tenofovir therapy in patients infected with lamivudine-resistant hepatitis B virus mutants. *J Viral Hepat* 2005; **12**: 364-372

S- Editor Wang GP L- Editor Wang XL E- Editor Lu W



Review

# Oxidative stress and hepatitis C viral infection

Kazuhiko Koike\*, Hideyuki Miyoshi

*Department of Internal Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan*

Received 17 October 2005; received in revised form 4 November 2005; accepted 4 November 2005

Available online 20 December 2005

## Abstract

The involvement of oxidative stress in the pathogenesis of hepatitis and hepatocellular carcinoma has been strongly suggested. Oxidative stress is produced by inflammatory processes that occur in hepatitis via immunological mechanisms. In addition, in hepatitis C virus (HCV) infectious disease, some role has been assigned to viral proteins in the induction of oxidative stress. In the presence of hepatic steatosis, insulin resistance and increased levels of some cytokines, all of which are also induced by viral protein expression, oxidative stress is enhanced in HCV infection. In this sense, the role of oxidative stress in the progression of chronic hepatitis and hepatocarcinogenesis is greater in hepatitis C than in other types of hepatitis such as hepatitis B or autoimmune hepatitis. The additive effects of oxidative stress caused by the inflammatory process and that induced by HCV proteins may, furthermore, exert synergistic effects with alterations in intracellular signaling systems such as mitogen-activated protein kinases (MAPK), which are also induced by HCV proteins. These synergistic effects may be responsible for rare characteristics, that is, the high incidence and multicentric nature of hepatocarcinogenesis in HCV infection.

© 2005 Elsevier Ireland Ltd. All rights reserved.

*Keywords:* Oxidative stress; Hepatitis C virus; Hepatocarcinogenesis; Lipid peroxidation; Steatosis; Insulin resistance

## Contents

1. Introduction .....	66
2. Oxidative stress, reactive oxygen, and the liver .....	66
2.1. Oxidative stress and reactive oxygen .....	66
2.2. Antioxidation system and oxidative stress markers .....	66
3. Viral infection and oxidative stress .....	67
3.1. ROS production associated with viral infection .....	67
3.2. Nitric oxide production associated with viral infection .....	67
4. Oxidative stress caused by viral protein .....	68
5. Relationship of HCV infection with insulin resistance .....	68
6. Relationship of HCV infection with hepatic steatosis .....	69
7. Iron and reactive oxygen .....	69
8. Interactions with alcohol .....	69
9. Hepatocarcinogenesis and oxidative stress .....	69
10. Conclusions .....	70
References .....	71

\* Corresponding author at: Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan. Tel.: +81 3 5800 8800; fax: +81 3 5800 8799.

*E-mail address:* [kkoike-tyk@umin.ac.jp](mailto:kkoike-tyk@umin.ac.jp) (K. Koike).

## 1. Introduction

There are approximately 200 million people infected with hepatitis C virus (HCV) worldwide, of which about 1.8 million are in Japan. It is one of the most serious causes of liver disease. It was reported that approximately 70% of those with HCV infection suffer from persistent infection, causing active or inactive chronic hepatitis and that about 30% of patients with chronic hepatitis are assumed to develop cirrhosis within their lifetime. Once HCV infection develops into cirrhosis, hepatocellular carcinoma (HCC) develops at an annual rate of 5–7% [1]. The strong association of oxidative stress with HCV infection has been demonstrated recently and it has become possible to explain at least part of the clinical progression of the disease. The pathogenesis of chronic hepatitis C is not merely ascribed to inflammation caused by viral infection, but the role of viral proteins in the pathogenesis was also reported [2–4]. Of proteins constituting HCV, the core protein, in particular, has various functions with respect to host cells [5] and is closely related to oxidative stress. In this overview, the relationship between HCV infection and oxidative stress is reviewed focusing on the pathological effect of the core protein of HCV, and the significance of oxidative stress in the pathogenesis of liver disease will be discussed.

## 2. Oxidative stress, reactive oxygen, and the liver

### 2.1. Oxidative stress and reactive oxygen

The main source of reactive oxygen species (ROS) in hepatocytes is the mitochondria. Outside of hepatocytes, ROS also originate from nicotinamide adenine dinucleotide phosphate

(NADPH) oxidase and xanthine oxidase in Kupffer cells and inflammatory cells. Several percent of consumed oxygen is constantly converted into ROS in the mitochondria accompanied by oxygen consumption in the electron transport system (ETS, Fig. 1). Hepatocytes contain many mitochondria and therefore have a high ROS production. Generated ROS are very unstable and highly reactive, and attack biomolecules such as DNA, lipids, and proteins. The liver not only produces much ROS but is also the center of the anti-oxidative effect in the form of protein synthesis. Oxidative stress refers to the oxidation-reaction-dominant state of the living body induced by an imbalance between the oxidation reaction caused by ROS and the anti-oxidation reaction. Main ROS include superoxide ( $^{\circ}\text{O}_2^-$ ), hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) and the hydroxyl radical ( $\text{HO}^{\circ}$ ). ROS are mainly produced from  $^{\circ}\text{O}_2^-$  and converted into stable  $\text{H}_2\text{O}_2$  through dismutation reaction.  $\text{H}_2\text{O}_2$  is converted into highly reactive  $\text{HO}^{\circ}$  in the presence of a transition metal.

### 2.2. Antioxidation system and oxidative stress markers

Antioxidants include glutathione (GSH), thioredoxin (TRX), vitamin E, vitamin C, and  $\beta$ -carotene. Reactive oxygen elimination enzymes include superoxide dismutase (SOD), glutathione peroxidase (GPx), and catalase. SOD is induced by oxidative stress and dismutates  $^{\circ}\text{O}_2^-$  to  $\text{H}_2\text{O}_2$  and oxygen. GSH is a compound belonging to the SH group and is highly abundant in the living body, and the SH group provides electrons to free radicals to stabilize the radicals. GSH exists in a reduced form in cells. Because it is converted into dimeric oxidized glutathione (GSSG) and becomes stable after donating electrons, GSSG prevents free radicals from continuously scrambling for electrons. GPx decomposes  $\text{H}_2\text{O}_2$  into water




Fig. 1. The electron transfer system (ETS) of the mitochondrion. Most of the oxygen, consumed by mammalian cells, is converted to water via the mitochondrial ETS. However, up to 5% of the electrons entering the mitochondrial ETS can become uncoupled and singly leak out onto oxygen to form superoxide. Therefore, if there is impairment in the mitochondrial ETS function, it can be a cause of the overproduction of reactive oxygen species (ROS). TCA, tricarboxylic acid; NADH, nicotinamide adenine dinucleotide phosphate; FMN, flavine mononucleotide; CoQ, coenzyme Q; cyt c, cytochrome c.




Fig. 2. Generation and scavenging of oxidative stress. SOD, super-oxide dismutase; GSH, reduced glutathione; GPx, glutathione peroxidase; TRX, thioredoxin.

and oxygen with GSH as an electron donor and reduces lipid peroxide to become neutralized. GSSG is converted back to GSH when glutathione reductase transfers an electron from NADPH to GSSG. Catalase in peroxisomes also decomposes  $H_2O_2$  to water and oxygen. TRX is also a protein induced by oxidative stress, and is reduced via the S–S binding of the substrate protein by two SH groups in TRX and acts on the  $H_2O_2$  elimination system via peroxiredoxins (Fig. 2).

ROS cause various forms of cellular damage. 4-Hydroxy-2-nonenal (HNE) and malondialdehyde (MDA) are the peroxidation reaction products of lipids, and 8-hydroxydeoxyguanosine (8-OHdG) is the product of DNA base modification (Fig. 3). These products serve as oxidative stress markers.




Fig. 3. Representative aldehyde species generated via lipid peroxidation reaction.

### 3. Viral infection and oxidative stress

#### 3.1. ROS production associated with viral infection

Upon viral infection, ROS are produced by NADPH oxidase and xanthine oxidase in neutrophils and macrophages. In particular, NS3, one of the non-structural proteins of HCV, was reported to induce ROS production by NADPH oxidase in neutrophils [6]. Furthermore, in viral hepatitis, ROS are also produced in hepatocytes through the release of inflammatory cytokines such as  $TNF-\alpha$  and  $IL-1\beta$  from inflammatory cells. Increased hepatic or serum 8-OHdG, HNE and MDA levels are observed in chronic hepatitis C, indicating an increase in ROS production [7–13]. Findings that indicate an increase in the activity of the ROS elimination system including decreased hepatic and blood GSH levels, an increased GSSG/GSH + GSSG ratio, and an increased serum TRX level have been reported [13–16]. The findings of markedly decreased HNE level following viral eradication with interferon [12] and decreased serum ALT and TRX levels following the administration of vitamin E, an antioxidant [17], also demonstrated that oxidative stress plays an important role in chronic hepatitis C.

#### 3.2. Nitric oxide production associated with viral infection

In the presence of an inflammation, inducible nitric oxide synthase (iNOS) is induced in macrophages and hepatocytes by  $TNF-\alpha$  and  $IFN-\gamma$  [18–20]. Other investigators reported that protein kinase (PKR) activated by double-stranded RNA formed during virus reproduction in turn activates the transcripts of  $NF-\kappa B$  and IRF-1 to induce iNOS [21]. In the case of HCV, it was reported that its constituent proteins (E2 and non-structural (NS) protein 5A) inhibit PKR activity [22,23], but iNOS induction by viral RNA via PKR is also suspected. Indeed, iNOS synthesis correlates with intrahepatic viral load in chronic hepatitis C [24].

NO is generally synthesized as a non-specific defense reaction to infectious diseases; however, in viral infection, antiviral activity may be present or absent in various viral types [20]. NO is reported to exhibit no antiviral activity against a tick-borne encephalitis virus (TBE-V), flavivirus [25], and NO may also have no antiviral activity against HCV. On the contrary, NO causes cellular damage upon its reaction to  $O_2$  or simultaneously produced  $O_2^{\cdot-}$  (reactive nitrogen species, RNS). Upon reaction to  $O_2^{\cdot-}$ , in particular, NO acts as a strong oxidant with the generation of peroxynitrous acid ( $ONOO^-$ ), and  $ONOO^-$  also produces nitrotyrosine through the nitration of aromatic amino acid residues in the presence of a transition metal. Nitrotyrosine accumulation was observed in correlation to inflammation severity in chronic hepatitis C tissue [26]; suggesting that the production of both NO and ROS increased. ROS and RNS are produced as defense factors for biological viral clearance, but these factors also have cytotoxic effects that

are assumed to contribute to the exacerbation of the disease state.

#### 4. Oxidative stress caused by viral protein

The HCV genome comprises the genes of four structural proteins and six non-structural proteins (Fig. 4), and it has been reported that at least two viral proteins cause oxidative stress in cells. The core protein, a structural protein, was found to have various actions, including the induction of oxidative stress and accumulation of lipids, in experimental studies using cultured cells and transgenic mice [2,27]. Experiments using mice transgenic for the core gene showed an increased ROS production, an increased intrahepatic catalase activity, a decreased intrahepatic GSH level and a decreased GSH/GSH – GSSG ratio indicating an anti-oxidation effect inhibition, although there was no increase in serum ALT level nor a histological finding of hepatitis. Increased levels of intrahepatic peroxide lipids in the core gene transgenic mice with aging as compared with those in the control mice also indicate increased oxidative stress. As a mechanism underlying oxidative stress induction by the core protein, mitochondrial damage is considered. Morphological abnormalities of the mitochondria were observed in the core gene transgenic mouse liver [2], and an increased ROS production caused by damage of the mitochondrial electron transport system was noted in core-protein-expressing cells [27]. Mitochondrial DNA, which has no protective proteins such as histone, is susceptible to damage by ROS [28,29]. Mitochondrial DNA in the core gene transgenic mice showed damage as early as 3-months old. This mitochondrial damage disrupts the synthesis of proteins constituting the electron transport system complex and could also increase oxidative stress caused by damage of the electron transport system.

A study using a cell culture system demonstrated that non-structural protein 5A (NS5A) also causes oxidative stress. NS5A induces endoplasmic reticulum calcium release via

endoplasmic reticulum stress, and this leads to an increased ROS production in the mitochondria [4]. Although the effect of NS5A has not been confirmed yet by other study groups, HCV has the direct action of increasing intracellular ROS production via its proteins, separate from oxidative stress induction by inflammation caused by viral infection. A report that oxidative stress was also observed in HCV carriers with a normal ALT level [13] indicates that it is caused by a direct oxidative stress induction without being mediating inflammatory reactions.

#### 5. Relationship of HCV infection with insulin resistance

The relationship of HCV infection with insulin resistance and type 2 diabetes has been suggested epidemiologically [30–32]. Insulin resistance was also observed in core gene transgenic mice before the onset of hepatic steatosis [33]. A disrupted tyrosine phosphorylation of the insulin receptor substrate (IRS-1) was observed in the liver of these transgenic mice. The analysis of hepatic tissues in patients with chronic hepatitis C not complicated by diabetes showed that insulin receptor and IRS-1 expression levels are elevated in patients with HCV infection, whereas the tyrosine phosphorylation of IRS-1 induced by insulin is inhibited. An excessive oxidative stress may be another potential cause of this insulin resistance. Oxidative stress indirectly blocks the phosphorylation of tyrosine residues of insulin receptors and IRS-1 and inhibits insulin signaling [34].

These reported results thus indicate an insulin signaling disorder in the liver infected with HCV [35]. There has been no report to date directly proving that hepatic insulin signaling disorder in patients with HCV infection is attributable to oxidative stress. However, because diabetes, which is the state of having abnormally high blood sugar levels that cannot be self-regulated by individual organisms, also induces oxidative stress [34], the close relationship between insulin resistance or diabetes and oxidative stress as the cause and the




Fig. 4. Structure of hepatitis C virus genome. The genome of HCV consists of two parts, structural and non-structural regions. The former comprises the core and envelope regions, and the latter consists of NS2 to NS5a, which regions chiefly code enzymes necessary for viral replication. NS, non-structural; ISDR, interferon sensitivity-determining region.



result, respectively, is a very interesting issue to investigate in the future.

## 6. Relationship of HCV infection with hepatic steatosis

Hepatic steatosis is frequently observed in patients with HCV infection. The relationships of HCV infection with intrahepatic viral loads and core protein levels, different prevalence of hepatic steatosis by viral genotype [higher incidence for genotype 3a], and improved steatosis following viral eradication were reported [36–38]. It is presumed from these reports that HCV itself causes hepatic steatosis. A similar hepatic steatosis caused even by the core protein alone was observed in a study using an expression system in cultured cells and transgenic mice, and it was thus suggested that the core protein plays a significant role in hepatic steatosis as the direct action of HCV [39,40]. Hyperinsulinemia induced by insulin resistance mentioned above causes the overloading of the liver with fatty acids from fat cells, and mitochondrial damage inhibits the  $\beta$ -oxidation of fatty acids [41]. Furthermore, the core protein was reported to inhibit microsomal triglyceride transfer protein (MTP) activity that is required when neutral fat is released as very low-density lipoproteins (VLDLs) [42]. All these actions could cause hepatic steatosis. In the liver of non-alcoholic steatohepatitis (NASH) patients, it was reported that  $\beta$ -oxidation in the mitochondria and peroxisomes or the metabolism of fatty acids by cytochrome P450 2E1 (CYP2E1) in microsomes is promoted under an excessive load of fatty acids, resulting in ROS production [43,44]. In HCV infection as well, intrahepatic fat accumulation possibly increases ROS production as in NASH. Because hepatic steatosis in chronic hepatitis C was reported to be a factor for disease progression [45–47], increased oxidative stress associated with hepatic steatosis is presumably involved in disease progression.

## 7. Iron and reactive oxygen

The iron content in the liver and spleen is high, and transition metals facilitate electron transfer and play an important role in the production of free radicals. Iron in combination with transferrin and ferritin is stable, but an unstable iron ion is freed when ferritin is decomposed by lysosomes [48]. ROS additionally promote iron release from ferritin [49]. A free iron ion catalyzes changes from relatively poor reactive  $O_2^-$  and  $H_2O_2$  to a highly reactive  $HO^\bullet$  (Fenton reaction) [50,51].  $HO^\bullet$  oxidizes membrane phospholipids, which compose cells and intracellular organelles, and iron forms radicals from produced peroxide lipids, thereby enhancing lipid peroxidation. Iron site-specifically combines with DNA and promotes DNA damage caused by ROS. Iron also increases ROS production by CYP2E1 [52]. A report that an enhanced peroxidation of intrahepatic lipids is attenuated by exsan-

guination in hemochromatosis also supports the involvement of iron in oxidative stress [53].

An excessively high iron content in the liver was observed in chronic hepatitis C [8,54]. Other investigators reported that iron removal therapy by exsanguination of chronic hepatitis C patients significantly improves serum ALT level without affecting viral load [55–57]. Another study showed that hepatic impairment is exacerbated following the administration of iron to chimpanzees with chronic hepatitis C [58]. Furthermore, oxidative stress is decreased by the iron removal therapy for chronic hepatitis C using intrahepatic 8-OHdG level as an index [57]. The above-mentioned reports show the close relationships of chronic hepatitis C with iron metabolism and oxidative stress.

## 8. Interactions with alcohol

Alcohol metabolism plays an important role in ROS production. Mainly alcohol dehydrogenase (ADH) in the cytosol and CYP2E1 (microsomal ethanol-oxidizing system) in microsomes are responsible for alcohol metabolism in the liver. When alcohol dehydrogenase oxidizes ethanol to acetaldehyde, the reduction from  $NAD^+$  to NADH simultaneously occurs. NADH accumulation causes stress on the mitochondrial electron transfer system, leading to an increased production of ROS [59]. NADH also inhibits xanthine dehydrogenase activity, and xanthine is thereby oxidized by xanthine oxidase with the production of ROS [60]. CYP2E1 is induced by chronic alcohol intake and ROS are produced when CYP2E1 oxidizes ethanol to acetaldehyde [52,61].

There is no significant difference in hepatic peroxide level between core gene transgenic mice at 3–6-months old and control transgenic mice, but hepatic peroxide level significantly increases following the administration of a low dose of alcohol in the core gene transgenic mice [2]. ROS production increases upon glutathione reduction in HepG2 cells, with the co-expression of the core protein and CYP2D1, the latter of which is induced by alcohol [62]. These findings show that the core protein and alcohol in combination increase oxidative stress. Indeed, it was reported that alcohol intake plays a role in promoting the progression of chronic hepatitis C [63,64] and that increased levels of oxidative stress markers such as HNE and lipid hydroperoxide also support these findings [65]. From the viewpoint of oxidative stress also, HCV infection and alcohol intake are both considered to promote hepatic impairment.

## 9. Hepatocarcinogenesis and oxidative stress

It has been demonstrated that oxidative stress plays a key role in carcinogenesis [66,67]. Animal experiments using hepatocarcinogenesis models with the administration of a chemical substance (diethyl-nitrosamine, peroxisome proliferators) and with the administration of a choline-deficient

amino acid diet also indicates the involvement of oxidative stress [68–72]. In Long Evans Cinnamon (LEC) rats, an animal model that spontaneously develops heritable hepatitis and HCC caused by an abnormal copper accumulation, a congenitally decreased glutathione peroxidase expression level was reported, and the close relationship between oxidative stress and hepatocarcinogenesis was indicated [73].

The epidemiological relationship between HCV infection and HCC is evident [74,75], but the mechanism underlying this relationship has not been fully elucidated yet. Among postulated hypotheses on the mechanism of HCV-associated hepatocarcinogenesis, that of the involvement of the viral protein, in particular, the core protein of HCV is attractive: HCC develops in core gene transgenic mice, and carcinogenesis starts with well-differentiated carcinoma with an excessively high fat content, similar to hepatocarcinogenesis in human chronic hepatitis C, and poorly differentiated carcinoma with a low fat content develops in the form of “nodules in nodules” [76]. Because oxidative stress is increased in the core gene transgenic mice as mentioned above, it is assumed that oxidative stress plays an important role in hepatocarcinogenesis in chronic hepatitis C. Because the development of HCC is also observed in transgenic mice carrying the full-length HCV protein gene, the non-structural protein may have an additive effect to the effect of the structural proteins including the core protein, contributing to hepatocarcinogenesis [77]. NS5A, which was also reported to induce ROS production [4], may also contribute to hepatocarcinogenesis, although ROS induction by NS5A is not unequivocally confirmed yet.

Mitochondrial DNA has no potent protective proteins such as histone and is near the electron transport system, the major ROS production site. Hence, it is 10 to 15 times more susceptible to mutation caused by ROS than nuclear DNA [28,29]. In an investigation of mitochondrial DNA mutation in the human normal liver, both cancerous and non-cancerous liver tissues in patients with HCC showed very high incidences of DNA mutations [78]; thus, a relationship between oxidative stress persistence and hepatocarcinogenesis is suggested.

In the core protein expression system in the hepatic tissue and cultured cells of core gene transgenic mice, the activation of transcription factor AP-1 via mitogen-activated protein (MAP) kinase was observed [79–83]. The activation of the transcription factors AP-1, NF- $\kappa$ B, and signal transducer and activator of transcription (STAT) 3 by NS5A were also reported [4,84]. The activation of these transcription factors may facilitate cell proliferation, contributing to tumorigenic transformation.

It was also reported that ROS facilitate apoptosis via c-Jun N-terminal kinase (JNK)/p38 MAP kinase or by directly attacking the mitochondria. Apoptosis is a protective mechanism of the host against viral infection and carcinogenesis. Some reports stated that the core protein facilitates apoptosis [85–88], whereas other reports stated that the core protein inhibits apoptosis [89–92]; thus, no fixed view has yet been established. If it indeed inhibits apoptosis, it is assumed that this inhibition proceeds by maintaining oxidative stress and

that the core protein has a beneficial effect against carcinogenesis and persistent viral infection.

In HCV infection, viral proteins such as the core protein and, possibly, NS5A protein induce oxidative stress, intracellular signaling, and transcription factors, which are not reflected in blood ALT level, contributing to the progression of carcinogenesis. Carcinogenesis, however, is slow as is observed in humans and core gene transgenic mice, the latter of which developed HCC in the latter half of their life. Recently, Okanoue et al. reported a long-term follow-up study of subjects with persistent HCV infection who had persistently normal ALT levels (PNAL) [93]. In their study, serum thioredoxin levels were not elevated in those with PNAL compared to those with chronic hepatitis. This may apparently seem contradictory to the results of our above-mentioned animal model studies. However, we should realize that anti-oxidant system is also instrumental in the liver. In these relatively younger people with PNAL than those with CH [93], active anti-oxidant system may erase the apparent elevation of ROS. Such a phenomenon was described in a mouse model by Moriya et al. [2], in which ROS was apparently normal in young core gene transgenic mice with the activation of catalase and reduction of GSH. Clinically, the presence of inflammation is thought to facilitate the process of hepatocarcinogenesis.

## 10. Conclusions

A very close pathological relationship between oxidative stress and HCV infection is observed, as shown by the above overview of relevant publications and discussion. The causes of oxidative stress in HCV infection are considered to include various factors such as mitochondrial damage, endoplasmic reticulum stress, iron accumulation, and lipid accumulation in the liver. Various study results demonstrated that even only viral proteins, mainly the HCV core protein, cause oxidative stress. When inflammation via immunoreactions to viral infection is added to oxidative stress, ROS production is expected to further increase, leading to a state in which the anti-oxidation system cannot cope with. In this sense, inflammation in chronic hepatitis C is considered to be qualitatively different from inflammation observed in other types of hepatitis such as autoimmune hepatitis or hepatitis B [94] (Fig. 5). As a treatment of chronic hepatitis C, the eradication of the virus is ideal. If it is not possible, however, the control of factors that exacerbate oxidative stress, such as inflammation via immune reaction and alcohol, and the relief of oxidative stress by the iron removal therapy and the administration of an anti-oxidation agent are considered to delay the progression of chronic hepatitis.

The development of such new anti-oxidation agents is being awaited. In further studies on the development of new therapies for hepatitis C and control methods for hepatocarcinogenesis in the future, the importance of those focusing on oxidative stress is expected to markedly increase.




Fig. 5. Oxidative stress and hepatocarcinogenesis in various types of hepatitis (hypothesis). Oxidative stress is generated in all types of hepatitis via inflammation accompanied by continual cell death and regeneration. In HCV infection, HCV itself causes the production of oxidative stress in a synergy with inflammation. In this sense, the quality of “inflammation” in HCV infection may be different from that in other types of hepatitis. Additional impact of HCV proteins on the intracellular signal transduction would provoke the development of HCC. These may explain the conspicuous properties of HCC development.

## References

- [1] Ikeda K, Saitoh S, Suzuki Y, et al. Disease progression and hepatocellular carcinogenesis in patients with chronic viral hepatitis: a prospective observation of 2215 patients. *J Hepatol* 1998;28:930–8.
- [2] Moriya K, Nakagawa K, Santa T, et al. Oxidative stress in the absence of inflammation in a mouse model for hepatitis C virus-associated hepatocarcinogenesis. *Cancer Res* 2001;61:4365–70.
- [3] Koike K, Moriya K, Ishibashi K, et al. Sialadenitis histologically resembling Sjögren syndrome in mice transgenic for hepatitis C virus envelope genes. *Proc Natl Acad Sci USA* 1997;94:233–6.
- [4] Gong G, Waris G, Tanveer R, Siddiqui A. Human hepatitis C virus NS5A protein alters intracellular calcium levels, induces oxidative stress, and activates STAT-3 and NF- $\kappa$ B. *Proc Natl Acad Sci USA* 2001;98:9599–604.
- [5] Suzuki R, Suzuki T, Ishii K, Matsuura Y, Miyamura T. Processing and functions of hepatitis C virus proteins. *Intervirology* 1999;42:145–52.
- [6] Bureau C, Bernad J, Chauouche N, et al. Non-structural 3 protein of hepatitis C virus triggers an oxidative burst in human monocytes via activation of NADPH oxidase. *J Biol Chem* 2001;276:23077–83.
- [7] Shimoda R, Nagashima M, Sakamoto M, et al. Increased formation of oxidative DNA damage, 8-hydroxydeoxyguanosine, in human livers with chronic hepatitis. *Cancer Res* 1994;54:3171–2.
- [8] Farinati F, Cardin R, Maria ND, et al. Iron storage, lipid peroxidation and glutathione turnover in chronic anti-HCV positive hepatitis. *J Hepatol* 1995;22:449–56.
- [9] De Maria N, Calantoni A, Fagivoli S, et al. Association between reactive oxygen species and disease activity in chronic hepatitis C. *Free Radic Biol Med* 1996;21:291–5.
- [10] Romero MJ, Bosch-Morell F, Romero B, Rodrigo JM, Serra MA, Romero FJ. Serum malondialdehyde: possible use for the clinical management of chronic hepatitis C patients. *Free Radic Biol Med* 1998;25:993–7.
- [11] Farinati F, Cardin R, Degan P, et al. Oxidative DNA damage in circulating leukocytes occurs as an early event in chronic HCV infection. *Free Radic Biol Med* 1999;27:1284–91.
- [12] Kageyama F, Kobayashi Y, Kawasaki T, Toyokuni S, Uchida K, Nakamura H. Successful interferon therapy reverses enhanced hepatic iron accumulation and lipid peroxidation in chronic hepatitis C. *Am J Gastroenterol* 2000;95:1041–50.
- [13] Vendemiale G, Grattagliano I, Portincasa P, Serviddio G, Palasciano G, Altomare E. Oxidative stress in symptom-free HCV carriers: relation with ALT flare-up. *Eur J Clin Invest* 2001;31:54–63.
- [14] Barbaro G, Di Lorenzo G, Ribersani M, et al. Serum ferritin and hepatic glutathione concentrations in chronic hepatitis C patients related to the hepatitis C virus genotype. *J Hepatol* 1999;30:774–82.
- [15] Sumida Y, Nakashima T, Yoh T, et al. Serum thioredoxin levels as an indicator of oxidative stress in patients with hepatitis C virus infection. *J Hepatol* 2000;33:616–22.
- [16] Jain SK, Pemberton PW, Smith A, et al. Oxidative stress in chronic hepatitis C: not just a feature of late stage disease. *J Hepatol* 2002;36:805–11.
- [17] Mahmood S, Yamada G, Niiyama G, et al. Effect of vitamin E on serum aminotransferase and thioredoxin levels in patients with viral hepatitis C. *Free Radic Res* 2003;37:781–5.
- [18] Schwarz KB. Oxidative stress during viral infection: a review. *Free Radic Biol Med* 1996;21:641–9.
- [19] Majano PL, Garcia-Monzon C, Lopez-Cabrera M, et al. Inducible nitric oxide synthase expression in chronic viral hepatitis. *J Clin Invest* 1998;101:1343–52.
- [20] Akaike T. Role of free radicals in viral pathogenesis and mutation. *Rev Med Virol* 2001;11:87–101.
- [21] Uetani K, Der SD, Zamanian-Daryoush M, et al. Central role of double-stranded RNA-activated protein kinase in microbial induction of nitric oxide synthase. *J Immunol* 2000;165:988–96.
- [22] Gale Jr MJ, Korth MJ, Tang NM, et al. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the non-structural 5A protein. *Virology* 1997;230:217–27.
- [23] Taylor DR, Shi ST, Romano PR, Barber GN, Lai MM. Inhibition of the interferon-inducible protein kinase PKR by HCV E2 protein. *Science* 1999;285:107–10.
- [24] Mihm S, Fayyazi A, Ramadori G. Hepatic expression of inducible nitric oxide synthase transcripts in chronic hepatitis C virus infection.

- tion: relation to hepatic viral load and liver injury. *Hepatology* 1997;26:451–8.
- [25] Kreil TR, Eibl MM. Nitric oxide and viral infection: no antiviral activity against a flavivirus in vitro, and evidence for contribution to pathogenesis in experimental infection in vivo. *Virology* 1996;219:304–6.
- [26] Garcia-Monzon C, Majano PL, Zubia I, Sanz P, Apolinario A, Moreno-Otero R. Intrahepatic accumulation of nitrotyrosine in chronic viral hepatitis is associated with histological severity of liver disease. *J Hepatol* 2000;32:331–8.
- [27] Okuda M, Li K, Beard MR, et al. Mitochondrial injury, oxidative stress, and antioxidant gene expression are induced by hepatitis C virus core protein. *Gastroenterology* 2002;122:366–75.
- [28] Shigenaga MK, Hagen TM, Ames BN. Oxidative damage and mitochondrial decay in aging. *Proc Natl Acad Sci USA* 1994;91:10771–8.
- [29] Lee CM, Weindruch R, Aiken JM. Age-associated alterations of the mitochondrial genome. *Free Radic Biol Med* 1997;22:1259–69.
- [30] Caronia S, Taylor K, Pagliaro L, et al. Further evidence for an association between non-insulin-dependent diabetes mellitus and chronic hepatitis C virus infection. *Hepatology* 1999;30:1059–63.
- [31] Alexander GJM. An association between hepatitis C virus infection and type 2 diabetes mellitus: what is the connection? *Ann Intern Med* 2000;133:650–2.
- [32] Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL. Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 2000;133:592–9.
- [33] Shintani Y, Fujie H, Miyoshi H, et al. Hepatitis C virus infection and diabetes: direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840–8.
- [34] Evans JL, Goldfine ID, Maddux BA, Grodsky GM. Are oxidative stress-activated signaling pathways mediators of insulin resistance and  $\beta$ -cell dysfunction? *Diabetes* 2003;52:1–8.
- [35] Aytug S, Reich D, Sapiro LE, Bernstein D, Begum N. Impaired IRS-1/PI3-kinase signaling in patients with HCV: a mechanism for increased prevalence of type 2 diabetes. *Hepatology* 2003;38:1384–92.
- [36] Fujie H, Yotsuyanagi H, Moriya K, et al. Steatosis and intrahepatic hepatitis C virus in chronic hepatitis. *J Med Virol* 1999;59:141–5.
- [37] Rubbia-Brandt L, Quadri R, Abid K, et al. Hepatocyte steatosis is a cytopathic effect of hepatitis C virus genome 3. *J Hepatol* 2000;33:106–15.
- [38] Kumar D, Farrell GC, Fung C, George J. Hepatitis C virus genotype 3 is cytopathic in hepatocytes. Genotype-specific reversal of hepatic steatosis after sustained response to antiviral therapy. *Hepatology* 2002;36:1266–72.
- [39] Moriya K, Yotsuyanagi H, Shintani Y, et al. Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J Gen Virol* 1997;78:1527–31.
- [40] Barba G, Harper F, Harada T, et al. Hepatitis C virus core protein shows a cytoplasmic localization and associates to cellular lipid storage droplets. *Proc Natl Acad Sci USA* 1997;94:1200–5.
- [41] Fromenty B, Pessayre D. Inhibition of mitochondrial  $\beta$ -oxidation as a mechanism of hepatotoxicity. *Pharmacol Ther* 1995;67:101–54.
- [42] Perlemuter G, Sabile A, Letteron P, et al. Hepatitis C virus core protein inhibits microsomal triglyceride transfer protein activity and very low density lipoprotein secretion: a model of viral-related steatosis. *FASEB J* 2002;16:185–94.
- [43] Day CP, James OFW. Steatohepatitis: a tale of two “hits”? *Gastroenterology* 1998;114:842–5.
- [44] Weltman MD, Farrell GC, Hall P, Ingelman-Sundberg M, Liddle C. Hepatic cytochrome P450 2E1 is increased in patients with non-alcoholic steatohepatitis. *Hepatology* 1998;27:128–33.
- [45] Czaja AJ, Carpenter HA, Santrach PJ, Moore SB. Host- and disease-specific factors affecting steatosis in chronic hepatitis C. *J Hepatol* 1998;29:198–206.
- [46] Hourigan LF, Macdonald GA, Purdie D, et al. Fibrosis in chronic hepatitis C correlates significantly with body mass index and steatosis. *Hepatology* 1999;29:1215–9.
- [47] Serfaty L, Poujol-Robert A, Carbonell N, Chazouilleres O, Poupon RE, Poupon R. Effect of the interaction between steatosis and alcohol intake on liver fibrosis progression in chronic hepatitis C. *Am J Gastroenterol* 2002;97:1807–12.
- [48] Radisky DC, Kaplan J. Iron in cytosolic ferritin can be recycled through lysosomal degradation in human fibroblasts. *Biochem J* 1998;336:201–5.
- [49] Kakhlon O, Cabantchik ZI. The labile iron pool: characterization, measurement, and participation in cellular processes. *Free Radic Biol Med* 2002;33:1037–46.
- [50] Halliwell B, Gutteridge JMC. Oxygen toxicity, oxygen radicals, transition metals and disease. *Biochem J* 1984;219:1–14.
- [51] Aust SD, Morehouse LA, Thomas CE. Roles of metals in oxygen radical reactions. *Free Radic Biol Med* 1985;1:3–25.
- [52] Cederbaum AI. Iron CYP2E1-dependent oxidative stress and toxicity. *Alcohol* 2003;30:115–20.
- [53] Houghlum K, Ramm G, Crawford DH, Witztum JL, Powell LW, Chojkier M. Excess iron induces hepatic oxidative stress and transforming growth factor  $\beta$ 1 in genetic hemochromatosis. *Hepatology* 1997;26:605–10.
- [54] Bonkovsky H, Banner BF, Rothman AL. Iron and chronic viral hepatitis. *Hepatology* 1997;25:759–68.
- [55] Hayashi H, Takikawa T, Nishimura N, Yano M, Isomura T, Sakamoto N. Improvement of serum aminotransferase levels after phlebotomy in patients with chronic active hepatitis C and excess hepatic iron. *Am J Gastroenterol* 1994;89:986–8.
- [56] Sartori M, Andorno S, Rigamonti C, Boldorini R. Chronic hepatitis C treated with phlebotomy alone: biochemical and histological outcome. *Dig Liver Dis* 2001;33:157–62.
- [57] Kato J, Kobune M, Nakamura T, et al. Normalization of elevated hepatic 8-hydroxy-2'-deoxyguanosine levels in chronic hepatitis C patients by phlebotomy and low iron diet. *Cancer Res* 2001;61:8697–702.
- [58] Bassett SE, Bisceglie AMD, Bacon BR, et al. Effects of iron loading on pathogenicity in hepatitis C virus-infected chimpanzees. *Hepatology* 1999;29:1884–92.
- [59] Cunningham CC, Bailey SM. Ethanol consumption and liver mitochondrial function. *Biol Signals Recept* 2001;10:271–82.
- [60] Kato S, Kawase T, Alderman J, et al. Role of xanthine oxidase in ethanol-induced lipid peroxidation in rats. *Gastroenterology* 1990;98:203–10.
- [61] Ingelman-Sundberg M, Johansson I. Mechanisms of hydroxyl radical formation and ethanol oxidation by ethanol-inducible and other forms of rabbit liver microsomal cytochrome P-450. *J Biol Chem* 1984;259:6447–58.
- [62] Wen F, Abdalla MY, Aloman C, et al. Increased prooxidant production and enhanced susceptibility to glutathione depletion in HepG2 cells co-expressing HCV core protein and CYP2E1. *J Med Virol* 2004;72:230–40.
- [63] Ostapowicz G, Watzon KJR, Locarnini SA, Desmond PV. Role of alcohol in the progression of liver disease caused by hepatitis C virus infection. *Hepatology* 1998;27:1730–5.
- [64] Hassan MM, Hwang LY, Hatten CJ, et al. Risk factors for hepatocellular carcinoma: synergism of alcohol with viral hepatitis and diabetes mellitus. *Hepatology* 2002;36:1206–13.
- [65] Rigamonti C, Mottaran E, Reale E, et al. Moderate alcohol consumption increases oxidative stress in patients with chronic hepatitis C. *Hepatology* 2003;38:42–9.
- [66] Cerutti PA, Trump BF. Inflammation and oxidative stress in carcinogenesis. *Cancer Cells (Cold Spring Harbor)* 1991;3:1–7.
- [67] Dreher D, Junod AF. Role of oxygen free radicals in cancer development. *Eur J Cancer* 1996;32:30–8.

- [68] Perera MI, Betschart JM, Virji MA, Katyal SL, Shinozuka H. Free radical injury and liver tumor promotion. *Toxicol Pathol* 1987;15:51–9.
- [69] Reddy JK, Rao MS. Oxidative DNA damage caused by persistent peroxisome proliferation: its role in hepatocarcinogenesis. *Mutat Res* 1989;214:63–8.
- [70] Nakae D, Kobayashi Y, Akai H, et al. Involvement of 8-hydroxyguanine formation in the initiation of rat liver carcinogenesis by low dose levels of *N*-nitrosodiethylamine. *Cancer Res* 1997;57:1281–7.
- [71] Nakae D, Kotake Y, Kishida H, et al. Inhibition by phenyl *N*-tert-butyl nitron on early phase carcinogenesis in the livers of rats fed a choline-deficient, L-amino acid-defined diet. *Cancer Res* 1998;58:4548–51.
- [72] Floyd RA, Kotake Y, Hensley K, Nakae D, Konishi Y. Reactive oxygen species in choline deficiency induced carcinogenesis and nitron inhibition. *Mol Cell Biochem* 2002;234–235:195–203.
- [73] Suemizu H, Yoshimura S, Takeichi N, Moriuchi T. Decreased expression of liver glutathione peroxidase in Long-Evans cinnamon mutant rats predisposed to hepatitis and hepatoma. *Hepatology* 1994;19:694–700.
- [74] Saito I, Miyamura T, Ohbayashi A, et al. Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990;87:6547–9.
- [75] Simonetti RG, Camma C, Fiorello F, et al. Hepatitis C virus infection as a risk factor for hepatocellular carcinoma in patients with cirrhosis. *Ann Intern Med* 1992;116:97–102.
- [76] Moriya K, Fujie H, Shintani Y, et al. The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065–7.
- [77] Lerat H, Honda M, Beard MR, et al. Steatosis and liver cancer in transgenic mice expressing the structural and non-structural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352–65.
- [78] Nishikawa M, Nishiguchi S, Shiomi S, et al. Somatic mutation of mitochondrial DNA in cancerous and non-cancerous liver tissue in individuals with hepatocellular carcinoma. *Cancer Res* 2001;61:1843–5.
- [79] Shrivastava A, Manna SK, Ray R, Aggarwal BB. Ectopic expression of hepatitis C virus core protein differentially regulates nuclear transcription factors. *J Virol* 1998;72:9722–8.
- [80] Kato N, Yoshida H, Kioko Ono-Nita S, et al. Activation of intracellular signaling by hepatitis B and C viruses: C-viral core is the most potent signal inducer. *Hepatology* 2000;32:405–12.
- [81] Erhardt A, Hassan M, Heintges T, Haussinger D. Hepatitis C virus core protein induces cell proliferation and activates ERK, JNK, and p38 MAP kinases together with the MAP kinase phosphatase MKP-1 in a HepG2 Tet-Off cell line. *Virology* 2002;292:272–84.
- [82] Tsutsumi T, Suzuki T, Moriya K, et al. Alteration of intrahepatic cytokine expression and AP-1 activation in transgenic mice expressing hepatitis C virus core protein. *Virology* 2002;304:415–24.
- [83] Tsutsumi T, Suzuki T, Moriya K, et al. Hepatitis C virus core protein activates ERK and p38 MAPK in cooperation with ethanol in transgenic mice. *Hepatology* 2003;38:820–8.
- [84] Qadri I, Iwahashi M, Capasso JM, Hopken MW, Flores S, Schaack J. Induced oxidative stress and activated expression of manganese superoxide dismutase during hepatitis C virus replication: role of JNK, p38 MAPK and AP-1. *Biochem J* 2004;378:919–28.
- [85] Ruggieri A, Harada T, Matsuura Y, Miyamura T. Sensitization to Fas-mediated apoptosis by hepatitis C virus core protein. *Virology* 1997;229:68–76.
- [86] Zhu N, Khoshnan A, Schneider R, et al. Hepatitis C virus core protein binds to the cytoplasmic domain of tumor necrosis factor (TNF) receptor 1 and enhances TNF-induced apoptosis. *J Virol* 1998;72:3691–7.
- [87] Hahn CS, Cho YG, Kang BS, Lester IM, Hahn YS. The HCV core protein acts as a positive regulator of fas-mediated apoptosis in a human lymphoblastoid T cell line. *Virology* 2000;276:127–37.
- [88] Honda M, Kaneko S, Shimazaki T, et al. Hepatitis C virus core protein induces apoptosis and impairs cell-cycle regulation in stably transformed Chinese hamster ovary cells. *Hepatology* 2000;31:1351–9.
- [89] Ray RB, Meyer K, Steele R, Shrivastava A, Aggarwal BB, Ray R. Inhibition of tumor necrosis factor (TNF- $\alpha$ )-mediated apoptosis by hepatitis C virus core protein. *J Biol Chem* 1998;273:2256–9.
- [90] Marusawa H, Hijikata M, Chiba T, Shimotohno K. Hepatitis C virus core protein inhibits Fas- and tumor necrosis factor  $\alpha$ -mediated apoptosis via NF- $\kappa$ B activation. *J Virol* 1999;73:4713–20.
- [91] Otsuka M, Kato N, Taniguchi H, et al. Hepatitis C virus core protein inhibits apoptosis via enhanced Bcl-xL expression. *Virology* 2002;296:84–93.
- [92] Sacco R, Tsutsumi T, Suzuki R, et al. Antiapoptotic regulation by hepatitis C virus core protein through up-regulation of inhibitor of caspase-activated DNase. *Virology* 2003;317:24–35.
- [93] Okanoue T, Makiyama A, Nakayama M, et al. A follow-up study to determine the value of liver biopsy and need for antiviral therapy for hepatitis C virus carriers with persistently normal serum aminotransferase. *J Hepatol* 2005;43:599–605.
- [94] Koike K. Molecular basis of hepatitis C virus-associated hepatocarcinogenesis: lessons from animal model studies. *Clin Gastroenterol Hepatol* 2005;3:S132–5.

# Hepatitis C Virus Infection Can Present with Metabolic Disease by Inducing Insulin Resistance

Kazuhiko Koike

Department of Infectious Diseases, Internal Medicine, Graduate School of Medicine, University of Tokyo, Tokyo, Japan

## Key Words

Diabetes · Hepatitis C virus · Insulin resistance · Insulin receptor substrate · Transgenic mouse

## Abstract

Although hepatitis C virus (HCV) targets the liver, it has become increasingly evident that HCV can induce diseases of many organs. Recently, much attention is drawn to metabolic disorders in HCV infection. First, hepatic steatosis and derangement in lipid metabolism have been found characteristic of HCV infection, and later on, a correlation was noted between HCV infection and diabetes as well as insulin resistance. We have demonstrated that HCV by itself can induce insulin resistance through disturbing the insulin signaling pathway by HCV proteins. The fact that HCV infection induces insulin resistance by the virus itself may influence the progression of chronic liver disease and open up novel therapeutic approaches. In conclusion, towards the future, HCV infection needs to be viewed not only as a liver disease but also as a metabolic disease.

Copyright © 2006 S. Karger AG, Basel

## Introduction

Hepatitis C virus (HCV) infects approximately 1.8 million people in Japan alone and as many as 200 million over the world and induces liver disease ranging from

chronic hepatitis through cirrhosis to hepatocellular carcinoma (HCC) [1, 2]. It has been noticed soon after the discovery that the infection with HCV does not exclusively involve the liver. In fact, type II cryoglobulinemia [3] and membranoproliferative glomerulonephritis [4] frequently occur in patients infected with HCV. Furthermore, strong associations of HCV infection with Sjogren's syndrome [5] and lichen planus [6] have been noted, which is verified in the animal model [7]. In addition, the relation between HCV infection and B cell lymphoma has attracted attention especially in Europe [8].

Recently, there have been increasing lines of evidence to indicate metabolic disturbances in HCV infection which, in turn, would influence the pathogenesis of chronic hepatitis C. The discovery of HCV in 1989 [9] enabled a comparison between chronic hepatitis C and other chronic liver diseases. As shown in the results, it has been repeatedly reported that steatosis is significantly associated with chronic hepatitis C [10, 11]. Steatosis in HCV infection is reproduced in animal models [12–14] to reinforce a pathologic role of HCV. Furthermore, patients infected with HCV have abnormalities in serum lipids, such as hypocholesterolemia and abnormal levels of apolipoproteins in serum [15, 16]; they are rectified in sustained virological responders to interferon (IFN) [16]. Thus, the association between HCV infection and a derangement in lipid metabolism has become increasingly strong, both in patients and experimental systems in animals. Finally, patients with chronic hepatitis C accompanied by severe steatosis develop hepatic fibrosis with an

## KARGER

Fax +41 61 306 12 34  
E-Mail [karger@karger.ch](mailto:karger@karger.ch)  
[www.karger.com](http://www.karger.com)

© 2006 S. Karger AG, Basel  
0300-5526/06/0492-0051\$23.50/0

Accessible online at:  
[www.karger.com/int](http://www.karger.com/int)

Kazuhiko Koike, MD  
Department of Infectious Diseases, Internal Medicine  
Graduate School of Medicine, University of Tokyo  
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655 (Japan)  
Fax +81 3 5800 8799, E-Mail [kkoike-tky@umin.ac.jp](mailto:kkoike-tky@umin.ac.jp)

increased velocity [17]. All in all, we could say that abnormal lipid metabolism in HCV infection is deeply involved in the pathogenesis of hepatitis C.

### HCV Infection and Diabetes


Diabetes is suggested as another metabolic disease in association with HCV infection. In 1994, Allison et al. [18] reported an epidemiological link between diabetes and HCV infection. However, doubts were cast on the association in view of a decreased glucose tolerance in advanced chronic hepatitis as well as an increased opportunity for HCV infection in diabetics who frequently receive determination of blood sugar. Several reports from the same group and others followed along this line. The trend to accept the solid association between diabetes and HCV infection seems to have been triggered in the United States by the population study by Metha et al. [19].

However, the association between diabetes and HCV infection is blemished by factors responsible for decreased glucose tolerance, such as advanced cirrhosis, obesity and ageing common in patients with hepatitis C; they make it difficult to prove this association. Hence, there is a need to evaluate the association by basic studies in experimental systems.

### HCV Infection Induces Insulin Resistance

We set out to demonstrate the association between HCV infection and diabetes using the animal model. Mice transgenic for the HCV core gene were employed to this end [12, 13]. These mice are engineered to have the HCV core gene of genotype 1b in the absence of other viral genes. They express HCV core protein of the expected size in the liver, in levels comparable with those of patients with chronic hepatitis C (fig. 1). Half of them develop HCC later during their lives [13]. These transgenic mice were fed with their normal littermates, and the glucose metabolism was compared between them [20].


Although mice transgenic for the core gene did not develop overt diabetes, they had markedly elevated serum levels of insulin. Plasma glucose levels were somewhat higher in transgenic mice than in their normal littermates, both in the fast and after ample feeding, with no significant differences between them. In remarkable contrast, serum insulin levels were significantly higher in transgenic than in normal mice in both conditions (fig. 2). Since such a combination of normal glucose levels and




**Fig. 1.** Expression of HCV core gene in transgenic mouse. It carries the core gene of HCV genotype 1b alone and expresses the core protein of expected size in the liver, at levels similar to those in human patients. Mice eventually develop HCC later in their lives.

hyperinsulinemia points to insulin resistance, glucose and insulin tolerance tests were conducted.

Mice transgenic for the HCV core gene exhibited glucose levels a little higher than those in normal littermates, without any significant differences between them. In insulin tolerance tests, glucose levels were significantly higher in transgenic than in normal mice, both 40 and 60 min after they were injected with insulin intraperitoneally (fig. 3). These results indicate suppression of the activity of insulin to decrease blood glucose levels for inducing insulin resistance in core-transgenic mice. Since only the HCV core gene had been incorporated into these transgenic mice, HCV core protein was able to induce insulin resistance *in vivo*.



**Fig. 2.** Altered homeostasis of glucose in mice transgenic for the HCV core gene. Body weight of 2-month-old mice (a), plasma glucose levels in fasting or fed mice (b) and serum insulin levels in fasting or fed mice (c) are shown. Values represent means  $\pm$  SE. NS = Not significant statistically; nTg = nontransgenic mice; Tg = transgenic mice.



**Fig. 3.** Insulin resistance in transgenic mice. Glucose tolerance in mice after overnight fasting (a). *D*-Glucose (1 g/kg body weight) was given intraperitoneally to conscious mice, and plasma glucose levels were determined at time points indicated. **b** Insulin tolerance in mice fasted overnight. Human insulin (1 U/kg body weight) was injected intraperitoneally, and glucose concentrations were determined sequentially. Values were normalized to the baseline glucose concentration at the time of insulin administration. NS = Not significant statistically; nTg = nontransgenic mice; Tg = transgenic mice.

By what mechanism does insulin resistance arise in this animal model? The insulin resistance is considered to involve two factors, namely central and peripheral insulin resistances (table 1) [21]. The hyperinsulinemic-euglycemic clamp method was employed to differentiate between them. In this method, hepatic glucose production (HGP) is calculated on the basis of amounts of glu-

cose required to keep plasma glucose levels within a certain range at serum insulin levels higher than physiological ones. In normal control mice, HGP was suppressed by 60% by the administration of insulin, in contrast to core-transgenic mice in which there was no appreciable suppression of HGP by insulin (fig. 4). These results indicate a hepatic (central) origin of the insulin resistance



**Table 1.** Two types of insulin resistance


Type	Mechanism
Peripheral	A shortage of insulin action in the muscle due to deficit in the insulin-induced uptake of glucose into muscles
Central	A shortage of insulin action in the liver due to deficit in the insulin-induced suppression of glucose production in hepatocytes

in transgenic mice. For further confirmation, an uptake of glucose into muscle was determined. There were no differences in the uptake in response to administration of insulin between normal and transgenic mice. Therefore, the insulin resistance in mice transgenic for the HCV core gene is central and hepatic.


### HCV Core Protein Suppresses the Transduction of Insulin Signaling in Hepatocytes

Next, we evaluated how insulin resistance elicits in mice transgenic for the HCV core gene. For this purpose, liver homogenate was immunoblotted with antiphosphotyrosine and antiphosphoserine antibodies after insulin receptor substrate (IRS)-1 and IRS-2 had been immunoprecipitated. Tyrosines in IRS-1 were weakly phosphorylated both in normal and transgenic mice before they received insulin, with no differences between them. However, after the administration of insulin, the phosphorylation of tyrosines in IRS-1 increased in normal but not in transgenic mice (fig. 5). Obtained results suggested disturbance in tyrosine phosphorylation as one of the factors responsible for insulin resistance in the liver. There were no differences in phosphorylation of serines in IRS-1 or tyrosines in IRS-2 between normal and transgenic mice. Combined, they provided experimental evidence for the development of insulin resistance by the presence of HCV in the liver that would occur by disturbing the transduction of insulin signaling in hepatocytes (fig. 6).


There remains a possibility for the HCV core protein to directly prohibit phosphorylation of tyrosines, or else, it might inhibit tyrosine phosphorylation via certain cytokines. In our extensive searches for the expression of cytokines in the liver of transgenic mice, only TNF- $\alpha$  and IL-1 $\beta$  have been found with an increased expression [22]. Therefore, for the purpose of evaluating the role of



**Fig. 4.** Characterization of glucose metabolism in transgenic mice. Glucose production in the liver was calculated using the hyperinsulinemic-euglycemic clamp method. NS = Not significant statistically; nTg = nontransgenic mice; Tg = transgenic mice.



**Fig. 5.** Phosphorylation of tyrosine in IRS-1 in response to insulin stimulation. Liver tissues from control mice, transgenic mice with or without anti-TNF- $\alpha$  antibody treatment, were analyzed before and 20 as well as 40 min after administration of insulin. Samples were subjected to immunoprecipitation with anti-IRS-1 antibody and then immunoblotted with indicated antibodies. Experiments were performed in triplicate, and a representative picture is exhibited. Immunoblotting with antiphosphotyrosine (PY) antibody (lane a) did not augment phosphorylation of tyrosine in IRS-1 after stimulation with insulin in the core gene transgenic mice (Tg), in contrast to tyrosine phosphorylation markedly enhanced in control mice (nTg). Insulin-stimulated tyrosine phosphorylation was restored 40 min after treatment with anti-TNF- $\alpha$  antibody. Note differences in the intensity of bands 40 min after the administration of insulin (arrowheads). Immunoblotting with anti-IRS-1 antibody (lane b) served as control for the IRS-1 load.



**Fig. 6.** A proposed mechanism for insulin resistance in HCV infection. HCV itself or elevated levels of cytokines such as TNF- $\alpha$  may inhibit tyrosine phosphorylation of IRS-1 in the liver, suppress intracellular transduction of insulin signal and lead to insulin resistance. PKC = Protein kinase C; MAPK = mitogen-activated protein kinase.

TNF- $\alpha$  in insulin resistance in transgenic mice, serum insulin was determined and an insulin tolerance test performed after they had received anti-TNF- $\alpha$  intraperitoneally. Pretreatment with anti-TNF- $\alpha$  partially improved insulin resistance in mice transgenic for the HCV core gene. Albeit a direct anti-insulin activity of core protein and direct or indirect factors for insulin resistance are not to be excluded, high levels of TNF- $\alpha$  in the liver would be one of the factors for expression of insulin resistance in this mouse model.

### Insulin Resistance in Patients with Chronic Hepatitis C

Concurrently with our report in experimental systems, Aytug et al. [23] investigated insulin signaling in biopsied liver specimens from patients with chronic hepatitis C. Specifically, they evaluated changes in IRS-1, IRS-2 and phosphatidylinositol (PI)3 kinase levels in the liver of patients. With insulin stimulation of biopsied liver samples, insulin receptor proteins and IRS-1 increased, while phosphorylation of tyrosines in IRS-1 decreased to one half of the baseline value, along with a diminished activity for PI3 kinase associated with IRS-1, in patients with chronic hepatitis C. The authors went on to propose a possibility for disturbed transduction of the insulin sig-

naling pathway in the liver to induce insulin resistance in patients with chronic hepatitis C [23]. Their report is quite intriguing in that it opens up the way for evaluating an association between HCV infection and insulin resistance in clinical samples at the molecular level.

The results of Aytug et al. [23] inadvertently coincide with ours in analyzing the mechanism of insulin resistance with the experimental system in mice (*vide supra*). They unanimously incriminate impaired tyrosine phosphorylation in IRS-1 in the induction of insulin resistance by HCV infection. It struck us as a surprise that the mechanism of insulin resistance induced by HCV infection has been in agreement between clinical samples and experimental animals, in spite of hepatic IRS-2 that was preferred to IRS-1 for its role in development of insulin resistance in former studies [24]. HCV infection is peculiar in that IRS-1 weighs heavier than IRS-2 in the induction of hepatic insulin resistance.

Although our data strongly indicate a hepatic character of insulin resistance in HCV infection, they by no means exclude roles of other factors in the induction of this resistance. There is little expression of the HCV core gene in muscles of our animal model; it is not known if HCV infects muscular cells in patients with chronic hepatitis C. Factors not intrinsic to the liver would have to be evaluated to sort this out, including dysfunction of mitochondria for induction of insulin resistance [25].

## Insulin Resistance for Advanced Hepatic Fibrosis

Insulin resistance in HCV infection may have an additional significant clinical implication. In 260 patients with chronic hepatitis C, Hui et al. [26] have tried to establish the relationship between liver histology and indicators of glucose metabolism, as well as insulin resistance represented by the homeostasis model assessment of insulin resistance. They have found that insulin resistance already exists in hepatitis C patients with stage 0 or 1 fibrosis in the liver. This indicates that insulin resistance in HCV infection is not attributable to advanced liver disease. In their study, independent predictors of insulin resistance in HCV infection were body mass index, non-response to antiviral treatment, intensity of portal inflammation and infection with HCV genotype 3 [26]. Furthermore, the homeostasis model assessment of insulin resistance was a significant and independent predictor of the stage and velocity of hepatic fibrosis. The results of the study are of much importance, because they implicate a role of insulin resistance and hyperinsulinemia by inference, in promoting the progression of hepatic fibrosis. Insulin has been proven as an aggravating factor not only in atherosclerosis, but also in systemic inflammation and fibrosis. The liver is no exception to this.

## Conclusions: Hepatitis C Viewed as a Metabolic Disease and Outlook for Therapeutic Strategies in the Future

We have demonstrated that HCV per se induces insulin resistance in the animal model. Superimposed high-fat diet and obesity may lead to overt diabetes. Since insulin resistance accelerates the progression of chronic hepatitis C, it would naturally influence the development of HCC. Although the association has not been established between nonalcoholic steatohepatitis and HCC, it needs to be energetically pursued in view of the histological homology of nonalcoholic steatohepatitis to chronic hepatitis C. Drugs for improving glucose metabolism and insulin resistance need to be kept in store in the treatment of hepatitis C patients who have failed to respond to antivirals, because they may well prevent progression of fibrosis and development of HCC in such patients. Traditional 'high-protein and high-calorie' diet, especially advocated in Japan after World War II, is obviously detrimental, except in some patients with advanced cirrhosis. Consultation on the dietary habit with hepatitis C patients should include iron restriction [27] as well as weight control, because high-calorie intakes are likely to accelerate hepatic fibrosis by aggravating insulin resistance.

## References

- 1 Saito I, Miyamura T, Ohbayashi A, Harada H, Katayama T, Kikuchi S, Watanabe Y, Koi S, Onji M, Ohta Y: Hepatitis C virus infection is associated with the development of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1990;87:6547-6549.
- 2 Simonetti RG, Camma C, Fiorello F, Cottone M, Rapicetta M, Marino L, Fiorentino G, Craxi A, Cicciaglione A, Giuseppetti R: Hepatitis C virus infection as a risk factor for hepatocellular carcinoma in patients with cirrhosis. A case-control study. *Ann Intern Med* 1992; 116:97-102.
- 3 Agnello V, Chung RT, Kaplan LM: A role for hepatitis C virus infection in type II cryoglobulinemia. *N Engl J Med* 1992;327:1490-1495.
- 4 Johnson RJ, Gretch DR, Yamabe H, Hart J, Bacchi CE, Hartwell P, Couser WG, Corey L, Wener MH, Alpers CE: Membranoproliferative glomerulonephritis associated with hepatitis C virus infection. *N Engl J Med* 1993;328: 465-470.
- 5 Haddad J, Deny P, Munz-Gotheil C, Ambrosini JC, Trinchet JC, Pateron D, Mal F, Callard P, Beaugrand M: Lymphocytic sialadenitis of Sjogren's syndrome associated with chronic hepatitis C virus liver disease. *Lancet* 1992; 339:321-323.
- 6 Pawlotsky JM, Benchiki H, Pellet C, Duval J, Dhumeaux D, Revuz J, Bagot M: Lichen planus and hepatitis C virus (HCV)-related chronic hepatitis: Evaluation of HCV genotypes. *Br J Dermatol* 1995;133:666-667.
- 7 Koike K, Moriya K, Ishibashi K, Yotsuyanagi H, Shintani Y, Fujie H, Kurokawa K, Matsuura Y, Miyamura T: Sialadenitis histologically resembling Sjogren syndrome in mice transgenic for hepatitis C virus envelope genes. *Proc Natl Acad Sci USA* 1997;94:233-236.
- 8 Gisbert JP, Garcia-Buey L, Pajares JM, Moreno-Otero R: Prevalence of hepatitis C virus infection in B-cell non-Hodgkin's lymphoma: Systematic review and meta-analysis. *Gastroenterology* 2003;125:1723-1732.
- 9 Choo QL, Kuo G, Weiner AJ, Overby LR, Bradley DW, Houghton M: Isolation of a cDNA clone derived from a blood-borne non-A, non-B viral hepatitis genome. *Science* 1989; 244:359-362.
- 10 Bach N, Thung SN, Schaffner F: The histological features of chronic hepatitis C and autoimmune chronic hepatitis: A comparative analysis. *Hepatology* 1992;15:572-577.
- 11 Lefkowitz JH, Schiff ER, Davis GL, Perrillo RP, Lindsay K, Bodenheimer HC Jr, Balart LA, Ortego TJ, Payne J, Dienstag JL, Gibas A, Jacobson IM, Tamburro CH, Carey W, O'Brien C, Sampliner R, van Thiel DH, Feit D, Albrecht J, Meschievitz C, Sanghvi V, Vaughan RD: Pathological diagnosis of chronic hepatitis C: A multicenter comparative study with chronic hepatitis B. *Gastroenterology* 1993; 104:595-603.
- 12 Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Ishibashi K, Matsuura Y, Miyamura T, Koike K: Hepatitis C virus core protein induces hepatic steatosis in transgenic mice. *J Gen Virol* 1997;78:1527-1531.
- 13 Moriya K, Fujie H, Shintani Y, Yotsuyanagi H, Tsutsumi T, Ishibashi K, Matsuura Y, Kimura S, Miyamura T, Koike K: The core protein of hepatitis C virus induces hepatocellular carcinoma in transgenic mice. *Nat Med* 1998;4:1065-1067.

- 14 Lerat H, Honda M, Beard MR, Loesch K, Sun J, Yang Y, Okuda M, Gosert R, Xiao SY, Weinman SA, Lemon SM: Steatosis and liver cancer in transgenic mice expressing the structural and nonstructural proteins of hepatitis C virus. *Gastroenterology* 2002;122:352–365.
- 15 Moriya K, Sintani Y, Fujie H, Miyoshi H, Tsutsumi T, Yotsuyanagi H, Yasuda K, Iino S, Kimura S, Koike K: Serum lipid profile of patients with genotype 1b hepatitis C viral infection in Japan. *Hepatol Res* 2003;25:369–374.
- 16 Naeem M, Bacon BR, Mistry B, Britton RS, Di Bisceglie AM: Changes in serum lipoprotein profile during interferon therapy in chronic hepatitis C. *Am J Gastroenterol* 2001;96:2468–2472.
- 17 Lonardo A, Adinolfi LE, Loria P, Carulli N, Ruggiero G, Day CP: Steatosis and hepatitis C virus: Mechanisms and significance for hepatic and extrahepatic disease. *Gastroenterology* 2004;126:586–597.
- 18 Allison ME, Wreghitt T, Palmer CR, Alexander GJ: Evidence for a link between hepatitis C virus infection and diabetes mellitus in a cirrhotic population. *J Hepatol* 1994;21:1135–1139.
- 19 Mehta SH, Brancati FL, Sulkowski MS, Strathdee SA, Szklo M, Thomas DL: Prevalence of type 2 diabetes mellitus among persons with hepatitis C virus infection in the United States. *Ann Intern Med* 2000;133:592–599.
- 20 Shintani Y, Fujie H, Miyoshi H, Tsutsumi T, Tsukamoto K, Kimura S, Moriya K, Koike K: Hepatitis C virus infection and diabetes: Direct involvement of the virus in the development of insulin resistance. *Gastroenterology* 2004;126:840–848.
- 21 Kahn BB: Type 2 diabetes: When insulin secretion fails to compensate for insulin resistance. *Cell* 1998;92:593–596.
- 22 Tsutsumi T, Suzuki T, Moriya K, Yotsuyanagi H, Shintani Y, Fujie H, Matsuura Y, Kimura S, Koike K, Miyamura T: Alteration of intrahepatic cytokine expression and AP-1 activation in transgenic mice expressing hepatitis C virus core protein. *Virology* 2002;304:415–424.
- 23 Aytug S, Reich D, Sapiro LE, Bernstein D, Begum N: Impaired IRS-1/PI3-kinase signaling in patients with HCV: A mechanism for increased prevalence of type 2 diabetes. *Hepatol* 2003;38:1384–1392.
- 24 Suzuki R, Tobe K, Aoyama M, Inoue A, Sakamoto K, Yamauchi T, Kamon J, Kubota N, Terauchi Y, Yoshimatsu H, Matsuhisa M, Nagasaka S, Ogata H, Tokuyama K, Nagai R, Kadowaki T: Both insulin signaling defects in the liver and obesity contribute to insulin resistance and cause diabetes in *Irs2(-/-)* mice. *J Biol Chem* 2004;279:25039–25049.
- 25 Petersen KF, Befroy D, Dufour S, Dziura J, Ariyan C, Rothman DL, DiPietro L, Cline GW, Shulman GI: Mitochondrial dysfunction in the elderly: Possible role in insulin resistance. *Science* 2003;300:1140–1142.
- 26 Hui JM, Sud A, Farrell GC, Bandara P, Byth K, Kench JG, McCaughan GW, George J: Insulin resistance is associated with chronic hepatitis C virus infection and fibrosis progression. *Gastroenterology* 2003;125:1695–1704.
- 27 Kato J, Kobune M, Nakamura T, Kuroiwa G, Takada K, Takimoto R, Sato Y, Fujikawa K, Takahashi M, Takayama T, Ikeda T, Niitsu Y: Normalization of elevated hepatic 8-hydroxy-2'-deoxyguanosine levels in chronic hepatitis C patients by phlebotomy and low iron diet. *Cancer Res* 2001;61:8697–8702.