

fatty acid intake with the subsequent risk of postpartum depression. The study failed to substantiate a clear inverse relationship between these factors but found several suggestive relationships. Consumption of n-3 polyunsaturated fatty acids, especially docosahexaenoic acid, in the third quartile was marginally independently associated with a decreased risk of postpartum depression, although the linear trends were not statistically significant.

The present findings are at variance with previous epidemiological studies showing an inverse relationship between fish intake and postpartum depression or major depressive symptoms (Hibbeln, 1998, 2002; Tanskanen *et al.* 2001*a,b*; Peet, 2004). The Alpha-Tocopherol Beta-Carotene Cancer Prevention study found no measurable association of consumption of fish or n-3 polyunsaturated fatty acids with self-reported depressed mood and hospital treatment for major depression, although a significant positive dose-response relationship between fish intake and self-reported depressed mood was identified (Hakkarainen *et al.* 2004). Peet and Horrobin (2002) demonstrated that a dosage of 1 g/day, but not 2 or 4 g/day, of eicopentaenoic acid for 12 weeks was effective in significantly improving depressive symptoms. The current observations are partially consistent with these results.

A significant inverse association between adipose tissue docosahexaenoic acid levels and depressive symptoms was found in a cross-sectional study among 139 healthy adults from the island of Crete (Mamalakis *et al.* 2002). The current results for docosahexaenoic acid, which showed borderline significance regarding the risk of postpartum depression in comparing two quartiles, are in partial agreement with this finding. An inverse association between docosahexaenoic acid intake and postpartum depression may be attributable to the regulation of serotonergic nervous system function. Plasma docosahexaenoic acid concentrations were significantly correlated with cerebrospinal fluid concentrations of 5-hydroxyindoleacetic acid, a metabolite of serotonin, among healthy subjects (Hibbeln *et al.* 1998). Docosahexaenoic acid may inhibit signal transduction mechanisms in the human central nervous system (Stoll *et al.* 1999). A laboratory study demonstrated that dietary docosahexaenoic acid altered T-cell

membrane microdomain composition and suppressed the protein kinase C θ signalling axis in mice (Fan *et al.* 2004).

The present results are incompatible with the hypothesis that a balance between n-3 and n-6 polyunsaturated fatty acid metabolism is important in the manifestation of depression (Hibbeln & Salem, 1995). The differences between Japanese and Western diets should be taken into account when interpreting our results. A clear protective association of docosahexaenoic acid and fish intake with postpartum depression may exist in populations with low fish intake. For example, in 20 Canadian pregnant women, mean intake of eicosapentaenoic and docosahexaenoic acids was estimated to be 35 and 82 mg/day respectively (Denomme *et al.* 2005). The corresponding figures for our population were 204 and 329 mg respectively. Unknown active agents in fish might have interfered with a beneficial effect of fish or docosahexaenoic acid in the development of postpartum depression in our population. For example, methylmercury and dioxins are accumulated in fish and shellfish through the marine food web.

A case-control study in China showed no relationship between saturated, monounsaturated or n-6 polyunsaturated fatty acid levels in red bloods cells and suicide attempts, although eicosapentaenoic and docosahexaenoic acid levels were significantly inversely associated with suicide attempt risk (Huan *et al.* 2004). Mental illnesses such as depression may be an important cause of suicide. In this context, the current observations were partially consistent with these results.

The present study had several methodological strengths. The prospective design and relatively high rate of follow-up (86.3%) in this study minimized the possibility of recall bias or bias caused by loss of follow-up. Although 135 subjects did not participate in the follow-up survey, there were no material differences in the distributions of age, gestation, changes in diet in the previous month, season when data at baseline were collected and intake of total fat, monounsaturated, n-3 polyunsaturated and n-6 polyunsaturated fatty acids, fish, meat and dairy products as compared to the 867 participants (data not shown). Compared with non-participants in the follow-up, participants were

more likely to report no history of smoking, high family income, high educational level and high intake of saturated fatty acids, cholesterol and eggs and were less likely to be thin, although these differences would not be so large as to cause a serious bias. Study subjects were homogeneous in terms of having the same residential background. We incorporated extensive information on potential confounding factors and adjustments were made for these in data analysis. However, we were not able to undertake a comprehensive assessment by controlling for personal and family psychiatric history, sociocultural factors and personal and family relations.

There are several limitations in the present study that deserve recognition. The diagnosis of postpartum depression was established with the EPDS, a self-report rating scale, rather than a clinician-administered structured diagnostic interview. The EPDS was originally designed as a screening test of depressive symptoms within the previous 7 days. Moreover, the second survey was conducted at 2–9 months postpartum even though 89.2% of the subjects took part in the second survey at 3–4 months postpartum. Thus, it is difficult to estimate the incidence and prevalence of postpartum depression accurately. The disadvantage could bias the magnitude of the observed effects towards the null due to non-differential outcome misclassification.

Dietary data were obtained using a self-administered semi-quantitative dietary assessment questionnaire. As we did not undertake the assessment of the dietary habits of the subjects, the possibility of misclassification should be considered. According to validation tests, the correlation coefficients for nutrient intake between those estimated from the diet history questionnaire and those observed by a 3-day dietary record were 0.75, 0.50, 0.37 and 0.49 for saturated fatty acids, monounsaturated fatty acids, polyunsaturated fatty acids and cholesterol respectively, in women (Sasaki *et al.* 1998). A highly positive correlation was also observed between marine-origin n-3 polyunsaturated fatty acid intake estimated by the diet history questionnaire and the corresponding concentration in the serum phospholipid fraction ($r=0.51$ and 0.69 in men and women respectively) (Sasaki *et al.* 2000). Subjects with postpartum depression might not be aware of the

ill effects of diet. Such a hypothesis would lead to bias towards the null. Our diet history questionnaire was designed to assess recent dietary intake, that is for 1 month prior to completing the questionnaire. However, this disadvantage is likely to be alleviated after adjustment for the season when data at baseline were collected. Changes in diet in the past month were controlled for because pregnant females are likely to change their diet for reasons such as nausea gravidarum. In this study, there was no significant interaction between changes in diet in the past month and exposures under investigation in relation to postpartum depression (data not shown). Potential misclassification associated with variability across subjects with regard to the stability of diet during pregnancy is likely to be negligible. Therefore, negative findings of our study were unlikely to be ascribed to dietary assessment during pregnancy.

In the baseline survey, the participation rate was low in Neyagawa City (17.2%). We were not able to evaluate the difference between participants and non-participants in Neyagawa City because data on personal characteristics among the non-participants were not available. In addition, we were not able to calculate the participation rate of subjects from other areas, nor were data available on the non-participants in those areas. The present study subjects were an unrepresentative sample of Japanese females in the general population and the present findings may not be generalized. In fact, educational levels were higher in the present study population than in the general population. According to the 2000 population census of Japan, the proportions of females aged 30 to 34 years in Osaka Prefecture with <13, 13–14, 15+ and unknown years of education were 49.2, 32.3, 13.6 and 4.9% respectively. The corresponding figures for the current study were 29.7, 42.4, 27.9 and 0.0% respectively. The absence of a relationship between intake of fish and n-3 polyunsaturated fatty acids in our data may be related to the fact that the current study did not have substantial statistical power.

In conclusion, the current prospective study failed to verify an inverse relationship between fish intake and the subsequent risk of postpartum depression. Larger investigations with more precise and detailed exposure and outcome measurements are needed to draw a

conclusion on whether consumption of fish and n-3 polyunsaturated fatty acids is preventive against postpartum depression.

APPENDIX

Space limitations preclude the inclusion as authors of the following members of the Osaka Maternal and Child Health Study Group: Ichiro Matsunaga and Dr Hajime Oda, Osaka Prefectural Institute of Public Health; Drs Hideharu Kanzaki and Mitsuyoshi Kitada, Department of Obstetrics and Gynaecology, Kansai Medical University; Dr Yorihiro Horikoshi, Department of Obstetrics and Gynaecology, Kansai Medical University Kori; Drs Osamu Ishiko, Yuichiro Nakai, Junko Nishio and Seiichi Yamamasu, Department of Obstetrics and Gynaecology, Osaka City University Graduate School of Medicine; Dr Jinsuke Yasuda, Department of Obstetrics and Gynaecology, Matsushita Memorial Hospital; Dr Seigo Kawai, Department of Obstetrics and Gynaecology, Hoshigaoka Koseinenkin Hospital; Dr Kazumi Yanagihara, Yanagihara Clinic; Dr Koji Wakuda, Department of Obstetrics and Gynaecology, Fujimoto Hospital; Dr Tokio Kawashima, Kyohritsu Women's Clinic; Dr Katsuhiko Narimoto, Ishida Hospital Obstetrics, Gynaecology; Dr Yoshihiko Iwasa, Iwasa Women's Clinic; Dr Katsuhiko Orino, Orino Lady's Clinic; Dr Itsuo Tsunetoh, Tsunetoh Obstetrics and Gynaecology; Dr Junichi Yoshida, Yoshida Clinic; Dr Junichi Ito, Ito Obstetrics and Gynaecology Clinic; Dr Takuzi Kaneko, Kaneko Sanfujinka; Dr Takao Kamiya, Kamiya Ladies Clinic; Dr Hiroyuki Kuribayashi, Kuribayashi Clinic; Dr Takeshi Taniguchi, Taniguchi Hospital; Dr Hideo Takemura, Kosaka Women's Hospital; Dr Yasuhiko Morimoto, Aizenbashi Hospital.

ACKNOWLEDGEMENTS

We thank the Neyagawa City Government, Hirakata City Government, Katano City Government, Shijonawate City Government, Kaizuka City Government, Takaishi City Government, Hannan City Government, Neyagawa City Medical Association, Hirakata City Medical Association and Kadoma City Medical Association for their valuable support;

Kumiko Shirata, Osaka City University School of Nursing for helpful discussions; and Tomoko Shibazaki, Nahoko Nishimura and Naomi Takaoka for their assistance. This study was supported by a Grant-in-Aid (13770206, 16790351) for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology, Health and Labour Sciences Research Grants, Research on Allergic Disease and Immunology from the Ministry of Health, Labour and Welfare and the Uehara Memorial Foundation, Japan.

DECLARATION OF INTEREST

None.

REFERENCES

- Adams, P. B., Lawson, S., Sanigorski, A. & Sinclair, A. J. (1996). Arachidonic acid to eicosapentaenoic acid ratio in blood correlates positively with clinical symptoms of depression. *Lipids* **31**, S157–S161.
- Cox, J. L., Holden, J. M. & Sagovsky, R. (1987). Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *British Journal of Psychiatry* **150**, 782–786.
- Denomme, J., Stark, K. D. & Holub, B. J. (2005). Directly quantitated dietary (n-3) fatty acid intakes of pregnant Canadian women are lower than current dietary recommendations. *Journal of Nutrition* **135**, 206–211.
- Edwards, R., Peet, M., Shay, J. & Horrobin, D. (1998). Omega-3 polyunsaturated fatty acid levels in the diet and in red blood cell membranes of depressed patients. *Journal of Affective Disorders* **48**, 149–155.
- Fan, Y. Y., Ly, L. H., Barhoumi, R., McMurray, D. N. & Chapkin, R. S. (2004). Dietary docosahexaenoic acid suppresses T cell protein kinase C theta lipid raft recruitment and IL-2 production. *Journal of Immunology* **173**, 6151–6160.
- Hakkarainen, R., Partonen, T., Haukka, J., Virtamo, J., Albanes, D. & Lönnqvist, J. (2004). Is low dietary intake of omega-3 fatty acids associated with depression? *American Journal of Psychiatry* **161**, 567–569.
- Hibbeln, J. R. (1998). Fish consumption and major depression. *Lancet* **351**, 1213.
- Hibbeln, J. R. (2002). Seafood consumption, the DHA content of mothers' milk and prevalence rates of postpartum depression: a cross-national, ecological analysis. *Journal of Affective Disorders* **69**, 15–29.
- Hibbeln, J. R., Linnoila, M., Umhau, J. C., Rawlings, R., George, D. T. & Salem Jr., N. (1998). Essential fatty acids predict metabolites of serotonin and dopamine in cerebrospinal fluid among healthy control subjects, and early- and late-onset alcoholics. *Biological Psychiatry* **44**, 235–242.
- Hibbeln, J. R. & Salem Jr., N. (1995). Dietary polyunsaturated fatty acids and depression: when cholesterol does not satisfy. *American Journal of Clinical Nutrition* **62**, 1–9.
- Huan, M., Hamazaki, K., Sun, Y., Itomura, M., Liu, H., Kang, W., Watanabe, S., Terasawa, K. & Hamazaki, T. (2004). Suicide attempt and n-3 fatty acid levels in red blood cells: a case control study in China. *Biological Psychiatry* **56**, 490–496.
- Jacka, F. N., Pasco, J. A., Henry, M. J., Kotowicz, M. A., Nicholson, G. C. & Berk, M. (2004). Dietary omega-3 fatty acids and depression in a community sample. *Nutritional Neuroscience* **7**, 101–106.

- Maes, M., Christophe, A., Delanghe, J., Altamura, C., Neels, H. & Meltzer, H. Y. (1999). Lowered omega3 polyunsaturated fatty acids in serum phospholipids and cholesteryl esters of depressed patients. *Psychiatry Research* **85**, 275–291.
- Maes, M., Smith, R., Christophe, A., Cosyns, P., Desnyder, R. & Meltzer, H. (1996). Fatty acid composition in major depression: decreased omega 3 fractions in cholesteryl esters and increased C20:4 omega 6/C20:5 omega 3 ratio in cholesteryl esters and phospholipids. *Journal of Affective Disorders* **38**, 35–46.
- Mamalakos, G., Tornaritis, M. & Kafatos, A. (2002). Depression and adipose essential polyunsaturated fatty acids. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* **67**, 311–318.
- Marangell, L. B., Martinez, J. M., Zboyan, H. A., Kertz, B., Kim, H. F. S. & Puryear, L. J. (2003). A double-blind, placebo-controlled study of the omega-3 fatty acid docosahexaenoic acid in the treatment of major depression. *American Journal of Psychiatry* **160**, 996–998.
- Nemets, B., Stahl, Z. & Belmaker, R. H. (2002). Addition of omega-3 fatty acid to maintenance medication treatment for recurrent unipolar depressive disorder. *American Journal of Psychiatry* **159**, 477–479.
- Okano, T., Murata, M., Masuji, F., Tamaki, R., Nomura, J., Miyaoka, H. & Kitamura, T. (1996). Validation and reliability of Japanese version of the EPDS. *Archives of Psychiatric Diagnostics and Clinical Evaluation* **7**, 525–533.
- Peet, M. (2004). International variations in the outcome of schizophrenia and the prevalence of depression in relation to national dietary practices: an ecological analysis. *British Journal of Psychiatry* **184**, 404–408.
- Peet, M. & Horrobin, D. F. (2002). A dose-ranging study of the effects of ethyl-eicosapentaenoate in patients with ongoing depression despite apparently adequate treatment with standard drugs. *Archives of General Psychiatry* **59**, 913–919.
- Peet, M., Murphy, B., Shay, J. & Horrobin, D. (1998). Depletion of omega-3 fatty acid levels in red blood cell membranes of depressive patients. *Biological Psychiatry* **43**, 315–319.
- Sasaki, S., Ushio, F., Amano, K., Morihara, M., Todoriki, T., Uehara, Y. & Toyooka, T. (2000). Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *Journal of Nutritional Science and Vitaminology* **46**, 285–296.
- Sasaki, S., Yanagibori, R. & Amano, K. (1998). Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *Journal of Epidemiology* **8**, 203–215.
- Stoll, A. L., Severus, W. E., Freeman, M. P., Rueter, S., Zboyan, H. A., Diamond, E., Cress, K. K. & Marangell, L. B. (1999). Omega 3 fatty acids in bipolar disorder: a preliminary double-blind, placebo-controlled trial. *Archives of General Psychiatry* **56**, 407–412.
- Su, K. P., Huang, S. Y., Chiu, C. C. & Shen, W. W. (2003). Omega-3 fatty acids in major depressive disorder. A preliminary double-blind, placebo-controlled trial. *European Neuropsychopharmacology* **13**, 267–271.
- Tanskanen, A., Hibbeln, J. R., Hintikka, J., Haatainen, K., Honkalampi, K. & Viinamäki, H. (2001a). Fish consumption, depression, and suicidality in a general population. *Archives of General Psychiatry* **58**, 512–513.
- Tanskanen, A., Hibbeln, J. R., Tuomilehto, J., Uutela, A., Haukkala, A., Viinamäki, H., Lehtonen, J. & Vartiainen, E. (2001b). Fish consumption and depressive symptoms in the general population in Finland. *Psychiatric Services* **52**, 529–531.
- Willett, W. & Stampfer, M. J. (1986). Total energy intake: implications for epidemiologic analyses. *American Journal of Epidemiology* **124**, 17–27.

Preliminary communication

Dietary folate and vitamins B₁₂, B₆, and B₂ intake and the risk of postpartum depression in Japan: The Osaka Maternal and Child Health Study

Yoshihiro Miyake^{a,*}, Satoshi Sasaki^b, Keiko Tanaka^a, Tetsuji Yokoyama^c,
Yukihiro Ohya^d, Wakaba Fukushima^e, Kyoko Saito^{c,f}, Satoko Ohfuji^e,
Chikako Kiyohara^g, Yoshio Hirota^c
The Osaka Maternal and Child Health Study Group¹

^a Department of Public Health, Faculty of Medicine, Fukuoka University, Fukuoka 814-0180, Fukuoka, Japan

^b Nutritional Epidemiology Program, National Institute of Health and Nutrition, Tokyo, Japan

^c Department of Technology Assessment and Biostatistics, National Institute of Public Health, Wako, Japan

^d Division of Allergy, Department of Medical Specialties, National Center for Child Health and Development, Tokyo, Japan

^e Department of Public Health, Osaka City University School of Medicine, Osaka, Japan

^f Center for Collaboration and Partnership, National Institute of Health and Nutrition, Tokyo, Japan

^g Department of Preventive Medicine, Division of Social Medicine, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan

Received 2 March 2006; received in revised form 19 May 2006; accepted 25 May 2006

Available online 11 July 2006

Abstract

Background: Previous studies showed an inverse association between folate intake and depression. However, epidemiological evidence for folate intake and postpartum depression is unavailable. This prospective study examined the relationship of dietary consumption of folate and B vitamins during pregnancy with the risk of postpartum depression.

Methods: Study subjects were 865 Japanese women. Dietary data were obtained during pregnancy from a validated self-administered diet history questionnaire. Postpartum depression was defined as present when subjects had an Edinburgh Postnatal Depression Scale score of 9 or higher between 2 and 9 months postpartum. Adjustment was made for age, gestation, parity, cigarette smoking, alcohol intake, family structure, family income, education, changes in diet in the previous 1 month, season when data at baseline were collected, body mass index, time of delivery before the second survey, medical problems in pregnancy, baby's sex, and baby's birth weight.

Results: Postpartum depression developed in 121 subjects (14.0%) 2 to 9 months postpartum. There was no measurable association between intake of folate, cobalamin, or pyridoxine and the risk of postpartum depression. Compared with riboflavin intake in the first quartile, only riboflavin consumption in the third quartile was independently related to a decreased risk of postpartum depression (multivariate odds ratio: 0.53, 95% CI: 0.29–0.95, *P* for trend = 0.55).

Limitations: Personal and family psychiatric history, sociocultural factors, and personal and family relations were not controlled for. The possibility of misclassification of dietary information during pregnancy should be considered.

* Corresponding author. Tel.: +81 92 801 1011x3311; fax: +81 92 863 8892.

E-mail address: miyake-y@fukuoka-u.ac.jp (Y. Miyake).

¹ Other members of the Study Group are listed in the Appendix.

Conclusions: Our results suggest that moderate consumption of riboflavin may be protective against postpartum depression.
© 2006 Elsevier B.V. All rights reserved.

Keywords: Dietary intake; Folate; Japan; Postpartum depression; Vitamin B₂

1. Introduction

Investigations on a possible role of nutritional factors in depressive symptoms are becoming more numerous. Components of 1-carbon metabolism such as folate and vitamin B₁₂ are involved in important processes related to central nervous system functions (Bjelland et al., 2003a). A prospective cohort study among Finnish middle-aged men demonstrated that a lower dietary intake of folate was significantly associated with an increased risk of a discharge diagnosis of a depressive disorder (Tolmunen et al., 2004). In a cross-sectional study using baseline data from a previously cited cohort study, a significant positive relationship was observed between lower intake of folate and depressive symptoms (Tolmunen et al., 2003). However, no associations were shown between intake of cobalamin (vitamin B₁₂), pyridoxine (vitamin B₆), or riboflavin (vitamin B₂) and depression (Tolmunen et al., 2003, 2004). Bjelland et al. (2003b) reported that hyperhomocysteinemia and the T/T methylenetetrahydrofolate reductase genotype, but not low plasma folate or vitamin B₁₂ levels, were significantly associated with the prevalence of depression in approximately 6000 middle-aged and elderly Norwegians; only in the subgroup of middle-aged women were plasma folate levels inversely related to depression. Some studies found low serum or red blood cell folate concentrations in patients with major depression (Abou-Saleh and Coppen, 1989; Bottiglieri et al., 2000; Carney et al., 1990; Ghadirian et al., 1980; Morris et al., 2003). On the other hand, a null (Hvas et al., 2004) or positive (Lee et al., 1998) relationship of erythrocyte folate levels with depression was found. Results regarding an association between serum vitamin B₁₂ levels and depression are equivocal (Bottiglieri et al., 2000; Lindeman et al., 2000; Penninx et al., 2000; Ramos et al., 2004; Tiemeier et al., 2002). One report suggested an inverse association between plasma vitamin B₆ levels and depressive symptoms (Hvas et al., 2004).

The mother is the only nutrient source for the fetus. Depletion of crucial nutrients during pregnancy can adversely affect both mother and infant. In view of the lack of epidemiological evidence of a relationship of dietary intake of folate and vitamins B₁₂, B₆, and B₂ with the risk of postpartum depression, we investigated this issue using data from the Osaka Maternal and Child Health Study (OMCHS).

2. Methods

2.1. Study population

The OMCHS is an ongoing prospective cohort study that investigates preventive and risk factors for maternal and child health problems. The principal objective of the OMCHS is to clarify risk factors for childhood allergic disorders. We conducted a baseline survey to obtain information on prenatal environmental factors from pregnant women between November 2001 and March 2003. Details of the baseline survey of the OMCHS have been described elsewhere (Miyake et al., 2005). In brief, eligible females were those who became pregnant in Neyagawa City, which is 1 of the 44 municipalities in Osaka Prefecture, a metropolis in Japan with a total population of approximately 8.8 million. Of the 3639 eligible females in Neyagawa City, 627 (17.2%) participated in this study between November 2001 and March 2003. Eight pregnant females who did not live in Neyagawa City but who had become aware of the present study at an obstetric clinic before August 2002 decided by themselves to participate in this study. Also, there were 77 participants who received explanations regarding the OMCHS from public health nurses in 6 other municipalities between August 2002 and March 2003. From October 2002 to March 2003, 290 participants were recruited from a university hospital and three obstetric hospitals in three other municipalities; these women were recommended for participation in the OMCHS by an obstetrician. A total of 1002 pregnant women gave their fully informed consent in writing and completed the baseline survey. Of the 1002 females, 867 mothers participated in the second survey at 2–9 months postpartum. Two mothers were excluded because of missing data on the baby's birth weight. The final analysis comprised 865 subjects. The ethics committee of the Osaka City University School of Medicine approved the OMCHS.

2.2. Measurements

At baseline, each participant filled out a set of 2 self-administered questionnaires. Also, a self-administered questionnaire was used in the second survey. Participants mailed these completed questionnaires to the data management center at the time of each survey. Research

technicians completed missing or illogical data by telephone interview.

In the baseline survey, we collected information on dietary habits during the previous 1 month by using a validated self-administered diet history questionnaire. The structure and validity of the questionnaire were described in detail elsewhere (Sasaki et al., 1998, 2000). In this instrument, intake of 147 food items was calculated using an ad-hoc computer algorithm developed to analyze the questionnaire. Energy-adjusted intake by the residual approach was used for the analyses (Willett and Stampfer, 1986). Because only a small number of participants used calcium (5.3%), vitamin C (5.4%), vitamin B complex (4.3%), and multivitamin (3.9%) supplements at least once a week, use of these dietary supplements was not incorporated into the analysis in this study.

Another self-administered questionnaire at baseline inquired about age, gestation, parity, smoking habits, family structure, occupation, family income, education, weight, height, and changes in diet in the previous 1 month. Body mass index was calculated by dividing self-reported body weight (kg) by the square of self-reported height (m).

A self-administered questionnaire in the second survey included the Japanese version of the Edinburgh Postnatal Depression Scale (EPDS). The EPDS is a 10-item self-reported scale especially designed to screen for postpartum depression in community samples (Cox et al., 1987). Each item is scored on a 4-point scale (from 0 to 3) and the total score ranged from a minimum of 0 to a maximum of 30. The scale rates the intensity of depressive symptoms present within the previous 7 days. Although Cox and colleagues (Cox et al., 1987) proposed a cutoff level of 10 if the test is to be used for screening purposes in the postpartum period, a threshold score of 8/9 was found to detect depression among Japanese women with a specificity of 93% and sensitivity of 75% (Okano et al., 1996). Therefore, in the present study postpartum depression was defined as present when subjects had an EPDS score of 9 or higher. The questionnaire also elicited information on medical problems in pregnancy and sex, birth weight, and date of birth of the infant born after the baseline survey.

2.3. Statistical analysis

Intake of selected nutrients was categorized at quartile points based on the distribution among 865 pregnant women. Age was classified into 3 categories (<29, 29–31, and 32+ years); gestation into 3 (<15, 15–20, and 21+ weeks); parity into 2 (0 and 1+); cigarette smoking into 3 (never, former, and current); family structure into 2

(nuclear and expanded); occupation into 2 (outside work and housewife); family income into 3 (<4,000,000, 4,000,000–5,999,999, and 6,000,000+ yen/year); education into 3 (<13, 13–14, and 15+ years); changes in diet in the previous 1 month into 3 (none or seldom, slight, and substantial); season when data at baseline were collected into 4 (spring, summer, fall, and winter); time of delivery before the second survey into 2 (<4 and 4+ months); medical problems in pregnancy into 2 (yes and no); baby's sex into 2 (male and female); and baby's birth weight into 2 (<2500 and 2500+ g). Body mass index and energy-adjusted alcohol intake were used as continuous variables.

Logistic regression analysis was used to estimate crude odds ratios (ORs) and 95% confidence intervals (CIs) of postpartum depression in relation to consumption of selected dietary factors under study. Multiple logistic regression analysis was used to control for the potential confounding factors. Trend of association was assessed by a logistic regression model assigning consecutive integers (1 to 4) to the levels of the independent variable. Two-sided *P*-values less than 0.05 were regarded as statistically significant. All computations were performed using the SAS software, version 9.1 (SAS Institute, Inc., Cary, NC).

3. Results

About 30% of subjects were from 29 to 31 years of age at baseline. Approximately 70% of the women took part in the baseline survey by the 20th week of gestation and around half had a parity of 1 or more at baseline. Infants of 6% of mothers had a birth weight less than 2500 g. There were 24 premature deliveries (24 to 36 weeks of gestation) and 816 babies were born at 37 to 41 weeks of gestation. Mean daily total energy and energy-adjusted consumption of folate, cobalamin, pyridoxine, and riboflavin were 7752 kJ, 286.1 µg, 5.7 µg, 1.0 mg, and 1.4 mg, respectively.

Of the 865 mothers, 121 subjects (14.0%) developed postpartum depression at 2–9 months postpartum. No measurable association was observed between intake of folate, cobalamin, or pyridoxine and the risk of postpartum depression (Table 1). Adjustment for age, gestation, parity, cigarette smoking, alcohol intake, family structure, family income, education, changes in diet in the previous 1 month, season when data at baseline were collected, body mass index, time of delivery before the second survey, medical problems in pregnancy, baby's sex, and baby's birth weight did not appreciably change these findings. Compared with riboflavin intake in the first quartile, riboflavin consumption in the second and third quartiles, but not the fourth quartile, was significantly related to a decreased risk of

Table 1

Odds ratios and 95% CIs for postpartum depression according to quartile of intake of selected vitamins in 865 women, the Osaka Maternal Child Health Study, Japan^a

Variable ^b	Quartile				P for trend
	1	2	3	4	
<i>Folate</i>					
Intake (μg/day)	204.8	255.4	294.5	377.3	
No. of cases	31	25	29	36	
Crude odds ratio (95% CI)	1.00	0.78 (0.44–1.37)	0.93 (0.53–1.60)	1.19 (0.71–2.01)	0.42
Multivariate odds ratio (95% CI)	1.00	0.81 (0.45–1.45)	0.99 (0.56–1.77)	1.20 (0.68–2.13)	0.40
<i>Cobalamin (vitamin B₁₂)</i>					
Intake (μg/day)	3.3	4.6	5.9	8.3	
No. of cases	37	29	27	28	
Crude odds ratio (95% CI)	1.00	0.75 (0.44–1.27)	0.69 (0.40–1.18)	0.72 (0.42–1.22)	0.20
Multivariate odds ratio (95% CI)	1.00	0.75 (0.43–1.28)	0.73 (0.41–1.28)	0.73 (0.42–1.29)	0.29
<i>Pyridoxine (vitamin B₆)</i>					
Intake (mg/day)	0.7	0.9	1.0	1.2	
No. of cases	32	23	28	38	
Crude odds ratio (95% CI)	1.00	0.69 (0.38–1.21)	0.86 (0.49–1.48)	1.22 (0.73–2.05)	0.32
Multivariate odds ratio (95% CI)	1.00	0.75 (0.41–1.35)	0.95 (0.53–1.69)	1.30 (0.75–2.28)	0.25
<i>Riboflavin (vitamin B₂)</i>					
Intake (mg/day)	1.0	1.2	1.4	1.7	
No. of cases	40	25	22	34	
Crude odds ratio (95% CI)	1.00	0.58 (0.33–0.98)	0.50 (0.28–0.87)	0.82 (0.49–1.35)	0.35
Multivariate odds ratio (95% CI)	1.00	0.61 (0.34–1.06)	0.53 (0.29–0.95)	0.88 (0.51–1.50)	0.55

^a Values for intake are medians for adjusted energy intake by using the residual method for each quartile.

^b The multivariate models included the following: age (<29, 29–31, and 32+ years), gestation (<15, 15–20, and 21+ weeks), parity (0 and 1+), cigarette smoking (never, former, and current), energy-adjusted alcohol intake (continuous), family structure (nuclear and expanded), family income (<4,000,000, 4,000,000–5,999,999, and 6,000,000+ yen/year), education (<13, 13–14, and 15+ years), changes in diet in the previous 1 month (none or seldom, slight, and substantial), season when data at baseline were collected (spring, summer, fall, and winter), body mass index (continuous), time of delivery before the second survey (<4 and 4+ months), medical problems in pregnancy (yes and no), baby's sex (male and female), and baby's birth weight (<2500 and 2500+ g).

postpartum depression. The inverse relationship was slightly attenuated after multivariate adjustment; only riboflavin consumption in the third quartile was independently inversely associated with the risk of postpartum depression (multivariate OR: 0.53, 95% CI: 0.29–0.95, *P* for trend=0.55).

4. Discussion

Folate and cobalamin are involved in the 1-carbon cycle (methylation) that furnishes *S*-adenosyl-methionine, the sole methyl donor for a broad range of reactions necessary for the synthesis of neuroactive substances, the formation of membrane phospholipids, and the metabolism of nucleic acids (Alpert et al., 2000; Bjelland et al., 2003a). Folate also appears to influence the biosynthesis of bipterin-dependent neurotransmitters (Alpert et al., 2000; Bjelland et al., 2003a). Methyltetrahydrofolate may have a neuromodulatory role at presynaptic sites by binding to glutamate receptors (Alpert et al., 2000;

Bjelland et al., 2003a). Elevated levels of homocysteine, resulting from folate deficiency, may play a role in mediating some of the neuropsychiatric complications due to folate deficiency (Alpert et al., 2000; Bjelland et al., 2003a). The synthesis of serotonin and the catecholamines is dependent on the metabolite of pyridoxine (Hvas et al., 2004). Riboflavin coenzymes are needed for both the remethylation and transsulfuration of homocysteine (Ganji and Kafai, 2004).

In the present study, an inverse relationship between the intake of folate, cobalamin, or pyridoxine and the risk of postpartum depression was not observed. The findings are at variance with previous studies that showed an inverse association between folate intake and depressive symptoms (Tolmunen et al., 2003, 2004). We also found an inverted J-shaped association between riboflavin intake and postpartum depression. Dietary intake of riboflavin and folate, but not cobalamin and pyridoxine, was independently related to decreased concentrations of serum homocysteine in the third National Health and

Examination Survey (Ganji and Kafai, 2004). Hyperhomocysteinemia may play a critical role in mediating the relationship between a lower intake of riboflavin and the risk of postpartum depression. In this context, our findings may be in partial agreement with this observation regarding riboflavin.

The present study had several methodological strengths. The prospective design and relatively high rate of follow-up (86.3%) in this study minimized the possibility of recall bias or bias caused by loss of follow-up. Study subjects were homogeneous in terms of having the same residential background. Extensive information on potential confounding factors was adjusted for. However, we were not able to control for personal and family psychiatric history, sociocultural factors, and personal and family relations.

There are several weaknesses. The diagnosis of postpartum depression was established with the EPDS. Moreover, although the second survey was conducted at 2–9 months postpartum, a large majority of subjects, 89.2%, participated in the second survey at 3–4 months postpartum. Thus, it is difficult to estimate the incidence of postpartum depression accurately. The consequence would introduce a bias toward the null because of nondifferential outcome misclassification. Dietary data were obtained using a self-administered semiquantitative dietary assessment questionnaire. Since we did not assess the actual dietary habits of the subjects over a long period of time, the possibility of misclassification should be considered. Our diet history questionnaire was designed to assess recent dietary intake, i.e. for 1 month prior to completing the questionnaire. This disadvantage is likely to be alleviated after adjustment for the season when baseline data were collected, however. Changes in diet in the past 1 month were controlled for because pregnant females are likely to change their diet for reasons such as nausea gravidarum. We found no significant interaction between changes in diet in the past 1 month or timing of the baseline assessment and exposures under investigation in relation to postpartum depression (data not shown). Potential misclassification associated with variability across subjects in terms of the stability of diet during pregnancy and the timing of the dietary assessment is likely to be negligible. Thus, negative findings of our study could not simply mean that diet was assessed at the wrong time. The study subjects were an unrepresentative sample of Japanese females in the general population, and the present findings may not be generalized. In fact, educational levels were higher in the present study population than in the general population. According to the 2000 population census of Japan, the proportions of females aged 30 to 34 years in Osaka Prefecture with

years of education of <13, 13–14, 15+, and unknown were 49.2, 32.3, 13.6, and 4.9%, respectively (Statistic Bureau, Ministry of Public Management, Home Affairs, Posts and Telecommunications, Japan, 2002). The corresponding figures for the current study were 29.7, 42.4, 27.9, and 0.0%, respectively.

This is the first prospective study on intake of folate and B vitamins and the risk of postpartum depression. Our results suggest that moderate riboflavin consumption may be protective against postpartum depression. However, these results are preliminary. Further studies with more precise exposure and outcome measurements are warranted to determine whether moderate consumption of riboflavin is protective against postpartum depression, taking into consideration additional factors such as psychiatric history and sociocultural status.

Acknowledgments

The authors would like to acknowledge the Neyagawa City Government, Hirakata City Government, Katano City Government, Shijonawate City Government, Kaizuka City Government, Takaishi City Government, Hannan City Government, Neyagawa City Medical Association, Hirakata City Medical Association, and Kadoma City Medical Association for their valuable support; Kumiko Shirata, PhD, Osaka City University School of Nursing for helpful discussions; and Ms. Tomoko Shibazaki, Nahoko Nishimura, and Naomi Takaoka for their assistance. This study is supported by a Grant-in-Aid (13770206, 16790351) for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology and Health and Labour Sciences Research Grants, Research on Allergic Disease and Immunology from the Ministry of Health, Labour, and Welfare, Japan.

Appendix A

Other members of the Osaka Maternal and Child Health Study Group were as follows: Mr. Shoichi Miyamoto, Mr. Ichiro Matsunaga, Dr. Hajime Oda, Dr. Hideharu Kanzaki, Dr. Mitsuyoshi Kitada, Dr. Yorihiro Horikoshi, Dr. Osamu Ishiko, Dr. Yuichiro Nakai, Dr. Junko Nishio, Dr. Seiichi Yamamasu, Dr. Jinsuke Yasuda, Dr. Seigo Kawai, Dr. Kazumi Yanagihara, Dr. Koji Wakuda, Dr. Tokio Kawashima, Dr. Katsuhiko Narimoto, Dr. Yoshihiko Iwasa, Dr. Katsuhiko Orino, Dr. Itsuo Tsunetoh, Dr. Junichi Yoshida, Dr. Junichi Iito, Dr. Takuzi Kaneko, Dr. Takao Kamiya, Dr. Hiroyuki Kuribayashi, Dr. Takeshi Taniguchi, Dr. Hideo Takemura, and Dr. Yasuhiko Morimoto.

References

- Abou-Saleh, M.T., Coppen, A., 1989. Serum and red blood cell folate in depression. *Acta Psychiatr. Scand.* 80, 78–82.
- Alpert, J.E., Mischoulon, D., Nierenberg, A.A., Fava, M., 2000. Nutrition and depression: focus on folate. *Nutrition* 16, 544–546.
- Bjelland, I., Ueland, P.M., Vollset, S.E., 2003a. Folate and depression. *Psychother. Psychosom.* 72, 59–60.
- Bjelland, I., Tell, G.S., Vollset, S.E., Refsum, H., Ueland, P.M., 2003b. Folate, vitamin B₁₂, homocysteine, and the MTHFR 677C→T polymorphism in anxiety and depression: the Hordaland Homocysteine Study. *Arch. Gen. Psychiatry* 60, 618–626.
- Bottiglieri, T., Laundy, M., Crellin, R., Toone, B.K., Carney, M.W.P., Reynolds, E.H., 2000. Homocysteine, folate, methylation, and monoamine metabolism in depression. *J. Neurol. Neurosurg. Psychiatry* 69, 228–232.
- Carney, M.W.P., Chary, T.K.N., Laundy, M., Bottiglieri, T., Chanarin, I., Reynolds, E.H., Toone, B., 1990. Red cell folate concentrations in psychiatric patients. *J. Affect. Disord.* 19, 207–213.
- Cox, J.L., Holden, J.M., Sagovsky, R., 1987. Detection of postnatal depression. Development of the 10-item Edinburgh Postnatal Depression Scale. *Br. J. Psychiatry* 150, 782–786.
- Ganji, V., Kafai, M.R., 2004. Frequent consumption of milk, yogurt, cold breakfast cereals, peppers, and cruciferous vegetables and intakes of dietary folate and riboflavin but not vitamins B-12 and B-6 are inversely associated with serum total homocysteine concentrations in the US population. *Am. J. Clin. Nutr.* 80, 1500–1507.
- Ghadirian, A.M., Ananth, J., Engelsmann, F., 1980. Folic acid deficiency and depression. *Psychosomatics* 21, 926–929.
- Hvas, A.M., Juul, S., Bech, P., Nexø, E., 2004. Vitamin B₆ level is associated with symptoms of depression. *Psychother. Psychosom.* 73, 340–343.
- Lee, S., Wing, Y.K., Fong, S., 1998. A controlled study of folate levels in Chinese inpatients with major depression in Hong Kong. *J. Affect. Disord.* 49, 73–77.
- Lindeman, R.D., Romero, L.J., Koehler, K.M., Liang, H.C., LaRue, A., Baumgartner, R.N., Garry, P.J., 2000. Serum vitamin B₁₂, C and folate concentrations in the New Mexico elder health survey: correlations with cognitive and affective functions. *J. Am. Coll. Nutr.* 19, 68–76.
- Miyake, Y., Sasaki, S., Ohya, Y., Miyamoto, S., Matsunaga, I., Yoshida, T., Hirota, Y., Oda, H., The Osaka Maternal and Child Health Study Group, 2005. Soy, isoflavones, and prevalence of allergic rhinitis in Japanese women: the Osaka Maternal and Child Health Study. *J. Allergy Clin. Immunol.* 115, 1176–1183.
- Morris, M.S., Fava, M., Jacques, P.F., Selhub, J., Rosenberg, I.H., 2003. Depression and folate status in the US population. *Psychother. Psychosom.* 72, 80–87.
- Okano, T., Murata, M., Masuji, F., Tamaki, R., Nomura, J., Miyaoka, H., Kitamura, T., 1996. Validation and reliability of Japanese version of the EPDS. *Arch. Psychiatr. Diag. Clin. Eval.* 7, 525–533.
- Penninx, B.W.J.H., Guralnik, J.M., Ferrucci, L., Fried, L.P., Allen, R. H., Stabler, S.P., 2000. Vitamin B₁₂ deficiency and depression in physically disabled older women: epidemiologic evidence from the Women's Health and Aging Study. *Am. J. Psychiatry* 157, 715–721.
- Ramos, M.I., Allen, L.H., Haan, M.N., Green, R., Miller, J.W., 2004. Plasma folate concentrations are associated with depressive symptoms in elderly Latina women despite folic acid fortification. *Am. J. Clin. Nutr.* 80, 1024–1028.
- Sasaki, S., Yanagibori, R., Amano, K., 1998. Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *J. Epidemiol.* 8, 203–215.
- Sasaki, S., Ushio, F., Amano, K., Morihara, M., Todoriki, T., Uehara, Y., Toyooka, T., 2000. Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J. Nutr. Sci. Vitaminol.* 46, 285–296.
- Statistic Bureau, Ministry of Public Management, Home Affairs, Posts and Telecommunications, Japan, 2002. 2000 population census of Japan, Vol. 3-2-27, labour force status of population, industry (major groups) of employed persons, and education. Osaka-fu, Statistic Bureau, Ministry of Public Management, Home Affairs, Posts and Telecommunications, Japan, Tokyo.
- Tiemeier, H., van Tuijl, H.R., Hofman, A., Meijer, J., Kiliaan, A.J., Breteler, M.M.B., 2002. Vitamin B₁₂, folate, and homocysteine in depression: the Rotterdam Study. *Am. J. Psychiatry* 159, 2099–2101.
- Tolmunen, T., Voutilainen, S., Hintikka, J., Rissanen, T., Tanskanen, A., Viinamaki, H., Kaplan, G.A., Salonen, J.T., 2003. Dietary folate and depressive symptoms are associated in middle-aged Finnish men. *J. Nutr.* 133, 3233–3236.
- Tolmunen, T., Hintikka, J., Ruusunen, A., Voutilainen, S., Tanskanen, A., Valkonen, V.P., Viinamaki, H., Kaplan, G.A., Salonen, J.T., 2004. Dietary folate and the risk of depression in Finnish middle-aged men. A prospective follow-up study. *Psychother. Psychosom.* 73, 334–339.
- Willett, W., Stampfer, M.J., 1986. Total energy intake: implications for epidemiologic analyses. *Am. J. Epidemiol.* 124, 17–27.

Magnesium intake is inversely associated with the prevalence of tooth loss in Japanese pregnant women: the Osaka Maternal and Child Health Study

K. Tanaka¹, Y. Miyake¹, S. Sasaki², Y. Ohya³, S. Miyamoto⁴, I. Matsunaga⁵, T. Yoshida⁵, Y. Hirota⁴, H. Oda⁵, the Osaka Maternal and Child Health Study Group⁶

¹ Department of Public Health, Faculty of Medicine, Fukuoka University, Fukuoka, Japan; ² Nutritional Epidemiology Program, National Institute of Health and Nutrition, Tokyo, Japan; ³ Division of Allergy, Department of Medical Specialties, National Center for Child Health and Development, Tokyo, Japan; ⁴ Department of Public Health, Osaka City University School of Medicine, Osaka, Japan; ⁵ Osaka Prefectural Institute of Public Health, Osaka, Japan; ⁶ Other members of the Osaka Maternal and Child Health Study Group are listed in the Appendix

Correspondence: K. Tanaka, DDS, PhD, Department of Public Health, Faculty of Medicine, Fukuoka University, Fukuoka 814-0180, Japan. Tel.: (+00 81) 92 801 1011 (ext. 3315); fax: (+00 81) 92 863 8892 <k-tanaka@fukuoka-u.ac.jp>

Abstract. There have only been a few studies on the role of mineral intake in tooth loss. We investigated the association between mineral intake and the prevalence of tooth loss in Japan. We used the baseline data on 1002 pregnant women who were enrolled in the Osaka Maternal and Child Health Study between November 2001 and March 2003. Tooth loss was defined as the previous extraction of one or more teeth. Nutrient intake was assessed by a validated diet history questionnaire. Prevalence *odds ratios* and confidence intervals were estimated by applying a multiple logistic regression model. The adjusted *odds ratio* upon comparison of the highest quartile with the lowest quartile of magnesium intake was 0.64 (95% confidence interval, 0.42-0.99), showing a tendency for an inverse dose-response relationship (*p* for linear trend = 0.05). There were no associations between the level of consumption of calcium, phosphate, iron, zinc, or copper and tooth loss. The present findings suggest that intake of magnesium is related to reduced prevalence of tooth loss among young Japanese women.

Key words: cross-sectional study, Japanese women, magnesium, tooth loss

There is a two-way relationship between diet and oral health: a balanced diet with adequate energy and nutrients is essential for oral health, and the condition of the oral cavity may impact an individual's diet and nutritional status [1]. Many previous studies on the association between tooth loss and nutrient intake have focused on nutritional deficiency such as decreased consumption of vegetables, fruits, dietary fiber and antioxidant nutrients

among the middle-aged and elderly who retain a few teeth or are edentulous [2-11]. There have been a few studies on the relationship between tooth loss and low mineral intake among elderly individuals [4, 6, 11]. A U.S. cross-sectional study of elderly people demonstrated that edentulism was associated with lower consumption of calcium and magnesium compared with the respective levels in the dentate [6].

A paucity of information exists on the possible role that mineral intake plays in oral diseases. In a longitudinal study of black South African children, low magnesium intake was significantly associated with caries in deciduous teeth, although no associations were found for consumption of calcium, iron, phosphorus, zinc, or copper [12]. A cross-sectional study among Swedish adolescents found that iron intake was inversely associated with the prevalence of dental caries [13]. The Third National Health and Nutrition Examination Survey in the U.S. showed that low dietary intake of calcium was significantly associated with severe periodontitis [14]. Freeland and colleagues [15] reported that dietary calcium intake was significantly and inversely associated with the severity of periodontitis, although no association with serum concentrations of magnesium was observed among 80 dental patients. One interventional study indicated that taking calcium and vitamin D supplements was significantly associated with a lower risk of tooth loss among the elderly [16]. There has been no epidemiologic study on the association between mineral intake and tooth loss in Japan. Therefore, we examined the relationship between the level of mineral intake and tooth loss among pregnant Japanese women using the baseline data from the Osaka Maternal and Child Health Study (OMCHS).

Methods

Study population

The subjects of the present study were pregnant Japanese women who were enrolled in the OMCHS between November 2001 and March 2003. Details of the OMCHS were described previously [17, 18]. Briefly, the OMCHS is an ongoing prospective cohort study that examines preventive factors and risk factors for maternal and child health problems, such as allergic disorders. In Japan, when women become pregnant, they notify the municipality of the domicile of the conception, and the municipality provides them with a maternal and child health handbook. Women who were eligible for the OMCHS were those who lived in Neyagawa City, a municipality in Osaka Prefecture, Japan. The Neyagawa City Government provided all pregnant women with a set of leaflets explaining our study, an application form, and a self-addressed and stamped return envelope together with the maternal and child health handbook. Among the 3639 women, 627 pregnant women (17.2%) consented to participate in the OMCHS. Eight pregnant

women who did not live in Neyagawa City but who became aware of the OMCHS at an obstetric clinic before August 2002, decided by themselves to participate in the OMCHS. Seventy-seven women received explanations of the OMCHS from public health nurses in 6 other municipalities from August 2002 to March 2003, and were enrolled in the OMCHS. From October 2002 to March 2003, 290 women were recruited from a university hospital and three obstetric hospitals in 3 other municipalities; it had been recommended that these women participate in the OMCHS by an obstetrician. Therefore, a total of 1002 pregnant women gave written informed consent to participate in the OMCHS between November 2001 and March 2003, and completed the two baseline questionnaires. The ethics committees of the Osaka City University School of Medicine and the Osaka Prefectural Institute of Public Health approved the OMCHS.

Questionnaire

At the time of enrollment in the OMCHS, each participant filled out two self-administered questionnaires. The participants mailed the questionnaires to the data management center. Research technicians completed missing and illogical data in the questionnaires by telephone interview.

The first questionnaire was a validated self-administered diet history questionnaire that was used to assess dietary habits over the previous one month. The structure and validity of this questionnaire were described previously [19, 20]. In this instrument, the intake of 147 food items is calculated using an *ad-hoc* computer algorithm that had been developed to analyze the questionnaire. Energy-adjusted intake was calculated by the residual method [21] and used in the analyses.

The second self-administered questionnaire elicited information on age, gestation, parity, smoking habit, passive smoking at home and at work, family income, education, height, weight, changes in diet in the previous one month, experience of extraction of one or more permanent teeth excluding third molars, and the number of remaining teeth. Tooth loss was defined as the extraction of one or more teeth. Body mass index was calculated by dividing body weight (kg) by the square of the height of the individual (m).

Statistical methods

The study population was divided into quartiles with regard to the intake of each selected nutrient. The logistic regression model was used to estimate crude *odds ratios* (ORs) and their 95% confidence intervals

(CIs) for tooth loss relative to intake of selected minerals. Multiple logistic regression analysis was used to control for potential confounders. Adjustment was made for age, gestation, parity, cigarette smoking, passive smoking at home and at work, family income, education, changes in diet in the previous one month, season when data were collected, and body mass index. Active and passive smoking exposure was controlled for because our previous study showed a significant positive association between active and passive smoking status and the prevalence of tooth loss [18]. Trend of association was assessed by a logistic model assigning scores to the levels of the independent variable. Analyses were conducted using SAS software (version 9.1, SAS Institute Inc., Cary, NC, USA). All p-values are 2-tailed. $p < 0.05$ was considered to be statistically significant.

Results

The 1002 pregnant women in this study ranged in age from 15 to 44 years (mean age, 29.8 years; median age, 30 years) at the time of enrollment in the OMCHS. Approximately 30% of the subjects were between 29 and 31 years old (*table 1*). Approximately 70% were enrolled by the 20th week of gestation, and about 50% had a parity of one or more. Seven hundred two pregnant women (70.1%) had experienced slight or substantial changes in diet in the previous one month due to nausea gravidarum (585 women), maternal and fetal health (107 women), or other reason (10 women). The mean daily intake of total energy was 6815 kJ. Among the 1002 pregnant women, 256 women (25.5%) had previously had one or more teeth extracted (*table 2*).

Table 3 shows the ORs and their 95% CIs for tooth loss according to the level of consumption of various minerals. Compared with magnesium intake in the lowest quartile, its consumption in the highest quartile was significantly associated with a decreased prevalence of tooth loss, showing a clear inverse dose-response relationship. After adjustment for the confounders under investigation, the inverse association remained significant although the linear trend fell just short of significance (adjusted OR in the highest quartile: 0.64, 95% CI: 0.42-0.99, p for linear trend = 0.05). There were no appreciable relationships between the level of intake of calcium, phosphate, iron, zinc, or copper and tooth loss.

Discussion

The current cross-sectional study found that intake of magnesium, but not calcium, phosphate, iron, zinc or copper, was significantly associated with a reduced prevalence of tooth loss. The present results are in partial agreement with a previous study that found an association between low magnesium intake and dental caries in children [12], but are not consistent with studies that found that calcium intake was inversely associated with periodontitis [14, 15]. The potential underlying mechanism for the observed inverse relationship between magnesium intake and the prevalence of tooth loss is not known. Magnesium *per se* may prevent tooth loss. A cross-sectional study among middle-aged and older U.S. women showed that magnesium intake was inversely associated with systemic inflammation [22]. Our previous study showed a tendency for an inverse relationship between magnesium intake and the prevalence of allergic rhinitis in the present subjects [23]. Magnesium intake might influence oral health thorough improving immunocompetence and reducing systemic inflammation. Magnesium deficiency caused loss of structure of periodontal tissues in adult rats [24]. On the other hand, the observed inverse association between magnesium intake and tooth loss may be ascribed to unknown nutritional factors related to bone metabolism. Magnesium is involved in bone and mineral homeostasis and is important in bone crystal growth and stabilization [25]. In a cross-sectional study of 62 healthy women aged 45-55 years in the U.S., intake of magnesium, potassium, and alcohol was significantly associated with increased bone mineral density [26]. There were significant linear relationships between the bone mineral density at the spine or radius and the total number of teeth among 329 healthy postmenopausal women aged 41-71 years [27]. Another possible explanation is that unknown factors related to magnesium intake produce a spurious inverse association between magnesium intake and tooth loss. Mineral intake in Japanese people may be associated with part of a larger pattern of traditional Japanese dietary customs or behaviors. Legumes and seaweed, which are the richest dietary sources of magnesium, are part of the traditional Japanese diet.

In this investigation, we used self-reported tooth loss as a measure of dental health status. We did not assess the validity of self-reported tooth loss in our subjects. However, in several validation studies of self-reported oral health, there was substantial agreement between the self-reported and observed number of remaining teeth regardless of age or sex

Table 1. Distribution of the baseline characteristics of the 1002 pregnant women, OMCHS^a, Japan^b.

Factor	Number (%) or mean (SD) ^c
Age (years)	
< 29	380 (37.9)
29–31	299 (29.8)
32+	323 (32.4)
Gestation (weeks)	
< 15	357 (35.6)
15–20	329 (32.8)
21+	316 (31.5)
Parity of one or more	513 (51.2)
Cigarette smoking	
Never	697 (69.6)
Former	121 (12.1)
Current	184 (18.4)
Passive smoking at home	
Never	284 (28.3)
Former	224 (22.4)
Current	494 (49.3)
Passive smoking at work	
Never	344 (34.3)
Former	538 (53.7)
Current	120 (12.0)
Family income (JPY^d/year)	
< 4,000,000	301 (30.0)
4,000,000–5,999,999	403 (40.2)
6,000,000+	298 (29.7)
Education (years)	
< 13	323 (32.2)
13–14	413 (41.2)
15+	266 (26.6)
Change in diet in the previous one month	
None or seldom	300 (29.9)
Slight	435 (43.4)
Substantial	267 (26.7)
Season when data were collected	
Spring	318 (31.7)
Summer	162 (16.2)
Fall	223 (22.3)
Winter	299 (29.8)
Body mass index (kg/m²)	21.4 (2.8)
Daily nutrient intake^e	
Total energy (kJ)	6815.3 (1793.8)
Calcium (mg)	556.0 (182.9)
Phosphate (mg)	890.0 (183.8)
Magnesium (mg)	194.4 (49.9)
Iron (mg)	7.5 (1.7)
Zinc (µg)	7222.8 (1320.7)
Copper (µg)	1094.9 (316.9)

^a OMCHS, Osaka Maternal and Child Health Study.^b Characteristics of the women at the time of entry into the OMCHS are shown.^c SD, Standard deviation.^d JPY, Japanese Yen, US \$1 = ¥118.^e Nutrient intake was adjusted for total energy intake by the residual method.

Table 2. Distribution of tooth loss among the 1002 pregnant women, OMCHS^a, Japan.

Number of teeth lost	Number of subjects (%)
0	746 (74.5)
1	85 (8.5)
2	57 (5.7)
3	29 (2.9)
4	43 (4.3)
5+	42 (4.2)

^a OMCHS, Osaka Maternal and Child Health Study.

[28-30]. These results indicated that self-reported oral status can provide reasonably valid data on the remaining teeth. Hence, the use of self-reported tooth loss as a key indicator in epidemiological studies on dental health can be considered satisfactory.

The strengths of the current study include the homogeneity of the study subjects in that they consisted of pregnant Japanese women, and the comprehensive assessment of potential confounders. All dietary data were self-reported. However, since we did not actually observe the dietary habits of the subjects, there is a possibility of misclassification. According to the validation study of Sasaki and colleagues [19], the correlation coefficients of nutrient intake between those estimated from the diet history questionnaire and those observed in a 3-day dietary record in women were 0.49, 0.59, and 0.40 for calcium, phosphorus, and iron, respectively. The subjects with tooth loss in our study might not be aware of the possible ill effects of diet. The consequence would have been underestimation of values in our results due to nondifferential exposure misclassification. Japanese people, as well as European people, generally like seasonal ingredients in cooking. The diet history questionnaire used in the OMCHS was designed to assess recent dietary intake, i.e., the subject's diet during the one-month period prior to filling out this questionnaire. Allowance for the season when data were collected likely eased this weakness, however. We controlled for changes in the diet in the past one month because pregnant women are likely to change their diet for reasons such as nausea gravidarum. There are other limitations that may be of concern. Only 627 (17.2%) of the 3639 eligible pregnant women in Neyagawa City participated in the OMCHS. Regarding the remaining 375 subjects in the OMCHS who live in municipalities other than Neyagawa City, we can not calculate the participation rate because data on the exact number of eligible subjects were not available. Our sample cannot

be considered representative of Japanese women. In fact, the educational level of our subjects was higher than that of the general population according to the 2000 population census of Japan [31]. However, the prevalence of tooth loss among our subjects (25.5%) was similar to that among Japanese women aged 25-30 years in a survey of dental diseases conducted in 1999 (27.3%) [32]. A study on the reasons for extraction of permanent teeth found that among Japanese individuals aged 9-35 years the proportion of extractions resulting from caries, periodontitis, eruption problems, orthodontic indications, trauma, and other causes were 51.5%, 6.2%, 21.9%, 5.1%, 0.1%, and 15.2%, respectively, and that more than 90% of extractions for eruption problems were third molars [33]. Among the 256 women with tooth loss, the tooth loss in more than 50% of the cases likely resulted from dental diseases although we do not have information on the reasons for tooth extraction because tooth loss was not a principal study outcome of the OMCHS. However, any outcome misclassification would likely to have been evenly distributed across the four categories of the exposures under study. Consequently, this would have given rise to an underestimation of our findings. Information regarding the point in time at which tooth loss occurred was not available. Therefore, the time sequence between mineral intake and tooth loss could not be determined. Although we adjusted our estimate for a wide range of potential confounders, we can not rule out the possibility that our findings were biased by unmeasured confounders such as the level of oral hygiene, caries, periodontitis, and oral health behavior including the frequency of toothbrushing and the pattern of dental visits. Because the outcome variable in our study was tooth loss, which is the ultimate endpoint of caries and periodontitis, early symptoms and subclinical symptoms of dental disease could not be taken into consideration in our estimate. The consequence would be underestimation of our results.

Table 3. Odds ratios for tooth loss by quartiles of intake of minerals, OMCHS^a, Japan.

Variable ^b	Prevalence	Crude odds ratio (95% CI ^c)		Adjusted odds ratio ^d (95% CI ^c)	
Calcium					
Q1 (368.0)	69/250 (27.6%)	1.00		1.00	
Q2 (491.7)	71/251 (28.3%)	1.04	0.70–1.53	1.12	0.74–1.69
Q3 (590.3)	59/250 (23.6%)	0.81	0.54–1.21	0.85	0.55–1.30
Q4 (739.4)	57/251 (22.7%)	0.77	0.51–1.16	0.81	0.53–1.25
p for linear trend		0.12		0.20	
Phosphate					
Q1 (713.1)	74/250 (29.6%)	1.00		1.00	
Q2 (828.7)	65/251 (25.9%)	0.83	0.56–1.23	0.83	0.55–1.26
Q3 (928.1)	55/250 (22.0%)	0.67	0.45–1.00	0.73	0.47–1.12
Q4 (1085.0)	62/251 (24.7%)	0.78	0.53–1.16	0.78	0.51–1.20
p for linear trend		0.13		0.21	
Magnesium					
Q1 (150.8)	76/250 (30.4%)	1.00		1.00	
Q2 (175.4)	64/251 (25.5%)	0.78	0.53–1.16	0.85	0.57–1.29
Q3 (199.4)	62/250 (24.8%)	0.76	0.51–1.12	0.83	0.55–1.26
Q4 (245.8)	54/251 (21.5%)	0.63	0.42–0.94	0.64	0.42–0.99
p for linear trend		0.03		0.05	
Iron					
Q1 (5.9)	73/250 (29.2%)	1.00		1.00	
Q2 (7.0)	63/251 (25.1%)	0.81	0.55–1.21	0.80	0.53–1.21
Q3 (7.9)	62/250 (24.8%)	0.80	0.54–1.19	0.83	0.55–1.26
Q4 (9.3)	58/251 (23.1%)	0.73	0.49–1.09	0.70	0.45–1.07
p for linear trend		0.13		0.14	
Zinc					
Q1 (5936.5)	77/250 (30.8%)	1.00		1.00	
Q2 (6816.2)	54/251 (21.5%)	0.62	0.41–0.92	0.68	0.44–1.03
Q3 (7493.8)	64/250 (25.6%)	0.77	0.52–1.14	0.84	0.56–1.28
Q4 (8608.4)	61/251 (24.3%)	0.72	0.49–1.07	0.78	0.51–1.19
p for linear trend		0.21		0.42	
Copper					
Q1 (819.2)	72/250 (28.8%)	1.00		1.00	
Q2 (977.7)	66/251 (26.3%)	0.88	0.60–1.31	0.88	0.59–1.33
Q3 (1104.0)	57/250 (22.8%)	0.73	0.49–1.09	0.72	0.47–1.10
Q4 (1415.8)	61/251 (24.3%)	0.79	0.53–1.18	0.78	0.50–1.20
p for linear trend		0.17		0.17	

^a OMCHS, Osaka Maternal and Child Health Study.^b Median in each quartile in mg/d (except for zinc and copper; µg/d) that had been adjusted for energy intake by the residual method, are given in parentheses.^c CI, Confidence interval.^d Odds ratios were calculated separately for each dietary variable and were adjusted for age (< 29, 29–31, and 32+ yr), gestation (< 15, 15–20, and 21+ weeks), parity (0 and 1+), cigarette smoking (never, former, and current), passive smoking at home (never, former and current), passive smoking at work (never, former, and current), family income (< 4,000,000, 4,000,000–5,999,999, and 6,000,000+ JPY/year), education (< 13, 13–14, and 15+ years), changes in diet in the previous one month (none or seldom, slight, and substantial), season when data were collected (spring, summer, fall, and winter), and body mass index as a continuous variable.

Appendix

Space limitations preclude the inclusion as authors of the following members of the Osaka Maternal and Child Health Study Group:

Hideharu Kanzaki, Mitsuyoshi Kitada, Department of Obstetrics and Gynecology, Kansai Medical University; Yorihiro Horikoshi, Department of Obstetrics and Gynecology, Kansai Medical University Kori; Osamu Ishiko, Yuichiro Nakai, Junko Nishio, Seiichi Yamamasu, Department of Obstetrics and Gynecology, Osaka City University Graduate School of Medicine; Jinsuke Yasuda, Department of Obstetrics and Gynecology, Matsushita Memorial Hospital; Seigo Kawai, Department of Obstetrics and Gynecology, Hoshigaoka Koseinenkin Hospital; Kazumi Yanagihara, Yanagihara Clinic; Koji Wakuda, Department of Obstetrics and Gynecology, Fujimoto Hospital; Tokio Kawashima, Kyohritsu Women's Clinic; Katsuhiko Narimoto, Ishida Hospital Obstetrics, Gynecology; Yoshihiko Iwasa, Iwasa Women's Clinic; Katsuhiko Orino, Orino Lady's Clinic; Itsuo Tsunetoh, Tsunetoh Obstetrics and Gynecology; Junichi Yoshida, Yoshida Clinic; Junichi Ito, Ito Obstetrics and Gynecology Clinic; Takuzi Kaneko, Kaneko Sanfujinka; Takao Kamiya, Kamiya Ladies Clinic; Hiroyuki Kuribayashi, Kuribayashi Clinic; Takeshi Taniguchi, Taniguchi Hospital; Hideo Takemura, Kosaka Women's Hospital; Yasuhiko Morimoto, Aizenbashi Hospital.

Conclusion

This is the first epidemiological study that assessed the effects of intake of various minerals on tooth loss among young adults using a well-validated diet history questionnaire. We found that magnesium intake was inversely associated with the prevalence of tooth loss in pregnant Japanese women. Since this study was a cross-sectional study, only associations rather than causal relationships between magnesium intake and tooth loss can be inferred. Further studies with detailed and objective indicators of dental health status such as caries and periodontitis are needed to assess the beneficial effects of magnesium intake on dental diseases.

Acknowledgments

The authors acknowledge the Neyagawa City Government, Hirakata City Government, Katano City Government, Shijonawate City Government, Kaizuka City Government, Takaishi City Government, Hannan City Government, Neyagawa City Medical Association, Hirakata City Medical Association, and the Kadoma City Medical Association for their valuable support; Dr. Tomomi Marugame at the Statistics and Cancer Control Division, National Cancer Centre Research Institute for helpful discussions; and Ms. Tomoko Shibazaki, Ms. Nahoko Nishimura, and Ms. Naomi Takaoka for their assistance.

This study was supported by Grants-in-Aid (13770206, 16790351) for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology and Health and Labour Sciences Research Grants, Research on Allergic Disease and Immunology from the Ministry of Health, Labour, and Welfare, Japan.

References

1. Touger-Decker R. Clinical and laboratory assessment of nutrition status in dental practice. *Dent Clin North Am* 2003; 47: 259-78.
2. Joshupura KJ, Willett WC, Douglass CW. The impact of edentulousness on food and nutrient intake. *J Am Dent Assoc* 1996; 127: 459-67.
3. Krall E, Hayes C, Garcia R. How dentition status and masticatory function affect nutrient intake. *J Am Dent Assoc* 1996; 129: 1261-9.
4. Sheiham A, Steele JG, Marcenes W, Lowe C, Finch S, Bates CJ, Prentice A, Walls AW. The relationship among dental status, nutrient intake, and nutritional status in older people. *J Dent Res* 2001; 80: 408-13.
5. Shinkai RS, Hatch JP, Sakai S, Mobley CC, Saunders MJ, Rugh JD. Oral function and diet quality in a community-based sample. *J Dent Res* 2001; 80: 1625-30.
6. Lee JS, Weyant RJ, Corby P, Kritchevsky SB, Harris TB, Rooks R, Rubin SM, Newman AB. Edentulism and nutritional status in a biracial sample of well-functioning, community-dwelling elderly: the health, aging, and body composition study. *Am J Clin Nutr* 2004; 79: 295-302.
7. Hung HC, Willett W, Ascherio A, Rosner BA, Rimm E, Joshupura KJ. Tooth loss and dietary intake. *J Am Dent Assoc* 2003; 134: 1185-92.
8. Nowjack-Raymer RE, Sheiham A. Association of edentulism and diet and nutrition in US adults. *J Dent Res* 2003; 82: 123-6.
9. Sahyoun NR, Lin CL, Krall E. Nutritional status of the older adult is associated with dentition status. *J Am Diet Assoc* 2003; 103: 61-6.
10. Kwok T, Yu CN, Hui HW, Kwan M, Chan V. Association between functional dental state and dietary intake of Chinese vegetarian old age home residents. *Gerodontology* 2004; 21: 161-6.
11. Marshall TA, Warren JJ, Hand JS, Xie XJ, Stumbo PJ. Oral health, nutrient intake and dietary quality in the very old. *J Am Dent Assoc* 2002; 133: 1369-79.

12. MacKeown JM, Cleaton-Jones PE, Fatti P. Caries and micronutrient intake among urban South African children: a cohort study. *Community Dent Oral Epidemiol* 2003; 31: 213-20.
13. Larsson B, Johansson I, Ericson T. Prevalence of caries in adolescents in relation to diet. *Community Dent Oral Epidemiol* 1992; 20: 133-7.
14. Nishida M, Grossi SG, Dunford RG, Ho AW, Trevisan M, Genco RJ. Calcium and the risk for periodontal disease. *J Periodontol* 2000; 71: 1057-66.
15. Freeland JH, Cousins RJ, Schwartz R. Relationship of mineral status and intake to periodontal disease. *Am J Clin Nutr* 1976; 29: 745-9.
16. Krall EA, Wehler C, Garcia RI, Harris SS, Dawson-Hughes B. Calcium and vitamin D supplements reduce tooth loss in the elderly. *Am J Med* 2001; 111: 452-6.
17. Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, Hirota Y, Oda H, Osaka Maternal and Child Health Study Group. Soy, isoflavones, and prevalence of allergic rhinitis in Japanese women: the Osaka Maternal and Child Health Study. *J Allergy Clin Immunol* 2005; 115: 1176-83.
18. Tanaka K, Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, Hirota Y, Oda H, Osaka Maternal and Child Health Study Group. Active and passive smoking and tooth loss in Japanese women: baseline data from the Osaka Maternal and Child Health Study. *Ann Epidemiol* 2005; 15: 358-64.
19. Sasaki S, Yanagibori R, Amano K. Self-administered diet history questionnaire developed for health education: a relative validation of the test-version by comparison with 3-day diet record in women. *J Epidemiol* 1998; 8: 203-15.
20. Sasaki S, Ushio F, Amano K, Morihara M, Todoriki O, Uehara Y, Toyooka E. Serum biomarker-based validation of a self-administered diet history questionnaire for Japanese subjects. *J Nutr Sci Vitaminol (Tokyo)* 2000; 46: 285-96.
21. Willett W, Stampfer MJ. Total energy intake: implications for epidemiologic analyses. *Am J Epidemiol* 1986; 124: 17-27.
22. Song Y, Ridker PM, Manson JE, Cook NR, Buring JE, Liu S. Magnesium intake, C-reactive protein, and the prevalence of metabolic syndrome in middle-aged and older U.S. women. *Diabetes Care* 2005; 28: 1438-44.
23. Miyake Y, Sasaki S, Ohya Y, Miyamoto S, Matsunaga I, Yoshida T, Hirota Y, Oda H, The Osaka Maternal And Child Health Study Group. Dietary intake of seaweed and minerals and prevalence of allergic rhinitis in Japanese pregnant females: baseline data from the Osaka maternal and child health study. *Ann Epidemiol* 2006; 16: 614-21.
24. Kleber BM, Fehlinger R. Dental and periodontal disturbances due to magnesium deficit. *Magnes Res* 1989; 2: 235-7.
25. Prentice A. Diet, nutrition and the prevention of osteoporosis. *Public Health Nutr* 2004; 7: 227-43.
26. New SA, Robins SP, Campbell MK, Martin JC, Garton MJ, Bolton-Smith C, Grubb DA, Lee SJ, Reid DM. Dietary influences on bone mass and bone metabolism: further evidence of a positive link between fruit and vegetable consumption and bone health? *Am J Clin Nutr* 2000; 71: 142-51.
27. Krall EA, Dawson-Hughes B, Papas A, Garcia RI. Tooth loss and skeletal bone density in healthy postmenopausal women. *Osteoporos Int* 1994; 4: 104-9.
28. Douglass CW, Berlin J, Tennstedt S. The validity of self-reported oral health status in the elderly. *J Public Health Dent* 1991; 51: 220-2.
29. Axelsson G, Helgadottir S. Comparison of oral health data from self-administered questionnaire and clinical examination. *Community Dent Oral Epidemiol* 1995; 23: 365-8.
30. Pitiphat W, Garcia RI, Douglass CW, Joshupura KJ. Validation of self-reported oral health measures. *J Public Health Dent* 2002; 62: 122-8.
31. Statistics Bureau, Ministry of Public Management, Home Affairs, Post and Telecommunications, Japan. In: *2000 population census of Japan, Vol. 3-2-27, Labour Force Status of Population, Industry (Major Groups) of Employed Persons, and Education*. Osaka-fu. Tokyo: Japan Statistical Association, 2002: 436-40.
32. Dental Health Division of Health Policy Bureau Ministry of Health. In: *Labour and Welfare Japan. Report on the Survey of Dental Diseases (1999)*. Tokyo: Oral Health Association, 2001: 142-4.
33. Ohishi K, Kitagawa E, Morita M, Watanabe T, Matsuura T, Ito K. Reasons for extraction of permanent teeth in Okayama prefecture. A comparison with the survey in 1986-87. *J Dent Health* 2001; 50: 57-62.