

**Table 1** Nucleotide Sequences of Probes Used in RT-PCR Assays and Amplification Conditions


Gene	Source (reference)	Sequence	Product (bp)	Amplification conditions
TGF- $\beta$ R	Mouse (Machida <i>et al.</i> , 2000)	Sense 5'-ATCCATCACTAGATCGCCCT-3'	824	94°C for 30 s, 57°C for 30 s, 72°C for 30 s
	Human (GenBank Accession Number, L11695)	Antisense 5'-CGATGGATCAGAAGGTACAAGA-3' Sense 5'-GATGGGCTCTGCTTTGTCTC-3'	214	94°C for 30 s, 54°C for 30 s, 72°C for 30 s
TGF- $\beta$ R	Mouse (Machida <i>et al.</i> , 2000)	Antisense 5'-CAAGGCCAGGTGATGACTTT-3' Sense 5'-CGTGTGGAGGAAGAACAAACA-3'	560	94°C for 30 s, 57°C for 30 s, 72°C for 30 s
	Human (GenBank Accession Number, E10743)	Antisense 5'-TCTCAAACCTGCTCTGAGGGTG-3' Sense 5'-TTTCCACCTGTGACCAACCA-3'	185	94°C for 30 s, 54°C for 30 s, 72°C for 30 s
		Antisense 5'-GGAGAAAGCAGCATCTTCCAG-3'		

189 parisons. The differences between means were considered to be significant when  
 190  $P$  values were less than 0.05.

191

RESULTS

192 To obtain molecular evidence for the expression of TGF- $\beta$  receptor I and II in  
 193 MBEC4 cells and human brain pericytes, RT-PCR was carried out with a primer  
 194 pair specific to each type of TGF- $\beta$  receptor, from either mouse (for use on MBEC4  
 195 cells) or human (for use on human brain pericytes). As shown in Fig. 1A, RT-PCR  
 196 with mRNA obtained from either MBEC4 cells or brain pericytes yielded a single  
 197 product. The size of these products was as expected from the primer positions.




**Fig. 1** (A) Expression of TGF- $\beta$  receptor I and II mRNA in MBEC4 cells and human brain pericytes by RT-PCR analysis. (B) Time-course of the effect of CsA ( $5 \mu\text{M}$ ) on the Na-F permeability of MBEC4 cells in MBEC4 monolayers and pericyte co-cultures. Transport experiments were performed after 1, 3, 6, 9 and 12 h of exposure to CsA. Results are expressed as % increase of each corresponding vehicle treatment (MBEC4 monolayers;  $2.78 \pm 0.10 \times 10^{-4}$  to  $3.47 \pm 0.14 \times 10^{-4}$  cm/min, pericyte co-cultures;  $3.24 \pm 0.10 \times 10^{-4}$  to  $4.47 \pm 0.20 \times 10^{-4}$  cm/min). Values are the means  $\pm$  SEM ( $n = 7-16$ ). (C) Effect of treatment with CsA ( $5 \mu\text{M}$ ) for 6 h on the Na-F permeability of MBEC4 cells in MBEC4 monolayers and pericyte co-cultures. Results are expressed as % of vehicle-treated MBEC4 monolayers (% of vehicle-monolayer ( $3.26 \pm 0.13 \times 10^{-4}$  cm/min)). Values are the means  $\pm$  SEM ( $n = 7$ ). (D) Effect of treatment with CsA ( $5 \mu\text{M}$ ) for 6 h on the rhodamine 123 accumulation of MBEC4 cells in MBEC4 monolayers and pericyte co-cultures. Results are expressed as % of vehicle-treated MBEC4 monolayers (% of vehicle-monolayer ( $0.51 \pm 0.12$  nmol/mg protein)). Values are the means  $\pm$  SEM ( $n = 8$ ).

When MBEC4 cells were co-cultured with pericytes, the Na-F permeability of MBEC4 cells was increased from  $3.26 \pm 0.13 \times 10^{-4}$  cm/min in monolayers to  $3.86 \pm 0.23 \times 10^{-4}$  cm/min (Fig. 1C). The presence of pericytes decreased rhodamine 123 accumulation in MBEC4 cells from  $0.51 \pm 0.12$  nmol/mg protein in monolayers to  $0.43 \pm 0.10$  nmol/mg protein (Fig. 1D). A 12 h-exposure to CsA at 5 or 10  $\mu$ M showed no effect on cell viability as determined by mitochondrial dehydrogenase activities (WST-8 assay) of MBEC4 cells in MBEC4 monolayers ( $100.0 \pm 1.8$  and  $103.1 \pm 7.8\%$  of vehicle, respectively,  $n = 4$  inserts) and pericyte co-cultures ( $99.2 \pm 2.5$  and  $106.8 \pm 2.3\%$  of vehicle, respectively,  $n = 4$  inserts). Figure 1C and D shows the effect of CsA (5  $\mu$ M) on the permeability of MBEC4 cells to Na-F and the accumulation of rhodamine 123 in MBEC4 cells, respectively, in both MBEC4 monolayers and pericyte co-cultures. The Na-F permeability of MBEC4 cells was time-dependently increased during a period of 1–6 h after the addition of CsA (5  $\mu$ M), reaching a peak at 6 h (Fig. 1B). A significant difference in CsA-induced hyperpermeability was observed between MBEC4 monolayers and pericyte co-cultures, with the effect being most apparent after a 6 h-exposure [ $F(1, 24) = 10.51, P < 0.01$ ] (Fig. 1C). At the peak time, CsA increased the Na-F permeability of MBEC4 cells by  $10.3 \pm 3.4$  and  $31.9 \pm 3.6\%$  of each corresponding vehicle treatment in MBEC4 monolayers and pericyte co-cultures, respectively (Fig. 1C). Following exposure of cells to CsA for 9–12 h, permeability of the MBEC4 monolayers was gradually increased. However, the effect of CsA on pericyte co-cultures permeability became more moderate, reducing to the same level seen in MBEC4 monolayers (Fig. 1B). To clarify the role of brain pericytes in CsA-induced dysfunction of BBB, a 6 h-exposure of cells to CsA (5  $\mu$ M) was employed in the following experiment. The accumulation of rhodamine 123 in MBEC4 cells was increased by  $13.9 \pm 5.9$  and  $43.4 \pm 8.5\%$  of each corresponding vehicle treatment in MBEC4 monolayers and pericyte co-cultures, respectively, after a 6 h-exposure to CsA (Fig. 1D). A significant difference in CsA-induced decrease in P-gp activity was observed between MBEC4 monolayers and pericyte co-cultures [ $F(1, 28) = 4.65, P < 0.05$ ]. CsA-induced inhibition of P-gp function was more potent in pericyte co-cultures than in MBEC4 monolayers.

When pericyte co-cultures were treated with CsA (5  $\mu$ M) for 6 h, the levels of TGF- $\beta$ 1 mRNA in brain pericytes were significantly decreased to  $81.0 \pm 7.3\%$  of vehicle (Fig. 2A) [ $F(2, 13) = 5.05, P < 0.05$ ]; however, a 12 h exposure to CsA failed to decrease levels of TGF- $\beta$ 1 mRNA in brain pericytes ( $103.8 \pm 10.8\%$  of vehicle). In pericyte co-cultures, TGF- $\beta$ 1 (0.01–1 ng/mL) dose-dependently inhibited the elevation of Na-F permeability (Fig. 2B), and rhodamine 123 accumulation (Fig. 2C), in MBEC4 cells, induced by CsA ( $132.9 \pm 11.4$  to  $105.4 \pm 2.5\%$  and  $140.3 \pm 9.9$  to  $98.6 \pm 6.4\%$  of vehicle, respectively).

## DISCUSSION

In the present study, CsA (5  $\mu$ M) time-dependently increased the Na-F permeability of MBEC4 cells in pericyte co-cultures, this effect reaching a peak at 6 h



**Fig. 2** (A) Effect of CsA ( $5 \mu\text{M}$ ) on TGF- $\beta$ 1 mRNA expression in brain pericytes at 6 and 12 h after the addition of CsA in pericyte co-cultures. Total RNA of brain pericytes was extracted and subjected to real-time PCR analysis. Fold changes in TGF- $\beta$ 1 mRNA are normalized to GAPDH and compared with each corresponding vehicle treatment. Values are the means  $\pm$  SEM ( $n = 3-5$ ). \* $p < 0.05$ , significant difference from vehicle. (B) Effect of TGF- $\beta$ 1 on CsA-increased Na-F permeability of MBEC4 cells in pericyte co-cultures. Results are expressed as % of vehicle (vehicle;  $3.28 \pm 0.22 \times 10^{-4}$  cm/min). Values are the means  $\pm$  SEM ( $n = 8-16$ ). \* $p < 0.05$ , significant differences from vehicle. # $p < 0.05$ , significant difference from CsA treatment. (C) Effect of TGF- $\beta$ 1 on CsA-increased rhodamine 123 accumulation of MBEC4 cells in pericyte co-cultures. Results are expressed as % of each corresponding vehicle treatment (vehicle;  $1.16 \pm 0.24$  nmol/mg protein). Values are the means  $\pm$  SEM ( $n = 4-12$ ). \* $p < 0.05$ , significant differences from vehicle. # $p < 0.05$ , significant difference from CsA treatment.

240 after the addition of CsA (Fig. 1B and C). A 6 h exposure to CsA also decreased P-  
 241 gp function in MBEC4 cells in pericyte co-cultures, this effect being more apparent  
 242 than that in MBEC4 monolayers (Fig. 1D). These findings suggest that CsA-induced  
 243 hyperpermeability of, and P-gp dysfunction in, MBEC4 cells, was aggravated in pericyte  
 244 co-cultures. We previously reported that treatment with CsA ( $0.5-10 \mu\text{M}$ ) for  
 245 24 h dose-dependently decreased the viability of MBEC4 cells (Kochi *et al.*, 1999).  
 246 The present study demonstrated that a 12 h-exposure to CsA at concentrations up to  
 247  $10 \mu\text{M}$  had no effect on the viability of MBEC4 cells in MBEC4 monolayers and  
 248 pericyte co-cultures. Thus, the exposure time (6 h) and the submaximum concentra-  
 249 tion ( $5 \mu\text{M}$ ) of CsA without cytotoxicity were selected here.

250 The significance of brain pericytes in the regulation of the BBB was suggested  
 251 by our previous study using a primary culture of rat brain pericytes. We reported

that brain pericyte-derived TGF- $\beta$ 1 contributed to the induction and up-regulation of BBB function (Dohgu *et al.*, 2004b, 2005). In the present study, the presence of human brain pericytes decreased the function of tight junctions and increased the function of P-gp, in MBEC4 cells, by about 20%. An interjunctional property of MBEC4 cells was lowered by co-culturing with human brain pericytes, this event being inconsistent with our previous report using a primary culture of rat brain pericytes. The different backgrounds of human brain pericytes commercially supplied and species-different cell type may be compounding factors in this discrepancy (Lai and Kuo, 2005); however, further studies are required to confirm this. Here, we employed a convenient *in vitro* model with MBEC4 cells and human brain pericytes.

To test whether TGF- $\beta$ 1 production of the brain pericytes participates in the mediation of CsA-induced dysfunction of the BBB, we examined the effect of CsA on the expression of TGF- $\beta$ 1 in brain pericytes. A 6 h exposure to CsA (5  $\mu$ M) significantly decreased the levels of TGF- $\beta$ 1 mRNA in brain pericytes in pericyte co-cultures (Fig. 2A). Treatment with TGF- $\beta$ 1 dose-dependently inhibited CsA-induced hyperpermeability and P-gp dysfunction in MBEC4 cells in pericyte co-cultures (Fig. 2B and C). These findings suggest that CsA decreases BBB function by inhibiting TGF- $\beta$ 1 production in the brain pericytes. In pericyte co-cultures, an aggravation of CsA-induced hyperpermeability of MBEC4 cells occurred with a peak at 6 h after the addition of CsA, becoming more moderate at 9–12 h (Fig. 1B). In parallel with these events, TGF- $\beta$ 1 mRNA levels were significantly decreased by a 6 h exposure to CsA but not by a 12 h exposure (Fig. 2A), suggesting that CsA-induced hyperpermeability is ameliorated by the compensatory secretion of TGF- $\beta$ 1 from brain pericytes during the later period of CsA exposure. These data further support a critical role of pericyte-derived TGF- $\beta$ 1 in mediating CsA-induced BBB dysfunction.

RT-PCR analysis demonstrated the expression of TGF- $\beta$  receptor I and II in MBEC4 cells and brain pericytes (Fig. 1A). TGF- $\beta$ 1 mRNA was also detected in brain pericytes (Fig. 2A) and MBEC4 cells (data not shown). Considering these observations, TGF- $\beta$  is likely to participate in the up-regulation of BBB function through an autocrine and/or paracrine pathway in brain endothelial cells and pericytes. Autoinduction of TGF- $\beta$ 1 is mediated by binding of the transcription factor activator protein-1 (AP-1) complex to homologous elements in two regions of the TGF- $\beta$ 1 promoter (Kim *et al.*, 1990). In this positive autoregulation process, CsA is known to directly inhibit activation of the JunD isoforms in the AP-1 complex responsible for TGF- $\beta$  signaling in lung fibroblasts (Eickelberg *et al.*, 2001). Calcineurin, a molecular target of CsA, has been shown to be involved in AP-1 activation in immune cells (Pfeuffer *et al.*, 1994; Tsuboi *et al.*, 1994). Based on this evidence, CsA may be interpreted as lowering TGF- $\beta$ 1 expression in brain pericytes due, at least in part, to an inhibition of the AP-1 activation step during TGF- $\beta$ 1 autoinduction. We previously reported that CsA increased NO production in MBEC4 cells (Dohgu *et al.*, 2004a) and elevated levels of NO have been known to reduce TGF- $\beta$ 1 production in the heart (Smith *et al.*, 2005). Therefore, CsA may also act on MBEC4 cells to lower TGF- $\beta$ 1 expression in brain pericytes.

297 In conclusion, CsA-induced hyperpermeability and P-gp dysfunction of  
298 MBEC4 cells were markedly aggravated in co-cultures with brain pericytes. This  
299 aggravation appears to occur due to CsA-induced inhibition of TGF- $\beta$ 1 expression  
300 in brain pericytes. These findings suggest that an inhibition of brain pericyte-derived  
301 TGF- $\beta$ 1 contributes to the occurrence of CsA-induced dysfunction of the BBB,  
302 thereby triggering neurotoxicity.

303

#### ACKNOWLEDGMENTS

304 This work was supported, in part, by Grants-in-Aid for Scientific Research  
305 ((B) 17390159) from JSPS, Japan and by a Grant-in-Aid for Exploratory Research  
306 (17659160) from MEXT, Japan. The authors thank Dr. Mária A. Deli (Institute of  
307 Biophysics, Biological Research Centre of the Hungarian Academy of Sciences) for  
308 pertinent comments on the manuscript.

309

#### REFERENCES

- 310 Antonelli-Orlidge, A., Saunders, K. B., Smith, S. R., and D'Amore, P. A. (1989). An activated form of  
311 transforming growth factor beta is produced by co-cultures of endothelial cells and pericytes. *Proc.*  
312 *Natl. Acad. Sci. U.S.A.* **86**:4544–4548.
- 313 Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of  
314 protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- 315 Dehouck, M.P., Jolliet-Riant, P., Bree, F., Fruchart, J. C., Cecchelli, R., and Tillement, J. P. (1992). Drug  
316 transfer across the blood-brain barrier: Correlation between *in vitro* and *in vivo* models. *J. Neu-*  
317 *rochem.* **58**:1790–1797.
- 318 Dohgu, S., Yamauchi, A., Nakagawa, S., Takata, F., Kai, M., Egawa, T., Naito, M., Tsuruo, T., Sawada,  
319 Y., Niwa, M., and Kataoka, Y. (2004a). Nitric oxide mediates cyclosporine-induced impairment of  
320 the blood-brain barrier in co-cultures of mouse brain endothelial cells and rat astrocytes. *Eur. J.*  
321 *Pharmacol.* **505**:51–59.
- 322 Dohgu, S., Yamauchi, A., Takata, F., Naito, M., Tsuruo, T., Higuchi, S., Sawada, Y., and Kataoka, Y.  
323 (2004b). Transforming growth factor-beta1 upregulates the tight junction and P-glycoprotein of  
324 brain microvascular endothelial cells. *Cell. Mol. Neurobiol.* **24**:491–497.
- 325 Dohgu, S., Takata, F., Yamauchi, A., Nakagawa, S., Egawa, T., Naito, M., Tsuruo, T., Sawada, Y., Niwa,  
326 M., and Kataoka, Y. (2005). Brain pericytes contribute to the induction and up-regulation of blood-  
327 brain barrier functions through transforming growth factor-beta production. *Brain Res.* **1038**:208-  
328 215.
- 329 Eickelberg, O., Pansky, A., Koehler, E., Bihl, M., Tamm, M., Hildebrand, P., Perruchoud, A. P., Kash-  
330 garian, M., and Roth, M. (2001). Molecular mechanisms of TGF-(beta) antagonism by interferon  
331 (gamma) and cyclosporine A in lung fibroblasts. *FASEB J.* **15**:797–806.
- 332 Fontaine, M., Elmquist, W. F., and Miller, D. W. (1996). Use of rhodamine 123 to examine the functional  
333 activity of P-glycoprotein in primary cultured brain microvessel endothelial cell monolayers. *Life*  
334 *Sci.* **59**:1521–1531.
- 335 Gijtenbeek, J. M., Van Den Bent, M. J., and Vecht, C. J. (1999). Cyclosporine neurotoxicity. *J. Neurol.*  
336 **246**:339–346.
- 337 Hori, S., Ohtsuki, S., Hosoya, K., Nakashima, E., and Terasaki, T. (2004). A pericyte-derived  
338 angiopoietin-1 multimeric complex induces occludin gene expression in brain capillary endothelial  
339 cells through Tie-2 activation *in vitro*. *J. Neurochem.* **89**:503–513.
- 340 Kahan, B. D. (1989). Cyclosporine. *N. Engl. J. Med.* **321**:1725–1738.
- 341 Kim, S. J., Angel, P., Lafyatis, R., Hattori, K., Kim, K. Y., Sporn, M. B., Karin, M., and Roberts, A. B.  
342 (1990). Autoinduction of transforming growth factor beta 1 is mediated by the AP-1 complex. *Mol.*  
343 *Cell. Biol.* **10**:1492–1497.

- Kochi, S., Takanaga, H., Matsuo, H., Naito, M., Tsuruo, T., and Sawada, Y. (1999). Effect of cyclosporin A or tacrolimus on the function of blood-brain barrier cells. *Eur. J. Pharmacol.* **372**:287–295. 344
- Kochi, S., Takanaga, H., Matsuo, H., Ohtani, H., Naito, M., Tsuruo, T., and Sawada, Y. (2000). Induction of apoptosis in mouse brain capillary endothelial cells by cyclosporin A and tacrolimus. *Life Sci.* **66**:2255–2260. 345
- Lai, C. H., and Kuo, K. H. (2005). The critical component to establish *in vitro* BBB model: Pericyte. *Brain Res. Brain Res. Rev.* **50**:258–265. 346
- Machida, H., Ogawa, K., Funaba, M., Mizutani, T., and Tsujimoto, M. (2000). mRNA expression of type I and type II receptors for activin, transforming growth factor-beta, and bone morphogenetic protein in the murine erythroleukemic cell line, F5-5.fl. *Eur. J. Endocrinol.* **143**:705–710. 347
- Pardridge, W. M. (1999). Blood-brain barrier biology and methodology. *J. Neurovirol.* **5**:556–569. 348
- Pirsch, J. D., Miller, J., Deierhoi, M. H., Vincenti, F., and Filo, R. S. (1997). A comparison of tacrolimus (FK506) and cyclosporine for immunosuppression after cadaveric renal transplantation. FK506 Kidney Transplant Study Group. *Transplantation* **63**:977–983. 349
- Pfeuffer, I., Klein-Hessling, S., Heinfling, A., Chuvpilo, S., Escher, C., Brabletz, T., Hentsch, B., Schwarzenbach, H., Matthias, P., and Serfling, E. (1994). Octamer factors exert a dual effect on the IL-2 and IL-4 promoters. *J. Immunol.* **153**:5572–5585. 350
- Ramsauer, M., Krause, D., and Dermietzel, R. (2002). Angiogenesis of the blood-brain barrier *in vitro* and the function of cerebral pericytes. *FASEB J.* **16**:1274–1276. 351
- Schinkel, A. H. (1999). P-Glycoprotein, a gatekeeper in the blood-brain barrier. *Adv. Drug Deliv. Rev.* **36**:179–194. 352
- Shuto, H., Kataoka, Y., Kanaya, A., Matsunaga, K., Sueyasu, M., and Oishi, R. (1998). Enhancement of serotonergic neural activity contributes to cyclosporine-induced tremors in mice. *Eur. J. Pharmacol.* **341**:33–37. 353
- Shuto, H., Kataoka, Y., Fujisaki, K., Nakao, T., Sueyasu, M., Miura, I., Watanabe, Y., Fujiwara, M., and Oishi, R. (1999). Inhibition of GABA system involved in cyclosporine-induced convulsions. *Life Sci.* **65**:879–887. 354
- Smith, R. S., Jr., Agata, J., Xia, C. F., Chao, L., and Chao, J. (2005). Human endothelial nitric oxide synthase gene delivery protects against cardiac remodeling and reduces oxidative stress after myocardial infarction. *Life Sci.* **76**:2457–2471. 355
- Tatsuta, T., Naito, M., Oh-hara, T., Sugawara, I., and Tsuruo, T. (1992). Functional involvement of P-glycoprotein in blood-brain barrier. *J. Biol. Chem.* **267**:20383–20391. 356
- The U.S. Multicenter FK506 Liver Study Group. (1994). A comparison of tacrolimus (FK 506) and cyclosporine for immunosuppression in liver transplantation. *N. Engl. J. Med.* **331**:1110–1115. 357
- Thomas, W. E. (1999). Brain macrophages: On the role of pericytes and perivascular cells. *Brain Res. Brain Res. Rev.* **31**:42–57. 358
- Tsuboi, A., Muramatsu, M., Tsutsumi, A., Arai, K., and Arai, N. (1994). Calcineurin activates transcription from the GM-CSF promoter in synergy with either protein kinase C or NF-kappa B/AP-1 in T cells. *Biochem. Biophys. Res. Commun.* **199**:1064–1072. 359
- Untergasser, G., Gander, R., Lilg, C., Lepperdinger, G., Plas, E., and Berger, P. (2005). Profiling molecular targets of TGF-b1 in prostate fibroblast-to-myofibroblast transdifferentiation. *Mech. Ageing Dev.* **126**:59–69. 360

---

Rapid Communication

---

**An Inhibitory Role of Nitric Oxide in the Dynamic Regulation of the Blood-Brain Barrier Function**

Atsushi Yamauchi,<sup>1</sup> Shinya Dohgu,<sup>1</sup> Tsuyoshi Nishioku,<sup>1</sup> Hideki Shuto,<sup>1</sup>  
Mikihiko Naito,<sup>2</sup> Takashi Tsuruo,<sup>2</sup> Yasufumi Sawada,<sup>3</sup> and Yasufumi Kataoka<sup>1,4</sup>

Received June 2, 2006; accepted June 19, 2006

---

**SUMMARY**

1. The present study aimed at elucidating the effects of nitric oxide (NO) on blood-brain barrier (BBB) function with mouse brain capillary endothelial (MBEC4) cells.

2. Histamine (20–100  $\mu$ M) evoked NO production (1.6–7  $\mu$ M) in MBEC4 cells in a dose-dependent manner.

3. The permeability coefficient of sodium fluorescein for MBEC4 cells and the cellular accumulation of rhodamine 123 in MBEC4 cells were increased dose-dependently by addition of NO solutions (14 and 28  $\mu$ M) every 10 min during a 30-min period.

4. The present study demonstrated that NO increased the permeability and inhibited the P-glycoprotein efflux pump of brain capillary endothelial cells, suggesting that NO plays an inhibitory role in the dynamic regulation of the BBB function.

---

**KEY WORDS:** nitric oxide; blood-brain barrier (BBB); permeability; P-glycoprotein; mouse brain endothelial cells.

**INTRODUCTION**

The blood-brain barrier (BBB) contributes to brain homeostasis and fulfills a protective function by regulating the access of solutes and toxic substances to the central nervous system. The BBB is formed by brain capillary endothelial cells, which are closely sealed by tight junctions (Pardridge, 1999). The tight junctions in the

---

<sup>1</sup>Department of Pharmaceutical Care and Health Sciences, Faculty of Pharmaceutical Sciences, Fukuoka University, 8-19-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan.

<sup>2</sup>Institute of Molecular and Cellular Biosciences, University of Tokyo, Bunkyo-ku, Tokyo 113-0032, Japan.

<sup>3</sup>Laboratory of Drug Informatics, Graduate School of Pharmaceutical Sciences, The University of Tokyo, 3-14-15 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan.

<sup>4</sup>To whom correspondence should be addressed at Department of Pharmaceutical Care and Health Sciences, Faculty of Pharmaceutical Sciences, Fukuoka University, 8-19-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan; e-mail: ykataoka@fukuoka-u.ac.jp.



BBB prevent significant passive movement of small hydrophilic molecules from the blood to the brain, but specialized transport systems mediate the entry of essential substances such as glucose, amino acids, choline, monocarboxylic acids, amines, thyroid hormones, purine bases, and nucleosides (Tsuji and Tamai, 1999; Kusunaga and Sugiyama, 2001). The efflux transporter P-glycoprotein (P-gp) is a key element of the molecular machinery that confers special permeability properties on the BBB. P-gp, which was initially recognized for its ability to excrete anticancer drugs from multidrug-resistant cancer cells, is strongly expressed in brain capillaries. Its expression in the BBB limits the accumulation of many hydrophobic molecules and potentially toxic substances in the brain.

Nitric oxide (NO) is a transient product of inflammatory processes, generated from L-arginine by the enzyme NO synthase (NOS). NO appears to be involved in numerous vital cellular functions including neurotransmission, blood-pressure control, and the regulation of vascular tone. The basal production of NO appears to be required for biological regulation, and yet an excess of this same molecule can be cytotoxic to organism. But the molecular mechanisms mediating NO-induced tissue injury and breakdown of the BBB are not completely understood.

In the present study, to clarify the role of NO in the dynamic regulation of the BBB, we examined effects of NO on the function of tight junctions and P-gp in mouse brain capillary endothelial (MBEC4) cells. MBEC4 cells show the highly specialized characteristics of brain microvascular endothelial cells including P-gp expression (Tatsuta *et al.*, 1992, 1994).

## MATERIALS AND METHODS

A saturated NO solution (typically containing approximately 1.4 mM NO) was prepared according to Ikesue *et al.* (2000). Deionized water (2 mL) was bubbled with argon for 20 min to remove oxygen. Then, the solution was bubbled with pure NO gas for 20 min and kept in a glass flask with a rubber septum under a NO atmosphere prior to use.

MBEC4 cells, which were isolated from BALB/c mouse brain cortices and immortalized by SV40-transformation (Tatsuta *et al.*, 1992), were cultured in Dulbecco's modified Eagle's medium (DMEM; GIBCO BRL, Life Technologies, Grand Island, NY) supplemented with 10% fetal bovine serum, 100 units/mL of penicillin, and 100  $\mu$ g/mL of streptomycin. They were grown in 2.5-cm<sup>2</sup> dish, 12-well Transwells (Costar, MA) and 24-well plates in a humidified atmosphere of 5% CO<sub>2</sub>/95% air at 37°C.

Direct and continuous electrochemical measurement of NO was performed with a three-electrode potentiostatic EMS-100 system (BIO-LOGIC, Grenoble, France), as previously described (Ikesue *et al.*, 2000; Trevin *et al.*, 1998). In brief, confluent MBEC4 cells in a 2.5-cm<sup>2</sup> dish were washed three times with Krebs-Ringer solution (143.0 mM NaCl, 4.7 mM KCl, 2.5 mM CaCl<sub>2</sub>, 1.0 mM NaH<sub>2</sub>PO<sub>4</sub>, and 11.0 mM D-glucose, pH 7.4). The dish was placed on the stage of an inverted microscope (ECLIPSE TE300, Nikon, Tokyo, Japan) mounted with an NO monitoring

70 system. The NO-biosensor (ASTEC, Fukuoka, Japan) was positioned about 10  $\mu\text{m}$   
71 above the cell surface. Ten minutes after treatment with 1-mM L-arginine (Sigma,  
72 St. Louis, MO), histamine (Wako, Osaka, Japan) in a volume of 10  $\mu\text{L}$  was added to  
73 the cells in 1 mL of Krebs–Ringer solution with a transient mixing step to give the  
74 final concentration indicated. The level of production of NO in MBEC4 cells was  
75 monitored for a 15-min period after the addition of histamine.

76 MBEC4 cells (42,000 cells/cm<sup>2</sup>) were cultured on the collagen-coated polycar-  
77 bonate membrane of the Transwell insert (3.0- $\mu\text{m}$  pore size, 12-well type). After  
78 3 days, they were washed three times with serum-free medium. Cells were exposed  
79 to 5 or 10  $\mu\text{L}$  of NO solution (final concentration, 14 and 28  $\mu\text{M}$ , respectively) in-  
80 jected into the inside of the insert (luminal side) every 10 min during a 30-min pe-  
81 riod. To initiate the transport experiments, the medium was removed, and cells were  
82 washed three times with Krebs–Ringer buffer (118 mM NaCl, 4.7 mM KCl, 1.3 mM  
83 CaCl<sub>2</sub>, 1.2 mM MgCl<sub>2</sub>, 1.0 mM NaH<sub>2</sub>PO<sub>4</sub>, 25 mM NaHCO<sub>3</sub>, and 11 mM D-glucose,  
84 pH 7.4). Krebs–Ringer buffer (1.5 mL) was added to the outside of the insert (ablu-  
85 minal side). Krebs–Ringer buffer (0.5 mL) containing 100  $\mu\text{g}/\text{mL}$  of sodium fluores-  
86 cein (Na-F; Sigma) was loaded on the luminal side of the insert. Samples (0.5 mL)  
87 were removed from the abluminal chamber at 10, 20, 30, and 60 min and were imme-  
88 diately replaced with fresh Krebs–Ringer buffer. Aliquots (5  $\mu\text{L}$ ) from the ablumi-  
89 nal chamber samples were mixed with 200  $\mu\text{L}$  of Krebs–Ringer buffer, and then  
90 the concentration of Na-F was determined using a multiwell fluorometer ( $E_x(\lambda)$   
91 485 nm;  $E_m(\lambda)$  530 nm; CytoFluor Series 4000, PerSeptive Biosystems, Framing-  
92 ham, MA). The permeability coefficient and clearance were calculated according  
93 to the method described by Dehouck *et al.* (1992). Clearance was expressed as mi-  
94 croliters of tracer diffusing from the luminal to abluminal chamber and was cal-  
95 culated from the initial concentration of tracer in the luminal chamber and final  
96 concentration in the abluminal chamber: Clearance ( $\mu\text{L}$ ) =  $[C]_A \times V_A/[C]_L$  where  
97  $[C]_L$  is the initial luminal tracer concentration,  $[C]_A$  is the abluminal tracer concen-  
98 tration, and  $V_A$  is the volume of the abluminal chamber. During a 60-min period  
99 of the experiment, the clearance volume increased linearly with time. The aver-  
100 age volume cleared was plotted versus time, and the slope was estimated by linear  
101 regression analysis. The slope of clearance curves for the MBEC4 monolayer was  
102 denoted by  $\text{PS}_{\text{app}}$ , where PS is the permeability-surface area product (in microliters  
103 per minute). The slope of the clearance curve with a control membrane was denoted  
104 by  $\text{PS}_{\text{membrane}}$ . The real PS value for the MBEC4 monolayer ( $\text{PS}_{\text{trans}}$ ) was calculated  
105 from  $1/\text{PS}_{\text{app}} = 1/\text{PS}_{\text{membrane}} + 1/\text{PS}_{\text{trans}}$ . The  $\text{PS}_{\text{trans}}$  values were divided by the sur-  
106 face area of the Transwell inserts to generate the permeability coefficient ( $P_{\text{trans}}$ , in  
107 centimeters per minute).

108 The functional activity of P-gp was determined by measuring the cellular ac-  
109 cumulation of rhodamine 123 (Sigma), according to the method of Fontaine *et al.*  
110 (1996). MBEC4 cells (21,000 cells/cm<sup>2</sup>) were cultured on collagen-coated 24-well  
111 plates. Three days after seeding, they were washed three times with serum-free  
112 medium and then exposed to 14–28  $\mu\text{M}$  of NO solution every 10 min during a 30-  
113 min period. The medium was removed, and the cells were washed three times with  
114 assay buffer (143 mM NaCl, 4.7 mM KCl, 1.3 mM CaCl<sub>2</sub>, 1.2 mM MgCl<sub>2</sub>, 1.0 mM  
115 NaH<sub>2</sub>PO<sub>4</sub>, 10 mM HEPES, and 11 mM D-glucose, pH 7.4). The cells were incubated

with 0.5 mL of assay buffer containing 5  $\mu$ M of rhodamine 123 for 60 min. Then, the solution was removed, and the cells were washed three times with ice-cold phosphate-buffered saline and solubilized in 1 M NaOH (0.2 mL). The solution was neutralized with 1-M HCl (0.2 mL), and the rhodamine 123 content was determined using a multiwell fluorometer ( $E_x(\lambda)$  485 nm;  $E_m(\lambda)$  530 nm). The cellular protein was measured by the method of Bradford (Bradford, 1976).

The effect of the NO solution on cell viability was assessed using a WST-8 assay (Cell Counting Kit, DOJINDO, Kumamoto, Japan). A highly water-soluble formazan dye, reduced by mitochondrial dehydrogenase, was measured by determining the absorbance of each sample with a 450-nm test wavelength and a 700-nm reference wavelength.

The values are expressed as means  $\pm$  SEM. Statistical analysis was performed using Student's *t*-test. One-way analysis of variance (ANOVA) followed by the Dunnett test was applied to multiple comparisons. The differences between means were considered to be significant when *P* values were less than 0.05.

## RESULTS

Figure 1A shows a representative current-time curve obtained with the NO biosensor in MBEC4 cells. When L-arginine (1 mM) was added to cells 10 min before the addition of histamine, the signal became stable within 1–2 min. The NO-biosensor signal increased rapidly to reach a peak within 1 min after the injection of histamine. Following this period, the signal decreased slowly to the baseline about 5–7 min post-injection. Histamine at concentrations of 20–100  $\mu$ M dose-dependently increased NO production ( $1.55 \pm 0.65$ – $6.94 \pm 1.43$   $\mu$ M) in MBEC4 cells (Fig. 1B).

To evaluate effect of NO on the BBB function, the concentration and the exposure time of NO were determined on the basis of NO production evoked by histamine. When the NO solution at the final concentrations of 14 and 28  $\mu$ M was added every 10 min during a 30-min period, the permeability coefficients of Na-F for MBEC4 cells increased dose-dependently to  $113.6 \pm 12.9$  and  $123.7 \pm 2.8\%$ , respectively (Fig. 2A). The exposure to NO solution had no effect on cell viability assessed with the WST-8 assay (14  $\mu$ M:  $93.7 \pm 3.3\%$ , 28  $\mu$ M:  $101.3 \pm 3.2\%$  of control).

As shown in Fig. 2B, the accumulation of rhodamine 123 in MBEC4 cells increased dose-dependently to  $108.3 \pm 4.8$  and  $164.5 \pm 14.6\%$  of the control value after a 30-min exposure to the NO solution at concentrations of 14 and 28  $\mu$ M, respectively.

## DISCUSSION

As shown in Fig. 1, direct and continuous electrochemical monitoring revealed that histamine stimulated MBEC4 cells to release NO over a short period (about 5 min). This phenomenon appears to be due to an activation of the constitutive endothelial form of NO synthase (eNOS) through a  $Ca^{2+}$ /calmodulin-dependent pathway in response to  $H_1$  receptor-mediated increases in intracellular  $Ca^{2+}$  (Daum




Fig. 1. Histamine-induced NO production in MBEC cells. (A) A representative differential pulse amperogram obtained using a NO biosensor shows the level of NO production evoked by histamine (100  $\mu\text{M}$ ). (B) Concentration-response relationship of histamine-induced NO production using direct electrochemical monitoring in MBEC cells. Values are means  $\pm$  SEM ( $n = 3$ ). \* $p < 0.05$ , significant difference from 20  $\mu\text{M}$  histamine.

158 *et al.*, 1983; Kishi *et al.*, 1996). Various endogenous substances including histamine  
 159 stimulate NO production in the microvascular endothelium and/or glial and neu-  
 160 ronal cells in the brain (Mayhan, 1996), probably contributing to dynamic regulation  
 161 of the BBB function.

162 The permeability of MBEC4 cells to Na-F was apparently increased by a brief  
 163 exposure to NO solutions at 2- to 4-fold the concentrations induced by histamine  
 164 (Fig. 2A). The NO solutions employed here are more prominent than convenient  
 165 NO donors such as sodium nitroprusside with respect to biological properties in-  
 166 cluding distribution and degradation. The cell viability was not influenced by NO  
 167 at the concentrations employed here (14–28  $\mu\text{M}$ ), suggesting that the increased  
 168 paracellular permeability of MBEC4 cells was not due to the cytotoxicity of NO.  
 169 Therefore, NO is highly likely to lower the functions of the tight junctions at the




Fig. 2. Changes in the permeability coefficient of Na-F (A) and the cellular accumulation of rhodamine 123 (B) in MBEC4 cell monolayers after addition of NO solutions every 10 min during a 30-min period. Data are expressed as a percentage of the corresponding control value (A:  $1.47 \pm 0.21 \times 10^{-4}$  cm/min, B:  $0.95 \pm 0.07$  nmol/mg protein). Values are shown as means  $\pm$  SEM ( $n = 3-18$ ). \*\* $P < 0.01$ , significant difference from control.

BBB, supporting early reports (Hurst and Fritz, 1996; Mayhan, 2000; Shukula *et al.*, 170  
 1996). The mechanisms by which NO donors increased vascular endothelial perme- 171  
 ability involved an increase in the level of cyclic guanosine monophosphate (cGMP) 172  
 A1 (Gimeno *et al.*, 1998) or the formation of peroxynitrite (Menconi *et al.*, 1998). These 173  
 substances conceivably influence the intrinsic tight junction proteins and the associ- 174  
 ated actin cytoskeleton through a direct or second signaling pathway. Further stud- 175  
 ies are required to clarify this mechanism. 176

In the present study, the most important finding was that the accumulation of 177  
 rhodamine 123, a substrate of P-gp (Fontaine *et al.*, 1996), in MBEC4 cells was 178  
 increased by treatment with NO solutions (Fig. 2B). P-gp is an energy-dependent 179  
 efflux pump mediating the transportation of substances from the basement mem- 180  
 brane to the luminal surface. NO induced a depletion of endothelial ATP by 181

182 inhibiting glyceraldehyde-3-phosphate dehydrogenase (glycolytic enzyme) activity  
 183 (Hurst *et al.*, 2001). This inhibition of energy metabolism is probably associated with  
 184 the NO-triggered inhibition of P-gp function.

185 In the present study, NO induced hyperpermeability in MBEC4 cells and inhi-  
 186 bition of the P-gp efflux pump. These findings suggest that NO in the brain plays an  
 187 inhibitory role in the dynamic regulation of the BBB function.

## 188 ACKNOWLEDGMENTS

189 This work was supported in part by Grants-in-Aid for Scientific Research  
 190 ((B)(2) 14370789) and ((C)(2) 15590475) from JSPS, Japan, by a Grant-in-Aid for  
 191 Exploratory Research (16659138) from MEXT, Japan, and by funds (No.: 031001)  
 from the Central Research Institute of Fukuoka University.

## 193 REFERENCES

- 194 Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of  
 195 protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**:248–254.
- 196 Daum, P. R., Downes, C. P., and Young, J. M. (1983). Histamine-induced inositol phospholipid break-  
 197 down mirrors H1-receptor density in brain. *Eur. J. Pharmacol.* **87**:497–498.
- 198 Dehouck, M.-P., Joliet-Riant, P., Brée, F., Fruchart J.-C., Cecchelli, R., and Tillement, J.-P. (1992). Drug  
 199 transfer across the blood-brain barrier: Correlation between in vitro and in vivo models. *J. Neu-  
 200 rochem.* **58**:1790–1797.
- 201 Fontaine, M., Elmquist, W. F., and Miller, D. W. (1996). Use of rhodamine 123 to examine the functional  
 202 activity of P-glycoprotein in primary cultured brain microvessel endothelial cell monolayers. *Life  
 203 Sci.* **59**:1521–1531.
- 204 Gimeno, G., Carpentier, P. H., Desquand-Billiald, S., Hanf, R., and Finet, M. (1998). L-arginine and NG-  
 205 nitro-L-arginine methyl ester cause macromolecular extravasation in the microcirculation of awake  
 206 hamsters. *Eur. J. Pharmacol.* **346**:275–282.
- 207 Hurst, R. D., and Fritz, I. B. (1996). Properties of an immortalized vascular endothelial/glioma cell co-  
 208 culture model of blood-brain barrier. *J. Cell. Physiol.* **167**:81–88.
- 209 Hurst, R. D., Azam, S., Hurst, A., and Clark, J. B. (2001). Nitric-oxide-induced inhibition of  
 210 glyceraldehyde-3-phosphate dehydrogenase may mediate reduced endothelial cell monolayer in-  
 211 tegrity in an in vitro model blood-brain barrier. *Brain Res.* **894**:181–188.
- 212 Ikesue, H., Kataoka, Y., Kawachi, R., Dohgu, S., Shuto, H., and Oishi, R. (2000). Cyclosporine enhances  
 213  $\alpha_1$ -adrenoceptor-mediated nitric oxide production in C6 glioma cells. *Eur. J. Pharmacol.* **407**:221–  
 214 226.
- 215 Kishi, F., Nakaya, Y., Takahashi, A., Miyoshi, H., Nomura, M., and Saito, K. (1996). Intracellular and  
 216 extracellular  $Ca^{2+}$  regulate histamine-induced release of nitric oxide in vascular endothelial cells as  
 217 shown with sensitive and selective nitric oxide electrodes. *Pharmacol. Res.* **33**:123–126.
- 218 Kusuvara, H., and Sugiyama, Y. (2001). Efflux transport systems for drugs at the blood-brain barrier and  
 219 blood-cerebrospinal fluid barrier (Part 1). *Drug. Discov. Today* **6**:150–156.
- 220 Mayhan, W. G. (1996). Role of nitric oxide in histamine-induced increases in permeability of the blood-  
 221 brain barrier. *Brain Res.* **743**:70–76.
- 222 Mayhan, W. G. (2000). Nitric oxide donor-induced increase in permeability of the blood-brain barrier.  
 223 *Brain Res.* **866**:101–108.
- 224 Menconi, M. J., Unno, N., Smith, M., Aguirre, D. E., and Fink, M. P. (1998). Nitric oxide donor-induced  
 225 hyperpermeability of cultured intestinal epithelial monolayers: Role of superoxide radical, hydroxyl  
 226 radical, and peroxynitrite. *Biochim. Biophys. Acta* **1425**:189–203.
- 227 Pardridge, W. M. (1999). Blood-brain barrier biology and methodology. *J. Neurovirol.* **5**:556–569.
- 228 Shukula, A., Dikshit, M., and Srimal, R. C. (1996). Nitric oxide-dependent blood-brain barrier perme-  
 229 ability alteration in the rat brain. *Experientia* **52**:136–140.

- Tatsuta, T., Naito, M., Oh-hara, T., Sugawara, I., and Tsuruo, T. (1992). Functional involvement of P-glycoprotein in blood-brain barrier. *J. Biol. Chem.* **267**:20383–20391. 230  
231
- Tatsuta, T., Naito, M., Mikami, K., and Tsuruo, T. (1994). Enhanced expression by the brain matrix of P-glycoprotein in brain capillary endothelial cells. *Cell Growth Differ.* **5**:1145–1152. 232  
233
- Trevin, S., Kataoka, Y., Kawachi, R., Shuto, H., Kumakura, K., and Oishi, R. (1998). Direct and continuous electrochemical measurement of noradrenaline-induced nitric oxide production in C6 glioma cells. *Cell. Mol. Neurobiol.* **18**:453–458. 234  
235  
236
- Tsuji, A., and Tamai, I. (1999). Carrier-mediated or specialized transport of drugs across the blood-brain barrier. *Adv. Drug. Deliv. Rev.* **36**:277–290. 237  
238

Short Communication

## Protective Action of Indapamide, a Thiazide-Like Diuretic, on Ischemia-Induced Injury and Barrier Dysfunction in Mouse Brain Microvascular Endothelial Cells

Tsuyoshi Nishioku<sup>1</sup>, Fuyuko Takata<sup>1</sup>, Atsushi Yamauchi<sup>1</sup>, Noriko Sumi<sup>1</sup>, Ikumi Yamamoto<sup>1</sup>, Akiko Fujino<sup>1</sup>, Mikihiro Naito<sup>2</sup>, Takashi Tsuruo<sup>2,3</sup>, Hideki Shuto<sup>1</sup>, and Yasufumi Kataoka<sup>1,\*</sup>

<sup>1</sup>Department of Pharmaceutical Care and Health Sciences, Faculty of Pharmaceutical Sciences, Fukuoka University, 8-19-1 Nanakuma, Jonan-ku, Fukuoka 814-0180, Japan

<sup>2</sup>Institute of Molecular and Cellular Biosciences, University of Tokyo, Bunkyo-ku, Tokyo 113-0032, Japan

<sup>3</sup>Cancer Chemotherapy Center, Japanese Foundation for Cancer Research, 3-10-6, Ariake, Koto-ku, Tokyo 135-8550, Japan

Received November 20, 2006; Accepted December 28, 2006

**Abstract.** The aim of the present study was to elucidate the effects of indapamide on ischemic damage to the blood-brain barrier (BBB) *in vitro*. The ischemia/reperfusion conditions employed here significantly decreased the viability of mouse brain capillary endothelial (MBEC4) cells, an effect ameliorated by indapamide. Ischemia increased the permeability of MBEC4 cells to two cellular transport markers, sodium fluorescein and Evan's blue-albumin. Indapamide reduced the ischemia-induced hyperpermeability of cells. These results suggest that indapamide may have a protective role against ischemia-induced injury and dysfunction of the BBB.

**Keywords:** indapamide, ischemia, blood-brain barrier

The blood-brain barrier (BBB) greatly restricts the transport of substances between the blood and the central nervous system. The BBB is primarily formed by brain capillary endothelial cells, which are sealed closely with tight junctions (1). Disruption of the BBB can lead to edema and central nervous system pathology in conditions such as stroke. *In vitro*, hypoxia has been shown to increase the permeability of brain microvascular endothelial cells (2, 3). Targeting the BBB may enhance our understanding of the mechanisms that mediate ischemic brain damage and might aid the development of future treatments for stroke (4, 5).

Indapamide is an indoline derivative of chlorosulfonamide that has both diuretic and antihypertensive properties. Indapamide protects against myocardial ischemia due to its antioxidant action (6). A randomized controlled clinical trial showed that in combination, therapy with perindopril, an angiotensin-converting-enzyme inhibitor, and indapamide reduced the risk of

stroke in hypertensive and non-hypertensive patients to a greater degree, compared to single drug therapy with perindopril alone (7). This clinical evidence suggests a possible involvement of indapamide in protecting against recurrent cerebrovascular disease. The present study was, therefore, aimed at evaluating the effect of indapamide on ischemia/reperfusion-induced damage to cerebral microvascular endothelial cells *in vitro*.

The mouse brain capillary endothelial cells used in the current study (MBEC4) show highly specialized characteristics of brain microvascular endothelial cells (8). MBEC4 cells were isolated from BALB/c mouse brain cortices and immortalized by SV40-transformation (8). Cells were cultured in Dulbecco's modified Eagle's medium (DMEM) (Invitrogen, Carlsbad, CA, USA) supplemented with 10% fetal bovine serum, 100 units/mL of penicillin, and 100 µg/mL of streptomycin. Cells were grown in collagen-coated 24-well plates (21,000 cells/cm<sup>2</sup>, 1.8 cm<sup>2</sup>/well; Corning, Acton, MA, USA) in a humidified atmosphere of 5% CO<sub>2</sub> / 95% air at 37°C. Ischemia was initiated *in vitro* by incubating cells with D-glucose-free Krebs-Ringer buffer (143 mM NaCl, 4.7 mM KCl, 1.3 mM CaCl<sub>2</sub>, 1.2 mM MgCl<sub>2</sub>, 1.0 mM

\*Corresponding author. ykataoka@fukuoka-u.ac.jp

Published online in J-STAGE: March 2, 2007

doi: 10.1254/jphs.SC0060222




NaH<sub>2</sub>PO<sub>4</sub>, 25 mM NaHCO<sub>3</sub>, and 11 mM sucrose, pH 7.4) in an anoxic incubator replaced with 5% CO<sub>2</sub>/95% N<sub>2</sub> for 10 h at 37°C (ischemia conditions). Subsequently, cells were incubated with serum-free DMEM in 5% CO<sub>2</sub>/95% air at 37°C for 1 h (reperfusion conditions). As a control, cells were incubated with normal Krebs-Ringer buffer (143 mM NaCl, 4.7 mM KCl, 1.3 mM CaCl<sub>2</sub>, 1.2 mM MgCl<sub>2</sub>, 1.0 mM NaH<sub>2</sub>PO<sub>4</sub>, 25 mM NaHCO<sub>3</sub>, and 11 mM D-glucose, pH 7.4) for 10 h and subsequently incubated with serum-free DMEM for 1 h in 5% CO<sub>2</sub>/95% air at 37°C (normal conditions). To test whether indapamide protects against ischemia/reperfusion-induced cell death, cells were exposed to 10–50 μM of indapamide (Kyoto Pharmaceutical Industries, Kyoto) or vehicle (0.25% dimethyl Sulfoxide) during conditions of ischemia and reperfusion or normal conditions. Cell viability was measured by the WST-1 assay (Cell Counting Kit-8; Dojindo, Kumamoto).

Endothelial barrier function was evaluated by measuring permeability of cells to sodium fluorescein (Na-F) and Evan's blue-albumin (EBA) as previously described (9). MBEC4 cells were grown on the collagen-coated membrane (1.1 cm<sup>2</sup>, 0.4-μm pore size) of a Transwells™ insert (42,000 cells/cm<sup>2</sup>, Corning) and subsequently exposed to the above-mentioned ischemia conditions for 7 h. As a control, MBEC4 cells were incubated with normal Krebs-Ringer buffer for 7 h (normal conditions). Cells were exposed to 10–100 μM of indapamide or vehicle during ischemia conditions or normal conditions. To initiate transport experiments, the medium was removed and cells were washed with normal Krebs-Ringer buffer. Krebs-Ringer buffer containing 100 μg/mL of Na-F (MW 376; Sigma, St. Louis, MO, USA) and 670 μg/mL EBA bound to 0.1% BSA (MW = 67 kDa) were loaded on to the luminal side of the insert. Samples were removed from the abluminal chamber at 30, 60, 90, and 120 min and immediately replaced with Krebs-Ringer buffer. The concentrations of Na-F and EBA were determined with a CytoFluor Series 4000 fluorescence multiwell plate reader [Ex(λ) 485 ± 10 nm and Em(λ) 530 ± 12.5 nm; PerSeptive Biosystems, Framingham, MA, USA] and an Opsys MR microplate reader (630 nm; DYNEX Technologies, Chantilly, VA, USA), respectively. The permeability coefficient and clearance were calculated according to the method of Dehouck et al. (10), as previously described (9).


Data are expressed as the mean ± S.E.M. Statistical analysis was performed using one-way analyses of variance (ANOVA) followed by Tukey-Kramer's post-hoc test. The difference in means was considered to be significant when the *P* value was less than 0.05.

As shown in Fig. 1, the ischemia (10 h)/reperfusion



**Fig. 1.** Effects of indapamide on the viability of MBEC4 cells subjected to normal (A) or ischemia/reperfusion (B) conditions. Results are expressed as % of cell viability under normal conditions with vehicle treatment (% of vehicle). The % recovery is shown on the right-side of panel B and was calculated using the following equation: [% of vehicle under ischemia/reperfusion conditions with indapamide treatment – % of vehicle under ischemia/reperfusion conditions with vehicle treatment] / [% of vehicle under normal conditions with vehicle treatment – % of vehicle under ischemia/reperfusion conditions with vehicle treatment] × 100. Data are expressed as the mean ± S.E.M. (n = 8–20). ##*P* < 0.01, significant difference from vehicle under normal conditions. \*\**P* < 0.01, significant difference from the vehicle under ischemia/reperfusion conditions.

(1 h) conditions significantly decreased the viability of MBEC4 cells grown on collagen-coated wells, by 26.5% of cells subjected to normal conditions. The effect of indapamide on ischemia/reperfusion-induced damage in MBEC4 cells was concentration-dependent: 20–50 μM indapamide improved recovery by 50%–86%. These concentrations have no effect on cell viability under normal conditions. As shown in Fig. 2, a 7-h period of ischemia resulted in increased permeability of MBEC4 cells grown on collagen-coated membranes to Na-F (paracellular transport marker) and EBA (transendothelial transport marker). Following ischemia, the perme-



**Fig. 2.** Effects of indapamide on Na-F (A) and EBA (B) permeability in MBEC4 cell monolayer after a 7-h exposure to normal or ischemic conditions. The inset in each panel shows the effects of indapamide on permeability of the MBEC4 cell monolayer to Na-F and EBA after a 7-h exposure to normal conditions. The permeability coefficients of Na-F and EBA in the MBEC4 monolayer, after a 7-h exposure to normal conditions with vehicle treatment, were  $3.109 \pm 0.012 \times 10^{-4}$  and  $0.798 \pm 0.003 \times 10^{-4}$  cm/min, respectively. Results are expressed as % of vehicle under normal conditions with vehicle treatment. The % recovery is shown on the right-side of each panel and was calculated by the following equation:  $[\% \text{ of vehicle under ischemia condition with indapamide treatment} - \% \text{ of vehicle under ischemia condition with vehicle treatment}] / [\% \text{ of vehicle under normal condition with vehicle treatment} - \% \text{ of vehicle under ischemic condition with vehicle treatment}] \times 100$ . Data are expressed as the mean  $\pm$  S.E.M. ( $n = 8$ ).  $##P < 0.01$ , significant difference from vehicle under normal conditions.  $*P < 0.05$ ,  $**P < 0.01$ , significant difference from vehicle under ischemic conditions.

ability coefficients of Na-F and EBA were significantly increased to  $223.7 \pm 9.8\%$  and  $518.9 \pm 42.5\%$  of the vehicle under normal conditions, respectively. The viability of MBEC4 cells grown on membranes after a 7-h exposure to ischemic conditions and following termination of the permeability test (2 h) were  $72.8 \pm 3.6\%$  and  $71.3 \pm 1.0\%$  of cells subjected to normal conditions, respectively; there was no difference in cell viability before and after the permeability test. Following a 7-h exposure to indapamide (50–100  $\mu\text{M}$ ) under ischemic

conditions, hyperpermeability of MBEC4 cells to Na-F and EBA was concentration-dependently reduced by 31.9%–47.4% and 41.4%–60.6% of the vehicle, respectively. This effect was not accompanied by a change in cell viability (vehicle:  $72.8 \pm 3.6\%$ , 100 mM:  $75.9 \pm 2.3\%$ ). Under normal conditions, these concentrations of indapamide had no effect on the permeability coefficients of Na-F and EBA in MBEC4 cells. These findings suggest that indapamide may efficiently inhibit ischemia-induced hyperpermeability rather than inhibit

ischemic cell death. MBEC4 cells that were grown on membranes exhibited a higher vulnerability to ischemic cytotoxicity and lower sensitivity to the cytoprotective action of indapamide than those grown on the smooth plastic surface of a well. Further experiments will be required to clarify these issues.

The *in vitro* ischemia/reperfusion conditions used in the present study significantly reduced the viability of MBEC4 cells, and this effect was ameliorated by indapamide. Ischemic conditions increased the permeability of MBEC4 cells to Na-F and EBA, but this hyperpermeability was reduced by indapamide.

Disruption of the BBB is a critical event during cerebral ischemia as it is followed by the passive diffusion of water, leading to vasogenic edema and secondary brain injury. Cerebral edema is a major and fatal complication of acute ischemic stroke. Free radical overproduction after brain ischemia and reperfusion contributes to brain edema and neuronal damage. Thus, an inhibition of free radical formation is likely to prevent the occurrence of brain edema and neuronal damage. Indapamide has an antioxidant effect and has the potential to scavenge oxygen free radicals and their derivatives. Boucher et al. reported that indapamide treatment has a protective effect on cardiac tissue during the early stages of postischemic reperfusion (6). The present findings suggest that indapamide may protect cerebral endothelial cells from ischemic damage due to antioxidation and/or free radical scavenging. This phenomenon may contribute, at least in part, to the mechanisms by which indapamide facilitated protection of perindopril against recurrent stroke in a recent clinical study (7).

Brain capillary endothelial cells form a metabolic and physical barrier that separates the periphery from the brain to maintain cerebral homeostasis. The lack of fenestrations and the presence of tight junctions differentiate brain microvessel endothelial cells from peripheral microvascular endothelium. While adherent junctions and other junctional proteins contribute to cell-to-cell contacts in the paracellular cleft, tight junctions are critical for restricting paracellular diffusion in the cerebral microvasculature. Increased cerebrovascular permeability is a principal factor in the development of cerebral edema after brain ischemia. Several studies have shown a relationship between cyclic AMP (cAMP) levels and permeability of endothelial cell monolayers. For example, elevation of intracellular cAMP concentrations induces high trans-endothelial electrical resistance and increases P-glycoprotein function in brain capillary endothelial cells (11, 12). In addition, when bovine aortic endothelial cells were cultured in hypoxic conditions, cellular cAMP

levels decreased and this phenomenon was associated with an increase in cellular permeability (13). Furthermore, a decrease in cAMP levels was detectable after brain microvascular endothelial cells were exposed to hypoxic conditions for 3 h (14). Moreover, indapamide was shown to augment cAMP production induced by forskolin, an adenylyl cyclase activator, but did not alter basal cAMP levels in cardiomyocytes alone (15). Therefore, indapamide may protect cerebral microvascular endothelial cells from ischemic dysfunction by increasing intracellular cAMP levels. The effects of indapamide on the expression of tight junction-related proteins and on intracellular messengers are now being investigated in rat brain primary cultured endothelial cells.

We present here *in vitro* evidence to suggest a possible protective action of indapamide against ischemia/reperfusion-induced injury and dysfunction of the BBB.

### Acknowledgments

Indapamide was kindly provided by Kyoto Pharmaceutical Industries Co., Ltd. This work was supported, in part, by Grants-in-Aid for Scientific Research ((B) 17390159) from JSPS, Japan and by a research fund from Fukuoka University (No. 003171PK; Sumitomo Pharma Co., Ltd.).

### References

- 1 Pardridge WM. Blood-brain barrier biology and methodology. *J Neurovirol.* 1999;5:556–569.
- 2 Mark KS, Davis TP. Cerebral microvascular changes in permeability and tight junctions induced by hypoxia-reoxygenation. *Am J Physiol Heart Circ Physiol.* 2002;282:H1485–H1494.
- 3 Hayashi K, Nakao S, Nakaoke R, Nakagawa S, Kitagawa N, Niwa M. Effects of hypoxia on endothelial/pericytic co-culture model of the blood-brain barrier. *Regul Pept.* 2004;123:77–83.
- 4 Brown RC, Davis TP. Calcium modulation of adherens and tight junction function: a potential mechanism for blood-brain barrier disruption after stroke. *Stroke.* 2002;33:1706–1711.
- 5 O'Donnell ME, Tran L, Lam TI, Liu XB, Anderson SE. Bumetanide inhibition of the blood-brain barrier Na-K-Cl cotransporter reduces edema formation in the rat middle cerebral artery occlusion model of stroke. *J Cereb Blood Flow Metab.* 2004;24:1046–1056.
- 6 Boucher FR, Schatz CJ, Guez DM, de Leiris JG. Beneficial effect of indapamide in experimental myocardial ischemia. *Am J Hypertens.* 1992;5:22–25.
- 7 PROGRESS Collaborative Group. Randomised trial of a perindopril-based blood-pressure-lowering regimen among 6,105 individuals with previous stroke or transient ischaemic attack. *Lancet.* 2001;358:1033–1041.
- 8 Tatsuta T, Naito M, Oh-hara T, Sugawara I, Tsuruo T. Functional involvement of P-glycoprotein in blood-brain barrier.

- J Biol Chem. 1992;267:20383–20391.
- 9 Dohgu S, Yamauchi A, Takata F, Naito M, Tsuruo T, Higuchi S, et al. Transforming growth factor-beta1 upregulates the tight junction and P-glycoprotein of brain microvascular endothelial cells. *Cell Mol Neurobiol.* 2004;24:491–497.
  - 10 Dehouck MP, Jolliet-Riant P, Bree F, Fruchart JC, Cecchelli R, Tillement JP. Drug transfer across the blood-brain barrier: correlation between in vitro and in vivo models. *J Neurochem.* 1992;58:1790–1797.
  - 11 Kis B, Deli MA, Kobayashi H, Abraham CS, Yanagita T, Kaiya H, et al. Adrenomedullin regulates blood-brain barrier functions in vitro. *Neuroreport.* 2001;12:4139–4142.
  - 12 Deli MA, Abraham CS, Takahata H, Niwa M. Tissue plasminogen activator inhibits P-glycoprotein activity in brain endothelial cells. *Eur J Pharmacol.* 2001;411:R3–R5.
  - 13 Ogawa S, Koga S, Kuwabara K, Brett J, Morrow B, Morris SA, et al. Hypoxia-induced increased permeability of endothelial monolayers occurs through lowering of cellular cAMP levels. *Am J Physiol.* 1992;262:C546–C554.
  - 14 Fischer S, Renz D, Schaper W, Karliczek GF. Effects of barbiturates on hypoxic cultures of brain derived microvascular endothelial cells. *Brain Res.* 1996;707:47–53.
  - 15 Rabkin SW. Indapamide alters the cyclic AMP signal transduction pathway in cardiomyocytes in culture. *Eur J Pharmacol.* 1994;266:117–123.