

18	FLJ10156	NM_019013	Hypothetical protein, cDNA clone MGC:961	Unknown	S16: GTSVRRRSIQHQEQL	Low	41	3	7	7	7:31156
19	ATP6Y0B	NM_004047	ATPase, H+ transporting, lysosomal, 21 kD, V0 subunit C'	A 23-kDa component of vacuolar ATPase that mediates acidification of intracellular organelles	T190: FRSPYSSTIEPLCSPS	Low	43	7	3	8	6:94023
20	FLJ25758	NM_001011541	Hypothetical protein, clone MGC:33355	Unknown	No sites	NA	48	7	7	4	3:3123
					No sites	NA	48	7	7	4	3:60565
					No sites	NA	48	7	7	8	4:18864
					No sites	NA	48	7	7	8	4:18122

Among 1752 proteins on the microarray, 20 were identified as showing a significant interaction, based on the signal intensity value exceeding the median plus three standard deviations of all the fluorescence intensities by analyzing with ProtoArray Prospector software. They are listed with the 14-3-3-binding consensus motif (putative phosphoserine and phosphothreonine indicated by underline) and its stringency level by the Scansite Motif Scanner, the position on the array, and the Z-score calculated as described in Section 2. Abbreviations: FLJ, the full-length long Japan; MGC, mammalian gene collection; NA, not available.

14-3-3-interacting domain is located in the N-terminal segment spanning amino acid residues 2-105 of STAC, and the interaction is independent of serine/threonine-phosphorylation of the binding domain of STAC.

4. Discussion

The present study was designed to rapidly and systematically identify 14-3-3-binding proteins by analyzing a high-density protein microarray. The array included 1752 proteins derived from multiple gene families of biological importance, including cell-signaling proteins, kinases, membrane-associated proteins, and metabolic proteins. In general, protein microarray has its own limitations associated with the expression and purification of a wide variety of target proteins. In the microarray we utilized, the target proteins were expressed in a baculovirus expression system, purified under native conditions, and spotted on the slides to ensure the preservation of native structure, posttranslational modifications, including glycosylation and serine phosphorylation (Culleré et al., 1998; Tennagels et al., 1999), and proper functionality. Immunolabeling of the array with anti-phosphotyrosine (pTyr) antibody indicated that approximately 10–20% of the proteins on the array are phosphorylated (the unpublished data of Invitrogen Technical Service). When this microarray was utilized for kinase substrate identification, most of known kinases immobilized on the array are enzymatically active with the capacity of some degree of autophosphorylation, suggesting that they are certainly phosphorylated on tyrosine, serine, and threonine residues (see the Protoarray application note on <http://www.invitrogen.com/protoarray>). However, we could not currently validate the precise levels of phosphorylation of individual proteins, because of a lack of anti-phosphoserine (pSer) and anti-phosphothreonine (pThr) antibodies suitable for detection of pSer and pThr residues of the proteins on glass slide.

The protein microarray utilized in the present study includes 11 known 14-3-3-binding proteins, such as PCTAIRE protein kinase 1 (PCTK1) (Graeser et al., 2002), protein kinase C zeta (PRKCZ) (van der Hoeven et al., 2000), keratin 18 (KRT18) (Ku et al., 1998), myosin light polypeptide kinase (MYLK) (Haydon et al., 2002), v-abl Abelson murine leukemia viral oncogene homolog 1 (ABL1) (Yoshida et al., 2005), v-akt murine thymoma viral oncogene homolog 1 (AKT1) (Powell et al., 2002), epidermal growth factor receptor (EGFR) (Oksvold et al., 2004), cell division cycle 2 (CDC2) (Chan et al., 1999), mitogen-activated protein kinase kinase 1 (MAP3K1) (Fanger et al., 1998), mitogen-activated protein kinase-activated protein kinase 2 (MAPKAPK2) (Powell et al., 2003), and stratifin (SFN) (Benzinger et al., 2005) (Table 3). All of these were not identified as a 14-3-3-binding protein in the present study. Therefore, the possibility could not be excluded that some 14-3-3 binding partners were not detected due to imperfect phosphorylation of the proteins on the array or due to 14-3-3 isoform-specific binding. Calmodulin, another known 14-3-3 interactor (Luk et al., 1999), was included as a negative control on the array and identified as negative in the present study, because the calcium-dependent interaction between 14-3-3 and calmodulin could not be detected under the calcium-free conditions we employed.

Table 3
Eleven known 14-3-3-binding proteins immobilized on the protein microarray utilized in the present study

Gene name	Database ID	Reference
PCTAIRE protein kinase 1 (PCTK1), transcript variant 3	NM_033019	Graeser et al. (2002)
Protein kinase C, zeta (PRKCZ)	NM_002744	van der Hoeven et al. (2000)
Keratin 18 (KRT18), transcript variant 1	NM_000224	Ku et al. (1998)
Myosin, light polypeptide kinase (MYLK), transcript variant 6	NM_005965	Haydon et al. (2002)
V-abl Abelson murine leukemia viral oncogene homolog 1 (ABL1), transcript variant a	NM_005157	Yoshida et al. (2005)
V-akt murine thymoma viral oncogene homolog 1 (AKT1), transcript variant 1	NM_005163	Powell et al. (2002)
Epidermal growth factor receptor (EGFR), transcript variant 1	NM_005228	Oksvold et al. (2004)
Cell division cycle 2 (CDC2), transcript variant 1	NM_001786	Chan et al. (1999)
Mitogen-activated protein kinase kinase kinase 1 (MAP3K1)	XM_042066	Fanger et al. (1998)
Mitogen-activated protein kinase-activated protein kinase 2 (MAPKAPK2), transcript variant 1	NM_004759	Powell et al. (2003)
14-3-3 Sigma, stratifin (SFN)	NM_006142	Benzinger et al. (2005)

The known 14-3-3-binding proteins, which were spotted on the protein microarray but were not detected in the present study are listed. The 14-3-3-binding proteins validated by definitive evidence are selected and shown with references.

Increasing studies indicate that 14-3-3-binding phosphorylation sites do not exactly fit the consensus motif (Aitken et al., 2002; Ku et al., 1998) and a second site is required to enhance a stable 14-3-3-target interaction (MacKintosh, 2004), and show that the 14-3-3 protein interacts with a set of target proteins in a phosphorylation-independent manner (Dai and Murakami, 2003; Henriksson et al., 2002; Zhai et al., 2001). Supporting the latter possibility, the present observations showed that the interaction of 14-3-3 with target proteins is independent of serine/threonine-phosphorylation of the binding sites of EAP30, DDX54, and STAC. This suggests that substantial numbers of 14-3-3 binding partners identified by protein microarray analysis, if not all, employ phosphorylation-independent binding domains.

All the procedure required for microarray analysis takes approximately 5 h. This analysis identified a set of 20 human proteins as 14-3-3 interactors, most of which were previously unreported except for glutathione *S*-transferase M3 (GSTM3) that was found as one of binding partners by 14-3-3 affinity purification of HeLa cell protein extracts (Pozuelo Rubio et al., 2004). Unexpectedly, the highly stringent 14-3-3-binding consensus motif was identified only in two, such as STAC and HNPRC, by the Scansite Motif Scanner search, while 15 of 20 proteins have one or several motifs when a query with the medium or low stringency was performed (Table 2). The specific binding to 14-3-3 of EAP30, DDX54, and STAC was verified by immunoprecipitation analysis of the recombinant proteins expressed in HEK293 cells. These results indicate that protein microarray is a powerful tool for rapid identification of protein–protein interactions, including those unpredicted by the Database search.

Among the 14-3-3-binding partners we identified, several proteins could be categorized as a component of multimolecular complexes involved in transcriptional regulation. ELL is a human oncogene encoding a RNA polymerase II (Pol II) transcription factor that promotes transcription elongation (Schmidt et al., 1999). EAP30 is a 30-kDa component of the ELL complex where EAP30 confers derepression of transcription by Pol II (Schmidt et al., 1999). A recent study showed that EAP30 could interact with the tumor susceptibility gene TSG101 product, a cellular factor that plays a key role in packaging of HIV

virions (von Schwedler et al., 2003). DDX54 is a 97-kDa RNA helicase (DP97) that interacts with estrogen receptor (ER) and represses the transcription of ER-regulated genes (Rajendran et al., 2003). A recent study by using chromatin immunoprecipitation (ChIP) assay combined with promoter microarray analysis showed that hepatocyte nuclear factor 4- α (HNF4 α), a master regulator of hepatocyte gene expression, interacts with the DDX54 gene promoter, together with Pol II (Odom et al., 2004). HNPRC is a member of heterogeneous nuclear ribonucleoproteins (hnRNPs) involved in pre-mRNA processing, mRNA metabolism and transport (Nakagawa et al., 1986). Increasing evidence indicates that the 14-3-3 protein and its targets are widely distributed in various subcellular compartments, including the nucleus (Dougherty and Morrison, 2004; Meek et al., 2004).

STAC is a 47-kDa cytosolic protein that has a cysteine-rich domain (CRD) of the protein kinase C family in the N-terminal half (NTF), and a src homology three (SH3) domain in the C-terminal half (CTF), suggesting its function as an adapter on which divergent signaling pathways converge (Hardy et al., 2005; Suzuki et al., 1996). STAC is expressed predominantly in the brain with the distribution in a defined population of neurons (Suzuki et al., 1996). IP analysis of mutant and truncated forms argued against an active involvement of the most stringent motif RYYSSP (*pS172*) of STAC in its binding to 14-3-3. The present observations indicated that the 14-3-3-interacting domain is located in the N-terminal segment spanning amino acid residues 2-105 of STAC and the interaction is serine/threonine phosphorylation-independent.

In conclusion, protein microarray is a useful tool for rapid and comprehensive profiling of 14-3-3-binding proteins, although the validation of the results by different methods is highly important.

Acknowledgements

This work was supported by grants from Research on Psychiatric and Neurological Diseases and Mental Health, the Ministry of Health, Labour and Welfare of Japan (H17-020), Research on Health Sciences Focusing on Drug Innovation, the Japan Health Sciences Foundation (KH21101), the Grant-in-Aid for Scientific

Research, the Ministry of Education, Science, Sports and Culture (B2-15390280 and PA007-16017320), and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO), Japan.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [10.1016/j.jneumeth.2005.09.015](https://doi.org/10.1016/j.jneumeth.2005.09.015).

References

- Aitken A, Baxter H, Dubois T, Clokie S, Mackie S, Mitchell K, Peden A, Zemlickova E. 14-3-3 proteins in cell regulation. *Biochem Soc Trans* 2002;30:351–60.
- Benzinger A, Muster N, Koch HB, Yates 3rd JR, Hermeking H. Targeted proteomic analysis of 14-3-3 sigma, a p53 effector commonly silenced in cancer. *Mol Cell Proteomics* 2005;4:785–95.
- Berg D, Holzmann C, Riess O. 14-3-3 proteins in the nervous system. *Nat Rev Neurosci* 2002;4:752–62.
- Chan SM, Ermann J, Su L, Fathman CG, Utz PJ. Protein microarrays for multiplex analysis of signal transduction pathways. *Nat Med* 2004;10:1390–6.
- Chan TA, Hermeking H, Lengauer C, Kinzler KW, Vogelstein B. 14-3-3 σ is required to prevent mitotic catastrophe after DNA damage. *Nature* 1999;401:616–20.
- Chen H-K, Fernandez-Funez P, Acevedo SF, Lam YC, Kaytor MD, Fernandez MH, Aitken A, Skoulakis EM, Orr HT, Botas J, Zoghbi HY. Interaction of Akt-phosphorylated ataxin-1 with 14-3-3 mediates neurodegeneration in spinocerebellar ataxia type 1. *Cell* 2003;113:457–68.
- Culleré X, Rose P, Thathamangalam U, Chatterjee A, Mullane KP, Pallas DC, Benjamin TL, Roberts TM, Schaffhausen BS. Serine 257 phosphorylation regulates association of polyomavirus middle T antigen with 14-3-3 proteins. *J Virol* 1998;72:558–63.
- Dai J-G, Murakami K. Constitutively and autonomously active protein kinase C associated with 14-3-3 ζ in the rodent brain. *J Neurochem* 2003;84:23–34.
- Dougherty MK, Morrison DK. Unlocking the code of 14-3-3. *J Cell Sci* 2004;117:1875–84.
- Fanger GR, Widmann C, Porter AC, Sather S, Johnson GL, Vaillancourt RR. 14-3-3 proteins interact with specific MEK kinases. *J Biol Chem* 1998;273:3476–83.
- Fu H, Subramanian RR, Masters SC. 14-3-3 proteins: structure, function, and regulation. *Annu Rev Pharmacol Toxicol* 2000;40:617–47.
- Graeser R, Gannon J, Poon RYC, Dubois T, Aitken A, Hunt T. Regulation of the CDK-related protein kinase PCTAIRE-1 and its possible role in neurite outgrowth in Neuro-2A cells. *J Cell Sci* 2002;115:3479–90.
- Hardy K, Mansfield L, Mackay A, Benvenuti S, Ismail S, Arora P, O'Hare MJ, Jat PS. Transcriptional networks and cellular senescence in human mammary fibroblasts. *Mol Biol Cell* 2005;16:943–53.
- Haydon CE, Watt PW, Morrice N, Knebel A, Gaestel M, Cohen P. Identification of a phosphorylation site on skeletal muscle myosin light chain kinase that becomes phosphorylated during muscle contraction. *Arch Biochem Biophys* 2002;397:224–31.
- Henriksson ML, Francis MS, Peden A, Aili M, Stefansson K, Palmer R, Aitken A, Hallberg B. A nonphosphorylated 14-3-3 binding motif on exoenzyme S that is functional in vivo. *Eur J Biochem* 2002;269:4921–9.
- Ichimura T, Yamamura H, Sasamoto K, Tominaga Y, Taoka M, Kakiuchi K, Shinkawa T, Takahashi N, Shimada S, Isobe T. 14-3-3 proteins modulate the expression of epithelial Na⁺ channels by phosphorylation-dependent interaction with Nedd4-2 ubiquitin ligase. *J Biol Chem* 2005;280:13187–94.
- Jin J, Smith FD, Stark C, Wells CD, Fawcett JP, Kulkarni S, Metalnikov P, O'Donnell P, Taylor P, Taylor L, Zougman A, Woodgett JR, Langeberg LK, Scott JD, Pawson T. Proteomic, functional, and domain-based analysis of in vivo 14-3-3 binding proteins involved in cytoskeletal regulation and cellular organization. *Curr Biol* 2004;14:1436–50.
- Kawamoto Y, Akiguchi I, Nakamura S, Honjo Y, Shibasaki H, Budka H. 14-3-3 proteins in Lewy bodies in Parkinson disease and diffuse Lewy body disease brains. *J Neuropathol Exp Neurol* 2002;61:245–53.
- Ku N-O, Liao J, Omary MB. Phosphorylation of human keratin 18 serine 33 regulates binding to 14-3-3 proteins. *EMBO J* 1998;17:1892–906.
- Layfield R, Fergusson J, Aitken A, Lowe J, Landon M, Mayer RJ. Neurofibrillary tangles of Alzheimer's disease brains contain 14-3-3 proteins. *Neurosci Lett* 1996;209:57–60.
- Luk SCW, Ngai S-M, Tsui SKW, Fung K-P, Lee C-Y, Waye MMY. In vivo and in vitro association of 14-3-3 epsilon isoform with calmodulin: implication for signal transduction and cell proliferation. *J Cell Biochem* 1999;73:31–5.
- MacBeath G, Schreiber SL. Printing proteins as microarrays for high-throughput function determination. *Science* 2000;289:1760–3.
- MacKintosh C. Dynamic interactions between 14-3-3 proteins and phosphoproteins regulate diverse cellular processes. *Biochem J* 2004;381:329–42.
- Malaspina A, Kaushik N, de Belleroche J. A 14-3-3 mRNA is up-regulated in amyotrophic lateral sclerosis spinal cord. *J Neurochem* 2000;75:2511–20.
- Meek SEM, Lane WS, Pivnicka-Worms H. Comprehensive proteomic analysis of interphase and mitotic 14-3-3-binding proteins. *J Biol Chem* 2004;279:32046–54.
- Michaud GA, Salcius M, Zhou F, Bangham R, Bonin J, Guo H, Snyder M, Predki PF, Schweitzer BI. Analyzing antibody specificity with whole proteome microarrays. *Nat Biotechnol* 2003;21:1509–12.
- Nakagawa TY, Swanson MS, Wold BJ, Dreyfuss G. Molecular cloning of cDNA for the nuclear ribonucleoprotein particle C proteins: a conserved gene family. *Proc Natl Acad Sci USA* 1986;83:2007–11.
- Obenaus JC, Cantley LC, Yaffe MB. Scansite 2.0: Proteome-wide prediction of cell signaling interactions using short sequence motifs. *Nucl Acids Res* 2003;31:3635–41.
- Odom DT, Zitlsperger N, Gordon DB, Bell GW, Rinaldi NJ, Murray HL, Volkert TL, Schreiber J, Rolfe PA, Gifford DK, Fraenkel E, Bell GI, Young RA. Control of pancreas and liver gene expression by HNF transcription factors. *Science* 2004;303:1378–81.
- Oksvold MP, Huitfeldt HS, Langdon WY. Identification of 14-3-3 ζ as an EGF receptor interacting protein. *FEBS Lett* 2004;569:207–10.
- Pozuelo Rubio M, Geraghty KM, Wong BH, Wood NT, Campbell DG, Morrice N, Mackintosh C. 14-3-3-affinity purification of over 200 human phosphoproteins reveals new links to regulation of cellular metabolism, proliferation and trafficking. *Biochem J* 2004;379:395–408.
- Powell DW, Rane MJ, Chen Q, Singh S, McLeish KR. Identification of 14-3-3 ζ as a protein kinase B/Akt substrate. *J Biol Chem* 2002;277:21639–42.
- Powell DW, Rane MJ, Joughin BA, Kalmukova R, Hong J-H, Tidor B, Dean WL, Pierce WM, Klein JB, Yaffe MB, McLeish KR. Proteomic identification of 14-3-3 ζ as a mitogen-activated protein kinase-activated protein kinase 2 substrate: role in dimmer formation and ligand binding. *Mol Cell Biol* 2003;23:5376–87.
- Rajendran RR, Nye AC, Frasor J, Balsara RD, Martini PG, Katzenellenbogen BS. Regulation of nuclear receptor transcriptional activity by a novel DEAD box RNA helicase (DP97). *J Biol Chem* 2003;278:4628–38.
- Satoh J, Kuroda Y. Differential gene expression between human neurons and neuronal progenitor cells in culture: an analysis of arrayed cDNA clones in NTERa2 human embryonal carcinoma cell line as a model system. *J Neurosci Meth* 2000;94:155–64.
- Satoh J, Yamamura T. Gene expression profile following stable expression of the cellular prion protein. *Cell Mol Neurobiol* 2004;24:793–814.
- Satoh J, Yamamura T, Arima K. The 14-3-3 protein ϵ isoform expressed in reactive astrocytes in demyelinating lesions of multiple sclerosis binds to vimentin and glial fibrillary acidic protein in cultured human astrocytes. *Am J Pathol* 2004;165:577–92.
- Schmidt AE, Miller T, Schmidt SL, Shiekhattar R, Shilatifard A. Cloning and characterization of the EAP30 subunit of the ELL complex that confers derepression of transcription by RNA polymerase II. *J Biol Chem* 1999;274:21981–5.
- Suzuki H, Kawai J, Taga C, Yaoi T, Hara A, Hirose K, Hayashizaki Y, Watanabe S. Stac, a novel neuron-specific protein with cysteine-rich and SH3 domains. *Biochem Biophys Res Commun* 1996;229:902–9.

- Tennagels N, Hube-Magg C, Wirth A, Noelle V, Klein HW. Expression, purification, and characterization of the cytoplasmic domain of the human IGF-1 receptor using a baculovirus expression system. *Biochem Biophys Res Commun* 1999;260:724–8.
- van der Hoeven PCJ, van der Wal JCM, Ruurs P, van Blitterswijk WJ. Protein kinase C activation by acidic proteins including 14-3-3. *Biochem J* 2000;347:781–5.
- van Hemert MJ, Steensma HY, van Heusden GPH. 14-3-3 proteins: key regulators of cell division, signaling and apoptosis. *Bioessays* 2001;23:936–47.
- Vidalain PO, Boxem M, Ge H, Li S, Vidal M. Increasing specificity in high-throughput yeast two-hybrid experiments. *Methods* 2004;32:363–70.
- von Mering C, Krause R, Snel B, Cornell M, Oliver SG, Fields S, Bork P. Comparative assessment of large-scale data sets of protein–protein interactions. *Nature* 2002;417:399–403.
- von Schwedler UK, Stuchell M, Müller B, Ward DM, Chung HY, Morita E, Wang HE, Davis T, He GP, Cimbara DM, Scott A, Kräusslich HG, Kaplan J, Morham SG, Sundquist WI. The protein network of HIV budding. *Cell* 2003;114:701–13.
- Yoshida K, Yamaguchi T, Natsume T, Kufe D, Miiki Y. JNK phosphorylation of 14-3-3 proteins regulates nuclear targeting of c-Abl in the apoptotic response to DNA damage. *Nat Cell Biol* 2005;7:278–85.
- Zerr I, Bodemer M, Gefeller O, Otto M, Poser S, Wiltfang J, Windl O, Kretzschmar HA, Weber T. Detection of 14-3-3 protein in the cerebrospinal fluid supports the diagnosis of Creutzfeldt–Jakob disease. *Ann Neurol* 1998;43:32–40.
- Zhai J, Lin H, Shamim M, Schlaepfer WW, Cañete-Soler R. Identification of a novel interaction of 14-3-3 with p190RhoGEF. *J Biol Chem* 2001;276:41318–24.
- Zhang LV, Wong SL, King OD, Roth FP. Predicting co-complexed protein pairs using genomic and proteomic data integration. *BMC Bioinform* 2004;5:38–52.
- Zhu H, Bilgin M, Bangham R, Hall D, Casamayor A, Bertone P, Lan N, Jansen R, Bidlingmaier S, Houfek T, Mitchell T, Miller P, Dean RA, Gerstein M, Snyder M. Global analysis of protein activities using proteome chips. *Science* 2001;293:2101–5.

T cell gene expression profiling identifies distinct subgroups of Japanese multiple sclerosis patients

Jun-ichi Satoh^a, Megumi Nakanishi^a, Fumiko Koike^a, Hiroyuki Onoue^a, Toshimasa Aranami^a,
Toshiyuki Yamamoto^b, Mitsuru Kawai^b, Seiji Kikuchi^c, Kyouichi Nomura^d,
Kazumasa Yokoyama^e, Kohéi Ota^f, Toshiro Saito^g, Masayuki Ohta^g, Sachiko Miyake^a,
Takashi Kanda^h, Toshiyuki Fukazawaⁱ, Takashi Yamamura^{a,*}

^a Department of Immunology, National Institute of Neuroscience, NCNP, 4-1-1 Ogawahigashi, Kodaira, Tokyo 187-8502, Japan

^b National Center Hospital for Mental, Nervous and Muscular Disorders, NCNP, Tokyo 187-8502, Japan

^c Department of Neurology, Hokkaido University Graduate School of Medicine, Sapporo 060-8638, Japan

^d Department of Neurology, Saitama Medical School, Saitama 350-0495, Japan

^e Department of Neurology, Juntendo University School of Medicine, Tokyo 113-841, Japan

^f Department of Neurology, Tokyo Women's Medical University, Tokyo 162-8666, Japan

^g Life Science Group, Hitachi Ltd., Saitama 350-1165, Japan

^h Department of Neurology, Tokyo Medical and Dental University, Tokyo 113-8519, Japan

ⁱ Hokuyukai Neurology Hospital, Sapporo 063-0802, Japan

Received 29 November 2005; received in revised form 30 January 2006; accepted 3 February 2006

Abstract

To clarify the molecular background underlying the heterogeneity of multiple sclerosis (MS), we characterized the gene expression profile of peripheral blood CD3⁺ T cells isolated from MS and healthy control (CN) subjects by using a cDNA microarray. Among 1258 cDNAs on the array, 286 genes were expressed differentially between 72 untreated Japanese MS patients and 22 age- and sex-matched CN subjects. When this set was used as a discriminator for hierarchical clustering analysis, it identified four distinct subgroups of MS patients and five gene clusters differentially expressed among the subgroups. One of these gene clusters was overexpressed in MS versus CN, and particularly enhanced in the clinically most active subgroup of MS. After 46 of the MS patients were treated with interferon-beta (IFN β -1b) for two years, IFN β responders were clustered in two of the four MS subgroups. Furthermore, the IFN β responders differed from nonresponders in the kinetics of IFN-responsive genes at 3 and 6 months after starting IFN β treatment. These results suggest that T-cell gene expression profiling is valuable to identify distinct subgroups of MS associated with differential disease activity and therapeutic response to IFN β .

© 2006 Elsevier B.V. All rights reserved.

Keywords: Gene expression profile; Hierarchical clustering analysis; IFN β responder; Microarray; Multiple sclerosis; T cells

1. Introduction

Multiple sclerosis (MS) is an inflammatory demyelinating disease of the central nervous system (CNS) white matter mediated by an autoimmune process whose deve-

lopment is triggered by a complex interplay of both genetic and environmental factors (Compston and Coles, 2002). Intravenous administration of interferon-gamma (IFN γ) to MS patients in a previous clinical trial provoked acute relapses accompanied by activation of the systemic immune response, indicating a central role of proinflammatory T helper type 1 (Th1) lymphocytes in the immunopathogenesis of MS (Panitch et al., 1987). In contrast, treatment with interferon-beta (IFN β) produced a

* Corresponding author. Tel.: +81 42 346 1723; fax: +81 42 346 1753.

E-mail address: yamamura@ncnp.go.jp (T. Yamamura).

beneficial effect on MS patients with a reduction of the relapse rate by approximately 30% (The IFN β Multiple Sclerosis Study Group, 1993; Jacobs et al., 1996; Saida et al., 2005). Recent studies indicated that an early initiation of IFN β delays the conversion to clinically definite MS in the patients who experienced a first demyelinating event (Jacobs et al., 2000).

MS exhibits a great range of phenotypic variability. It is classified into relapsing–remitting MS (RRMS), secondary progressive MS (SPMS), or primary progressive MS (PPMS) with respect to the disease course, conventional MS (CMS) or opticospinal MS (OSMS) in terms of the lesion distribution (Saida et al., 2005), and IFN β responder or nonresponder based on the therapeutic response to IFN β (Waubant et al., 2003). MS brain lesions show a remarkable heterogeneity in the degree of inflammation, complement activation, antibody deposition, demyelination and remyelination, oligodendrocyte apoptosis, and axonal degeneration (Lucchinetti et al., 2000). These observations suggest that MS is a kind of neurological syndrome caused by different immunopathological mechanisms leading to the final common pathway that provokes inflammatory demyelination. Therefore, it is not surprising to find that individual MS patients show highly variable responses to IFN β treatment. Currently, very little is known about the molecular background underlying clinical and pathological heterogeneity of MS.

DNA microarray technology is a novel approach that allows us to systematically monitor the expression of a large number of genes in disease-affected tissues (Staudt, 2001). This approach has discovered therapeutically relevant targets and prognostic markers for cancers (Alizadeh et al., 2000; van de Vijver et al., 2000), and has given new insights into the complexity of molecular interactions promoting the autoimmune process in MS (Steinman and Zamvil, 2003). Importantly, the comprehensive gene expression profiling of MS brain tissues and peripheral blood lymphocytes identified a battery of genes deregulated in MS, whose role has not been previously predicted in its pathogenesis (Lock et al., 2002; Graumann et al., 2003; Tajouri et al., 2003; Stürzbecher et al., 2003; Achiron et al., 2004). However, most of previous studies have focused on gene expression in heterogeneous populations of unfractionated lymphocytes and brain cells. Recently, by using microarray we showed that IFN β treatment elevates the expression of 7 IFN-responsive genes in highly purified peripheral blood CD3⁺ T cells of 13 Japanese RRMS patients (Koike et al., 2003). More recently, we found that the majority of differentially expressed genes in CD3⁺ T cells between 72 untreated MS patients and 22 healthy control (CN) subjects were categorized into apoptosis signaling-related genes (Satoh et al., 2005).

To extend our previous studies, we conducted hierarchical clustering analysis of differentially expressed genes between MS and CN in peripheral blood CD3⁺ T cells. Here we report that T-cell gene expression profiling classifies a

heterogeneous population of Japanese MS into four subgroups that differ in the disease activity and therapeutic response to IFN β , suggesting that this analysis could be applied for designing tailor-made treatment of MS.

2. Subjects and methods

2.1. The study population

The Research Group for IFN β treatment of Japanese MS, sponsored by the Ministry of Health, Labour and Welfare of Japan, conducted the present study. It enrolled 72 clinically active Japanese MS patients, including 65 RRMS and 7 SPMS cases composed of 55 women and 17 men with the mean age of 36.1 ± 10.3 years, and 22 healthy control (CN) subjects composed of 16 women and 6 men with the mean age of 38.6 ± 12.3 years. The members of this research group (SK, KN, KY, KO, TK, TF and TY), all of who are certified neurologists, diagnosed individual cases according to the established criteria (McDonald et al., 2001), and followed up the patients for at least two years after entry. The patients showed the mean Expanded Disability Status Scale (EDSS) score of 2.8 ± 2.0 upon entry. No patients had a history of treatment with interferons, glatiramer acetate or mitoxantrone before enrollment, or received corticosteroids and other immunosuppressants during at least one month before blood sampling. MS patients were divided into two groups according to their own determination upon entry: one treated with IFN β and the other without IFN β . The IFN β -treated group included 46 patients who started to receive an administration of 8 million units of IFN β -1b (Betaferon, Schering, Osaka, Japan) for two years given subcutaneously on alternate days, while the IFN β -untreated group included 26 patients who were followed up without IFN β treatment for successive two years. From the IFN β -treated group, blood samples were taken at three time points: before starting IFN β treatment (designated Pre) and at 3 and 6 months after starting the treatment. In the IFN β -untreated group, they were collected twice: at enrollment and at 6 months after the enrollment. In case of acute relapse, the patients in both groups were given intravenous methylprednisolone pulse (IVMP) following the standard protocol, although none received glatiramer acetate, mitoxantrone, or other immunosuppressants. The samples obtained during clinically obvious relapses or episodes of infection were omitted. Written informed consent was obtained from all the subjects. The present study was approved by the Ethics Committee of National Center of Neurology and Psychiatry (NCNP).

2.2. IFN β responder/nonresponder score

To evaluate the therapeutic response to IFN β , we monitored the following six parameters during four years spanning two years before and after initiation of IFN β

treatment; the number of clinical relapse, the day of IVMP treatment, the day of hospitalization, EDSS score, the number of lesions on T2-weighted MRI, and the patient's satisfaction on the treatment (Table 1). When compared before and after IFN β treatment, these parameters have given three ranks and scores; good (+1), intermediate (0), and poor (–1). The total score was calculated for each patient, ranging from the maximum value of +6 to the minimum value of –6. The patients with the total score equal to or greater than +3 were considered as being the responder (R), the score from 0 to +2 as one with the undetermined response (UD), and the score equal to or smaller than –1 as the nonresponder (NR) (Table 1).

2.3. cDNA microarray analysis

The present study utilized a custom microarray containing duplicate spots of 1258 cDNA immobilized on a poly-L-lysine-coated slide glass. They were composed of well annotated genes of various functional classes, including cytokines/growth factors and their receptors, apoptosis regulators, oncogenes, transcription factors, signal transducers, cell cycle regulators and housekeeping genes (Hitachi Life Science, Kawagoe, Saitama, Japan: <http://www.hitachi.co.jp/LS>). Peripheral blood mononuclear cells (PBMC) were isolated from 30 ml of heparinized blood by centrifugation on a Ficoll density gradient. They were labeled with anti-CD3 antibody-coated magnetic microbeads (#130-050-101, Miltenyi Biotec, Auburn, CA), and CD3⁺ T cells were separated by AutoMACS (Miltenyi Biotec). The remaining cells after the positive selection of CD3⁺ T cells were harvested as CD3[–] non-T cell fraction as described previously (Koike et

al., 2003; Satoh et al., 2005). Total RNA was isolated from the cells by using RNeasy Mini Kit (Qiagen, Valencia, CA). Five micrograms of purified RNA was in vitro amplified, and the antisense RNA (aRNA) of MS patients and CN subjects was labeled with a fluorescent dye Cy5, while pooled aRNA of three independent healthy volunteers who were not included in the present study was labeled with Cy3 for a universal reference to standardize the gene expression levels throughout the experiments. The arrays were hybridized at 62 °C for 10 h in the hybridization buffer containing equal amounts of Cy3- or Cy5-labeled cDNA, and they were then scanned by the ScanArray 5000 scanner (GSI Lumonics, Boston, MA). The data were analyzed by using the QuantArray software (GSI Lumonics). The average of fluorescence intensities (FI) of duplicate spots was obtained after global normalization between Cy3 and Cy5 signals. The gene expression level (GEL) was calculated according to the formula: GEL = FI (Cy5) of the sample / FI (Cy3) of the universal reference.

2.4. Hierarchical clustering analysis, principal component analysis, and statistical analysis

The genes whose expression was significantly different between MS and CN groups were identified by using *perre* of the “R” statistical software system (www.cran.r-project.org) based on a Bayesian framework for analysis of microarray expression data (Baldi and Long, 2001). The error rate of this test smaller than 0.25 following the Bonferroni correction was considered as significant. Hierarchical clustering analysis and principal component analysis (PCA) were performed on a set of 286 genes differentially expressed between MS and CN groups, which were selected

Table 1
IFN β responder/nonresponder score

Category	The parameters	Rank and score of the therapeutic response		
		Poor	Intermediate	Good
#1	Number of relapse after 2 years/number of relapse before 2 years Score	≥ 1.5 (–1)	1.5–0.5 0	≤ 0.5 (+1)
#2	Number of IVMP treatment after 2 years/number of IVMP treatment before 2 years Score	≥ 1.5 (–1)	1.5–0.5 0	≤ 0.5 (+1)
#3	Day of hospitalization after 2 years/day of hospitalization before 2 years Score	≥ 1.5 (–1)	1.5–0.5 0	≤ 0.5 (+1)
#4	EDSS score before treatment-EDSS score in 2 years after treatment Score	≥ -0.5 (–1)	0.5–(–0.5) 0	≥ 0.5 (+1)
#5	Number of lesions on T2-weighted MRI in 2 years after treatment/number of lesions on T2-weighted MRI before treatment Score	≥ 1.2 (–1)	1.2–0.8 0	≤ 0.8 (+1)
#6	Patient's satisfaction Score	Unsatisfied (–1)	Neither satisfied nor unsatisfied 0	Satisfied (+1)

The total responder/nonresponder score of six categories ranges from the maximum value of +6 to the minimum value of –6. The patients with the score equal to +3 or greater than +3 were classified as responder (R), the score ranging from 0 to +2 as undetermined (UD), and the score equal to –1 or smaller than –1 as nonresponder (NR). Abbreviations: IVMP, intravenous methylprednisolone pulse.

as a discriminator for a standard \times standard algorithm on GeneSpring 7.2 (Agilent Technologies, Palo Alto, CA). The differences in clinical parameters among MS subgroups were evaluated by multiple comparison test following the Bonferroni correction.

3. Results

3.1. Microarray analysis identified 286 genes differentially expressed in peripheral blood T cells between MS and control subjects

Among 1258 genes on the microarray, 286 genes were expressed differentially in peripheral blood CD3⁺ T cells between 72 untreated MS patients and 22 CN subjects. Among them, 78 genes were upregulated, while 208 genes

downregulated in MS versus CN (Supplementary Table 1 online for all datasets). We also conducted the microarray analysis of CD3⁻ non-T cells, composed of B cells, monocytes/macrophages and NK cells, and found that 96 genes were differentially expressed in the non-T cell fraction between MS and CN (data not shown).

3.2. Hierarchical clustering analysis identified four distinct subgroups of MS and five gene classes

Hierarchical clustering analysis was performed on CD3⁺ T-cell samples of 72 untreated MS patients and 22 CN subjects, by using the set of 286 differentially expressed genes described above as a discriminator. This unsupervised approach, which arranged the genes and samples with a similar expression pattern to make a cluster in the dendrogram, identified four distinct




Fig. 1. Hierarchical clustering analysis of 286 genes differentially expressed between untreated MS patients and control subjects. The gene expression profile of peripheral blood CD3⁺ T cells was studied in 72 untreated MS patients and 22 age- and sex-matched healthy control (CN) subjects, by using a 1258 cDNA microarray. Hierarchical clustering analysis was performed by selecting a set of 286 genes differentially expressed between MS and CN as a discriminator. The results are expressed in a matrix format, with each row representing the gene expression level (GEL) of a single gene in all the subjects and each column representing GEL of 286 genes in an individual subject. The matrix is shown by a pseudo-color, with red expressing upregulation, green expressing downregulation, and the color intensity representing the magnitude of the deviation from GEL 1.0 as shown on the upper right. Hierarchical clustering analysis separated MS (purple) from CN (dark blue), and classified the former into four subgroups named A (green), B (light blue), C (red) and D (yellow). The 286 genes were categorized into five classes numbered #1 (pink) to #5 (light blue).




Fig. 2. Principal component analysis of 286 discriminator genes. Principal component analysis (PCA), which reduces all of the variance in the original dataset to three dimensions accounting for a significant fraction of the variance, verified a clear separation of the CN group (dark blue) and four MS subgroups named A (green), B (light blue), C (red) and D (yellow) identified by hierarchical clustering analysis.

subgroups of MS, clearly separated from the CN group (Fig. 1). We operationally designated each subgroup of MS as A, B, C and D, following the relative location in the dendrogram (Fig. 1). Principal component analysis (PCA) verified a clear discrimination of four MS subgroups and CN group (Fig. 2). Among 94 subjects examined, two MS patients and three CN subjects were considered as being unclassifiable (UC). In contrast, the clustering analysis of CD3⁺ non-T cells did not clearly separate MS subgroups from CN (data not shown). Hierarchical clustering analysis categorized 281 of 286 differentially expressed genes into five distinct classes numbered #1 to #5 (Fig. 1 and Supplementary Table 1 online for all datasets). The remaining five, including TOP1, CHST4, SLC35A1, ST1B2, and TAF2H, were unable to be categorized into any classes. All the class #5 genes were upregulated in MS, whereas the genes of classes #1 to #4 were downregulated in MS, when compared with CN (Fig. 1). Upregulation of several class #5 genes in MS was validated by quantitative real-time RT-PCR analysis (data not shown).

3.3. Association of MS subgroups with gene clusters

Expression of the class #5 genes were elevated in all MS subgroups, whereas the classes #1 to #4 genes were downregulated in all of them, although the present study could not identify the marker genes specific for each MS subgroup. The subgroup A showed the gene expression pattern that is the most similar to CN. The similarity was supported by a partial overlap between A and CN in PCA (Fig. 2), and by the observations that one CN subject was incorporated in A, while two MS patients of A were included in CN (Fig. 1). Notably, the subgroup B showed the greatest upregulation of class #5 genes and the most prominent suppression of classes #1 to #4 genes (Fig. 1).

The class #5 genes ($n=78$) contain nine chemokines (11.5%), including CCL1, CCL3, CCL13, CCL18, CCL24, CXCL1, CXCL2, CXCL9, and CXCL14. In contrast, the classes #1 to #4 genes ($n=203$) contained only two chemokines (1.0%), such as CXCL5 and CXCL10. These observations suggest that the class #5 gene cluster is highly enriched in chemokine genes.

Fig. 3. Clinical characteristics of microarray-determined four MS subgroups. MS patients were classified into four distinct subgroups named A, B, C, and D by hierarchical clustering analysis. The bar indicates the data of individual patients. The number of relapse, the day of IVMP treatment, the day of hospitalization, and the number of lesions on T2-weighted MRI represent the data of 2 years before enrollment. Abbreviations: EDSS, Expanded Disability Status Scale; IVMP, intravenous methylprednisolone pulse; R/NR, responder/nonresponder.

MS subgroup




Table 2
The therapeutic response to IFN β in microarray-determined four MS subgroups

	Total	A	B	C	D	UC
IFN β -treated patients (<i>n</i>)	46	14	14	11	5	2
Age of IFN β -treated patients (average, SD)	34.9 \pm 9.2	33.2 \pm 7.6	36.5 \pm 10.4	33.1 \pm 8.3	36.2 \pm 13.3	41.5
Male to female ratio of IFN β -treated patients	8 to 38	1 to 13	3 to 11	3 to 8	0 to 5	1 to 1
IFN β responder/nonresponder score (average, SD)	1.9 \pm 2.6	2.5 \pm 2.3	2.1 \pm 2.6	1.3 \pm 2.1	-0.3 \pm 4.0	3
Dropout during a follow-up (<i>n</i>)	7	2	0	3	2	0
IFN β responder (<i>n</i> , %)	19 (41.3%)	8 (57.1%)	8 (57.1%)	2 (18.2%)	0 (0%)	1 (50%)
IFN β nonresponder (<i>n</i>)	7	2	3	1	1	0
Undetermined group (<i>n</i>)	13	2	3	5	2	1
The patients with IFN β -related adverse effects (<i>n</i> , %)	29 (63.0%)	8 (57.1%)	9 (64.3%)	7 (63.6%)	4 (80%)	1 (50%)
Increase in the number of lesions on T2-weighted MRI during a follow-up (average, SD)	1.7 \pm 1.9	-2.0 \pm 1.7	2.8 \pm 6.6	7.6 \pm 15.8	-0.7 \pm 8.1	-3.5
The patients satisfied with IFN β treatment (<i>n</i> , %)	17 (37.0%)	8 (57.1%)	6 (42.9%)	2 (18.2%)	0 (0%)	1 (50%)
The patients neither satisfied nor unsatisfied with IFN β treatment (<i>n</i>)	21	4	7	7	2	1
The patients unsatisfied with IFN β treatment (<i>n</i>)	8	2	1	2	3	0

Among 72 MS patients, 46 patients were treated with IFN β for two years after enrollment. The therapeutic response was evaluated by IFN β responder/nonresponder score shown in Table 1. Abbreviations: UC, unclassifiable.

3.4. Clinical characteristics of microarray-determined MS subgroups

Next, we investigated clinical characteristics of four MS subgroups (Supplementary Table 2 online and Fig. 3). No statistically significant differences were found among the subgroups in the age, disease duration, EDSS score, and the number of lesions on T2-weighted MRI at enrollment. However, there was a trend that the subgroup D showed a greater EDSS score and had a larger number of MRI lesions, suggestive of an advanced stage of the disease (Supplementary Table 2 online). The female outnumbered the male in all the subgroups. The male to female ratio was relatively higher in C, while no male patient was included in D. The patients with RRMS outnumbered those with SPMS in all the subgroups, although there was a mild bias for SPMS in B. The number of relapse, the day of IVMP treatment, and the day of hospitalization during preceding two years before enrollment were the largest and longest in subgroup B, and this difference was statistically significant, when compared between subgroups B and C ($p=0.0128$, 0.0183 , and 0.0329 for each parameter) (Supplementary Table 2 online and Fig. 3). These observations indicate that the subgroup B included the patients who were the clinically most active before starting IFN β .

In all MS subgroups, the conventional form of MS (CMS) greatly outnumbered non-CMS, the latter was composed of the optico-spinal form (OSMS) and multifocal recurrent myelitis without optic nerve involvement. No obvious association was identified between a particular MS subgroup and the spinal cord involvement. However, 5 of 6 patients having the lesions restricted to the cerebrum (CBR) were included in subgroup C (Supplementary Table 2 online). These observations suggest that the status of T-cell gene expression might affect the lesion distribution in this subgroup.

3.5. IFN β responders were clustered in subgroups A and B

Based on the patient's own determination at enrollment, 72 MS patients were separated into two groups; 46 who started to receive IFN β treatment for following two years, and 26 who were followed up without IFN β treatment for successive two years (Supplementary Table 3 online). All the IFN β -treated patients were evaluated by the IFN β responder/nonresponder score (Table 1) at the end of the two year-treatment. They were classified into 19 IFN β responders, 7 nonresponders, 13 undetermined subjects, and 7 dropouts (Table 2). The difference in the score among the subgroups (A: 2.5 ± 2.3 ; B: 2.1 ± 2.6 ; C: 1.3 ± 2.1 ; and D: -0.3 ± 4.3) did not reach the level of statistical significance (Table 2). However, there existed a trend that IFN β responders were clustered either in subgroup A or B. Because the subgroup A contains the greatest proportion of IFN β responders (57.1%), the patients of A were judged as being the most IFN β responsive (Table 2). All the responders of A expressed a satisfaction on IFN β treatment. The patients of the subgroup B also showed a good response equivalent to A (57.1%), although the number of satisfied patients was smaller. In contrast, only 2 of 11 IFN β -treated patients in subgroup C (18.2%) and none of the patients in subgroup D were judged as IFN β responders. The patients of C showed a trend for great increase in the number of MRI lesions during IFN β treatment, consistent with the poor response to IFN β (Table 2). A battery of IFN β treatment-related adverse effects, including skin reactions, flu-like symptoms, leukocytopenia, depression, and amenorrhea, were observed in more than 50% of IFN β -treated patients in all the subgroups (Table 2). Seven patients of the IFN β -treated group discontinued the treatment: five due to adverse effects, one due to a severe relapse, and another by a personal reason.

We also studied T-cell gene expression profile of IFN β -treated MS patients at 3 or 6 months after starting the

treatment. Although hierarchical clustering analysis classified these patients into several subgroups, they did not match with the subgroup A, B, C, or D determined at pretreatment (data not shown). Furthermore, no significant association was identified between these new clusters and the response to IFN β . These observations suggest that T-cell gene expression profiling at pretreatment is the most valuable to predict the clinical outcome, whereas the analysis after starting IFN β treatment is less informative.

3.6. Temporal profile of IFN-responsive gene expression in the first six months discriminated responders and nonresponders

Finally, we investigated the temporal expression profile of the genes with IFN-responsive promoter elements named IFN-responsive genes (IRGs) following IFN β treatment. As we previously reported (Koike et al., 2003), IFN β treatment for 6 months enhanced the expression of a battery of IRGs in T cells (Fig. 4). A remarkable difference was found between IFN β responders (R) and nonresponders (NR) in the kinetics of several IRGs, such as IFN-stimulated protein 15 (ISG15), small inducible cytokine A2 (SCYA2, CCL2, or MCP-1), TNF receptor subfamily member 1B (TNFRSF1B, TNFRp75),

and IFN α -inducible protein 27 (IFI27) (Fig. 5). The IFN β responders exhibited a pattern of persistent upregulation during 6 months of the treatment. In contrast, the nonresponders showed a seesaw pattern, i.e. higher upregulation at 3 months than the responders, followed by substantial downregulation at 6 months. The differences between R and NR in the kinetics of both TNFRSF1B and IFI27 levels from 3 to 6 months were statistically significant ($p=0.0092$ and 0.0307 , respectively) (Fig. 5). These observations suggest that IFN β nonresponders also well respond to IFN β at 3 months, but they could not maintain the responsiveness until 6 months.

4. Discussion

To elucidate the molecular basis underlying clinicopathological variability of MS, we conducted a comprehensive study that combines T-cell gene expression profiling and clinical characteristics of Japanese MS patients. Hierarchical clustering analysis of 286 genes differentially expressed between 72 untreated MS patients and 22 CN subjects classified a clinically heterogeneous population of MS into four distinct subgroups, named A, B, C, and D, and identified five gene classes numbered #1 to #5. The class




Fig. 4. Induction of IFN-responsive genes in IFN β -treated MS patients. Seventy-two MS patients were divided into IFN β -treated group (IFN β +; $n=46$) and untreated group (IFN β -; $n=26$). A cluster of known IFN-responsive genes (IRGs) indicated by the star were significantly upregulated exclusively in IFN β -treated patients at 6 months after starting the treatment.




Fig. 5. Temporal profile of induction of IFN-responsive genes in IFN β responders and nonresponders during IFN β treatment. T-cell gene expression profiling was examined in 46 IFN β -treated MS patients at three time points: before starting IFN β treatment (Pre), and at 3 months (3M) and 6 months (6M) after starting the treatment. Based on the IFN β responder/nonresponder score, they were separated into 19 IFN β responders (R: the left) and 7 nonresponders (NR: the right) as shown in Table 2. The temporal expression profile of IFN-responsive genes (IRGs), such as IFN-stimulated protein 15 (ISG15), small inducible cytokine A2 (SCYA2), TNF receptor subfamily member 1B (TNFRSF1B), and IFN α -inducible protein 27 (IFI27), is shown with the statistical differences in the kinetics during 3 to 6 months between R and NR. The vertical axis represents the gene expression level (GEL), while the horizontal axis indicates the time course.

#5 genes containing nine chemokines were upregulated exclusively in MS. The most clinically active subgroup B showed the highest upregulation of the class #5 genes. These observations suggest that the higher disease activity of B is in part attributable to overproduction of chemokines which promote lymphocyte and macrophage trafficking into the CNS (Balashov et al., 1999; Opdenakker et al., 2003). Recently, using this database as a training set for support vector machine (SVM) analysis of T cell gene expression, we found that the great majority of active RRMS patients were classified into MS subgroups, while clinically stable patients without obvious relapses for several years were occasionally classified into CN group (unpublished observations).

The class #5 genes upregulated in MS T cells include various cytokines, growth factors and their receptors, whose expression was detected at high levels in demyelinating lesions of MS, such as IL-12p40 (Windhagen et al., 1995), IL-10 (Hulshof et al., 2002), granulocyte colony-stimulating factor (G-CSF) (Lock et al., 2002), platelet-derived growth factor receptor- α (PDGFRA) (Maeda et al., 2001), transforming growth factor- β 2 (TGFB2) (Peress et al., 1996), and insulin-like growth factor-II (IGF-II) (Gveric et al., 1999). The class #5 genes also contain many apoptosis-signaling regulators pivotal for T cell development. It is worthy to note that nuclear receptor subfamily 4, group A, member 2 (NR4A2) in the class #5 was the most strongly upregulated gene in MS T cells. NR4A2 encodes an orphan member of the steroid-thyroid hormone receptor superfamily transcription factors designated Nurrl. Nurrl is induced

in T cells during apoptosis (Okabe et al., 1995) and the members of this family regulate clonal deletion of self-reactive T cells in the thymus (Zhou et al., 1996). Nurrl activates the transcription of osteopontin (Lammi et al., 2004), a Th1 cytokine that plays a key role in progression of inflammatory demyelination in MS (Steinman and Zamvil, 2003). In contrast, a previous study showed that NR4A2 is downregulated in unfractionated PBMC of MS patients (Achiron et al., 2004), although the following study from the same group indicated a significant upregulation of NR4A2 in PBMC of both MS and SLE patients (Mandel et al., 2004). We validated upregulation of NR4A2 mRNA levels in MS T cells by quantitative real-time RT-PCR analysis (unpublished observations).

The present study suggests that the microarray-based classification of MS is useful to predict therapeutic response to IFN β . The proportion of IFN β responders greatly differed among MS subgroups: 57.1% in A, 57.1% in B, 18.2% in C and 0% in D. Furthermore, the responders are significantly different from the non-responders in the kinetics of IFN-responsive genes (IRGs). A panel of IRGs were upregulated persistently in IFN β responders, whereas they were downregulated to some extent in nonresponders by 6 months. The precise reason for downregulation of IRGs in nonresponders after a long-term treatment remains unknown, because our study did not determine neutralizing antibody (NAb) development in individual patients. Unexpectedly, the patients of subgroup B exhibited a good response to IFN β , despite its highest clinical activity. A recent study showed that IFN β res-

ponders are characterized by higher relapse rates during the year prior to initiation of IFN β treatment (Waubant et al., 2003), supporting our observations. In contrast, the patients of C with the poor response to IFN β showed a trend for great increase in the number of MRI lesions during IFN β treatment. A different study indicated that the number of on-treatment new T2 MRI lesions correlates with poor response to IFN β -1a (Rudick et al., 2004), being consistent with our study. Although the subgroup D did not include any IFN β responders, the number of MRI lesions did not increase much during the treatment, suggesting that this subgroup undergoes a neurodegenerative process independent of active inflammation (Steinman, 2001).

A number of differentially expressed genes between MS and CN include those having IFN-responsive elements in the promoter regions: p300 (EP300) and IFN α receptor 1 (IFNAR1) in the class #1, CXCL10, ATP-binding cassette (ABC) subfamily E member 1 (ABCE1 or RNS4I) (Bisbal et al., 1995), IFN γ -inducible protein 16 (IFI16), and STAT1 in the class #2, myxovirus resistance 2 (MX2) in the class #3, IFN-regulatory factors IRF9 and IRF2 in the class #4, and IFN α -16 (IFNA16) and CXCL9 (MIG) in the class #5. These observations suggest that T cells of MS patients have a constitutive defect in regulation of IFN signaling. We previously showed that CXCL9 expression was suppressed in T cells of 13 RRMS patients by a long-term treatment with IFN β (Koike et al., 2003), suggesting that IFN β produces a beneficial effect on MS by correcting the preexisting disturbance in the IFN-signaling pathway.

Finally, T-cell gene expression profiling identified an aberrant expression of key regulators for drug metabolism, whose role has not previously been proposed in MS (Nguyen et al., 2000). A panel of cytochrome P450 (CYP) family, which regulates Ca²⁺ influx in activated T cells (Aussel et al., 1994), were overexpressed in MS T cells. On the other hand, a wide range of ABC transporters in the classes #2, #3, and #4 were downregulated in MS T cells (see Supplementary Table 1 for all datasets). The ABC transporter superfamily regulates the transport of amino acids, ions, sugars, lipids and drugs across the cell membrane by consuming the energy derived from ATP hydrolysis. The downregulated genes in MS include ABCB1 and ABCG2 expressed on brain endothelial cells, which act as a main transporter in the blood–brain barrier and determine bioavailability of corticosteroids and mitoxantrone in the brain (Zhang et al., 2003). The clinicopathological relevance of opposing changes in CYP family enzymes and ABC transporters to MS remains to be further investigated.

In conclusion, T-cell gene expression profiling is highly valuable to identify distinct subgroups of MS associated with differential disease activity and therapeutic response to IFN β . This approach could be applied for designing tailor-made treatment of MS.

Acknowledgements

This work was supported by grants from Research on Intractable Diseases, and Research on Psychiatric and Neurological Diseases and Mental Health, from the Ministry of Health, Labour and Welfare of Japan, Research on Health Sciences Focusing on Drug Innovation, the Japan Health Sciences Foundation (KH21101), and the Program for Promotion of Fundamental Studies in Health Sciences of the National Institute of Biomedical Innovation (NIBIO), Japan. The authors thank Drs. Masaaki Niino and Hikoaki Fukaura, Department of Neurology, Hokkaido University Graduate School of Medicine, and Dr. Masanaga Yamawaki, Department of Neurology, Tokyo Medical and Dental University for their invaluable help.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.jneuroim.2006.02.004.

References

- Achiron, A., Gurevich, M., Friedman, N., Kaminski, N., Mandel, M., 2004. Blood transcriptional signatures of multiple sclerosis: unique gene expression of disease activity. *Ann. Neurol.* 55, 410–417.
- Alizadeh, A.A., Eisen, M.B., Davis, R.E., Ma, C., Lossos, J.S., Rosenwald, A., Boldrick, J.C., Sabet, H., Tran, T., Yu, X., Powell, J.L., Yang, L., Marti, G.E., Moore, T., Hudson, J. Jr., Lu, L., Lewis, D.B., Tibshirani, R., Sherlock, G., Chan, W.C., Greiner, T.C., Weisenburger, D.D., Armitage, J.O., Warnke, R., Levy, R., Wilson, W., Grever, M.R., Byrd, J.C., Botstein, D., Brown, P.O., Staudt, L.M., 2000. Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling. *Nature* 403, 503–511.
- Aussel, C., Breittmayer, J.P., Ticchioni, M., Pelassy, C., Bernard, A., 1994. Regulation of T cell activation by cytochrome P450 inhibitors. *Cell. Immunol.* 155, 436–445.
- Balashov, K.E., Rottman, J.B., Weiner, H.L., Hancock, W.W., 1999. CCR5⁺ and CXCR3⁺ T cells are increased in multiple sclerosis and their ligands MIP-1 α and IP-10 are expressed in demyelinating brain lesions. *Proc. Natl. Acad. Sci. U. S. A.* 96, 6873–6878.
- Baldi, P., Long, A.D., 2001. A Bayesian framework for the analysis of microarray expression data: regularized t-test and statistical inferences of gene changes. *Bioinformatics* 17, 509–519.
- Bisbal, C., Marinand, C., Silhol, M., Lebleu, B., Salehzada, T., 1995. Cloning and characterization of a RNase I inhibitor. A new component of the interferon-regulated 2-5A pathway. *J. Biol. Chem.* 270, 13308–13317.
- Compston, A., Coles, A., 2002. Multiple sclerosis. *Lancet* 359, 1221–1231.
- Graumann, U., Reynolds, R., Steck, A.J., Schaeren-Wiemers, N., 2003. Molecular changes in normal appearing white matter in multiple sclerosis are characteristic of neuroprotective mechanisms against hypoxic insult. *Brain Pathol.* 13, 554–573.
- Gveric, D., Cuzner, M.L., Newcombe, J., 1999. Insulin-like growth factors and binding proteins in multiple sclerosis plaques. *Neuropathol. Appl. Neurobiol.* 25, 215–225.
- Hulshof, S., Montagne, L., De Groot, C.J., Van Der Valk, P., 2002. Cellular localization and expression patterns of interleukin-10.

- interleukin-4, and their receptors in multiple sclerosis lesions. *Glia* 38, 24–35.
- Jacobs, L.D., Cookfair, D.L., Rudick, R.A., Herndon, R.M., Richert, J.R., Salazar, A.M., Fischer, J.S., Goodkin, D.E., Granger, C.V., Simon, J.H., Alan, J.J., Bartoszak, D.M., Bourdette, D.N., Brainan, J., Brownscheidle, C.M., Coats, M.E., Cohan, S.L., Dougherty, D.S., Kinkel, R.P., Mass, M.K., Munschauer, F.E. III, Priore, R.L., Pullicino, P.M., Scherolman, B.J., Weinstock-Guttman, B., Whitham, R.H., 1996. Intramuscular interferon beta-1a for disease progression in relapsing multiple sclerosis. The Multiple Sclerosis Collaborative Research Group (MSCRG). *Ann. Neurol.* 39, 285–294.
- Jacobs, L.D., Beck, R.W., Simon, J.H., Kinkel, R.P., Brownscheidle, C.M., Murray, T.J., Simonian, N.A., Slator, P.J., Sandrock, A.W., 2000. Intramuscular interferon beta-1a therapy initiated during a first demyelinating event in multiple sclerosis. CHAMPS study group. *N. Engl. J. Med.* 343, 898–904.
- Koike, F., Satoh, J., Miyake, S., Yamamoto, T., Kawai, M., Kikuchi, S., Nomura, K., Yokoyama, K., Ota, K., Kanda, T., Fukazawa, T., Yamamura, T., 2003. Microarray analysis identifies interferon β -regulated genes in multiple sclerosis. *J. Neuroimmunol.* 139, 109–118.
- Lammi, J., Hupponen, J., Aarnisalo, P., 2004. Regulation of the osteopontin gene by the orphan nuclear receptor NURR1 in osteoblasts. *Mol. Endocrinol.* 18, 1546–1557.
- Lock, C., Iemmans, G., Pedotti, R., Brendolan, A., Schadt, E., Garen, H., Langer-Gould, A., Strober, S., Cannella, B., Allard, J., Klonowski, P., Austin, A., Lad, N., Kaminski, N., Galli, S.J., Oksenberg, J.R., Raine, C.S., Heller, R., Steinman, L., 2002. Gene-microarray analysis of multiple sclerosis lesions yields new targets validated in autoimmune encephalomyelitis. *Nat. Med.* 8, 500–508.
- Lucchinetti, C., Brück, W., Parisi, J., Scheithauer, B., Rodriguez, M., Lassmann, H., 2000. Heterogeneity of multiple sclerosis lesions: implications for the pathogenesis of demyelination. *Ann. Neurol.* 47, 707–717.
- Maeda, Y., Solanky, M., Menonna, J., Chapin, J., Li, W., Dowling, P., 2001. Platelet-derived growth factor- α receptor-positive oligodendroglia are frequent in multiple sclerosis lesions. *Ann. Neurol.* 49, 776–785.
- Mandel, M., Gurevich, M., Pazuner, R., Kaminski, N., Achiron, A., 2004. Autoimmunity gene expression portrait: specific signature that intersects or differentiates between multiple sclerosis and systemic lupus erythematosus. *Clin. Exp. Immunol.* 138, 164–170.
- McDonald, W.I., Compston, A., Edan, G., Goodkin, D., Hartung, H.P., Lublin, F.D., McFarland, H.F., Paty, D.W., Polman, C.H., Reingold, S.C., Sandberg-Wollheim, M., Sibley, W., Thompson, A., van den Noort, S., Weinshenker, B.Y., Wolinsky, J.S., 2001. Recommended diagnostic criteria for multiple sclerosis: guidelines from the international panel on the diagnosis of multiple sclerosis. *Ann. Neurol.* 50, 121–127.
- Nguyen, L.T., Ramanathan, M., Weinstock-Guttman, B., Dole, K., Miller, C., Planter, M., Patrick, K., Brownscheidle, C., Jacobs, L.D., 2000. Detection of cytochrome P450 and other drug-metabolizing enzyme mRNAs in peripheral blood mononuclear cells using DNA arrays. *Drug Metab. Dispos.* 28, 987–993.
- Okabe, T., Takayanagi, R., Iwasaki, K., Haji, M., Nawata, H., Watanabe, T., 1995. cDNA cloning of a NGF1-B/nur77-related transcription factor from an apoptotic human T cell line. *J. Immunol.* 154, 3871–3879.
- Odenakker, G., Nelissen, I., Van Damme, J., 2003. Functional roles and therapeutic targeting of gelatinase B and chemokines in multiple sclerosis. *Lancet Neurol.* 2, 747–756.
- Panitch, H.S., Hirsch, R.L., Schindler, J., Johnson, K.P., 1987. Treatment of multiple sclerosis with gamma interferon: exacerbations associated with activation of the immune function. *Neurology* 37, 1097–1102.
- Peress, N.S., Perillo, E., Seidman, R.J., 1996. Glial transforming growth factor (TGF)- β isotypes in multiple sclerosis: differential glial expression of TGF- β 1, 2 and 3 isotypes in multiple sclerosis. *J. Neuroimmunol.* 71, 115–123.
- Rudick, R.A., Lee, J.-C., Simon, J., Ransohoff, R.M., Fisher, E., 2004. Defining interferon β response status in multiple sclerosis patients. *Ann. Neurol.* 56, 548–555.
- Saida, T., Tashiro, K., Itoyama, Y., Sato, T., Ohashi, Y., Zhao, Z., the Interferon Beta-1b Multiple Sclerosis Study Group of Japan, 2005. Interferon beta-1b is effective in Japanese RRMS patients. A randomized, multicenter study. *Neurology* 64, 621–630.
- Satoh, J., Nakanishi, M., Koike, F., Miyake, S., Yamamoto, T., Kawai, M., Kikuchi, S., Nomura, K., Yokoyama, K., Ota, K., Kanda, T., Fukazawa, T., Yamamura, T., 2005. Microarray analysis identifies an aberrant expression of apoptosis and DNA damage-regulatory genes in multiple sclerosis. *Neurobiol. Dis.* 18, 537–550.
- Staudt, L.M., 2001. Gene expression physiology and pathophysiology of the immune system. *Trends Immunol.* 22, 35–40.
- Steinman, L., 2001. Multiple sclerosis: a two-stage disease. *Nature Immunol.* 2, 762–764.
- Steinman, L., Zamvil, S., 2003. Transcriptional analysis of targets in multiple sclerosis. *Nat. Rev., Immunol.* 3, 483–492.
- Stürzbecher, S., Wandinger, K.P., Rosenwald, A., Sathymoorthy, M., Tzou, A., Mattar, P., Frank, J.A., Staudt, L., Martin, R., McFarland, H.F., 2003. Expression profiling identifies responder and non-responder phenotypes to interferon- β in multiple sclerosis. *Brain* 126, 1219–1429.
- Tajouri, L., Mellick, A.S., Ashton, K.J., Tannenbaum, A.E., Nagra, R.M., Tourtellotte, W.W., Griffiths, L.R., 2003. Quantitative and qualitative changes in gene expression patterns characterize the activity of plaques in multiple sclerosis. *Mol. Brain Res.* 119, 170–183.
- The IFNB Multiple Sclerosis Study Group, 1993. Interferon beta-1b is effective in relapsing–remitting multiple sclerosis. I. Clinical results of a multicenter, randomized, double-blind, placebo-controlled trial. *Neurology* 43, 655–661.
- van de Vijver, M.J., He, Y.D., van't Veer, L.J., Dai, H., Hart, A.A., Voskuil, D.W., Schreiber, G.J., Peterse, J.L., Roberts, C., Marton, M.J., Parrish, M., Atsma, D., Witteveen, A., Glas, A., Delahaye, L., van der Velde, T., Bartelink, H., Rodenhuis, S., Rutgers, E.T., Friend, S.H., Bernards, R., 2000. A gene-expression signature as a predictor of survival in breast cancer. *N. Engl. J. Med.* 347, 1999–2009.
- Waubant, E., Vukusic, S., Gignoux, L., Dubief, F.D., Achiti, I., Blanc, S., Renoux, C., Confavreux, C., 2003. Clinical characteristics of responders to interferon therapy for relapsing MS. *Neurology* 61, 184–189.
- Windhagen, A., Newcombe, J., Dangond, F., Strand, C., Woodroose, M.N., Cuzner, M.L., Hafler, D.A., 1995. Expression of costimulatory molecules B7-1 (CD80), B7-2 (CD86), and interleukin 12 cytokine in multiple sclerosis lesions. *J. Exp. Med.* 182, 1985–1996.
- Zhou, T., Cheng, J., Yang, P., Wang, Z., Liu, C., Su, X., Blüthmann, H., Mountz, J., 1996. Inhibition of Nur77/Num1 leads to inefficient clonal deletion of self-reactive T cells. *J. Exp. Med.* 183, 1879–1892.
- Zhang, W., Mojsilovic-Petrovic, J., Andrade, M.F., Zhang, H., Ball, M., Staninovic, D.B., 2003. The expression and functional characterization of ABCG2 in brain endothelial cells and vessels. *FASEB J.* 17, 2085–2087.

Research article

Open Access

Microarray analysis identifies a set of CXCR3 and CCR2 ligand chemokines as early IFN β -responsive genes in peripheral blood lymphocytes *in vitro*: an implication for IFN β -related adverse effects in multiple sclerosis

Jun-ichi Satoh*^{1,2}, Yusuke Nanri², Hiroko Tabunoki¹ and Takashi Yamamura²

Address: ¹Department of Bioinformatics, Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo 204-8588, Japan and ²Department of Immunology, National Institute of Neuroscience, NCNP, 4-1-1 Ogawahigashi, Kodaira, Tokyo 187-8502, Japan

Email: Jun-ichi Satoh* - satoj@ncnp.go.jp; Yusuke Nanri - nanriyu@cc.saga-u.ac.jp; Hiroko Tabunoki - tabunoki@my-pharm.ac.jp; Takashi Yamamura - yamamura@ncnp.go.jp

* Corresponding author

Published: 19 May 2006

Received: 20 March 2006

BMC Neurology 2006, 6:18 doi:10.1186/1471-2377-6-18

Accepted: 19 May 2006

This article is available from: <http://www.biomedcentral.com/1471-2377/6/18>

© 2006 Satoh et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: A substantial proportion of multiple sclerosis (MS) patients discontinue interferon-beta (IFN β) treatment due to various adverse effects, most of which emerge at the early phase after initiation of the treatment and then diminish with time. At present, the molecular mechanism underlying IFN β -related adverse effects remains largely unknown. The aim of this study is to identify a comprehensive list of early IFN β -responsive genes (IRGs) in peripheral blood mononuclear cells (PBMC) that may play a key role in induction of adverse effects.

Methods: Total RNA of PBMC exposed to 50 ng/ml recombinant human IFN β for 3 to 24 hours *in vitro* was processed for cDNA microarray analysis, followed by quantitative real-time RT-PCR analysis.

Results: Among 1,258 genes on the array, IFN β elevated the expression of 107 and 87 genes, while it reduced the expression of 22 and 23 genes at 3 and 24 hours, respectively. Upregulated IRGs were categorized into conventional IFN-response markers, components of IFN-signaling pathways, chemokines, cytokines, growth factors, and their receptors, regulators of apoptosis, DNA damage, and cell cycle, heat shock proteins, and costimulatory and adhesion molecules. IFN β markedly upregulated CXCR3 ligand chemokines (SCYB11, SCYB10 and SCYB9) chiefly active on effector T helper type 1 (Th1) T cells, and CCR2 ligand chemokines (SCYA8 and SCYA2) effective on monocytes, whereas it downregulated CXCR2 ligand chemokines (SCYB2, SCYB1 and IL8) primarily active on neutrophils.

Conclusion: IFN β immediately induces a burst of gene expression of proinflammatory chemokines *in vitro* that have potential relevance to IFN β -related early adverse effects in MS patients *in vivo*.

Background

Multiple sclerosis (MS) is an inflammatory demyelinating disease of the central nervous system (CNS) white matter mediated by an autoimmune process, whose development is triggered by a complex interplay of both genetic and environmental factors [1]. Administration of interferon-gamma (IFN γ) induced acute relapses, along with activation of the systemic immune response [2], suggesting that T-lymphocytes producing proinflammatory T helper type 1 (Th1) cytokines play a pivotal role in the immunopathogenesis of MS. In contrast, interferon-beta (IFN β) significantly reduces the frequency of clinical exacerbations and delays the progression of disability in relapsing-remitting MS (RRMS), accompanied by a reduction in the number of new brain lesions on MRI [3,4]. Furthermore, an early initiation of IFN β delays the conversion to clinically definite MS in the patients who experienced a first demyelinating event [5]. Although the precise mechanism underlying therapeutic effects of IFN β on MS remains to be fully elucidated, previous studies proposed several possibilities, including the inhibition of Th1 cell development [6], induction of Th2 immune deviation [7], restoration of function of the disrupted blood-brain barrier [8], and downregulation of IFN γ -induced expression of class II major histocompatibility complex (MHC) molecules [9].

Although clinical benefits of IFN β in MS are meaningful, approximately one-third of the patients receiving IFN β therapy suffered from a higher or identical annual relapse rate on treatment [10]. New lesion formation on MRI during the treatment correlates with poor response to IFN β [11]. Furthermore, a substantial proportion of the patients discontinued IFN β treatment because of various adverse effects, including skin reactions, flu-like symptoms, leukocytopenia, liver dysfunction, depression and amenorrhea [12]. The molecular mechanisms accounting for IFN β -related adverse effects remain unknown, although most of these emerge at the early phase after initiation of the treatment, and then diminish with time [12]. At present, no biologically relevant markers capable of predicting either therapeutic or detrimental responses of IFN β in MS are available [13].

DNA microarray technology is a novel approach that allows us to systematically monitor the expression of a large number of genes. It has given new insights into the complexity of molecular interactions promoting the autoimmune process in MS [14]. IFN β induces a complex pattern of gene regulation in peripheral blood mononuclear cells (PBMC) of MS [15]. Recently, we studied the gene expression profile of CD3 $^+$ T cells isolated from PBMC of 13 Japanese MS patients before and after treatment with IFN β -1b by analyzing a custom cDNA microarray containing a set of well-annotated, immunologically

relevant genes. IFN β upregulates the expression of 7 IFN β -responsive genes (IRGs) during the treatment [16]. A following study performed on RRMS patients receiving IFN β -1a supported our observations [17]. More recently, we found that the vast majority of genes expressed in CD3 $^+$ T cells differentially between 72 untreated MS patients and 22 healthy controls are categorized into apoptosis regulators [18]. Regarding the gene expression profile of IFN β responders in MS, baseline levels of IL-12p35 mRNA are significantly lower in the responders [19]. Downregulation of IL-8 expression in PBMC during IFN β treatment distinguishes the responders from nonresponders in RRMS [20]. IFN β responders differ from nonresponders in the kinetics of expression of IRGs at 3 and 6 months after starting the treatment [21]. A three-dimensional model of gene triplets detected by RT-PCR analysis predicts IFN β response status in RRMS [22]. However, all of previous observations do not clearly illustrate the molecular basis of complex biological effects of IFN β in MS. Furthermore, no databases of immediate early IRGs in PBMC are currently available.

The present study is designed to identify a comprehensive list of immediate early IRGs in PBMC with potential relevance to IFN β -related early adverse effects in MS.

Methods

Treatment of peripheral blood lymphocytes with IFN β

PBMC were isolated from heparinized blood by centrifugation on a Ficoll density gradient. PBMC were suspended at 5×10^6 cells/ml in RPMI 1640 medium containing 10% fetal bovine serum, 2 mM L-glutamine, 55 μ M 2-mercaptoethanol, 100 U/ml penicillin, and 100 μ g/ml streptomycin (culture medium). The cells were incubated in a 5%CO $_2$ /95% air incubator at 37°C for 3 hours to characterize the immediate response or for 24 hours to detect the early response, in the culture medium with or without inclusion of 50 ng/ml recombinant human IFN β (a specific activity of = 2×10^7 units/mg, PeptoTech, London, UK), 50 ng/ml recombinant human IFN γ (a specific activity of = 2×10^7 units/mg, PeptoTech), 50 ng/ml recombinant human TNF α (a specific activity of = 2×10^7 units/mg, PeptoTech), or 50 ng/ml recombinant human IL-1 β (a specific activity of = 1×10^7 units/mg, PeptoTech). They were then processed for RNA preparation as described previously [16,18,21]. Written informed consents were obtained from the subjects involved in the present study according to the form approved by the Ethics Committee of National Center of Neurology and Psychiatry (NCNP), Tokyo, Japan. The samples of the subjects #1, #2 and #4 were processed for both microarray and real-time RT-PCR analysis, while those of the subject #3 were studied only by real-time RT-PCR analysis.

Table 1: Primers utilized for real-time RT-PCR analysis

Genes	GenBank accession No.	Sense primers	Antisense primers	PCR product (bp)
ISG15 (GIP2)	NM_005101 (M13755)	5'aagcccctgacacccgtgttc3'	5'tgatcctgctcggatgctggg3'	102
SCYB10 (CXCL10, IP-10)	NM_001565 (X02530)	5'tcgatgcagtgctccaaggatgg3'	5'ccttctacaggagtagtagcagc3'	162
SCYA8 (CCL8, MCP2)	NM_005623 (Y10802)	5'tctgtctgaccccaaggagat3'	5'taatgtcacactgcacctggggga3'	164
SCYA2 (CCL2, MCP1)	NM_002982 (S71513)	5'ctagcttccccagacacccctgt3'	5'ccagggtagaactgtggtcaag3'	197
SCYB2 (CXCL2, GRO2)	NM_002089	5'cccgcacgcccattggttaagaa3'	5'tcttctgctcctgtaaggcagggc3'	131
FOS	NM_005252	5'gagctggctcattacagaggag3'	5'ggacttgagccacacatggatgc3'	140
RGS14	NM_006480	5'tgacagctaccaacagtcaggga3'	5'agggattgggggtgagctgttga3'	222
G3PDH	NM_002046	5'ccatgttcgtcaggggtgaacca3'	5'gccagtagaggcaggatgatgttc3'	251

Abbreviations: ISG15, interferon-stimulated gene 15; SCYB10, small inducible cytokine subfamily B, member 10; SCYA8, small inducible cytokine subfamily A, member 8; SCYA2, small inducible cytokine subfamily A, member 2; SCYB2, small inducible cytokine subfamily B, member 2; FOS, cellular oncogene *c-fos*; RGS14, regulator of G-protein signaling 14; and G3PDH, glyceraldehyde-3-phosphate dehydrogenase

Quantitative real-time RT-PCR analysis

DNase-treated total RNA was processed for cDNA synthesis using oligo(dT)₁₂₋₁₈ primers and SuperScript II reverse transcriptase (Invitrogen, Carlsbad, CA). cDNA was amplified by PCR in LightCycler ST300 (Roche Diagnostics, Tokyo, Japan) using SYBR Green I dye and the primer sets listed in Table 1. To calibrate the concentration of mRNA levels in test cDNA samples, serially-diluted purified PCR products generated by conventional RT-PCR (a 10-fold dilution from 1 pg/ml to 0.0001 pg/ml) were amplified in parallel. The levels of expression of target genes were standardized against those of the glyceraldehyde-3-phosphate dehydrogenase (G3PDH) gene detected in the identical cDNA samples. The assays were performed in triplicate measurements of the same sample and the results were expressed as the average with standard error.

cDNA microarray analysis

The present study utilized a custom microarray containing duplicate spots of 1,258 cDNA immobilized on a poly-L-lysine-coated slide glass (Hitachi Life Science, Kawagoe, Saitama, Japan) [16,18,21]. They were prepared by PCR of well-annotated genes, selected from cytokines, growth factors and their receptors, apoptosis regulators, oncogenes, transcription factors, cell cycle regulators and housekeeping genes. The complete gene list is shown in Additional file 1. Five µg of purified RNA was *in vitro* amplified and antisense RNA (aRNA) was processed for microarray analysis. aRNA derived from IFNβ-treated PBMC was labeled with a fluorescent dye Cy5, while aRNA of untreated PBMC was labeled with Cy3 by reverse transcriptase reaction. The arrays were hybridized at 62°C for 17 hours in the hybridization buffer containing equal amounts of Cy3- or Cy5-labeled cDNA, and they were then scanned at two different photomultiplier tube (PMT) gains by the

ScanArray 5000 scanner (GSI Lumonics, Boston, MA). The data were analyzed by using the QuantArray software (GSI Lumonics). The fluorescence intensities (FI) of individual spots were quantified following global normalization between Cy3 and Cy5 signals. The average of FI of duplicate spots was calculated, then the ratio of FI of Cy5 signal versus FI of Cy3 signal exceeding 2.0 was defined as significant upregulation, whereas the ratio smaller than 0.5 was considered as substantial downregulation. The impact of inter-experiment variability was validated by analyzing a scatter plot (see Additional file 2).

The IFN-regulated expression of the genes we identified was verified by computerized search of PubMed database and IFN Stimulated Gene (ISG) database [23].

Results

Microarray analysis identified immediate early IFNβ-responsive genes in PBMC

Among 1,258 genes on the array, IFNβ treatment for 3 hours elevated the expression of 107 genes in PBMC isolated from a 46 year-old healthy man (the subject #1), while it reduced the expression of 22 genes (see Additional files 3 and 4 for the complete list). IFNβ treatment for 24 hours upregulated 87 genes and downregulated 23 genes (see Additional files 5 and 6 for the complete list). Sixty-nine genes were upregulated at both 3 and 24 hours, while only two genes such as FOS and IL1A were downregulated at both. The IRGs upregulated at both time points contained 11 *in vivo* IRGs reported previously by us [16], including IFIT1 (IFI56), ISG15 (GIP2), IFIT4 (IFI60), IFI27, G1P3 (IFI6-16), IRF7, ABCB2 (TAP1), ATF3, IFITM1 (IFI17), SULT1C1, and TNFAIP6, whose expression was elevated in T cells and non-T cells *ex vivo*, isolated from 13 RRMS patients during IFNβ treatment for 3 to 6 months. Top 20 most significant genes, either

Table 2: Top 20 upregulated genes in PBMC following exposure to IFN β

No.	Cy5/Cy3 ratio	Symbol	GenBank	Gene name
at 3 hours				
1	188.50	<u>IFIT1</u>	<u>X03557</u>	interferon-induced protein with tetratricopeptide repeats 1 (IFI56)
2	149.82	SCYB11	<u>AF030514</u>	chemokine (C-X-C motif) ligand 11 (CXCL11, IP-9, I-TAC)
3	81.81	<u>ISG15</u>	<u>M13755</u>	interferon-stimulated gene ISG15 (GIP2)
4	67.84	<u>IFIT4</u>	<u>AF083470</u>	interferon-induced protein with tetratricopeptide repeats 4 (IFI60)
5	66.38	MX1	<u>M30817</u>	myxovirus resistance protein 1 (MXA)
6	64.43	SCYB10	<u>X02530</u>	chemokine (C-X-C motif) ligand 10 (CXCL10, IP-10)
7	55.01	SCYA8	<u>Y10802</u>	chemokine (C-C motif) ligand 8 (CCL8, MCP2)
8	36.88	SCYB9	<u>X72755</u>	chemokine (C-X-C motif) ligand 9 (CXCL9, MIG)
9	35.14	TNFSF10	<u>U37518</u>	tumor necrosis factor superfamily, member 10 (TRAIL)
10	32.86	MX2	<u>M30818</u>	myxovirus resistance protein 2 (MXB)
11	21.99	<u>IFI27</u>	<u>X67325</u>	interferon alpha-inducible protein 27
12	21.98	GIP3	<u>X02492</u>	interferon alpha-inducible protein (IFI6-16), isoform a
13	19.26	ISG20	<u>U88964</u>	interferon-stimulated gene ISG20
14	19.12	STAT1	<u>M97935</u>	signal transducer and activator of transcription 1, isoform alpha
15	13.71	<u>IRE7</u>	<u>U53830</u>	interferon-regulatory factor 7, isoform alpha
16	13.44	CCNA1	<u>U66838</u>	cyclin A1
17	12.72	IL6	<u>M14584</u>	interleukin 6
18	10.86	IL1RN	<u>X53296</u>	interleukin 1 receptor antagonist, isoform 1
19	10.47	SCYA2	<u>S71513</u>	chemokine (C-C motif) ligand 2 (CCL2, MCP1)
20	9.91	STAT2	<u>M97934</u>	signal transducer and activator of transcription 2
at 24 hours				
1	193.19	SCYA8	<u>Y10802</u>	chemokine (C-C motif) ligand 8 (CCL8, MCP2)
2	124.77	<u>IFIT1</u>	<u>X03557</u>	interferon-induced protein with tetratricopeptide repeats 1 (IFI56)
3	107.21	<u>IFI27</u>	<u>X67325</u>	interferon alpha-inducible protein 27
4	70.51	<u>ISG15</u>	<u>M13755</u>	interferon-stimulated gene ISG15 (GIP2)
5	53.22	SCYB11	<u>AF030514</u>	chemokine (C-X-C motif) ligand 11 (CXCL11, IP-9, I-TAC)
6	38.07	MX1	<u>M30817</u>	myxovirus resistance protein 1 (MXA)
7	37.95	<u>IFIT4</u>	<u>AF083470</u>	interferon-induced protein with tetratricopeptide repeats 4 (IFI60)
8	31.56	SCYB10	<u>X02530</u>	chemokine (C-X-C motif) ligand 10 (CXCL10, IP-10)
9	23.65	<u>GIP3</u>	<u>X02492</u>	interferon alpha-inducible protein (IFI6-16), isoform a
10	23.10	TNFSF10	<u>U37518</u>	tumor necrosis factor superfamily, member 10 (TRAIL)
11	21.31	MX2	<u>M30818</u>	myxovirus resistance protein 2 (MXB)
12	20.68	PDGFRL	<u>D37965</u>	platelet-derived growth factor receptor-like protein
13	15.95	CD80	<u>M27533</u>	CD80 antigen (B7-1)
14	13.99	<u>IRE7</u>	<u>U53830</u>	interferon-regulatory factor 7, isoform alpha
15	13.93	IL1RN	<u>X53296</u>	interleukin 1 receptor antagonist, isoform 1
16	13.31	ISG20	<u>U88964</u>	interferon-stimulated gene ISG20
17	11.81	<u>IFITM1</u>	<u>J04164</u>	interferon induced transmembrane protein 1 (IFI17)
18	8.67	SCYA2	<u>S71513</u>	chemokine (C-C motif) ligand 2 (CCL2, MCP1)
19	8.54	IL6	<u>M14584</u>	interleukin 6
20	8.51	IL8RB	<u>L19593</u>	interleukin 8 receptor beta (CXCR2)

Top 20 upregulated genes in PBMC of the subject #1 (a 46 year-old healthy man) by exposure to 50 ng/ml recombinant human IFN β for 3 or 24 hours are listed with Cy5/Cy3 signal intensity ratio, gene symbol, GenBank accession number, and gene name. In vivo IRGs in T cells and non-T cells of RRMS patients reported previously (Ref. 16) are underlined.

upregulated or downregulated, are listed in Table 2 and Table 3, respectively. All of top 20 upregulated genes were found as known IRGs identified in various cell types by searching through PubMed and ISG databases.

The upregulated IRGs in the complete lists (see Additional files 3 and 5) were classified into several functional categories following; (i) conventional IFN-response markers (n = 12), (ii) components of classical and Toll-like receptor (TLR)-dependent IFN-signaling pathways (n = 12),

(iii) chemokines and their receptors (n = 11), (iv) cytokines, growth factors and their receptors (n = 17), (v) apoptosis, DNA damage, and cell cycle regulators (n = 29), (vi) heat shock proteins (n = 9), and (vii) costimulatory and adhesion molecules (n = 7) (Table 4). The chemokine and chemokine receptor group included both CXC and CC chemokines and their receptors, such as SCYB11 (CXCL11, I-TAC), SCYB10 (CXCL10, IP-10), SCYA8 (CCL8, MCP2), SCYB9 (CXCL9, MIG), SCYA2 (CCL2, MCP1), CCR5, SCYA4 (CCL4, MIP1B), IL8RB (CXCR2),

Table 3: Top 20 downregulated genes in PBMC following exposure to IFN β

No.	Cy5/Cy3 ratio	Symbol	GenBank	Gene name
at 3 hours				
1	0.29	RGS14	NM_006480	regulator of G-protein signaling 14
2	0.30	FOS	NM_005252	cellular oncogene c-fos
3	0.31	SCYB2	NM_002089	chemokine (C-X-C motif) ligand 2 (GRO2, MIP2A)
4	0.41	PPARG2	U63415	peroxisome proliferative activated receptor gamma, isoform 2
5	0.41	TNFSF11	NM_003701	tumor necrosis factor ligand superfamily, member 11 (RANKL), isoform 1
6	0.42	ABCC4	NM_005845	ATP-binding cassette, subfamily C, member 4 (MRP4)
7	0.42	MERTK	NM_006343	c-mer proto-oncogene tyrosine kinase
8	0.42	TGFBR3	NM_003243	transforming growth factor beta receptor III (betaglycan)
9	0.44	ATP2B1	J04027	ATPase, Ca ⁺⁺ transporting, plasma membrane 1, isoform 1b
10	0.45	SCYB1	NM_001511	chemokine (C-X-C motif) ligand 1 (GRO1, MGSA)
11	0.45	IL8	NM_000584	interleukin 8 (SCYB8, CXCL8)
12	0.45	TRAF5	NM_004619	TNF receptor-associated factor 5, variant 1
13	0.47	CD3G	NM_000073	CD3G antigen, gamma polypeptide
14	0.47	IL1A	NM_000575	interleukin 1 alpha
15	0.47	TCFL5	NM_006602	transcription factor-like 5
16	0.48	LAT	AF036905	linker for activation of T cells, isoform b
17	0.48	HNMT	U08092	histamine N-methyltransferase, isoform 1
18	0.48	SLC3A1	NM_000341	solute carrier family 3, member 1
19	0.49	MGST1	NM_145764	microsomal glutathione S-transferase 1, variant 1d
20	0.49	TNFSF8	NM_001244	tumor necrosis factor ligand superfamily, member 8 (CD30L)
at 24 hours				
1	0.18	ABCC3	AF083552	ATP-binding cassette, subfamily C, member 3, isoform MRP3
2	0.20	FOS	NM_005252	cellular oncogene c-fos
3	0.30	ALDH2	NM_000690	aldehyde dehydrogenase 2 family
4	0.30	TPST1	NM_003596	tyrosylprotein sulfotransferase 1
5	0.31	CD9	NM_001769	CD9 antigen
6	0.34	NRG1	NM_013962	neuregulin 1, isoform GGF2
7	0.35	TLR5	NM_003268	toll-like receptor 5
8	0.36	PTGS1	NM_000962	prostaglandin-endoperoxide synthase 1 (COX1), isoform 1
9	0.37	MGST2	NM_002413	microsomal glutathione S-transferase 2
10	0.38	ITGAM	NM_000632	integrin, alpha M (CD11B, MAC-1, CR3A)
11	0.38	SCYA24	NM_002991	chemokine (C-C motif) ligand 24 (eotaxin 2)
12	0.38	CYP27A1	NM_000784	cytochrome P450, family 27, subfamily A, polypeptide 1 (CTX)
13	0.39	ALDH1A1	NM_000689	aldehyde dehydrogenase 1 family, member A1
14	0.39	MYCL1	NM_005376	v-myc myelocytomatosis viral oncogene homolog 1 (LMYC)
15	0.41	ATP6AP2	NM_005765	ATPase, H ⁺ transporting, lysosomal accessory protein 2 (ATP6M8-9)
16	0.44	PRKCG	NM_002739	protein kinase C gamma
17	0.45	ATP2B1	J04027	ATPase, Ca ⁺⁺ transporting, plasma membrane 1, isoform 1b
18	0.46	AP1S2	NM_003916	adaptor-related protein complex 1, sigma 2 subunit
19	0.47	NR6A1	NM_001489	nuclear receptor subfamily 6, group A, member 1 (RTR), isoform 2
20	0.47	GCG	NM_173916	glucagon

Top 20 downregulated genes in PBMC of the subject #1 (a 46 year-old healthy man) by exposure to 50 ng/ml recombinant human IFN β for 3 or 24 hours are listed with Cy5/Cy3 signal intensity ratio, gene symbol, GenBank accession number, and gene name.

SCYA3 (CCL3, MIP1A), SCYA19 (CCL19, MIP3B) and SCYA13 (CCL13, MCP4). It is worthy to note that both CXCR3 ligand chemokines (SCYB11, SCYB10 and SCYB9) and CCR2 ligand chemokines (SCYA10 and SCYA2) were clustered in top 20 genes greatly elevated at 3 and 24 hours of IFN β treatment (Table 2). With respect to top 20 downregulated genes, four genes such as SCYB2 (CXCL2, GRO2), SCYB1 (CXCL1, GRO1), IL8 (SCYB8, CXCL8), and SCYA24 (eotaxin-2) were categorized into the chemokine group (Table 3). Among them, SCYB2, SCYB1 and IL8, whose expression was reduced immedi-

ately at 3 hours of IFN β treatment, belong to CXCR2 ligand and chemokines. Although the analysis in the present study was a single microarray for each sample design, the results from two additional subjects, including a 28 year-old healthy man (the subject #2) and a 27 year-old woman with RRMS who was a dropout of IFN β treatment due to induction of frequent severe relapses (the subject #4), verified the observations of immediate early induction of CXCR3 ligand and CCR2 ligand chemokine genes in PBMC by exposure to IFN β , supporting the reproducibility of these results (see Additional file 7).