

causes B-cell lymphomas and myelogenous leukemia in mice [8]. Multiple proline-rich stretches are found in the C-terminal domain of Cbl, enabling it to bind to several src homology 2/3 (SH2/SH3) adaptor proteins, such as Grb2 and Nck [9–11]. In addition, the C-terminus of Cbl contains several tyrosine phosphorylation sites. Tyrosine 700, 731 and 774 have been identified as major phosphorylation sites of Cbl by protein tyrosine kinases (PTKs) [12]. Cbl-b is a distinct Cbl-related gene with similar overall domain structure to Cbl [13]. Cbl is a major target of tyrosine phosphorylation in response to stimulation through a wide variety of cell surface receptors [14]. Cbl-deficient mice manifest tissue hyperplasia and abnormal duct formation in mammary glands, and Zap-70 is hyperphosphorylated in the thymocytes [15,16]. Overexpression of Cbl in mast cells suppresses Syk tyrosine kinase activity [17]. The aims of this study were to confirm and extend roles for Cbl and CD45 and determine a role for Cbl-b and associated proteins in the intracellular cascade of events set off by SDF-1 α /CXCL12 actions in induction of chemotaxis.

Materials and methods

Reagents and antibody

Recombinant human SDF-1 α /CXCL12, anti-phosphotyrosine monoclonal antibody (mAb, 4G10) and anti-Crk-L mAb were purchased from Upstate Biotechnology (Lake Placid, NY). Rabbit anti-Cbl antibody (Ab), anti-Cbl-b Ab, GST agarose, GST-fyn agarose and protein A/G agarose were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). Anti-phosphotyrosine mAb (PY20), anti-Cbl mAb, anti-Nck mAb anti-Zap-70 mAb, anti-Fyb mAb and anti-PLC- γ mAb were obtained from Transduction Laboratories (Lexington, KY). Anti-phospho-Cbl Ab was from Cell Signaling Technology (Beverly, MA). Src kinase inhibitor PP2 was from Calbiochem-Novabiochem Corporation (San Diego, CA). Other reagents were from Sigma (St. Louis, MO).

Cell culture, transfection and infection

Human leukemic T cell line Jurkat, Lck-deficient T cell line J.CaM1.6 and CD45-deficient T cell line J45.01 were obtained from American Type Culture Collection (ATCC, Manassas, VA). The J.CaM1.6 cell line is a derivative mutant of Jurkat, and J45.01 cell line is a CD45-deficient variant of the E6-1 clone of Jurkat. These cell lines were maintained in RPMI1640 supplemented with 10% heat-inactivated fetal bovine serum (FBS, Hyclone, Logan UT) with 1% penicillin/streptomycin in a humidified incubator at 37°C.

Immunoprecipitation and Western blot analysis

Jurkat, J.CaM1.6 and J45.01 cells were factor-starved overnight and treated with 100 ng/ml SDF-1 α /CXCL12 (a pre-established maximal concentration for induction of chemotaxis for Jurkat and other T cells) at the indicated times and washed once with ice-cold phosphate-buffered saline (PBS). Cells were lysed in lysis buffer containing 20 mM Tris-HCl, pH

8.0; 137 mM NaCl; 10% glycerol; 1 mM phenylmethylsulfonyl fluoride (PMSF); 10 μ M EDTA; 10 μ g/ml leupeptin; 100 mM sodium fluoride; 2 mM sodium orthovanadate; 1% NP-40, for 20 min on ice. The lysates were centrifuged at 12,000 rpm for 20 min at 4°C. Protein content of lysates was determined with protein assay kit (Bio-Rad Laboratories, Hercules, CA). Equivalent amounts of protein of cell lysates were boiled with 2 \times SDS sample buffer for 5 min. For immunoprecipitation, cell lysates were incubated at 4°C overnight with the indicated precipitating Ab. Immunoprecipitates were collected using 40 μ l of protein A/G agarose for 2 h at 4°C. After washing four times in lysis buffer, immunocomplexes were eluted and boiled with 5 min in 2 \times sample buffer. Proteins or immunocomplexes were loaded onto polyacrylamide gels (BioWhittaker, Rockland, ME) and then transferred to polyvinylidene difluoride (PVDF) membranes (Millipore, Bedford, MA). The membranes were blocked by 3% skim milk PBS-Tween 20 (PBST) or 1% BSA PBST and probed with the indicated primary antibody at appropriate dilution for 2 h at room temperature (RT) or 4°C overnight. Blots were probed with secondary antibodies conjugated horseradish peroxidase and developed using the




Fig. 1. SDF-1/CXCL12 enhances tyrosine phosphorylation of Cbl and Cbl-b. Jurkat cells were left unstimulated (–) or stimulated with SDF-1 α /CXCL12 (100 ng/ml) for 5 min. Total cell lysates (TCL) were immunoprecipitated with anti-Cbl Ab (A) or anti Cbl-b Ab (B). Immunoprecipitates (IP) were immunoblotted with anti-phosphotyrosine Ab, anti-Cbl Ab (A) or anti Cbl-b Ab (B). WB: Western blot. Results shown in panels A and B are representative of at least 3 complete experiments.




Fig. 2. Src kinase inhibitor PP2 blocks SDF-1/CXCL12 enhanced tyrosine phosphorylation of Cbl and Cbl-b. Jurkat cells were pretreated with src kinase inhibitor PP2 (10 μ M) for 30 min and left unstimulated or stimulated with SDF-1 α /CXCL12 at 100 ng/ml for 5 min. Cell lysates were immunoprecipitated with anti-Cbl Ab and immunoblotted with anti-phosphotyrosine or anti-Cbl Ab (A) or immunoprecipitated with anti-Cbl-b Ab and then immunoblotted with anti-phosphotyrosine or anti-Cbl-b Ab (B). Total cell lysates were incubated with GST or GST-fyn and then immunoblotted with anti-Cbl, anti-phospho-Cbl or anti-Cbl-b Ab (C). Results in panels A–C are one representative each of at least three similar experiments.

enhanced chemiluminescence (ECL, Amersham Pharmacia Biotech, Bucks, UK) system with ECL film according to the manufacturer's specification.

Chemotaxis assay

Chemotaxis assays were performed using a Boyden chamber with a 5 μ m pore size (Coming Incorporated, Coming NY) as described previously [18]. After incubation of the apparatus at 37°C for 2 h in humidified air with 5% CO₂, the filter was removed and the cells were counted by FACScan (Becton Dickinson, Mountain View) for 20 s.

Flow cytometric analysis

Cells were fixed with using Cytofix/Cytoperm kit (BD Pharmingen, San Diego, CA) or 1% paraformaldehyde PBS. Cells were incubated with anti-CXCR4 mAb (R&D systems, Minneapolis, MN) for 30 min at 4°C and then incubated with fluorescein-conjugated secondary antibody. The cells were monitored by flow cytometric analysis.

Electroporation of Jurkat cells

Jurkat cells were transfected by electroporation with siRNA for Cbl and Cbl-b. Cells were electroporated in 400 μ l cultured medium at a density of 2×10^7 /ml in electroporation cuvettes and mixed with 1 nmol siRNA. Cells were electroporated by using the Bio-Rad gene pulser II system at a setting of 0.2 kV




Fig. 3. SDF-1/CXCL12 enhances Cbl and Cbl-b association with Zap-70, Nck, PLC- γ , Crk-L and Fyb in Jurkat cells. Cbl (A) or Cbl-b (B) were immunoprecipitated from cell lysates of unstimulated or SDF-1 α /CXCL12-stimulated cells and analyzed by Western blotting with anti-Zap-70, Nck, PLC- γ , Crk-L and Fyb Ab, or anti-Cbl Ab (A) or anti-Cbl-b Ab (B). Results in panels A and B are for 1 of at least 3 reproducible experiments.

and 950 μ F capacitance. Fifteen minutes after electroporation, cells were diluted in culture medium and incubated at 37°C, 5% CO₂ and 92% humidity. After 48 h, cells were analyzed by chemotaxis assay and immunoblotting.

Results

SDF-1 α /CXCL12 enhances tyrosine phosphorylation of Cbl and Cbl-b

Cbl is tyrosine-phosphorylated in response to stimulation via immune, growth factor, integrin and cytokine receptors [12] and SDF-1/CXCL12 [6]. In order to characterize signaling pathways activated by SDF-1/CXCL12, we used the Jurkat T cell line, which expresses the SDF-1 α /CXCL12 receptor CXCR4. Jurkat T cells were serum-starved and stimulated with 100 ng/ml of SDF-1 α /CXCL12 for the indicated times (Fig. 1). SDF-1 α /CXCL12 enhanced tyrosine phosphorylation of Cbl (Fig. 1A), confirming the results of others [6]. Another Cbl family member Cbl-b was also tyrosine-phosphorylated in response to

SDF-1/CXCL12 (Fig. 1B) extending information to a Cbl family member. Enhanced tyrosine phosphorylation of Cbl and Cbl-b was detected from 2 min to 30 min after treatment of cells with SDF-1/CXCL12. Determination of the loading of equal amounts of protein in all lanes was accomplished by stripping and blotting with Cbl and Cbl-b antibody.

SDF-1/CXCL12-enhanced phosphorylation of Cbl and Cbl-b is regulated by src family kinases

Tyrosine phosphorylation of Cbl appears to be mediated by several protein tyrosine kinases (PTKs), such as those of the src family PTKs [19,20]. Serum-starved Jurkat cells were pre-treated with 10 μ M of the specific src kinase inhibitor PP2 for 30 min and then stimulated with 100 ng/ml SDF-1 α /CXCL12 for 5 min. Pretreatment of cells with PP2 completely or greatly blocked Cbl and Cbl-b phosphorylation (Figs. 2A and B). These results suggest that phosphorylation of Cbl and Cbl-b is regulated by src family kinases. We then assessed if Cbl and Cbl-b bound to the Src family member Fyn. SDF-1/CXCL12




Fig. 4. SDF-1/CXCL12 does not enhance tyrosine phosphorylation of Cbl and Cbl-b in J.CaM1.6 and J45.01 cells. Cell surface expression of CXCR4. Jurkat, J.CaM1.6 and J45.01 cells were fixed by 1% paraformaldehyde PBS and stained with anti-CXCR4 Ab. After staining with a secondary Ab, surface CXCR4 levels were analyzed by FACSscan (A). Tyrosine phosphorylation of Cbl or Cbl-b in J.CaM1.6 (B) or J45.01 (C). Cbl and Cbl-b were immunoprecipitates and analyzed by Western blotting with anti-phosphotyrosine or anti-Cbl or Cbl-b Ab. Results in panels A–C are for 1 of at least 3 reproducible experiments each.




Fig. 5. SDF-1/CXCL12-enhanced MAPK activity is the same in J45.01 and parental Jurkat cells. Jurkat (A) and J45.01 (B) cells were left unstimulated (–) or stimulated with SDF-1 α /CXCL12 (100 ng/ml) for the indicated times. Total cell lysates were immunoblotted with anti-phospho-ERK Ab or anti-ERK Ab. Results in panels A and B are representative of at least 3 complete experiments.

stimulation enhanced the binding of Cbl and Cbl-b to the Src family member Fyn as assessed by GST “pull-down” (Fig. 2C).

SDF-1/CXCL12 enhances the association of tyrosine-phosphorylated Cbl and Cbl-b with Crk-L, Zap-70, Nck, PLC- γ and Fyb

Cbl has been shown to form complexes with many proteins via its various functional domains. Tyrosine kinase binding domains of Cbl bind to several receptor PTKs, including Syk, Zap-70 and Src, and proline-rich regions of Cbl interact with SH3-containing proteins, such as Nck and Crk family adaptor proteins. Fyn binding protein (Fyb), also called SLAP-130 (SLP-76 associated phosphorylation of 130 kDa), includes several proline-rich regions and multiple tyrosine motifs [21]. SDF-1/CXCL12 enhanced the association of tyrosine-phosphorylated Cbl (Fig. 3A) and Cbl-b (Fig. 3B) with Crk-L, Zap-70, PLC- γ , Nck and Fyb.

Enhanced tyrosine phosphorylation of Cbl and Cbl-b in response to SDF-1/CXCL12 is not apparent in J.CaM1.6 and J45.01 cells

We showed above that tyrosine phosphorylation of Cbl and Cbl-b is regulated by src kinase family members. Lck is a member of the src kinase family and is important to T cell function. Lck regulates T cell surface receptors, such as CD2 and CD4 [22,23]. J.CaM1.6 cells are derived from Jurkat cells and are defective in expression of Lck. J45.01 cells are also derived from Jurkat cells but are defective in the phosphatase CD45. Surface expression of CXCR4 was similar between J.

CaM1.6 and J45.01 cells and their parental cell line Jurkat (Fig. 4A). CD45 is a receptor-like protein tyrosine phosphatases (PTPs) expressed on all nucleated hematopoietic cells. CD45 serves as a positive regulator of src family kinases [24]. SDF-1/CXCL12 did not enhance tyrosine phosphorylation of Cbl and Cbl-b in J.CaM1.6 (Fig. 4B) or J45.01 (Fig. 4C) cells. These results demonstrate that Lck and CD45 are involved in regulation of tyrosine phosphorylation of Cbl and Cbl-b in response to SDF-1 α /CXCL12.

MAPK activity is not decreased in J45.01 cells

MAPK is involved in signal transduction events mediating proliferation, differentiation and apoptosis in eukaryotic cells [25]. Extracellular-signal-regulated kinases 1/2 (ERK1/2) are typically stimulated by growth-related stimuli. We recently reported that MAPK activity is not different in J.CaM1.6 cells and parental Jurkat cells, suggesting that Lck is not involved in regulation of MAPK activity in response to SDF-1/CXCL12 [26]. We now evaluated whether or not the tyrosine phosphatase activity of CD45 was involved in SDF-1/CXCL12 effects on MAPK activity. MAPK activity was not decreased in J45.01 cells compared to Jurkat cells (Fig. 5), suggesting that CD45 does not regulate MAPK activity in response to SDF-1/CXCL12.

Chemotaxis response decreased in J45.01 cells

Jurkat cells respond chemotactically to SDF-1 α /CXCL12 [6,18]. We chose to assess the chemotactic response of Jurkat (Parent), J.CaM1.6 (Lck-deficient) and J45.01 (CD45-deficient) cells to a concentration of SDF-1 (200 ng/ml) that is a plateau dose that induces maximum migration of Jurkat cells [18]. J. CaM1.6 cells were previously shown by us to be responsive to SDF-1/CXCL12-induced chemotaxis [26] and were used here as a control. As shown in Fig. 6, the chemotactic response to SDF-1 α /CXCL12 was significantly decreased in J45.01 cells, but not in J.CaM1.6 cells, compared to Jurkat cells. This confirms the results of others [7] that CD45 is involved in cell migration in response to SDF-1 α /CXCL12.




Fig. 6. J45.01 but not J.CaM1.6 cells are reduced in response to the chemotaxis activity of SDF-1/CXCL12 compared to parental Jurkat cells. Jurkat, J.CaM1.6 or J45.01 cells (1×10^5 each) were placed in the upper part of chemotaxis chambers. Cells that migrate to the lower chambers after 2 h were counted using FACScan. These experiments were done four times with similar results. * $P < 0.05$, compared to Jurkat.

Chemotaxis response, but not MAPK activity, decreased after Cbl and Cbl-b siRNA transfection

To determine the effect of Cbl and Cbl-b in SDF-1 α /CXCL12-induced migration, Cbl, Cbl-b or control siRNA was transfected into Jurkat cells by electroporation. Cbl and Cbl-b siRNAs reduced Cbl and Cbl-b protein levels 48 h after transfection (Fig. 7A). At this time, we examined the effect of Cbl and Cbl-b siRNA on chemotactic response to SDF-1 α /CXCL12. The chemotactic response to SDF-1 α /CXCL12 was significantly reduced in cells transfected with siRNA to Cbl or Cbl-b compared to that of control siRNA-transfected cells (Fig. 7B), but MAPK activity was not reduced (Fig. 7C). These results




Fig. 7. Jurkat cells transfected with Cbl or Cbl-b siRNA are reduced in response to the chemotaxis induced by SDF-1/CXCL12 but are not reduced in MAPK activity. Jurkat cells were electroporated with 1 nmol siRNA for Cbl, Cbl-b or with control. (A) Cells were stimulated with SDF-1 α /CXCL12 (100 ng/ml) and immunoblotted 48 h after electroporation. (B) Cells were placed in the upper part of chemotaxis chambers, and cells moving to the lower chamber were counted using FACScan. These experiments were done four times with similar results. * $P < 0.05$, compared to control. (C) Cells were stimulated by SDF-1/CXCL12 and evaluated for phosphorylation of Erk.

suggest that Cbl and Cbl-b are involved, at least in part, in SDF-1/CXCL12-induced chemotaxis of Jurkat cells, but MAPK activity is not influenced by this decrease in Cbl or Cbl-b.

Discussion

SDF-1 α /CXCL12 plays a key role in the regulation of migration and homing of hematopoietic cells. Cell migration is mediated by multiple signaling mechanisms. In the present study, we investigated the phosphorylation of Cbl and Cbl-b and the association of these adaptor molecules with other signaling molecules in T cell lines. We also assessed the functional effects of decreased Cbl and Cbl-b expression, after transfection of siRNA for Cbl and Cbl-b, on SDF-1/CXCL12-induced migration of the Jurkat T cell line. Our results both confirm and extend the findings of others [6] that Cbl family members function as adaptor proteins downstream of CXCR4 in response to SDF-1 α /CXCL12.

Cbl and Cbl-b contain substrate recognition domains that interact with activated Src protein tyrosine kinases. Cbl interacts with the SH3 domains of src kinases, such as Lck, Fyn and Lyn [27]. We demonstrate here that Fyn associates with Cbl and Cbl-b and that Lck also regulates Cbl family phosphorylation in response to SDF-1 α /CXCL12. Our observations that the Src kinase inhibitor PP2 blocks SDF-1 α /CXCL12-induced tyrosine phosphorylation of Cbl and Cbl-b suggest that src kinases are upstream effectors of Cbl and Cbl-b in SDF-1 α /CXCL12 actions.

Cbl is a negative regulator in the immune system of Syk/Zap-70 PTKs [28]. In Cbl-b knockout mice, Zap-70 is hyperphosphorylated in thymocytes and ERK is activated compared to wild type mice [15,16]. Cbl has no known catalytic function, but it contains several domains that are able to interact with a wide variety of substrates. We demonstrated that Cbl and Cbl-b interact with the Zap-70 tyrosine kinase and that phosphorylated Cbl and Cbl-b are enhanced in their capacity to associate with the adaptor proteins Crk-L, Nck, PLC- γ and Fyb in response to SDF-1/CXCL12. This extends the studies of others [6] on Cbl associations in response to SDF-1/CXCL12 and demonstrates additional effects on Cbl-b and associated proteins.

A functional role for both Cbl and Cbl-b in SDF-1/CXCL12-induced chemotaxis was identified in our studies in which Jurkat T cells, which were transfected with siRNA for either Cbl or Cbl-b and was decreased in expression of both these molecules, were also decreased in response to SDF-1/CXCL12-induced chemotaxis. Of interest was that MAPK activity in response to SDF-1/CXCL12 was not influenced by the decreased Cbl and Cbl-b expression in the Cbl- and Cbl-b-transfected cells.

CD45 is an important regulator of src family kinases [24]. The phosphatase activity of CD45 appears to be required for activation of src family kinase members. In contrast, in some reports, CD45 inhibits activation of src family kinases in macrophages and some lymphocytes. Thus, CD45 function while important is still somewhat controversial. In our studies, we found using a CD45-deficient cell line that SDF-1 α /CXCL12 did not enhance tyrosine phosphorylation of Cbl or

Cbl-b, demonstrating that CD45 is required for the enhanced phosphorylation of both these intracellular signals. Moreover, CD45 was, but MAPK was not, required for the chemotactic response of cells to SDF-1 α /CXCL12, in part adding valuable confirmation and extension to the work of others [7] in this still somewhat controversial area. Thus, our results implicate Cbl and Cbl-b as intracellular and functional mediators in cell migration induced by SDF-1 α /CXCL12, effects that likely involve interactions of Cbl and Cbl-b with Crk-L, Zap70, Nck, PLC- γ , Fyb and Fyn and regulation by CD45 and Src kinases.

Acknowledgments

These studies were supported by US Public Health Service NIH RO1 Grants: RO1 HL67384 and RO1 DK53674 from the NIH to HEB.

References

- [1] T. Nagasawa, K. Tachibana, T. Kishimoto, A novel CXC chemokine PBSF/SDF-1 and its receptor CXCR4: their functions in development, hematopoiesis and HIV infection, *Semin. Immunol.* 10 (1998) 179–185.
- [2] M. Loetscher, T. Geiser, T. O'Reilly, R. Zwahlen, M. Baggiolini, B. Moser, Cloning of a human seven-transmembrane domain receptor, LESTR, that is highly expressed in leukocytes, *J. Biol. Chem.* 269 (1994) 232–237.
- [3] H.E. Broxmeyer, Regulation of hematopoiesis by chemokine family members, *Int. J. Hematol.* 74 (2001) 9–17.
- [4] B.S. Youn, C. Mantel, H.E. Broxmeyer, Chemokines, chemokine receptors and hematopoiesis, *Immunol. Rev.* 177 (2000) 150–174.
- [5] Q. Ma, D. Jones, P.R. Borghesani, R.A. Segal, T. Nagasawa, T. Kishimoto, R.T. Bronson, T.A. Springer, Impaired B-lymphopoiesis, myelopoiesis, and derailed cerebellar neuron migration in CXCR4- and SDF-1-deficient mice, *Proc. Natl. Acad. Sci. U. S. A.* 95 (1998) 9448–9453.
- [6] R.D. Chemoek, R.P. Cherla, R.K. Ganju, SHP2 and cbl participate in α -chemokine receptor CXCR4-mediated signaling pathways, *Blood* 97 (2001) 608–615.
- [7] A.Z. Fernandis, R.P. Cherla, R.K. Ganju, Differential regulation of CXCR4-mediated T-cell chemotaxis and mitogen-activated protein kinase activation by the membrane tyrosine phosphatase, CD45, *Biol. Chem.* 278 (2003) 9536–9543.
- [8] W.Y. Langdon, J.W. Hartley, S.P. Klinken, S.K. Ruscetti, H.C. Morse III, v-cbl, an oncogene from a dual-recombinant murine retrovirus that induces early B-lineage lymphomas, *Proc. Natl. Acad. Sci. U. S. A.* 86 (1989) 1168–1172.
- [9] L. Buday, A. Khwaja, S. Sipeki, A. Farago, J. Downward, Interactions of Cbl with two adapter proteins, Grb2 and Crk, upon T cell activation, *J. Biol. Chem.* 271 (1996) 6159–6163.
- [10] H. Meisner, B.R. Conway, D. Hartley, M.P. Czech, Interactions of Cbl with Grb2 and phosphatidylinositol 3'-kinase in activated Jurkat cells, *Mol. Cell. Biol.* 15 (1995) 3571–3578.
- [11] O.M. Rivero-Lezcano, J.H. Sameshima, A. Marcilla, K.C. Robbins, Physical association between Src homology 3 elements and the protein product of the c-cbl proto-oncogene, *J. Biol. Chem.* 269 (1994) 17363–17366.
- [12] E.A. Feshchenko, W.Y. Langdon, A.Y. Tsygankov, Fyn, Yes, and Syk phosphorylation sites in c-Cbl map to the same tyrosine residues that become phosphorylated in activated T cells, *J. Biol. Chem.* 273 (1998) 8323–8331.
- [13] M.M. Keane, O.M. Rivero-Lezcano, J.A. Mitchell, K.C. Robbins, S. Lipkowitz, Cloning and characterization of cbl-b: a SH3 binding protein with homology to the c-cbl proto-oncogene, *Oncogene* 10 (1995) 2367–2377.
- [14] Y.C. Liu, A. Altman, Cbl: complex formation and functional implications, *Cell. Signal.* 10 (1998) 377–385.
- [15] M.A. Murphy, R.G. Schnell, D.J. Venter, L. Barnet, I. Bertoncello, C.B. Thien, W.Y. Langdon, D.D. Bowtell, Tissue hyperplasia and enhanced T-cell signalling via ZAP-70 in c-Cbl-deficient mice, *Mol. Cell. Biol.* 18 (1998) 4872–4882.
- [16] C.B. Thien, D.D. Bowtell, W.Y. Langdon, Perturbed regulation of ZAP-70 and sustained tyrosine phosphorylation of LAT and SLP-76 in c-Cbl-deficient thymocytes, *J. Immunol.* 162 (1999) 7133–7139.
- [17] Y. Ota, L.E. Samelson, The product of the proto-oncogene c-cbl: a negative regulator of the Syk tyrosine kinase, *Science* 276 (1997) 418–420.
- [18] S. Okabe, S. Fukuda, H.E. Broxmeyer, Activation of Wiskott–Aldrich syndrome protein and its association with other proteins by stromal cell-derived factor-1 α is associated with cell migration in a T-lymphocyte line, *Exp. Hematol.* 30 (2002) 761–766.
- [19] S. Tanaka, L. Neff, R. Baron, J.B. Levy, Tyrosine phosphorylation and translocation of the c-cbl protein after activation of tyrosine kinase signaling pathways, *J. Biol. Chem.* 270 (1995) 14347–14351.
- [20] J.A. Donovan, Y. Ota, W.Y. Langdon, L.E. Samelson, Regulation of the association of p120cbl with Grb2 in Jurkat T cells, *J. Biol. Chem.* 271 (1996) 26369–26374.
- [21] A.J. da Silva, Z. Li, C. de Vera, E. Canto, P. Findell, C.E. Rudd, Cloning of a novel T-cell protein FYB that binds FYN and SH2-domain-containing leukocyte protein 76 and modulates interleukin 2 production, *Proc. Natl. Acad. Sci. U. S. A.* 94 (1997) 7493–7498.
- [22] A. Veillette, M.A. Bookman, E.M. Horak, J.B. Bolen, The CD4 and CD8 T cell surface antigens are associated with the internal membrane tyrosine-protein kinase p56lck, *Cell* 55 (1988) 301–308.
- [23] A.M. Carmo, D.W. Mason, A.D. Beyer, Physical association of the cytoplasmic domain of CD2 with the tyrosine kinases p56lck and p59fyn, *Eur. J. Immunol.* 23 (1993) 2196–2201.
- [24] M.L. Thomas, E.J. Brown, Positive and negative regulation of Src-family membrane kinases by CD45, *Immunol. Today* 20 (1999) 406–411.
- [25] J. English, G. Pearson, J. Wilsbacher, J. Swantek, M. Karandikar, S. Xu, M.H. Cobb, New insights into the control of MAP kinase pathways, *Exp. Cell Res.* 253 (1999) 255–270.
- [26] S. Okabe, S. Fukuda, H.E. Broxmeyer, Src kinase, but not the src kinase family member p56lck, mediates stromal cell-derived factor 1 α /CXCL12 induced chemotaxis of a T cell line, *J. Hematother. Stem Cell Res.* 11 (2002) 923–928.
- [27] A.Y. Tsygankov, A.M. Teckchandani, E.A. Feshchenko, G. Swaminathan, Beyond the RING: CBL proteins as multivalent adapters, *Oncogene* 20 (2001) 6382–6402.
- [28] P.S. Ohashi, T-cell signalling and autoimmunity: molecular mechanisms of disease, *Nat. Rev., Immunol.* 2 (2002) 427–438.

Clonally Expanded T-Cells in the Peripheral Blood of Patients with Idiopathic Thrombocytopenic Purpura and *Helicobacter pylori* Infection

Midori Ishiyama,^a Masanao Teramura,^a Koji Iwabe,^a Tomohiro Kato,^b Toshiko Motoji^a

^aDepartment of Haematology, Tokyo Women's Medical University, Tokyo; ^bRheumatology, Immunology and Genetics Program, Institute of Medical Science, St. Marianna University School of Medicine, Kanagawa, Japan

Received August 1, 2005; received in revised form November 2, 2005; accepted November 14, 2005

Abstract

Eradication of *Helicobacter pylori* leads to platelet recovery in some patients with idiopathic thrombocytopenic purpura (ITP). Therefore, the pathogenesis of a subgroup of ITP is probably associated with *H pylori* infection (*H pylori*-related ITP). If *H pylori*-related ITP is a definite subgroup of ITP, specific oligoclonal T-cells might accumulate in the peripheral blood (PB). To address this issue, we performed single-strand conformation polymorphism analysis of complementarity-determining region 3 (CDR3) of the T-cell receptor β -chain genes of PB T-cells. Fourteen ITP patients with *H pylori* infection and 12 age-adjusted healthy volunteers were studied. Of the 14 patients, 8 patients (responders) exhibited a platelet response after successful *H pylori* eradication therapy, but 6 patients (nonresponders) did not. V β 5.2, V β 15, and V β 19 gene usage by clonally expanded T-cells in PB obtained before *H pylori* eradication therapy was significantly higher in responders than in nonresponders or healthy volunteers (V β 5.2, $P = .023$; V β 15, $P = .004$; V β 19, $P = .036$). Furthermore, an abrogation of clonally expanded T-cells was observed after therapy in some responders. These findings suggest that specific T-cell clones accumulate in *H pylori*-related ITP and that such clones may be associated with immune-mediated destruction of platelets.

Int J Hematol. 2006;83:147-151. doi: 10.1532/IJH97.05119

©2006 The Japanese Society of Hematology

Key words: Idiopathic thrombocytopenic purpura (ITP); *Helicobacter pylori*; T-cell clonality; T-cell receptor; Single-strand conformation polymorphism (SSCP)

1. Introduction

Idiopathic thrombocytopenic purpura (ITP) is an autoimmune disorder caused by autoantibodies against platelets. Platelet membrane glycoproteins (GPs) such as GP IIb-IIIa and GP Ib have been identified as target autoantigens. Recently, many investigators reported that eradication of *Helicobacter pylori* leads to platelet recovery in patients with ITP (response rate, 33%-100%) [1-11]. However, a few investigators have shown the opposite result [12-14]. Therefore, *H pylori* infection may play an important pathophysiological role in a subgroup of ITP (*H pylori*-related ITP).

Oligoclonal T-cells that respond to antigenic stimulation accumulate in the peripheral blood (PB) in autoimmune diseases. Single-strand conformation polymorphism (SSCP) analysis of complementarity-determining region 3 (CDR3)

of the T-cell receptor (TCR) β chain has demonstrated that oligoclonal T-cells accumulate in the PB of ITP patients [15]. This result suggests that specific oligoclonally expanded T-cells that drive B-cells to produce autoantibodies against platelets may be present in the PB of patients with ITP.

H pylori infection is prevalent among healthy people. In Japan, the frequency of *H pylori* infection is greater than 50% in healthy adults and 70% to 80% among elderly people [16]. Therefore, the occurrence of complicating *H pylori* infection in ITP patients does not necessarily mean that these patients have *H pylori*-related ITP. The diagnosis of *H pylori*-related ITP is made retrospectively according to the platelet response following *H pylori* eradication therapy. If a patient with *H pylori* infection recovers from thrombocytopenia after successful eradication therapy (ie, a responder), a diagnosis of *H pylori*-related ITP can be made. On the other hand, if a patient fails to recover from thrombocytopenia even after *H pylori* has successfully been eradicated (ie, a nonresponder), a diagnosis of *H pylori*-unrelated ITP can be made. If specific clonally expanded T-cells that are different from those of nonresponders and healthy volunteers can be demonstrated to be present in the PB of responders, it may

Correspondence and reprint requests: Masanao Teramura, MD, Department of Haematology, Tokyo Women's Medical University, 8-1, Kawada-cho, Shinjuku-ku, Tokyo, 162-8666, Japan; 81-3-3353-8111; fax: 81-3-5269-7363 (e-mail: teramura@dh.twmu.ac.jp).

Table 1.

Clinical and Laboratory Characteristics of 14 Patients with Idiopathic Thrombocytopenic Purpura*

Patient No.	Age, y	Sex	Disease Duration, mo	Previous Treatment	¹³ C-Urea Breath Test, ‰			Platelets, ×10 ⁹ /L		Response
					Before <i>H pylori</i> Eradication	After <i>H pylori</i> Eradication	<i>H pylori</i> Eradication	Before <i>H pylori</i> Eradication	After <i>H pylori</i> Eradication	
1	50	M	38	PSL	38	0.9	Yes	14	138	+
2	52	M	6	—	16	2.1	Yes	14	239	+
3	56	M	18	PSL	4	1.3	Yes	54	95	+
4	56	M	10	PSL	14.9	1.8	Yes	18	105	+
5	63	F	23	PSL, Sp	10.2	1.2	Yes	20	92	+
6	74	F	58	Vitamin C	20	1.1	Yes	49	182	+
7	74	F	6	—	19.8	1.4	Yes	35	125	+
8	76	M	11	—	3	1.2	Yes	23	95	+
9	34	F	9	PSL	43.8	0.2	Yes	52	67	-
10	43	F	133	mPSL, Dan	4.3	1	Yes	19	11	-
11	50	F	78	—	49.3	0.6	Yes	73	66	-
12	53	F	16	—	10.3	1.2	Yes	70	92	-
13	58	F	60	PSL	20.1	1.3	Yes	31	18	-
14	61	F	63	Vitamin C	39	0.8	Yes	38	43	-

*Eight patients (nos. 1-8) showed a platelet response after *Helicobacter pylori* eradication therapy, and 6 patients (nos. 9-14) did not show a platelet response after eradication therapy. PSL indicates prednisolone; Sp, splenectomy; mPSL, methylprednisolone; Dan, danazol.

be possible to show that *H pylori*-related ITP is indeed a definite subgroup of ITP. To address this issue, we performed SSCP analysis of TCR V β -chain genes of PB T-cells in ITP patients with *H pylori* infection and investigated T-cell repertoire usage by the clonally expanded T-cells.

2. Materials and Methods

2.1. Patients

Fourteen patients with chronic ITP complicated by *H pylori* infection were studied. The patients comprised 5 men and 9 women with a median age of 56 years (range, 35-72 years). Chronic ITP was defined as thrombocytopenia (platelets <100 × 10⁹/L) lasting for at least 6 months, normal or increased numbers of megakaryocytes in the bone marrow, and absence of other apparent causes of thrombocytopenia. *H pylori* infection was diagnosed by a positive value of greater than 2.5% in the breath test using carbon 13 (¹³C)-labeled urea [17].

H pylori infection was initially eradicated by treatment with amoxicillin (750 mg twice daily), clarithromycin (200 mg twice daily), and lansoprazole (30 mg twice daily) for 7 days. After a minimum of 2 months, the ¹³C-urea breath test was again performed to evaluate the effect of *H pylori* eradication therapy. *H pylori* was eradicated in 12 of the 14 patients. The 2 patients (patients 13 and 14) who failed to respond were then successfully treated with metronidazole (250 mg twice daily), amoxicillin (750 mg twice daily), and lansoprazole (30 mg twice daily). Therefore, *H pylori* infection was eventually eradicated in all 14 patients.

The platelet response due to *H pylori* eradication was evaluated 6 months after treatment. A response was defined as an absolute increase in the platelet count of >30 × 10⁹/L from the baseline. The response criteria are based on the previous report by Vianelli et al [18]. As shown in Table 1, eradication of *H pylori* led to a platelet response in 8 patients

(patients 1-8) but no response in 6 patients (patients 9-14). All of the responders are now in remission with a platelet count of >100 × 10⁹/L for at least 34 months after treatment.

2.2. SSCP Analysis of TCR β -Chain Genes

Samples of heparinized PB were obtained from all patients before eradication therapy. Samples were also obtained 3 to 6 months after eradication therapy for some patients. As a control, PB samples were also obtained from 12 age-adjusted healthy volunteers (5 men and 7 women with a median age of 57.5 years [range, 35-72 years]) with no history of recent infection. All patients gave informed consent.

SSCP analysis of TCR β -chain genes was performed as described elsewhere [19]. In brief, mononuclear cells were separated from the PB samples by density-gradient centrifugation over Ficoll-Paque (Amersham Biosciences, Uppsala, Sweden), and total RNA was extracted by the acid guanidinium thiocyanate-phenol-chloroform method [20]. RNA was converted to complementary DNA in a solution containing 200 U reverse transcriptase (SuperScript; Gibco BRL, Gaithersburg, MD, USA) and 100 pmol random hexamer oligonucleotide primer (Gibco BRL). For amplification of each TCR V β gene family, a biotinylated primer for the constant region of the β chain and a V β -specific primer were used. The sequences of the V β -specific primers were as described elsewhere [21]. The polymerase chain reaction was performed with deoxynucleoside triphosphates and *Taq* DNA polymerase (TaKaRa Bio, Shiga, Japan) for 35 cycles in a thermocycler (PerkinElmer, Norwalk, CT, USA). Following dilution and heat denaturation, amplified DNA fragments were separated on the basis of differences in their single-strand conformation by electrophoresis on nondenaturing 4% polyacrylamide gels containing 10% glycerol. The electrophoresed DNA fragments were transferred to membranes (GeneScreen; NEN Life Science Products, Boston, MA, USA) and visualized by subsequent incubations with strep-

tavidin, biotinylated alkaline phosphatase, and a chemiluminescent substrate system (Phototope-Star Detection kit; New England BioLabs, Beverly, MA, USA). We counted distinct bands as corresponding to clonal T-cell expansions. Although bands usually were easily recognized visually, a densitometer (ACD-25DX; ATTO Technology, Tokyo, Japan) was also used to confirm the presence of the bands. If one or more distinct bands were present in a certain V β gene family, the existence of clonally expanded T-cells was demonstrated. The presence of clonally expanded T-cells in each TCR V β gene family was individually assessed, and then the proportion of patients (or healthy volunteers) with clonally expanded T-cells was calculated for each TCR V β gene.

2.3. DNA Sequencing

TCR V β gene transcripts obtained before and after eradication therapy from one patient who had recovered from thrombocytopenia after *H pylori* eradication were extracted from the SSCP gel and cloned with a TA cloning kit (Invitrogen, Carlsbad, CA, USA). Approximately 20 plaques were randomly chosen and subjected to dideoxy direct sequencing.

2.4. Statistical Analysis

Differences in proportions among the groups were evaluated by the Kruskal-Wallis test, the Student *t* test, or the chi-square test. A *P* value <.05 was considered statistically significant.

3. Results

3.1. Analysis of the Clonally Expanded T-Cells in ITP Patients with *H pylori* Infection

We performed SSCP analyses of 20 major TCR V β gene families in patients with *H pylori* infection to investigate whether clonally expanded T-cells were present in PB. The median number of TCR V β gene families with clonally expanded T-cells in responders, nonresponders, and healthy subjects was 7 (range, 1-21), 3.5 (range, 2-11), and 4 (range, 1-6), respectively. The number of V β gene families with clonally expanded T-cells was significantly greater in responders (*P* = .032).

TCR V β subfamilies that frequently (>50% of cases) revealed clonally expanded T-cells were observed in 7 families (V β 1, V β 2, V β 5.2, V β 7, V β 11, V β 15, and V β 19) in responders, 2 families (V β 1 and V β 5.1) in nonresponders, and 3 families (V β 7, V β 10, and V β 11) in healthy volunteers (Figure 1). We analyzed the differences in V β usage of clonally expanded T-cells among responders, nonresponders, and healthy volunteers and found that the usage of V β 5.2, V β 15, and V β 19 genes of clonally expanded T-cells was significantly higher in responders than in nonresponders or healthy volunteers (V β 5.2, *P* = .023; V β 15, *P* = .004; V β 19, *P* = .036). We investigated whether the distinct bands in the SSCP analysis of V β 5.2 (patients 1, 2, 3, and 6), V β 15 (patients 3, 4, and 8), or V β 19 (patients 3, 6, and 8) genes that were found in some responders before therapy disappeared after therapy. Disappearance of distinct bands from V β 5.2, V β 15, and V β 19




Figure 1. T-cell receptor (TCR) V β subfamily usage by clonally accumulated T-cells for responders, nonresponders, and healthy volunteers. Peripheral blood samples from patients with idiopathic thrombocytopenic purpura were obtained before eradication therapy and analyzed. If one or more distinct bands for a given V β gene were present in the single-strand conformation polymorphism analysis, the existence of clonally expanded T-cells was confirmed. The presence of clonally expanded T-cells for each TCR V β gene was individually assessed, and then the proportion of patients (or healthy volunteers) with clonal T-cell expansion was calculated for each TCR V β gene. Responders indicates patients who recovered from thrombocytopenia after successful *Helicobacter pylori* eradication; nonresponders, patients who failed to recover from thrombocytopenia even after *H pylori* was successfully eradicated. *V β 5.2, V β 15, and V β 19 gene usage was significantly higher in responders than in nonresponders or healthy volunteers (*P* = .023, .004, and .036, respectively).

genes was observed in none of 4 responders, 1 (patient 3) of 3 responders, and 1 (patient 3) of 3 responders, respectively (data not shown).

We compared nonresponders and healthy volunteers with respect to V β usage by clonally expanded T-cells and found V β 3, V β 4, V β 5.1, and V β 8 gene usage to be significantly higher in nonresponders (*P* = .034, .034, .034, and .034, respectively).

3.2. DNA Sequencing of TCR CDR3

In a patient (patient 1) who recovered from thrombocytopenia after *H pylori* eradication, the distinct band that was seen in the SSCP analysis of the V β 8 gene before eradication therapy disappeared after eradication therapy (Figure 2). To confirm the disappearance of clonally expanded T-cells in V β 8 following *H pylori* eradication, we determined the CDR3 DNA sequences of the TCR V β genes. In the sample obtained before *H pylori* eradication, all 18 subcloned genes showed the same sequence. However, in the sample obtained after *H pylori* eradication, all 19 subcloned genes showed different sequences, and none of these sequences were identical to the sequence seen before *H pylori* eradication (Table 2).

4. Discussion

We performed an SSCP analysis of TCR V β -chain genes of PB T-cells from ITP patients with *H pylori* infection and




Figure 2. Single-strand conformation polymorphism analysis of T-cell receptor (TCR) V β genes in peripheral blood T-cells. The number at the top of each lane indicates the TCR V β gene subfamily. Results obtained from a typical healthy control subject (A) and patient 1 (responder) (B) are shown. In patient 1, a distinct band for the V β 8 gene segment present before *Helicobacter pylori* eradication (B-1) disappeared after *H pylori* eradication (B-2) (arrows).

investigated T-cell repertoire usage by clonally expanded T-cells. The number of TCR V β gene families with clonally expanded T-cells was significantly higher in the patients with *H pylori* infection who subsequently recovered from thrombocytopenia after successful eradication therapy (responders) than in the patients who failed to recover from thrombocytopenia after successful eradication therapy (nonresponders). In addition, the usage of V β 5.2, V β 15, and

Table 2.

Deduced Amino Acid Sequences of the T-Cell Receptor (TCR) β Chains Carrying the V β Gene Segment Derived from the Peripheral Blood of Patient 1

V β 8	nDn	J β (Gene Segment)	TCR β Chains, n
Before <i>Helicobacter pylori</i> eradication			
CASS	FSYCSA	NYGYT (J1S2)	18 (100%)
After <i>Helicobacter pylori</i> eradication			
CASSL	AWSGRY	TGELF (J2S2)	1 (5.3%)
CAS	RTTGG	SYEQY (J2S7)	1 (5.3%)
CASS	FSGGR	ETQYF (J2S5)	1 (5.3%)
CASS	KTGYE	QYFGP (J2S3)	1 (5.3%)
CAS	SRLAGGHPPT	QYFGP (J2S7)	1 (5.3%)
CAS	TRPEGGT	YNEQFF (J2S1)	1 (5.3%)
CAS	EEG	NTEAF (J1S1)	1 (5.3%)
CAS	SRFPAGA	YEQYF (J2S7)	1 (5.3%)
CA	SRPLAP	QETQYF (J2S5)	1 (5.3%)
CASS	SATV	SYEQY (J2S7)	1 (5.3%)
CASS	PRLDG	SYEQY (J2S7)	1 (5.3%)
CASS	RDFRA	NYGYT (J1S2)	1 (5.3%)
CASS	FGGTAR	QETQYF (J2S5)	1 (5.3%)
CASS	GTGTTSD	EQFFGPG (J2S1)	1 (5.3%)
CASSL	RPY	QPQHFG (J1S5)	1 (5.3%)
CAS	QGQH	NSPLHF (J1S6)	1 (5.3%)
CAS	NRLAGGHP	DTQYFGP (J2S3)	1 (5.3%)
CASSL	ELQDGYA	FGSGTRL (J1S2)	1 (5.3%)
CAS	RL	SGANVLT (J2S6)	1 (5.3%)

V β 19 genes by clonally expanded T-cells was significantly higher in responders than in nonresponders. This difference notably does not derive from the presence or absence of *H pylori* infection, because all of the patients had been infected with *H pylori*. These results suggest that some clonally expanded T-cells with specific TCR V β subfamily usage are present in patients with *H pylori*-related ITP. Distinct bands seen in the SSCP analysis of V β genes before *H pylori* eradication therapy in some of the responders disappeared after *H pylori* eradication therapy. Furthermore, we confirmed the disappearance of clonally expanded T-cells after *H pylori* eradication therapy in a patient with *H pylori*-related ITP by analyzing the DNA sequences of CDR3 of V β genes. Our results indicate that clonally expanded T-cells were abrogated by *H pylori* eradication and suggest that the disappearance of clonally expanded T-cells was responsible for platelet recovery.

H pylori infection is associated with various autoimmune diseases, including rheumatoid arthritis, Sjögren syndrome, and autoimmune hypothyroidism [22-25]. Clinical data from patients with these disorders raise the possibility that immune reactions against *H pylori* have pivotal roles in the onset of autoimmune diseases. As for *H pylori*-related ITP, why *H pylori* eradication is able to induce platelet recovery is unknown. However, one possible explanation is that anti-*H pylori* antibodies bind to platelets in the presence of cross-mimicry between platelet surface antigens and *H pylori* antigens, resulting in platelet destruction. This speculation is supported by recent work demonstrating that platelet-associated immunoglobulin possesses cross-reactivity to

H pylori cytotoxin-associated gene A (CagA) [26]. If this mechanism operates in *H pylori*-related ITP, it strongly suggests that *H pylori* eradication induces a reduction in the T-cell clones against CagA that drive B-cells to produce cross-reactive antibodies, resulting in the reduction of cross-reactive antibodies. Our data demonstrating the subsequent disappearance of clonally expanded T-cells after *H pylori* eradication support this speculation.

Cytotoxic T-cell-mediated lysis of autologous platelets has recently been demonstrated in active ITP, and T-cell-mediated cytotoxicity has been suggested to be an alternative mechanism of platelet destruction in ITP [27]. Therefore, it is also possible that clonally expanded T-cells observed in *H pylori*-related ITP are cytotoxic T-cells against *H pylori* with cross-reactivity to platelets.

In conclusion, our findings suggest that specific T-cell clones accumulate in *H pylori*-related ITP and that these clones may be associated with immune-mediated platelet destruction. Further studies are needed to elucidate the role of the clonally expanded T-cells observed in *H pylori*-related ITP.

References

- Gasbarrini A, Franceschi F, Tartaglione R, Landolfi R, Pola P, Gasbarrini G. Regression of autoimmune thrombocytopenia after eradication of *Helicobacter pylori*. *Lancet*. 1998;352:878.
- Emilia G, Longo G, Luppi M, et al. *Helicobacter pylori* eradication can induce platelet recovery in idiopathic thrombocytopenic purpura. *Blood*. 2001;97:812-814.
- Kohda K, Kuga T, Kogawa K, et al. Effect of *Helicobacter pylori* eradication on platelet recovery in Japanese patients with chronic idiopathic thrombocytopenic purpura and secondary autoimmune thrombocytopenic purpura. *Br J Haematol*. 2002;118:584-588.
- Veneri D, Franchini M, Gottardi M, et al. Efficacy of *Helicobacter pylori* eradication in raising platelet count in adult patients with idiopathic thrombocytopenic purpura. *Haematologica*. 2002;87:1177-1179.
- Hino M, Yamane T, Park K, et al. Platelet recovery after eradication of *Helicobacter pylori* in patients with idiopathic thrombocytopenic purpura. *Ann Hematol*. 2003;82:30-32.
- Hashino S, Mori A, Suzuki S, et al. Platelet recovery in patients with idiopathic thrombocytopenic purpura after eradication of *Helicobacter pylori*. *Int J Hematol*. 2003;77:188-191.
- Ando T, Tsuzuki T, Mizuno T, et al. Characteristics of *Helicobacter pylori*-induced gastritis and the effect of *H. pylori* eradication in patients with chronic idiopathic thrombocytopenic purpura. *Helicobacter*. 2004;9:443-452.
- Sato R, Murakami K, Watanabe K, et al. Effect of *Helicobacter pylori* eradication on platelet recovery in patients with chronic idiopathic thrombocytopenic purpura. *Arch Intern Med*. 2004;164:1904-1907.
- Fujimura K, Kuwana M, Kurata Y, et al. Is eradication therapy useful as the first line of treatment in *Helicobacter pylori*-positive idiopathic thrombocytopenic purpura? Analysis of 207 eradicated chronic ITP cases in Japan. *Int J Hematol*. 2005;81:162-168.
- Inaba T, Mizuno M, Take S, et al. Eradication of *Helicobacter pylori* increases platelet count in patients with idiopathic thrombocytopenic purpura in Japan. *Eur J Clin Invest*. 2005;35:214-219.
- Stasi R, Rossi Z, Stipa E, Amadori S, Newland AC, Provan D. *Helicobacter pylori* eradication in the management of patients with idiopathic thrombocytopenic purpura. *Am J Med*. 2005;118:414-419.
- Jarque I, Andrew R, Llopi I, et al. Absence of platelet response after eradication of *Helicobacter pylori* infection in patients with chronic idiopathic thrombocytopenic purpura. *Br J Haematol*. 2001;115:1002-1003.
- Michel M, Khellaf M, Desforges L, et al. Autoimmune thrombocytopenic purpura and *Helicobacter pylori* infection. *Arch Intern Med*. 2002;162:1033-1036.
- Michel M, Cooper N, Jean C, Frizzera C, Busnel JB. Does *Helicobacter pylori* initiate or perpetuate immune thrombocytopenic purpura? *Blood*. 2004;103:890-896.
- Shimomura T, Fujimura K, Takafuta T, et al. Oligoclonal accumulation of T cells in peripheral blood from patients with idiopathic thrombocytopenic purpura. *Br J Haematol*. 1996;95:732-737.
- Asaka M, Kimura T, Kudo M, et al. Relationship of *Helicobacter pylori* to serum pepsinogen in an asymptomatic Japanese population. *Gastroenterology*. 1992;102:760-766.
- Ohara S, Kato M, Asaka M, Toyota T. Studies of ¹³C-urea breath test for diagnosis of *Helicobacter pylori* infection in Japan. *J Gastroenterol*. 1998;33:6-13.
- Vianelli N, Valdre L, Fiacchini M, et al. Long-term follow-up of idiopathic thrombocytopenic purpura in 310 patients. *Haematologica*. 2001;86:504-509.
- Yamamoto K, Sakoda H, Nakajima T, et al. Accumulation of multiple T cell clonotypes in the synovial lesions of patients with rheumatoid arthritis revealed by a novel clonality analysis. *Int Immunol*. 1992;4:1219-1223.
- Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem*. 1987;162:156-159.
- Choi YW, Kotzin B, Herron L, Callahan J, Marrack P, Kappler J. Interaction of *Staphylococcus aureus* toxin "superantigens" with human T cells. *Proc Natl Acad Sci U S A*. 1989;86:8941-8945.
- Gasbarrini A, Franceschi F. Autoimmune disease and *Helicobacter pylori* infection. *Biomed Pharmacother*. 1999;53:223-226.
- Zentilin P, Serio B, Dulbecco P, et al. Eradication of *Helicobacter pylori* may reduce disease severity in rheumatoid arthritis. *Aliment Pharmacol Ther*. 2002;16:1291-1299.
- Figura N, Giordano N, Burroni D, et al. Sjögren's syndrome and *Helicobacter pylori* infection. *Eur J Gastroenterol Hepatol*. 1994;6:321-322.
- de Luis DA, Varela C, de La Calle H, et al. *Helicobacter pylori* infection is markedly increased in patients with autoimmune atrophic thyroiditis. *J Clin Gastroenterol*. 1998;26:259-263.
- Takahashi T, Yujiri T, Shinohara K, et al. Molecular mimicry by *Helicobacter pylori* CagA protein may be involved in the pathogenesis of *H. pylori*-associated chronic idiopathic thrombocytopenic purpura. *Br J Haematol*. 2004;124:91-96.
- Olsson B, Andersson PO, Jernas M, et al. T-cell-mediated cytotoxicity toward platelets in chronic idiopathic thrombocytopenic purpura. *Nat Med*. 2003;9:1123-1124.

ORIGINAL ARTICLE

Hyperactivation of the RAS signaling pathway in myelodysplastic syndrome with *AML1/RUNX1* point mutations

H Niimi^{1,5}, H Harada^{1,5}, Y Harada^{2,5}, Y Ding¹, J Imagawa¹, T Inaba³, T Kyo⁴ and A Kimura¹

¹Department of Hematology/Oncology, Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan; ²International Radiation Information Center, Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan; ³Department of Molecular Oncology, Research Institute for Radiation Biology and Medicine, Hiroshima University, Hiroshima, Japan and ⁴Department of Internal Medicine, Hiroshima Red Cross Hospital and Atomic Bomb Survivors Hospital, Hiroshima, Japan

AML1/RUNX1 mutations have been reported frequently in myelodysplastic syndrome (MDS) patients, especially those diagnosed with refractory anemia with excess blast (RAEB), RAEB in transformation (RAEBt), or AML following MDS (these categories are defined as MDS/AML). Although AML1 mutations are suspected to play a pivotal role in the development of MDS/AML, acquisition of additional genetic alterations is also necessary. We analyzed gene alterations in MDS/AML patients with AML1 mutations, comparing them to alterations in those without an AML1 mutation. AML1 mutations were significantly associated with $-7/7q-$, whereas MDS/AML patients without AML1 mutations showed a high frequency of $-5/5q-$ and a complex karyotype. Patients with AML1 mutations showed more mutations of their *FLT3*, *N-RAS*, *PTPN11*, and *NF1* genes, resulting in a significantly higher mutation frequency for receptor tyrosine kinase (RTK)–RAS signaling pathways in AML1-mutated MDS/AML patients compared to AML1-wild-type MDS/AML patients (38% versus 6.3%, $P < 0.0001$). Conversely, *p53* mutations were detected only in patients without AML1 mutations. Furthermore, blast cells of the AML1-mutated patients expressing surface c-KIT, and SHP-2 mutants contributed to prolonged and enhanced extracellular signal-regulated kinase activation following stem cell factor stimulation. Our results suggest that MDS/AML arising from AML1/RUNX1 mutations has a significant association with $-7/7q-$ alteration, and frequently involves RTK–RAS signaling pathway activation.

Leukemia (2006) 20, 635–644. doi:10.1038/sj.leu.2404136; published online 9 February 2006

Keywords: AML1/RUNX1; MDS/AML; point mutation; second hit; RTK/RAS

Introduction

Somatically acquired point mutations of critical genes have been demonstrated to contribute to the development of acute myeloid leukemia (AML) and myelodysplastic syndrome (MDS). Genes encoding key regulatory factors for cell division, differentiation, or cell survival of hematopoietic progenitors, as well as transcription factors, are frequent mutation targets. The *AML1/RUNX1* gene was also found to be altered by point mutations in AML and MDS, and unique features associated with these mutations have been revealed by several studies.

Correspondence: Dr H Harada, Department of Hematology/Oncology, Research Institute for Radiation Biology and Medicine, Hiroshima University, 1-2-3 Kasumi, Minami-ku, Hiroshima 734-8553, Japan. E-mail: herf1@hiroshima-u.ac.jp

⁵These authors contributed equally to this work.

Received 12 August 2005; revised 22 December 2005; accepted 30 December 2005; published online 9 February 2006

First, although the frequency of AML1 mutations in *de novo* AML is low, they have been detected with a substantially higher frequency in poorly differentiated AML M0,^{1–3} and in radiation-associated and therapy-related AML.^{4,5} Second, germ-line mutations of *AML1* have been shown to occur in a rare autosomal dominant disorder, familial platelet disorder with predisposition to AML (FPD/AML).^{6,7} Third, we previously reported a high frequency of AML1 mutations in MDS patients, especially those with refractory anemia with excess blast (RAEB), RAEB in transformation (RAEBt), and AML following MDS (these categories are defined as MDS/AML).⁸ This suggests that AML1 point mutations are strongly associated with these specific types of hematopoietic malignancy. In the molecular pathogenesis, *de novo* AML is often associated with oncogenic chimeras such as AML1–ETO (MTG8), which are considered to be the major cause of malignant transformation of hematopoietic progenitors, while point mutations are likely a complement for chimeras. However, AML1 point mutations should be considered one of the major decisive factors of the development of MDS/AML, similar to chimeras in *de novo* AML.

All of the AML1 mutants detected so far, with a few exceptions, showed a lowering or loss of their *trans*-activation potential, regardless of their DNA-binding potential. This suggests that various AML1 mutants act through a loss-of-function mechanism, and contribute to development of the same type of myeloid malignancy, MDS/AML. However, loss of AML1 function caused by AML1 mutations in hematopoietic stem cells is not sufficient for an individual to develop MDS/AML, and this explains the long latency period before development of AML among people having FPD/AML pedigrees with congenital AML1 mutations. Moreover, AML1^{+/-} mice, conditional AML1^{-/-} mice or conditional heterozygous AML1–ETO knockin mice do not develop leukemia without mutagenic reagents, in spite of the increase in number and the self-renewal capacity of their hematopoietic stem cells.^{9,10} This in turn suggests that the acquisition of some additional genetic alterations that cooperate with the AML1 mutations is needed for development of MDS/AML. Recently, c-kit activating mutations were identified in 20–30% of CBF leukemia patients who had blasts containing either AML1–ETO or CBF β –MYH11 fusion proteins.¹¹ This was not observed in any of the non-CBF leukemias, suggesting that the expression of CBF fusion proteins provides a selective advantage for acquisition of a c-kit mutation, and that these events function together in the development of leukemia. This theory supports a ‘two-hit’ model for leukemogenesis.¹² The basis for the hypothesis is that AML is the consequence of a collaboration between at least two broad classes of mutation: class I (proliferative) mutations that

confer a proliferative and/or survival advantage to cells, including gene alterations of *RAS*, *c-KIT*, *FLT3*, *PTPN11* and *NF1*, and class II (blocking) mutations that primarily impair hematopoietic differentiation and subsequent cellular apoptosis, including *AML1-ETO* and *CBF β -MYH11* fusion genes. *AML1* point mutations are classified as class II mutations, and *FLT3* mutations were frequently observed in *AML1*-mutated AML M0 patients.¹³

It is hard to explain the molecular mechanisms in MDS that contribute to the transformation of hematopoietic progenitors by a simple 'two-hit' model. Many cases of MDS are generally considered to develop as a result of accumulated gene deletions and point mutations. Genetic alterations associated with MDS have been identified, including receptors for hematopoietic growth factors, RAS signaling molecules, cell cycle regulators and transcriptional factors.^{14,15} However, these are all relatively rare (<10%) and none are specific to MDS. Chromosomal abnormalities are also associated with MDS, and they vary from single numerical or structural changes to complex genomic lesions.^{14,15} Unbalanced numeric chromosomal alterations, including partial and complete chromosome loss (especially 5q-, -7, -Y and 20q-) or chromosome gain (most frequently +8), predominate in MDS. A model of stepwise genetic progression has emerged, based on genetic and cytogenetic findings, to explain the development and evolution of MDS.¹⁶ In this model, a primary genetic event incites the initial DNA damage and subsequently increases the susceptibility to further damage. Secondary genetic events promote acquisition of molecular-genetic or cytogenetic abnormalities common to MDS and precipitate additional abnormalities.

Myelodysplastic syndrome is a quite heterogeneous disease category. We have been trying to re-classify MDS based on the molecular pathogenesis, and have proposed a new disease entity, MDS/AML with *AML1* point mutations. In this subgroup of MDS, an *AML1* mutation is considered to be the primary genetic event. Similar to CBF leukemia, patients with MDS/AML with *AML1* mutations probably also have some selective secondary genetic events. Here, we report a high frequency of mutations affecting the class III receptor tyrosine kinase (RTK)-RAS signaling pathway in *AML1*-mutated MDS/AML patients. Our data suggest that *AML1* point mutations affect activation of the RTK-RAS signaling pathway, and that this may be one of the molecular mechanisms to develop MDS/AML.

Materials and methods

Patients

We examined 625 patients with hematologic diseases, including MDS, AML, acute lymphoid leukemia (ALL), chronic myeloid leukemia (CML) and myeloproliferative disorder (MPD), all of whom were diagnosed at Hiroshima University Hospital and its affiliated hospitals between 1990 and 2004. Diagnosis was based on morphologic and immunophenotypic studies according to the French-American-British (FAB) classification. Three disease categories corresponding to subgroups of MDS and AML were identified (i.e., RAEB, RAEBt and AML following MDS) as MDS/AML.⁸ Cytogenetic analyses using standard procedures were performed according to the International System of Human Cytogenetic Nomenclature (1995).¹⁷ The majority of the patients in this study were treated at Hiroshima University Hospital or Hiroshima Red Cross Hospital and Atomic Bomb Survivors Hospital using a protocol involving intensive chemotherapy and bone marrow transplantation.

Patient samples were taken after obtaining informed consent and approval from the institutional review board at Hiroshima University. Mononuclear cells were isolated from bone marrow or peripheral blood samples by Ficoll-Conray density gradient centrifugation. Genomic DNA was extracted with a Puregene Kit (Gentra, Minneapolis, MN, USA) and total RNA was extracted using a TRIzol Kit (Invitrogen, Carlsbad, CA, USA), according to the manufacturers' instructions.

Identification of AML1 mutations

Mutation analysis of *AML1* exons 3–8 was performed by polymerase chain reaction–single-strand conformation polymorphism (PCR–SSCP) of all 625 patients as described previously.⁸ Briefly, 100 ng of genomic DNA was amplified by PCR using the flanking intronic, forward/reverse primers (Supplementary Table). To identify *AML1* mutations, SSCP analysis was performed on a GenePhor system (Amersham Pharmacia Biotech, Buckinghamshire, UK). PCR products that showed abnormal bands were sequenced in both directions. To confirm the mutations, PCR products from cDNA were also sequenced.

We selected patients for further studies according to their diagnosis and their *AML1* mutations, and they were divided into three disease categories as follows: (1) 34 cases of MDS/AML with an *AML1* mutation (10 RAEB, 16 RAEBt and 8 AML following MDS); (2) 80 cases of MDS/AML without an *AML1* mutation (32 RAEB, 21 RAEBt and 27 AML following MDS) and (3) 25 cases of CBF leukemia (19 with 8; 21 translocation and 6 with inversion 16).

Polymerase chain reaction of N-RAS, K-RAS, c-KIT, PTPN11, NF1 and p53 fragments

The PCR amplified the DNA sequence of interest: exons 1 and 2 of *N-RAS* and *K-RAS*;¹⁸ exons 8 and 17 of *c-KIT*;¹⁹ exons 3 and 13 of *PTPN11*;²⁰ exons 1–49 of *NF1*²¹ and exons 5–8 of *p53*.²² Genomic DNA (100 ng) was amplified by PCR in a total volume of 25 μ l containing 1 \times PCR buffer II (Applied Biosystems, Foster City, CA, USA), 1.5 mM MgCl₂ (Applied Biosystems), 0.2 mM dNTP (deoxynucleotide triphosphate; Roche, Mannheim, Germany), 0.2 μ M of each primer, and 0.5 unit of AmpliTaq Gold (Applied Biosystems). PCR was performed on genes of interest using the flanking intronic, forward/reverse primers listed in the Supplementary Table. Amplicons were checked by agarose gel electrophoresis to make sure that only the specific product was amplified.

Denaturing high performance liquid chromatography analysis

PCR-amplified samples were warmed to 95°C, then cooled to 25°C over approximately 45 min to promote heteroduplex formation. Denaturing high performance liquid chromatography (DHPLC) was performed on a WAVE DNA fragment analysis system (Transgenomic, Omaha, NE, USA). Briefly, 5 μ l of each DNA sample was injected into a high-throughput DNasep column and eluted through a 260 nm photodetector with concentrations of buffer A and B (Transgenomic) adjusted automatically as calculated by the Navigator software package (Transgenomic). All samples were run at the oven temperature listed in the Supplementary Table. The Navigator software predicted that each segment of the exonic component of the amplicon would be under partially denaturing conditions at these temperatures. For each abnormal elution profile, genomic

DNA was re-amplified and the PCR products were directly sequenced.

Direct sequencing

The PCR-amplified product was electrophoresed through a 2% agarose gel. The bands of interest were excised from the agarose gel and purified using the QIAquick Gel Extraction Kit (Qiagen, Hilden, Germany). Purified DNA fragments were sequenced in both directions using a BigDye Terminator Cycle Sequencing Kit (Applied Biosystems) and were analyzed on an ABI PRISM 310 Genetic Analyzer (Applied Biosystems). To confirm the mutations, a second independent PCR amplification and subsequent bidirectional sequencing were performed.

Screening for the internal tandem duplication and the D835 mutation of the *FLT3* gene

The *FLT3* internal tandem duplication (ITD) was examined by amplification of the juxtamembrane (JM) domain using forward/reverse primers (Supplementary Table), followed by electrophoresis in an agarose gel, as reported previously.²³ Additional bands, indicative of ITD, were cut out for direct sequencing. Screening for the D835 mutation was carried out by amplification of genomic DNA using forward/reverse primers (Supplementary Table). PCR products were digested with *EcoRV* and were then resolved on an agarose gel, as described previously.²⁴ Direct sequencing was carried out on samples with an undigested band.

Plasmid constructions

PCR-generated fragments of *PTPN11* cDNA encoding SHP-2 or SHP-2 mutants with the C-terminus c-myc epitope tag were subcloned into the pcDNA3.1 expression vector (Invitrogen). The wild-type *PTPN11* cDNA was mutated at nucleotides 181 (G>T) or 226 (G>A) using PCR-based mutagenesis to yield amino-acid changes D61Y or E76K, respectively. The integrity of the amplified sequence was confirmed by DNA sequencing.

Cell culture and transfection

HEL cells were maintained in RPMI1640 (Invitrogen) with 10% fetal calf serum (FCS), 2 mM glutamine, 100 U/ml penicillin and 100 µg/ml streptomycin and grown at 37°C in a humidified atmosphere with 5% CO₂. The plasmids were transfected into HEL cells by electroporation (300 V, 1050 µF) using a Gene Pulser (Bio-Rad, Hercules, CA, USA). Transfected cells were selected with 1 mg/ml G418 (Invitrogen), cloned by limiting dilution, and analyzed for SHP-2 expression by Western blot. Cells were washed with RPMI1640 containing 10% FCS three times, resuspended in the same medium alone or supplemented with 100 ng/ml of stem cell factor (SCF) (PeproTech, Rocky Hill, NJ, USA).

Immunoblotting

For immunoblot analysis, cells were solubilized in NP-40 lysis buffer (50 mM Tris-HCl, pH 8.0; 150 mM NaCl; 1% NP-40) containing a protease inhibitor mixture (Complete; Roche). In order to detect phosphorylation, a phosphatase inhibitor mixture (50 mM sodium fluoride, 10 mM sodium pyrophosphate, 2 mM sodium orthovanadate) was added to the lysis buffer. Protein concentrations were determined with Bradford reagents (Bio-Rad). For Western blot, the lysates were boiled in Laemmli

buffer and then separated by 15% SDS-PAGE gel and transferred to Hybond enhanced chemiluminescence (ECL) (Amersham Pharmacia Biotech). The membrane was blocked in 5% non-fat milk in PBS containing 0.1% Tween-20 and incubated sequentially with primary antibodies and a horseradish peroxidase-conjugated secondary antibody (Amersham Pharmacia Biotech). The primary antibodies used in this study were anti-Myc antibody, anti-c-kit polyclonal antibody, anti-SHP-2 polyclonal antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA), anti-p44/42 MAP kinase polyclonal antibody and anti-phospho-p44/42 MAP kinase polyclonal antibody (Cell Signaling Technology, Beverly, MA, USA). Bound antibodies were detected by ECL using a Western blotting kit (Amersham Pharmacia Biotech).

Results

*The difference of cytogenetic findings in MDS/AML patients with or without an *AML1* point mutation*

We previously reported 32 cases of hematological disease associated with somatic mutations of the *AML1* gene.⁸ We then extended the mutation analysis by PCR-SSCP assay using genomic DNA, and thereby found 13 new cases. The clinical findings of 10 patients with MDS/AML are summarized in Table 1 (cases 33 – 42). Two patients with secondary AML had frame shift and nonsense mutations, and one patient with reactive leukocytosis had the same silent mutation we previously reported in an MDS RA patient (case 1).⁸

We selected the patients according to their diagnosis and *AML1* mutations, and divided them into three disease categories: (A) MDS/AML patients with an *AML1* mutation; (B) MDS/AML patients without an *AML1* mutation; and (C) CBF leukemia patients (8;21 translocation and inversion 16). Two patients (cases 2 and 15) who had silent mutations were excluded from the group of patients with *AML1* mutations because these mutations represent a simple polymorphism, and indeed we found one silent mutation in a reactive leukocytosis patient as well. We analyzed 34 MDS/AML patients with *AML1* mutations summarized in Table 1, including the previously published 24 cases designated as nos. 3–9 and 16–32,⁸ 80 MDS/AML patients without an *AML1* mutation and 25 CBF leukemia patients. This analysis included 28 cases of secondary MDS/AML: 13 with *AML1* mutations (six atomic-bomb survivors, five patients who received previous chemotherapy and/or radiotherapy, one atomic-bomb survivor who also received previous chemotherapy and one case of exposure to mustard gas), and 15 patients without *AML1* mutations (seven atomic-bomb survivors and eight therapy-related patients).

The cytogenetic findings of *AML1*-mutated patients were compared with those of *AML1*-wild-type patients to see whether the *AML1*-mutated patients had distinct cytogenetic characteristics (Table 2). Five previously reported patients who showed a normal karyotype (cases 9, 19, 25, 26 and 29) had minor but clonal karyotypic abnormalities; case 9: 47, XX, +1, der(1;7)(q10;p10), i(21)(q10), case 19: 47, XY, del(7)(q22q32), +8, case 25: 47, XY, dup(1)(q21q44), +8, case 26: 45, XY, -7 and case 29: 44, XX, -7, -17. Clonal cytogenetic abnormalities could be detected in more than half of the patients with MDS/AML, 19 (56%) of the 34 patients with *AML1* mutations and 47 (59%) of the 80 patients without *AML1* mutations. A complex karyotype (five or more aberrations) was seen in 20 (25%) of the 80 patients without an *AML1* mutation, whereas it was not seen in the patients with an *AML1* mutation ($P=0.0007$).

Table 1 Clinical features and mutation characteristics of the MDS/AML patients with AML1 mutations

Case no.	Age (years)/sex	Diagnosis	History of exposure	Chromosome	AML1 mutation	Other mutations
3	69/M	RAEBt	A-bomb (2.7 km)	Others	G42R	
4	59/F	RAEBt	A-bomb (1.7 km)	Others	L71fsX94	N-RAS(G60E) SHP2(N58Y)
5	68/M	RAEBt	A-bomb (0.8 km)	-7, others	D171N	N-RAS(G13D)
6	80/F	AML following MDS	A-bomb (2.5 km)	+8	G42R	
7	80/M	RAEBt	Mustard gas	Normal	Y113X	
8	43/F	RAEBt	AML(M3); Chem	-7	D171N	N-RAS(G12V)
9	39/F	RAEBt	Astrocytoma; Rad/ Chem	7q-	R177Q	
16	74/M	AML following MDS	—	Normal	S114fsX119	
17	65/M	RAEB	—	+8	R130fsX148	
18	47/F	RAEBt	—	Normal	R142fsX151	
19	79/M	RAEBt	—	7q-, +8	G143_K144insRRG	
20	64/F	RAEB	—	Normal	I166T	N-RAS(Q61 K)
21	75/M	RAEBt	—	-7	D171G	FLT3(ITD 7aa) SHP2(F71L)
22	41/M	RAEB	AML(M5b); Chem	Others	T84_L85insP	
23	62/M	RAEBt	Esophageal cancer; Rad	Others	V137_G138insG	
24	62/M	AML following MDS	—	Normal	A224fsX228	SHP2(A72 T)
25	75/M	RAEBt	—	+8	P232fsX567	SHP2(N58Y)
26	73/M	RAEBt	—	-7	D242fsX287	
27	55/F	AML following MDS	—	Normal	S291fsX300	
28	75/M	AML following MDS	—	Normal	R292fsX574	NF1(S382fsX390)
29	70/F	RAEBt	—	-7	T296fsX305	FLT3(ITD 7aa)
30	54/M	RAEB	—	Normal	T296fsX338 T296Sins(36aa)	
31	64/M	RAEB	—	7q-, +8	A364fsX570	
32	87/F	RAEBt	A-bomb (2.0 km)	Normal	L378fsX573	
33	65/F	AML following MDS	—	+21	D171N	
34	83/M	RAEB	—	Normal	G172R	
35	76/M	AML following MDS	—	Others	Q247fsX283	
36	71/F	RAEBt	Thyroid cancer; Chem	Normal	P248fsX283	N-RAS(G13D)
37	80/M	RAEB	—	Normal	S266fsX288	
38	68/M	RAEB	—	Normal	Y322fsX569	NF1(C118fsX164)
39	63/F	RAEBt	A-bomb (2.5 km)	+8	G340fsX576	FLT3(ITD 22aa)
40	64/M	RAEB	—	Normal	Y379fsX573	
41	66/M	AML following MDS	—	Normal	G394fsX562	FLT3(ITD 11aa)
42	72/M	RAEB	A-bomb (1.1 km) AML(M5); Chem	7q-, +8	R396_P398del	

AML = acute myeloid leukemia; MDS = myelodysplastic syndrome; RAEB = refractory anemia with excess blast; RAEBt = RAEB in transformation. A-bomb indicates atomic bomb, and distances in parentheses indicate how far from the center of the explosion the patient was. Chem = chemotherapy; Rad = radiation therapy.

Most chromosomal aberrations were unbalanced numeric alterations, including partial and complete chromosome loss or chromosome gain, in both groups of patients. -5/5q- was found with a significantly higher frequency in the AML1-wild-type group compared with the AML1-mutated group (17/80 versus 0/34, $P=0.0027$), but no significant association was observed between AML1 mutations and -7/7q-, 20q- or +8. However, half of the patients with a complex karyotype had more than 10 aberrations, and many of them had both -5/5q- and -7/7q- abnormalities. Therefore, we compared the incidence of -7/7q- with a normal chromosome 5 between AML1-mutated patients and AML1-wild-type patients (Table 2). This 'simple' -7/7q- abnormality was seen in nine (26%) of the 34 patients with AML1 mutations, and in three (4%) of the 80 patients without AML1 mutations ($P=0.0009$). Of the four patients with AML1-mutated MDS/AML (cases 8, 9, 22 and 42) who had a history of treatment with alkylating agents, three patients presented the -7/7q- cytogenetic abnormalities.

-7/7q- and N-RAS mutation are closely related to AML1 mutations in patients developing MDS after chemotherapy for AML

We identified two patients who developed MDS/AML with AML1 mutations after receiving chemotherapy for AML (Figure 1). At the point of AML diagnosis, they did not show AML1 mutations. Several years after they achieved complete remission, they developed MDS with AML1 mutations. Interestingly, the -7/7q- chromosomal abnormality and the N-RAS mutation also developed at the time of MDS diagnosis. One patient (case 8) had a history of receiving chemotherapy, including alkylating agents and all-trans retinoic acid, for AML (M3). This patient had a normal karyotype and a PML-RAR α oncogenic chimeric gene. Continuous complete remission (CCR) was achieved for 3 years, and then she developed MDS (RAEBt). No mutation was found in either the AML1 or the N-RAS gene before she developed MDS. At the time of MDS diagnosis, she had a karyotypic abnormality of monosomy 7. Moreover, she gained not only an AML1 mutation but also an

Table 2 Cytogenetic abnormalities in MDS/AML patients with or without AML1 point mutations

Karyotype	With AML1 mutation (N = 34)	Without AML1 mutation (N = 80)	P-value*
Normal	15	33	0.8371
Simple karyotype ^a	19	27	0.0369
Complex karyotype ^b	0	20	0.0007
-7/7q- with normal chromosome 5	9	3	0.0009
-5/5q-	0	17	0.0027
20q-	1	7	0.4323
+8	7	13	0.5970
21 abnormality	1 ^c	5 ^d	0.6673

*Fisher's exact probability test was used to determine the P-value.

^aLess than five aberrations.

^bFive or more aberrations.

^cA patient with trisomy 21.

^dPatients including one case with +21, one case with +21q and three cases with balanced translocations with the break point of 21q22.

N-RAS mutation, and lost the PML-RAR α chimeric gene. The other patient (case 42) had a history of atomic bomb exposure. He received chemotherapy including alkylating agents for AML (M4) with a t(2;11) karyotype. Continuous complete remission was achieved for 10 years, during which time he showed a normal karyotype. Then he developed MDS (RAEB) with an AML1 mutation with a karyotypic abnormality of 7q- and +8. At the time of MDS diagnosis, the t(2;11) clone was not detected.

Genetic abnormalities in MDS/AML patients with AML1 point mutation

To investigate the gene mutations in patients with MDS/AML or CBF leukemia, we analyzed the N-RAS, K-RAS, PTPN11, NF1, FLT3, c-KIT and p53 genes (Table 3). We found a total of nine N-RAS mutations, four PTPN11 mutations, two NF1 mutations, eight FLT3 mutations, six c-KIT mutations and 12 p53 mutations. Two patients with an AML1 mutation had two additional mutations (N-RAS and PTPN11, PTPN11 and FLT3). No mutation of the K-RAS gene was detected in these patients with MDS/AML or CBF leukemia. Activating RAS mutations, loss-of-function mutations in NF1 and gain-of-function PTPN11 mutations imply that there is upregulation of the RAS signaling pathway. Furthermore, activating mutations of c-KIT and FLT3 also induce activation of the RAS signaling pathway. N-RAS, PTPN11, NF1 and FLT3 mutations were more frequent in AML1-mutated MDS/AML patients compared with AML1-wild-type MDS/AML patients (15 versus 3.8%, 12 versus 0%, 6.5 versus 0%, and 12 versus 2.5%, respectively), but not significant except for PTPN11 ($P=0.0069$). However, the frequency of all these mutations affecting the RTK-RAS signaling pathway was significantly higher in AML1-mutated MDS/AML patients than in AML1-wild-type MDS/AML patients (38 versus 6.3%, $P<0.0001$). Conversely, p53 mutations were not detected in the 34 patients with AML1 mutations, but were detected in 12 (15%) of the 80 patients without AML1 mutations ($P=0.0170$). Mutations in c-KIT were frequently detected in CBF leukemia patients, but were not detected in any of the patients with MDS/AML. Furthermore, similar to MDS/AML patients with AML1 point mutations, a high frequency (nine of 25, 36%) of RTK-RAS

signaling pathway mutations was observed in patients with CBF leukemia.

SHP-2 mutants contribute to prolonged and enhanced extracellular signal-regulated kinase activation following stem cell factor stimulation

Somatic mutations in PTPN11 are known to upregulate SHP-2 physiologic activation, leading to RAS pathway activation. This gain of function in mutated SHP-2 depends on ligand stimulation. Epidermal growth factor (EGF),^{25,26} and granulocyte-macrophage colony stimulating factor (GM-CSF)²⁷ are known as ligands that upregulate SHP-2, but so far SCF has not been known as such as a ligand. In AML1-mutated MDS/AML patients, genes involved in the RTK-RAS signaling pathway were frequently mutated, and blast cells of these patients expressed surface c-KIT (data not shown). Therefore, we suspected that cells bearing SHP-2 mutants would send strong stimulation signals through the RTKs, especially c-KIT. To investigate the effect of the mutant SHP-2 molecules on the RTK-RAS signaling pathway activation in response to SCF, we used HEL cells, which express the SCF receptor protein c-KIT. We established cell lines stably expressing Myc-tagged wild-type or mutant SHP-2 proteins (Figure 2). D61Y and E76K mutant SHP-2 proteins were previously identified in samples from patients with juvenile myelomonocytic leukemia (JMML), and their molecular characteristics have been thoroughly analyzed.^{25,27,28} Cells were stimulated with SCF for various amounts of time, and we examined the activation of a downstream effector of RAS, extracellular signal-regulated kinase (ERK). Levels of phospho-ERK (p-ERK) at baseline were not elevated in cultures transduced with any SHP-2 constructs compared with parental HEL cells (compare lane 1 with lane 8 in Figure 2b). The HEL cells expressing wild-type SHP-2 showed activation of ERK at 5 min after SCF stimulation, a maximum activation level at 10 min, and p-ERK levels returned to baseline after 30 min. In cells expressing mutant SHP-2, SCF stimulation induced a prolonged and more intense signal of p-ERK lasting up to 60 min (Figure 2b, c). Although previous reports showed that the D61Y mutant was less hypersensitive than E76K in response to GM-CSF and IL-3,^{27,28} our data showed that hyperactivity to SCF stimulation was detected slightly more in the D61Y-expressing cells than in the E76K-expressing cells (Figure 2b, c). The discrepancies between these studies and our data might be due to differences in the cytokine-dependent or independent cell systems and/or stimulating cytokines. The mechanism of different reactivities between these mutants is unclear, but may be caused by a minute structural difference.

Discussion

In this study, we have demonstrated correlations between AML1 point mutations and -7/7q- or gene mutations affecting the RAS signaling pathway in patients with MDS/AML (i.e., RAEB, RAEBt and AML following MDS). This correlation was also shown in the clinical course of two patients who developed therapy-related MDS/AML with a combination of an AML1 mutation, a -7/7q- abnormality and an N-RAS mutation induced by chemotherapy for AML (Figure 1).

Our study revealed a significant association between AML1 mutations and -7/7q-, and between normal AML1 and -5/5q- in both sporadic and secondary cases of MDS/AML patients. Previous studies have shown that the genetic pathways for development of MDS/AML may be classified by the cytogenetic




Figure 1 Clinical course of the patients developing MDS after chemotherapy for AML. (a) Case 8 was a female diagnosed as AML (M3) at the age of 38. She was treated with enocitabine, daunorubicin, vincristine, 6-mercaptopurine, prednisolone, mitoxantrone, etoposide, cyclophosphamide, aclalubichin, methotrexate and all-*trans* retinoic acid for a year and a half, leading to CCR. She developed MDS (RAEBt) at the age of 43. (b) Case 42 was a male diagnosed as AML (M4) at the age of 62. He was treated with enocitabine, daunorubicin, vincristine, 6-mercaptopurine, prednisolone, mitoxantrone, etoposide, cyclophosphamide, aclalubichin and methotrexate for two years, leading to CCR. He developed MDS (RAEB) at the age of 72. CR indicates complete remission; Chem Tx, chemotherapy; ATRA, all-*trans* retinoic acid.

Table 3 Frequency of genetic abnormalities in patients with MDS/AML or CBF leukemia

Genes	MDS/AML patients		P*	CBF leukemia patients (N = 25)
	With AML1 mutation (N = 34)	Without AML1 mutation (N = 80)		
N-RAS	5	3	0.0501	1
K-RAS	0	0	1	0
PTPN11	4	0	0.0069	0
NF1	2	0	0.0871	0
FLT3	4	2	0.0638	2
c-KIT	0	0	1	6
RTK-RAS pathway	13 (38%)	5 (6.3%)	<0.0001	9 (36%)
p53	0	12 (15%)	0.0170	0

RTK = receptor tyrosine kinase. *Fisher's exact probability test was used to determine the *P*-value. *N-RAS* mutations were G12V, G13D, G60E, Q61K and Q61H; *PTPN11* mutations were N58Y, F71L and A72T; *NF1* mutations were C118fsX164 and S382fsX390; *FLT3* mutations were tandem duplications ranged in size from 7 to 24 amino acids and D835Y; *c-KIT* mutations were TYD417_419FG, D816A, D816Y and D816V; and p53 mutations were Y126C, S127C, V143M, R175H, H179Y, H214R, Y234C, R248Q, Y236X and C275F.




Figure 2 Effects of wild-type and mutant SHP-2 proteins on ERK activation in response to SCF. (a) HEL cells were transfected with an expression vector containing either wild-type SHP-2 or a mutated SHP-2 construct and stimulated with SCF (100 ng/ml) for 0, 5 or 10 min. The equivalent expression levels of wild-type, D61Y or E76K SHP-2 protein in total cell lysates were detected by immunoblot analysis with an anti-Myc antibody (Myc). Cell lysates were immunoblotted with an anti-p-ERK specific antibody to monitor ERK activation, or with an anti-ERK antibody (ERK) to confirm equivalent levels of ERK expression. (b) SHP2-overexpressed HEL cells were stimulated with SCF for up to 90 min. Parental HEL cells (labeled P) were also analyzed as a control. (c) The relative p-ERK/ERK ratio compared with parental HEL cells is presented.

abnormalities or by the patient's previous exposure to chemotherapy. Two main cytogenetic pathways have been proposed to explain the malignant transformation step in

patients with therapy-related MDS (t-MDS):²⁹⁻³¹ pathway I is '-7/7q-' with normal chromosome 5', and pathway II is '-5/5q-'. The patients belonging to the pathway I group frequently show mutations of RAS genes³² and methylation of the *p15^{INK4b}* gene promoter.³³ A more recent study has shown that AML1 mutations in patients with t-MDS were highly significantly associated with -7/7q-.⁵ On the other hand, patients belonging to pathway II frequently present a complex karyotype and a mutation of *p53*, whereas pathway I has a very low frequency of *p53* mutations.^{22,34} However, we have focused on the pivotal role of AML1 mutations, rather than cytogenetic abnormalities, during the development of MDS/AML. We are trying to define the disease entity of 'MDS/AML with AML1 mutation', which could be considered for inclusion in the recurrent genetic abnormalities under the WHO classification. Thus, we have attempted to classify the genetic pathway of developing MDS/AML with or without AML1 mutations.

Many reports have shown that genes involved in the RAS signaling pathway are mutated in hematopoietic diseases. In MDS, mutations of the *N-RAS* gene have been known to be a frequent (~10%) genetic alteration,³⁵ and these typically occur during transformation to AML. However, mutations of *PTPN11* have been reported to be rare in adult MDS (2 of 189 MDS/AML, 1.1%),³⁶⁻³⁸ although they have been frequently found in childhood MDS.²⁵ Mutations of the *NF1* gene are also very rare in adult MDS.³⁹ Mutations of the *FLT3* gene are found in 5% of MDS patients, and are associated with leukemic transformation.^{40,41} In this study, we found a significantly higher mutation frequency of these genes in MDS/AML patients with AML1 mutations than in those without an AML1 mutation (38 versus 6.3%, $P < 0.0001$). The high frequency of these mutations in patients with AML1 mutations leads us to consider that mutations in the RAS pathway, besides being valuable deciding factors for prognosis, may function also as genetic partners of the AML1 mutations. Mutations of *RAS*, *NF1* or *PTPN11* are seen in most patients with JMML, a MPD of young children.²⁸ All mouse models bearing a heterozygous *Nf1* (*Nf1*^{+/+}),⁴² a conditional inactivation of *Nf1* (*Nf1*^{-/-}),⁴³ a conditional expression of activating *K-ras* mutation (*K-ras*^{G12D/+}),^{44,45} or a heterozygous *Ptpn11* mutation (*Ptpn11*^{D61G/+})⁴⁶ developed a myeloproliferative disease. Thus, we suggest that these gene alterations might work as 'proliferative' partners of the AML1 mutations.

Which ligand/receptor system would be involved in stimulation of this pathway? Most previous studies of gene mutations affecting the RAS pathway have revealed the hypersensitive proliferation of hematopoietic progenitor cells by using a stimulating ligand of GM-CSF, especially in the studies of JMML.^{44,46,47} However, GM-CSF seems not to be essential for proliferation of hematopoietic stem cells expressing CD34 in patients with MDS. CD34-positive hematopoietic stem cells express the c-KIT receptor protein, and AML1 is required for the generation of these cells.⁴⁸ Moreover, c-KIT mutations and/or overexpression are frequently seen in patients with CBF leukemia in which AML1 functions are impaired (Table 3).^{11,49,50} These gain-of-function mutations of c-KIT result not only in constitutive activation but also in receptor hyperactivation in response to SCF stimulation.⁵¹ In the patients with AML1-mutated MDS/AML, a disease characterized by impaired AML1 function that is similar to CBF leukemia, we did not find any c-KIT mutation. However, blast cells of these patients expressed c-KIT, and SCF stimulation of c-KIT expressing HEL cells bearing SHP-2 mutants yielded hyperactivation of the RAS pathway (Figure 2). Taken together, these results suggest that SCF stimulation via the c-KIT receptor might be important




Figure 3 Genetic pathways in MDS/AML and in CBF leukemia.

for activating the RAS pathway in *AML1*-mutated MDS/AML patients, and hyperactivation of the RTK- (especially c-KIT-) RAS pathway may be a common genetic event in hematological malignancies with impaired *AML1*. In further support of this hypothesis, the *AML1*-ETO chimeric protein represses *NF1* expression.⁵² The somatic mutations associated with some additional factors in the RTK-RAS pathway still remain unknown. Genes encoding these factors also might be altered in hematological diseases characterized by *AML1* impairment. To clarify the cooperative functions of *AML1* mutations and hyperactivation of the RAS pathway in development of MDS/AML, mice carrying both gene mutations are currently being established in our laboratory, and we will address these questions in future studies.

Here, we propose two genetic pathways in the development of MDS/AML (Figure 3). One is a pathway 'with *AML1* mutation'. In this pathway, *AML1* mutations, similar to chimeras in *de novo* AML, would be considered to be the major decision factor to develop MDS/AML and would inhibit the differentiation of hematopoietic stem cells. A hematopoietic stem cell that has acquired *AML1* gene mutation apparently requires a long latency period to be transformed and to be conferred a proliferative and/or survival advantage by gene alterations belonging to the RTK-RAS signal transduction pathways, leading to development of MDS/AML. During this process, loss of the long arm of chromosome 7, the critical region at 7q22.1 possibly related to DNA repair genes,⁵³ would also play an important role in this pathway. It is suspected that some more patients with MDS/AML who have an *AML1* haploinsufficiency so far undetected (such as micro-deletion of 21q22, methylation of the promoter, etc.) still exist, and they should be included in this pathway. The other pathway is one without an *AML1* mutation. In this pathway, the master genetic events that inhibit the differentiation of hematopoietic stem cells are still unknown. However, patients without *AML1* mutation frequently show the 5q- chromosomal abnormality. A recent paper showed that mice carrying heterozygous *Nucleophosmin 1* gene, which maps to 5q35 in human, developed a hematological syndrome with features of human MDS.⁵⁴ Haploinsufficiency of nucleophosmin might be one of the master genetic events of the

pathogenesis of MDS/AML in this pathway. Furthermore, *p53* gene alterations are considered to work as a genetic partner leading to development of MDS/AML in this pathway. Identification of a relevant genetic pathway in MDS/AML pathogenesis may lead us to develop new therapies based on a clearer understanding of disease biology.

Acknowledgements

This work was supported in part by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Osaka Cancer Research Foundation, and the Tsuchiya Foundation. We are grateful to Ryoko Matsumoto-Yamaguchi for excellent technical support in carrying out these experiments. We also thank Dr Ross Fisher and Dr Hideaki Nakajima for critical reading of the article.

References

- Osato M, Asou N, Abdalla E, Hoshino K, Yamasaki H, Okubo T *et al*. Biallelic and heterozygous point mutations in the runt domain of the *AML1/PEBP2alphaB* gene associated with myeloblastic leukemias. *Blood* 1999; **93**: 1817-1824.
- Preudhomme C, Warot-Loze D, Roumier C, Gardel-Duflos N, Garand R, Lai JL *et al*. High incidence of biallelic point mutations in the Runt domain of the *AML1/PEBP2 alpha B* gene in Mo acute myeloid leukemia and in myeloid malignancies with acquired trisomy 21. *Blood* 2000; **96**: 2862-2869.
- Silva FP, Morolli B, Storlazzi CT, Anelli L, Wessels H, Bezroukove V *et al*. Identification of *RUNX1/AML1* as a classical tumor suppressor gene. *Oncogene* 2003; **22**: 538-547.
- Harada H, Harada Y, Tanaka H, Kimura A, Inaba T. Implications of somatic mutations in the *AML1* gene in radiation-associated and therapy-related myelodysplastic syndrome/acute myeloid leukemia. *Blood* 2003; **101**: 673-680.
- Christiansen DH, Andersen MK, Pedersen-Bjergaard J. Mutations of *AML1* are common in therapy-related myelodysplasia following therapy with alkylating agents and are significantly associated with deletion or loss of chromosome arm 7q and with subsequent leukemic transformation. *Blood* 2004; **104**: 1474-1481.
- Song WJ, Sullivan MG, Legare RD, Hutchings S, Tan X, Kufria D *et al*. Haploinsufficiency of *CBFA2* causes familial thrombocytopenia

- with propensity to develop acute myelogenous leukaemia. *Nat Genet* 1999; **23**: 166–175.
- 7 Osato M. Point mutations in the RUNX1/AML1 gene: another actor in RUNX leukemia. *Oncogene* 2004; **23**: 4284–4296.
 - 8 Harada H, Harada Y, Niimi H, Kyo T, Kimura A, Inaba T. High incidence of somatic mutations in the AML1/RUNX1 gene in myelodysplastic syndrome and low blast percentage myeloid leukemia with myelodysplasia. *Blood* 2004; **103**: 2316–2324.
 - 9 Ichikawa M, Asai T, Saito T, Yamamoto G, Seo S, Yamazaki I et al. AML-1 is required for megakaryocytic maturation and lymphocytic differentiation, but not for maintenance of hematopoietic stem cells in adult hematopoiesis. *Nat Med* 2004; **10**: 299–304.
 - 10 Higuchi M, O'Brien D, Kumaravelu P, Lenny N, Yeoh EJ, Downing JR. Expression of a conditional AML1-ETO oncogene bypasses embryonic lethality and establishes a murine model of human t(8;21) acute myeloid leukemia. *Cancer Cell* 2002; **1**: 63–74.
 - 11 Beghini A, Peterlongo P, Ripamonti CB, Larizza L, Cairoli R, Morra E et al. C-kit mutations in core binding factor leukemias. *Blood* 2000; **95**: 726–727.
 - 12 Gilliland DG. Hematologic malignancies. *Curr Opin Hematol* 2001; **8**: 189–191.
 - 13 Matsuno N, Osato M, Yamashita N, Yanagida M, Nanri T, Fukushima T et al. Dual mutations in the AML1 and FLT3 genes are associated with leukemogenesis in acute myeloblastic leukemia of the M0 subtype. *Leukemia* 2003; **17**: 2492–2499.
 - 14 Fenaux P. Chromosome and molecular abnormalities in myelodysplastic syndromes. *Int J Hematol* 2001; **73**: 429–437.
 - 15 Hirai H. Molecular mechanisms of myelodysplastic syndrome. *Jpn J Clin Oncol* 2003; **33**: 153–160.
 - 16 Rosenfeld C, List A. A hypothesis for the pathogenesis of myelodysplastic syndromes: implications for new therapies. *Leukemia* 2000; **14**: 2–8.
 - 17 Mitelman F (ed). *An International System for Human Cytogenetic Nomenclature (1995): Recommendations of the International Standing Committee on Human Cytogenetic Nomenclature*. Basel, Switzerland: S Karger, 1995.
 - 18 Bowen DT, Frew ME, Rollinson S, Roddam PL, Dring A, Smith MT et al. CYP1A1*2B (Val) allele is overrepresented in a subgroup of acute myeloid leukemia patients with poor-risk karyotype associated with NRAS mutation, but not associated with FLT3 internal tandem duplication. *Blood* 2003; **101**: 2770–2774.
 - 19 Pullarkat VA, Bueso-Ramos C, Lai R, Kroft S, Wilson CS, Pullarkat ST et al. Systemic mastocytosis with associated clonal hematological non-mast-cell lineage disease: analysis of clinicopathologic features and activating c-kit mutations. *Am J Hematol* 2003; **73**: 12–17.
 - 20 Tartaglia M, Kalidas K, Shaw A, Song X, Musat DL, van der Burgt I et al. PTPN11 mutations in Noonan syndrome: molecular spectrum, genotype-phenotype correlation, and phenotypic heterogeneity. *Am J Hum Genet* 2002; **70**: 1555–1563.
 - 21 De Luca A, Buccino A, Gianni D, Mangino M, Giustini S, Richetta A et al. NF1 gene analysis based on DHPLC. *Hum Mutat* 2003; **21**: 171–172.
 - 22 Christiansen DH, Andersen MK, Pedersen-Bjergaard J. Mutations with loss of heterozygosity of p53 are common in therapy-related myelodysplasia and acute myeloid leukemia after exposure to alkylating agents and significantly associated with deletion or loss of 5q, a complex karyotype, and a poor prognosis. *J Clin Oncol* 2001; **19**: 1405–1413.
 - 23 Kiyoi H, Naoe T, Nakano Y, Yokota S, Minami S, Miyawaki S et al. Prognostic implication of FLT3 and N-RAS gene mutations in acute myeloid leukemia. *Blood* 1999; **93**: 3074–3080.
 - 24 Yamamoto Y, Kiyoi H, Nakano Y, Suzuki R, Kodaera Y, Miyawaki S et al. Activating mutation of D835 within the activation loop of FLT3 in human hematologic malignancies. *Blood* 2001; **97**: 2434–2439.
 - 25 Tartaglia M, Niemeyer CM, Fragale A, Song X, Buechner J, Jung A et al. Somatic mutations in PTPN11 in juvenile myelomonocytic leukemia, myelodysplastic syndromes and acute myeloid leukemia. *Nat Genet* 2003; **34**: 148–150.
 - 26 Fragale A, Tartaglia M, Wu J, Gelb BD. Noonan syndrome-associated SHP2/PTPN11 mutants cause EGF-dependent prolonged GAB1 binding and sustained ERK2/MAPK1 activation. *Hum Mutat* 2004; **23**: 267–277.
 - 27 Chan RJ, Leedy MB, Munugalavadla V, Voorhorst CS, Li Y, Yu M et al. Human somatic PTPN11 mutations induce hematopoietic cell hypersensitivity to granulocyte-macrophage colony stimulating factor. *Blood* 2005; **105**: 3737–3742.
 - 28 Loh ML, Vattikuti S, Schubert S, Reynolds MG, Carlson E, Lieu KH et al. Mutations in PTPN11 implicate the SHP-2 phosphatase in leukemogenesis. *Blood* 2004; **103**: 2325–2331.
 - 29 Pedersen-Bjergaard J, Andersen MK, Christiansen DH, Nerlov C. Genetic pathways in therapy-related myelodysplasia and acute myeloid leukemia. *Blood* 2002; **99**: 1909–1912.
 - 30 Pedersen-Bjergaard J, Christiansen DH, Andersen MK, Skovby F. Causality of myelodysplasia and acute myeloid leukemia and their genetic abnormalities. *Leukemia* 2002; **16**: 2177–2184.
 - 31 Qian Z, Fernald AA, Godley LA, Larson RA, Le Beau MM. Expression profiling of CD34+ hematopoietic stem/progenitor cells reveals distinct subtypes of therapy-related acute myeloid leukemia. *Proc Natl Acad Sci USA* 2002; **99**: 14925–14930.
 - 32 Stephenson J, Lizhen H, Mufti GJ. Possible co-existence of RAS activation and monosomy 7 in the leukaemic transformation of myelodysplastic syndromes. *Leuk Res* 1995; **19**: 741–748.
 - 33 Christiansen DH, Andersen MK, Pedersen-Bjergaard J. Methylation of p15INK4B is common, is associated with deletion of genes on chromosome arm 7q and predicts a poor prognosis in therapy-related myelodysplasia and acute myeloid leukemia. *Leukemia* 2003; **17**: 1813–1819.
 - 34 Horiike S, Misawa S, Kaneko H, Sasai Y, Kobayashi M, Fujii H et al. Distinct genetic involvement of the TP53 gene in therapy-related leukemia and myelodysplasia with chromosomal losses of Nos 5 and/or 7 and its possible relationship to replication error phenotype. *Leukemia* 1999; **13**: 1235–1242.
 - 35 Hirai H, Kobayashi Y, Mano H, Hagiwara K, Maru Y, Omine M et al. A point mutation at codon 13 of the N-ras oncogene in myelodysplastic syndrome. *Nature* 1987; **327**: 430–432.
 - 36 Johan MF, Bowen DT, Frew ME, Goodeve AC, Wilson GA, Peake IR et al. Mutations in PTPN11 are uncommon in adult myelodysplastic syndromes and acute myeloid leukaemia. *Br J Haematol* 2004; **124**: 843–844.
 - 37 Watkins F, Fidler C, Boulwood J, Wainscoat JS. Mutations in PTPN11 are rare in adult myelodysplastic syndromes and acute myeloid leukemia. *Am J Hematol* 2004; **76**: 417.
 - 38 Loh ML, Martinelli S, Cordeddu V, Reynolds MG, Vattikuti S, Lee CM et al. Acquired PTPN11 mutations occur rarely in adult patients with myelodysplastic syndromes and chronic myelomonocytic leukemia. *Leuk Res* 2005; **29**: 459–462.
 - 39 Kaneko H, Horiike S, Nakai H, Ueda Y, Nakao M, Hirakawa K et al. Neurofibromatosis 1 gene (NF1) mutation is a rare genetic event in myelodysplastic syndrome regardless of the disease progression. *Int J Hematol* 1995; **61**: 113–116.
 - 40 Yokota S, Kiyoi H, Nakao M, Iwai T, Misawa S, Okuda T et al. Internal tandem duplication of the FLT3 gene is preferentially seen in acute myeloid leukemia and myelodysplastic syndrome among various hematological malignancies. A study on a large series of patients and cell lines. *Leukemia* 1997; **11**: 1605–1609.
 - 41 Horiike S, Yokota S, Nakao M, Iwai T, Sasai Y, Kaneko H et al. Tandem duplications of the FLT3 receptor gene are associated with leukemic transformation of myelodysplasia. *Leukemia* 1997; **11**: 1442–1446.
 - 42 Jacks T, Shih TS, Schmitt EM, Bronson RT, Bernards A, Weinberg RA. Tumour predisposition in mice heterozygous for a targeted mutation in Nf1. *Nat Genet* 1994; **7**: 353–361.
 - 43 Le DT, Kong N, Zhu Y, Lauchle JO, Aiyigari A, Braun BS et al. Somatic inactivation of Nf1 in hematopoietic cells results in a progressive myeloproliferative disorder. *Blood* 2004; **103**: 4243–4250.
 - 44 Braun BS, Tuveson DA, Kong N, Le DT, Kogan SC, Rozmus J et al. Somatic activation of oncogenic Kras in hematopoietic cells initiates a rapidly fatal myeloproliferative disorder. *Proc Natl Acad Sci USA* 2004; **101**: 597–602.
 - 45 Chan IT, Kutok JL, Williams IR, Cohen S, Kelly L, Shigematsu H et al. Conditional expression of oncogenic K-ras from its endogenous promoter induces a myeloproliferative disease. *J Clin Invest* 2004; **113**: 528–538.
 - 46 Araki T, Mohi MG, Ismat FA, Bronson RT, Williams IR, Kutok JL et al. Mouse model of Noonan syndrome reveals cell type- and