

| | | | | | |
|--|---|-------------------------------|-----|---------|------|
| Kawanishi S. | capsaicin, a principal ingredient of hot chili pepper. | | | | |
| Kawanishi S., Murata M. | Mechanism of DNA damage induced by bromate differs from general types of oxidative stress. | Toxicology | 221 | 172-178 | 2006 |
| Abe M, Suzuki N, Yoshida M, Igarashi M, Usuda K, Furukawa S, Maekawa A, Juneja LR, Okubo T and Nakae D, | Preliminary evaluation of toxicologic and carcinogenic risks of copper gluconate in rats given multiple carcinogens. | J Toxicol Pathol, | 19 | 129-135 | 2006 |
| Kuribayashi, M., Asamoto, M., Suzuki, S., Hokaiwado, N., Ogawa, K. and Shirai, T. | Lack of modification of 2-amino-3,8-dimethylimidazo[4,5-f]quinoxaline (MeIQx) rat hepatocarcinogenesis by caffeine, a CYP1A2 inducer, points to complex counteracting influences. | Cancer Lett. | 232 | 289-299 | 2006 |
| Matsubara T, Noracharttiyapot W, Toriyabe T, Yoshinari K, Nagata K, and Yamazoe Y. | Assessment of human pregnane X receptor-involvement in pesticide-mediated activation of CYP3A4 gene. | <i>Drug Metab Dispos.</i> | 35 | 728-733 | 2007 |