

Fig.1. Immunohistochemistry of estrogen receptor α in the uteri of female offspring at postnatal days (PNDs) 10 (A, C, E) and 14 (B, D, F) in 0 mg/kg (A, B) and 0.006 mg/kg (C, D) and 6 mg/kg (E, F) bisphenol A (BPA)-treated groups. At PND 10 ER α was expressed in stromal cells of the endometrium but not luminal epithelial cells. At PND 14, uterine glands developed into the endometrium, and ER α was expressed in both luminal and glandular epithelium. There were no differences in ER α expression of the uterus and gland genesis at PNDs 10 and 14 among the control and BPA-treated groups.




Fig. 2. Sequential development of persistent estrus in female offspring perinatally exposed to 0, 0.006 or 6 mg/kg bisphenol A (BPA). White, grey or black column indicates offspring treated with BPA at doses 0 mg/kg/day, 0.006 mg/kg/day and 6 mg/kg/day, respectively.

all animals were affected by persistent estrus at 11 months of age (Fig. 2). In endocrine tissues and representative organs such as those of the alimentary, urinary, respiratory and nervous systems, treatment-related lesions were not morphologically detected.

The average numbers and SD values of ova at 8 weeks of age were 13.3 ± 1.0 , 13.0 ± 2.8 and 12.7 ± 1.2 in control, 0.006 and 6 mg/kg BPA-treated

groups, respectively, with no significant differences.

Serum gonadotropin levels for BPA-treated and control rats in the immature period are shown in Fig. 3. During the immature period, serum FSH and LH levels were comparable among the BPA and control groups, the differences at each PND being not significant.

The incidences of uterine preneoplastic and




Fig. 3. Sequential change in gonadotropin levels (A, FSH; B, LH) of female offspring up to puberty in 0 mg/kg (white circle), 0.006 mg/kg bisphenol A (BPA) (grey circle) and 6 mg/kg/day BPA (black circle) groups.

Table 3. Proliferative lesions in the uteri and histopathology of the ovary at 15 months of age

| Dose | 0 mg/kg/day | 0.006 mg/kg/day | 6 mg/kg/day |
|--|----------------------|-----------------|-------------|
| Number of female offspring examined | 24 | 30 | 30 |
| Uterus: | | | |
| Hyperplasia | | | |
| Slight | 4 (21) ^{a)} | 6 (20) | 4 (13) |
| Moderate | 5 (21) | 6 (20) | 14 (47) |
| Severe | 3 (13) | 5 (17) | 5 (17) |
| Endometrial adenocarcinoma | 8 (33) | 10 (33) | 6 (20) |
| Sub-classification of adenocarcinoma | | | |
| Differentiation | | | |
| Well-differentiated | 8 | 10 | 6 |
| Moderately- or poorly-differentiated | 0 | 0 | 0 |
| Invasion | | | |
| Limited to the uterus | 8 | 10 | 6 |
| Invading into the serosa | 0 | 0 | 0 |
| Distant metastasis | 0 | 0 | 0 |
| Ovary: | | | |
| Atrophy with cystic follicles and absence of corpus luteum | 24 | 30 | 29 |

a) Values in parentheses indicate percentage of incidence.

neoplastic lesions at the termination of the experiment are shown in Table 3. There were no significant differences or treatment-related tendencies among the groups. Sub-classification of adenocarcinomas with regard to their differentiation and invasion status also revealed no inter-group variation. Most of the ovaries in all groups showed atrophy with small cystic atretic follicles and an absence of any corpus luteum (Table 3). Various non-neoplastic and neoplastic

lesions including mammary or pituitary tumors were observed of animals in all groups, but again the incidences were not significant among the three groups. In addition, the animals found dead or euthanized when moribund showed no treatment related changes histopathologically.

Serum and tissue concentrations of BPA are shown in Table 4. BPA was detected in all serum and tissues examined in all groups of the present study. In dams, serum BPA was significantly

Table 4. BPA concentration (ppb) in the serum, milk and liver of dams and pups

| Sample | Age of examination | Doses of BPA (ppb) | | | |
|-------------------|---------------------|--------------------|-------------|--------------|--------|
| | | 0 mg/kg | 0.006 mg/kg | 6 mg/kg | |
| Dam: | | | | | |
| Serum | PND 21 of offspring | 3 ± 0 (5) | 4 ± 0 (5) | 11 ± 4* (5) | |
| Milk | PND 10 of offspring | 28 ± 9 (3) | 8 ± 21 (3) | 8 ± 3 (3) | |
| | PND 14 of offspring | 255 ± 78 (4) | 205 ± 7 (3) | 185 ± 50 (4) | |
| Offspring: | | | | | |
| Serum | PND 10 | Females | 4 (6) | 10 (3) | 23 (4) |
| | | Males | 15 (3) | 5 (3) | 7 (3) |
| | PND 14 | Females | 5 (2) | 4 (3) | 3 (4) |
| | | Males | 4 (2) | 5 (6) | 4 (6) |
| | PND 21 | Females | 9 (4) | 3 (4) | 9 (4) |
| | | Males | 14 (2) | 9 (4) | 20 (5) |
| Liver | PND 10 | Females | 13 (6) | 12 (3) | 17 (4) |
| | | Males | 9 (3) | 9 (3) | 14 (3) |
| | PND 14 | Females | 22 (2) | 100 (3) | 18 (4) |
| | | Males | 45 (2) | 14 (6) | 16 (6) |
| | PND 21 | Females | 60 (4) | 70 (4) | 37 (4) |
| | | Males | 69 (2) | 9 (4) | 60 (5) |

PND, post natal day. Values are means ± SD. Values in parentheses are numbers of animals examined. The serum and liver tissues obtained from offspring were pooled for analysis. *p<0.05.

Table 5. Environmental BPA

| Instruments or diet | Concentration of BPA |
|---|-----------------------|
| Fresh tap water | 0 ± 0 ng/ml (3) |
| Drinking water stored in plastic containers | 2.56 ± 2.51 ng/ml (3) |
| Pellet diet (on opening) | 40.06 ± 2.59 ng/g (3) |
| Pellet diet (several days after opening) | 39.70 ± 0.56 ng/g (3) |

Values are means ± SD. Values in parentheses indicate numbers of samples.

elevated in the 6 mg/kg group as compared to the controls. BPA levels in the milk, however, did not significantly vary among the BPA-treated and control groups. In the offspring, there were no differences in BPA levels in the serum and liver among BPA-treated and control animals from PND 10 up to PND 21.

The concentrations of environmental BPA in the animal room specimens are shown in Table 5. BPA was not detected in fresh tap water, but was identified in drinking water stored in the plastic containers. BPA was also detected in fresh pellet diet at levels several times higher than in the stocked water.

Discussion

The present study was performed to investigate

the effects of maternal exposure to low doses of BPA, at levels comparable with human exposure, on growth and development of the female reproductive system and uterine carcinogenesis in Donryu rats. Unappropriate maternal and/or neonatal exposure to estrogens has been well known to exert irreversible influence directly and indirectly on the reproductive system. The typical influences called 'androgenization' are characterized by lowering of the gonadotropin levels in prepuberty, anovulation, polycystic ovary and persistent estrus immediately after vaginal opening, resulting from direct modulation of the hypothalamic-pituitary-gonadal axis [21, 23]. Androgenized uteri showed abnormal development such as inhibition of uterine glandogenesis or abnormal expression of ER α , and these effects were detectable until maturation [20, 29]. In addition, perinatal exposure to high-doses

estrogens or EDCs with estrogenic activity induced a 'delayed' influence with a different phenotype from that of typical androgenization and is probably caused by delayed modulation of the hypothalamic-pituitary-ovarian control system [25–27]. For instance, first 5 days exposure after birth to 100 mg/kg *p-tert*-octylphenol, this dose is estrogenic [42] and extremely higher (about $\times 10^6$) than waste water, caused a 'delayed' influence which was characterized by accelerated appearance of atrophic ovary compared to controls. This was manifested by an early occurring and a long-term continuing persistent estrus status, whereas no abnormalities could be found with regard to growth and development of the reproductive organs and the hypothalamic-pituitary-gonadal control system up to maturation [27]. In the present study, the treatment did not exert any influences on the reproductive ability of the dams and also on the uterine growth and development of BPA-treated offspring with reference to ER α expression, cell proliferating activity and gland genesis in the uterus, estrous cyclicity, vaginal opening and hormonal secretion up to sexual maturation. These results demonstrate no influence of low doses of BPA on the hypothalamic-pituitary-gonadal control system and the reproductive system up to puberty. After maturation, no disruption of ovarian function reflected by vaginal cytology was also noted, indicating no 'delayed' modulation effects on the reproductive system under the present experimental conditions. Vaginal cytology or its morphological feature in the vagina might be useful for assessment of the individual hormone milieu including dysfunction of the hypothalamic-pituitary-ovarian axis, as previously reported [43].

The most striking examples of changes caused by prenatal exposure to EDCs in the reproductive system of humans and rodents are uterine or vaginal cancers [29, 30, 44, 45]. Many studies have also demonstrated the induction of uterine endometrial adenocarcinomas in rats by perinatal treatment with estrogenic compounds [27, 46]. Uterine endometrial adenocarcinoma is one of the most common malignant tumors in women and has increased in number in recent years, although some epidemiological aspects remain unclear [47, 48]. The Donryu strain rat is a high-incidence strain for spontaneous endometrial adenocarcinoma development with aging, and the tumors have morphological and biological similarities to those

found in humans [30, 31]. In this strain, earlier occurrence of ovarian atrophy with cystic atretic follicles but without corpus luteum leads to ovarian hormonal imbalance, resulting in prolonged elevation of the serum estrogen/progesterone ratio and then early onset of persistent estrus [33]. Under such characteristics, spontaneous uterine cancer development is ascribed to the age-dependent modulation of the ovarian hormonal control, as similarly evidenced in humans [49]. We also reported that neonatal exposure of Donryu rats to high-dose *p-tert* octylphenol enhanced uterine carcinogenesis with prolonged persistent estrus status [27]. The present study clearly demonstrated that maternal treatment with low doses BPA did not affect ovarian function manifesting as vaginal cytology throughout the experiment or susceptibility to uterine carcinogenesis. It might be recommended that relatively long-term comprehensive studies of endocrinological and morphological aspects are necessary for determination of perinatal effects of EDCs.

When considering about effects of maternal exposure to low doses of EDCs on the offspring, the subject of most concern is transfer of EDCs from dams to pups through the placenta and/or milk and subsequent modification of toxicokinetics [50], however, the data about transfer of test compounds are quite limited. BPA is known to form its major metabolite, bisphenol A glucuronide in the liver and is excreted very quickly via feces and urine [51–53]. In the present study, serum BPA levels were elevated in the dams given 6 mg/kg BPA, but BPA was not elevated in the milk, serum or liver of offspring. Surprisingly, however, BPA was detected in the serum and tissues of all the animals examined, including the controls. Furthermore, environmental BPA was found in the drinking water and more prominently in the diet. The increase in the serum of dams at 6 mg/kg group might be related to the BPA-treatment; however the influences of biotransfer of 0.006 mg/kg BPA could not be decided in the present study due to the environmental BPA. In the animal room, there were a number of instruments made with BPA such as plastic cages and water containers, and pellet diet and wood chips are packed in plastic wrapping. Therefore, the presence of environmental BPA in the present study is not considered to have been an incidental contamination, and indicates the possibility that

animal studies using rats or mice are always exposed to environmental BPA. While the influence of long-term exposure to environmental BPA on experimental animals remains to be determined, the possibility that major effects on the female reproductive system occur is unlikely, since the present data were similar to those of relevant studies previously reported in rats [10, 54] and our background data [55, 56]. In addition, any abnormalities, which are defined as effects of perinatal exposure to estrogens or EDCs with estrogenic activity on the female reproductive system, or suggested disruption of the hypothalamic-pituitary-gonadal axis were not detected in the present study [10, 20–27, 29].

In conclusion, transplacental and lactational exposure to BPA at levels comparable to human exposure did not exert any adverse influence on the

female reproductive system such as uterine growth and development, and uterine carcinogenesis. Maternal exposure to BPA did not result in appreciable transfer to the offspring, although serum BPA levels in dams treated with 6 mg/kg BPA group were significantly elevated. The situation, however, is complicated, because low doses of environmental BPA were detected in the drinking water and diet.

Acknowledgements

This study was supported by a Grant-in Aid from the Ministry of Health, Labor and Welfare of Japan. We sincerely thank Mr. Murakami K and Mr. Ochiai K for their expert animal technical assistance and Ms. Ichihara H for the histopathology.

References

- Olea N, Pulgar R, Perez P, Olea-Serrano F, Rivas A, Novillo-Fertrell A, Pedraza N, Soto AM, Sonnenschein C. Estrogenicity of resin-based composites and sealants used in dentistry. *Environ Health Perspect* 1991; 104: 298–305.
- Krishnan AV, Stathis P, Permuth SF, Tokes L, Feldman D. Bisphenol-A: an estrogenic substance is released from polycarbonate flasks during autoclaving. *Endocrinol* 1993; 132: 2279–2286.
- Gaido KW, Leonard LS, Lovell S, Gould JC, Babai D, Portier CJ, McDonnell DP. Evaluation of chemicals with endocrine modulating activity in a yeast-based steroid hormone receptor gene transcription assay. *Toxicol Appl Pharmacol* 1997; 143: 205–212.
- Kuiper GG, Lemmen JG, Carlsson B, Corton JC, Safe SH, van der Saag PT, van der Burg B, Gustafsson JA. Interaction of estrogenic chemicals and phytoestrogens with estrogen receptor beta. *Endocrinol* 1998; 139: 4252–4263.
- Ashby J, Tinwell H. Uterotrophic activity of bisphenol A in the immature rats. *Environ Health Perspect* 1998; 106: 719–720.
- Paraconstantinou AD, Umbreit TH, Fisher BR, Goering PL, Lappas NT, Brown KM. Bisphenol A-induced increase in uterine weight and alterations in uterine morphology in ovariectomized B6C3F1 mice: role of the estrogen receptor. *Toxicol Sci* 2000; 56: 332–339.
- Morrissey RE, Lamb JC IV, Morris RW, Chapin RE, Gulati DK, Heindel JJ. Results and evaluations of 48 continuous breeding reproduction studies conducted in mice. *Fund Appl Toxicol* 1989; 13: 747–777.
- Kwon S, Stedman DB, Ilswick RC, Cattley RC, Welsch F. Pubertal development and reproductive functions of Crl:CD BR Sprague-Dawley rats exposed to bisphenol A during prenatal and postnatal development. *Toxicol Sci* 2000; 55: 399–406.
- Morrissey RE, George JJ, Price CJ, Tyl RW, Marr MC, Kimmel CA. The developmental toxicity of bisphenol A in rats and mice. *Fund Appl Toxicol* 1987; 8: 1294–1302.
- Nagao T, Saito Y, Usumi K, Kuwagata M, Imai K. Reproductive function in rats exposed neonatally to bisphenol A and estradiol benzoate. *Reprod Toxicol* 1999; 13: 303–311.
- Fisher JS, Turner KJ, Brown D, Sharpe RM. Effect of neonatal exposure to estrogenic compounds on development of the excurrent ducts of the rat testis through puberty to adulthood. *Environ Health Perspect* 1999; 107: 397–405.
- Nagel SC, vom Saal FS, Thayer KA, Dhar MG, Boechler M, Welshons WV. Relative binding affinity-serum modified access (RBA-SMA) assay predicts the relative in vivo bioactivity of the xenoestrogens bisphenol A and octylphenol. *Environ Health Perspect* 1997; 105: 70–76.
- Sharpe RM, Fisher J, Milar MR, Jobling S, Sumpter JS. Gestational and/or neonatal exposure of rats to environmental estrogenic chemicals results in reduced testis size and daily sperm production in adulthood. *Environ Health Perspect* 1995; 103: 1136–1143.

14. vom Saal FS, Cooke PS, Buchanan DL, Palanza P, Thayer KA, Nagel SC, Parmigiani S, Welshons WV. A physiologically based approach to the study of bisphenol A and other estrogenic chemicals on the size of reproductive organs, daily sperm production, and behavior. *Toxicol Ind Health* 1998; 14: 239-260.
15. Howdeshell KL, Hotchkiss AK, Thayer KA, Vanderbergh JG, vom Saal FS. Exposure to bisphenol A advances puberty. *Nature* 1999; 401: 763-764.
16. Ashby J, Tinwell H, Haseman J. Lack of effects for low dose levels of bisphenol A and diethylstilbestrol on the prostate gland of CF1 mice exposed in utero. *Regul Toxicol Pharmacol* 1999; 30: 156-166.
17. Cagen SZ, Waechter JM Jr, Dimond SS, Breslin WJ, Butala JH, Jekat FW, Joiner RL, Shiotsuka RN, Veenstra GE, Harris LR. Normal reproductive organ development in CF-1 mice following prenatal exposure to bisphenol A. *Toxicol Sci* 1999; 50: 36-44.
18. Cagen SZ, Waechter JM Jr, Dimond SS, Breslin WJ, Butala JH, Jekat FW, Joiner RL, Shiotsuka RN, Veenstra GE, Harris LR. Normal reproductive organ development in Wistar rats exposed to bisphenol A in the drinking water. *Regul Toxicol Pharmacol* 1999; 30: 130-139.
19. Tyl RW, Myers CB, Marr MC, Thomas BF, Keimowitz AR, Brine DR, Veselica MM, Fail PA, Chang TY, Seely JC, Joiner RL, Butala JH, Dimond SS, Cagen SZ, Shiotsuka RN, Stropp GD, Waechter JM. Three-generation reproductive toxicity study of dietary bisphenol A in CD Sprague-Dawley rats. *Toxicol Sci* 2002; 68: 121-146.
20. Branham WS, Sheehan DM, Zehr DR, Medlock KL, Nelson CJ, Ridlon E. Inhibition of rat uterine gland genesis by tamoxifen. *Endocrinol* 1985; 17: 2238-2248.
21. Branham WS, Sheehan DM, Zehr DR, Ridlon E, Nelson CJ. The postnatal ontogeny of rat uterine glands and age-related effects of 17 β -estradiol. *Endocrinol* 1985; 17: 2229-2237.
22. Cheng HC, Johnson DC. Serum estrogens and gonadotropins in developing androgenized and normal female rats. *Neuroendocrinol* 1974; 13: 357-365.
23. Katsuda S, Yoshida M, Watanabe G, Taya K, Maekawa A. Irreversible effects on the reproductive tract of female rats by neonatal exposure to p-tert-octylphenol. *Toxicol Appl Pharmacol* 2000; 165: 217-226.
24. Takasugi N, Bern HA, DeOme KB. Persistent vaginal cornification in mice. *Science* 1962; 138: 438-439.
25. Herath CB, Watanabe G, Katsuda S, Yoshida M, Suzuki A, Taya K. Exposure to neonatal female rats to p-tert-octylphenol disrupts afternoon surges of luteinizing hormone, follicle-stimulating hormone, and prolactin secretion, and interferes with sexual receptive behavior in adulthood. *Biol Reprod* 2001; 64: 1216-1224.
26. MacLusky NJ, Naftolin F. Sexual differentiation of the central nervous system. *Science* 1981; 211: 1294-1303.
27. Yoshida M, Katsuda S, Tanimoto T, Asai S, Nakae D, Kurokawa Y, Taya K, Maekawa A. Induction of different types of uterine adenocarcinomas in Donryu rats due to neonatal exposure to high-dose p-tert-octylphenol for different periods. *Carcinogenesis* 2002; 23: 1745-1750.
28. Yokota H, Tsuruda Y, Maeda M, Oshima Y, Tadokoro H, Nakazono A, Honjo T, Kobayashi K. Effect of bisphenol A on the early life stage in Japanese medaka (*Oryzias Latipes*). *Environ Toxicol Chem* 2000; 19: 1925-1930.
29. Yoshida M, Takenaka A, Katsuda S, Kurokawa Y, Maekawa A. Neonatal exposure to p-tert-octylphenol causes abnormal expression of estrogen receptor α and subsequent alteration of cell proliferating activity in the developing rat uteri. *Toxicol Pathol* 2002; 30: 357-364.
30. Newbold RR, Bullock BC, McLachlan JA. Uterine adenocarcinoma in mice following developmental treatment with estrogens: A model for hormonal carcinogenesis. *Cancer Res* 1990; 50: 7677-7681.
31. Maekawa A, Onodera H, Tanigawa H, Furuta K, Kanno J, Matsuoka C, Ogiu T, Hayashi Y. Spontaneous neoplastic and non-neoplastic lesions in aging Donryu rats. *Gann* 1986; 77: 882-890.
32. Maekawa A, Takahashi M, Ando J, Yoshida M. Uterine carcinogenesis by chemicals/hormones in rodents (review). *J Toxicol Pathol* 1999; 12: 1-11.
33. Nagaoka T, Onodera H, Matsushima Y, Todate A, Shibusatani M, Ogasawara H, Maekawa A. Spontaneous uterine adenocarcinomas in aged rats and their relation to endocrine imbalance. *J Cancer Res Clin Oncol* 1990; 116: 623-628.
34. Katsuda S, Yoshida M, Isagawa S, Asagawa Y, Kuroda H, Watanabe T, Ando J, Takahashi M, Maekawa A. Dose- and treatment duration-dependent effects of p-tert-octylphenol on female rats. *Reprod Toxicol* 2000; 14: 119-126.
35. Kawamura Y, Inoue K, Nakazawa H, Yamada T, Maitani T. Cause of bisphenol A migration from cans for drinks and assessment of improved cans. *Syokuhin Eiseigaku Zasshi* 2001; 42: 13-17 (In Japanese).
36. Taya K, Sasamoto S. Age-related changes in the prepubertal female rats. *Endocrinol Japon* 1988; 35: 349-355.
37. Watanabe G, Taya K, Sasamoto S. Dynamics of ovarian inhibin secretion during the oestrous cycle of the rat. *J Endocrinol* 1990; 126: 151-157.
38. Ando-Lu J, Takahashi M, Imai S, Ishihara R, Kitamura T, Iijima T, Takano S, Nishiyama K,

- Suzuki K, Maekawa A. High-yield induction of endometrial adenocarcinomas in Donryu rats by a single intra-uterine administration of N-ethyl-N'-nitro-N-nitrosoguanidine. *Jpn J Cancer Res* 1994; 85: 789-793.
39. Nagaoka T, Takeuchi M, Onodera H, Matsushima Y, Ando-Lu J, Maekawa A. Sequential observation of spontaneous endometrial adenocarcinoma development in Donryu rats. *Toxicol Pathol* 1994; 22: 261-269.
 40. Pecorelli S, Benedet JL, Creasman WT, Shepher JH. FIGO staging of gynecologic cancer. *Int J Gynecol Obstetrics* 1999; 64: 5-10.
 41. Usuki Y, Kondo M, Oguri H, Kodaira T, Kato I, Yanaihara N. Comparison of bisphenol-A level in human urine measured by ELISA and HPLC. *BIO Clinica* 2000; 15: 145-148 (In Japanese).
 42. White R, Jobling S, Hoare SA, Sumpter JP, Parker MG. Environmental persistent alkylphenolic compounds are estrogenic. *Endocrinol* 1993; 135: 175-182.
 43. Nagaoka T, Takegawa K, Maekawa A. The relationship between endocrine imbalance and alteration of rat vaginal epithelium. *J Toxicol Pathol* 2002; 15: 103-109.
 44. Herbst AL, Ulfelder H, Poskanzer DC. Adenocarcinoma of the vagina. Association of maternal stilbestrol therapy with tumor appearance in young women. *New Eng J Med* 1971; 184: 878-881.
 45. Kitamura T, Nishimura S, Sasahara K, Yoshida M, Ando J, Takahashi M, Shirai T, Maekawa A. Transplacental administration of diethylstilbestrol (DES) causes lesions in female reproductive organs of Donryu rats, including endometrial neoplasia. *Cancer Lett* 1999; 141: 219-228.
 46. Carthew R, Edwards RE, Nolan BM, Martin EA, Heydon RT, White IN, Tucker MJ. Tamoxifen induces endometrial and vaginal cancer in rats in the absence of endometrial hyperplasia. *Carcinogenesis* 2000; 21: 793-797.
 47. Klave G, Heuch I, Ursin G. Reproductive factors and risk of cancer of the uterine corpus: a prospective study. *Cancer Res* 1988; 48: 6217-6221.
 48. Noumoff JS, Faruqi S. Endometrial adenocarcinoma. *Micro Res Tech* 1993; 25: 246-254.
 49. Wood GP, Toronow RC. Endometrial adenocarcinoma and the polycystic ovary syndrome. *Am J Obst Gynecol* 1976; 124: 140-142.
 50. Takahashi O, Oishi S. Disposition of orally administered 2,2-bis(4-hydroxyphenyl)propane (bisphenol A) in pregnant rats and the placental transfer to fetuses. *Environ Health Perspect* 2000; 108: 931-935.
 51. Pottenger LH, Domoradzki JY, Markham DA, Hansen SC, Cagen SZ, Waechter JM Jr. The relative bioavailability and metabolism of bisphenol A in rats is dependent upon the route of administration. *Toxicol Sci* 2000; 54: 3-18.
 52. Snyder RW, Maness SC, Gaido KW, Welsch F, Sumner SCJ, Fennell TR. Metabolism and disposition of bisphenol A in female rats. *Toxicol Appl Pharmacol* 2000; 168: 225-234.
 53. Yoo SD, Shin BS, Kwack SJ, Lee BM, Park KL, Han SY, Kim HS. Pharmacokinetics, disposition and tissue distribution of bisphenol A in rats after intravenous administration. *J Toxicol Environ Health* 2000; 29: 130-131.
 54. Dühler KD, Wuttke W. Serum LH, FSH, prolactin and progesterone from birth to puberty in female and male rats. *Endocrinol* 1974; 94: 1003-1008.
 55. Katsuda S, Yoshida M, Kuroda H, Ando J, Takahashi M, Kurokawa Y, Watanabe G, Taya K, Maekawa A. Uterine adenocarcinoma in N-ethyl-N'-nitro-N-nitrosoguanidine-treated rats with high-dose exposure to p-tert-octylphenol during adulthood. *Jpn J Cancer Res* 2002; 93: 117-124.
 56. Nishiyama K, Ando-Lu J, Nishimura S, Takahashi M, Yoshida M, Sasahara K, Miyajima K, Maekawa A. Initiating and promoting effects of concurrent oral administration of ethylenethiourea and sodium nitrite on uterine endometrial adenocarcinoma development in Donryu rats. *In Vivo* 1998; 12: 363-368.

Review

Toxicologic/carcinogenic Effects of Endocrine Disrupting Chemicals on the Female Genital Organs of Rodents

Akihiko Maekawa¹, Midori Yoshida², Sin-ichi Katsuda³, and Kiyoshi Imai⁴

¹Sasaki Institute, Sasaki Foundation, 2–2 Kanda-Surugadai, Chiyoda-ku, Tokyo 101–0062, Japan

²Department of Pathology, Sasaki Institute, Sasaki Foundation, 2–2 Kanda-Surugadai, Chiyoda-ku, Tokyo 101–0062, Japan

³Department of Biological Safety Research, Japan Food Research Laboratories, 6–11–10 Nagayama, Tama-shi, Tokyo 206–0025, Japan

⁴Biosafety Research Center, Foods, Drugs and Pesticides, 582–2 Shiohinden, Fukude-cho, Iwata-gun, Shizuoka 437–1213, Japan

Abstract: Toxicologic/carcinogenic effects of some representative endocrine disrupting chemicals (EDCs) having estrogenic activity, such as alkylphenols, on the female genital organs of rodents, especially rats, are reviewed and discussed, focusing on our recent research. Neonatal treatment of high-dose p-tert octylphenol (t-OP, 100 mg/kg s.c. injection every other day from postnatal day 1 (PND 1) to PND 15) induced various long-term persistent irreversible effects on the female reproductive system of Donryu rats, such as lower gonadotropin levels at prepuberty, inhibition of uterine gland genesis, persistent estrus and polycystic ovaries. The result indicates that neonatal high-dose treatment of estrogenic EDCs can affect gonadotropin secretion during the developmental period of sexual maturation with direct masculinization of the hypothalamic function. Exposure limited to the first 5 days after birth (PNDs 1–5) to 100 mg/kg t-OP, however, caused delayed influence which was characterized by accelerated appearance of atrophic ovary, manifested by early-occurring and long-term continuing persistent estrus after puberty, whereas no abnormalities could be found with regard to growth and differentiation of the reproductive organs and the hypothalamo-pituitary-gonadal control system up to maturation, the influence being caused by delayed modulation of the hypothalamo-pituitary-gonadal control system. The most notable effect on the female reproductive system when normal cycling rats were exposed to high-doses t-OP for 28 days, was disappearance of the estrous cycle, but no clear changes were detected in other parameters such as uterine weight and morphology. These results indicate that the most serious issue with EDCs is the potential effects of prenatal and/or neonatal exposure on rodents. Well or moderately differentiated adenocarcinomas increased in Donryu rats initiated with N-ethyl-N'-nitro-N-nitrosoguanidine, when high-dose t-OP was given subcutaneously during adulthood. Neonatal exposure for PNDs 1–5 to high-dose t-OP also showed promoting effects on uterine adenocarcinoma development. However, in rats given t-OP for PNDs 1–15, uterine tumor malignancy was clearly increased, although there was no significant alteration in the total incidence of adenocarcinomas. The results are very interesting in consideration of the histogenesis of uterine adenocarcinomas. However, maternal exposure to low doses of EDCs such as nonylphenol and bisphenol A at actual human exposure levels by the oral route showed no effects on growth and development of the female reproductive system or uterine carcinogenesis. These results indicate that dietary exposure to low doses of EDCs might not induce any adverse effects on the female genital system in mammals including humans. (*J Toxicol Pathol* 2004; 17: 69–83)

Key words: endocrine disrupting chemicals (EDCs), toxicity/carcinogenicity, female genital organs, rodents

Introduction

Recently, the possible adverse consequences arising from the release of man-made substances with estrogenic, anti-estrogenic or androgenic properties, so called endocrin

disrupting chemicals (EDCs), into the environment have become an important environmental problem. There is much concern that these EDCs may have the potential to disturb normal sexual differentiation and development in wild life and mammals, including humans, and exert various deleterious effects on many organs, with carcinogenic effects being particularly important in mammals. The genital organs are the obvious target organs of various EDCs, and various toxicologic changes have been reported to be induced in both male and female genital organs of rodents. Unfortunately, however, there is less information

Received: 15 March 2004, Accepted: 12 April 2004

Mailing address: Akihiko Maekawa, Sasaki Institute, Sasaki Foundation, 2–2 Kanda-Surugadai, Chiyoda-ku, Tokyo 101–0062, Japan

TEL: 81-3-3294-3286 FAX: 81-3-3294-3290

E-mail: amaekawa@sasaki.or.jp

on females than males, although many EDCs have estrogenic properties.

In the present report, toxicologic/carcinogenic effects of some representative EDCs, such as alkylphenols, on the female genital organs of rodents, especially rats, are briefly reviewed and discussed from the point of view of extrapolation to humans, mainly focusing on our recent research. In our studies of the effects of EDCs, Donryu rats were mainly used. The Donryu rat is a unique domestic strain having a regular 4-day estrous cycle at the juvenile stage. After 5 months of age, however, persistent estrus appears and increases age-dependently, and a high occurrence of spontaneous uterine adenocarcinomas is observed at about 2 years of age or thereafter (Table 1)¹. In

this rat strain, the early appearance of persistent estrus results in an increase in the estrogen (E2):progesterone(P) ratio (E2:P ratio). In humans, it has been reported that relatively high E2:P values increase the endometrial cancer risk^{2,3}. Using this strain, we recently demonstrated effects of reproduction on uterine carcinogenesis, in line with the known lower risk of uterine adenocarcinoma in multiparous as compared to nulliparous or infertile women. The incidence of spontaneous endometrial adenocarcinomas showed a tendency to decrease in animals having three reproductive experiences, compared to the nulliparous case, although the incidence was not influenced by a single pregnancy⁴. Thus, this rat strain is a good animal model for endometrial adenocarcinoma development due to the imbalance of endogenous steroid hormones, as found in humans. We also succeeded in obtaining high induction of tumors in this strain by single intra-uterine administration of N-ethyl-N'-nitro-N-nitrosoguanidine (ENNG), thereby establishing a two-stage uterine carcinogenesis model (Fig. 1)^{1,5}. Quite recently, Vollmer⁶ reviewed experimental endometrial cancer models, including Donryu rats, and considered them useful for studies on molecular aspects of endometrial cancer and carcinogenesis.

Table 1. Persistent Estrus and Spontaneous Uterine Tumors in Donryu and F344 Rats*

Sequential Changes in Persistent Estrus Incidences in Female Donryu and F344 Rats

| Strain | Incidence (%) | | | | | | |
|--------|---------------|----|----|----|----|----|--------------------|
| | 4 | 5 | 6 | 8 | 10 | 12 | 15 (Months of age) |
| Donryu | 0 | 17 | 32 | 64 | 87 | 88 | 85 |
| F344 | 0 | 0 | 2 | 0 | 6 | 11 | 4 |

Main Spontaneous Uterine Tumors in Donryu and F344 Rats

| Uterine tumors | Incidences (%) | |
|----------------------------|----------------|----------------|
| | Donryu | F344 |
| Mean survival time (weeks) | 108.8 (62-120) | 114.1 (60-131) |
| Endometrial adenocarcinoma | 35 | 1 |
| Endometrial stromal polyp | 1 | 21 |

*: Maekawa *et al.*, J Toxicol Pathol 1999; 12: 1-11.

Classification of EDCs

Various chemicals have been shown to have endocrine disrupting effects not only on wildlife but also mammals including humans. Major representative EDCs are as follows, according to their use, chemical structures and/or chemical characteristics.

1. DDT (1,1,1-trichloro-2,2-bis(4-chlorophenyl)ethane), DDE (1,1'-(dichloro-ethenylidene)bis(4-chlorobenzene)), dieldrin: agricultural chemicals (insecticides) with properties of high-accumulation and resistance to degradation.




Fig. 1. Two-stage rat uterine carcinogenesis model

2. PCBs (polychlorinated biphenyl), PBB (polybrominated biphenyl): industrial chemicals (insulators etc.) which accumulate and are difficult to degrade.
3. DEHP (di(2-ethylhexyl)phthalate), alkylphenols such as nonylphenol and octylphenol, bisphenol A: industrial chemicals widely used as plasticizers or surfactants.
4. Dioxins such as TCDD (2,3,7,8-tetrachlorodibenzo-p-dioxin), dibenzofuran: chemicals naturally produced by dust-incineration.
5. TBT (tributyltin), TBTO (tributyltin oxide): industrial chemicals used for coating of ships' bottom.
6. DBCP (1,2-dibromo-3-chloropropane), atrazine, vinclozolin: agricultural chemicals.
7. DES (diethylstilbestrol), tamoxifen, oral contraceptives: medical drugs.

Development of the Female Genital Organs and Profile of Hormonal Secretions in Rodents

In general, development of the female genital organs in rats is roughly classified into 3 stages, prenatal (embryonic), neonatal/juvenile and adult/aged. The prenatal stage is from the day of fertilization, i.e., day 1 post-coitum till birth (about 20–22 days in rats). The embryonic bipotential gonad develops from mesoderm in the gonadal ridge located on the dorsal coelomic walls. The primordial germ cells and the gonadal ridge are visualized as condensations of cells localized ventral to the mesonephrons by gestational day 12 in rodents. In the embryonic developmental stage, two sets of paired tubular organs develop: the Wolffian ducts and the Mullerian ducts. It is well established that the presence or absence of functioning embryonic testes plays a major role in determining which duct system undergoes further development. In the rat, the critical time period for Mullerian duct development covers days 14–18 of gestation. On day 18, in the absence of testicular hormones including anti-Mullerian hormone (AMH) from Sertoli cells, the Mullerian ducts undergo further development and the Wolffian ducts degenerate.

The female reproductive tract of rodents is immature at birth and the developing uterus undergoes a period of rapid growth and differentiation during the first 2 weeks of postnatal life. In rats, the uterus at birth corresponds developmentally to the fetal uterus at gestation day 100 in human beings⁷. During this period, luminal epithelial cells invaginate into the underlying stroma to form uterine glands⁸. The uterine growth phase in this period coincides with an elevation of serum estradiol levels beginning on postnatal day (PND) 9. Thus, the role of endogenous estrogen (17β -estradiol, E2) and its receptor (ER) are very important for uterine growth and differentiation. In normal rats, ER expression in the uterine epithelium appears at various days from PND 7 to PND 15⁹.

In female rats, serum FSH (follicle-stimulating hormone) levels rise to a peak at PNDs 15–16 followed by an abrupt nadir, while LH (luteinizing hormone) concentrations are high at PNDs 2–10 followed by gradual decline during

sexual maturation; E2 levels also rise to a peak during the first 2 weeks of age^{10,11}. On the other hand, α -fetoprotein, the estrogen-binding protein produced in the liver, is found in very high concentrations for several weeks after birth¹². It is well known that the increase of FSH before puberty is caused by nullification of the negative feedback of estrogen because of the high concentrations of serum α -fetoprotein¹³.

A striking sexual dimorphism in gross morphology of the medial preoptic area (sexually dimorphic nucleus of the preoptic area: SDN-POA) has been recognized in the rat brain¹⁴. The development of this nucleus starts during late fetal life and depends on the hormonal environment at the critical period of sexual differentiation^{14–16}. In genetic males, the relatively high levels of perinatal testosterone are aromatized to estradiol in the nervous cells of SDN-POA and the estrogenic signals may be directly responsible for the increase of SDN-POA volume. In genetic females, in contrast, estrogenic effects on SDN-POA are prevented because estrogen is bound to the serum binding protein, α -fetoprotein, during the late embryonic and early neonatal periods. The female SDN-POA is smaller than that of males as a result of an orchestrated pattern of decreased cell proliferation and/or increased programmed cell death^{17,18}. It is well known that the SDN-POA volume of genetic females becomes larger than normal on perinatal exposure to testosterone or high amounts of some estrogenic compounds¹⁹. Analogues of SDN-POA have also been identified in various animal species such as the gerbil, Guinea pig, ferret, quail and human²⁰. Recently, another sexual dimorphism had been demonstrated in the anteroventral periventricular nucleus of the preoptic area (AVPvN-POA) and the locus coeruleus^{21,22}. The volumes of these are larger in females than males, but a direct correlation with the hormonal environment has not yet been clarified.

In the rat brain, ER α expression is found primarily in ventral midline structures such as bed nucleus of the stria terminalis, hypothalamic medial preoptic area, hypothalamic ventromedial nucleus, hypothalamic arcuate nucleus, septohypothalamic nucleus, septum and central gray area of the midbrain. ER β is similarly distributed in the brain and is additionally detected in the paraventricular nucleus of the hypothalamus and the hippocampus^{23,24}.

After weaning, the first estrous cycle starts at about 36 days of age, and the minimum breeding age is about 84 days of age²⁵. The estrous cycle is characterized by cyclic changes of the epithelial surface in the vagina and the uterus. The estrous cycle in the rat lasts 4–5 days and is divided into proestrus, estrus, metestrus and diestrus. Uterine weights increase with luminal excretion from diestrus to proestrus, showing a peak at proestrus, but decrease at estrus and metestrus.

In normal cycling rats, the E2 and P levels are highest at proestrus, corresponding with the increased uterine weights. Thereafter, the E2 level drops toward estrus and slightly increases again at diestrus. The P value increases slightly at metestrus, although it is low at estrus and diestrus. At

proestrus and diestrus, and especially the former, the luminal and glandular epithelial cells along with stromal cells beneath the luminal epithelium are strongly positive for ER- α mRNA expression. At estrus, the expression is slightly diminished in luminal cells, but is almost completely lacking in glandular cells. At metestrus, positive signals appear again in the latter. In the myometrium, expression is constant in all estrous stages. Thus cell-type specific patterns of ER-mRNA expression characterize the uteri of normal estrous cycling rats²⁶.

In general, the adult stage in rats is from minimum breeding age to maximum age (about 360–450 days), and thereafter the aged stage starts²⁵. In Donryu rats, however, estrous cycle abnormalities increase age-dependently after 26 weeks of age, and almost all animals show persistent estrus at 52 weeks of age. In contrast, vaginal smears of F344 rats indicate a normal estrous cycle until 52 weeks of age²⁷. In our studies, various histological changes such as follicular cysts and atrophic changes such as absence of corpora lutea in the ovary and cornification of epithelium in the vagina in Donryu rats were observed to be linked with persistent estrus, and increased with time, especially after 10 months of age. In F344 rats, in contrast, atrophy of the ovary is observed in only a few animals at 15 months of age. As a result of ovarian atrophy, in Donryu rats, the plasma values of E2 and P, and especially the latter, decrease with age, the E2:P ratio becoming elevated, and the bromodeoxyuridine (BrdU)-labeling indices of uterine endometrial cells are age-dependently increased²⁸. This age-related hormonal imbalance and the constant high level of proliferating activity of epithelial cells are considered to play important roles in high yield development of spontaneous uterine endometrial adenocarcinomas in this rat strain^{1,28}.

Toxicologic and/or Carcinogenic Effects of EDCs on the Female Genital Organs of Rodents

Rodents in the first 2 weeks of postnatal life, termed “a critical point” or “a window of vulnerability”, are very sensitive to exogenous estrogens and androgens including EDCs, because the reproductive tract undergoes rapid growth and differentiation within this period, as mentioned above. Thus, the OECD (Organization for Economic Cooperation and Development) recently proposed the immature rat uterotrophic assay as one of the screening test methods for the detection of estrogenic or anti-estrogenic properties of chemicals²⁹. In studies using adult animals, the ovariectomized (OVX) animal model is also effective for the detection of estrogen agonists, because the effects of endogenous estrogen can be minimized³⁰. In various toxicity studies using adult animals, oral administration has generally been used to assess the toxicity of chemicals, and the OECD has also proposed a new 28-day repeated oral-dosing toxicity test protocol using adult rats, the enhanced OECD TG407 protocol, for the assessment of the toxic effects of EDCs. For the detection of endocrine disrupting activity of direct-acting chemicals, however, other administration routes such

as subcutaneous injection may provide greater sensitivity than oral administration, because this eliminates the direct effects of metabolism of the chemicals during first passage through the liver.

As mechanisms for the biological effects of EDCs on their target organs, their binding to growth factor receptors and arylhydrocarbon (Ah) receptors as well as steroid receptors has been considered to be very important. Furthermore, some chemicals have been shown to have effects on endogenous estrogen metabolism, resulting in disturbance of the hormonal milieu.

Effects of High-doses of EDCs Effects on Growth and Development of Female Reproductive Organs Prenatal and/or Neonatal Exposure

Inappropriate exposure to estrogens and also EDCs in the prenatal and/or neonatal period has been well established to exert irreversible influence directly and indirectly on the female reproductive system^{31,32}. “Androgenization” is characterized by direct modulation of the hypothalamo-pituitary-gonadal control system, resulting in lowering of gonadotropin levels and persistent estrus as an indirect effect, and abnormal uterine/vaginal development and/or growth as direct influences.

Alkylphenolic compounds are derived from biodegradation of nonionic surfactants, alkylphenol ethoxylates, which are widely used as detergents in many industries. Alkylphenol ethoxylates are also broken down in the process of sewage-treatment or in rivers into alkylphenols, such as nonyl or octylphenol (NP or OP), which are well known representative EDCs with weak estrogenic activity, acting via binding to ER. In vitro data indicate that OP has the most potent estrogenic activity of the alkylphenols (approximately 1000 times less estrogenic than E2), although NP is detected with higher levels than OP in the environment.

In our studies of the toxicologic/carcinogenic effects of EDCs on the female genital organs, OP was selected as a representative compound. It has already been reported that neonatal treatment with OP disrupts estrous cyclicity after weaning in female rats³³. We also examined the effects of neonatal exposure to a high dose of p-tert octylphenol (t-OP) on the female genital organs of Donryu rats³⁴, and the results were in line with those of other papers: long-term persistent irreversible effects such as lower gonadotropin levels at prepuberty, inhibition of uterine gland genesis, persistent estrus shown by vaginal cytology, and polycystic ovaries. In our recent study, newborn female pups were injected with 100 mg/kg t-OP subcutaneously within 24 h of birth. Administration was repeated every other day until PND 15 (PNDs 1–15), and animals were observed till PND 77. Histologically, inhibition of uterine gland genesis was apparent during the immature period before weaning. The day of vaginal opening was about 4 days earlier in OP-treated animals than in controls, and after vaginal opening,




Fig. 2. Serum gonadotropins and sex steroid hormones in control and PNDs 1–15 OP-treated rats. Open circles (○), controls; black circles (●), PNDs 1–15 OP-treated. Katsuda *et al.*, *Toxicol Appl Pharmacol* 2000; 165: 217–226.

none of the OP-treated rats showed a regular estrous cycle, and persistent estrus was ultimately observed in these animals. Atrophic and polycystic ovaries without corpora lutea were anovular. In the endometrium, cell-proliferative activity and cell-death were increased and decreased, respectively, and expression of estrogen receptor alpha mRNA was apparent on in situ hybridization. At 8 weeks of age, treated animals exhibited luminal epithelial hyperplasia with overexpression of ER-mRNA. During the immature period, serum FSH and LH levels were consistently lower in OP-treated rats than in controls. In particular, serum FSH levels remained uniformly low. Serum E2 levels demonstrated essentially the same pattern as in controls, being elevated at PND 14, and then falling to low levels. After weaning before sexual maturation, FSH values in treated rats remained low, while those of control animals decreased rapidly and were maintained at the same levels as in the OP-treated case. In contrast, LH levels of treated animals increased after weaning and remained high until the end of the experiment (PND 77). Serum P levels of both OP-treated and control rats were constant, but the level in the former was only half of the latter value (Fig. 2). Serum inhibin levels of OP-treated rats were nearly the same as in controls at PND 28. The results resembled those of male or

androgenized female rats in the secretory pattern of gonadotropins at this age^{10,11}, indicating that neonatal treatment with high-dose t-OP affects gonadotropin secretion during the developmental period of sexual maturation with direct masculinization of hypothalamic function.

In another of our studies, neonatal exposure for the first 2 weeks (PNDs 1–15) to 100 mg/kg t-OP induced an early and enhanced ER expression in the luminal epithelium compared with age-matched controls, and increased proliferating cell nuclear antigen (PCNA) positive cells, though expression in the glandular epithelium was suppressed in relation to inhibited gland-gensis. Therefore neonatal exposure to high doses of EDCs with estrogenic activity can induce abnormal differentiation in the developing rat uteri via abnormal ER expression and subsequent alteration of cell proliferating activity³⁵.

Recently, however, it has been reported that prenatal and/or neonatal exposure to high doses of estrogens or EDCs with estrogenic activity also exerts a “delayed” influence, different from that of typical androgenization. The delayed influence is probably caused by delayed modulation of the hypothalamo-pituitary-ovarian control system³⁶. A number of investigators have described effects of neonatal exposure

Table 2. Uterine Gland Genesis before Puberty in Control and PNDs 1–5 or PNDs 1–15 OP-treated Rats*

| | No. of uterine gland / section (Mean \pm SD) | | |
|--------|--|----------------|------------------|
| | Control | PNDs 1–5 | PNDs 1–15 |
| PND 10 | 0 | 0 | 0 |
| PND 14 | 3.94 \pm 0.5 | 4.05 \pm 1.5 | 0.1 \pm 0.13** |
| PND 21 | 4.58 \pm 0.6 | 5.57 \pm 1.7 | 2.55 \pm 1.5** |
| PND 28 | 6.42 \pm 1.5 | 7.83 \pm 1.3 | 3.14 \pm 2.2** |

*: Yoshida *et al.*, Carcinogenesis 2002; 23: 1745–1750.

** : Significantly different from the control value ($P < 0.05$).

Table 3. Sequential Changes in Incidences of Persistent Estrus in Control and PNDs 1–5 or PNDs 1–15 OP-treated Rats*

| Group | Incidence of persistent estrus (%) | | | | | | | | |
|------------------------|------------------------------------|------|--------|--------|--------|-------|-------|------|--------------------|
| | 1.5 | 2 | 3 | 4 | 5 | 6 | 8 | 10 | 11 (Months of age) |
| Control | 0 | 0 | 0 | 2.6 | 17.9 | 30.8 | 64.1 | 85.7 | 100 |
| OP-treated (PNDs 1–5) | 4.9 | 12.2 | 53.7** | 70.1** | 87.8** | 100** | 100** | 100 | 100 |
| OP-treated (PNDs 1–15) | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 | 100 |

*: Yoshida *et al.*, Carcinogenesis 2002; 23: 1745–1750.

** : Significantly different from the control value ($P < 0.05$).

to EDCs including estrogens or androgens, but information on such delayed effects is limited. In our recent study, exposure after birth to 100 mg/kg t-OP for the first 5 days (PNDs 1–5) caused a “delayed” influence which was characterized by accelerated appearance of atrophic ovary, manifested by early-occurring and long-term continuing persistent estrus, whereas no abnormalities could be found with regard to growth and development of the reproductive organs and the hypothalamo-pituitary-gonadal control system up to maturation³⁷, thus differing from the case of exposure for PNDs 1–15 to the same dose of t-OP³⁴ (Tables 2 and 3). Previously, we confirmed neonatal OP-treatment of 50 mg/kg/day every other day for PNDs 1–15 did not affect estrous cyclicity³⁴, the total administration-dose (400 mg/kg) being higher than that (300 mg/kg) in the PNDs 1–5 study. This result indicates that the differences were due to the treatment period, rather than the total dosing volume.

Postnatal Exposure

Chronic administration of OP to adult male rats causes alteration in hormonal secretions³⁸, and also induces atrophies of the testis and other genital organs³⁹. We therefore tested estrogenic effects of t-OP using adult OVX Donryu rats given daily subcutaneous injections of 6.25, 12.5, 25, 50 or 100 mg/kg for 2 or 14 days. t-OP was detected in serum at doses of 25 mg/kg and above for 2 days and of 12.5 mg/kg and above for 14 days, and uterine weights and luminal epithelial heights were increased dose-dependently. OP-treatment for 2 days caused a dose-related increase in proliferation of uterine luminal, glandular and stromal cells and vaginal epithelial cells, and the effects were fundamentally related to the serum OP levels⁴⁰.

Effects of t-OP on the female reproductive tract of

normal cycling rats were also investigated. F344 and Donryu rats were used, and t-OP was subcutaneously injected for 28 days at similar concentrations to those applied to OVX rats. The most notable changes were disappearance of normal cyclicity in 50 mg/kg or more OP-treated rats and appearance of persistent estrus in the 100 mg/kg group. In rats showing abnormal cyclicity, the uterine morphology deviated from the normal at each estrous stage of cycling rats, and cell proliferation in the endometrium was slightly increased. However, the data for uterine weights, luminal epithelial cell heights and/or numbers of epithelial cells in the endometrium demonstrated only equivocal alteration. In treated rats, the serum E2 levels were decreased with 50 mg/kg of OP or more. Donryu and F344 rats showed similar sensitivity to estrogenic effects of OP, no strain difference being evident. The results indicate that vaginal cytology may be the most sensitive endpoint for the detection of estrogenic activity of potential EDCs in studies using adult female rats⁴¹. It was also demonstrated that vaginal cytology or its morphological features might be very useful in animal toxicity studies for assessment of the individual hormonal milieu including dysfunction of the hypothalamo-pituitary-gonadal control system⁴².

The suitability of the 28-day repeated oral-dosing study for risk assessment of EDCs or strain differences was investigated in adult SD, F344 and Donryu female rats given 60 or 250 (150) mg/kg/day of NP, or 5 or 50 mg/kg/day of atrazine by stomach tube for 28 days. No morphological changes were noted in any reproductive organs of the treated animals, although abnormal estrous cycles were detected in high-dose groups of all strains, without any strain differences⁴³. The results also indicate that vaginal smear is the most sensitive parameter for detection of effects of estrogenic or anti-estrogenic chemicals, when normal

cycling animals are used. Although atrazine is an agrochemical having weak estrogen-antagonistic activity, an anti-estrogenic property was not clear in the study. However, effects were detected in the immature rat uterotrophic assay, in which atrazine alone was not associated with any changes in uterine weight, but co-treatment with atrazine and E2 reduced E2-induced increase of uterine weight⁴⁴.

Effects on Uterine Carcinogenesis

While the etiology of uterine adenocarcinomas in women is still inconclusive, hormones such as estrogens are considered to be of essential importance^{2,3}. The carcinogenic effects on the female genital tract in mammals, including humans, are considered to be one of the most important adverse consequences of EDCs with estrogenic activity. However, there have been only a few reports of unequivocal induction of carcinomas in experimental animals by EDCs, except with diethylstilbestrol (DES), as reviewed previously¹. In humans, the causation of vaginal and uterine cancers by prenatal exposure to DES is a striking example of environmental carcinogenesis⁴⁵. In experimental animals also, the effects of prenatal DES exposure have been studied in rats and mice, as reviewed by Marselos and Tomatis⁴⁶. Vaginal and uterine adenocarcinomas were induced in mice exposed prenatally to DES^{47,48}. In rats following in utero DES exposure, however, mammary and vaginal tumors, rather than uterine tumors, were observed⁴⁶. Thereafter, uterine carcinomas were also induced in Donryu rats by transplacental administration of DES⁴⁹. In the study, interestingly, data for persistent estrus incidence indicate a "delayed" influence in offspring exposed prenatally, similar to our recent report³⁷.

Tamoxifen (TAM) is a non-steroidal anti-estrogen which competes with estrogen for binding to ER. However, its pharmacology is very complex, and both estrogen agonistic and antagonistic properties have been found, depending on the species, age, exposure duration, dose, route and organs in experimental studies⁵⁰. It has been pointed out that the risk of endometrial cancer may be increased in postmenopausal women exposed to TAM for mammary cancer therapy, the agent acting on the uterus as a weak estrogen agonist^{51,52}. In experimental studies using adult rats and mice, however, it has been impossible to cause endometrial cancers by TAM treatment, although endometrial carcinomas were induced in mice treated neonatally⁵³. Also the incidences of uterine and cervical/vaginal cancers increased in rats, in the absence of any estrogen agonistic effect, when tamoxifen was administered orally on days 2–5 after birth⁵⁴. Previously we reported that TAM showed potent anti-estrogenic effects on the adult rat uterus and inhibited the development of endometrial adenocarcinomas in our two-stage uterine carcinogenesis model⁵⁵. In that study, however, the dose levels used might have been high. Quite recently, we also reported that TAM showed promotion, but not progression, effects on mouse

uterine carcinogenesis, so that the influence in the progression stage appears to be different from the estrogen agonism reported for human beings, although TAM did show estrogen agonistic effects in the promotion stage⁵⁶.

In one study, atrazine slightly increased the incidence of endometrial adenocarcinomas in female F344 rats, when given in the diet⁵⁷. Quite recently, however, it was reported that atrazine administered in diet has no modifying effects on uterine carcinogenesis in ICR mice initiated with N-ethyl-N-nitrosourea⁵⁸. Vinclozoline, a pesticide also showing an anti-estrogenic effect, induced uterine adenocarcinomas in female Wistar rats, as well as ovarian sex cord-stromal tumors, when given orally⁵⁹. The carcinogenic mechanisms of these chemicals with anti-estrogenic activity are not clear and further studies are needed to elucidate them.

Dioxin (2,3,7,8-TCDD) is known to exert its modulatory actions through the Ah receptor, and there is experimental evidence suggesting that it can also act in both estrogenic and anti-estrogenic manners, depending on the dose, species, and organ system involved. In rodents, TCDD induces mainly hepatocellular tumors. In addition, in an initiation-promotion study, morphological changes were also noted in both the uterus and the ovary. Although there is no evidence that TCDD can induce tumors in the female genital tract of rodents, it was reported to cause endometriosis in monkeys⁶⁰.

Another interesting example is ethylenethiourea (ETU), a metabolic product of ethylenebisthiocarbamate fungicides such as maneb and zineb, which are also listed as EDCs. ETU itself is a well established carcinogen, inducing thyroid tumors in rats and hepatic and lymphoid tumors in mice. In addition, it reacts with nitrite under acidic conditions in vitro and in vivo to form a mutagenic and carcinogenic compound, N-nitroso ETU⁶¹. Concurrent oral administration of ETU and sodium nitrite is reported to induce uterine endometrial adenocarcinomas in mice⁶². In our two-stage uterine carcinogenesis model using Donryu rats, concurrent oral administration of ETU (80 mg/kg) and sodium nitrite (56 mg/kg) resulted in uterine endometrial carcinomas without initiation by intrauterine administration of ENNG, and also promoted development of the tumors in animals initiated by ENNG, presumably by influencing the hormonal balance⁶³. Both ETU and nitrite are known environmental chemicals which are included in foods. Our confirmation that endometrial adenocarcinomas can be induced in this way in rats as well as mice, thus points to an importance of the oral route of exposure to these chemicals, although the doses used in the study were much higher than those in the diet.

The effects of high-dose t-OP on uterine carcinogenesis were investigated using adult Donryu rats initiated with a single intrauterine treatment of ENNG at 11 weeks of age and exposed thereafter to 100 mg/kg/day t-OP by s.c. injections until 15 months of age. Adult OVX rats were also treated in the same way. t-OP had no effect on the occurrence of persistent estrus in non-OVX rats, although uterotrophic effects were obvious in the OVX case. At the

Table 4. Uterine Adenocarcinomas in ENNG-initiated Rats with Exposure to High-dose OP*

| Group | No. of rats examined | Incidence of endometrial lesions | | | | | | |
|---------------------------|----------------------|----------------------------------|----|-----|-------|-------------------|----|------|
| | | Hyperplasia | | | total | Adenocarcinoma | | |
| | | + | ++ | +++ | | differentiation** | | |
| | | | | | G1 | G2 | G3 | |
| 1. Control | 23 | 2 | 8 | 7 | 4 | 4 | 0 | 0 |
| 2. OP-treated (Adulthood) | 26 | 1 | 8 | 5 | 12*** | 9 (G1 and/or G2) | | |
| 3. Control | 23 | 3 | 7 | 5 | 6 | 6 | 0 | 0 |
| 4. OP-treated (PNDs 1–5) | 28 | 1 | 3 | 5 | 18*** | 17 | 1 | 0 |
| 5. OP-treated (PNDs 1–15) | 22 | 2 | 2 | 1 | 8 | 1*** | 3 | 4*** |

*: Groups 1–2: Katsuda *et al.*, *Jpn J Cancer Res* 2002; 93: 117–124.

Groups 3–5: Yoshida *et al.*, *Carcinogenesis* 2002; 23: 1745–1750.

** : Histological grades of uterine adenocarcinomas by tumor differentiation.

G1: well differentiated; G2: moderately differentiated; G3: poorly differentiated.

***: Significantly different from the control value ($p < 0.05$).

end of the experiment, however, development of uterine adenocarcinomas was significantly increased in animals exposed to t-OP during adulthood, but no tumors developed in OVX rats. This finding suggests that high-dose t-OP has tumor-promoting effects on the ENNG-treated endometrium of rats, possibly due to direct action on the uterus, as indicated by the uterotrophic effect of OP⁶⁴ (Table 4).

Uterine carcinogenesis in Donryu rats treated neonatally with a high-dose of t-OP has also been investigated. Female pups were subcutaneously administered 100 mg/kg/day t-OP every other day for the first 5 days after birth (PNDs 1–5), or the first 2 weeks (PNDs 1–15). Thereafter, they received a single intra-uterine injection of 20 mg/kg ENNG at 11 weeks of age and were observed until 15 months of age. PNDs 1–5 OP-treated animals showed normal development of the female reproductive system, including uterine gland genesis before weaning and normal estrous cycling immediately after vaginal opening. However, the treatment accelerated the occurrence of persistent estrus after 6 weeks of age, and increased the number of well-differentiated uterine adenocarcinomas at the end of the experiment (15 months of age), as compared with controls. This indicates that PNDs 1–5 OP-treatment resulted in delayed modulation of the hypothalamus-pituitary-ovarian hormonal control system, and thus increased the serum E2:P ratio, leading to promotion of uterine carcinoma development. On the other hand, PNDs 1–15 OP-treatment demonstrated immediate and irreversible influences on the control system, called “androgenization”, and induced suppression of uterine gland genesis as well as abnormal uterine development manifested by prolonged persistent estrus immediately after vaginal opening, similar to our previous report³⁴. In addition, at the end of the experiment, uterine tumor malignancy as assessed by morphological and biological properties was clearly increased, although there was no significant alteration in the total incidence of adenocarcinomas. The total incidence of hyperplasias was significantly lowered, probably related to suppression of uterine gland genesis (Table 4). That study

provided evidence that neonatal exposure during PNDs 1–5 or 1–15 to high-dose t-OP enhances uterine carcinogenesis in ENNG-initiated rats, and that the type of uterine tumor is changed by the period of neonatal treatment³⁷.

Concerning the histogenesis of endometrial adenocarcinomas in Donryu rats, the tumors are considered to arise from hyperplasias of the luminal or glandular epithelium, especially the latter²⁸. In humans, it has been pointed out that the presence or absence of hyperplasia as the background is important for the biological behavior of endometrial adenocarcinomas. High-dose OP treatment at PNDs 1–15 induced luminal epithelial hyperplasia in the uteri of rats at 8 weeks of age³⁴, and finally increased development of undifferentiated adenocarcinomas, although the incidence of hyperplasias was decreased³⁷. Carthew *et al.* also reported that tamoxifen induced uterine adenocarcinomas, including biologically malignant examples, in rats in the absence of endometrial hyperplasia, when given on days 2–5 after birth⁵⁴. These results are very interesting in consideration of the histogenesis of uterine adenocarcinomas.

As mentioned above, estrogen and related compounds are reported to increase the risk of endometrial adenocarcinoma development in women. Estrogens occur naturally within the normal body, and are mainly metabolized in the liver by two separate pathways, producing either catechol estrogens (2- or 4-hydroxylated products) or 16 α - or 16 β -hydroxylated products. 2-Hydroxylation of estradiol or estrone to a catechol is a major metabolic pathway, and the catechol estrogens 2-OHE2 and 2-OHE1 have much weaker hormonal potency than their parent hormones, and lack carcinogenic potency when given to adult animals. On the other hand, 4-hydroxyestradiol (4-OHE2) and the two 16 α -hydroxylated forms, 16 α -OHE1 and 16 α -OHE2, retain potent hormonal activity by acting on classical estrogen receptor and also are tumorigenic⁶⁵. In fact, induction of preneoplastic and neoplastic lesions by estrogen and its steroid metabolites (16 steroids) were studied with our two-stage mouse uterine carcinogenesis

model, and 2-OHE1 or 2-OHE2 exerted promoting, but not progressing, effects, while 16 α - and 16 β -OHE1 caused both promotion and progression⁶⁶.

It is known that indole-3-carbinol binds to the Ah receptor, similar to TCDD, and induces cytochrome p450 metabolic enzymes mainly in the liver. It has been reported that this chemical shows a chemopreventive effect on spontaneous endometrial adenocarcinoma development in Donryu rats when given orally, the effect being speculated to be due to enhanced 2-hydroxylation⁶⁷. We also assessed the effect of indole-3-carbinol on uterine carcinogenesis using our two-stage rat uterine carcinogenesis model. Contrary to expectation, however, the incidences of endometrial carcinomas were increased. In rats given indole-3-carbinol, elevated liver weights and centrilobular enlargement of hepatocytes were also observed, the results indicating an effect on estrogen metabolism in the liver, and further studies are now under way, to clarify the discrepancy (Yoshida *et al.* unpublished data).

Effects of Low-doses of EDCs

The concentrations of EDCs including OP in the environment are very low, and the main exposure route is oral, rather than cutaneous, in humans. In general, the toxicokinetics of chemicals including EDCs in animals is known to be influenced by the method of administration. It has been reported that low doses of estrogens and EDCs such as OP might be removed from the blood during the first passage through the liver, when given orally^{68,69}. For risk assessment of EDCs, it is very important to investigate oral dose effects at human exposure levels and thus we have also focused on relatively low doses of OP (t-OP or n-OP) by oral administration. Female Donryu rats initiated by intrauterine administration of ENNG were given diets containing 100 or 1000 ppm t-OP (about 5 or 50 mg/kg/day) or 100 ppm n-OP (about 5 mg/kg/day) from 11 weeks of age to 15 months of age. Although the concentrations are higher than those in the environment, no significant increase in the incidences of uterine adenocarcinomas was observed in any treated group at the end of the experiment, and also there was no difference in tumor malignancy among the groups (Yoshida *et al.* unpublished data).

As detailed above, exposure to high doses of estrogens or EDCs in the fetal or new born period exerts irreversible androgenization of the female reproductive organs, because of heightened sensitivity. In addition, "delayed" influences on these organs may occur after puberty or sexual maturation. Therefore, relatively long-term comprehensive studies on the endocrinological and morphological aspects may be necessary for determination of prenatal and/or neonatal effects of low doses of EDCs regarding toxicity/carcinogenicity in the female genital organs. Low doses of EDCs such as NP or bisphenol A (BPA) were given orally to pregnant rats, and offspring were observed until 15 months of age, to investigate the prenatal and neonatal effects on growth and development of the female reproductive system

and uterine carcinogenesis. In the reproductive toxicity studies reported by others, high doses of NP caused estrogenic effects on pubertal development in male and female rats^{70,71}. However, maternal or neonatal exposure to relatively low doses demonstrated no adverse influence on the reproductive tract⁷². In our study with NP, dams were administered 0.1, 10 and 100 mg/kg daily by gavage from gestation day 2 up to the day before weaning of their offspring. Then, all female pups at 11 weeks of age were administered a single dose of 20 mg/kg ENNG into a uterine horn, and observed until 15 months of age. The low level, 0.1 mg/kg, was selected as a dose relevant to human daily intake (1 mg/kg) of isoflavones, uterotrophic activity of NP being reported to be 10 times stronger than that of daidzein, one of the major isoflavones, and the middle-dose, 10 mg/kg, was selected as near the no observed effect level in a multi-generation reproductive study using rats⁷¹. None of the treated groups demonstrated any alteration in reproductive ability. In their offspring also, uterine growth and development, vaginal opening and hormonal secretion until puberty were not changed and there were no effects on estrous cyclicity and morphology of the reproductive organs after maturation, or on uterine carcinogenesis in animals initiated with ENNG⁷³.

BPA, a volume chemical used in the manufacture of polycarbonate plastics and found in canned foods, lacquered containers and composite dental sealant, is one of the most representative EDCs with weak estrogenic activity, and uterotrophic potential has been demonstrated in the immature rat assay⁷⁴. A study conducted by the National Toxicity Program (NTP) in the USA demonstrated that maternal exposure to high doses of BPA at 0.5 or 1.0% in the diet (approximately daily intakes of 875 and 1750 mg/kg/day) reduced the number of live pups per litter and litters per pair in first generation mice⁷⁵, although pre- and/or postnatal high-dose BPA exposure did not have any apparent adverse effects on pubertal development in female rats or reproductive functions in rats and mice⁷⁶⁻⁷⁸. Recently, however, perinatal treatment with BPA at much lower doses has been described to influence male reproductive organ parameters such as weight of the testis, prostate, preputial gland and epididymis, and the efficiency of sperm production in rodents⁷⁹⁻⁸¹, and neonatal treatment advanced puberty in mice⁸², although there are also some reports of no treatment-related effects at low dose levels when given to pregnant mice and rats⁸³⁻⁸⁵, and to rats in a three-generation reproductive toxicity study⁸⁶. To further assess the risk, we also investigated effects of maternal exposure to low-doses of BPA, including a human exposure-level, on growth and development of the female reproductive system, and also uterine carcinogenesis in Donryu rats. Dams were administered BPA (0, 0.006 and 6 mg/kg/day) daily by gavage from gestation day 2 up to the day before weaning (PND 21). The concentration of 0.006 mg/kg was selected as consistent with the 63 ppb defined as the average daily intake from canned food in human beings, and 6 mg/kg was selected as appropriate to simulate the maximum dose level

Table 5. Proliferative Uterine Endometrial Lesions in Rats Given Low-doses of BPA*

| Dose | Incidence of lesions (%) | | | Adenocarcinoma** |
|-----------------|--------------------------|----|-----|------------------|
| | Hyperplasia | | | |
| | + | ++ | +++ | |
| 0 mg/kg/day | 21 | 21 | 13 | 33 |
| 0.006 mg/kg/day | 20 | 20 | 17 | 33 |
| 6 mg/kg/day | 13 | 47 | 17 | 20 |

*: Yoshida *et al.*, J Reprod Dev 2004; 50: 349–360.

** : All adenocarcinomas in the three groups were well differentiated and limited to the uterus.

(80 ppm) detected in plastic plates⁸⁷. The treatment did not exert any influences on the reproductive system of female offspring in either treated group, in terms of prepubertal uterine growth and gland-genesis, vaginal opening and gonadotropin secretion. After maturation also, no effects were evident with regard to estrous cyclicity, age-matched sequential changes of the reproductive organs, and uterine carcinogenesis until 15 months of age (Table 5). The results demonstrated that maternal exposure to BPA at human exposure-levels did not have any adverse effects on the female reproductive organs of offspring in rats⁸⁸.

For determination of effects of EDCs on offspring by maternal treatment, biotransfer of the chemicals from dam to offspring is crucial, because the impact is fundamentally related to the serum EDC level⁴⁰. However, data for transfer of the test chemical via the placenta or milk to offspring, or toxicokinetics of low-dose EDCs are very limited⁸⁹. In one of our studies, NP at 10 and 100 mg/kg doses was transferred from dams to their offspring via the milk, but the compound could not be detected in their serum or liver⁷³. Furthermore, BPA levels in the milk of dams, and those in the serum and liver in offspring were comparable between control and treated groups, although the serum level of BPA in dams receiving 6 mg/kg was significantly elevated⁸⁸.

Quite recently, it was reported that cadmium has potent estrogen-like activity *in vivo*⁹⁰. Thus, exposure to low-dose cadmium (a single ip injection at a dose of 5 µg/kg) increased uterine net weight accompanied by proliferation of the endometrium, promoted growth and development of the mammary glands, and induced hormone-regulated genes in ovariectomized rats. *In utero* exposure to the metal (i.p. injections of 0.5 or 5 µg/kg on days 12 and 17 gestation) mimicked also the effects of estrogens, and female offspring experienced an earlier onset of puberty and an increase in the epithelial area and the number of terminal end buds in the mammary gland. The amounts of cadmium used in the study were environmentally relevant, because the WHO-recommended Provisional Tolerable Weekly Intake Level is 7 µg/kg/week. Although the administration route was intraperitoneal, not oral, the ability of environmentally relevant amounts of cadmium to mimic the effects of estradiol is very important and the metal may represent a new class of EDC.

Species Differences in Toxicologic/carcinogenic Effects of EDCs, Effects of EDCs based on the Molecular Biology and Extrapolation of the Effects to Humans

It is well known that the toxicokinetics of chemicals in animals are influenced by many factors, including species, strain, sex, age, dosage and/or administration method, as mentioned above. In particular, species differences are very important for the risk assessment in humans. Species differences in occurrence of toxicologic/carcinogenic effects of EDCs may be an indication of variation in endogenous hormonal factors, in addition to susceptibility to exogenous agents. Mice are generally more sensitive to estrogens than rats, and uterine adenocarcinomas can be induced in mice by estrogen alone, but not in rats. In mice, perinatal exposure to estrogens was found to induce ovary-independent proliferation of the vaginal epithelium, which could not be abolished by ovariectomy⁹¹. On the other hand, the vagina in rats neonatally exposed to high-dose t-OP became atrophic immediately after ovariectomy³⁴. Adenocarcinoma development in the ENNG-initiated endometrium of Donryu rats exposed to high-dose t-OP was also ovary-dependent⁶⁴. In mice, however, E2 promoted uterine adenocarcinoma development ovary-independently⁵⁶. Differences in the reaction to estrogens or EDCs with estrogenic activity may help to explain species differences in toxicity/carcinogenicity, though further studies on this point, focusing on metabolism of estrogens or EDCs and localization of ER expression, are needed.

The prenatal and/or neonatal periods are more sensitive to estrogens, and also EDCs, than the adult period. As mentioned above, it has been pointed out that tamoxifen increases the risk of endometrial cancer in women. In rodents, TAM can induce uterine carcinomas when given to newborn animals, but not adults. Similarly, 2-OHE2 has weak estrogenic, but not carcinogenic effects, although 4-OHE2 is a potent estrogenic catechol causing uterine tumors in adult mice. However, both catechols induced tumors when given on days 1–5 of neonatal life, although carcinogenic activity of 4-OHE2 is stronger than that of 2-OHE2⁹². Further studies on age-dependent differences in the mechanism of EDCs' effects on the female genital organs are also needed.

Various methods such as DNA micro-array techniques based on gene expression levels have recently been used for the evaluation of hazardous effects of various chemicals, because they should reveal very early changes. Up till now, however, there have only been a few reports concerning EDCs. The fact that changes in ontogenic expression of ER alpha and not of ER beta occur in the fetal female rat reproductive tract, provides fundamental information critical for clarifying species-specific physiological roles of ER subtypes during fetal development and for investigating the tissue-specific mechanisms underlying prenatal responses to estrogen and E2 agonists⁹³. Genome-wide analysis of early gene expression has furthermore suggested a basis for the




Fig. 3. Hypothesis of the scheme for "androgenization effects" or "delayed modulation effects" on the hypothalamo-pituitary-gonadal system in rats exposed neonatally to high-dose OP.

drastic uterotrophic effects following estrogen administration⁹⁴. Although DNA micro-array techniques presently demonstrate problems with reliability and reproducibility, future precise analysis should facilitate understanding of the mechanisms underlying effects of estrogenic EDCs⁹⁵.

Almost all EDCs exist at only very low concentrations in the environment, but humans may be exposed for long periods. In many animal studies, various toxicologic effects of EDCs on the female genital organs were demonstrated when very high doses were given, but no obvious effects were detected with low-doses. In humans also, low doses may show no adverse effects, because of homeostasis, although there are notable exceptions in animal and human studies. In fish, the ova-testis is known to be a good indicator of estrogenic effects of EDCs in females, but in female rodents, we are still lacking a consensus regarding equivalent reliable endpoint markers. Thus, more comprehensive studies of the endocrinological, morphological and also biomolecular aspects are necessary in animal studies using rodents for extrapolation of EDCs' effects to humans.

Conclusion

It is well known that the prenatal and/or neonatal period is particularly sensitive to various chemicals, including EDCs, in humans and rodents. Inappropriate exposure may exert irreversible influence, resulting in androgenization of the female genital system. In addition, it has also been reported that a delayed influence may be exerted. Neonatal exposure to a high dose of t-OP (100 mg/kg s.c. injection

every other day from PND 1 to PND 15) induced various long-term persistent irreversible effects on the female reproductive system of Donryu rats, such as lower gonadotropin levels at prepuberty, inhibition of uterine gland genesis, persistent estrus shown by vaginal cytology and polycystic ovaries. Neonatal treatment of high-dose EDCs having estrogenic activity can thus affect gonadotropin secretion during the developmental period of sexual maturation with direct masculinization of the hypothalamic function. Abnormal differentiation in the developing rat uteri may be induced via abnormal ER expression and subsequent alteration of cell proliferating activity. However, exposure limited to the first 5 days after birth to 100 mg/kg t-OP caused "delayed" influence which was characterized by accelerated appearance of atrophic ovary, manifested by an early-occurring and long-term continuing persistent estrus status after puberty, whereas no abnormalities could be found with regard to growth and development of the reproductive organs and the hypothalamo-pituitary-gonadal control system up to maturation. The hypothetical scheme for "androgenization effects" or "delayed modulation effects" on the hypothalamo-pituitary-gonadal system in rats exposed neonatally to high-dose OP is shown in Fig. 3.

On the other hand, the most notable effect on the female reproductive system when normal cycling rats were exposed to a high-dose of t-OP for a short time (28 days), was disappearance of the estrous cycle, and no clear changes were detected in other parameters such as uterine weight and morphology. These results indicate that the vaginal smear is the most sensitive parameter for the detection of effects of EDCs in normal cycling rats.

Well or moderately differentiated adenocarcinomas

were increased in Donryu rats initiated by ENNG, when high dose t-OP was given subcutaneously during adulthood. Neonatal exposure to a high dose of t-OP also showed promoting effects on uterine adenocarcinoma development in a two-stage rat uterine carcinogenesis model using Donryu rats, with slight to higher malignancy with more prolonged treatment.

For the risk assessment of EDCs to human health, it is very important to investigate the effects of low doses at actual human exposure levels, because the concentrations of agents, including alkylphenols and BPA, in the environment are very low. In addition, the main exposure route to EDCs is oral, not subcutaneous, in humans. Thus, we have focused on effects of maternal exposure to low doses of EDCs, such as NP and BPA, by the oral route, which have shown no effects on growth and development of the female reproductive system or uterine carcinogenesis. Transfer of low doses of BPA from dams to offspring via the placenta and/or milk was not unequivocal, although NP was transferred when relatively high doses were given.

These results indicate that dietary exposure to low doses of EDCs might not induce any adverse effects on the female genital system in mammals, including humans, because of the effects of homeostasis and clearance from the blood stream on first passage through the liver.

References

1. Maekawa A, Takahashi M, Ando J, and Yoshida M. Uterine carcinogenesis by chemicals/hormones in rodents. *J Toxicol Pathol* 1999; **12**: 1–11.
2. Zeil HK. Estrogen's role in endometrial cancer. *Obstet Gynecol* 1982; **60**: 509–515.
3. Fox H. Endometrial carcinogenesis and its relation to estrogens. *Pathol Res Pract* 1984; **179**: 13–19.
4. Nagaoka T, Takegawa K, Takeuchi M, and Maekawa A. Effects of reproduction on spontaneous development of endometrial adenocarcinomas and mammary tumors in Donryu rats. *Jpn J Cancer Res* 2000; **91**: 375–382.
5. Ando-Lu J, Takahashi M, Imai S, Ishihara R, Kitamura T, Iijima T, Takano S, Nishiyama K, Suzuki K, and Maekawa A. High-yield induction of uterine endometrial adenocarcinomas in Donryu rats by a single intra-uterine administration of N-ethyl-N'-nitro-N-nitrosoguanidine via the vagina. *Jpn J Cancer Res* 1994; **85**: 789–793.
6. Vollmer G. Endometrial cancer: experimental models useful for studies on molecular aspects of endometrial cancer and carcinogenesis. *Endocrine-Related Cancer* 2003; **10**: 23–42.
7. Hoare RM. Comparative developmental aspects of selected organ system. II. Gastrointestinal and urogenital systems. *Environ Health Perspect* 1976; **18**: 61–66.
8. Branham WS, Sheehan DM, Zehr DR, Ridlon E, and Nelson CJ. The postnatal ontogeny of rat uterine glands and age-related effects of 17 β -estradiol. *Endocrinology* 1985; **117**: 2229–2237.
9. Fishman RB, Branham WS, Streck RD, and Sheehan DM. Ontogeny of estrogen receptor messenger ribonucleic acid expression in the postnatal rat uterus. *Biol Reprod* 1996; **55**: 1221–1230.
10. Cheng HC and Johnson DC. Serum estrogens and gonadotropins in developing androgenized and normal female rats. *Neuroendocrinology* 1974; **13**: 357–365.
11. Dussault JH, Walker P, Dubois JD, and Labrie F. The development of the hypothalamo-pituitary axis in the neonatal rat: Sexual maturation in male and female rats assessed by hypothalamic LHRH and pituitary and serum LH and FSH concentrations. *Can J Physiol Pharmacol* 1977; **55**: 1091–1097.
12. Sell S, Nichols M, Becker FF, and Leffert HL. Hepatocyte proliferation and α 1-fetoprotein in pregnant, neonatal, and partially hepatectomized rats. *Cancer Res* 1974; **34**: 865–871.
13. Meijs-Roelofs HM and Kramer P. Maturation of the inhibitory feedback action of oestrogen on follicle-stimulating hormone secretion in the immature female rat: A role for alpha-fetoprotein. *J Endocrinol* 1979; **81**: 199–208.
14. Gorski RA, Gordon JH, Shryne JE, and Southam AM. Evidence for a morphological sex difference within the medial preoptic area of rat brain. *Brain Res* 1978; **148**: 333–346.
15. Jacobson CD, Csernus VJ, Shryne JE, and Gorski RA. The influence of gonadectomy, androgen exposure or a gonadal graft in the neonatal rat on the volume of the sexually dimorphic nucleus of the preoptic area. *J Neurosci* 1981; **1**: 1142–1147.
16. Doehler KD. The pre- and postnatal influence of hormones and neurotransmitters on sexual differentiation of the mammalian hypothalamus. *Int Rev Cytology* 1991; **131**: 1–57.
17. Houtsmuller EJ, Brand T, de Jonge FH, Joosten RN, Van de Poll NE, and Slob AK. SDN-POA volume, sexual behavior and preference of male rats affected by perinatal treatment with ATD. *Physiol Behav* 1994; **56**: 535–541.
18. Davis EC, Popper P, and Gorski RA. The role of apoptosis in sexual differentiation of the rat sexually dimorphic nucleus of the preoptic area. *Brain Res* 1996; **734**: 10–18.
19. Furguson SA, Scallet AC, Flynn KM, Meredith JM, and Schwetz BA. Developmental neurotoxicity of endocrine disruptors: focus on estrogen. *Neurotoxicol* 2000; **21**: 947–956.
20. Doehler KD and Jarzab B. The influence of hormone antagonists on sexual differentiation of the brain. *J Clean Technol Environ Toxicol & Occup Med* 1998; **7**: 195–220.
21. Pinos H, Collado P, Rodriguez-Zafra M, Rodriguez C, Segovia S, and Guillamon A. The development of sex difference in the locus coeruleus of the rats. *Brain Res Bull* 2001; **56**: 73–78.
22. Orikasa C, Kondo Y, Hayashi S, McEwen BS, and Sakuma Y. Sexually dimorphic expression of estrogen receptor β in the anteroventral periventricular nucleus of the rat preoptic area: Implication in luteinizing hormone surge. *Proc Natl Acad Sci USA* 2002; **99**: 3306–3311.
23. Henry WW Jr, Medlock KL, Sheehan DM, and Scallet AC. Detection of estrogen receptor (ER) in the hypothalamus using rat anti-ER monoclonal IgG with the unlabeled antibody method. *Histochem* 1991; **96**: 157–162.
24. Kuiper GGJM, Carlsson B, Grandien K, Enmark E, Haggblad J, Nilsson S, and Gustafsson JA. Comparison of the ligand binding specificity and transcript tissue distribution of estrogen receptors α and β . *Endocrinol* 1997; **138**: 863–870.

25. Del Vecchio FR. Normal development, growth, and aging of the female genital tract. In: Pathology of the Aging Rat (Vol. 1), Mohr U, Dungworth DL and Capen CC (eds), Washington DC: ILSI Press, 331–336, 1992.
26. Katsuda S, Yoshida M, Watanabe T, Kuroda H, Ando-Lu J, Takahashi M, Hayashi H, and Maekawa A. Estrogen receptor mRNA in uteri of normal estrous cycling and ovariectomized rats by in situ hybridization. *Proc Soc Exp Biol Med* 1999; **221**: 207–214.
27. Ando-Lu J, Sasahara K, Nishiyama K, Takano S, Takahashi M, Yoshida M, and Maekawa A. Strain-differences in proliferative activity of uterine endometrial cells in Donryu and Fischer 344 rats. *Exp Toxic Pathol* 1998; **50**: 185–190.
28. Nagaoka T, Takeuchi M, Onodera H, Matsushima Y, Ando-Lu J, and Maekawa A. Sequential observation of spontaneous endometrial adenocarcinoma development in Donryu rats. *Toxicol Pathol* 1994; **22**: 261–269.
29. OECD. OECD guideline and protocol for the prevalidation of the rat uterotrophic screening assay. ENV/JM/TG/EDTA(99)2/ANN1. Joint Meeting of the Chemical Committee and the Working Party on Chemicals, Paris, 1999.
30. Dodge JA, Glasebrook AL, Magee DE, Phillips DL, Sato M, Short LL, and Bryant HU. Environmental estrogens: effects on cholesterol lowering and bone in the ovariectomized rat. *J Steroid Biochem Mol Biol* 1996; **59**: 155–161.
31. Iguchi T and Takasugi N. Occurrence of permanent changes in vaginal and uterine epithelia in mice treated neonatally with progestin, estrogen, and aromatizable or non-aromatizable androgens. *Endocrinol Jpn* 1976; **23**: 327–332.
32. Colborn T and Clement C. Chemically-induced alterations in sexual and functional development: the wildlife/human connection. In: *Advances in Modern Environmental Toxicology*. Colborn T and Clement C (eds), Princeton, NJ: Princeton Scientific Publishing, 1–403, 1992.
33. Blake CA and Ashiru OA. Disruption of rat estrous cyclicity by the environmental estrogen 4-tert-octylphenol. *Proc Soc Exp Biol Med* 1997; **216**: 446–451.
34. Katsuda S, Yoshida M, Watanabe G, Taya K, and Maekawa A. Irreversible effects of neonatal exposure to p-tert-octylphenol on the reproductive tract in female rats. *Toxicol Appl Pharmacol* 2000; **165**: 217–226.
35. Yoshida M, Takenaka A, Katsuda S, Kurokawa Y, and Maekawa A. Neonatal exposure to p-tert-octylphenol causes abnormal expression of estrogen receptor α and subsequent alteration of cell proliferating activity in the developing Donryu rat uterus. *Toxicol Pathol* 2002; **30**: 357–364.
36. MacLusky NJ and Naftolin F. Sexual differentiation of the central nervous system. *Science* 1981; **211**: 1294–1303.
37. Yoshida M, Katsuda S, Tanimoto T, Asai S, Nakae D, Kurokawa Y, Taya K, and Maekawa A. Induction of different types of uterine adenocarcinomas in Donryu rats due to neonatal exposure to high-dose p-t-octylphenol for different periods. *Carcinogenesis* 2002; **23**: 1745–1750.
38. Blake CA and Boockfor FR. Chronic administration of the environmental pollutant 4-tert-octylphenol to adult male rats interferes with the secretion of luteinizing hormone, follicle-stimulating hormone, prolactin, and testosterone. *Biol Reprod* 1997; **57**: 255–266.
39. Boockfor FR and Blake CA. Chronic administration of 4-tert-octylphenol to adult male rats causes shrinkage of the testes and male accessory sex organs, disrupts spermatogenesis, and increases the incidence of sperm deformities. *Biol Reprod* 1997; **57**: 267–277.
40. Katsuda S, Yoshida M, Isagawa S, Asagawa Y, Kuroda H, Watanabe T, Ando J, Takahashi M, and Maekawa A. Dose- and treatment duration-related effects of p-tert-octylphenol on female rats. *Reprod Toxicol* 2000; **14**: 119–126.
41. Yoshida M, Katsuda S, Ando J, Kuroda H, Takahashi M, and Maekawa A. Subcutaneous treatment of p-tert-octylphenol exerts estrogenic activity on the female reproductive tract in normal cycling rats of two different strains. *Toxicol Lett* 2000; **116**: 89–101.
42. Nagaoka T, Takegawa K, and Maekawa A. The relationship between endocrine imbalance and alteration of rat vaginal epithelium. *J Toxicol Pathol* 2002; **15**: 103–109.
43. Aso S, Anai M, Noda S, Imatanaka N, Yamasaki K, and Maekawa A. Twenty-eight-day repeated-dose toxicity studies for detection of weak endocrine disrupting effects of nonylphenol and atrazine in female rats. *J Toxicol Pathol* 2000; **13**: 13–20.
44. Yamasaki K, Sawaki M, Noda S, Muroi T, and Maekawa A. Immature rat uterotrophic assay of diethylstilbestrol, ethynyl estradiol and atrazine. *J Toxicol Pathol* 2000; **13**: 145–149.
45. Herbst AL, Ulfelder H, and Poskanzer DC. Adenocarcinoma of the vagina. Association of maternal stilbestrol therapy with tumor appearance in young women. *N Engl J Med* 1971; **184**: 878–881.
46. Marselos M and Tomatis L. Diethylstilbestrol. II. Pharmacology, toxicology and carcinogenicity in experimental animals. *Eur J Cancer* 1993; **29**: 149–155.
47. Newbold RR and McLachlan JA. Vaginal adenosis and adenocarcinoma in mice exposed prenatally or neonatally to diethylstilbestrol. *Cancer Res* 1982; **42**: 2003–2011.
48. Walker BE. Uterine tumors in old female mice exposed prenatally to diethylstilbestrol. *J Natl Cancer Inst* 1983; **70**: 477–484.
49. Kitamura T, Nishimura S, Sasahara K, Yoshida M, Ando J, Takahashi M, Shirai T, and Maekawa A. Transplacental administration of diethylstilbestrol (DES) causes lesions in female reproductive organs of Donryu rats, including endometrial neoplasia. *Cancer Lett* 1999; **141**: 219–228.
50. Pasqualini JR, Sumida C, and Giambiagi N. Pharmacodynamic and biological effects of anti-estrogens in different models. *J Steroid Biochem* 1988; **31**: 613–643.
51. Killackey MA, Hakes TB, and Pierce VK. Endometrial adenocarcinoma in breast cancer patients receiving antiestrogens. *Cancer Treat Rep* 1985; **69**: 237–238.
52. IARC. Tamoxifen. In: *IARC Monographs on the Evaluation of Carcinogenic Risks to Humans* (Vol. 66), Lyon: IARC, 253–365, 1996.
53. Newbold RR, Jefferson WN, Padilla-Burgos E, and Bullock BC. Uterine carcinoma in mice treated neonatally with tamoxifen. *Carcinogenesis* 1997; **18**: 2293–2298.
54. Carthew P, Edwards RE, Nolan BM, Martin EA, Heydon RT, White INH, and Tucker MJ. Tamoxifen induces endometrial and vaginal cancer in rats in the absence of endometrial hyperplasia. *Carcinogenesis* 2000; **21**: 793–797.
55. Yoshida M, Kudoh K, Katsuda S, Takahashi M, Ando J, and Maekawa A. Inhibitory effects of uterine endometrial carcinogenesis in Donryu rats by tamoxifen. *Cancer Lett* 1998; **134**: 43–51.
56. Takahashi M, Shimomoto M, Miyajima K, Iizuka S, Watanabe T, Yoshida M, Kurokawa Y, and Maekawa A.