

pimarinic acid and dehydroabiatic acid act as BK channel openers via direct interaction with the α -subunit of BK channels.²⁴ Several possible mechanisms can explain the effect of the activators on BK channels: (i) modulation of the gating (=open/close) kinetics of the α subunit, (ii) modulation of the Ca^{2+} binding site at the C-terminal end of the α -subunit, (iii) strengthening of the interaction between the α - and β -subunits, and (iv) mimicking the interaction of β - with the α -binding site. Novel synthetic channel modulators would be useful tools to reveal the mechanism of the channel gating at the atomic level, in addition to having therapeutic potential.

We have already found that pimarinic acid (**2**) and dehydroabiatic acid (**4**) exhibit BK channel opening activities, while the structurally related abiatic acid (**3**) has weak activity (Chart 2).²⁴ In order to explore new structural scaffolds for BK channel openers, we carried out molecular design on the basis of two concepts: (1) introduction of a heteroatom into the dehydroabiatic acid skeleton would allow easier introduction of substituents. (2) Because of the fourfold symmetrical structure of the BK channels, dimeric compounds in which two pharmacophores are linked through a tether may have a greater binding probability to the channels, resulting in increased channel opening activity. Herein we describe our exploration of the hexahydrodibenzazepinone derivatives (**5**), which can be generated from dehydroabiatic


acid (**4**), to examine the value of the hexahydrodibenzazepinone structure as a new scaffold for BK channel openers.

The synthesized monomer compounds were evaluated by means of electrophysiological patch-clamp studies in HEK293 cells expressing BK channels, followed by the Magnus contraction-relaxation assay with rabbit urinary bladder smooth muscle strips to evaluate overall actions on smooth muscle. Dimeric compounds were designed on the basis of the monomeric scaffolds, and their channel opening activities were evaluated by electrophysiological methods. Our results demonstrate that the hexahydrodibenzazepinone structure is a potential new scaffold for novel BK channel openers.


2. Chemistry

2.1. Hexahydrodibenzazepinone derivatives


The hexahydrodibenzazepinone derivatives (**11a** and **b**) were synthesized as shown in Scheme 1. The benzylic oxidation of the dehydroabiatic acid derivatives (**6a** and **b**) was carried out with CrO_3 in $\text{Ac}_2\text{O}/\text{AcOH}$ to give the corresponding ketone compounds (**7a** and **b**). The ketones were converted to the oximes (**8a** and **b**), and the hydroxyl group of the oximes was tosylated. These tosylated oximes (**9a** and **b**) were subjected to the Beck-



Scheme 1.



Scheme 2.



Scheme 4.

the amide NH group with NaH (Scheme 2). Hydrolysis of the methyl ester was carried out in a similar manner to that described for **10**, to give the carboxylic acids (**13a** and **b**). The lactam structure of hexahydrodibenzazepinone derived from the Beckmann rearrangement was reported previously,²⁵ and our ¹H NOE analysis between the N-CH₃ group and the aromatic protons of the compound **12a** confirmed the structure.

The syntheses of the dimeric compounds (**27–30**) are illustrated in Scheme 3. Sonogashira coupling was the key reaction. With respect to the linker moiety, 4-pentyn-1-ol (**14**) was tosylated and converted to the iodide (**16**) with NaI. N-Alkylation was carried out with the iodide **16** to afford the alkyne derivatives (**17a,b**), which were coupled with *m*-diiodobenzene or *p*-diiodobenzene in the presence of Pd(PPh₃)₂Cl₂/CuI in Et₃N at rt (the Sonogashira coupling reaction). The acetylene disubstituted products (**20–22**) and/or the acetylene monosubstituted products (**24–26**) were obtained. In the case of the monosubstituted products, a second Sonogashira coupling reaction was carried out to obtain the acetylene disubstituted products (**20–22**). Finally, the ester group was hydrolyzed with KOH and crown ether in methanol to afford the corresponding dimeric compounds (**27–29**). The alkyne (**17**) and *o*-diiodobenzene did not readily undergo the Sonogashira coupling reaction with Pd(PPh₃)₂Cl₂/CuI in Et₃N at rt. Under these reaction conditions, in addition to the acetylene monosubstituted product (**23**), the acetylene homodimeric product (**18**) was obtained, probably because of steric hindrance. Thus, on coupling of the alkyne (**17a**) with *o*-diiodobenzene, Pd(PPh₃)₄ was used as a catalyst and purified CuI

was used under reflux in Et₃N, to give the desired ortho dimeric compound (**19**). Hydrolysis of the ester provided the final carboxylic acid (**27**). The monomeric compounds (**31** and **33a,b**), which bear an alkynyl substituent on the lactam amide, were also synthesized as shown in Scheme 4.

2.2. Molecular design of dimers

The X-ray structures of several K⁺ channels revealed that the diameters of the pores (the gate region) in the closed state are 6 Å (for the KcsA channel) and 8 Å (KirBac channel), and those in open state are 25 Å (for the MthK channel) and 19 Å (KvAP channel). In the cases of other ion channels, the binding sites of small organic molecules, such as channel blockers, are located on the inner side of the pore.²⁶ Moreover, the dichlorodehydroabiatic acid (**4a**) was shown to activate the BK channel through interaction with the α subunit.^{24c} It seems reasonable to assume that the hexahydrodibenzazepinone derivatives (**13**) also interact with the α subunits of the BK channels.

We thus designed several dimeric compounds, **27–29**, in which two units of the pharmacophore, dichlorohexahydrodibenzazepinone **13a**, are linked through the rigid diacetylenebenzene unit. Because we have little information about the arrangement and topology of the binding sites, we added adjustable molecular-dynamic flexibility by using an *n*-propyl chain as a tether between the nitrogen atom of the hexahydrodibenzazepinone core and the rigid diacetylenyl benzene core. Ortho, meta, and para substitution can change the direction of extension of




Figure 1. Conformational spaces of isomeric dimer compounds energy-minimized structures of the dimers in an *n*-octanol environment.

the two pharmacophores, and this molecular design allowed us to change the distance between the two pharmacophores in the dimer. In fact, molecular dynamic calculations on the basis of conformational search with OPLS2001 force field parameters in an *n*-octanol environment (MacroModel 8.5) generated a range of dimer structures.²⁷ We obtained various numbers of independent structures as follows: ortho (**27**): 2852 structures; meta (**28**): 2581 structures; para (**29**): 2838 structures, when the number of generated trial structures was set to 3000 in each case. The minimum-energy structures in each case are shown in Figure 1. The structural space was not significantly different in the gas-phase or in water as a solvent (data not shown). Because the carboxylic acid group of dehydroabiatic acid **4** is known to be essential for BK channel opening activity,²⁴ this group presumably plays an important role in the interaction with the channel proteins. A similar role of the carboxylic acid group was expected for the hexahydrodibenzazepinone derivatives. The average (with standard deviation) minimum and maximum distances between the two carbonyl oxygen atoms of the individual carboxylic acids in the calculated dimer structures are as follows: ortho (**27**): 6.54 ± 4.02 Å (minimum: 3.01 Å; maximum: 15.37 Å); meta (**28**): 7.04 ± 5.54 Å (minimum: 2.99 Å; maximum: 18.42 Å); para (**29**): 22.23 ± 1.66 Å (minimum: 4.40 Å; maximum: 25.09 Å). The distances between the pharmacophores, particularly those of the carboxylic acids of **27–29**, accord well with the size of the pores of the K^+ channels, which are well conserved among the K^+ channel superfamily. The present designed dimeric compounds should be able to bind to the two putative binding sites of a single BK channel, if the binding sites are located close to the pore regions,

and this should result in an increase of channel opening activity.

3. Results and discussion

3.1. Electrophysiological studies of hexahydrodibenzazepinones derivatives

The activity of the monomeric hexahydrodibenzazepinone derivatives as BK channel openers was evaluated by the patch-clamp technique with inside-out configuration. Human BK channel subunits, hSlo α and β_1 , were transiently expressed in tsA201 cells, a derivative of the human embryonic kidney (HEK) cell line 293. In the patch-clamp measurement, when a compound has channel opening activity, the conductance–voltage curve shifts towards the negative voltage direction (leftward), that is, a larger BK channel current is activated by the same voltage step under the voltage-clamp condition. The BK channel opening activities of the tested compounds at 10 μ M are represented as the shift of $V_{1/2}$ value (mV, $n = 3–5$), the voltage activating the half-maximum current, in comparison with the negative control in the presence of 100 μ M Ca^{2+} on the intracellular side of the membrane patch (Fig. 2). In this work, we used dichlorodehydroabiatic acid **4a** as a reference compound which has a related structure.²⁴

The voltage shift of $V_{1/2}$ of **4a** was -5.3 ± 3.7 mV, and that of the *N*-methyl derivative of the hexahydrodibenzazepinone **13a** was -9.2 ± 3.7 mV. Compound **13a** can open the BK channel, while **13b**, its counterpart lacking the dichloro-substituent, did not show channel




Figure 2. Effects of test compounds on the BK channel current. (A) Effect of 33a on BK channel currents recorded in the inside-out macro patch-clamp configuration. BK channel currents were activated by test pulses to 100 mV from a holding potential at -100 mV. Current zero level is indicated by a dotted line. Outward BK channel currents are shown in upward direction. (B) Conductance–voltage relationship (G – V curve) of BK channel currents ($n = 4$). Compound 33a shifted G – V curve toward hyperpolarized direction. (C) Shifts of $V_{1/2}$ value (mV) of G – V curve by test compounds ($n = 3$ – 5). A negative value indicates a shift to hyperpolarizing voltages.

opening activity. In this context, the results of the smooth muscle relaxation assay are consistent with those of the patch-clamp experiments.

Thus, the results of the electrophysiological study supported the idea that the dichloro-substituted *N*-methyl-hexahydrodibenzazepinone **13a** can serve as a pharmacophore for BK channel openers.

3.2. Electrophysiological study of the dimers and *N*-substituent effects

These dimeric compounds bearing dichloro substituents on the aromatic ring (**27**–**29**) were studied by means of an inside-out patch-clamp configuration. They showed BK channel opening activities, the left-shift of $V_{1/2}$ being -7.7 ± 1.2 mV, -6.3 ± 3.2 mV, and -5.4 ± 1.6 mV, respectively. While the channel opening activity of the ortho derivative **27** is likely to be stronger than those of the other isomers (**28** and **29**), the differences are not significant. In addition, the magnitude of the shift of **27** was as large as that of the monomer **13a**. These results imply that a part of the structures common to these dimeric isomers could be responsible for the activity, rather than the dual binding of the dimeric compounds to multiple binding sites of the BK channel. Thus, we synthesized monomeric fragments of the dimers, that is, **31** and **33a**, the former bearing a *N*-alkyl acetylene substituent and the latter bearing an additional terminal benzene moiety. We examined the BK channel opening activity of these compounds in patch clamp experiments (Fig. 2). Unexpectedly, we found that **33a** showed significant channel opening activity (-23.9 ± 1.4 mV left-shift of $V_{1/2}$), while **31** showed activity (-6.6 ± 1.5 mV) comparable in magnitude with that of the *N*-methyl derivative **13a**.

The prototype *N*-methyl hexahydrodibenzazepinone **13b**, lacking the aromatic dichloro-substituents, did not show BK channel opening activity in patch-clamp experiments (Fig. 2). However, the *m*-dimeric compound **30** showed channel opening activity with a hyperpolarizing shift of $V_{1/2}$ by -11.9 ± 0.9 mV. This activity is also probably derived from the fragment structure of the dimer. In fact, the monomeric compound **33b** also showed a strong opening activity, with a leftward shift of $V_{1/2}$ by -18.9 ± 3.8 mV, which is less than the shift caused by **33a**, but comparable in magnitude. The most potent compound **33a** expanded the width of the tail current in the patch-clamp measurements (data not shown), which suggests that this compound stabilizes the BK channel in the open state.

3.3. Isolated rabbit detrusor smooth muscle assay as overall activity

We evaluated the overall activities of hexahydrodibenzazepine derivatives on detrusor smooth muscle. Figure 3 shows the results of the rabbit detrusor smooth muscle assay. The K^+ channel-dependent muscle dilating effect was evaluated as follows.²⁸ Prior to the application of test compounds, detrusor smooth muscle strips were precontracted by depolarization caused by raising the extracellular K^+ concentration to either 30 mM or 120 mM. With the extracellular K^+ concentration at 30 mM, the small depolarization allows Ca^{2+} influx through Ca^{2+} channels to produce contraction. If a test compound activates the BK channel, the K^+ efflux will shift the membrane potential toward the equilibrium potential for K^+ (E_K) of approximately -40 mV where Ca^{2+} channels are closed, thus causing relaxation of the muscle strip. In contrast, with the extracellular K^+ concentration at 120 mM, the opening




Figure 3. Detrusor smooth muscle relaxation assay. The relative magnitude of relaxation was normalized to the precontraction caused by high K⁺ (30 mM or 120 mM) as 100%. The relaxant activities of the test compounds were compared with that of 0.1% DMSO as the vehicle (control).

of the BK channel will shift the membrane potential toward E_K of around 0 mV, where Ca²⁺ channels are open, and the dilating activity of the BK channel opener should be abolished. The relative magnitude of relaxation was normalized to the contraction caused by high K⁺ precontraction as 100%. The concentration of DMSO as a vehicle was set to be no more than 0.1% (v/v), because DMSO at high concentration caused relaxation by itself. Moreover, the concentration of the test compounds was fixed at 10 μM because of their relatively poor solubility in DMSO–water. The relaxant activities of the test compounds were compared with that of vehicle (0.1% DMSO) measured in the same muscle strip. All values are expressed as means ± SEM. Statistical significance of differences was determined by paired Student's *t*-test, and *P* values less than 0.05 were considered to be significant.

The results of the isolated rabbit detrusor smooth muscle relaxation study of the dehydroabietic acid derivatives were consistent with what we had found in the

previous²⁴ and present electrophysiological studies. The dichloro-substituted **4a** showed a statistically significant relaxation activity, while the unsubstituted **4b** showed weak activity (data not shown). The smooth muscle relaxation study showed that the *N*-methyl derivative **13a**, bearing aromatic dichloro substituents, showed a statistically significant relaxation effect, while the prototype NH amide **11a** showed less potent activity (data not shown). Even in the absence of the aromatic dichloro substituents, the NH derivative **11b** showed distinct relaxation activity, while the *N*-methyl derivative **11b** did not.

The monomeric compounds **31** and **33a** showed distinct rabbit urinary bladder relaxation activities in detrusor smooth muscle strips ($54.3 \pm 2.3\%$ and $48.68 \pm 5.8\%$, respectively, $n = 5$), the activity of **33a** being statistically significant (Fig. 3). Reasonably, the *N*-alkylated monomeric compound **33b** also exhibited potent relaxation activities ($48.68 \pm 5.8\%$, $n = 5$).

4. Conclusion

In this work, we first demonstrated that the hexahydro-dibenzazepinone structure represents a new scaffold for BK channel openers. Second, we carried out molecular design based on our hypothesis that dual binding of linked pharmacophores might provide strong activation even if the single pharmacophore is a rather weak opener. We found that the dimerization retained (in the case of **27**) or increased (in the case of **30**) the channel opening activity as compared with the corresponding monomeric pharmacophores (**13a** and **13b**, respectively). In the present design, ortho, meta, and para isomers (**27–29**) of the dimeric compounds showed similar, not increased, magnitudes of activity as compared with the minimal monomer **13a**, probably because the two pharmacophores cannot bind two putative binding sites at the same time. However, a portion of the molecules clearly can gain access to the binding site. Because of the highly hydrophobic nature of **33** and the dimers **27–30**, and retention of the opening activities of the dimers **27–30**, it is possible that there are hydrophobic binding sites in the transmembrane region of the channels which allow these hydrophobic molecules to gain access in a well-ordered conformation (at least partially) via a membrane bilayer pathway, rather than a direct aqueous pathway.²⁹ Finally, we found that the phenyl-bearing alkynyl substituent on the lactam amide nitrogen atom is crucial for BK channel opening activity. Our findings provide a new basis for development of novel BK channel openers.

5. Experimental

5.1. Chemistry

Commercially available reagents were purchased and were used without further purification unless otherwise stated. ¹H NMR spectra were recorded on a Bruker Avance (400 MHz) spectrometer. The chemical shifts (ppm) were shown by using tetramethylsilane (TMS) as an internal standard in deuterated chloroform (CDCl₃), and the peak of the solvent was used as an internal standard in the case of deuterated methanol (CD₃OD). Mass spectra were recorded on a JEOL MStation JMS-700 spectrometer. Elemental analyses were recorded on a Yanaco CHN CORDER spectrometer and were carried out at the Analysis Center of the Graduate School of Pharmaceutical Sciences, the University of Tokyo. Flash column chromatography was carried out with silica gel (Kanto Chemical 60 N, particle size 40–50 mm). Melting points were measured on a Yanaco hot-stage microscope and are uncorrected. Yields are shown in terms of those of isolated pure materials.

12,14-Dichlorodehydroabiatic acid (4a). To a suspension of dehydroabiatic acid **4b** (2.0 g, 6.66 mmol), 2% w/w FeCl₃ on SiO₂ (1 g), and 1% w/w DDQ on SiO₂ (0.1 g) in 25 mL CCl₄ was added a solution of 2.8 M Cl₂ in 35.7 mL CCl₄ (99.9 mmol, 15 equiv) at 0 °C. The reaction mixture was vigorously stirred at rt for one day in

a sealed tube. Then to this mixture was added a solution of 2.8 M Cl₂ in 10 mL CCl₄ (28.0 mmol, 4 equiv) at rt, and the whole was stirred for 4 days. The insoluble materials were filtered off and the filtrate was then evaporated in vacuum. The residue was recrystallized from *n*-hexane to afford **4a** (574 mg, 1.554 mmol, yield 23%). Compound **4a**: colorless fine needles. Mp 218–222 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.20 (3H, s, CH₃), 1.28 (3H, s, CH₃), 1.39 (6H, d, *J* = 7.3 Hz, 2CH₃CH), 1.61–1.66 (2H, m, 2CH), 1.72–1.86 (5H, m, 5CH), 2.13 (1H, dd, *J* = 12.6 Hz, 2.2 Hz, CH), 2.24 (1H, d, *J* = 11.9 Hz, CH), 2.70–2.79 (1H, m, CH), 2.94 (1H, dd, *J* = 18.4 Hz, 6.0 Hz, CH), 3.91 (1H, br s, CH₃CH), 7.15 (1H, s, ArH). MS (FAB⁺) *m/z* 369 ([M(³⁵Cl₂)+H]⁺), 371 ([M(³⁵Cl³⁷Cl)+H]⁺), 373 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₀H₂₆Cl₂O₂: C, 65.04; H, 7.10. Found: C, 64.95; H, 7.16.

12,14-Dichlorodehydroabiatic acid methyl ester (6a). To a solution of 12,14-dichlorodehydroabiatic acid **4a** (442 mg, 1.197 mmol) in MeOH (2 mL) and PhMe (4 mL) was dropwise added 2.0 M TMSCHN₂ in Et₂O (0.78 mL, 1.557 mmol, 1.3 equiv) at rt over 5 min, and the whole was stirred at rt for 30 min. Excess TMSCHN₂ was quenched with Et₂O, then the reaction mixture was evaporated in vacuum, and the residue was purified by flash chromatography (*n*-hexane only to *n*-hexane/AcOEt = 10:1) to afford **6a** (453 mg, 1.181 mol, yield 99%) as a colorless solid. **6a**: colorless needles. Mp 126–129 °C. ¹H NMR (CDCl₃) 1.19 (3H, s, CH₃), 1.26 (3H, s, CH₃), 1.39 (6H, d, *J* = 7.1 Hz, 2CH₃CH), 1.47–1.53 (2H, m, 2CH), 1.65–1.75 (5H, m, 5CH), 2.12 (1H, d, *J* = 10.6 Hz, CH), 2.23 (1H, d, *J* = 11.7 Hz, CH), 2.71–2.73 (1H, m, CH), 2.92 (1H, did, *J* = 18.3 Hz, 6.2 Hz, CH), 3.67 (3H, s, CO₂CH₃), 3.91 (1H, br s, CH₃CH), 7.15 (1H, s). LRMS (FAB⁺) *m/z* 383 ([M(³⁵Cl₂)+H]⁺), 385 ([M(³⁵Cl³⁷Cl)+H]⁺), 387 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₁H₂₈Cl₂O₂: C, 65.79; H, 7.36. Found: C, 65.86; H, 7.14.

Dehydroabiatic acid methyl ester (6b). A mixture of dehydroabiatic acid **4b** (5.1 g, 0.017 mol) and concd H₂SO₄ (6 mL) in MeOH (200 mL) was stirred at 85 °C for 19 h. Then concd H₂SO₄ (2 mL) was carefully added to the reaction mixture. Stirring was continued for 2 days, then the mixture was cooled, the solvent was evaporated in vacuum, and the residue was diluted with water (300 mL). The whole was extracted with CHCl₃ (1 × 200 mL, 2 × 100 mL). The combined organic layer was washed with brine (1 × 100 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 10:1) to afford **6b** (4.4 g, 0.014 mol, yield 83%) as a colorless solid. Compound **6b**: colorless needles. Mp 45–49 °C (*n*-hexane). ¹H NMR (CDCl₃) 1.21–1.23 (9H, m, 2CH₃CH and CH₃), 1.27 (3H, s, CH₃), 1.42 (1H, m, CH), 1.50 (1H, m, CH), 1.61–1.70 (5H, m, 5CH₂), 2.24 (1H, dd, *J* = 12.5 Hz, 2.1 Hz, CH), 2.30 (1H, d, *J* = 12.3 Hz, CH), 2.80–2.84 (1H, m, CH₃CH), 2.86–2.90 (1H, m, CH), 3.66 (3H, s, COCH₃), 6.88 (1H, s, ArH), 6.99 (1H, d, *J* = 6.4 Hz, ArH), 7.16 (1H, d, *J* = 8.1 Hz, ArH). LRMS (FAB⁺) *m/z* 315

([M+H]⁺). Anal. Calcd for C₂₁H₃₀O₂: C, 80.21; H, 9.62. Found: C, 80.07; H, 9.82.

13-Isopropyl-12,14-dichloro-7-oxopodocarpe-8,11,13-triene-15-carboxylic acid methyl ester (7a). To a solution of CrO₃ (287 mg, 2.867 mmol, 1.1 equiv) in Ac₂O (9 mL) and AcOH (4 mL) was dropwise added a suspension of the methyl ester **6a** (999 mg, 2.606 mmol) in AcOH (15 mL) at 0 °C over 10 min. The reaction mixture was stirred at 50 °C for 9 h, the whole was cooled, and poured into ice-water (40 mL) and then the whole was extracted with CHCl₃ (3 × 30 mL). The combined organic layer was washed with water (1 × 30 mL), satd NaHCO₃ (2 × 30 mL), and brine (1 × 30 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **7a** (847 mg, 2.130 mmol, yield 82%) as a pale yellow amorphous solid. Compound **7c**: ¹H NMR (CDCl₃) 1.17 (3H, s, CH₃), 1.34 (3H, s, CH₃), 1.41 (6H, d, *J* = 7.3 Hz, 2CH₃CH), 1.75–1.79 (5H, m, 5CH), 2.20 (1H, d, *J* = 12.5 Hz, CH), 2.51 (1H, dd, *J* = 16.9 Hz, 5.2 Hz, CH), 2.64 (2H, m, 2CH), 3.66 (3H, s, CO₂CH₃), 3.94 (1H, br s, CH₃CH), 7.19 (1H, br s, ArH).

12,14-Dichloro-13-isopropyl-7-hydroxyiminepodocarpe-8,11,13-triene-15-carboxylic acid methyl ester (8a). A mixture of the methyl ester **7a** (198 mg, 0.498 mmol), pyridine (0.06 mL), and NH₂OH·HCl (54 mg, 0.772 mmol, 1.6 equiv) in EtOH (2 mL) was stirred at 100 °C for 3.5 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **8a** (207 mg, quantitative yield) as a colorless amorphous solid. Compound **8a**: ¹H NMR (CDCl₃) 1.06 (3H, s, CH₃), 1.22 (6H, d, *J* = 8.4 Hz, 2CH₃CH), 1.26 (1H, dd, *J* = 18.7 Hz, 6.4 Hz, CH), 1.72–1.77 (5H, m, 5CH), 2.15 (2H, dd, *J* = 12.1 Hz, 6.1 Hz, CH), 2.38 (1H, dd, *J* = 18.7 Hz, 6.4 Hz, CH), 3.09 (1H, dd, *J* = 18.7 Hz, 13.0 Hz, CH), 3.66 (3H, s, CO₂CH₃), 3.69 (1H, br s, CH₃CH), 7.15 (1H, s, ArH).

12,14-Dichloro-13-isopropyl-7-tosyloxyiminepodocarpe-8,11,13-triene-15-carboxylic acid methyl ester (9a). To a solution of the methyl ester **8a** (207 mg, 0.503 mmol) in pyridine (1 mL) was added TsCl (144 mg, 0.754 mmol, 1.5 equiv) at rt. The reaction mixture was stirred for 17 h at rt, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **9a** (282 mg, 0.497 mmol, yield 99%) as a colorless oil. Compound **9a**: ¹H NMR (CDCl₃) 1.03 (3H, s, CH₃), 1.25 (3H, d, *J* = 7.1 Hz, CH₃CH), 1.27 (3H, d, *J* = 7.1 Hz, CH₃CH), 1.37 (3H, s, CH₃), 1.74–1.77 (5H, m, 5CH), 2.06–2.15 (2H, m, 2CH), 2.46 (3H, s, CH₃), 2.51 (1H, dd, *J* = 13.8 Hz, 5.0 Hz, CH), 3.02 (1H, dd, *J* = 19.0 Hz, 12.9 Hz, CH), 3.67 (3H, s, CO₂CH₃), 3.94 (1H, br s, CH₃CH), 7.12 (1H, br s, ArH), 7.35 (2H, d, *J* = 8.4 Hz, ArH), 7.94 (2H, d, *J* = 8.2 Hz, ArH).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid methyl ester (10a). The mixture of the methyl ester **9a** (282 mg, 0.497 mmol) in TFA (2 mL) was stirred at rt for 1 h, and then the solvent was evaporated in vacuum.

The residue was purified by flash chromatography (*n*-hexane/AcOEt = 1:1) to afford **10a** (166 mg, 0.403 mmol, yield 81%) as a colorless amorphous solid. Compound **10a**: ¹H NMR (CDCl₃) 1.42 (6H, d, *J* = 7.1 Hz, 2CH₃CH), 1.45 (3H, s, CH₃), 1.58 (3H, s, CH₃), 1.67–1.76 (2H, m, 2CH), 1.81–1.83 (2H, m, 2CH), 1.90–1.93 (2H, m, 2CH), 2.29 (1H, d, *J* = 8.6 Hz, CH), 2.92 (1H, dd, *J* = 15.9 Hz, 6.8 Hz, CH), 3.47–3.55 (1H, m, CH), 3.64 (3H, s, CO₂CH₃), 3.96 (1H, br s, CH₃CH), 7.26 (1H, br s, ArH), 7.85 (1H, br s, NH). LRMS (FAB⁺) *m/z* 412 ([M(³⁵Cl₂)+H]⁺), 414 ([M(³⁵Cl³⁷Cl)+H]⁺), 416 ([M(³⁷Cl₂)+H]⁺).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (11a). A mixture of the methyl ester **10a** (80 mg, 0.194 mmol), KOH (109 mg, 3.068 mmol, 10.0 equiv), and 18-crown ether-6 (128 mg, 0.485 mmol, 2.5 equiv) in MeOH (2 mL) was stirred at 80 °C for 7.5 h, the whole was cooled, and then the solvent was evaporated in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and the whole was extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt only to AcOEt/MeOH = 10:1) to afford **11a** (35 mg, 0.087 mmol, yield 45%) as a colorless solid. Compound **11a**: colorless powder. Mp 192–197 °C (*n*-hexane/AcOEt/MeOH). ¹H NMR (CD₃OD): 1.32 (6H, d, *J* = 7.1 Hz, 2CH₃CH), 1.36 (3H, s, CH₃), 1.47 (3H, s, CH₃), 1.60 (2H, m, 2CH), 1.70 (2H, m, 2CH), 1.70 (2H, m, 2CH), 1.85 (2H, m, 2CH), 2.10 (1H, d, *J* = 8.4 Hz, CH), 2.86 (1H, dd, *J* = 15.9 Hz, CH), 3.27 (1H, m, CH), 3.90 (1H, br s, CH₃CH), 7.23 (1H, br s, ArH). LRMS (FAB⁺) *m/z* 398 ([M(³⁵Cl₂)+H]⁺), 400 ([M(³⁵Cl³⁷Cl)+H]⁺), 402 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₀H₂₅Cl₂NO₃·3/4H₂O: C, 58.33; H, 6.30; N, 3.40. Found: C, 58.29; H, 6.39; N, 3.39.

13-Isopropyl-7-oxopodocarpe-8,11,13-triene-15-carboxylic acid methyl ester (7b). To a solution of CrO₃ (790 mg, 7.886 mmol, 1.1 equiv) in Ac₂O (28 mL) and AcOH (14 mL) was dropwise added a suspension of dehydroabiatic acid methyl ester **6b** (2.35 g, 7.456 mmol) in AcOH (6 mL) at rt for 11 min. The reaction mixture was stirred for 15 min at rt, then stirred at 50 °C for 8.5 h. The mixture was cooled, and the whole was poured into icewater (100 mL) and then extracted with CHCl₃ (1 × 200 mL, 2 × 100 mL). The combined organic layer was washed with brine (1 × 100 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford the desired compound **7b** (570 mg, 1.746 mmol, yield 23%) as a pale yellow oil and the undesired overreacted compound (550 mg, 1.431 mmol, yield 19%) as a yellow oil. Compound **7b**: ¹H NMR (CDCl₃) 1.24–1.26 (9H, m, 2CH₃CH and CH₃), 1.34 (3H, s, CH₃), 1.61–1.80 (5H, m, 5CH), 2.32–2.38 (2H, m, 2CH), 2.71–2.73 (2H, m, 2CH), 2.91–2.95 (1H, m, CH₃CH), 3.65 (3H, s, CO₂CH₃), 7.29 (1H, d, *J* = 8.1 Hz, ArH), 7.40 (1H, dd, *J* = 8.2 Hz, 2.1 Hz, ArH), 7.87 (1H, d, *J* = 2.0 Hz,

ArH). LRMS (FAB⁺) *m/z* 329 ([M+H]⁺). *By-product*: (13-(1-Acetoxy-1-methylethyl)-7-oxopodocarpe-8,11,13-triene-15-carboxylic acid methyl ester): ¹H NMR (CDCl₃) 1.26 (3H, s, CH₃), 1.34 (3H, s, CH₃), 1.72–1.79 (1H, m, 5CH and 2CH₃CPh), 2.10 (3H, s, CH₃CO), 2.32–2.36 (2H, m, 2CH), 2.70–2.74 (2H, m, 2× CH), 3.66 (3H, s, CO₂CH₃), 7.33 (1H, d, *J* = 8.2 Hz, ArH), 7.52 (1H, dd, *J* = 8.3 Hz, 2.3 Hz, ArH), 7.97 (1H, d, *J* = 2.4 Hz, ArH).

13-Isopropyl-7-hydroxyimino-podocarpe-8,11,13-triene-15-carboxylic acid methyl ester (8b). A mixture of the methyl ester **7b** (598 mg, 1.667 mmol), pyridine (0.2 mL), and NH₂OH·HCl (180 mg, 2.585 mmol, 1.6 equiv) in EtOH (4 mL) was stirred at 100 °C for 3 h, the mixture was cooled, and the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **8b** (390 mg, 1.137 mmol, yield 68%) as a colorless amorphous solid. Compound **8b**: ¹H NMR (CDCl₃) 1.12 (3H, s, CH₃), 1.25 (6H, d, *J* = 7.0 Hz, 2CH₃CH), 1.38 (3H, s, CH₃), 1.42–1.76 (5H, m, 5CH), 2.28–2.35 (2H, m, 2CH), 2.64–2.67 (2H, m, 2CH), 2.88–2.91 (1H, m, CH₃CH), 3.65 (3H, s, CO₂CH₃), 7.21 (2H, m, ArH), 7.70 (1H, s, ArH). LRMS (FAB⁺) *m/z* 344 ([M+H]⁺).

13-Isopropyl-7-tosyloxyimino-podocarpe-8,11,13-triene-15-carboxylic acid methyl ester (9b). To a solution of the methyl ester **8b** (390 mg, 1.137 mmol) in pyridine (1 mL) was added TsCl (325 mg, 1.705 mmol, 1.5 equiv) at rt. The reaction mixture was stirred for 14 h at rt, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **9b** (545 mg, 1.095 mmol, yield 96%) as a colorless solid. Compound **9b**: colorless powder. Mp 158–160 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.04 (3H, s, CH₃), 1.20–1.28 (6H, m, 2CH₃CH), 1.34 (3H, s, CH₃), 1.68–1.74 (5H, m, 5CH), 2.26 (2H, t, *J* = 9.3 Hz, CH₂), 2.45 (3H, s, PhCH₃), 2.66 (2H, d, *J* = 8.6 Hz, CH₂), 2.85–2.88 (1H, m, CH₃CH), 3.66 (3H, s, CO₂CH₃), 7.18 (1H, d, *J* = 8.2 Hz, ArH), 7.25 (1H, dd, *J* = 8.2 Hz, 2.0 Hz, ArH), 7.35 (2H, d, *J* = 8.6 Hz, ArH), 7.58 (1H, d, *J* = 1.8 Hz, ArH), 7.95 (2H, d, *J* = 8.4 Hz, ArH). LRMS (FAB⁺) *m/z* 498 ([M+H]⁺). Anal. Calcd for C₂₈H₃₅NO₅S: C, 67.58; H, 7.09; N, 2.81. Found: C, 69.39; H, 7.21; N, 2.91.

13-Isopropyl-8,11a-dimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid methyl ester (10b). A mixture of the methyl ester **9b** (505 mg, 1.015 mmol) in TFA (3 mL) was stirred at rt for 40 min, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 1:1) to afford **10b** (392 mg, quantitative yield) as a colorless amorphous solid. Compound **10b**: colorless needles. Mp 148–149 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.24 (6H, d, *J* = 7.0 Hz, 2CH₃CH), 1.42 (3H, s, CH₃), 1.45 (3H, s, CH₃), 1.69–1.93 (6H, m, 6CH), 2.18 (1H, d, *J* = 14.8 Hz, CH), 2.57 (1H, dd, *J* = 14.7 Hz, 8.2 Hz, CH), 2.78–2.81 (1H, dd, *J* = 8.2 Hz, 3.6 Hz, CH), 2.85–2.88 (1H, m, CH₃CH), 3.65 (3H, s, CO₂CH₃), 6.76 (1H, d, *J* = 2.0 Hz, ArH), 7.04 (1H, d, *J* = 8.2 Hz, ArH), 7.33 (1H, d, *J* = 8.4 Hz, ArH), 8.62 (1H, s, NH).

LRMS (FAB⁺) *m/z* 343 ([M+H]⁺). Anal. Calcd for C₂₁H₂₉NO₃: C, 73.44; H, 8.51; N, 4.08. Found: C, 73.21; H, 8.29; N, 4.10.

13-Isopropyl-8,11a-dimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (11b). A mixture of the methyl ester **10b** (100 mg, 0.291 mmol), KOH (163 mg, 2.911 mmol, 10.0 equiv) and 18-crown ether-6 (192 mg, 0.728 mmol, 2.5 eq.) in MeOH (2 mL) was stirred at 80 °C for 15 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and then the mixture was extracted with CHCl₃ (3× 20 mL). The combined organic layer was washed with brine (1× 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 1:2) to afford **11b** (75 mg, 0.228 mmol, yield 78%) as a colorless oil. Compound **11b**: colorless powder (*n*-hexane/AcOEt); Mp 135–140 °C; ¹H NMR (CDCl₃) 1.22 (6H, d, *J* = 6.8 Hz, 2CH₃CH), 1.42 (3H, s, CH₃), 1.45 (3H, s, CH₃), 1.74–1.88 (5H, m, 5CH), 1.95 (1H, d, *J* = 11.3 Hz, CH), 2.28 (1H, dd, *J* = 14.9 Hz, 3.4 Hz, CH), 2.67 (1H, dd, *J* = 14.9 Hz, 7.8 Hz, CH), 2.80 (1H, dd, *J* = 7.9 Hz, 3.7 Hz, CH), 2.83–2.86 (1H, m, CH₃CH), 6.70 (1H, d, *J* = 1.8 Hz, ArH), 7.00 (1H, dd, *J* = 8.0 Hz, 1.8 Hz, ArH), 7.31 (1H, d, *J* = 8.4 Hz, ArH), 7.99 (1H, s, ArH). LRMS (FAB⁺) *m/z* 330 ([M+H]⁺). Anal. Calcd for C₂₀H₂₇NO₃·1/6H₂O: C, 72.26; H, 8.28; N, 4.21. Found: C, 72.21; H, 8.15; N, 4.16.

12,14-Dichloro-3-isopropyl-5,8,11a-trimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid methyl ester (12a). To a suspension of NaH (60% in mineral oil, washed twice with *n*-hexane) (10 mg, 0.244 mmol, 1.1 equiv) in DMF (1 mL) was dropwise added a solution of the methyl ester **10a** (91 mg, 0.222 mmol) in DMF (1 mL) at 0 °C for 2 min in a flask fitted with a CaCl₂ tube. After stirring at 0 °C for 15 min, MeI (0.07 mL, 1.108 mmol, 5.0 equiv) was added to the reaction mixture at 0 °C. Stirring was continued at rt for 7 h, the solvent was evaporated in vacuum, and the residue was diluted with 1N HCl (10 mL) and then extracted with CHCl₃ (3× 20 mL). The combined organic layer was washed with brine (1× 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **12a** (73 mg, 0.171 mmol, yield 77%) as a colorless oil. Compound **12a**: ¹H NMR (CDCl₃) 1.36 (3H, s, CH₃), 1.42 (6H, d, *J* = 7.1 Hz, 2CH₃CH), 1.46 (3H, s, CH₃), 1.72–1.73 (2H, m, CH₂), 1.82–1.91 (4H, m, 4CH), 2.21 (1H, d, *J* = 8.6 Hz, CH), 2.74 (1H, d, *J* = 16.3 Hz, CH), 3.19 (3H, s, NCH₃), 3.64 (3H, s, CO₂CH₃), 3.81 (1H, dd, *J* = 16.2 Hz, 8.8 Hz, CH), 4.09 (1H, br s, CH₃CH), 7.20 (1H, br s, ArH).

12,14-Dichloro-13-isopropyl-5,8,11a-trimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (13a). A mixture of the methyl ester **12a** (73 mg, 0.171 mmol), KOH (96 mg, 1.705 mmol, 10.0 equiv) and 18-crown ether-6 (113 mg, 0.426 mmol, 2.5 equiv) in MeOH (1 mL) was stirred at 80 °C for 13 h, the whole was cooled, and the solvent was evaporat-

ed in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and the whole was extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt only) to afford **13a** (65 mg, 0.156 mmol, yield 92%) as a colorless solid. Compound **13a**: colorless powder. Mp 155–159 °C (*n*-hexane). ¹H NMR (CDCl₃) 1.34 (3H, s, CH₃), 1.39 (6H, d, *J* = 7.1 Hz, 2CH₃CH), 1.43 (3H, s, CH₃), 1.54 (2H, m, 2CH), 1.73–1.88 (4H, m, 4CH), 2.18 (1H, d, *J* = 8.6 Hz, CH), 2.83 (1H, d, *J* = 16.3 Hz, CH), 3.19 (3H, s, NCH₃), 3.79–3.85 (1H, m, CH), 3.94 (1H, br s, CH₃CH), 7.19 (1H, br s, ArH), 10.03 (1H, br s, CO₂H). LRMS (FAB⁺) *m/z* 412 ([M(³⁵Cl₂)+H]⁺), 414 ([M(³⁵Cl³⁷Cl)+H]⁺), 416 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₁H₂₇Cl₂NO₂·1/4H₂O: C, 60.51; H, 6.64; N, 3.36. Found: C, 60.72; H, 6.82; N, 3.26.

13-Isopropyl-5,8,11a-trimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (12b). To a suspension of NaH (60% in mineral oil, washed twice with *n*-hexane) (13 mg, 0.032 mmol, 1.1 equiv) in DMF (1 mL) was dropwise added a solution of the carboxylic acid **10b** (100 mg, 0.291 mmol) in DMF (1 mL) at 0 °C for 1 min, in a flask fitted with a CaCl₂ tube. After stirring at 0 °C for 15 min, MeI (0.091 mL, 1.456 mmol, 5.0 equiv) was added to the reaction mixture at 0 °C. Stirring was continued at rt for 14 h, the solvent was evaporated in vacuum, and the residue was diluted with 1NHCl and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **12b** (90 mg, 0.252 mmol, yield 87%) as a pale orange oil. Compound **12b**: colorless plates. Mp 149–150 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.26 (6H, d *J* = 7.0 Hz, 2CH₃CH), 1.31 (3H, s, CH₃), 1.48 (3H, s, CH₃), 1.65–1.68 (2H, m, 2CH), 1.74–1.77 (2H, m, 2CH), 1.90 (1H, d, *J* = 13.6 Hz, CH), 1.89–1.98 (3H, m, 3CH), 2.50–2.63 (2H, m, 2CH), 2.88–2.92 (1H, m, CH₃CH), 3.27 (3H, s, NCH₃), 3.60 (3H, s, CO₂CH₃), 7.01 (1H, d, *J* = 1.8 Hz, ArH), 7.07 (1H, dd, *J* = 8.2 Hz, 1.9 Hz, ArH), 7.28 (1H, d, *J* = 8.2 Hz, ArH). LRMS (FAB⁺) *m/z* 358 ([M+H]⁺). Anal. Calcd for C₂₂H₃₁NO₃: C, 73.91; H, 8.74; N, 3.92. Found: C, 73.78; H, 8.89; N, 3.97.

13-Isopropyl-5,8,11a-trimethyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid (13b). A mixture of the methyl ester **12b** (50 mg, 0.141 mmol), KOH (79 mg, 1.410 mmol, 10.0 equiv) and 18-crown ether-6 (93 mg, 0.352 mmol, 2.5 equiv) in MeOH (1 mL) was stirred at 80 °C for 18 h, the mixture was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 1:2) to afford **13b** (42 mg, 0.121 mmol, yield 86%) as a colorless amorphous solid. Compound **13b**: colorless flakes. Mp 123–

127 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.24 (6H, d, *J* = 7.0 Hz, 2CH₃CH), 1.31 (3H, s, CH₃), 1.50 (3H, s, CH₃), 1.67–1.78 (4H, m, 2CH), 1.98–2.04 (3H, m, 3CH), 2.50–2.62 (2H, m, 2CH), 2.86–2.90 (1H, m, CH₃CH), 3.28 (3H, s, NCH₃), 6.97 (1H, d, *J* = 1.8 Hz, ArH), 7.06 (1H, dd, *J* = 8.2 Hz, 1.9 Hz, ArH), 7.26 (1H, d, *J* = 8.2 Hz, ArH). LRMS (FAB⁺) *m/z* 344 ([M+H]⁺). Anal. Calcd for C₂₁H₂₉NO₃: C, 73.44; H, 8.51; N, 4.08. Found: C, 73.28; H, 8.31; N, 4.03.

Toluene-4-sulfonic acid pent-4-ynyl ester (15). To a solution of 4-pentyn-1-ol **14** (1.10 mL, 0.012 mol) and Et₃N (1.81 mL, 0.013 mol, 1.1 equiv) in CH₂Cl₂ (8 mL) was dropwise added a solution of TsCl (2.49 g, 0.0113 mol, 1.1 equiv) in CH₂Cl₂ (20 mL) at 0 °C for 20 min, in a flask fitted with a CaCl₂ tube. Stirring was continued at rt for 20 h, and the whole was poured into icewater (50 mL). The separated aqueous layer was extracted with CHCl₃ (3 × 30 mL). Then the combined organic layer was washed with water (1 × 30 mL) and brine (1 × 30 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 4:1) to afford **15** (2.37 g, 0.010 mol, yield 84%) as a colorless liquid. Compound **15**: ¹H NMR (CDCl₃) 1.85–1.89 (2H, m, CH₂CH₂CH₂), 1.89 (1H, t, *J* = 2.7 Hz, HCC), 2.26 (2H, dt, *J* = 6.9 Hz, 2.7 Hz, CCH₂), 2.45 (3H, s, CH₃), 4.15 (2H, t, *J* = 6.1 Hz, CH₂O), 7.35 (2H, d, *J* = 8.2 Hz, ArH), 7.80 (2H, d, *J* = 8.4 Hz, ArH).

5-Iodopent-1-yne (16). A mixture of toluene-4-sulfonic acid pent-4-ynyl ester **15** (817 mg, 3.430 mmol) and NaI (1.234 g, 8.232 mmol, 2.4 equiv) in acetone (8 mL) was stirred at 70 °C for 1 h, the mixture was cooled, and poured into ice-water (20 mL). The whole was extracted with Et₂O (3 × 20 mL). The combined organic layer was washed with satd NaHCO₃ (1 × 20 mL) and brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum to afford **16** (600 mg, 3.093 mmol, yield 90%) as a yellow liquid. Compound **16**: ¹H NMR (CDCl₃) 2.00 (1H, t, *J* = 2.6 Hz, HCC), 2.01 (2H, tt, *J* = 6.7 Hz, 6.7 Hz, CH₂CH₂I), 2.35 (2H, dt, *J* = 6.7 Hz, 2.6 Hz, CCH₂), 3.32 (2H, t, *J* = 6.7 Hz, CH₂).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-pent-4-ynyl-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (17a). To a suspension of NaH (60% in mineral oil, washed twice with *n*-hexane) (6 mg, 0.138 mmol, 1.1 equiv) in DMF (1 mL) was dropwise added a solution of the methyl ester **10a** (52 mg, 0.125 mmol) in DMF (1.5 mL) at 0 °C for 2 min, in a flask fitted with a CaCl₂ tube. After stirring at 0 °C for 30 min, a solution of 5-iodopent-1-yne **16** (122 mg, 0.627 mmol, 5.0 equiv) in DMF (1.5 mL) was added to the reaction mixture at 0 °C. Stirring was continued at rt for 17 h, the solvent was evaporated in vacuum, and the residue was diluted with water (20 mL) and the whole was extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 4:1) to afford **17a** (44 mg, 0.093 mmol, yield 74%) as a colorless

oil. Compound **17a**: ^1H NMR (CDCl_3) 1.42 (3H, s, CH_3), 1.42 (6H, d, $J = 7.0$ Hz, $2\text{CH}_3\text{CH}$), 1.47 (3H, s, CH_3), 1.72 (2H, m, 2CH), 1.83–1.91 (5H, m, 5CH), 2.01 (1H, t, $J = 2.3$ Hz, HCC), 2.21 (1H, d, $J = 8.5$ Hz, CH), 2.30 (2H, t, $J = 7.0$ Hz, CCH_2), 2.90 (1H, d, $J = 16.2$ Hz, CH), 3.05–3.12 (1H, m, CH), 3.66 (3H, s, CO_2CH_3), 3.70 (1H, dd, $J = 16.4$ Hz, 8.9 Hz, CH), 3.96 (1H, br s, CH_3CH), 4.11–4.28 (1H, m, CH), 7.21 (1H, br s, ArH). LRMS (FAB^+) m/z 478 ($[\text{M}(^{35}\text{Cl}_2)+\text{H}]^+$), 480 ($[\text{M}(^{35}\text{Cl}^{37}\text{Cl})+\text{H}]^+$), 482 ($[\text{M}(^{37}\text{Cl}_2)+\text{H}]^+$).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-[5-(4-iodophenyl)pent-4-ynyl]-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (23) and 1,4-bis[[5-(2,4-dichloro-3-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (21). A mixture of the methyl ester **17a** (44 mg, 0.093 mmol), *p*-diiodobenzene (34 mg, 0.102 mmol, 1.1 equiv), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (1 mg, 0.001 mmol, 0.02 equiv), and CuI (1 mg, 0.005 mmol, 0.06 equiv) in Et_3N (2 mL) was stirred at rt for 39 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford the mixture of the monosubstituted compound **25** and *p*-diiodobenzene (55 mg) as pale yellow oil and the disubstituted compound **21** (14 mg, 0.013 mmol, yield 28%) as pale yellow amorphous solid. Then the former mixture was purified by flash chromatography (*n*-hexane/AcOEt = 4:1) again to afford the monosubstituted compound **25** (32 mg, 0.046 mmol, yield 50%) as a pale orange amorphous solid. Compound **25**: ^1H NMR (CDCl_3) 1.41 (6H, d, $J = 5.3$ Hz, $2\text{CH}_3\text{CH}$), 1.42 (3H, s, CH_3), 1.46 (3H, s, CH_3), 1.70 (2H, m, 2CH), 1.81–1.97 (5H, m, 5CH), 2.02–2.09 (1H, m, CH), 2.21 (1H, d, $J = 8.5$ Hz, CH), 2.50 (2H, t, $J = 6.9$ Hz, CH_2), 2.86 (1H, d, $J = 16.2$ Hz, CH), 3.06–3.50 (1H, m, CH), 3.51 (3H, s, CO_2CH_3), 3.72 (1H, dd, $J = 16.1$ Hz, 8.6 Hz, CH), 3.95 (1H, br s, CH_3CH), 4.33–4.39 (1H, m, CH), 7.14 (2H, d, $J = 8.1$ Hz, ArH), 7.20 (1H, br s, ArH), 7.62 (2H, d, $J = 8.1$ Hz, ArH). LRMS (FAB^+) m/z 680 ($[\text{M}(^{35}\text{Cl}_2)+\text{H}]^+$), 682 ($[\text{M}(^{35}\text{Cl}^{37}\text{Cl})+\text{H}]^+$), 684 ($[\text{M}(^{37}\text{Cl}_2)+\text{H}]^+$). Compound **21**: ^1H NMR (CDCl_3) 1.43 (12H, d, $J = 6.0$ Hz, $4\text{CH}_3\text{CH}$), 1.44 (6H, s, 2CH_3), 1.48 (6H, s, 2CH_3), 1.71 (4H, m, 4CH), 1.83–1.99 (10H, m, 10CH), 2.04–2.09 (2H, m, 2CH), 2.22 (2H, d, $J = 8.5$ Hz, 2CH), 2.53 (4H, d, $J = 6.9$ Hz, 4CH), 2.89 (2H, d, $J = 16.3$ Hz, 2CH), 3.09–3.16 (2H, m, 2CH), 3.53 (6H, s, $2\text{CO}_2\text{CH}_3$), 3.74 (2H, dd, $J = 16.1$ Hz, 8.6 Hz, 2CH), 3.97 (2H, br s, $2\text{CH}_3\text{CH}$), 4.35–4.41 (2H, m, 2CH), 7.22 (2H, br s, ArH), 7.35 (4H, s, ArH). LRMS (FAB^+) m/z 1029 ($[\text{M}(^{35}\text{Cl}_4)+\text{H}]^+$), 1031 ($[\text{M}(^{35}\text{Cl}_3^{37}\text{Cl})+\text{H}]^+$), 1033 ($[\text{M}(^{35}\text{Cl}_2^{37}\text{Cl}_2)+\text{H}]^+$), 1035 ($[\text{M}(^{35}\text{Cl}^{37}\text{Cl}_3)+\text{H}]^+$).

1,4-Bis[[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (21). The mixture of the alkyne **17a** (17 mg, 0.036 mmol), the iodide **25** (32 mg, 0.047 mmol, 1.3 equiv), $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (1 mg, 0.001 mmol, 0.04 equiv), and CuI (1 mg, 0.005 mmol, 0.14 equiv) in Et_3N (2 mL) was stirred at rt for 72 h under Ar atmosphere. The reaction mixture

was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **21** (32 mg, 0.031 mmol, yield 84%) as a colorless oil.

1,4-Bis[[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-hydroxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (29). A mixture of the methyl ester **21** (46 mg, 0.045 mmol), KOH (25 mg, 0.445 mmol, 10.0 equiv) and 18-crown ether-6 (59 mg, 0.223 mmol, 5.0 equiv) in MeOH (2 mL) was stirred at 80 °C for 19 h, the mixture was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (20 mL), acidified with 2 N HCl (2 mL) and then extracted with CHCl_3 (3×20 mL). The combined organic layer was washed with brine (1×20 mL), dried over Na_2SO_4 , filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt/MeOH = 10:1) to afford **29** (29 mg, 0.029 mmol, yield 65%) as a colorless solid. Compound **29**: colorless powder. Mp 211–214 °C (*n*-hexane/AcOEt). ^1H NMR (CD_3OD) 1.31 (6H, s, 2CH_3), 1.43–1.49 (12H, m, $2\text{CH}_3\text{CH}$ and 2CH_3), 1.76–2.18 (14H, m, 14CH), 2.27 (2H, d, $J = 8.6$ Hz, 2CH), 2.54–2.55 (4H, m, 4CH), 3.15 (2H, d, $J = 16.5$ Hz, 2CH), 3.65–3.72 (2H, m, 2CH), 4.02 (2H, br s, $2\text{CH}_3\text{CH}$), 4.30 (2H, m, 2CH), 7.32 (4H, s, ArH), 7.35 (2H, br s, ArH). LRMS (FAB^+) m/z 1001 ($[\text{M}(^{35}\text{Cl}_4)+\text{H}]^+$), 1003 ($[\text{M}(^{35}\text{Cl}_3^{37}\text{Cl})+\text{H}]^+$), 1005 ($[\text{M}(^{35}\text{Cl}_2^{37}\text{Cl}_2)+\text{H}]^+$), 1007 ($[\text{M}(^{35}\text{Cl}^{37}\text{Cl}_3)+\text{H}]^+$), 1009 ($[\text{M}(^{37}\text{Cl}_4)+\text{H}]^+$). HRMS (FAB^+) found: 1001.3432. Calcd for $\text{C}_{56}\text{H}_{64}^{35}\text{Cl}_4\text{N}_2\text{O}_6$: 1001.3449. Anal. Calcd for $\text{C}_{56}\text{H}_{64}\text{Cl}_4\text{N}_2\text{O}_6 \cdot 3/2\text{H}_2\text{O}$: C, 65.30; H, 6.56; N, 2.72. Found: C, 65.34; H, 6.59; N, 2.60.

13-Isopropyl-8,11a-dimethyl-6-oxo-5-pent-4-ynyl-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (17d). To a suspension of NaH (60% in mineral oil, washed twice with *n*-hexane) (12 mg, 0.292 mmol, 1.1 equiv) in DMF (1 mL) was dropwise added a solution of the methyl ester **10b** (91 mg, 0.265 mmol) in DMF (1.5 mL) at 0 °C over 1 min, in a flask fitted with a CaCl_2 tube. After stirring at 0 °C for 30 min, a solution of 5-iodopent-1-yne **16** (257 mg, 1.326 mmol, 5.0 equiv) in DMF (1.5 mL) was added to the reaction mixture at 0 °C. Stirring was continued at rt for 18 h, and the solvent was evaporated in vacuum, and the residue was diluted with water (20 mL) and then extracted with CHCl_3 (3×20 mL). The combined organic layer was washed with brine (1×20 mL), dried over Na_2SO_4 , filtered and the solvent was evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 4:1) to afford **17b** (76 mg, 0.85 mmol, yield 70%) as a colorless oil. Compound **17b**: ^1H NMR (CDCl_3) 1.27 (6H, d, $J = 6.9$ Hz, $2\text{CH}_3\text{CH}$), 1.32 (3H, s, CH_3), 1.48 (3H, s, CH_3), 1.66–1.68 (2H, m, 2CH), 1.75–1.78 (2H, m, 2CH), 1.90 (1H, d, $J = 13.3$ Hz, CH), 1.95–2.04 (2H, m, 2CH), 1.96 (1H, t, $J = 2.6$ Hz, HCC), 2.10–2.17 (1H, m, CH), 2.19–2.37 (2H, m, 2CH), 2.50–2.59 (2H, m, 2CH), 2.88–2.96 (1H, m, CH_3CH), 3.48–3.56 (1H, m, CH), 3.60 (3H, s, CO_2CH_3), 3.90–3.97 (1H, m, CH), 7.08 (1H, dd, $J = 8.2$ Hz, 1.9 Hz, ArH), 7.11 (1H, d,

$J = 1.9$ Hz, ArH), 7.29 (1H, d, $J = 8.2$ Hz, ArH). LRMS (FAB⁺) m/z 410 ([M+H]⁺).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-[5-(3-iodophenyl)pent-4-ynyl]-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (24) and 1,3-bis[[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (20). A mixture of the alkyne **17b** (62 mg, 0.129 mmol), *m*-diiodobenzene (21 mg, 0.064 mmol, 0.5 equiv), Pd(PPh₃)₂Cl₂ (1 mg, 0.001 mmol, 0.01 equiv), and CuI (1 mg, 0.005 mmol, 0.04 equiv) in Et₃N (3 mL) was stirred at rt for 49 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum. The resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 4:1) to afford the monosubstituted compound **24** (32 mg, 0.046 mmol, yield 36%) as an orange oil and the disubstituted compound **20** (31 mg, 0.030 mmol, yield 47%) as an orange oil. Compound **24**: ¹H NMR (CDCl₃) 1.41 (6H, d, $J = 5.9$ Hz, 2CH₃CH), 1.42 (3H, s, CH₃), 1.47 (3H, s, CH₃), 1.70–1.71 (2H, m, 2CH), 1.82–1.93 (5H, m, 5CH), 2.03–2.10 (1H, m, CH), 2.20 (1H, d, $J = 8.5$ Hz, CH), 2.51 (2H, t, $J = 7.0$ Hz, CH₂), 2.87 (1H, d, $J = 16.2$ Hz, CH), 3.07–3.14 (1H, m, CH), 3.52 (3H, s, CO₂CH₃), 3.72 (1H, dd, $J = 16.1$ Hz, 8.6 Hz, CH), 3.96 (1H, br s, CH₃CH), 4.32–4.39 (1H, m, CH), 7.02 (1H, t, $J = 7.9$ Hz, ArH), 7.20 (1H, br s, ArH), 7.37 (1H, td, $J = 7.8$ Hz, 1.2 Hz, ArH), 7.61 (1H, td, $J = 8.5$ Hz, 1.4 Hz, ArH), 7.77 (1H, t, $J = 1.6$ Hz, ArH). LRMS (FAB⁺) m/z 680 ([M(³⁵Cl₂)+H]⁺), 682 ([M(³⁵Cl³⁷Cl)+H]⁺), 684 ([M(³⁷Cl₂)+H]⁺). Compound **20**: ¹H NMR (CDCl₃) 1.41 (6H, s, 2CH₃), 1.41 (12H, d, $J = 6.3$ Hz, 4CH₃CH), 1.47 (6H, s, 2CH₃), 1.69–1.70 (4H, m, 4CH), 1.81–1.93 (10H, m, 10CH), 2.03–2.08 (2H, m, 2CH), 2.20 (2H, d, $J = 8.5$ Hz, 2CH), 2.50 (4H, d, $J = 7.0$ Hz, 4CH), 2.88 (2H, d, $J = 16.2$ Hz, 2CH), 3.07–3.14 (2H, m, 2CH), 3.51 (6H, s, 2CO₂CH₃), 3.73 (2H, dd, $J = 16.2$ Hz, 8.7 Hz, 2CH), 3.96 (2H, br s, 2CH₃CH), 4.32–4.39 (2H, m, 2CH), 7.20 (2H, br s, ArH), 7.21 (2H, t, $J = 7.7$ Hz, ArH), 7.32 (2H, d, $J = 7.8$ Hz, ArH), 7.44 (1H, s, ArH). LRMS (FAB⁺) m/z 1029 ([M(³⁵Cl₄)+H]⁺), 1031 ([M(³⁵Cl₃³⁷Cl)+H]⁺), 1033 ([M(³⁵Cl₂³⁷Cl₂)+H]⁺), 1035 ([M(³⁵Cl³⁷Cl₃)+H]⁺).

1,3-Bis[[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (20). A mixture of the alkyne **17a** (22 mg, 0.046 mmol), the iodide **24** (32 mg, 0.046 mmol), Pd(PPh₃)₂Cl₂ (1 mg, 0.001 mmol, 0.03 equiv), and CuI (1 mg, 0.005 mmol, 0.1 equiv) in Et₃N (2 mL) was stirred at rt for 48 h under Ar atmosphere. The reaction mixture was filtered, and the solvent was evaporated in vacuum, and the residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **20** (33 mg, 0.032 mmol, yield 70%) as an orange oil.

1,3-Bis[[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-hydroxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (28). A mixture of the methyl ester **20** (69 mg, 0.067 mmol),

KOH (38 mg, 0.672 mmol, 10.0 equiv), and 18-crown ether-6 (89 mg, 0.336 mmol, 5.0 equiv) in MeOH (2 mL) was stirred at 80 °C for 18 h, the whole was cooled, and the solvent was evaporated in vacuum. The resultant residue was diluted with water (20 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt/MeOH = 10:1) to afford the desired compound **28** (50 mg, 0.050 mmol, yield 75%) as a colorless solid. Compound **28**: colorless powder. Mp 207–213 °C (*n*-hexane/CHCl₃). ¹H NMR (CDCl₃) 1.29 (6H, s, 2CH₃), 1.32 (12H, d, $J = 7.1$ Hz, 4CH₃CH), 1.33 (6H, s, 2CH₃), 1.61 (4H, m, 4CH), 1.75–1.89 (10H, m, 10CH), 2.13 (2H, d, $J = 8.6$ Hz, 2CH), 2.21 (4H, m, 4CH), 2.92–3.00 (2H, m, 2CH), 2.98 (2H, d, $J = 16.7$ Hz, 2CH), 3.50–3.57 (2H, m, 2CH), 3.91 (2H, br s, 2CH₃CH), 4.13–4.20 (2H, m, 2CH), 7.09 (3H, m, ArH), 7.24 (2H, br s, ArH), 7.30 (1H, s, ArH). LRMS (FAB⁺) m/z 1001 ([M(³⁵Cl₄)+H]⁺), 1003 ([M(³⁵Cl₃³⁷Cl)+H]⁺), 1005 ([M(³⁵Cl₂³⁷Cl₂)+H]⁺), 1007 ([M(³⁵Cl³⁷Cl₃)+H]⁺), 1009 ([M(³⁷Cl₄)+H]⁺). HRMS (FAB⁺) found: 1001.4380. Calcd for C₅₆H₆₅³⁵Cl₄N₂O₆: 1001.4302. Anal. Calcd for C₅₆H₆₄Cl₄N₂O₆·3/2H₂O: C, 65.30; H, 6.56; N, 2.72. Found: C, 65.03; H, 6.78; N, 2.62.

13-Isopropyl-8,11a-dimethyl-6-oxo-5-[5-(3-iodophenyl)pent-4-ynyl]-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (26) and 1,3-bis[[5-(13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (22). A mixture of the alkyne **17b** (66 mg, 0.161 mmol), *m*-diiodobenzene (53 mg, 0.161 mmol), Pd(PPh₃)₂Cl₂ (1 mg, 0.001 mmol, 0.01 equiv), and CuI (1 mg, 0.005 mmol, 0.03 equiv) in Et₃N (2 mL) was stirred at rt for 19 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 4:1–2:1) to afford the monosubstituted compound **26** (63 mg, 0.103 mmol, yield 64%) as yellow oil and the disubstituted compound **22** (26 mg, 0.029 mmol, yield 36%) as a yellow oil. Compound **26**: ¹H NMR (CDCl₃) 1.23 (6H, d, $J = 6.9$ Hz, 2CH₃CH), 1.34 (3H, s, CH₃), 1.50 (3H, s, CH₃), 1.68–1.69 (2H, m, 2CH), 1.77–1.79 (2H, m, 2CH), 1.92 (1H, d, $J = 8.3$ Hz, CH), 1.97–2.00 (2H, m, 2CH), 2.09 (1H, m, CH), 2.22–2.23 (1H, m, CH), 2.46–2.64 (4H, m, 4CH), 2.83–2.90 (1H, m, CH₃CH), 3.55–3.61 (1H, m, CH), 3.62 (3H, s, CO₂CH₃), 3.95–4.02 (1H, m, CH), 7.00 (1H, t, $J = 7.9$ Hz, ArH), 7.08 (1H, dd, $J = 8.2$ Hz, 1.8 Hz, ArH), 7.13 (1H, d, $J = 1.8$ Hz, ArH), 7.30 (1H, t, $J = 8.3$ Hz, ArH), 7.32 (1H, dt, $J = 8.0$ Hz, 1.4 Hz, ArH), 7.60 (1H, td, $J = 7.9$ Hz, 1.2 Hz, ArH), 7.73 (1H, t, $J = 1.6$ Hz, ArH). LRMS (FAB⁺) m/z 612 ([M+H]⁺). Compound **22**: ¹H NMR (CDCl₃) 1.19 (12H, d, $J = 6.9$ Hz, 4CH₃CH), 1.33 (6H, s, 2CH₃), 1.48 (6H, s, 2CH₃), 1.67–1.68 (4H, m, 4CH), 1.75–1.77 (4H, m, 4CH), 1.90 (2H, d, $J = 7.3$ Hz, 2CH), 2.01–2.11 (2H, m, 2CH), 2.17–2.24 (2H, m, 2CH), 2.40–2.62 (8H, m, 8CH), 2.80–2.90 (2H, m, 2CH₃CH), 3.42–3.60 (2H, m, 2CH), 3.61 (6H, s, 2CO₂CH₃), 3.92–4.00 (2H, m, 2CH), 7.05 (2H, dd, $J = 8.2$ Hz, 1.9 Hz, ArH), 7.11 (2H, d, $J = 1.9$ Hz, ArH), 7.16 (1H, d,

$J = 8.4$ Hz, ArH), 7.24 (2H, dd, $J = 8.6$ Hz, 1.5 Hz, ArH), 7.28 (2H, d, $J = 8.3$ Hz, ArH), 7.36 (1H, m, ArH). LRMS (FAB⁺) m/z 893 ([M+H]⁺).

1,3-Bis[5-(13-isopropyl-8,11a-dimethyl-8-hydroxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (30). A mixture of the methyl ester **22** (26 mg, 0.029 mmol), KOH (16 mg, 0.289 mmol, 10.0 equiv), and 18-crown ether-6 (38 mg, 0.145 mmol, 5.0 equiv) in MeOH (2 mL) was stirred at 80 °C for 17 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (20 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt/MeOH = 10:1) to afford **30** (9 mg, 0.010 mmol, yield 36%) as a colorless amorphous solid. Compound **30**: colorless powder. Mp 159–163 °C (*n*-hexane/CHCl₃). ¹H NMR (CD₃OD) 1.04 (12H, d, $J = 6.9$ Hz, 4CH₃CH), 1.22 (6H, s, 2CH₃), 1.33 (6H, s, 2CH₃), 1.57–1.68 (8H, m, 8CH), 1.84–1.88 (8H, m, 8CH), 2.21–2.27 (2H, m, 2CH), 2.38–2.48 (8H, m, 8CH), 2.73–2.78 (2H, m, 2CH₃CH), 3.30–3.38 (2H, m, 2CH), 3.80 (2H, m, 2CH), 7.01–7.08 (6H, m, ArH), 7.23–7.25 (4H, m, ArH). LRMS (FAB⁺) m/z 866 ([M+H]⁺). Anal. Calcd for C₅₆H₆₈N₂O₆·2H₂O: C, 74.64; H, 8.05; N, 3.11. Found: C, 74.45; H, 8.31; N, 2.83.

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-[5-(2-iodophenyl)pent-4-ynyl]-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepine-8-carboxylic acid methyl ester (23) and 1,10-bis[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]decane-4,6-diyne (18). A mixture of the alkyne **17a** (77 mg, 0.161 mmol), *o*-diiodobenzene (0.010 mL, 0.080 mmol, 0.5 equiv), Pd(PPh₃)₂Cl₂ (1 mg, 0.001 mmol, 0.01 equiv), and CuI (1 mg, 0.005 mmol, 0.03 equiv) in Et₃N (2 mL) was stirred at rt for 48 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford the monosubstituted compound **23** (17 mg, 0.025 mmol, yield 15%) as orange oil and the homodimeric compound **18** (52 mg, 0.054 mmol, yield 67%) as an orange oil. Compound **23**: ¹H NMR (CDCl₃) 1.43 (6H, d, $J = 7.5$ Hz, 2CH₃CH), 1.44 (3H, s, CH₃), 1.49 (3H, s, CH₃), 1.70 (2H, m, 2CH), 1.85–1.92 (4H, m, 2CH), 2.02–2.03 (1H, m, CH), 2.12–2.17 (1H, m, CH), 2.22 (1H, d, $J = 8.5$ Hz, CH), 2.61 (2H, t, $J = 6.9$ Hz, CH₂), 2.91 (1H, d, $J = 16.3$ Hz, CH), 3.20–3.27 (1H, m, CH), 3.52 (3H, s, CO₂CH₃), 3.77 (1H, dd, $J = 16.1$ Hz, 8.6 Hz, CH), 3.98 (1H, br s, CH₃CH), 4.34–4.38 (1H, m, CH), 6.98 (1H, dt, $J = 7.7$ Hz, 1.7 Hz, ArH), 7.22 (1H, br s, ArH), 7.29 (1H, dt, $J = 7.6$ Hz, 1.2 Hz, ArH), 7.44 (1H, dd, $J = 7.7$ Hz, 1.6 Hz, ArH), 7.84 (1H, dd, $J = 8.0$ Hz, 1.6 Hz, ArH). LRMS (FAB⁺) m/z 680 ([M(³⁵Cl₂)+H]⁺), 682 ([M(³⁵Cl³⁷Cl)+H]⁺), 684 ([M(³⁷Cl₂)+H]⁺). Compound **18**: ¹H NMR (CDCl₃) 1.40 (6H, s, 2CH₃), 1.41 (12H, d, $J = 5.8$ Hz, 4CH₃CH), 1.46 (6H, s, 2CH₃), 1.71 (4H, m, 4CH), 1.79–1.92 (10H, m, 10CH), 1.95–

2.04 (2H, m, 2CH), 2.20 (2H, d, $J = 8.5$ Hz, 2CH), 2.36 (4H, d, $J = 7.1$ Hz, 4CH), 2.83 (2H, d, $J = 16.2$ Hz, 2CH), 2.98–3.05 (2H, m, 2CH), 3.67 (6H, s, 2CO₂CH₃), 3.69 (2H, dd, $J = 16.0$ Hz, 8.6 Hz, 2CH), 3.94 (2H, br s, 2CH₃CH), 4.26–4.32 (2H, m, 2CH), 7.19–7.23 (2H, br s, ArH). LRMS (FAB⁺) m/z 1029 ([M(³⁵Cl₄)+H]⁺), 1031 ([M(³⁵Cl₃³⁷Cl)+H]⁺), 1033 ([M(³⁵Cl₂³⁷Cl₂)+H]⁺), 1035 ([M(³⁵Cl³⁷Cl₃)+H]⁺), 1037 ([M(³⁷Cl₄)+H]⁺).

1,2-Bis[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-methoxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (19). A mixture of the alkyne **17a** (15 mg, 0.032 mmol), the iodide **23** (22 mg, 0.032 mmol), Pd(PPh₃)₄ (2 mg, 0.002 mmol, 0.05 equiv), and CuI (1 mg, 0.010 mmol, 0.3 equiv) in Et₃N (1 mL) was stirred at rt for 19 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **19** (11 mg, 0.011 mmol, yield 33%) as an orange oil. Compound **19**: ¹H NMR (CDCl₃) 1.41 (6H, s, 2CH₃), 1.42 (12H, d, $J = 6.6$ Hz, 4CH₃CH), 1.46 (6H, s, 2CH₃), 1.68 (4H, m, 4CH), 1.81–1.92 (8H, m, 8CH), 1.95–2.00 (2H, m, 2CH), 2.07–2.14 (2H, m, 2CH), 2.20 (2H, d, $J = 8.0$ Hz, 2CH), 2.57 (4H, t, $J = 6.9$ Hz, 2CH₂), 2.89 (2H, d, $J = 16.2$ Hz, 2CH), 3.13–3.20 (2H, m, 2CH), 3.47 (6H, s, 2CO₂CH₃), 3.74 (2H, dd, $J = 16.2$ Hz, 8.6 Hz, 2CH), 3.96 (2H, br s, 2CH₃CH), 4.33–4.38 (2H, m, 2CH), 7.20 (2H, dd, $J = 5.7$ Hz, 3.3 Hz, ArH), 7.19–7.20 (2H, m, ArH), 7.41 (2H, dd, $J = 5.7$ Hz, 3.4 Hz, ArH). LRMS (FAB⁺) m/z 1029 ([M(³⁵Cl₄)+H]⁺), 1031 ([M(³⁵Cl₃³⁷Cl)+H]⁺), 1033 ([M(³⁵Cl₂³⁷Cl₂)+H]⁺), 1035 ([M(³⁵Cl³⁷Cl₃)+H]⁺).

1,2-Bis[5-(12,14-dichloro-13-isopropyl-8,11a-dimethyl-8-hydroxycarbonyl-6-oxo-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[*b,d*]azepin-5-yl)]pent-1-ynyl]benzene (27). A mixture of the methyl ester **19** (11 mg, 0.011 mmol), KOH (60 mg, 0.108 mmol, 10.0 equiv), and 18-crown ether-6 (14 mg, 0.054 mmol, 5.0 equiv) in MeOH (2 mL) was stirred at 80 °C for 16 h, the whole was cooled, and the solvent was evaporated in vacuum, and the resultant residue was diluted with water (20 mL), acidified with 2 N HCl (2 mL) and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt/MeOH = 20:1) to afford **27** (4 mg, 0.004 mmol, yield 41%) as a colorless solid. Compound **27**: pale yellow powder. Mp 181–184 °C (*n*-hexane/AcOEt). ¹H NMR (CD₃OD) 1.28 (6H, s, 2CH₃), 1.34 (12H, m, 2CH₃CH and 2CH₃), 1.73 (14H, m, 14CH), 2.21 (6H, m, 6CH), 2.98 (4H, m, 4CH), 3.54 (2H, m, 2CH), 3.91 (2H, br s, 2CH₃CH), 4.15 (2H, m, 2CH), 7.06 (2H, s, ArH), 7.27 (4H, m, ArH). LRMS (FAB⁺) m/z 1001 ([M(³⁵Cl₄)+H]⁺), 1003 ([M(³⁵Cl₃³⁷Cl)+H]⁺), 1005 ([M(³⁵Cl₂³⁷Cl₂)+H]⁺), 1007 ([M(³⁵Cl³⁷Cl₃)+H]⁺), 1009 ([M(³⁷Cl₄)+H]⁺). Anal. Calcd for C₅₆H₆₄Cl₄N₂O₆·3/2H₂O: C, 65.30; H, 6.56; N, 2.72. Found: C, 65.17; H, 6.79; N, 2.69.

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-pent-4-ynyl-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (31). A mixture of the methyl ester **17a** (41 mg, 0.086 mmol), KOH (48 mg, 0.863 mmol, 10.0 equiv) and 18-crown ether-6 (57 mg, 0.216 mmol, 2.5 equiv) in MeOH (1 mL) was stirred at 80 °C for 17 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then the solvent was evaporated in vacuum. The residue was purified by flash chromatography (AcOEt/MeOH = 10:1) to afford **31** (29 mg, 0.0624 mmol, yield 72%) as a colorless solid. Compound **31**: colorless flakes. Mp 122–126 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.39 (3H, s, CH₃), 1.40 (6H, d, *J* = 5.5 Hz, 2CH₃CH), 1.44 (3H, s, CH₃), 1.69–1.74 (2H, m, 2CH), 1.73 (1H, t, *J* = 2.4 Hz, HCC), 1.81–1.89 (5H, m, 5CH), 2.16 (1H, d, *J* = 8.5 Hz, CH), 2.50–2.53 (2H, m, 2CH), 2.94–3.01 (1H, s, CH), 3.04 (1H, d, *J* = 16.4 Hz, CH), 3.74 (1H, dd, *J* = 15.9 Hz, 9.0 Hz, CH), 3.95 (1H, br s, CH₃CH), 4.36–4.43 (1H, m, CH), 7.19 (1H, br s, ArH). LRMS (FAB⁺) *m/z* 464 ([M(³⁵Cl₂)+H]⁺), 466 ([M(³⁵Cl³⁷Cl)+H]⁺), 468 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₅H₃₁Cl₂NO₃: C, 64.65; H, 6.73; N, 3.02. Found: C, 64.63; H, 6.87; N, 2.94.

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-(5-phenylpent-4-ynyl)-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid methyl ester (32c). A mixture of the alkyne **17a** (20 mg, 0.063 mmol), iodobenzene (0.008 mL, 0.075 mmol, 1.2 equiv), Pd(PPh₃)₂Cl₂ (1 mg, 0.001 mmol, 0.02 equiv), and CuI (1 mg, 0.005 mmol, 0.08 equiv) in Et₃N (1 mL) was stirred at rt for 14 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 5:1) to afford **32a** (37 mg, quantitative yield) as a pale yellow amorphous solid. Compound **32a**: ¹H NMR (CDCl₃) 1.42 (6H, d, *J* = 6.7 Hz, 2CH₃CH), 1.42 (3H, s, CH₃), 1.47 (3H, s, CH₃), 1.69 (2H, m, 2CH), 1.81–1.98 (5H, m, 5CH), 2.06–2.11 (1H, m, CH), 2.20 (1H, d, *J* = 8.5 Hz, CH), 2.52 (2H, t, *J* = 7.0 Hz, CH₂), 2.89 (1H, d, *J* = 16.3 Hz, CH), 3.10–3.17 (1H, m, CH), 3.48 (3H, s, CO₂CH₃), 3.73 (1H, dd, *J* = 16.2 Hz, 8.5 Hz, CH), 3.96 (1H, br s, CH₃CH), 4.32–4.39 (1H, m, CH), 7.20 (1H, br s, ArH), 7.27–7.29 (3H, m, ArH), 7.41 (2H, dd, *J* = 7.9 Hz, 1.9 Hz, ArH). LRMS (FAB⁺), *m/z* 554 ([M(³⁵Cl₂) + H]⁺), 556 ([M(³⁵Cl³⁷Cl)+H]⁺), 558 ([M(³⁷Cl₂)+H]⁺).

12,14-Dichloro-13-isopropyl-8,11a-dimethyl-6-oxo-5-(5-phenylpent-4-ynyl)-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (33a). A mixture of the methyl ester **32a** (37 mg, 0.067 mmol), KOH (38 mg, 0.669 mmol, 10.0 equiv), and 18-crown ether-6 (44 mg, 0.167 mmol, 2.5 equiv) in MeOH (1 mL) was stirred at 80 °C for 15 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (10 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then evaporated in vacuum. The

residue was purified by flash chromatography (*n*-hexane/AcOEt = 10:1) to afford **33a** (31 mg, 0.058 mmol, yield 87%) as a colorless oil. Compound **33a**: colorless powder. Mp 127–130 °C (*n*-hexane/AcOEt). ¹H NMR (CDCl₃) 1.37–1.44 (12H, m, 4CH₃), 1.70 (2H, m, 2CH), 1.79–1.88 (5H, m, 5CH), 1.90–2.04 (1H, m, CH), 2.11 (1H, d, *J* = 8.5 Hz, CH), 2.47 (2H, t, *J* = 6.9 Hz, CH₂), 2.89 (1H, d, *J* = 16.3 Hz, CH), 3.07–3.14 (1H, m, CH), 3.48 (1H, dd, *J* = 16.1 Hz, 8.6 Hz, CH), 3.94 (1H, br s, CH₃CH), 4.27–4.36 (1H, m, CH), 7.13–7.36 (6H, m, ArH). LRMS (FAB⁺) *m/z* 540 ([M(³⁵Cl₂)+H]⁺), 542 ([M(³⁵Cl³⁷Cl)+H]⁺), 544 ([M(³⁷Cl₂)+H]⁺). Anal. Calcd for C₂₂H₂₉Cl₂NO₃: C, 68.88; H, 6.53; N, 2.59. Found: C, 68.62; H, 6.65; N, 2.58.

13-Isopropyl-8,11a-dimethyl-6-oxo-5-(5-phenylpent-4-ynyl)-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid methyl ester (32b). A mixture of the alkyne **17b** (88 mg, 0.215 mmol), iodobenzene (0.029 mL, 0.258 mmol, 1.2 equiv), Pd(PPh₃)₂Cl₂ (2 mg, 0.002 mmol, 0.01 equiv), and CuI (1 mg, 0.005 mmol, 0.02 equiv) in Et₃N (1 mL) was stirred at rt for 21 h under Ar atmosphere. The reaction mixture was filtered, the solvent was evaporated in vacuum, and the resultant residue was purified by flash chromatography (*n*-hexane/AcOEt = 2:1) to afford **32b** (81 mg, 0.167 mmol, yield 78%) as a pale orange amorphous solid. Compound **32b**: ¹H NMR (CDCl₃) 1.20 (6H, d, *J* = 6.9 Hz, 2CH₃CH), 1.33 (3H, s, CH₃), 1.49 (3H, s, CH₃), 1.66–1.68 (2H, m, 2CH), 1.74–1.77 (2H, m, 2CH), 1.91 (1H, d, *J* = 13.3 Hz, CH), 1.92–1.98 (2H, m, 2CH), 2.05–2.12 (1H, m, CH), 2.16–2.27 (1H, m, CH), 2.41–2.62 (4H, m, 4CH), 2.79–2.90 (1H, m, CH₃CH), 3.56–3.63 (1H, m, CH), 3.60 (3H, s, CO₂CH₃), 3.95–4.02 (1H, m, CH), 7.06 (1H, dd, *J* = 8.2 Hz, 1.7 Hz, ArH), 7.13 (1H, d, *J* = 1.9 Hz, ArH), 7.24–7.26 (3H, m, ArH), 7.28 (1H, d, *J* = 8.3 Hz, ArH), 7.34–7.37 (2H, m, ArH). LRMS (FAB⁺) *m/z* 486 ([M+H]⁺).

13-Isopropyl-8,11a-dimethyl-6-oxo-5-(5-phenylpent-4-ynyl)-6,7,7a,8,9,10,11,11a-octahydro-5H-dibenzo[b,d]azepine-8-carboxylic acid (33b). A mixture of the methyl ester **32b** (67 mg, 0.139 mmol), KOH (93 mg, 0.167 mmol, 1.2 equiv), and 18-crown ether-6 (92 mg, 0.347 mmol, 2.5 equiv) in MeOH (1 mL) was stirred at 80 °C for 15 h, the whole was cooled, and the solvent was evaporated in vacuum. The residue was diluted with water (20 mL), acidified with 2 N HCl (2 mL), and then extracted with CHCl₃ (3 × 20 mL). The combined organic layer was washed with brine (1 × 20 mL), dried over Na₂SO₄, filtered, and then evaporated in vacuum. The residue was purified by flash chromatography (*n*-hexane/AcOEt = 1:1) to afford the unreacted compound **32b** (24 mg, 0.049 mmol, yield 35% after recrystallization) as a colorless amorphous solid and the desired compound **33b** (32 mg, 0.068 mmol, yield 49%) as a colorless amorphous solid. Compound **33b**: pale yellow powder. Mp 76–79 °C (*n*-hexane). ¹H NMR (CDCl₃) 1.24 (6H, d, *J* = 7.0 Hz, 2CH₃CH), 1.26 (3H, s, CH₃), 1.46 (3H, s, CH₃), 1.65–1.77 (4H, m, 4CH), 1.88–2.03 (2H, m, 2CH), 1.97 (1H, d, *J* = 13.3 Hz, CH), 2.05–2.12 (1H, m, CH), 2.46–2.57 (4H, m, 4CH), 2.84–2.91 (1H, m, CH₃CH), 3.23–3.31 (1H, m, CH), 3.74–3.80

(1H, m, CH), 7.05 (1H, dd, $J = 8.2$ Hz, 1.8 Hz, ArH), 7.08 (1H, d, $J = 1.8$ Hz, ArH), 7.18 (2H, d, $J = 8.2$ Hz, ArH), 7.21–7.25 (2H, m, ArH), 7.29 (1H, d, $J = 7.4$ Hz, ArH). LRMS (FAB⁺) m/z 472 ([M+H]⁺). Anal. Calcd for C₃₁H₃₇Cl₂NO₃: C, 78.95; H, 7.91; N, 2.97. Found: C, 78.65; H, 8.07; N, 2.80.

5.2. Biological assays

5.2.1. Electrophysiological measurements. Human BK channel subunits (hslo α and hslo β 1) were subcloned into pcDNA3 (Invitrogen, CA, USA) and pTracer-CMV2 (Invitrogen), respectively, and transiently co-expressed in TSA201 cells with FuGENE6 transfection reagent (Roche Diagnostics Corporation, IN, USA). BK channel currents were measured with the inside-out patch clamp technique (Patch/Whole Cell Clamp Amplifier CEZ-2400, Nihon Koden). Experiments were performed within 24–48 h after the transfection. All the experiments were carried out at room temperature (20–25 °C). Test compounds were diluted in the intracellular solution (in mM; 140 K-Mes (Alfa Aesar, MA, USA), 10 HEPES (Sigma–Aldrich, Inc., MO, USA), 5 HEDTA (Nacalai Tesque, Inc., Kyoto, Japan), and 100 nM free-Ca²⁺, pH 7.4) at the final concentration of 10 μ M and applied to the intracellular side of the patch membrane. The resistance of the patch pipette was 2–5 M Ω when filled with the pipette solution. Test voltages from –100 mV to 190 mV were applied at 10 mV steps. The BK channel conductance was determined by measuring the tail current at each test voltage, and plotted as conductance–voltage curves. $V_{1/2}$ (half-maximal activation voltage) values were calculated based on Boltzmann's fit of the conductance–voltage curves. The BK channel opening activities of tested compounds were evaluated as the hyperpolarizing shift of $V_{1/2}$.

5.2.2. Measurement of isolated detrusor smooth muscle relaxation. The experimental protocol complied with the Guidelines for Animal Experiments approved by Toho University. Urinary bladders were isolated from male Japanese white rabbits (3–4 kg, 3–4 months old) that had been euthanized with an overdose of pentobarbital. The isolated tissue was immediately immersed in Krebs-bicarbonate solution (in mM; 118 NaCl, 4.7 KCl, 1.2 MgSO₄, 2.6 CaCl₂, 1.2 KH₂PO₄, 25 NaHCO₃, and 11 glucose). The longitudinal strips of detrusor smooth muscle (5–7 mm length, 2–3 mm width), after removal of mucosa by dissection, were suspended in an organ bath chamber containing 10 mL of Krebs solution, maintained at 37 °C and bubbled with 95% O₂/5% CO₂ gas. Muscle tension was increased isotonicly using a force–displacement transducer (TB-612T, Nihon Koden, Tokyo, Japan) connected to a carrier amplifier (AP-601G, Nihon Koden) with a basal preload of ca. 2 gW as initial tension. The tissues were allowed to equilibrate for 1 h. The strips were precontracted with high-K⁺ (30 mM or 120 mM) Krebs solution by replacement of the whole bath solution and allowed to equilibrate for 5–30 min (typically) before the addition of test compounds. When high-K⁺ solutions were used to contract preparations, equimolar concentrations of

Na⁺ in Krebs solution were replaced with K⁺ to maintain isotonicity. The relaxant activities of the test compounds were compared with that of vehicle (0.1% DMSO) measured in the same muscle strip. All values are expressed as means \pm SEM. Statistical significance of differences was determined by paired Student's *t*-test, and *P* values less than 0.05 were considered to be significant.

Acknowledgments

We are grateful to Prof. Ligia Toro (University of California at Los Angeles, U.S.A.) for the gifts of hslo α and hslo β 1. This work was supported by the Health and Labour Sciences Research Grants, Research on Risk of Chemical Substances, from Ministry of Health, Labour and Welfare to T.O. and S.A.-A.

References and notes

- Marty, A. *Nature* **1981**, *291*, 497–500.
- Srinivas Ghatta, S.; Nimmagadda, D.; Xu, X.; Stephen, T.; O'Rourke, S. T. *Pharmacol. Ther.* **2006**, *110*, 103–116.
- Jones, E. M. C.; Gray-Keller, M.; Fettiplace, R. *J. Physiol.* **1999**, *518*, 653–665.
- Ahluwalia, J.; Tinker, A.; Clapp, L. H.; Duchon, M. R.; Abramov, A. Y.; Pope, S.; Nobles, M.; Segal, A. W. *Nature* **2004**, *427*, 853–858.
- Kotlikoff, M.; Hall, I. *J. Clin. Invest.* **2003**, *112*, 654–656.
- (a) Shen, K. Z.; Lagrutta, A.; Davies, N. W.; Standen, N. B.; Adelman, J. P.; North, R. A. *Pflugers Arch.* **1994**, *426*, 440–445; (b) Garcia-Calvo, M.; Knaus, H. G.; McManus, O. B.; Giangiacomo, K. M.; Kaczorowski, G. J.; Garcia, M. L. *J. Biol. Chem.* **1994**, *269*, 676–682; (c) McManus, O. B.; Helms, L. M.; Pallanck, L.; Ganetzky, B.; Swanson, R.; Leonard, R. *J. Neurosci.* **1995**, *14*, 645–650.
- Wallner, M.; Meera, P.; Toro, L. *Proc. Natl. Acad. Sci. U.S.A.* **1996**, *93*, 14922–14927.
- Papazian, D. M.; Timpe, L. C.; Jan, Y. N.; Jan, L. Y. *Nature* **1991**, *349*, 305–310.
- Brenner, R.; Jegla, T. J.; Wickenden, A.; Liu, Y.; Aldrich, R. W. *J. Biol. Chem.* **2000**, *275*, 6453–6461.
- (a) Meredith, A. L.; Thorneloe, K. S.; Werner, M. E.; Nelson, M. T.; Aldrich, R. W. *J. Biol. Chem.* **2004**, *279*, 36746–36752; (b) Petkov, G. V.; Bonev, A. D.; Heppner, T. J.; Brenner, R.; Aldrich, R. W.; Nelson, M. T. *J. Physiol.* **2001**, *537*, 443–452; (c) Thorneloe, K. S.; Meredith, A. L.; Knorn, A. M.; Aldrich, R. W.; Nelson, M. T. *Am. J. Physiol. Renal Physiol.* **2003**, *289*, F604–F610.
- Ruttiger, L.; Sausbier, M.; Zimmermann, U.; Winter, H.; Braig, C.; Engel, J.; Knirsch, M.; Arntz, C.; Langer, P.; Hirt, B.; Müller, M.; Köpfschall, I.; Pfister, M.; Münkner, S.; Rohbock, K.; Pfaff, I.; Rüscher, A.; Ruth, P.; Knipper, M. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 12922–12927.
- Sausbier, M.; Hu, H.; Arntz, C.; Feil, S.; Kamm, S.; Adelsberger, H.; Sausbier, U.; Sailer, C. A.; Feil, R.; Hofmann, F.; Korth, M.; Shipston, M. J.; Knaus, H.-G.; Wolfer, D. P.; Pedroarena, C. M.; Storm, J. F.; Ruth, P. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101*, 9474–9478.
- Sato, T.; Saito, T.; Saegusa, N.; Nakaya, H. *Circulation* **2005**, *111*, 198–203.
- Olesen, S.-P.; Munchm, E.; Watjen, F.; Drejer, J. *Neuroreport* **1994**, *5*, 1001–1004.

15. Olesen, S. P.; Munch, E.; Moldt, P.; Drejer, J. *Eur. J. Pharmacol.* **1994**, *251*, 53–59.
16. Ottolia, M.; Toro, L. *Biophys. J.* **1994**, *67*, 2272–2279.
17. Strøbak, D.; Christopherso, P.; Holm, N. R.; Moldt, P.; Ahring, P. K.; Johansen, T. E.; Olesen, S.-P. *Neuropharmacology* **1996**, *35*, 903–914.
18. (a) Gribkoff, V. K.; Starrrett, J. E., Jr.; Dworetzky, S. I.; Hewawasam, P.; Boissard, C. G.; Cook, D. A.; Frantz, S. W.; Heman, K.; Hibbard, J. R.; Huston, K.; Johnson, G.; Krishnan, B. S.; Kinney, G. G.; Lombardo, L. A.; Meanwell, N. A.; Molinoff, P. B.; Myers, R. A.; Moon, S. L.; Oritiz, A.; Pajor, L.; Pieschl, R. L.; Post-Munson, D. J.; Signor, L. J.; Srinivas, N.; Taber, M. T.; Thalody, G.; Trojnacki, J. T.; Wiener, H.; Yeleswaram, K.; Yeola, S. W. *Nat. Med.* **2001**, *7*, 471–477; (b) Hewawasam, P.; Erway, M.; Moon, S. L.; Knipe, J.; Weiner, H.; Boissard, C. G.; Post-Munson, D. J.; Gao, Q.; Huang, S.; Gribkoff, V. K.; Nicholas, A.; Meanwell, N. A. *J. Med. Chem.* **2002**, *45*, 1487–1499.
19. Tanaka, M.; Sasaki, Y.; Hukui, T.; Kyotani, J.; Hayashi, S.; Hamada, K.; Kimura, Y.; Ukai, Y.; Kitano, M.; Kimura, K. *BJU Int.* **2003**, *92*, 1031–1036.
20. Hu, S.; Fink, C. A.; Lappe, R. W. *Drug Dev. Res.* **1997**, *41*, 10–21, and *Drug Dev. Res.* **1997**, *41*, 109.
21. McManus, O. B.; Harris, G. H.; Giangiacomo, K. M.; Feigenbaum, P.; Reuben, J. P.; Addy, M. E.; Burka, J. F.; Kaczorowski, G. J.; Garcia, M. L. *Biochemistry* **1993**, *32*, 6128–6133.
22. (a) Singh, S. B.; Goetz, M. A.; Zink, D. L.; Dombrowski, A. W.; Polishook, J. D.; Garcia, M. L.; Schemallhofer, W.; McManus, O. B.; Kaczorowski, G. J. *J. Chem. Soc., Perkin Trans. 1* **1994**, 3349–3352; (b) Kaczorowski, G. J.; Knaus, H. G.; Leonard, R. J.; McManus, O. B.; Garcia, M. L. *J. Bioenerg. Biomembr.* **1996**, *28*, 255–267.
23. Coghlan, M. J.; Carroll, W. A.; Gopalakrishnan, M. *J. Med. Chem.* **2001**, *44*, 1628–1653.
24. (a) Ohwada, T.; Nonomura, T.; Maki, K.; Sakamoto, K.; Ohya, S.; Muraki, K.; Imaizumi, Y. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3971–3974; (b) Imaizumi, Y.; Sakamoto, K.; Yamada, A.; Hotta, A.; Ohya, S.; Muraki, K.; Uchiyama, M.; Ohwada, T. *Mol. Pharmacol.* **2002**, *62*, 836–846; (c) Sakamoto, K.; Nonomura, T.; Ohya, S.; Muraki, K.; Ohwada, T.; Imaizumi, Y. *J. Pharmacol. Exp. Ther.* **2006**, *316*, 144–153.
25. Nelson, W. L.; Miller, D. D.; Wilson, R. S. *J. Heterocycl. Chem.* **1969**, *6*, 131–133.
26. (a) Bruhova, I.; Zhorov, B. Z. *Biophys. J.* **2005**, *89*, 1020–1029; (b) Lipkind, G. M.; Fozzard, H. A. *Mol. Pharmacol.* **2005**, *68*, 1611–1622.
27. Macromolecule 8.5, 2003, Schrödinger, L.L.C., Portland, Oregon, USA.
28. (a) Seth, A.; Capello, E.; Chou, C.-L.; Longhurst, P. A. *BJU Int.* **2005**, *95*, 157–162; (b) Uchida, W.; Masuda, M.; Shirai, Y.; Shibasaki, K.; Satoh, N.; Takenaka, T. *Naunyn-Schmiedeberg's Arch. Pharmacol.* **1994**, *350*, 398–402; (c) Kobayashi, H.; Satomi Adachi-Akahane, S.; Nagao, T. *Eur. J. Pharmacol.* **2000**, *404*, 231–238.
29. (a) Mason, R. P.; Rhodes, D. G.; Herbetette, L. G. *J. Med. Chem.* **1991**, *34*, 869–877; (b) Yamaguchi, S.; Zhorov, B. S.; Yoshioka, K.; Nagao, T.; Ichijo, H.; Adachi-Akahane, S. *Mol. Pharmacol.* **2003**, *64*, 235–248.

Molecular Mechanisms for Large Conductance Ca^{2+} -Activated K^+ Channel Activation by a Novel Opener, 12,14-Dichlorodehydroabiatic Acid

Kazuho Sakamoto, Taro Nonomura, Susumu Ohya, Katsuhiko Muraki, Tomohiko Ohwada, and Yuji Imaizumi

Department of Molecular and Cellular Pharmacology, Graduate School of Pharmaceutical Sciences, Nagoya City University, Nagoya, Japan (K.S., S.O., K.M., Y.I.); and Laboratory of Organic and Medicinal Chemistry, Graduate School of Pharmaceutical Sciences, University of Tokyo, Tokyo, Japan (T.N., T.O.)

Received August 4, 2005; accepted September 28, 2005

ABSTRACT

Our recent study has revealed that 12,14-dichlorodehydroabiatic acid (diCl-DHAA), which is synthetically derived from a natural product, abiatic acid, is a potent opener of large conductance Ca^{2+} -activated K^+ (BK) channel. Here, we examined, by using a channel expression system in human embryonic kidney 293 cells, the mechanisms underlying the BK channel opening action of diCl-DHAA and which subunit of the BK channel (α or β 1) is the site of action for diCl-DHAA. BK channel activity was significantly enhanced by diCl-DHAA at concentrations of 0.1 μM and higher in a concentration-dependent manner. diCl-DHAA enhanced the activity of $\text{BK}\alpha$ by increasing sensitivity to both Ca^{2+} and membrane potential without changing the single channel conductance. It is notable that the

increase in BK channel open probability by diCl-DHAA showed significant inverse voltage dependence, i.e., larger potentiation at lower potentials. Since coexpression of β 1 subunit with $\text{BK}\alpha$ did not affect the potency of diCl-DHAA, the site of action for diCl-DHAA is suggested to be $\text{BK}\alpha$ subunit. Moreover, kinetic analysis of single channel currents indicates that diCl-DHAA opens $\text{BK}\alpha$ mainly by decreasing the time staying in a long closed state. Although reconstituted voltage-dependent Ca^{2+} channel current was significantly reduced by 1 μM diCl-DHAA, BK channels were selectively activated at lower concentrations. These results indicate that diCl-DHAA is one of the most potent BK channel openers acting on $\text{BK}\alpha$ and a useful prototype compound to develop a novel BK channel opener.

Large conductance Ca^{2+} -activated K^+ (BK) channels are expressed in many different types of excitable cells and have significant physiological roles in the regulation of frequency of firing, action potential repolarization, and/or afterhyperpolarization (for reviews, see Vergara et al., 1998; Kaczorowski and Garcia, 1999). BK channel activation by the Ca^{2+} -induced Ca^{2+} release during excitation-contraction coupling significantly contributes to action potential repolarization/afterhyperpolarization in some smooth muscle cells (Imaizumi et al., 1998; Ohi et al., 2001).

In addition, the negative feedback control of intracellular

Ca^{2+} concentration ($[\text{Ca}^{2+}]_i$) by BK channels works to protect cells from Ca^{2+} overload during pathophysiological conditions (Lawson, 2000). Hyperpolarization of neuronal cells by BK channel activation down-regulates the activity of voltage-dependent Na^+ and Ca^{2+} channels and may prevent cell death, which is mainly caused by excess intracellular Ca^{2+} in the setting of brain ischemia following stroke. In smooth muscle cells, BK channels are also activated by spontaneous Ca^{2+} release from sarcoplasmic reticulum (Nelson and Quayle, 1995; Bolton and Imaizumi, 1996; Imaizumi et al., 1999) and are thought to be one of the essential regulators of resting membrane potential. Accumulated evidence indicates that the control of BK channel activity in arterial smooth muscle is one of the major determinants of vascular tone and that its abnormality can be a cause of hypertension (Brenner

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.
doi:10.1124/jpet.105.093856.

ABBREVIATIONS: BK, large conductance Ca^{2+} -activated K^+ ; $[\text{Ca}^{2+}]_i$, intracellular Ca^{2+} concentration; BMS-204352, (3S)-(+)-(5-chloro-2-methoxyphenyl)-1,3-dihydro-3-fluoro-6-(trifluoromethyl)-2H-indole-2-one; L-735,334, 14-hydroxy 8-daucene-3,4-diol oleate; diCl-DHAA, 12,14-dichlorodehydroabiatic acid; CaV, voltage-dependent Ca^{2+} ; HEK, human embryonic kidney; I-V, current-voltage; SK, small conductance Ca^{2+} -activated K^+ ; IK, intermediate conductance Ca^{2+} -activated K^+ ; CGS-7181, ethyl 2-hydroxy-1-[[[4-methylphenyl]amino]oxo]-6-trifluoromethyl-1H-indole-3-carboxylate; CGS-7184, ethyl 1-[[[4-chlorophenyl]amino]oxo]-2-hydroxy-6-trifluoromethyl-1H-indole-3-carboxylate; NS-1608, N-(3-(trifluoromethyl)phenyl)-N'-(2-hydroxy-5-chlorophenyl)urea; NS-1619, 1,3-dihydro-1-[2-hydroxy-5-(trifluoromethyl)phenyl]-5-(trifluoromethyl)-2H-benzimidazol-2-one; NS-8, 2-amino-3-cyano-5-(2-fluorophenyl)-4-methylpyrrole.

et al., 2000b; Wellman and Nelson, 2003; Fernández-Fernández et al., 2004).

Agents that enhance BK channel activity (BK channel openers) may therefore be effective in protecting neurons from damage following an ischemic stroke and/or in suppressing excess activity of smooth muscle tissues (Lawson, 2000). Many compounds that were found from natural products or were synthesized have been reported to be BK channel openers (Coghlan et al., 2001). Most of these BK channel openers, including BMS-204352 (Gribkoff et al., 2001), are not highly potent activators (EC_{50} values >300 nM) under the resting cellular conditions where intracellular Ca^{2+} concentration is 50 to 150 nM (Schröder et al., 2003). Terpenoids derived from natural products—dehydrosoyasaponin-I, maxikdiol, and L-735,334—have been reported as BK channel openers (Kaczorowski and Garcia, 1999). In addition, 17 β -estradiol (Valverde et al., 1999) and epoxyeicosatrienoic acids (Fukao et al., 2001) may be endogenous BK channel openers, and some transmitters and hormones can enhance BK channel activity via kinase activation (Vergara et al., 1998).

In our previous study (Imaizumi et al., 2002), novel compounds, including pimelic acid, were discovered from terpenoids, which have chemical structures similar to that of maxikdiol, a moderate BK channel opener (Singh et al., 1994). Moreover, our recent study (Ohwada et al., 2003) has revealed that chemical modification of abietic acid, an inactive compound of resin acid derivatives, to dehydroabietic acid resulted in BK channel opening, and further chemical modification to 12,14-dichlorodehydroabietic acid (diCl-DHAA) led to finding of a potent BK channel opener. However, the underlying mechanisms of diCl-DHAA-induced activation of BK channel and the selectivity against voltage-dependent Ca^{2+} (CaV) channel have not been defined. The present study was therefore undertaken to identify molecular mechanisms of diCl-DHAA-induced activation of BK channels and to examine the selectivity against inhibition of CaV channels by using human embryonic kidney (HEK) 293 cells as an expression system.

Materials and Methods

Vector Constructs, Cell Culture, and Transfection. Restriction enzyme-digested DNA fragments of BK α (KpnI/XbaI-double digested) and BK β 1 (EcoRI/XbaI-double digested) were ligated into mammalian expression vectors pcDNA3.1(+) and pcDNA3.1/Zeo(+), (Invitrogen, Carlsbad, CA), respectively, using the TaKaRa ligation kit version 1 (TaKaRa, Osaka, Japan) (Yamada et al., 2001). HEK293 cell lines were obtained from Health Science Research Resources Bank (Tokyo, Japan) and maintained in minimal essential medium (Invitrogen) supplemented with 10% heat-inactivated fetal calf serum (JRH Biosciences, Lenexa, KS), 100 units/ml penicillin (Wako Pure Chemicals, Osaka, Japan), and 100 μ g/ml streptomycin (Meiji Seika, Tokyo, Japan). Stable expression of BK α and BK β was achieved by using calcium phosphate coprecipitation transfection techniques as reported previously (Imaizumi et al., 2002). G418- and G418/zeocin-resistant cells were selected as those which were BK α -expressing and BK β 1-coexpressing, respectively.

The cDNAs encoding voltage-dependent Ca^{2+} channel α 1C subunit of the rabbit (rCaV α 1C) and β 3 subunit of the mouse (mCaV β 3) were kind gifts from Dr. Veit Flockerzi (Institut für Pharmakologie und Toxikologie, Universität des Saarlandes, Hamburg, Germany) and were ligated into mammalian expression vectors pcDNA3.1(+) and pTracer(+), respectively (Murakami et al., 2003). These plasmid vectors were transfected into HEK293 cells for transient expression.

The functional coexpression of rCaV α 1C and mCaV β 3 was successfully determined by the appearance of the inward currents and green fluorescent protein fluorescence.

Solutions. The standard HEPES-buffered solution for electrophysiological recording had an ionic composition of 137 mM NaCl, 5.9 mM KCl, 2.2 mM $CaCl_2$, 1.2 mM $MgCl_2$, 14 mM glucose, and 10 mM HEPES. The pH of the solution was adjusted to 7.4 with NaOH. The pipette solution for whole-cell recordings of K^+ currents contained 140 mM KCl, 1 mM $MgCl_2$, 10 mM HEPES, 2 mM Na_2ATP , and 5 mM EGTA. The pCa and pH of the pipette solution were adjusted to 6.5 and 7.2 by adding $CaCl_2$ and KOH, respectively. For recordings of single BK channel currents in the excised inside-out patch configuration, the pipette solution contained the standard HEPES-buffered solution or K^+ -rich HEPES-buffered solution that was prepared by replacement of 134.1 mM NaCl in the standard HEPES-buffered solution with equimolar KCl. The bathing solution contained 140 mM KCl, 1.2 mM $MgCl_2$, 14 mM glucose, 10 mM HEPES, and 5 mM EGTA. Selected pCa of the bathing solution was obtained by adding adequate amount of $CaCl_2$, and the pH was adjusted to 7.2 with NaOH. The pipette solution for whole-cell recording of Ca^{2+} inward currents had an ionic composition of 140 mM CsCl, 1 mM $MgCl_2$, 10 mM HEPES, 2 mM Na_2ATP , and 5 mM EGTA. The pH of the pipette solution was adjusted to 7.2 by adding CsOH.

Electrophysiological Experiments. The whole-cell and inside-out patch clamps were applied to single cells using CEZ-2400 amplifier (Nihon Kohden, Tokyo, Japan) and EPC-7 amplifier (List Electronics, Darmstadt, Germany), respectively. The procedures of electrophysiological recordings and data acquisition/analysis for whole-cell recording have been described previously (Imaizumi et al., 1989). The resistance of the pipette was 1.5 to 3 M Ω for whole-cell and 15 to 25 M Ω for inside-out patch configurations when filled with the pipette solutions. The series resistance was partly compensated electrically under whole-cell voltage clamp. Whole-cell and single channel recordings were carried out at room temperature ($24 \pm 1^\circ C$). Single channel current analyses were done using software PAT V7.0C (developed by Dr. J. Dempster, University of Strathclyde, Glasgow, Scotland). The open probability (P_o) was measured from the event histogram plotted against current amplitude. The number of channels in a patch was determined from recordings at pCa = 3.5, and the analyses were performed only when the number of channels in a patch was less than six.

Chemicals. Most of pharmacological agents were obtained from Sigma-Aldrich (St. Louis, MO), unless mentioned otherwise. Iberiotoxin was obtained from Peptide Institute Inc. (Osaka, Japan). Pimelic acid, abietic acid, dehydroabietic acid, and diCl-DHAA were obtained from Helix Biotech (New Westminster, BC, Canada). The test compounds were dissolved with dimethyl sulfoxide. The final concentration of dimethyl sulfoxide was 0.03% or lower.

Statistics. Data are expressed as means \pm S.E.M. Statistical significance between two groups and among multiple groups was evaluated using Student's *t* test and Scheffé's test after F-test or one-way analysis of variance, respectively.

Results

Effects of diCl-DHAA on Macroscopic BK Channel

Currents. Effects of diCl-DHAA on BK channel currents were examined in single HEK293 cells under whole-cell voltage-clamp mode. The Ca^{2+} concentration in the pipette solution was fixed at pCa = 6.5 using a Ca^{2+} -EGTA buffer (see *Materials and Methods*). Depolarization from -60 to $+10$ mV induced outward currents in both native HEK and HEK293 cells, whereas the current density was approximately 4 times larger in the latter cells as reported previously (Imaizumi et al., 2002). Application of diCl-DHAA in a concentration range of 0.1 to 1.0 μ M increased the outward




Fig. 1. Effects of diCl-DHAA on macroscopic membrane currents in HEKKBK $\alpha\beta$ 1. A and B, a single HEKKBK $\alpha\beta$ 1 was depolarized from -60 to $+10$ mV for 150 ms under whole-cell voltage-clamp mode. diCl-DHAA was applied in a concentration range of 0.1 and 1 μ M. Original current recordings at different concentrations were superimposed and are shown in A. The peak outward current amplitude at $+10$ mV was measured and plotted against time in B. The original traces were obtained at the time indicated by vertical arrows in the time course. It is notable that effects of 0.1 to 1 μ M diCl-DHAA were completely removed by washout. C, concentration-response relationships for diCl-DHAA. Experiments were carried out in a manner typically shown in A. Data about current density at $+10$ mV, which was obtained by dividing peak current amplitude with cell capacitance in each cell (pA/pF), are summarized as open columns. The relative amplitude of peak outward current at $+10$ mV in the presence of diCl-DHAA ($I_{diCl-DHAA}/I_{control}$) was also determined by taking the amplitude in the absence of diCl-DHAA as unity (closed columns). Means \pm S.E.M. are shown by columns and vertical bars, respectively. #/#, $p < 0.05$ and **/##, $p < 0.01$ versus control.

currents in HEKKBK $\alpha\beta$ 1 in a dose-dependent manner (Fig. 1, A and B) but not in native HEK cells (data not shown). The enhancement of the outward currents in HEKKBK $\alpha\beta$ 1 was completely removed by washout of diCl-DHAA (Fig. 1B). The relationship between concentrations of diCl-DHAA and corresponding responses is summarized in Fig. 1C. The increase

in outward current density by diCl-DHAA was significant at a concentration of 0.1 μ M and higher. Taking the current density at $+10$ mV in the control as unity, the relative amplitude of peak outward currents in the presence of 0.1, 0.3, and 1 μ M diCl-DHAA was also plotted against concentration (Fig. 1C).




Fig. 2. Effects of diCl-DHAA on I-V relationship in BK $\alpha\beta$ 1. A, each single HEKKBK $\alpha\beta$ 1 was depolarized from -60 mV by 10-mV steps for 150 ms under whole-cell voltage-clamp mode. The outward currents elicited by depolarization was markedly enhanced by application of 1 μ M diCl-DHAA and then reduced by the addition of 100 nM iberiotoxin. B, I-V relationships were obtained in the control (open circles), in the presence of diCl-DHAA (closed circles), and after the addition of 100 nM iberiotoxin (closed triangles) in experiments such as typically shown in A. Number of examples is seven. C, voltage dependence of the potentiation of the outward current by diCl-DHAA was reevaluated from the data shown in B. The relative potentiation of the outward currents by 1 μ M diCl-DHAA was plotted against test potentials by taking that at $+30$ mV as unity. *, $p < 0.05$ and **, $p < 0.01$ versus unity at $+30$ mV.




Fig. 3. Effects of diCl-DHAA on single BK α channel currents recorded using an inside-out patch-clamp techniques. **A**, single channel currents were recorded at +40 mV in a patch from HEK BK α under symmetrical 140 mM K $^+$ conditions. The free Ca $^{2+}$ in the bathing solution was adjusted to pCa = 7.0. The original current traces were recorded before and after application of 1 μM diCl-DHAA and after the washout of diCl-DHAA. A closed triangle on the left side of each trace indicates the zero current level. **B**, amplitude histograms in the control, in the presence of 1 μM diCl-DHAA, and after the washout were obtained from the recordings shown in **A**. The ordinate expresses the relative area (percentage) at the corresponding amplitude in each bin (0.2 pA). **C**, summarized data demonstrate the relationship between concentrations of diCl-DHAA and P_o of BK α . P_o was calculated from the histogram shown in **B** as the relative time spent at open state based on the total number of BK α channels in the patch, which was determined by elevating Ca $^{2+}$ concentration to pCa = 3.5 (open columns). The relative P_o was obtained taking the P_o in the absence of diCl-DHAA as unity (closed columns). Number of examples is six. */#, $p < 0.05$ and **/##, $p < 0.01$ versus control.

Voltage Dependence of diCl-DHAA-Induced Enhancement of BK Channel Currents. In Fig. 2, the voltage-dependent enhancement of BK channel currents was examined by analyzing effects of diCl-DHAA on the current-voltage (I-V) relationship. HEK BK α 1 was depolarized from a holding potential of -60 mV to test potentials in the range between -50 and +30 mV with 10-mV steps (Fig. 2A). Application of 1 μM diCl-DHAA increased the currents at any test potential. Addition of 100 nM iberiotoxin, a specific BK channel blocker, removed the enhancement of outward currents, supporting that the action of diCl-DHAA was selective to BK channel currents. Figure 2B summarizes the relationships between current density of peak outward currents and test potentials in the absence and presence of 1 μM diCl-DHAA and after addition of 100 nM iberiotoxin. The current




Fig. 4. Structure-activity relationships of abietic acid derivatives on P_o of BK α . **A**, chemical structures of test compounds pimaric acid, abietic acid, dehydroabietic acid, and diCl-DHAA. **B**, concentration-response relationships for compounds listed in **A**. Effects of abietic acid (open triangles), DHAA (open squares), and diCl-DHAA (closed circles) were examined in experiments identical to that shown in Fig. 3 for diCl-DHAA. The data for pimaric acid (closed squares) are taken from a previous study (Imaizumi et al., 2002). The relative P_o was determined taking the P_o in the absence of compounds as unity (a dotted line). Means \pm S.E.M. are shown by symbols and vertical bars, respectively. Number of experiments is four to six for each compound.

density at +30 mV was increased from 39.13 ± 7.34 to 142.55 ± 30.81 pA/pF ($n = 7$; $p < 0.01$). In Fig. 2C, the voltage dependence of diCl-DHAA-induced enhancement of BK channel currents was determined as the relationship between the relative potentiation of outward currents and test potentials by taking the potentiation at +30 mV in the presence of 1 μM diCl-DHAA as unity. The potentiation at -20 and -10 mV was significantly greater than that at +30 mV (4.48 ± 1.15 and 3.80 ± 1.09 times at -20 and -10 mV, respectively, $p < 0.01$, versus unity at +30 mV).

Activation of Single BK α Channel Current by diCl-DHAA and Related Compounds. Effects of diCl-DHAA on single BK α channel currents were examined in excised inside-out patch configuration. The bathing and pipette solution contained symmetrical 140 mM K $^+$. The free Ca $^{2+}$ concentration in the bathing solution was pCa7. Under these conditions, the unitary current amplitude and open probability (P_o) at +40 mV was 10.1 ± 0.2 pA and 0.0028 ± 0.0005 ($n = 6$), respectively. The application of 1 μM diCl-DHAA increased channel activity without change in the unitary