

- Wera, S., and Hemmings, B. A. (1995). Serine/threonine protein phosphatases. *Biochem. J.* **311**, 17–29.
- Werries, E., and Thurn, I. (1989). Breakdown of glucopolysaccharides in *Entamoeba histolytica* by phosphorylase. *J. Protozool.* **36**, 607–612.
- Werries, E., Franz, A., and Geisemeyer, S. (1990). Detection of glycogen-debranching system in trophozoites of *Entamoeba histolytica*. *J. Protozool.* **37**, 576–580.
- West, C. M. (2003). Comparative analysis of spore coat formation, structure, and function in *Dictyostelium*. *Int. Rev. Cytol.* **222**, 237–293.
- West, C. M., van der Wel, H., Coutinho, P. M., and Henrissat, B. (2005). Glycosyltransferase genomics in *Dictyostelium*. In “*Dictyostelium Genomics*” (W. F. Loomis and A. Kuspa, eds.), pp. 235–264. Horizon Bioscience, Norfolk, UK.
- Whitney, J. A., Gomez, M., Sheff, D., Kreis, T. E., and Mellman, I. (1995). Cytoplasmic coat proteins involved in endosome function. *Cell* **83**, 703–713.
- WHO (1998). “The world health report 1998. Life in the 21st century: A vision for all.” World Health Organization, Geneva, Switzerland.
- Willhoeft, U., Buss, H., and Tannich, E. (1999). Analysis of cDNA expressed sequence tags from *Entamoeba histolytica*: Identification of two highly abundant polyadenylated transcripts with no overt open reading frames. *Protist* **150**, 61–70.
- Willhoeft, U., and Tannich, E. (1999). The electrophoretic karyotype of *Entamoeba histolytica*. *Mol. Biochem. Parasitol.* **99**, 41–53.
- Willhoeft, U., Buss, H., and Tannich, E. (2002). The abundant polyadenylated transcript 2 DNA sequence of the pathogenic protozoan parasite *Entamoeba histolytica* represents a nonautonomous non-long-terminal-repeat retrotransposon-like element which is absent in the closely related nonpathogenic species *Entamoeba dispar*. *Infect. Immun.* **70**, 6798–6804.
- Winder, S. J., and Ayscough, K. R. (2005). Actin-binding proteins. *J. Cell Sci.* **118**, 651–654.
- Winkelmann, J., Leippe, M., and Bruhn, H. (2006). A novel saposin-like protein of *Entamoeba histolytica* with membrane-fusogenic activity. *Mol. Biochem. Parasitol.* **147**, 85–94.
- Wittenberg, C., Sugimoto, K., and Reed, S. I. (1990). G1-specific cyclins of *S. cerevisiae*: Cell cycle periodicity, regulation by mating pheromone, and association with the p34CDC28 protein kinase. *Cell* **62**, 225–237.
- Wohlschlegel, J. A., Dhar, S. K., Prokhorova, T. A., Dutta, A., and Walter, J. C. (2002). Xenopus Mcm10 binds to origins of DNA replication after Mcm2–7 and stimulates origin binding of Cdc45. *Mol. Cell* **9**, 233–240.
- Wood, Z. A., Poole, L. B., and Karplus, P. A. (2003a). Peroxiredoxin evolution and the regulation of hydrogen peroxide signaling. *Science* **300**, 650–653.
- Wood, Z. A., Schroder, E., Robin Harris, J., and Poole, L. B. (2003b). Structure, mechanism and regulation of peroxiredoxins. *Trends Biochem. Sci.* **28**, 32–40.
- Woods, S. A., Schwartzbach, S. D., and Guest, J. R. (1988). Two biochemically distinct classes of fumarase in *Escherichia coli*. *Biochim. Biophys. Acta* **954**, 14–26.
- Wostmann, C., Tannich, E., and Bakker-Grunwald, T. (1992). Ubiquitin of *Entamoeba histolytica* deviates in 6 amino acid residues from the consensus of all other known ubiquitins. *FEBS Lett.* **308**, 54–58.
- Wu, G., and Müller, M. (2003). Glycogen phosphorylase sequences from the amitochondriate protists, *Trichomonas vaginalis*, *Mastigamoeba balamuthi*, *Entamoeba histolytica* and *Giardia intestinalis*. *J. Eukaryot. Microbiol.* **50**, 366–372.
- Xu, Y., Moseley, J. B., Sagot, I., Poy, F., Pellman, D., Goode, B. L., and Eck, M. J. (2004). Crystal structures of a Formin Homology-2 domain reveal a tethered dimer architecture. *Cell* **116**, 711–723.
- Yadava, N., Chandok, M. R., Prasad, J., Bhattacharya, S., Sopory, S. K., and Bhattacharya, A. (1997). Characterization of EhCaBP, a calcium binding protein of *Entamoeba histolytica* and its binding proteins. *Mol. Biochem. Parasitol.* **84**, 69–82.

- Yamawaki-Kataoka, Y., Tamaoki, T., Choe, H. R., Tanaka, H., and Kataoka, T. (1989). Adenylate cyclases in yeast: A comparison of the genes from *Schizosaccharomyces pombe* and *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* **86**, 5693–5697.
- Yang, B., Steegmaier, M., Gonzalez, L. C., Jr., and Scheller, R. H. (2000). nSec1 binds a closed conformation of syntaxin1A. *J. Cell Biol.* **148**, 247–252.
- Yang, W., Li, E., Kairong, T., and Stanley, S. L., Jr. (1994). *Entamoeba histolytica* has an alcohol dehydrogenase homologous to the multifunctional *adhE* gene product of *Escherichia coli*. *Mol. Biochem. Parasitol.* **64**, 253–260.
- Yasui, K., Ishiguro, A., and Ishihama, A. (1998). Location of subunit-subunit contact sites on RNA polymerase II subunit 3 from the fission yeast *Schizosaccharomyces pombe*. *Biochemistry* **37**, 5542–5548.
- Yeo, M., Lin, P. S., Dahmus, M. E., and Gill, G. N. (2003). A novel RNA polymerase II C-terminal domain phosphatase that preferentially dephosphorylates serine 5. *J. Biol. Chem.* **278**, 26078–26085.
- Yin, Q. Y., de Groot, P. W., Dekker, H. L., de Jong, L., Klis, F. M., and de Koster, C. G. (2005). Comprehensive proteomic analysis of *Saccharomyces cerevisiae* cell walls: Identification of proteins covalently attached via glycosylphosphatidylinositol remnants or mild alkali-sensitive linkages. *J. Biol. Chem.* **280**, 20894–20901.
- Yoshihisa, T., Barlowe, C., and Schekman, R. (1993). Requirement for a GTPase-activating protein in vesicle budding from the endoplasmic reticulum. *Science* **259**, 1466–1468.
- Young, R. A. (1991). RNA polymerase II. *Annu. Rev. Biochem.* **60**, 689–715.
- Yu, X., and Warner, J. R. (2001). Expression of a micro-protein. *J. Biol. Chem.* **276**, 33821–33825.
- Zerial, M., and McBride, H. (2001). Rab proteins as membrane organizers. *Nat. Rev. Mol. Cell Biol.* **2**, 107–117.
- Zhang, J. (2000). Protein-length distributions for the three domains of life. *Trends Genet.* **16**, 107–109.
- Zhang, W. W., Shen, P. S., Descoteaux, S., and Samuelson, J. (1994). Cloning and expression of the gene for an NADP⁺-dependent aldehyde dehydrogenase of *Entamoeba histolytica*. *Mol. Biochem. Parasitol.* **63**, 157–161.
- Zhang, Z., Wang, L., Seydel, K. B., Li, E., Ankri, S., Mirelman, D., and Stanley, S. L., Jr. (2000). *Entamoeba histolytica* cysteine proteinases with interleukin-1 beta converting enzyme (ICE) activity cause intestinal inflammation and tissue damage in amoebiasis. *Mol. Microbiol.* **37**, 542–548.
- Zhang, Z., Duchene, M., and Stanley, S. L., Jr. (2002). A monoclonal antibody to the amebic lipophosphoglycan-proteophosphoglycan antigens can prevent disease in human intestinal xenografts infected with *Entamoeba histolytica*. *Infect. Immun.* **70**, 5873–5876.
- Zhang, Z. Y., and VanEtten, R. L. (1991). Pre-steady-state and steady-state kinetic analysis of the low molecular weight phosphotyrosyl protein phosphatase from bovine heart. *J. Biol. Chem.* **266**, 1516–1525.
- Zuo, X., and Coombs, G. H. (1995). Amino acid consumption by the parasitic, amoeboid protists *Entamoeba histolytica* and *E. invadens*. *FEMS Microbiol. Lett.* **130**, 253–258.